Asymmetric Synthesis of the Tetraponerine Alkaloids

Stephen G. Davies, Ai M. Fletcher, Ian T. T. Houlsby, Paul M. Roberts, and James E. Thomson

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.7b00837 • Publication Date (Web): 24 May 2017

Downloaded from <http://pubs.acs.org> on June 6, 2017

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

Asymmetric Synthesis of the Tetraponerine Alkaloids

Stephen G. Davies,* Ai M. Fletcher, Ian T. T. Houlsby

Paul M. Roberts, and James E. Thomson

Department of Chemistry, Chemistry Research Laboratory, University of Oxford,

Mansfield Road, Oxford, OX1 3TA, U.K.

steve.davies@chem.ox.ac.uk

The asymmetric syntheses of all eight tetraponerine alkaloids (T1–T8) were achieved using the diastereoselective conjugate additions of lithium amide reagents in the key stereodefining steps. Conjugate addition of either lithium (*R*)-*N*-allyl-*N*-(α -methylbenzyl)amide or lithium (*R*)-*N*-(but-3-en-1-yl)-*N*-(α -methylbenzyl)amide to *tert*-butyl sorbate was followed by ring-closing metathesis of the resultant *N*-alkenyl β -amino esters, reduction to the corresponding aldehydes, and reaction with *tert*-butyl (triphenylphosphoranylidene)acetate. Subsequent conjugate addition of the requisite antipode of lithium *N*-benzyl-*N*-(α -methylbenzyl)amide to the resultant α,β -unsaturated esters gave a range of diamines for elaboration to T1–T8 via a sequence involving reduction of the ester moiety to give the corresponding aldehyde, olefination, tandem hydrogenation/hydrogenolysis, and cyclisation upon reaction with 4-bromobutanal to give the tricyclic skeleton.

Introduction

Tetraponerines 1–8 (which are also known as T1–T8, respectively) are a series of tricyclic aminals first isolated from the paralyzing venom of the New Guinean pseudomyrmecine ant *Tetraponera* *sp.*¹ These alkaloids represent the major constituents of the contact poison secreted by host ants as a defence against enemy colonies: when smeared with the contact poison enemies display signs of paralysis and ultimately death, presumably by inhibition of nicotinic acetylcholine receptors.² The structures of the tetraponerine alkaloids are based on either a decahydrodipyrrolo[1,2-*a*:1',2'-*c*]pyrimidine skeleton **1** (for tetraponerines 1, 2, 5 and 6) or a decahydro-5*H*-pyrido[1,2-*c*]pyrrolo[1,2-*a*]pyrimidine skeleton **2** (for tetraponerines 3, 4, 7

and 8), with a C(5)-alkyl substituent being present in all cases. Within each heterocyclic series, the alkaloids differ in their configurations at C(5) and also in the identity of the C(5)-alkyl substituent (Fig. 1). (+)-Tetraponerine 8 is the only member of this family for which an X-ray crystal structure has been obtained to confirm both its gross structure and relative configuration.¹ The absolute configurations of tetraponerines 3–8 have been established by separate chemical syntheses and comparison of the specific rotations for the natural and synthetic material.^{3–5} Synthetic samples of (+)-tetraponerine 1 and (+)-tetraponerine 2 have also been prepared,^{5,6} although the absolute configurations of the natural materials have not been unambiguously established as only very small amounts were obtained from natural sources.^{4,7} The unusual tricyclic aminal structure of the tetraponerine alkaloids, as well as their potent insecticidal activity, has meant that they have become popular targets for total synthesis. Since their isolation in 1987, there have been several syntheses of these alkaloids in both racemic⁸ and enantiopure^{3–6,9} form.^{10,11} Herein, we report total syntheses of all eight tetraponerine alkaloids using our diastereoselective lithium amide conjugate addition methodology¹² in the key stereodefining steps.

FIGURE 1. The structures of the tetraponerine alkaloids T1–T8.

Our divergent synthetic strategy towards the tetraponerine alkaloids involved two iterative lithium amide conjugate additions to install the requisite stereochemistry. In the first instance, conjugate addition of either lithium (*R*)-*N*-allyl-*N*-(α -methylbenzyl)amide (*R*)-**4** ($n = 1$) or lithium (*R*)-*N*-(but-3-en-1-yl)-*N*-(α -methylbenzyl)amide (*R*)-**5** ($n = 2$) to *tert*-butyl sorbate **3**,^{13–15} followed by ring-closing metathesis of the resultant *N*-alkenyl β -amino ester **6** ($n = 1$ or 2) and homologation would give the key α,β -unsaturated ester **7** ($n = 1$ or 2). Subsequent conjugate addition of the requisite antipode of lithium *N*-benzyl-*N*-(α -methylbenzyl)amide **8** to α,β -unsaturated ester **7** ($n = 1$ or 2) would then provide a range of diamines **9** and **10** ($n = 1$ or 2 in both cases) for elaboration. In each case, a sequence involving reduction of the ester moiety to the corresponding aldehyde, olefination, and tandem hydrogenation/hydrogenolysis would produce the corresponding diamines **11** or **12** ($n = 1$ or 2 in both cases), which have previously been converted into the

tetraponerine alkaloids by Gonzalez-Gomez *et al.*^{6,9d} upon reaction with 4-bromobutanol (Fig. 2). As part of their study, Gonzalez-Gomez *et al.* proposed that the configuration at the aminal functionality within each of the tetraponerine alkaloids is susceptible to equilibration via the corresponding iminium ion and that the natural products may therefore possess the most thermodynamically favourable relative configurations with respect to epimerisation at the aminal centre.

FIGURE 2. Proposed synthetic strategy to access all eight tetraponerine alkaloids.

Results and Discussion

The tetraponerine alkaloids based upon a decahydro-5*H*-pyrido[1,2-*c*]pyrrolo[1,2-*a*]pyrimidine skeleton **2** (i.e., tetraponerines **3**, **4**, **7** and **8**) were targeted initially. Tetrahydropyridine **14** was prepared via a known sequence of reactions,^{13,14} involving conjugate addition of lithium (*R*)-*N*-(but-3-en-1-yl)-*N*-(α -methylbenzyl)amide (*R*)-**5** to *tert*-butyl sorbate **3**,¹⁶ which gave β -amino ester **13** as a single diastereoisomer (>99:1 dr) in 69% yield. Subsequent ring-closing metathesis of **13** gave tetrahydropyridine **14** in 83% yield.^{13,14} A superior overall yield of **14** from *tert*-butyl sorbate **3** was obtained by omitting the purification of the intermediate β -amino ester **13**; following this sequence, tetrahydropyridine **14** was isolated in 73% yield (from **3**) and >99:1 dr. Reduction of **14** with LiAlH_4 gave the corresponding alcohol **15** in 94% yield as a single diastereoisomer. Swern oxidation of primary alcohol **15** followed by Wittig reaction of the resultant aldehyde with $\text{Ph}_3\text{P}=\text{CHCO}_2^t\text{Bu}$ gave a 93:7 mixture of diastereoisomeric products **16** and **17**, respectively. Purification of the crude reaction mixture gave the major product **16** in 46% yield and >99:1 dr [(*E*):(*Z*)], and the minor product **17** in 3% yield and >99:1 dr [(*Z*):(*E*)]. However, a superior yield of **16** was obtained from

the reduction of β -amino ester **14** with DIBAL-H followed by in situ Wittig reaction of the intermediate aldehyde with $\text{Ph}_3\text{P}=\text{CHCO}_2^t\text{Bu}$, which gave an 80:20 mixture of **16** and **17**, respectively, from which **16** was isolated in 72% yield and >99:1 dr [(*E*):(*Z*)], and **17** was isolated in 14% yield and >99:1 dr [(*Z*):(*E*)] (Scheme 1).

SCHEME 1. Preparation of tetrahydropyridine **16**

The inherent level of substrate control offered by chiral α,β -unsaturated ester **16** upon reaction with a lithium amide reagent was probed upon conjugate addition of lithium *N*-isopropyl-*N*-benzylamide **18**, which we have previously employed as an achiral model for the conjugate addition of either antipode of enantiopure lithium *N*-benzyl-*N*-(α -methylbenzyl)amide **8**.¹⁷ Reaction of **16** with the achiral lithium amide reagent **18** gave a 50:50 mixture of the diastereoisomeric β -amino ester adducts **19** and **20**, which were isolated in 96% combined yield. This result suggests that α,β -unsaturated ester **16** elicits essentially no diastereoselectivity in the conjugate addition reaction, and is consistent with the low levels of substrate control observed upon conjugate addition of lithium amides to substrates bearing remote stereogenic centres.¹⁸ The conjugate addition of lithium (*R*)-*N*-benzyl-*N*-(α -methylbenzyl)amide (*R*)-**8** to α,β -unsaturated ester **16** gave β -amino ester **21** in >95:5 dr, and conjugate addition of lithium (*S*)-*N*-benzyl-*N*-(α -methylbenzyl)amide (*S*)-**8** to **16** gave β -amino ester **22** in >95:5 dr. Following purification of the crude reaction mixtures, **21** was isolated in 97% yield and >99:1 dr, and **22** was isolated in 94% yield and >99:1 dr (Scheme 2). The configurations of the newly formed stereogenic centres within **21** and **22** were assigned on the assumption that conjugate addition proceeds under the dominant stereocontrol of the lithium amide

reagent **8** (given the total lack of substrate control observed upon conjugate addition of lithium *N*-isopropyl-*N*-benzylamide **18** to α,β -unsaturated ester **16**), and is consistent with the formation of **21** and **22** as single diastereoisomers in both cases. These stereochemical assignments were later confirmed upon derivatisation to the target tetraoponerine alkaloids.

SCHEME 2. Conjugate additions to **16**

The elaboration of **21** to (+)-tetraoponerine 4 (**27**) was achieved via reduction of the ester moiety within **21** with DIBAL-H followed by Wittig reaction of the resultant aldehyde with $\text{Ph}_3\text{P}=\text{CH}_2$, which gave olefin **23** in 62% and >99:1 dr. Tandem Pd-mediated hydrogenation/hydrogenolysis of **23** in a mixture of MeOH, H_2O and AcOH gave diamine **25** in 73% isolated yield and >99:1 dr. Finally, reaction of **25** with 4-bromobutanol and K_2CO_3 in CH_2Cl_2 , according to the literature protocol,^{6,9d} gave (+)-tetraoponerine 4 (**27**) in 68% yield and >99:1 dr. The spectroscopic data and specific rotation for this sample of (+)-tetraoponerine 4 (**27**) were in good agreement with literature values { $[\alpha]_{\text{D}}^{25} +90.8$ (c 1.0 in CHCl_3); lit.⁴ for a sample isolated from the natural source $[\alpha]_{\text{D}}^{20} +94$ (c 0.2 in CHCl_3); lit.^{9b} $[\alpha]_{\text{D}}^{20} +96$ (c 2.0 in CHCl_3)}, thereby confirming the assigned configurations of the synthetic precursors **21**, **23** and **25**. Repetition of this hydrogenation/hydrogenolysis and ring-closure reaction sequence, but omitting the purification of the intermediate diamine **25**, gave (+)-tetraoponerine 4 (**27**) in 54% overall yield (from **23**), representing a marginal improvement over the stepwise protocol (i.e., 50% overall yield from **23**). (+)-Tetraoponerine 8 (**28**) was also prepared from **21** in a directly analogous manner: one-pot reduction/Wittig reaction of **21** gave

olefin **24** in 62% yield and >95:5 dr [(Z):(E)].¹⁹ Subsequent tandem hydrogenation/hydrogenolysis of **24** gave diamine **26** (although in this case only it was necessary to re-subject the hydrogenation/hydrogenolysis reaction using Pd(OH)₂ as the catalyst to effect complete *N*-debenzylation). Diamine **26** was then immediately converted into (+)-tetraponerine 8 (**28**) upon treatment with 4-bromobutanol.^{6,9d} Following purification of the crude reaction mixture, (+)-tetraponerine 8 (**28**) was isolated as a single diastereoisomer (>99:1 dr) in 67% yield from **24** (Scheme 3). The spectroscopic data, melting point and specific rotation for this sample of (+)-tetraponerine 8 (**28**) were all in good agreement with literature values {mp 38–40 °C; lit.⁵ mp 40 °C; [α]_D²⁵ +93.0 (c 1.0 in CHCl₃); lit.⁴ for a sample isolated from the natural source [α]_D²⁰ +102 (c 0.2 in CHCl₃); lit.^{9b} [α]_D²⁰ +101 (c 2.0 in CHCl₃)}, thereby confirming the assigned configurations of the synthetic precursors **21**, **24** and **26**. The asymmetric syntheses of (+)-tetraponerine 4 (**27**) and (+)-tetraponerine 8 (**28**) were therefore achieved in 15.5 and 19.3% overall yield, respectively, in eight steps from commercially available starting materials.

SCHEME 3. Preparation of (+)-tetraponerine 4 and (+)-tetraponerine 8

Elaboration of the diastereoisomeric substrate **22** to (+)-tetraponerine 3 (**33**) and (+)-tetraponerine 7 (**34**) was also evaluated. Reduction of **22** with DIBAL-H followed by Wittig reaction of the resultant aldehyde with the requisite ylid reagent Ph₃P=CHR gave either **29** (R = H) or **30** (R = Et); after chromatographic purification of the crude reaction mixtures, **29** was isolated in 54% yield as a single diastereoisomer and **30** was isolated in 47% yield and >95:5 dr [(Z):(E)].²⁰ Tandem hydrogenation/hydrogenolysis of **29** and **30** gave the corresponding diamines **31** and **32**, which were immediately reacted with 4-bromobutanol^{6,9d} to give (+)-tetraponerine 3 (**33**) and (+)-tetraponerine 7 (**34**) in 60 and 44% yield, respectively, as single diastereoisomers (>99:1 dr) in each case (Scheme 4). The

spectroscopic data and specific rotations for these samples of (+)-tetraponerine 3 (**33**) and (+)-tetraponerine 7 (**34**) were in good agreement with literature values {for (+)-tetraponerine 3 (**33**): $[\alpha]_{\text{D}}^{25} +30.9$ (c 1.0 in CHCl_3); lit.⁴ for a sample isolated from the natural source $[\alpha]_{\text{D}}^{20} +27$ (c 0.07 in CHCl_3); lit.⁵ $[\alpha]_{\text{D}}^{20} +31$ (c 3.1 in CHCl_3); lit.^{9d} $[\alpha]_{\text{D}}^{20} +35$ (c 0.49 in CHCl_3); for (+)-tetraponerine 7 (**34**): $[\alpha]_{\text{D}}^{25} +30.4$ (c 0.8 in CHCl_3); lit.⁴ for a sample isolated from the natural source $[\alpha]_{\text{D}}^{20} +30$ (c 0.22 in CHCl_3); lit.^{9b} $[\alpha]_{\text{D}}^{20} +29.5$ (c 2.2 in CHCl_3)}, thereby confirming the assigned configurations of the synthetic precursors **22** and **29–32**. The asymmetric syntheses of (+)-tetraponerine 3 (**33**) and (+)-tetraponerine 7 (**34**) were therefore achieved in 14.6 and 9.3% overall yield, respectively, in eight steps from commercially available starting materials.

SCHEME 4. Preparation of (+)-tetraponerine 3 and (+)-tetraponerine 7

The remaining tetraponerine alkaloids in the series, based on a decahydrodipyrrolo[1,2- α :1',2'-c]pyrimidine scaffold **1** (i.e., tetraponerines 1, 2, 5 and 6), were targeted next. Dihydropyrrole **36** was prepared via a known sequence of reactions,^{13,15} involving conjugate addition of lithium (*R*)-*N*-allyl-*N*-(α -methylbenzyl)amide (*R*)-**4** to *tert*-butyl sorbate **3**,¹⁶ which gave β -amino ester **35** in 89% yield as a single diastereoisomer ($>99:1$ dr). Subsequent ring-closing metathesis of **35** gave dihydropyrrole **36** in 86% yield and $>99:1$ dr.^{13,15} In this case, it was found that hydrogenation of **36** with Wilkinson's catalyst $[\text{RhCl}(\text{PPh}_3)_3]$ ^{13,15a,b} to give pyrrolidine **37** was necessary to avoid oxidative side reactions, leading to various pyrrole containing species, in the subsequent steps.²¹ Reduction of the ester moiety within pyrrolidine **37** with DIBAL-H followed by in situ Wittig reaction of the corresponding aldehyde with $\text{Ph}_3\text{P}=\text{CHCO}_2^t\text{Bu}$ gave a 75:25 mixture of **38** and **39**, respectively. Following purification of the crude reaction mixture, an 86:14 mixture of **38** and **39**, respectively, was isolated in 66% combined yield in addition to a 9:91 mixture of **38** and **39** in 13% combined yield. Upon repetition of the reaction sequence, hydrogenation of dihydropyrrole **36** followed by immediate reduction of the crude reaction mixture with

DIBAL-H and Wittig reaction of the resultant aldehyde gave **38** in 50% yield (from **36**) and 98:2 dr (Scheme 5).

SCHEME 5. Preparation of pyrrolidine **38**

The inherent level of substrate control offered by α,β -unsaturated ester **38** (98:2 dr [(E):(Z)]) was probed upon conjugate addition of lithium *N*-isopropyl-*N*-benzylamide **18**, which gave a 50:50 mixture of the diastereoisomeric β -amino ester adducts **40** and **41**. Following purification of the crude reaction mixture, **40** and **41** were isolated as a 50:50 mixture in 70% combined yield. This result suggests that α,β -unsaturated ester **38** elicits essentially no diastereoselectivity in the conjugate addition reaction, as was also the case for the homologous tetrahydropyridine containing substrate **16**, and is again consistent with the low levels of substrate control observed upon conjugate addition of lithium amides to other substrates bearing remote stereogenic centres.¹⁸ The conjugate addition of lithium (*R*)-*N*-benzyl-*N*-(α -methylbenzyl)amide (*R*)-**8** to α,β -unsaturated ester **38** (98:2 dr [(E):(Z)]) gave β -amino ester **42** in 98% yield and >99:1 dr, whereas conjugate addition of lithium (*S*)-*N*-benzyl-*N*-(α -methylbenzyl)amide (*S*)-**8** to **38** (98:2 dr [(E):(Z)]) gave β -amino ester **43** in 92% isolated yield and >99:1 dr (Scheme 6). The configurations of the newly formed stereogenic centres within the conjugate addition products **42** and **43** were assigned, as before, on the assumption that conjugate addition proceeds under the dominant stereocontrol of the lithium amide reagent in both cases; these assignments were later confirmed upon derivatisation to the target alkaloids. Conjugate addition of lithium amide (*R*)-**8** to an 86:14 mixture of (*E*)-**38** and (*Z*)-**39**, respectively, gave an 86:14 mixture of β -amino ester **42** and a species which was tentatively assigned as the corresponding β,γ -

unsaturated derivative **44**; there was no evidence of either (*E*)-**38** or (*Z*)-**39** in the ^1H NMR spectrum of the crude reaction mixture. As β -amino ester **42** and β,γ -unsaturated ester **44** were only partially separable upon attempted chromatographic purification, this result confirmed that it was necessary to enrich the diastereoisomeric purity of the α,β -unsaturated ester prior to attempting the conjugate addition reaction.

SCHEME 6. Conjugate additions to **38**

Reduction of **42** with DIBAL-H followed by Wittig reaction of the resultant aldehyde with the requisite ylid reagent $\text{Ph}_3\text{P}=\text{CHR}$ gave either **45** ($\text{R} = \text{H}$) or **46** ($\text{R} = \text{Et}$); after chromatographic purification of the crude reaction mixtures, **45** was isolated in 76% yield and >99:1 dr, and **46** was isolated in 75% yield and >95:5 dr [(*Z*):(*E*)].²⁰ Tandem hydrogenation/hydrogenolysis of **45** and **46** gave the corresponding diamines **47** and **48**, which were immediately reacted with 4-bromobutanol^{6,9d} to give (+)-tetraoponerine 2 (**49**) and (+)-tetraoponerine 6 (**50**) in 37 and 54% yield, respectively, as single diastereoisomers (>99:1 dr) in each case (Scheme 7). For (+)-tetraoponerine 2 (**49**), the specific rotation was consistent with literature values $\{[\alpha]_{\text{D}}^{25} + 44.2$ (c 1.0 in CHCl_3); lit.⁵ $[\alpha]_{\text{D}}^{20} + 36$ (c 1.79 in CHCl_3); lit.⁶ $[\alpha]_{\text{D}}^{20} + 47$ (c 0.232 in CHCl_3)}; however, for (+)-tetraoponerine 6 (**50**) the specific rotations of our synthetic samples were consistently larger than literature values $\{[\alpha]_{\text{D}}^{25} + 66.3$ (c 1.0 in CHCl_3); lit.⁴ for a sample isolated from the natural source $[\alpha]_{\text{D}}^{20} + 35$ (c 0.15 in CHCl_3); lit.⁶ $[\alpha]_{\text{D}}^{20} + 40$ (c 0.75 in CHCl_3)}. Despite this discrepancy,²² the spectroscopic data for both samples of (+)-tetraoponerine 2 (**49**) and (+)-tetraoponerine 6 (**50**) were in good agreement with literature values, thereby confirming the assigned configurations of the synthetic precursors **42** and **45–48**. The

asymmetric syntheses of (+)-tetraponerine 2 (**49**) and (+)-tetraponerine 6 (**50**) were therefore achieved in 9.6 and 13.8% overall yield, respectively, in nine steps from commercially available starting materials.

SCHEME 7. Preparation of (+)-tetraponerine 2 and (+)-tetraponerine 6

Reduction of the diastereoisomeric substrate **43** with DIBAL-H followed by Wittig reaction of the resultant aldehyde with the requisite ylid reagent $\text{Ph}_3\text{P}=\text{CHR}$ gave either **51** (R = H) or **52** (R = Et); after chromatographic purification of the crude reaction mixtures, **51** was isolated in 40% yield as a single diastereoisomer and **52** was isolated in 48% yield and >95:5 dr [(Z):(E)].²⁰ Tandem hydrogenation/hydrogenolysis of **51** and **52** gave the corresponding diamines **53** and **54**, which were immediately reacted with 4-bromobutanol^{6,9d} to give (+)-tetraponerine 1 (**55**) and (+)-tetraponerine 5 (**56**) in 15 and 14% yield, respectively, as single diastereoisomers (>99:1 dr) in each case (Scheme 8). The spectroscopic data and specific rotations for these samples of (+)-tetraponerine 1 (**55**) and (+)-tetraponerine 5 (**56**) were in good agreement with literature values {for (+)-tetraponerine 1 (**55**): $[\alpha]_{\text{D}}^{25} +14.4$ (c 0.13 in CHCl_3); lit.⁵ $[\alpha]_{\text{D}}^{20} +11$ (c 0.14 in CHCl_3); lit.⁶ $[\alpha]_{\text{D}}^{20} +14$ (c 0.498 in CHCl_3); for (+)-tetraponerine 5 (**56**): $[\alpha]_{\text{D}}^{25} +12.4$ (c 0.13 in CHCl_3); lit.⁴ for a sample isolated from the natural source $[\alpha]_{\text{D}}^{20} +10$ (c 0.2 in CHCl_3); lit.⁵ $[\alpha]_{\text{D}}^{20} +10$ (c 0.24 in CHCl_3); lit.⁶ $[\alpha]_{\text{D}}^{20} +14$ (c 1.6 in CHCl_3)}, thereby confirming the assigned configurations of the synthetic precursors **43** and **51–54**. The asymmetric syntheses of (+)-tetraponerine 1 (**55**) and (+)-tetraponerine 5 (**56**) were therefore achieved in 1.9 and 2.2% overall yield, respectively, in nine steps from commercially available starting materials.

SCHEME 8. Preparation of (+)-tetraponerine 1 and (+)-tetraponerine 5

Conclusion

In conclusion, the conjugate addition of either lithium (*R*)-*N*-allyl-*N*-(α -methylbenzyl)amide or lithium (*R*)-*N*-(but-3-en-1-yl)-*N*-(α -methylbenzyl)amide to *tert*-butyl sorbate was followed by ring-closing metathesis of the resultant *N*-alkenyl β -amino esters and homologation. Subsequent conjugate addition of the requisite antipode of lithium *N*-benzyl-*N*-(α -methylbenzyl)amide to the resultant α,β -unsaturated esters gave a range of diamines for elaboration to all eight tetraponerine alkaloids (T1–T8) via a sequence involving reduction of the ester moiety to the corresponding aldehyde, olefination, tandem hydrogenation/hydrogenolysis, and reaction with 4-bromobutanol to give the tricyclic skeleton. In each case, the target alkaloids were isolated enantiomerically pure as single diastereoisomers in nine steps or fewer from commercially available sorbic acid.

Experimental Section

General experimental details. All reactions involving organometallic or other moisture-sensitive reagents were carried out under a nitrogen atmosphere using standard vacuum line techniques and glassware that was flame dried and cooled under nitrogen before use. Solvents were dried according to the procedure outlined by Grubbs and co-workers.²³ BuLi was purchased as a solution in hexanes and titrated against diphenylacetic acid before use. All other reagents were used as supplied without prior purification. Organic layers were dried over Na₂SO₄. Thin layer chromatography was performed on aluminium plates coated with 60 F₂₅₄ silica. Plates were visualised using UV light (254 nm), 1% aq KMnO₄ or Dragendorff's reagent. Flash column chromatography was performed on Kieselgel 60 silica. Melting points are uncorrected. Specific rotations are reported in 10⁻¹ deg cm² g⁻¹ and concentrations in g/100 mL. IR spectra were recorded

using an ATR module. Selected characteristic peaks are reported in cm^{-1} . NMR spectra were recorded in the deuterated solvent stated. Spectra were recorded at rt unless otherwise stated. The field was locked by external referencing to the relevant deuterium resonance. ^1H - ^1H COSY, ^1H - ^{13}C HMQC, and ^1H - ^{13}C HMBC analyses were used to establish atom connectivity. Accurate mass measurements were run on a TOF spectrometer internally calibrated with polyalanine.

***tert*-Butyl (*E,E*)-hexa-2,4-dienoate [*tert*-butyl sorbate] 3.** Condensed isobutylene (180 mL) at -78°C was added to a stirred solution of sorbic acid (30.0 g, 268 mmol) and conc aq H_2SO_4 (3.00 mL) in CH_2Cl_2 (600 mL) at 0°C , and the resultant mixture was allowed to warm to rt and stirred at rt for 48 h. The reaction mixture was washed with satd aq NaHCO_3 (5×300 mL) and the combined aqueous washings were extracted with CH_2Cl_2 (2×300 mL). The combined organic extracts were washed with brine (500 mL), then dried and concentrated in vacuo to give **3** as a pale yellow oil (41.1 g, 91%, $>99:1$ dr); ^{13}C δ_{H} (400 MHz, CDCl_3) 1.48 (9H, s, CMe_3), 1.84 (3H, d, J 5.9, C(6) H_3), 5.70 (1H, d, J 15.4, C(2) H), 6.04–6.21 (2H, m, C(4) H , C(5) H), 7.15 (1H, dd, J 15.4, 10.0, C(3) H).

***tert*-Butyl (3*S*, α *R*,*E*)-3-[*N*-(but-3'-en-1'-yl)-*N*-(α -methylbenzyl)amino]hex-4-enoate 13.** BuLi (2.3 M in hexanes, 2.00 mL, 4.61 mmol) was added dropwise to a stirred solution of (*R*)-*N*-(but-3-en-1-yl)-*N*-(α -methylbenzyl)amine (833 mg, 4.76 mmol, $>99:1$ er) in THF (5 mL) at -78°C and the resultant mixture was stirred at -78°C for 30 min. A solution of **3** (500 mg, 2.97 mmol, $>99:1$ dr) in THF (2 mL) at -78°C was added via cannula and the resultant mixture was stirred at -78°C for 2 h. Satd aq NH_4Cl (2 mL) was added and the reaction mixture was allowed to warm to rt, then concentrated in vacuo. The residue was partitioned between CH_2Cl_2 (10 mL) and 10% citric acid (10 mL), and the organic layer was washed with satd aq NaHCO_3 (10 mL) and brine (10 mL), then dried and concentrated in vacuo to give **13** in $>95:5$ dr. Purification via flash column chromatography (eluent 30 – 40°C petrol/ Et_2O , 20:1) gave **13** as a pale yellow oil (708 mg, 69%, $>99:1$ dr); ^{13}C $[\alpha]_{\text{D}}^{25}$ -15.4 (c 2.0 in CHCl_3); {lit.¹⁴ $[\alpha]_{\text{D}}^{20}$ -14.6 (c 1.0 in CHCl_3)}; δ_{H} (400 MHz, CDCl_3) 1.36 (3H, d, J 6.7, C(α) Me), 1.40 (9H, s, CMe_3), 1.69 (3H, d, J 5.0, C(6) H_3), 1.96–2.08 (2H, m, C(2') H_2), 2.28 (1H, dd, J 14.1, 8.2, C(2) H_{A}), 2.40 (1H, dd, J 14.1, 6.6, C(2) H_{B}), 2.44–2.58 (2H, m, C(1') H_2), 3.75–3.81 (1H, m, C(3) H), 3.93 (1H, q, J 6.7, C(α) H), 4.86–4.94 (2H, m, C(4') H_2), 5.45–5.58 (2H, m, C(4) H , C(5) H), 5.64 (1H, ddt, J 17.0, 10.3, 6.8, C(3') H), 7.17–7.23 (1H, m, *Ph*), 7.25–7.30 (2H, m, *Ph*), 7.32–7.37 (2H, m, *Ph*).

***tert*-Butyl (2'*S*, α *R*)-2-[*N*(1')-(α -methylbenzyl)-1',2',5',6'-tetrahydropyridin-2'-yl]acetate 14.**
Method A (from 13): Grubbs I catalyst (301 mg, 0.366 mmol) was added to a stirred, degassed solution of **13** (3.14 g, 9.14 mmol, $>99:1$ dr) in anhydrous CH_2Cl_2 (EtOH stabilised, 300 mL) at rt. The resultant mixture

was stirred at rt for 16 h, then concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:10:1) gave **14** as a pale yellow oil (2.30 g, 83% from **13**, >99:1 dr);¹³ $[\alpha]_{\text{D}}^{25} +41.5$ (*c* 1.0 in CHCl₃); {lit.¹³ $[\alpha]_{\text{D}}^{25} +41.8$ (*c* 1.5 in CHCl₃)}; δ_{H} (400 MHz, CDCl₃) 1.38 (3H, d, *J* 6.6, C(α)Me), 1.48 (9H, s, CMe₃), 1.66–1.75 (1H, m, C(5')H_A), 2.04–2.16 (1H, m, C(5')H_B), 2.37 (1H, dd, *J* 14.2, 6.8, C(2)H_A), 2.43–2.50 (1H, m, C(6')H_A), 2.59 (1H, dd, *J* 14.2, 7.3, C(2)H_B), 2.84 (1H, ddd, *J* 13.9, 9.5, 4.7, C(6')H_B), 3.70–3.77 (1H, m, C(2')H), 3.89 (1H, q, *J* 6.6, C(α)H), 5.62–5.67 (1H, m, C(3')H), 5.78–5.85 (1H, m, C(4')H), 7.20–7.34 (5H, m, Ph).

Method B (from 3) – Step 1: BuLi (2.3 M in hexanes, 20.0 mL, 46.1 mmol) was added dropwise to a stirred solution of (*R*)-*N*-(but-3-en-1-yl)-*N*-(α -methylbenzyl)amine (8.33 g, 47.6 mmol, >99:1 er) in THF (70 mL) at –78 °C and the resultant mixture was stirred at –78 °C for 30 min. A solution of **3** (5.00 g, 29.7 mmol, >99:1 dr) in THF (10 mL) at –78 °C was then added and the resultant mixture was stirred at –78 °C for 2 h. Satd aq NH₄Cl (20 mL) was added and the reaction mixture was allowed to warm to rt, then concentrated in vacuo. The residue was partitioned between CH₂Cl₂ (50 mL) and 10% citric acid (50 mL), and the organic layer was washed with satd aq NaHCO₃ (50 mL) and brine (50 mL), then dried and concentrated in vacuo to give **13** in >95:5 dr.

Method B (from 3) – Step 2: Grubbs I catalyst (854 mg, 1.04 mmol) was added to a stirred, degassed solution of the residue of **13** from the previous step in anhydrous CH₂Cl₂ (EtOH stabilised, 900 mL) at rt. The resultant solution was stirred at rt for 16 h, then concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:10:1) gave **14** as a pale yellow oil (6.55 g, 73% from **3**, >99:1 dr); $[\alpha]_{\text{D}}^{25} +44.8$ (*c* 0.8 in CHCl₃).

(2'*S*, α *R*)-2-[*N*(1')-(α -Methylbenzyl)-1',2',5',6'-tetrahydropyridin-2'-yl]ethan-1-ol 15. LiAlH₄ (2.4 M in THF, 3.18 mL, 7.63 mmol) was added to a stirred solution of **14** (2.30 g, 7.63 mmol, >99:1 dr) in THF (70 mL) at 0 °C. The resultant mixture was allowed to warm to rt and stirred at rt for 16 h. 2.0 M aq NaOH (7 mL) was then added and the resultant mixture was stirred at rt for 3 h. The reaction mixture was filtered through Celite® (eluent CH₂Cl₂), then concentrated in vacuo to give **15** as a pale yellow oil (1.66 g, 94%, >99:1 dr); $[\alpha]_{\text{D}}^{25} -3.9$ (*c* 1.0 in CHCl₃); ν_{max} (ATR) 3393 (O–H), 3026, 2970, 2839, (C–H), 1454 (C=C); δ_{H} (400 MHz, CDCl₃) 1.49 (3H, d, *J* 6.6, C(α)Me), 1.73–1.88 (2H, m, C(2)H_A, C(5')H_A), 1.88–1.97 (1H, m, C(2)H_B), 2.04–2.14 (1H, m, C(5')H_B), 2.34 (1H, app dt, *J* 13.4, 4.5, C(6')H_A), 3.02 (1H, ddd, *J* 13.4, 8.9, 4.7, C(6')H_B), 3.59–3.66 (1H, m, C(2')H), 3.85–3.96 (2H, m, C(1)H₂), 4.05 (1H, q, *J* 6.6, C(α)H), 5.50–5.56 (1H, m, C(3')H), 5.84–5.97 (1H, m, C(4')H), 6.36 (1H, br s, OH), 7.24–7.39 (5H, m, Ph); δ_{C} (100 MHz, CDCl₃) 21.3, 21.4 (C(5'), C(α)Me), 32.9 (C(2)), 40.0 (C(6')), 56.2 (C(2')), 57.6 (C(α)), 63.0 (C(1)), 126.4 (C(4')),

127.4 (*p-Ph*), 127.8, 128.5 (*o,m-Ph*), 128.5 (*C(3')*), 142.6 (*i-Ph*); m/z (ESI^+) 232 ($[\text{M} + \text{H}]^+$, 100%); HRMS (ESI^+) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{15}\text{H}_{22}\text{NO}^+$ 232.1696; Found 232.1696.

***tert*-Butyl (2'*S*,*aR*,*E*)-4-[*N*(1')-(*a*-methylbenzyl)-1',2',5',6'-tetrahydropyridin-2'-yl]but-2-enoate**
16 and *tert*-Butyl (2'*S*,*aR*,*Z*)-4-[*N*(1')-(*a*-methylbenzyl)-1',2',5',6'-tetrahydropyridin-2'-yl]but-2-enoate

17. Method A (from 15): DMSO (0.61 mL, 8.65 mmol) was added to a stirred solution of $(\text{COCl})_2$ (0.37 mL, 4.32 mmol) in CH_2Cl_2 (50 mL) at -78°C . After 20 min, **15** (500 mg, 2.16 mmol, >99:1 dr) was added and the resultant mixture was stirred at -78°C for 40 min, then Et_3N (1.81 mL, 13.0 mmol) was added. The reaction mixture was allowed to warm to rt, $\text{Ph}_3\text{P}=\text{CHCO}_2^t\text{Bu}$ (814 mg, 2.16 mmol) was added, and the resultant mixture was stirred at rt for 16 h. Satd aq K_2CO_3 (300 mL) was then added and the reaction mixture was extracted with CH_2Cl_2 (3×300 mL). The combined organic extracts were washed with brine (500 mL), then dried and concentrated in vacuo to give a 93:7 mixture of **16** and **17**, respectively. Purification via flash column chromatography (eluent $30\text{--}40^\circ\text{C}$ petrol/ $\text{Et}_2\text{O}/\text{NH}_4\text{OH}$, 100:10:1) gave **17** as a pale yellow oil (19 mg, 3%, >99:1 dr); $[\alpha]_{\text{D}}^{25} +57.4$ (c 1.0 in CHCl_3); ν_{max} (ATR) 2975, 2931 (C-H), 1712 (C=O), 1639 (C=C); δ_{H} (400 MHz, CDCl_3) 1.34 (3H, d, J 6.8, $\text{C}(\alpha)\text{Me}$), 1.47 (9H, s, CMe_3), 1.84–1.94 (1H, m, $\text{C}(5')\text{H}_{\text{A}}$), 1.96–2.05 (1H, m, $\text{C}(5')\text{H}_{\text{B}}$), 2.23 (1H, ddd, J 12.1, 7.2, 5.7, $\text{C}(6')\text{H}_{\text{A}}$), 2.65–2.73 (1H, m, $\text{C}(4)\text{H}_{\text{A}}$), 2.92 (1H, app dt, J 12.1, 5.6, $\text{C}(6')\text{H}_{\text{B}}$), 3.16–3.26 (2H, m, $\text{C}(4)\text{H}_{\text{B}}$, $\text{C}(2')\text{H}$), 4.00 (1H, q, J 6.8, $\text{C}(\alpha)\text{H}$), 5.46–5.51 (1H, m, $\text{C}(3')\text{H}$), 5.73 (1H, d, J 11.4, $\text{C}(2)\text{H}$), 5.73–5.78 (1H, m, $\text{C}(4')\text{H}$), 6.30 (1H, app dt, J 11.4, 7.5, $\text{C}(3)\text{H}$), 7.17–7.34 (5H, m, *Ph*); δ_{C} (100 MHz, CDCl_3) 20.9 ($\text{C}(\alpha)\text{Me}$), 24.3 ($\text{C}(5')$), 28.4 (CMe_3), 32.1 ($\text{C}(4)$), 40.8 ($\text{C}(6')$), 54.9 ($\text{C}(2')$), 57.1 ($\text{C}(\alpha)$), 80.1 (CMe_3), 122.0 ($\text{C}(2)$), 126.5 ($\text{C}(4')$), 126.9 (*p-Ph*), 128.1, 128.1 (*o,m-Ph*), 130.1 ($\text{C}(3')$), 142.6 (*i-Ph*), 146.7 ($\text{C}(3)$), 166.4 ($\text{C}(1)$); m/z (ESI^+) 328 ($[\text{M} + \text{H}]^+$, 100%); HRMS (ESI^+) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{30}\text{NO}_2^+$ 328.2271; Found 328.2273. Further elution gave **16** as a yellow oil (322 mg, 46%, >99:1 dr); $[\alpha]_{\text{D}}^{25} +65.9$ (c 1.0 in CHCl_3); ν_{max} (ATR) 2979, 2932 (C-H), 1713 (C=O), 1651 (C=C); δ_{H} (400 MHz, CDCl_3) 1.38 (3H, d, J 6.7, $\text{C}(\alpha)\text{Me}$), 1.49 (9H, s, CMe_3), 1.82–1.93 (1H, m, $\text{C}(5')\text{H}_{\text{A}}$), 2.03–2.15 (1H, m, $\text{C}(5')\text{H}_{\text{B}}$), 2.36–2.55 (3H, m, $\text{C}(4)\text{H}_2$, $\text{C}(6')\text{H}_{\text{A}}$), 2.88 (1H, ddd, J 12.7, 7.8, 4.8, $\text{C}(6')\text{H}_{\text{B}}$), 3.23–3.31 (1H, m, $\text{C}(2')\text{H}$), 3.95 (1H, q, J 6.7, $\text{C}(\alpha)\text{H}$), 5.53–5.58 (1H, m, $\text{C}(3')\text{H}$), 5.77 (1H, d, J 15.5, $\text{C}(2)\text{H}$), 5.77–5.83 (1H, m, $\text{C}(4')\text{H}$), 6.90 (1H, ddd, J 15.5, 8.0, 6.7, $\text{C}(3)\text{H}$), 7.21–7.35 (5H, m, *Ph*); δ_{C} (100 MHz, CDCl_3) 21.4 ($\text{C}(\alpha)\text{Me}$), 23.9 ($\text{C}(5')$), 28.3 (CMe_3), 35.9 ($\text{C}(4)$), 40.7 ($\text{C}(6')$), 54.6 ($\text{C}(2')$), 57.9 ($\text{C}(\alpha)$), 80.1 (CMe_3), 124.3 ($\text{C}(2)$), 126.5 ($\text{C}(4')$), 127.0 (*p-Ph*), 127.9, 128.3 (*o,m-Ph*), 129.5 ($\text{C}(3')$), 143.4 (*i-Ph*), 145.9 ($\text{C}(3)$), 166.1 ($\text{C}(1)$); m/z (ESI^+) 328 ($[\text{M} + \text{H}]^+$, 100%); HRMS (ESI^+) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{30}\text{NO}_2^+$ 328.2271; Found 328.2271.

Method B (from 14): DIBAL-H (1.0 M in PhMe, 7.30 mL, 7.30 mmol) was added dropwise to a stirred solution of **14** (2.00 g, 6.64 mmol, >99:1 dr) in PhMe (20 mL) at -78°C and the resultant mixture was stirred at -78°C for 1 h. MeOH (1.34 mL, 33.2 mmol) and $\text{Ph}_3\text{P}=\text{CHCO}_2^t\text{Bu}$ (2.50 g, 6.64 mmol) were added sequentially, and the resultant mixture was allowed to warm to rt and stirred at rt for 16 h. The reaction mixture was then concentrated in vacuo to give an 80:20 mixture of **16** and **17**, respectively. Purification via flash column chromatography (eluent $30-40^{\circ}\text{C}$ petrol/ $\text{Et}_2\text{O}/\text{NH}_4\text{OH}$, 100:10:1) gave **17** as a pale yellow oil (313 mg, 14%, >99:1 dr); $[\alpha]_{\text{D}}^{25} +60.2$ (c 0.5 in CHCl_3). Further elution gave **16** as a yellow oil (1.57 g, 72%, >99:1 dr); $[\alpha]_{\text{D}}^{25} +63.4$ (c 1.0 in CHCl_3).

***tert*-Butyl (3*R*,2'*S*, α *R*)-3-[*N*-benzyl-*N*-isopropylamino]-4-[*N*(1')-(α -methylbenzyl)-1',2',5',6'-tetrahydropyridin-2'-yl]butanoate and *tert*-butyl (3*S*,2'*S*, α *R*)-3-[*N*-benzyl-*N*-isopropylamino]-4-[*N*(1')-(α -methylbenzyl)-1',2',5',6'-tetrahydropyridin-2'-yl]butanoate **19** and **20**.** BuLi (2.3 M in hexanes, 0.44 mL, 1.00 mmol) was added dropwise to a stirred solution of *N*-benzyl-*N*-isopropylamine (155 mg, 1.04 mmol) in THF (2 mL) at -78°C and the resultant mixture was stirred at -78°C for 30 min. A solution of **16** (212 mg, 0.647 mmol, >99:1 dr) in THF (1 mL) at -78°C was added via cannula and the resultant mixture was stirred at -78°C for 2 h. Satd aq NH_4Cl (0.5 mL) was added and the reaction mixture was allowed to warm to rt, then concentrated in vacuo. The residue was partitioned between CH_2Cl_2 (10 mL) and 10% citric acid (10 mL), and the organic layer was washed with satd aq NaHCO_3 (10 mL) and brine (10 mL), then dried and concentrated in vacuo to give a 50:50 mixture of **19** and **20** as a colourless oil (296 mg, 96%). Data for **19**: δ_{H} (400 MHz, CDCl_3) 1.05–1.11 (6H, m, NCHMe_2), 1.35 (3H, d, J 6.5, $\text{C}(\alpha)\text{Me}$), 1.36–1.63 (2H, m, $\text{C}(4)\text{H}_{\text{A}}$, $\text{C}(5')\text{H}_{\text{A}}$), 1.50 (9H, s, CMe_3), 1.84–1.98 (1H, m, $\text{C}(4)\text{H}_{\text{B}}$), 2.04–2.17 (1H, m, $\text{C}(5')\text{H}_{\text{B}}$), 2.42–2.57 (3H, m, $\text{C}(2)\text{H}_2$, $\text{C}(6')\text{H}_{\text{A}}$), 2.72–2.84 (1H, m, $\text{C}(6')\text{H}_{\text{B}}$), 2.91–3.02 (1H, m, NCHMe_2), 3.32–3.43 (1H, m, $\text{C}(2')\text{H}$), 3.52–3.61 (1H, m, $\text{C}(3)\text{H}$), 3.65 (1H, d, J 14.5, $\text{NCH}_{\text{A}}\text{H}_{\text{B}}\text{Ph}$), 3.73 (1H, d, J 14.5, $\text{NCH}_{\text{A}}\text{H}_{\text{B}}\text{Ph}$), 3.84 (1H, q, J 6.5, $\text{C}(\alpha)\text{H}$), 5.33–5.43 (1H, m, $\text{C}(3')\text{H}$), 5.69–5.79 (1H, m, $\text{C}(4')\text{H}$), 7.19–7.43 (10H, m, *Ph*). Data for **20**: δ_{H} (400 MHz, CDCl_3) 1.05–1.11 (6H, m, NCHMe_2), 1.35 (3H, d, J 6.5, $\text{C}(\alpha)\text{Me}$), 1.36–1.63 (2H, m, $\text{C}(4)\text{H}_{\text{A}}$, $\text{C}(5')\text{H}_{\text{A}}$), 1.46 (9H, s, CMe_3), 1.68 (1H, app dt, J 14.2, 5.2, $\text{C}(4)\text{H}_{\text{B}}$), 2.04–2.17 (1H, m, $\text{C}(5')\text{H}_{\text{B}}$), 2.29 (1H, dd, J 13.7, 6.2, $\text{C}(2)\text{H}_{\text{A}}$), 2.40–2.57 (2H, m, $\text{C}(2)\text{H}_{\text{B}}$, $\text{C}(6')\text{H}_{\text{A}}$), 2.72–2.84 (1H, m, $\text{C}(6')\text{H}_{\text{B}}$), 2.91–3.02 (1H, m, NCHMe_2), 3.32–3.43 (1H, m, $\text{C}(2')\text{H}$), 3.52–3.61 (1H, m, $\text{C}(3)\text{H}$), 3.66 (2H, s, NCH_2Ph), 3.84 (1H, q, J 6.5, $\text{C}(\alpha)\text{H}$), 5.49–5.55 (1H, m, $\text{C}(3')\text{H}$), 5.69–5.79 (1H, m, $\text{C}(4')\text{H}$), 7.19–7.43 (10H, m, *Ph*). Data for mixture of **19** and **20**: ν_{max} (ATR) 3024, 2967, 2931, 2837 (C–H), 1724 (C=O), 1602 (C=C); δ_{C} (100 MHz, CDCl_3) 20.2, 20.4, 20.8, 20.9, 21.1, 21.8, 22.7, 23.0, 28.3, 28.3, 36.5, 37.4, 39.8, 39.9, 40.2, 40.4, 47.8, 49.3, 49.4, 50.0, 51.2, 51.9, 52.4, 53.9, 58.0, 58.2, 80.0, 80.1, 125.4, 125.7, 126.6, 126.6, 126.8, 126.8,

127.5, 127.6, 128.1, 128.1, 128.3, 128.3, 128.5, 129.0, 129.8, 130.2, 141.8, 142.3, 145.9, 146.2, 172.3, 172.6; m/z (ESI⁺) 477 ([M + H]⁺, 100%); HRMS (ESI⁺) m/z : [M + H]⁺ Calcd for C₃₁H₄₅N₂O₂⁺ 477.3476; Found 477.3471.

tert-Butyl (3*R*,2'*S*, α *R*, α' *R*)-3-[*N*-benzyl-*N*-(α -methylbenzyl)amino]-4-[*N*(1')-(α' -methylbenzyl)-1',2',5',6'-tetrahydropyridin-2'-yl]butanoate 21. BuLi (2.3 M in hexanes, 0.58 mL, 1.33 mmol) was added dropwise to a stirred solution of (*R*)-*N*-benzyl-*N*-(α -methylbenzyl)amine (289 mg, 1.37 mmol, >99:1 er) in THF (2 mL) at –78 °C and the resultant mixture was stirred at –78 °C for 30 min. A solution of **16** (280 mg, 0.855 mmol, >99:1 dr) in THF (1 mL) at –78 °C was added and the resultant mixture was stirred at –78 °C for 2 h. Satd aq NH₄Cl (0.5 mL) was added and the reaction mixture was allowed to warm to rt, then concentrated in vacuo. The residue was partitioned between CH₂Cl₂ (5 mL) and 10% citric acid (5 mL), and the organic layer was washed with satd aq NaHCO₃ (5 mL) and brine (5 mL), then dried and concentrated in vacuo to give **21** in >95:5 dr. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:10:1) gave **21** as a pale yellow oil (449 mg, 97%, >99:1 dr); [α]_D²⁵ +23.8 (*c* 1.0 in CHCl₃); $\nu_{\max}$ (ATR) 3025, 2973, 2931, 2837 (C–H), 1722 (C=O), 1618 (C=C); δ_{H} (400 MHz, CDCl₃) 1.28 (3H, d, *J* 6.8, C(α')Me), 1.29 (3H, d, *J* 6.9, C(α)Me), 1.32 (9H, s, CMe₃), 1.39–1.50 (2H, m, C(4)*H*_A, C(5')*H*_A), 1.90–2.00 (1H, m, C(4)*H*_B), 2.06 (1H, dd, *J* 14.1, 7.7, C(2)*H*_A), 2.03–2.19 (1H, m, C(5')*H*_B), 2.18 (1H, dd, *J* 14.1, 5.4, C(2)*H*_B), 2.48 (1H, app dd, *J* 13.8, 5.3, C(6')*H*_A), 2.81 (1H, ddd, *J* 13.8, 11.0, 4.7, C(6')*H*_B), 3.47–3.59 (2H, m, C(3)*H*, C(2')*H*), 3.58 (1H, d, *J* 14.7, NCH_AH_BPh), 3.83 (1H, d, *J* 14.7, NCH_AH_BPh), 3.84 (1H, q, *J* 6.8, C(α')*H*), 3.92 (1H, q, *J* 6.9, C(α)*H*), 5.40–5.47 (1H, m, C(3')*H*), 5.71–5.77 (1H, m, C(4')*H*), 7.17–7.43 (15H, m, *Ph*); δ_{C} (100 MHz, CDCl₃) 20.5 (C(α)Me), 20.9 (C(5')), 22.9 (C(α')Me), 28.2 (CMe₃), 38.3, 38.5 (C(2), C(4)), 40.3 (C(6')), 50.4 (NCH₂Ph), 50.9 (C(2')), 52.7 (C(3')), 58.2 (C(α')), 58.6 (C(α)), 80.1 (CMe₃), 125.7 (C(4')), 126.6, 126.8, 127.0 (*p*-*Ph*), 127.6, 128.2, 128.2, 128.3, 128.3, 128.6 (*o,m*-*Ph*), 129.6 (C(3')), 142.1, 144.2, 146.2 (*i*-*Ph*), 172.1 (C(1)); m/z (ESI⁺) 539 ([M + H]⁺, 100%); HRMS (ESI⁺) m/z : [M + H]⁺ Calcd for C₃₆H₄₇N₂O₂⁺ 539.3632; Found 539.3648.

tert-Butyl (3*S*,2'*S*, α *S*, α' *R*)-3-[*N*-benzyl-*N*-(α -methylbenzyl)amino]-4-[*N*(1')-(α' -methylbenzyl)-1',2',5',6'-tetrahydropyridin-2'-yl]butanoate 22. BuLi (2.3 M in hexanes, 0.64 mL, 1.47 mmol) was added dropwise to a stirred solution of (*S*)-*N*-benzyl-*N*-(α -methylbenzyl)amine (318 mg, 1.51 mmol, >99:1 er) in THF (2 mL) at –78 °C and the resultant mixture was stirred at –78 °C for 30 min. A solution of **16** (308 mg, 0.941 mmol, >99:1 dr) in THF (1 mL) at –78 °C was added and the resultant mixture was stirred at –78 °C for 2 h. Satd aq NH₄Cl (0.5 mL) was added and the reaction mixture was allowed to warm to rt, then concentrated in vacuo. The residue was partitioned between CH₂Cl₂ (5 mL) and 10% citric acid (5 mL), and

the organic layer was washed with satd aq NaHCO₃ (5 mL) and brine (5 mL), then dried and concentrated in vacuo to give **22** in >95:5 dr. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:10:1) gave **22** as a pale yellow oil (476 mg, 94%, >99:1 dr); [α]_D²⁵ +68.1 (*c* 1.0 in CHCl₃); $\nu_{\max}$ (ATR) 3064, 3025, 2973, 2932 (C–H), 1722 (C=O), 1609 (C=C); δ_{H} (400 MHz, CDCl₃) 1.28 (3H, d, *J* 6.6, C(α')Me), 1.40 (3H, d, *J* 6.9, C(α)Me), 1.44 (9H, s, CMe₃), 1.55–1.73 (3H, m, C(4)H₂, C(5')H_A), 2.03–2.14 (1H, m, C(5')H_B), 2.17 (1H, dd, *J* 14.4, 6.6, C(2)H_A), 2.30 (1H, dd, *J* 14.4, 6.6, C(2)H_B), 2.41 (1H, app dd, *J* 13.7, 3.5, C(6')H_A), 2.76 (1H, ddd, *J* 13.7, 10.4, 4.8, C(6')H_B), 3.28–3.35 (1H, m, C(2')H), 3.55–3.63 (1H, m, C(3)H), 3.63 (1H, d, *J* 15.0, NCH_AH_BPh), 3.76 (1H, d, *J* 15.0, NCH_AH_BPh), 3.82 (1H, q, *J* 6.6, C(α')H), 4.01 (1H, q, *J* 6.9, C(α)H), 5.55–5.61 (1H, m, C(3')H), 5.71–5.77 (1H, m, C(4')H), 7.17–7.39 (15H, m, *Ph*); δ_{C} (100 MHz, CDCl₃) 19.6 (C(α)Me), 21.5 (C(5')), 22.5 (C(α')Me), 28.3 (CMe₃), 36.6 (C(4)), 39.8, 39.9 (C(2), C(6')), 49.8 (NCH₂Ph), 52.9 (C(2')), 54.3 (C(3)), 58.1 (C(α')), 58.9 (C(α)), 80.0 (CMe₃), 125.3 (C(4')), 126.6, 126.8, 126.9 (*p-Ph*), 127.7, 128.1, 128.2, 128.3, 128.3, 128.5 (*o,m-Ph*), 130.4 (C(3')), 141.9, 143.9, 145.7 (*i-Ph*), 172.3 (C(1)); *m/z* (ESI⁺) 539 ([M + H]⁺, 100%); HRMS (ESI⁺) *m/z*: [M + H]⁺ Calcd for C₃₆H₄₇N₂O₂⁺ 539.3632; Found 539.3631.

(2*S*,2'*S*, α *R*, α' *R*)-1-[N(1')-(α -Methylbenzyl)-1',2',5',6'-tetrahydropyridin-2'-yl]-2-[N-benzyl-N-(α' -methylbenzyl)amino]pent-4-ene 23. DIBAL-H (1.0 M in PhMe, 0.20 mL, 0.20 mmol) was added dropwise to a stirred solution of **21** (100 mg, 0.19 mmol, >99:1 dr) in PhMe (2 mL) at –78 °C and the resultant mixture was stirred at –78 °C for 1 h. MeOH (38 μ L, 0.93 mmol) and Ph₃P=CH₂ (0.31 M in PhMe/THF, 6.58 mL, 2.04 mmol)²⁴ were added sequentially, and the resultant mixture was allowed to warm to rt and stirred at rt for 16 h, then concentrated in vacuo. The residue was dissolved in Et₂O (10 mL) and the resultant solution was filtered through a short plug of Celite[®] (eluent Et₂O) and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:5:1) gave **23** as a colourless oil (54 mg, 62%, >99:1 dr); [α]_D²⁵ +28.2 (*c* 1.0 in CHCl₃); $\nu_{\max}$ (ATR) 3358, 2961 (C–H), 1596 (C=C); δ_{H} (400 MHz, CDCl₃) 1.26 (3H, d, *J* 6.4, C(α)Me), 1.37 (3H, d, *J* 6.9, C(α')Me), 1.46–1.57 (2H, m, C(1)H_A, C(5')H_A), 1.89–2.01 (2H, m, C(3)H_A, C(1)H_B), 2.05–2.16 (2H, m, C(3)H_B, C(5')H_B), 2.49 (1H, app dd, *J* 14.0, 5.2, C(6')H_A), 2.86 (1H, ddd, *J* 14.0, 11.2, 4.6, C(6')H_B), 3.12 (1H, app quintet, *J* 6.4, C(2)H), 3.38–3.46 (1H, m, C(2')H), 3.73 (1H, d, *J* 14.7, NCH_AH_BPh), 3.82 (1H, q, *J* 6.4, C(α)H), 3.84 (1H, d, *J* 14.7, NCH_AH_BPh), 3.96 (1H, q, *J* 6.9, C(α')H), 4.85–4.93 (2H, m, C(5)H₂), 5.46–5.52 (1H, m, C(3')H), 5.59 (1H, ddt, *J* 17.0, 10.2, 7.0, C(4)H), 5.72–5.79 (1H, m, C(4')H), 7.18–7.34 (11H, m, *Ph*), 7.36–7.41 (2H, m, *Ph*), 7.42–7.46 (2H, m, *Ph*); δ_{C} (100 MHz, CDCl₃) 17.9 (C(α')Me), 20.6 (C(5')), 22.9 (C(α)Me), 36.4 (C(3)), 37.9 (C(1)), 40.2 (C(6')), 50.3 (NCH₂Ph), 51.3 (C(2')), 53.8 (C(2)), 57.3 (C(α')Me), 58.1 (C(α)Me), 115.3 (C(5)),

125.5 (C(4')), 126.5, 126.6, 126.6 (*p*-Ph), 127.4, 127.9, 128.1, 128.2, 128.2, 128.5 (*o,m*-Ph), 130.0 (C(3')), 137.9 (C(4)), 142.2, 144.7, 146.1 (*i*-Ph); m/z (ESI⁺) 465 ([M + H]⁺, 100%); HRMS (ESI⁺) m/z : [M + H]⁺ Calcd for C₃₃H₄₁N₂⁺ 465.3264; Found 465.3258.

(2*S*,2'*S*, α *R*, α' *R*,*Z*)-1-[N(1')-(α -Methylbenzyl)-1',2',5',6'-tetrahydropyridin-2'-yl]-2-[N-benzyl-N-(α' -methylbenzyl)amino]hept-4-ene **24**. DIBAL-H (1.0 M in PhMe, 0.53 mL, 0.53 mmol) was added dropwise to a stirred solution of **21** (260 mg, 0.483 mmol, >99:1 dr) in PhMe (4 mL) at -78 °C and the resultant mixture was stirred at -78 °C for 1 h. MeOH (98 μ L, 2.41 mmol) and Ph₃P=CHCH₂CH₃ (0.31 M in PhMe/THF, 15.6 mL, 4.83 mmol)²⁵ were added sequentially, and the resultant mixture was allowed to warm to rt and stirred at rt for 16 h, then concentrated in vacuo. The residue was dissolved in Et₂O (20 mL) and the resultant solution was filtered through a short plug of Celite[®] (eluent Et₂O) and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:5:1) gave **24** as a pale yellow oil (148 mg, 62%, >95:5 dr [(*Z*):(*E*)]); $[\alpha]_D^{25}$ +36.5 (*c* 1.0 in CHCl₃); $\nu_{\max}$ (ATR) 3024, 2931, 2838 (C–H), 1580, 1493, 1453 (C=C); δ_H (400 MHz, CDCl₃) 0.91 (3H, t, *J* 7.5, C(7)*H*₃), 1.26 (3H, d, *J* 6.4, C(α')*Me*), 1.36 (3H, d, *J* 6.9, C(α)*Me*), 1.46–1.58 (2H, m, C(1)*H*_A, C(5')*H*_A), 1.89–2.16 (6H, m, C(1)*H*_B, C(3)*H*₂, C(6)*H*₂, C(5')*H*_B), 2.49 (1H, dd, *J* 14.0, 5.2, C(6')*H*_A), 2.87 (1H, ddd, *J* 14.0, 11.2, 4.7, C(6')*H*_B), 3.06 (1H, app quintet, *J* 6.4, C(2)*H*), 3.40–3.47 (1H, m, C(2')*H*), 3.73 (1H, d, *J* 14.8, NCH_AH_BPh), 3.81 (1H, q, *J* 6.4, C(α')*H*), 3.86 (1H, d, *J* 14.8, NCH_AH_BPh), 3.98 (1H, q, *J* 6.9, C(α)*H*), 5.10–5.19 (1H, m, C(4)*H*), 5.25–5.33 (1H, m, C(5)*H*), 5.44–5.50 (1H, m, C(3')*H*), 5.71–5.78 (1H, m, C(4')*H*), 7.19–7.35 (11H, m, *Ph*), 7.37–7.41 (2H, m, *Ph*), 7.42–7.46 (2H, m, *Ph*); δ_C (100 MHz, CDCl₃) 14.3 (C(7)), 18.5 (C(α)*Me*), 20.8 (C(5')), 20.9 (C(6)), 23.0 (C(α')*Me*), 29.6 (C(3)), 38.3 (C(1)), 40.3 (C(6')), 50.5 (NCH₂Ph), 51.6 (C(2')), 54.5 (C(2)), 57.7 (C(α)), 58.2 (C(α')), 125.5 (C(4')), 126.5, 126.7, 126.7 (*p*-Ph), 127.6 (*o,m*-Ph), 128.0 (C(4)), 128.0, 128.2, 128.3, 128.3, 128.6 (*o,m*-Ph), 130.2 (C(3')), 131.9 (C(5)), 142.6, 145.2, 146.3 (*i*-Ph); m/z (ESI⁺) 493 ([M + H]⁺, 100%); HRMS (ESI⁺) m/z : [M + H]⁺ Calcd for C₃₅H₄₅N₂⁺ 493.3577; Found 493.3568.

(2*S*,2'*R*)-1-(Piperidin-2'-yl)-2-aminopentane **25**. Palladium black (15 mg, 20% w/w) was added to a stirred, degassed solution of **23** (75 mg, 0.16 mmol) in MeOH/H₂O/AcOH (10:1:1, 2 mL). The resultant mixture was stirred under an atmosphere of H₂ (5 atm) at rt for 48 h. The reaction mixture was then filtered through a short plug of Celite[®] (eluent MeOH/H₂O/AcOH, 10:1:1) and concentrated in vacuo. Purification via flash column chromatography (eluent CHCl₃/MeOH/NH₄OH 10:1:1) gave **25** as a colourless oil (20 mg, 73%, >99:1 dr); $[\alpha]_D^{25}$ -3.3 (*c* 1.0 in CHCl₃); $\nu_{\max}$ (ATR) 3292 (N–H), 2930, 2860 (C–H); δ_H (400 MHz, CDCl₃) 0.89 (3H, t, *J* 6.9, C(5)*H*₃), 1.02–1.13 (1H, m, C(3')*H*_A), 1.17–1.46 (8H, m, C(1)*H*₂, C(3)*H*₂, C(4)*H*₂, C(4')*H*_A, C(5')*H*_A), 1.53–1.68 (2H, m, C(3')*H*_B, C(4')*H*_B), 1.72–1.80 (1H, m, C(5')*H*_B), 2.55–2.66 (2H, m,

C(2')H, C(6')H_A), 2.75–2.82 (1H, m, C(2)H), 3.01–3.07 (1H, m, C(6')H_B); δ_C (100 MHz, CDCl₃) 14.3 (C(5)), 19.2 (C(4)), 25.0 (C(5')), 26.7 (C(4')), 33.4 (C(3')), 42.1 (C(3)), 45.5 (C(1)), 47.1 (C(6')), 49.9 (C(2)), 56.1 (C(2')); m/z (ESI⁺) 171 ([M + H]⁺, 100%); HRMS (ESI⁺) m/z : [M + H]⁺ Calcd for C₁₀H₂₃N₂⁺ 171.1856; Found 171.1856.

(5S,6aR,11aS)-5-Propyldecahydro-5H-pyrido[1,2-c]pyrrolo[1,2-a]pyrimidine [(+)-tetraponerine

4] 27. *Method A (from 25):* K₂CO₃ (88 mg, 0.63 mmol) and 4-bromobutanal (64 mg, 0.42 mmol) were added to a stirred solution of **25** (36 mg, 0.21 mmol, >99:1 dr) in CH₂Cl₂ (2 mL) at rt, and the resultant mixture was stirred at rt for 2 h. H₂O (2 mL) and satd aq NaHCO₃ (10 mL) were added, and the resultant mixture was extracted with CH₂Cl₂ (3 × 5 mL), then dried and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Me₂CO/NH₄OH, 100:10:1) gave (+)-tetraponerine 4 (**27**) as a colourless oil (32 mg, 68% from **25**, >99:1 dr); $[\alpha]_D^{25} +90.8$ (c 1.0 in CHCl₃); {lit.⁴ for a sample isolated from the natural source $[\alpha]_D^{20} +94$ (c 0.2 in CHCl₃); lit.^{9b} $[\alpha]_D^{20} +96$ (c 2.0 in CHCl₃)}; $\nu_{\max}$ (ATR) 2933, 2855, 2793 (C–H);²⁶ δ_H (400 MHz, C₆D₆) 0.89 (3H, t, *J* 7.1, C(3')H₃), 1.13–1.29 (2H, m, C(9)H_A, C(2')H_A), 1.30–1.56 (9H, m, C(2)H_A, C(6)H₂, C(7)H₂, C(8)H_A, C(1')H₂, C(2')H_B), 1.59–1.77 (7H, m, C(1)H₂, C(2)H_B, C(6a)H, C(8)H_B, C(9)H_B, C(10)H_A), 2.02 (1H, dd, *J* 16.3, 8.6, C(3)H_A), 2.11 (1H, ddd, *J* 14.6, 7.4, 3.6, C(5)H), 2.31 (1H, dd, *J* 7.8, 5.6, C(11a)H), 2.80–2.85 (1H, m, C(10)H_B), 3.13 (1H, app td, *J* 8.6, 2.2, C(3)H_B); δ_C (100 MHz, C₆D₆) 14.8 (C(3')), 18.7 (C(2')), 20.2 (C(2)), 25.1 (C(9)), 26.3 (C(8)), 29.7 (C(1)), 33.0 (C(7)), 36.9 (C(1')), 38.0 (C(6)), 49.0 (C(3)), 51.6 (C(10)), 61.1 (C(5)), 62.7 (C(6a)), 85.6 (C(11a)); m/z (ESI⁺) 223 ([M + H]⁺, 100%); HRMS (ESI⁺) m/z : [M + H]⁺ Calcd for C₁₄H₂₇N₂⁺ 223.2169; Found 223.2169.

Method B (from 23): Palladium black (24 mg, 20% w/w) was added to a stirred, degassed solution of **23** (121 mg, 0.26 mmol, >99:1 dr) in MeOH/H₂O/AcOH (10:1:1, 2 mL). The resultant mixture was stirred under an atmosphere of H₂ (5 atm) at rt for 48 h. The reaction mixture was then filtered through a short plug of Celite® (eluent MeOH/H₂O/AcOH, 10:1:1), and concentrated in vacuo. The residue was dissolved in CH₂Cl₂ (2 mL), then K₂CO₃ (108 mg, 0.78 mmol) and 4-bromobutanal (79 mg, 0.52 mmol) were added to the resultant solution. The reaction mixture was stirred at rt for 2 h, then H₂O (2 mL) and satd aq NaHCO₃ (10 mL) were added. The reaction mixture was then extracted with CH₂Cl₂ (3 × 5 mL) and the combined organic extracts were dried and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Me₂CO/NH₄OH, 100:10:1) gave (+)-tetraponerine 4 (**27**) as a colourless oil (31 mg, 54% from **23**, >99:1 dr); $[\alpha]_D^{25} +88.6$ (c 1.0 in CHCl₃).

(5S,6aR,11aS)-5-Pentyldecahydro-5H-pyrido[1,2-c]pyrrolo[1,2-a]pyrimidine [(+)-tetraponerine

8] 28. Palladium black (26 mg, 20% w/w) was added to a stirred, degassed solution of **24** (130 mg, 0.26

mmol, >95:5 dr) in MeOH/H₂O/AcOH (10:1:1, 2 mL). The resultant mixture was stirred under an atmosphere of H₂ (5 atm) at rt for 48 h. The reaction mixture was then filtered through a short plug of Celite® (eluent MeOH/H₂O/AcOH, 10:1:1) and concentrated in vacuo. The residue was dissolved in MeOH/H₂O/AcOH (10:1:1, 2 mL) and the resultant solution was degassed. Pd(OH)₂/C (65 mg, 50% w/w) was added and the resultant mixture was stirred under an atmosphere of H₂ (5 atm) at rt for 24 h. The reaction mixture was then filtered through a short plug of Celite® (eluent MeOH/H₂O/AcOH, 10:1:1) and concentrated in vacuo. The residue was dissolved in CH₂Cl₂ (2 mL), then K₂CO₃ (110 mg, 0.792 mmol) and 4-bromobutanal (80 mg, 0.53 mmol) were added to the resultant solution. The reaction mixture was stirred at rt for 2 h, then H₂O (2 mL) and satd aq NaHCO₃ (10 mL) were added. The reaction mixture was then extracted with CH₂Cl₂ (3 × 5 mL) and the combined organic extracts were dried and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Me₂CO/NH₄OH, 100:10:1) gave (+)-tetraponerine 8 (**28**) as a white solid (44 mg, 67% from **24**, >99:1 dr); mp 38–40 °C; {lit.⁵ mp 40 °C}; [α]_D²⁵ +93.0 (c 1.0 in CHCl₃); {lit.⁴ for a sample isolated from the natural source [α]_D²⁰ +102 (c 0.2 in CHCl₃); lit.^{9b} [α]_D²⁰ +101 (c 2.0 in CHCl₃)}; ν_{max} (ATR) 2931, 2857, 2780 (C–H);²⁶ δ_H (400 MHz, C₆D₆) 0.90 (3H, t, *J* 7.0, C(5')H₃), 1.16–1.67 (17H, m, C(2)H_A, C(6)H₂, C(7)H₂, C(8)H₂, C(9)H₂, C(1')H₂, C(2')H₂, C(3')H₂, C(4')H₂), 1.68–1.78 (5H, m, C(1)H₂, C(2)H_B, C(6a)H, C(10)H_A), 2.04 (1H, app dd, *J* 16.3, 8.5, C(3)H_A), 2.12 (1H, app ddd, *J* 14.6, 7.3, 3.6, C(5)H), 2.31 (1H, dd, *J* 7.9, 5.7, C(11a)H), 2.81–2.86 (1H, m, C(10)H_B), 3.15 (1H, app td, *J* 8.5, 2.2, C(3)H_B); δ_C (100 MHz, C₆D₆) 14.4 (C(5')), 20.2 (C(2)), 23.2 (C(4')), 25.2 (C(9)), 25.2 (C(2')), 26.3 (C(8)), 29.7 (C(1)), 32.8 (C(3')), 33.0 (C(7)), 34.6 (C(1')), 38.0 (C(6)), 49.0 (C(3)), 51.6 (C(10)), 61.4 (C(5)), 62.7 (C(6a)), 85.6 (C(11a)); *m/z* (ESI⁺) 251 ([M + H]⁺, 100%); HRMS (ESI⁺) *m/z*: [M + H]⁺ Calcd for C₁₆H₃₁N₂⁺ 251.2482; Found 251.2472.

(2*R*,2'*S*,α*S*,α'*R*)-1-[N(1')-(α-Methylbenzyl)-1',2',5',6'-tetrahydropyridin-2'-yl]-2-[N-benzyl-N-(α'-methylbenzyl)amino]pent-4-ene 29. DIBAL-H (1.0 M in PhMe, 0.78 mL, 0.78 mmol) was added dropwise to a stirred solution of **22** (383 mg, 0.711 mmol, >99:1 dr) in PhMe (6 mL) at –78 °C and the resultant mixture was stirred at –78 °C for 1 h. MeOH (144 μL, 3.55 mmol) and Ph₃P=CH₂ (0.31 M in PhMe/THF, 22.9 mL, 7.11 mmol)²⁴ were added sequentially, and the resultant mixture was allowed to warm to rt and stirred at rt for 16 h, then concentrated in vacuo. The residue was dissolved in Et₂O (30 mL) and the resultant solution was filtered through a short plug of Celite® (eluent Et₂O) and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:5:1) gave **29** as a pale yellow oil (180 mg, 54%, >99:1 dr); [α]_D²⁵ +52.5 (c 1.0 in CHCl₃); ν_{max} (ATR) 3061, 3025, 2933, 2836 (C–H), 1601, 1492, 1452 (C=C); δ_H (400 MHz, CDCl₃) 1.26 (3H, d, *J* 6.6, C(α)Me), 1.34 (3H, d, *J* 6.9,

C(α')Me), 1.58–1.76 (3H, d, C(1) H_2 , C(5') H_A), 1.90–1.99 (1H, m, C(3) H_A), 2.00–2.15 (2H, m, C(3) H_B , C(5') H_B), 2.46 (1H, ddd, J 13.4, 5.4, 2.6, C(6') H_A), 2.83 (1H, ddd, J 13.4, 9.9, 4.7, C(6') H_B), 2.93 (1H, app quintet, J 6.2, C(2) H), 3.24–3.30 (1H, m, C(2') H), 3.73 (1H, d, J 14.8, NCH_AH_BPh), 3.81 (1H, d, J 14.8, NCH_AH_BPh), 3.82 (1H, q, J 6.6, C(α) H), 3.96 (1H, q, J 6.9, C(α') H), 4.86–4.88 (1H, m, C(5) H_A), 4.89–4.92 (1H, m, C(5) H_B), 5.54–5.65 (2H, m, C(4) H , C(3') H), 5.72–5.77 (1H, m, C(4') H), 7.18–7.37 (13H, m, Ph), 7.42–7.46 (2H, m, Ph); δ_C (100 MHz, $CDCl_3$) 18.2 (C(α')Me), 22.2 (C(5')), 22.5 (C(α)Me), 36.8 (C(1)), 37.1 (C(3)), 40.1 (C(6')), 50.1 (NCH_2Ph), 53.1 (C(2')), 55.1 (C(2)), 57.6 (C(α')), 58.3 (C(α)), 115.2 (C(5)), 125.2 (C(4')), 126.6, 126.8, 126.8 ($p-Ph$), 127.7, 128.1, 128.3, 128.3, 128.3, 128.5 ($o,m-Ph$), 130.6 (C(3')), 138.1 (C(4)), 142.3, 144.4, 145.5 ($i-Ph$); m/z (ESI^+) 465 ($[M + H]^+$, 100%); HRMS (ESI^+) m/z : $[M + H]^+$ Calcd for $C_{33}H_{41}N_2^+$ 465.3264; Found 465.3259.

(2*R*,2'*S*, α *S*, α' *R*,*Z*)-1-[*N*(1')-(α -Methylbenzyl)-1',2',5',6'-tetrahydropyridin-2'-yl]-2-[*N*-benzyl-*N*-(α' -methylbenzyl)amino]hept-4-ene 30. DIBAL-H (1.0 M in PhMe, 0.77 mL, 0.77 mmol) was added dropwise to a stirred solution of **22** (377 mg, 0.700 mmol, >99:1 dr) in PhMe (6 mL) at $-78^\circ C$ and the resultant mixture was stirred at $-78^\circ C$ for 1 h. MeOH (142 μ L, 3.50 mmol) and $Ph_3P=CHCH_2CH_3$ (0.31 M in PhMe/THF, 22.6 mL, 7.00 mmol)²⁵ were added sequentially, and the resultant mixture was allowed to warm to rt and stirred at rt for 16 h, then concentrated in vacuo. The residue was dissolved in Et_2O (30 mL) and the resultant solution was filtered through a short plug of Celite[®] (eluent Et_2O) and concentrated in vacuo. Purification via flash column chromatography (eluent $30-40^\circ C$ petrol/ Et_2O/NH_4OH , 100:5:1) gave **30** as a pale yellow oil (162 mg, 47%, >95:5 dr [(*Z*):(*E*)]); $[\alpha]_D^{25} +59.2$ (c 1.0 in $CHCl_3$); ν_{max} (ATR) 2970 (C–H), 1493, 1453 (C=C); δ_H (400 MHz, $CDCl_3$) 0.90 (3H, t, J 7.5, C(7) H_3), 1.26 (3H, d, J 6.6, C(α)Me), 1.34 (3H, d, J 6.9, C(α')Me), 1.60–1.80 (3H, m, C(1) H_2 , C(5') H_A), 1.91 (2H, app quintet, J 7.5, C(6) H_2), 1.97 (2H, app t, J 6.8, C(3) H_2), 2.04–2.15 (1H, m, C(5') H_B), 2.47 (1H, ddd, J 13.1, 5.3, 3.1, C(6') H_A), 2.80–2.92 (2H, m, C(2) H , C(6') H_B), 3.20–3.26 (1H, m, C(2') H), 3.73 (1H, d, J 14.8, NCH_AH_BPh), 3.81 (1H, d, J 14.8, NCH_AH_BPh), 3.83 (1H, q, J 6.6, C(α) H), 3.98 (1H, q, J 6.9, C(α') H), 5.12–5.19 (1H, m, C(4) H), 5.25–5.32 (1H, m, C(5) H), 5.56–5.61 (1H, m, C(3') H), 5.71–5.77 (1H, m, C(4') H), 7.17–7.44 (15H, m, Ph); δ_C (100 MHz, $CDCl_3$) 14.3 (C(7)), 18.3 (C(α')Me), 20.9 (C(6)), 22.3 (C(α)Me), 22.6 (C(5')), 30.0 (C(3)), 36.8 (C(1)), 40.2 (C(6')), 50.0 (NCH_2Ph), 53.2 (C(2')), 55.5 (C(2)), 57.6 (C(α')), 58.2 (C(α)), 125.1 (C(4')), 126.6, 126.7, 126.8 ($p-Ph$), 127.8, 128.0 ($o,m-Ph$), 128.1 (C(4)), 128.2, 128.2, 128.3, 128.5 ($o,m-Ph$), 130.8 (C(3')), 131.8 (C(5)), 142.4, 144.7, 145.2 ($i-Ph$); m/z (ESI^+) 493 ($[M + H]^+$, 100%); HRMS (ESI^+) m/z : $[M + H]^+$ Calcd for $C_{35}H_{45}N_2^+$ 493.3577; Found 493.3565.

(5*R*,6*aR*,11*aS*)-5-Propyldecahydro-5*H*-pyrido[1,2-*c*]pyrrolo[1,2-*a*]pyrimidine**[(+)-**

tetraponerine 3] 33. Palladium black (23 mg, 20% w/w) was added to a stirred, degassed solution of **29** (115 mg, 0.25 mmol, >99:1 dr) in MeOH/H₂O/AcOH (10:1:1, 2 mL). The resultant mixture was stirred under an atmosphere of H₂ (5 atm) at rt for 48 h. The reaction mixture was then filtered through a short plug of Celite[®] (eluent MeOH/H₂O/AcOH, 10:1:1) and concentrated in vacuo. The residue was dissolved in CH₂Cl₂ (2 mL), then K₂CO₃ (102 mg, 0.741 mmol) and 4-bromobutanal (75 mg, 0.49 mmol) were added to the resultant solution. The reaction mixture was stirred at rt for 2 h, then H₂O (2 mL) and satd aq NaHCO₃ (10 mL) were added. The reaction mixture was then extracted with CH₂Cl₂ (3 × 5 mL) and the combined organic extracts were dried and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Me₂CO/NH₄OH, 100:10:1) gave (+)-tetraponerine 3 (**33**) as a colourless oil (33 mg, 60% from **29**, >99:1 dr); $[\alpha]_{\text{D}}^{25} +30.9$ (*c* 1.0 in CHCl₃); {lit.⁴ for a sample isolated from the natural source $[\alpha]_{\text{D}}^{20} +27$ (*c* 0.07 in CHCl₃); lit.⁵ $[\alpha]_{\text{D}}^{20} +31$ (*c* 3.1 in CHCl₃); lit.^{9d} $[\alpha]_{\text{D}}^{20} +35$ (*c* 0.49 in CHCl₃)}; ν_{max} (ATR) 2954, 2928, 2859, 2802 (C–H);²⁶ δ_{H} (400 MHz, C₆D₆) 0.95 (3H, t, *J* 7.2, C(3')H₃), 1.09 (1H, app dt, *J* 12.7, 1.9, C(6)H_A), 1.15–1.28 (1H, m, C(9)H_A), 1.27–1.48 (5H, m, C(7)H₂, C(1')H_A, C(2')H₂), 1.48–1.84 (9H, m, C(1)H₂, C(2)H₂, C(8)H₂, C(9)H_B, C(10)H_A, C(1')H_B), 1.91 (1H, ddd, *J* 12.7, 11.9, 5.3, C(6)H_B), 1.99–2.07 (1H, m, C(6a)H), 2.72–2.86 (3H, m, C(3)H_A, C(5)H, C(10)H_B), 3.17 (1H, app q, *J* 7.5, C(3)H_B), 3.30 (1H, app t, *J* 3.5, C(11a)H); δ_{C} (100 MHz, C₆D₆) 14.5 (C(3')), 20.7 (C(2')), 22.2 (C(2)), 25.3 (C(9)), 26.6 (C(8)), 30.6 (C(1)), 32.2 (C(6)), 33.4 (C(1')), 34.3 (C(7)), 50.6 (C(3)), 51.0 (C(10)), 53.0 (C(5)), 56.7 (C(6a)), 75.4 (C(11a)); *m/z* (ESI⁺) 223 ([M + H]⁺, 100%); HRMS (ESI⁺) *m/z*: [M + H]⁺ Calcd for C₁₄H₂₇N₂⁺ 223.2169; Found 223.2168.

(5*R*,6*aR*,11*aS*)-5-Pentyldecahydro-5*H*-pyrido[1,2-*c*]pyrrolo[1,2-*a*]pyrimidine [(+)-tetraponerine

7] 34. Palladium black (22 mg, 20% w/w) was added to a stirred, degassed solution of **30** (108 mg, 0.22 mmol, >95:5 dr) in MeOH/H₂O/AcOH (10:1:1, 2 mL). The resultant mixture was stirred under an atmosphere of H₂ (5 atm) at rt for 48 h. The reaction mixture was then filtered through a short plug of Celite[®] (eluent MeOH/H₂O/AcOH, 10:1:1) and concentrated in vacuo. The residue was dissolved in CH₂Cl₂ (2 mL), then K₂CO₃ (91 mg, 0.66 mmol) and 4-bromobutanal (66 mg, 0.44 mmol) were added to the resultant solution. The reaction mixture was stirred at rt for 2 h, then H₂O (2 mL) and satd aq NaHCO₃ (10 mL) were added. The reaction mixture was then extracted with CH₂Cl₂ (3 × 5 mL) and the combined organic extracts were dried and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Me₂CO/NH₄OH, 100:10:1) gave (+)-tetraponerine 7 (**34**) as a colourless oil (24 mg, 44% from **30**, >99:1 dr); $[\alpha]_{\text{D}}^{25} +30.4$ (*c* 0.8 in CHCl₃); {lit.⁴ for a sample isolated from the natural source $[\alpha]_{\text{D}}^{20} +30$ (*c* 0.22

in CHCl_3); lit.^{9b} $[\alpha]_{\text{D}}^{20} +29.5$ (c 2.2 in CHCl_3); ν_{max} (ATR) 2928, 2856, 2790 (C–H);²⁶ δ_{H} (400 MHz, C_6D_6) 0.93 (3H, t, J 6.9, C(5') H_3), 1.13 (1H, app dt, J 12.9, 1.9, C(6) H_{A}), 1.17–1.86 (19H, m, C(1) H_2 , C(2) H_2 , C(7) H_2 , C(8) H_2 , C(9) H_2 , C(10) H_{A} , C(1') H_2 , C(2') H_2 , C(3') H_2 , C(4') H_2), 1.93 (1H, ddd, J 12.9, 11.9, 5.4, C(6) H_{B}), 2.02–2.09 (1H, m, C(6a) H), 2.74–2.86 (3H, m, C(3) H_{A} , C(5) H , C(10) H_{B}), 3.15–3.21 (1H, m, C(3) H_{B}), 3.33 (1H, app t, J 3.6, C(11a) H); δ_{C} (100 MHz, C_6D_6) 14.4 (C(5')), 22.2 (C(2)), 23.2 (C(4')), 25.3 (C(8)), 26.6 (C(9)), 27.4 (C(2')), 30.6 (C(1)), 31.1 (C(1')), 32.3 (C(6)), 32.5 (C(3')), 34.3 (C(7)), 50.7 (C(3)), 51.0 (C(10)), 53.3 (C(5)), 56.7 (C(6a)), 75.5 (C(11a)); m/z (ESI^+) 251 ($[\text{M} + \text{H}]^+$, 100%); HRMS (ESI^+) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{16}\text{H}_{31}\text{N}_2^+$ 251.2482; Found 251.2480.

tert-Butyl (3*S*, α *R*,*E*)-3-[*N*-allyl-*N*-(α -methylbenzyl)amino]hex-4-enoate 35. BuLi (2.30 M in hexanes, 14.0 mL, 32.3 mmol) was added dropwise to a stirred solution of (*R*)-*N*-benzyl-*N*-(α -methylbenzyl)amine (5.36 g, 33.3 mmol, >99:1 er) in THF (30 mL) at -78°C and the resultant mixture was stirred at -78°C for 30 min. A solution of **3** (3.50 g, 20.8 mmol, >99:1 dr) in THF (10 mL) at -78°C was added dropwise via cannula and the resultant mixture was stirred at -78°C for 2 h. Satd aq NH_4Cl (10 mL) was added and the reaction mixture was allowed to warm to rt, then concentrated in vacuo. The residue was partitioned between CH_2Cl_2 (100 mL) and 10% aq citric acid (100 mL), and the organic layer was washed with satd aq NaHCO_3 (100 mL) and brine (100 mL), then dried and concentrated in vacuo to give **35** in >95:5 dr. Purification via flash column chromatography (eluent 30–40 °C petrol/EtOAc, 20:1) gave **35** as a pale yellow oil (6.10 g, 89%, >99:1 dr);^{15b} $[\alpha]_{\text{D}}^{25} -3.5$ (c 2.0 in CHCl_3); {lit.^{15c} $[\alpha]_{\text{D}}^{20} -4.7$ (c 1.0 in CHCl_3)}; δ_{H} (400 MHz, CDCl_3) 1.36 (3H, d, J 6.8, C(α) Me), 1.40 (9H, s, CMe_3), 1.69 (3H, d, J 5.3, C(6) H_3), 2.26 (1H, dd, J 14.2, 8.4, C(2) H_{A}), 2.40 (1H, dd, J 14.2, 6.4, C(2) H_{B}), 3.10–3.14 (2H, m, C(1') H_2), 3.78–3.85 (1H, m, C(3) H), 4.00 (1H, q, J 6.8, C(α) H), 4.99 (1H, dd, J 10.3, 1.6, C(3') H_{A}), 5.07 (1H, dd, J 17.0, 1.6, C(3') H_{B}), 5.43–5.58 (2H, m, C(4) H , C(5) H), 5.77 (1H, ddt, J 17.0, 10.3, 6.3, C(2') H), 7.17–7.22 (1H, m, *Ph*), 7.25–7.31 (2H, m, *Ph*), 7.35–7.39 (2H, m, *Ph*).

tert-Butyl (2'*S*, α *R*)-2-[*N*(1')-(α -methylbenzyl)-2',5'-dihydro-1*H*-pyrrol-2'-yl]acetate 36. Grubbs I catalyst (230 mg, 0.36 mmol) was added to a stirred, degassed solution of **35** (3.00 g, 9.11 mmol, >99:1 dr) in anhydrous CH_2Cl_2 (EtOH stabilised, 300 mL) at rt. The resultant mixture was stirred at rt for 16 h, then concentrated in vacuo. The residue was dissolved in CH_2Cl_2 (80 mL), then $\text{P}(\text{CH}_2\text{OH})_3$ (4.52 g, 36.4 mmol) and Et_3N (2.54 mL, 18.2 mmol) were added sequentially. The resultant mixture was stirred at rt for 5 min, then excess silica gel (10 g) was added and the resultant mixture was stirred at rt for 12 h before being concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/ Et_2O / NH_4OH , 100:10:1) gave **36** as a pale yellow oil (2.25 g, 86%, >99:1 dr);^{15a} $[\alpha]_{\text{D}}^{25} +99.0$ (c 1.0 in CHCl_3); {lit.^{15b} for

ent-36: $[\alpha]_{\text{D}}^{24} -132.0$ (*c* 0.99 in CHCl_3); δ_{H} (400 MHz, CDCl_3) 1.43 (3H, d, *J* 6.7, C(α)Me), 1.46 (9H, s, CMe₃), 2.33 (1H, dd, *J* 14.5, 8.9, C(2)H_A), 2.59 (1H, dd, *J* 14.5, 4.0, C(2)H_B), 3.34–3.41 (1H, m, C(5')H_A), 3.57–3.63 (1H, m, C(5')H_B), 3.85 (1H, q, *J* 6.7, C(α)H), 4.10–4.18 (1H, m, C(2')H), 5.68–5.76 (2H, m, C(3')H, C(4')H), 7.19–7.34 (5H, m, *Ph*).

tert-Butyl (R,R)-2-[N(1')-(α -methylbenzyl)pyrrolidin-2'-yl]acetate 37. RhCl(PPh₃)₃ (100 mg, 0.11 mmol) was added to a degassed solution of **36** (1.00 g, 3.48 mmol, >99:1 dr) in EtOAc (5 mL) and the resultant mixture was stirred under H₂ (2 atm) at rt for 16 h. The reaction mixture was then filtered through Celite[®] (eluent EtOAc) and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O, 10:1) gave **37** as a colourless oil (860 mg, 86%, >99:1 dr);^{15b} $[\alpha]_{\text{D}}^{25} +57.0$ (*c* 1.0 in CHCl_3); {lit.^{15b} for **ent-37**: $[\alpha]_{\text{D}}^{23} -56.1$ (*c* 1.6 in CHCl_3)}; δ_{H} (400 MHz, CDCl_3) 1.44 (3H, d, *J* 6.8, C(α)Me), 1.44 (9H, s, CMe₃), 1.55–1.64 (2H, m, C(3')H_A, C(4')H_A), 1.65–1.77 (1H, m, C(4')H_B), 1.79–1.89 (1H, m, C(3')H_B), 2.18 (1H, dd, *J* 14.2, 9.9, C(2)H_A), 2.39 (1H, td, *J* 8.8, 6.9, C(5')H_A), 2.58 (1H, dd, *J* 14.2, 3.5, C(2)H_B), 2.76–2.82 (1H, m, C(5')H_B), 3.00–3.09 (1H, m, C(2')H), 3.76 (1H, q, *J* 6.8, C(α)H), 7.21–7.35 (5H, m, *Ph*).

tert-Butyl (R,R,E)-4-[N(1')-(α -methylbenzyl)pyrrolidin-2'-yl]but-2-enoate 38. Method A (from **37**): DIBAL-H (1.0 M in PhMe, 0.57 mL, 0.57 mmol) was added dropwise to a stirred solution of **37** (151 mg, 0.522 mmol, >99:1 dr) in PhMe (2 mL) at –78 °C and the resultant mixture was stirred at –78 °C for 1 h. MeOH (0.11 mL, 2.61 mmol) and Ph₃P=CHCO₂^tBu (196 mg, 0.522 mmol) were added sequentially, and the resultant mixture was allowed to warm to rt and stirred at rt for 16 h. The reaction mixture was then concentrated in vacuo to give a 75:25 mixture of **38** and **39**, respectively. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:20:1) gave a 9:91 mixture of **38** and **39**, respectively, as a pale yellow oil (22 mg, 13%). Data for **39**: δ_{H} (400 MHz, CDCl_3) 1.44 (3H, d, *J* 6.8, C(α)Me), 1.46–1.60 (2H, m, C(3')H_A, C(4')H_A), 1.50 (9H, s, CMe₃), 1.65–1.77 (2H, m, C(3')H_B, C(4')H_B), 2.37 (1H, app q, *J* 8.2, C(5')H_A), 2.65–2.79 (2H, m, C(4)H_A, C(2')H), 2.84–2.90 (1H, m, C(5')H_B), 2.94–3.03 (1H, m, C(4)H_B), 3.87 (1H, q, *J* 6.8, C(α)H), 5.75 (1H, dt, *J* 11.6, 1.7, C(2)H), 6.18 (1H, dt, *J* 11.6, 7.2, C(3)H), 7.21–7.37 (5H, m, *Ph*); δ_{C} (100 MHz, CDCl_3) 22.0 (C(α)Me), 22.6 (C(4')), 28.4 (CMe₃), 29.9 (C(3')), 32.6 (C(4)), 49.0 (C(5')), 58.9 (C(2')), 59.2 (C(α)), 80.1 (CMe₃), 122.5 (C(2)), 126.9 (*p*-Ph), 128.1, 128.2 (*o,m*-Ph), 142.4 (*i*-Ph), 146.2 (C(3)), 166.2 (C(1)). Data for mixture: ν_{max} (ATR) 2972, 2873 (C–H), 1713 (C=O), 1651 (C=C); *m/z* (ESI⁺) 316 ([M + H]⁺, 100%); HRMS (ESI⁺) *m/z*: [M + H]⁺ Calcd for C₂₀H₃₀NO₂⁺ 316.2271; Found 316.2269. Further elution (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:20:1) gave an 86:14 mixture of **38** and **39**, respectively, as a colourless oil (108 mg, 66%).

Method B (from 36) – Step 1: RhCl(PPh₃)₃ (320 mg, 0.348 mmol) was added to a degassed solution of **36** (5.00 g, 17.4 mmol) in EtOAc (100 mL) and the resultant mixture was stirred under H₂ (2 atm) at rt for 16 h. The reaction mixture was then filtered through Celite[®] (eluent EtOAc) and concentrated in vacuo to give **37**.

Method B (from 36) – Step 2: DIBAL-H (1.0 M in PhMe, 19.1 mL, 19.1 mmol) was added dropwise to a stirred solution of the residue of **37** from the previous step in PhMe (70 mL) at –78 °C and the resultant mixture was stirred at –78 °C for 1 h. MeOH (3.57 mL, 87.0 mmol) and Ph₃P=CHCO₂^tBu (6.55 g, 17.4 mmol) were added sequentially, and the resultant mixture was allowed to warm to rt and stirred at rt for 16 h. The reaction mixture was then concentrated in vacuo to give a 75:25 mixture of **38** and **39**, respectively. Purification via exhaustive flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:20:1) gave **38** as a pale yellow oil (2.76 g, 50% from **36**, 98:2 dr); [α]_D²⁵ +82.7 (*c* 1.0 in CHCl₃); ν_{max} (ATR) 2972, 2873 (C–H), 1713 (C=O); δ_H (400 MHz, CDCl₃) 1.44 (3H, d, *J* 6.8, C(α)*Me*), 1.48 (9H, s, *CMe*₃), 1.48–1.81 (4H, m, C(3')*H*₂, C(4')*H*₂), 2.18 (1H, app dt, *J* 14.2, 9.0, C(4)*H*_A), 2.40 (1H, app td, *J* 8.7, 7.2, C(5')*H*_A), 2.50 (1H, dddd, *J* 14.2, 7.0, 3.3, 1.5, C(4)*H*_B), 2.70–2.78 (1H, m, C(2')*H*), 2.80–2.86 (1H, m, C(5')*H*_B), 3.80 (1H, q, *J* 6.8, C(α)*H*), 5.76 (1H, dt, *J* 15.6, 1.5, C(2)*H*), 6.83 (1H, app dt, *J* 15.6, 7.0, C(3)*H*), 7.21–7.34 (5H, m, *Ph*); δ_C (100 MHz, CDCl₃) 22.2 (C(α)*Me*), 22.7 (C(4')), 28.3 (*CMe*₃), 30.2 (C(3')), 37.6 (C(4)), 49.9 (C(5')), 58.7 (C(2')), 60.3 (C(α)), 80.2 (*CMe*₃), 124.4 (C(2)), 127.0 (*p-Ph*), 128.0, 128.2 (*o,m-Ph*), 142.9 (*i-Ph*), 145.9 (C(3)), 166.1 (C(1)); *m/z* (ESI⁺) 316 ([M + H]⁺, 100%); HRMS (ESI⁺) *m/z*: [M + H]⁺ Calcd for C₂₀H₃₀NO₂⁺ 316.2271; Found 316.2270.

tert-Butyl (R,R,R)-3-[N-benzyl-N-isopropylamino]-4-[N(1')-(α-methylbenzyl)pyrrolidin-2'-yl]butanoate and *tert*-butyl (3S,2'R,αR)-3-[N-benzyl-N-isopropylamino]-4-[N(1')-(α-methylbenzyl)pyrrolidin-2'-yl]butanoate **40** and **41**. BuLi (2.3 M in hexanes, 0.11 mL, 0.253 mmol) was added dropwise to a stirred solution of *N*-benzyl-*N*-isopropylamine (38 mg, 0.25 mmol) in THF (0.5 mL) at –78 °C and the resultant mixture was stirred at –78 °C for 30 min. A solution of **38** (50 mg, 0.159 mmol, 98:2 dr [(*E*):(*Z*)] in THF (0.5 mL) at –78 °C was added via cannula and the resultant mixture was stirred at –78 °C for 2 h. Satd aq NH₄Cl (0.2 mL) was added and the reaction mixture was allowed to warm to rt, then concentrated in vacuo. The residue was partitioned between CH₂Cl₂ (5 mL) and 10% citric acid (5 mL), and the organic layer was washed with satd aq NaHCO₃ (5 mL) and brine (5 mL), then dried and concentrated in vacuo to give a 50:50 mixture of **40** and **41**. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:20:1) gave a 50:50 mixture of **40** and **41** as a colourless oil (52 mg, 70%). Data for **40**: δ_H (400 MHz, CDCl₃) 0.96–1.06 (6H, m, NCHMe₂), 1.07–1.31 (2H, m, C(4)*H*_A, C(3')*H*_A), 1.34–1.41 (1H, m, C(3')*H*_B), 1.39 (3H, d, *J* 6.7, C(α)*Me*), 1.45 (9H, s, *CMe*₃), 1.48–1.62 (2H, m, C(4')*H*₂), 1.87 (1H,

ddd, J 13.5, 10.2, 3.8, C(4) H_B), 2.10–2.18 (1H, m, C(2) H_A), 2.33–2.42 (1H, m, C(5') H_A), 2.64 (1H, dd, J 13.8, 3.8, C(2) H_B), 2.70 (1H, ddd, J 9.2, 7.4, 3.7, C(5') H_B), 2.87–2.97 (2H, m, C(2') H , NCHMe₂), 3.11 (1H, app tt, J 14.5, 3.8, C(3) H), 3.43 (1H, d, J 14.1, NCH_AH_BPh), 3.70 (1H, d, J 14.1, NCH_AH_BPh), 3.74 (1H, q, J 6.7, C(α) H), 7.17–7.40 (10H, m, *Ph*). Data for **41**: δ_H (400 MHz, CDCl₃) 0.96–1.06 (6H, m, NCHMe₂), 1.45 (3H, d, J 6.9, C(α)*Me*), 1.45 (9H, s, CMe₃), 1.46–1.63 (3H, m, C(4) H_A , C(3') H_A , C(4') H_A), 1.67–1.87 (3H, m, C(4) H_B , C(3') H_B , C(4') H_B), 2.10–2.18 (1H, m, C(2) H_A), 2.27–2.43 (3H, m, C(2) H_B , C(2') H , C(5') H_A), 2.82–2.97 (2H, m, C(5') H_B , NCHMe₂), 3.22–3.29 (1H, m, C(3) H), 3.62 (1H, d, J 14.6, NCH_AH_BPh), 3.75 (1H, d, J 14.6, NCH_AH_BPh), 3.84 (1H, q, J 6.9, C(α) H), 7.17–7.40 (10H, m, *Ph*). Data for mixture of **40** and **41**: $\nu_{\max}$ (ATR) 3026, 2966, 2932, 2871 (C–H), 1724 (C=O); δ_C (100 MHz, CDCl₃) 18.8, 19.8, 21.4, 21.7, 22.5, 22.8, 22.9, 23.0, 28.2, 28.2, 29.6, 30.5, 37.4, 37.8, 39.2, 40.0, 46.8, 48.7, 48.9, 49.1, 49.3, 50.4, 52.6, 52.9, 57.5, 58.2, 59.4, 61.0, 80.1, 80.2, 126.6, 126.6, 126.7, 127.0, 127.9, 128.0, 128.0, 128.1, 128.1, 128.2, 128.6, 129.0, 141.1, 141.8, 142.1, 144.2, 172.4, 172.4; m/z (ESI⁺) 465 ([M + H]⁺, 100%); HRMS (ESI⁺) m/z : [M + H]⁺ Calcd for C₃₀H₄₅N₂O₂⁺ 465.3476; Found 465.3475.

tert-Butyl

(*R,R,R,R*)-3-[*N*-benzyl-*N*-(α -methylbenzyl)amino]-4-[*N*(1')-(α '-

methylbenzyl)pyrrolidin-2'-yl]butanoate **42**. *Method A*: BuLi (2.3 M in hexanes, 0.85 mL, 1.96 mmol) was added dropwise to a stirred solution of (*R*)-*N*-benzyl-*N*-(α -methylbenzyl)amine (429 mg, 2.03 mmol, >99:1 *er*) in THF (3 mL) at –78 °C and the resultant mixture was stirred at –78 °C for 30 min. A solution of **38** (400 mg, 1.27 mmol, 98:2 *dr* [(*E*):(*Z*)] in THF (2 mL) at –78 °C was added and the resultant mixture was stirred at –78 °C for 2 h. Satd aq NH₄Cl (1 mL) was added and the reaction mixture was allowed to warm to rt, then concentrated in vacuo. The residue was partitioned between CH₂Cl₂ (10 mL) and 10% citric acid (10 mL), and the organic layer was washed with satd aq NaHCO₃ (10 mL) and brine (10 mL), then dried and concentrated in vacuo to give **42** in >95:5 *dr*. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:20:1) gave **42** as a pale yellow oil (655 mg, 98%, >99:1 *dr*); [α]_D²⁵ +45.1 (*c* 1.0 in CHCl₃); $\nu_{\max}$ (ATR) 3027, 2971, 2932, 2872 (C–H), 1723 (C=O); δ_H (400 MHz, CDCl₃) 1.13 (1H, ddd, J 13.7, 10.9, 2.7, C(4) H_A), 1.32 (3H, d, J 7.0, C(α)*Me*), 1.38 (3H, obscd, C(α')*Me*), 1.40 (9H, s, CMe₃), 1.42–1.50 (1H, m, C(3') H_A), 1.53–1.74 (3H, m, C(3') H_B , C(4') H_2), 1.75–1.83 (2H, m, C(2) H_2), 1.92 (1H, ddd, J 13.7, 10.8, 2.7, C(4) H_B), 2.42 (1H, app q, J 8.1, C(5') H_A), 2.73–2.79 (1H, m, C(5') H_B), 3.10–3.18 (1H, m, C(2') H), 3.34–3.41 (1H, m, C(3) H), 3.44 (1H, d, J 14.8, NCH_AH_BPh), 3.73–3.81 (2H, m, C(α) H , C(α') H), 3.87 (1H, d, J 14.8, NCH_AH_BPh), 7.16–7.40 (15H, m, *Ph*); δ_C (100 MHz, CDCl₃) 20.8 (C(α)*Me*), 23.0 (C(α')*Me*), 23.1 (C(4')), 28.2 (CMe₃), 29.8 (C(3')), 37.9 (C(2)), 38.8 (C(4)), 50.4 (NCH₂Ph), 50.7 (C(5')), 51.8 (C(3)), 57.1 (C(α)), 57.5 (C(2')), 61.0 (C(α')), 80.1 (CMe₃), 126.7, 126.8, 127.1 (*p*-*Ph*), 127.9, 128.2,

128.2, 128.3, 128.4, 128.5 (*o,m-Ph*), 141.1, 142.5, 144.2 (*i-Ph*), 172.1 (*C*(1)); m/z (ESI^+) 527 ($[\text{M} + \text{H}]^+$, 100%); HRMS (ESI^+) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{35}\text{H}_{47}\text{N}_2\text{O}_2^+$ 527.3632; Found 527.3626.

Method B: BuLi (2.3 M in hexanes, 0.22 mL, 0.50 mmol) was added dropwise to a stirred solution of (*R*)-*N*-benzyl-*N*-(α -methylbenzyl)amine (109 mg, 0.52 mmol, >99:1 er) in THF (2 mL) at -78°C and the resultant mixture was stirred at -78°C for 30 min. A solution of **38** and **39** (102 mg, 0.32 mmol, 86:14 dr [(*E*):(*Z*)]) in THF (1 mL) at -78°C was added and the resultant mixture was stirred at -78°C for 2 h. Satd aq NH_4Cl (0.5 mL) was added and the reaction mixture was allowed to warm to rt, then concentrated in vacuo. The residue was partitioned between CH_2Cl_2 (5 mL) and 10% citric acid (5 mL), and the organic layer was washed with satd aq NaHCO_3 (5 mL) and brine (5 mL), then dried and concentrated in vacuo to give an 86:14 mixture of **42** and **44**, respectively. Purification via flash column chromatography (eluent 30–40 $^\circ\text{C}$ petrol/ $\text{Et}_2\text{O}/\text{NH}_4\text{OH}$, 100:20:1) gave a 90:10 mixture of **42** and **44**, respectively, as a pale yellow oil (143 mg). Data for **44**: δ_{H} (400 MHz, CDCl_3) [selected peaks] 2.28–2.36 (1H, m, $\text{C}(5')\text{H}_{\text{A}}$) 2.85–2.95 (2H, m, $\text{C}(2')\text{H}$, $\text{C}(5')\text{H}_{\text{B}}$), 2.99 (1H, dd, J 15.9, 6.9, $\text{C}(2)\text{H}_{\text{A}}$), 3.05 (1H, dd, J 15.9, 6.9, $\text{C}(2)\text{H}_{\text{B}}$), 5.52 (1H, dd, J 15.5, 8.3, $\text{C}(4)\text{H}$), 5.65 (1H, app dt, J 15.5, 6.9, $\text{C}(3)\text{H}$); m/z (ESI^+) 527 ($[\text{M}(\textbf{42}) + \text{H}]^+$, 100%); 316 ($[\text{M}(\textbf{44}) + \text{H}]^+$, 32%).

***tert*-Butyl (3*S*,2'*R*, α *S*, α' *R*)-3-[*N*-benzyl-*N*-(α -methylbenzyl)amino]-4-[*N*(1')-(α' -methylbenzyl)pyrrolidin-2'-yl]butanoate **43**.** BuLi (2.3 M in hexanes, 0.85 mL, 1.97 mmol) was added dropwise to a stirred solution of (*S*)-*N*-benzyl-*N*-(α -methlbenzyl)amine (429 mg, 2.03 mmol, >99:1 er) in THF (3 mL) at -78°C and the resultant mixture was stirred at -78°C for 30 min. A solution of **38** (400 mg, 1.27 mmol, 98:2 dr [(*E*):(*Z*)]) in THF (2 mL) at -78°C was added and the resultant mixture was stirred at -78°C for 2 h. Satd aq NH_4Cl (1 mL) was added and the reaction mixture was allowed to warm to rt, then concentrated in vacuo. The residue was partitioned between CH_2Cl_2 (10 mL) and 10% citric acid (10 mL), and the organic layer was washed with satd aq NaHCO_3 (10 mL) and brine (10 mL), then dried and concentrated in vacuo to give **43** in >95:5 dr. Purification via flash column chromatography (eluent 30–40 $^\circ\text{C}$ petrol/ $\text{Et}_2\text{O}/\text{NH}_4\text{OH}$, 100:20:1) gave **43** as a pale yellow oil (615 mg, 92%, >99:1 dr); $[\alpha]_{\text{D}}^{25} +72.8$ (c 1.0 in CHCl_3); ν_{max} (ATR) 3027, 2972, 2932, 2801 (C–H), 1725 (C=O); δ_{H} (400 MHz, CDCl_3) 1.34 (3H, d, J 6.9, $\text{C}(\alpha)\text{Me}$), 1.38 (9H, s, CMe_3), 1.42 (3H, d, J 6.9, $\text{C}(\alpha')\text{Me}$), 1.41–1.47 (1H, m, $\text{C}(3')\text{H}_{\text{A}}$), 1.52–1.78 (4H, m, $\text{C}(4)\text{H}_{\text{A}}$, $\text{C}(3')\text{H}_{\text{B}}$, $\text{C}(4')\text{H}_2$), 1.84 (1H, ddd, J 13.0, 9.8, 2.5, $\text{C}(4)\text{H}_{\text{B}}$), 1.93 (1H, dd, J 15.1, 6.6, $\text{C}(2)\text{H}_{\text{A}}$), 2.10 (1H, dd, J 15.1, 5.8, $\text{C}(2)\text{H}_{\text{B}}$), 2.32–2.47 (2H, m, $\text{C}(2')\text{H}$, $\text{C}(5')\text{H}_{\text{A}}$), 2.79–2.85 (1H, m, $\text{C}(5')\text{H}_{\text{B}}$), 3.31–3.39 (1H, m, $\text{C}(3)\text{H}$), 3.62 (1H, d, J 14.8, $\text{NCH}_\text{A}\text{H}_\text{B}\text{Ph}$), 3.75 (1H, d, J 14.8, $\text{NCH}_\text{A}\text{H}_\text{B}\text{Ph}$), 3.80 (1H, q, J 6.9, $\text{C}(\alpha')\text{H}$), 3.91 (1H, q, J 6.9, $\text{C}(\alpha)\text{H}$), 7.17–7.40 (15H, m, *Ph*); δ_{C} (100 MHz, CDCl_3) 19.8 ($\text{C}(\alpha)\text{Me}$), 21.9

(C(α')Me), 22.7 (C(4')), 28.2 (CMe₃), 30.9 (C(3')), 38.8 (C(4)), 39.0 (C(2)), 49.2 (C(5')), 50.4 (NCH₂Ph), 53.4 (C(3)), 58.2 (C(2')), 58.6 (C(α)), 59.8 (C(α')), 80.1 (CMe₃), 126.6, 126.9, 127.0 (*p*-Ph), 128.1, 128.1, 128.1, 128.2, 128.2, 128.5 (*o,m*-Ph), 141.8, 142.5, 144.1 (*i*-Ph), 172.1 (C(1)); *m/z* (ESI⁺) 527 ([M + H]⁺, 100%); HRMS (ESI⁺) *m/z*: [M + H]⁺ Calcd for C₃₅H₄₇N₂O₂⁺ 527.3632; Found 527.3627.

(2*S*,2'*R*, α *R*, α' *R*)-1-[N(1')-(α -Methylbenzyl)pyrrolidin-2'-yl]-2-[N-benzyl-N-(α' -methylbenzyl)amino]pent-4-ene 45. DIBAL-H (1.0 M in PhMe, 0.59 mL, 0.59 mmol) was added dropwise to a stirred solution of **42** (284 mg, 0.539 mmol, >99:1 dr) in PhMe (5 mL) at -78 °C and the resultant mixture was stirred at -78 °C for 1 h. MeOH (109 μ L, 2.70 mmol) and Ph₃P=CH₂ (0.31 M suspension in PhMe/THF 17.4 mL, 5.39 mmol)²⁴ were added sequentially, and the resultant mixture was allowed to warm to rt and stirred at rt for 16 h, then concentrated in vacuo. The residue was dissolved in Et₂O (40 mL) and the resultant solution was filtered through a short plug of Celite[®] (eluent Et₂O) and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:10:1) gave **45** as a colourless oil (185 mg, 76%, >99:1 dr); [α]_D²⁵ +76.9 (*c* 1.0 in CHCl₃); $\nu_{\max}$ (ATR) 3026, 2970, 2933, 2870, 2803 (C–H), 1638, 1601, 1493, 1452 (C=C); δ_{H} (400 MHz, CDCl₃) 1.21–1.31 (2H, m, C(1)*H*_A, C(3')*H*_A), 1.27 (3H, d, *J* 7.0, C(α')Me), 1.37 (3H, d, *J* 6.8, C(α)Me), 1.48–1.72 (4H, m, C(3)*H*_A, C(3')*H*_B, C(4')*H*₂), 1.72–1.80 (1H, m, C(3)*H*_B), 1.85 (1H, ddd, *J* 13.7, 9.9, 3.3, C(1)*H*_B), 2.36 (1H, app q, *J* 8.2, C(5')*H*_A), 2.73–2.81 (2H, m, C(2)*H*, C(5')*H*_B), 2.99–3.08 (1H, m, C(2')*H*), 3.57 (1H, d, *J* 15.0, NCH_AH_BPh), 3.79 (1H, q, *J* 6.8, C(α)*H*), 3.83 (1H, q, *J* 7.0, C(α')*H*), 3.92 (1H, d, *J* 15.0, NCH_AH_BPh), 4.81 (1H, d, *J* 17.1, C(5)*H*_A), 4.87 (1H, d, *J* 10.3, C(5)*H*_B), 5.46–5.57 (1H, m, C(4)*H*), 7.16–7.36 (13H, m, *Ph*), 7.39–7.43 (2H, m, *Ph*); δ_{C} (100 MHz, CDCl₃) 20.8 (C(α')Me), 22.7 (C(α)Me), 22.9 (C(4')), 30.3 (C(3')), 35.9 (C(3)), 37.6 (C(1)), 50.1 (C(5')), 50.4 (NCH₂Ph), 55.0 (C(2)), 57.4 (C(α')), 57.9 (C(2')), 60.5 (C(α)), 116.1 (C(5)), 126.6, 126.8, 127.0 (*p*-Ph), 128.1, 128.1, 128.1, 128.2, 128.4, 128.4 (*o,m*-Ph), 137.4 (C(4)), 141.8, 143.6, 143.7 (*i*-Ph); *m/z* (ESI⁺) 453 ([M + H]⁺, 100%); HRMS (ESI⁺) *m/z*: [M + H]⁺ Calcd for C₃₂H₄₁N₂⁺ 453.3264; Found 453.3259.

(2*S*,2'*R*, α *R*, α' *R,Z*)-1-[N(1')-(α -Methylbenzyl)pyrrolidin-2'-yl]-2-[N-benzyl-N-(α' -methylbenzyl)amino]hept-4-ene 46. DIBAL-H (1.0 M in PhMe, 0.53 mL, 0.53 mmol) was added dropwise to a stirred solution of **42** (255 mg, 0.484 mmol, >99:1 dr) in PhMe (5 mL) at -78 °C and the resultant mixture was stirred at -78 °C for 1 h. MeOH (98 μ L, 2.42 mmol) and Ph₃P=CHCH₂CH₃ (0.31 M suspension in PhMe/THF 15.6 mL, 4.84 mmol)²⁵ were added sequentially, and the resultant mixture was allowed to warm to rt and stirred at rt for 16 h, then concentrated in vacuo. The residue was dissolved in Et₂O (40 mL) and the resultant solution was filtered through a short plug of Celite[®] (eluent Et₂O) and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:10:1) gave

46 as a colourless oil (174 mg, 75%, >95:5 dr [(Z):(E)]); $[\alpha]_{\text{D}}^{25} +78.7$ (*c* 1.0 in CHCl_3); ν_{max} (ATR) 3061, 3026, 2966, 2932, 2872 (C–H), 1602, 1493, 1453 (C=C); δ_{H} (400 MHz, CDCl_3) 0.90 (3H, t, *J* 7.5, C(7)*H*₃), 1.16–1.30 (2H, m, C(1)*H*_A, C(3')*H*_A), 1.27 (3H, d, *J* 7.0, C(α')*Me*), 1.38 (3H, d, *J* 6.8, C(α)*Me*), 1.51–1.65 (3H, m, C(3')*H*_B, C(4')*H*₂), 1.72–1.95 (5H, m, C(1)*H*_B, C(3)*H*₂, C(6)*H*₂), 2.39 (1H, app q, *J* 7.9, C(5')*H*_A), 2.69–2.81 (2H, m, C(2)*H*, C(5')*H*_B), 2.98–3.06 (1H, m, C(2')*H*), 3.60 (1H, d, *J* 15.1, $\text{NCH}_A\text{H}_B\text{Ph}$), 3.80 (1H, q, *J* 6.8, C(α)*H*), 3.86 (1H, q, *J* 7.0, C(α')*H*), 3.96 (1H, d, *J* 15.1, $\text{NCH}_A\text{H}_B\text{Ph}$), 5.03–5.12 (1H, m, C(4)*H*), 5.24–5.32 (1H, m, C(5)*H*), 7.18–7.37 (13H, m, *Ph*), 7.40–7.43 (2H, m, *Ph*); δ_{C} (100 MHz, CDCl_3) 14.3 (C(7)), 20.8 (C(6)), 21.1 (C(α')*Me*), 22.6 (C(α)*Me*), 22.8 (C(4')), 28.7 (C(3)), 30.2 (C(3')), 37.5 (C(1)), 50.0 (C(5')), 50.5 (NCH_2Ph), 55.8 (C(2)), 57.8 (C(α')), 57.8 (C(2')), 60.3 (C(α)), 126.5, 126.8, 126.9 (*p-Ph*), 127.5 (C(4)), 128.0, 128.1, 128.1, 128.2, 128.3, 128.4 (*o,m-Ph*), 132.5 (C(5)), 142.1, 143.7, 144.3 (*i-Ph*); *m/z* (ESI^+) 481 ($[\text{M} + \text{H}]^+$, 100%); HRMS (ESI^+) *m/z*: $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{34}\text{H}_{45}\text{N}_2^+$ 481.3577; Found 481.3574.

(5S,6aR,10aS)-5-Propyldecahydrodipyrrolo[1,2-*a*:1',2'-*c*]pyrimidine [(+)-tetraponerine 2] 49.

Palladium black (37 mg, 20% w/w) was added to a stirred, degassed solution of **45** (183 mg, 0.404 mmol, >99:1 dr) in MeOH/H₂O/AcOH (10:1:1, 2 mL). The resultant mixture was stirred under an atmosphere of H₂ (5 atm) at rt for 48 h. The reaction mixture was then filtered through a short plug of Celite® (eluent MeOH/H₂O/AcOH, 10:1:1) and concentrated in vacuo. The residue was dissolved in CH_2Cl_2 (2 mL), then K₂CO₃ (168 mg, 1.21 mmol) and 4-bromobutanal (123 mg, 0.81 mmol) were added to the resultant solution. The reaction mixture was stirred at rt for 2 h, then H₂O (2 mL) and satd aq NaHCO₃ (10 mL) were added. The reaction mixture was then extracted with CH_2Cl_2 (3 × 5 mL) and the combined organic extracts were dried and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Me₂CO/NH₄OH, 100:10:1) gave (+)-tetraponerine 2 (**49**) as a colourless oil (31 mg, 37% from **45**, >99:1 dr); $[\alpha]_{\text{D}}^{25} +44.2$ (*c* 1.0 in CHCl_3); {lit.⁵ $[\alpha]_{\text{D}}^{20} +36$ (*c* 1.79 in CHCl_3); lit.⁶ $[\alpha]_{\text{D}}^{20} +47$ (*c* 0.232 in CHCl_3)}; ν_{max} (ATR) 2957, 2931, 2871, 2786 (C–H);²⁶ δ_{H} (400 MHz, C_6D_6) 0.92 (3H, t, *J* 7.2, C(3')*H*₃), 1.27–1.83 (14H, m, C(1)*H*₂, C(2)*H*₂, C(6)*H*₂, C(7)*H*₂, C(8)*H*₂, C(1')*H*₂, C(2')*H*₂), 1.83–1.97 (2H, m, C(9)*H*_A, C(6a)*H*), 2.29–2.36 (1H, m, C(3)*H*_A), 2.39–2.46 (1H, m, C(5)*H*), 2.86 (1H, app t, *J* 5.3, C(10a)*H*), 2.93 (1H, app td, *J* 8.5, 2.5, C(9)*H*_B), 3.01–3.07 (1H, m, C(3)*H*_B); δ_{C} (100 MHz, C_6D_6) 14.7 (C(3')), 19.3 (C(2')), 21.0 (C(2)), 21.3 (C(8)), 29.3 (C(1)), 30.6 (C(7)), 33.4 (C(6)), 37.0 (C(1')), 45.8 (C(3)), 49.2 (C(9)), 59.4 (C(5)), 64.2 (C(6a)), 83.4 (C(10a)); *m/z* (ESI^+) 209 ($[\text{M} + \text{H}]^+$, 100%); HRMS (ESI^+) *m/z*: $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{13}\text{H}_{25}\text{N}_2^+$ 209.2012; Found 209.2012.

(5S,6aR,10aS)-5-Pentyldecahydrodipyrrolo[1,2-*a*:1',2'-*c*]pyrimidine [(+)-tetraponerine 6] 50.

Palladium black (31 mg, 20% w/w) was added to a stirred, degassed solution of **46** (155 mg, 0.322 mmol,

>95:5 dr [(E):(Z)] in MeOH/H₂O/AcOH (10:1:1, 2 mL). The resultant mixture was stirred under an atmosphere of H₂ (5 atm) at rt for 48 h. The reaction mixture was then filtered through a short plug of Celite[®] (eluent MeOH/H₂O/AcOH, 10:1:1) and concentrated in vacuo. The residue was dissolved in CH₂Cl₂ (2 mL), then K₂CO₃ (134 mg, 0.97 mmol) and 4-bromobutanal (98 mg, 0.65 mmol) were added to the resultant solution. The reaction mixture was stirred at rt for 2 h, then H₂O (2 mL) and satd aq NaHCO₃ (10 mL) were added. The reaction mixture was then extracted with CH₂Cl₂ (3 × 5 mL) and the combined organic extracts were dried and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Me₂CO/NH₄OH, 100:10:1) gave (+)-tetraponerine 6 (**50**) as a pale yellow oil (41 mg, 54% from **46**, >99:1 dr); [α]_D²⁵ +66.3 (c 1.0 in CHCl₃); {lit.⁴ for a sample isolated from the natural source [α]_D²⁰ +35 (c 0.15 in CHCl₃); lit.⁶ [α]_D²⁰ +40 (c 0.75 in CHCl₃)}; ν_{max} (ATR) 2955, 2928, 2857, 2785 (C–H);²⁶ δ_H (400 MHz, C₆D₆) 0.91 (3H, t, *J* 7.0, C(5')H₃), 1.23–1.54 (11H, m, C(6)H₂, C(7)H_A, C(8)H_A, C(1')H_A, C(2')H₂, C(3')H₂, C(4')H₂), 1.54–1.98 (9H, m, C(1)H₂, C(2)H₂, C(6a)H, C(7)H_B, C(8)H_B, C(9)H_A, C(1')H_B), 2.31–2.38 (1H, m, C(3)H_A), 2.39–2.46 (1H, m, C(5)H), 2.87 (1H, app t, *J* 5.3, C(10a)H), 2.94 (1H, app td, *J* 8.5, 2.5, C(9)H_B), 3.03–3.10 (1H, m, C(3)H_B); δ_C (100 MHz, C₆D₆) 14.4 (C(5')), 21.0 (C(2)), 21.3 (C(8)), 23.2 (C(4')), 25.9 (C(2')), 29.3 (C(1)), 30.6 (C(7)), 32.7 (C(3')), 33.5 (C(6)), 34.8 (C(1')), 45.9 (C(3)), 49.2 (C(9)), 59.7 (C(5)), 64.2 (C(6a)), 83.5 (C(10a)); *m/z* (ESI⁺) 237 ([M + H]⁺, 100%); HRMS (ESI⁺) *m/z*: [M + H]⁺ Calcd for C₁₅H₂₉N₂⁺ 237.2325; Found 237.2326.

(2*R*,2'*R*,α*S*,α'*R*)-1-[*N*(1')-(α-Methylbenzyl)pyrrolidin-2'-yl]-2-[*N*-benzyl-*N*-(α'-methylbenzyl)amino]pent-4-ene 51. DIBAL-H (1.0 M in PhMe, 0.65 mL, 0.65 mmol) was added dropwise to a stirred solution of **43** (200 mg, 0.380 mmol, >99:1 dr) in PhMe (5 mL) at –78 °C and the resultant mixture was stirred at –78 °C for 1 h. MeOH (46 μL, 1.1 mmol) and Ph₃P=CH₂ (0.31 M suspension in PhMe/THF, 6.1 mL, 1.9 mmol)²⁴ were added sequentially, and the resultant mixture was allowed to warm to rt and stirred at rt for 16 h, then concentrated in vacuo. The residue was dissolved in Et₂O (40 mL) and the resultant solution was filtered through a short plug of Celite[®] (eluent Et₂O) and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:20:1) gave **51** as a colourless oil (68 mg, 40%, >99:1 dr); [α]_D²⁵ +56.3 (c 1.0 in CHCl₃); ν_{max} (ATR) 3026, 2968, 2932, 2872 (C–H), 1638, 1602, 1493, 1452 (C=C); δ_H (400 MHz, CDCl₃) 1.31 (3H, d, *J* 6.9, C(α)Me), 1.43 (3H, d, *J* 6.8, C(α')Me), 1.42–1.48 (1H, m, C(3')H_A), 1.52–1.76 (4H, m, C(1)H_A, C(3')H_B, C(4')H₂), 1.82–1.90 (2H, m, C(1)H_B, C(3)H_A), 1.99–2.08 (1H, m, C(3)H_B), 2.36–2.48 (2H, m, C(2')H, C(5')H_A), 2.62–2.70 (1H, m, C(2)H), 2.81–2.87 (1H, m, C(5')H_B), 3.74 (1H, d, *J* 14.7, NCH_AH_BPh), 3.79 (1H, d, *J* 14.7, NCH_AH_BPh), 3.83 (1H, q, *J* 6.8, C(α')H), 3.97 (1H, q, *J* 6.9, C(α)H), 4.80–4.86 (2H, m, C(5)H₂), 5.52 (1H, ddt, *J* 15.8,

11.4, 7.0, C(4)*H*), 7.16–7.35 (13*H*, m, *Ph*), 7.39–7.43 (2*H*, m, *Ph*); δ_{C} (100 MHz, CDCl₃) 17.9 (C(α)*Me*), 21.9 (C(α')*Me*), 22.4 (C(4')), 30.5 (C(3')), 36.7 (C(3)), 37.6 (C(1)), 48.8 (C(5')), 50.2 (NCH₂Ph), 55.4 (C(2)), 57.1 (C(α)), 58.5 (C(2')), 59.3 (C(α')), 115.3 (C(5)), 126.6, 126.7, 127.0 (*p-Ph*), 128.0, 128.0, 128.1, 128.2, 128.3, 128.5 (*o,m-Ph*), 138.1 (C(4)), 142.1, 142.3, 144.8 (*i-Ph*); m/z (ESI⁺) 453 ([M + H]⁺, 100%); HRMS (ESI⁺) m/z : [M + H]⁺ Calcd for C₃₂H₄₁N₂⁺ 453.3264; Found 453.3259.

(2*R*,2'*R*, α *S*, α' *R*,*Z*)-1-[N(1')-(α -Methylbenzyl)pyrrolidin-2'-yl]-2-[N-benzyl-N-(α' -methylbenzyl)amino]hept-4-ene 52. DIBAL-H (1.0 M in PhMe, 1.0 mL, 1.0 mmol) was added dropwise to a stirred solution of **43** (242 mg, 0.459 mmol, >99:1 dr) in PhMe (5 mL) at –78 °C and the resultant mixture was stirred at –78 °C for 1 h. MeOH (93 μ L, 2.3 mmol) and Ph₃P=CHCH₂CH₃ (0.31 M suspension in PhMe/THF, 14.8 mL, 4.59 mmol)²⁵ were added sequentially, and the resultant mixture was allowed to warm to rt and stirred at rt for 16 h, then concentrated in vacuo. The residue was dissolved in Et₂O (40 mL) and the resultant solution was filtered through a short plug of Celite[®] (eluent Et₂O) and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Et₂O/NH₄OH, 100:20:1) gave **52** as a pale yellow oil (106 mg, 48%, >95:5 dr [(*Z*):(*E*)]); $[\alpha]_{\text{D}}^{25}$ +53.9 (*c* 1.0 in CHCl₃); v_{max} (ATR) 3025, 2964, 2932, 2872 (C–H), 1602, 1493, 1453 (C=C); δ_{H} (400 MHz, CDCl₃) 0.86 (3*H*, t, *J* 7.5, C(7)*H*₃), 1.32 (3*H*, d, *J* 6.8, C(α)*Me*), 1.41–1.47 (1*H*, m, C(1)*H*_A), 1.44 (3*H*, d, *J* 6.8, C(α')*Me*), 1.53–1.77 (4*H*, m, C(1)*H*_B, C(3')*H*_A, C(4')*H*₂), 1.79–1.92 (4*H*, m, C(3)*H*_A, C(6)*H*₂, C(3')*H*_B), 1.93–2.03 (1*H*, m, C(3)*H*_B), 2.36–2.45 (2*H*, m, C(2')*H*, C(4')*H*_A), 2.58–2.66 (1*H*, m, C(2)*H*), 2.84 (1*H*, ddd, *J* 9.1, 7.8, 2.4, C(4')*H*_B), 3.73 (1*H*, d, *J* 14.8, NCH_AH_BPh), 3.77 (1*H*, d, *J* 14.8, NCH_AH_BPh), 3.85 (1*H*, q, *J* 6.8, C(α')*H*), 3.99 (1*H*, q, *J* 6.8, C(α)*H*), 5.09 (1*H*, app dtt, *J* 10.8, 6.8, 1.4, C(4)*H*), 5.23 (1*H*, app dtt, *J* 10.8, 7.1, 1.6, C(5)*H*), 7.15–7.41 (15*H*, m, *Ph*); δ_{C} (100 MHz, CDCl₃) 14.2 (C(7)), 18.0 (C(α)*Me*), 20.9 (C(6)), 21.9 (C(α')*Me*), 22.4 (C(4')), 29.7 (C(3)), 30.6 (C(1)), 37.4 (C(3')), 48.7 (C(5')), 50.1 (NCH₂Ph), 55.9 (C(2)), 57.0 (C(α)), 58.6 (C(2')), 59.2 (C(α')), 126.6, 126.6, 127.0 (*p-Ph*), 128.0, 128.1 (*o,m-Ph*), 128.2 (C(4)), 128.2, 128.2, 128.3, 128.6 (*o,m-Ph*), 131.7 (C(5)), 142.2, 142.3, 145.0 (*i-Ph*); m/z (ESI⁺) 481 ([M + H]⁺, 100%); HRMS (ESI⁺) m/z : [M + H]⁺ Calcd for C₃₄H₄₅N₂⁺ 481.3577; Found 481.3571.

(5*R*,6*aR*,10*aS*)-5-Propyldecahydrodipyrrolo[1,2-*a*:1',2'-*c*]pyrimidine [(+)-tetraponerine 1] 55. Palladium black (29 mg, 20% w/w) was added to a stirred, degassed solution of **51** (145 mg, 0.320 mmol, >99:1 dr) in MeOH/H₂O/AcOH (10:1:1, 2 mL). The resultant mixture was stirred under an atmosphere of H₂ (5 atm) at rt for 48 h. The reaction mixture was then filtered through a short plug of Celite[®] (eluent MeOH/H₂O/AcOH, 10:1:1) and concentrated in vacuo. The residue was dissolved in CH₂Cl₂ (2 mL), then K₂CO₃ (133 mg, 0.96 mmol) and 4-bromobutanal (96 mg, 0.64 mmol) were added to the resultant solution.

The reaction mixture was stirred at rt for 2 h, then H₂O (2 mL) and satd aq NaHCO₃ (10 mL) were added. The reaction mixture was then extracted with CH₂Cl₂ (3 × 5 mL) and the combined organic extracts were dried and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Me₂CO/NH₄OH, 100:10:1) gave (+)-tetraponerine 1 (**55**) as a colourless oil (10 mg, 15% from **51**, >99:1 dr); [α]_D²⁵ +14.4 (*c* 0.13 in CHCl₃); {lit.⁵ [α]_D²⁰ +11 (*c* 0.14 in CHCl₃); lit.⁶ [α]_D²⁰ +14 (*c* 0.498 in CHCl₃)}; $\nu_{\max}$ (ATR) 2927, 2801 (C–H);²⁶ δ_{H} (400 MHz, C₆D₆) 0.96 (3H, t, *J* 7.3, C(3')H₃), 1.23–1.45 (6H, m, C(1)H_A, C(6)H_A, C(7)H_A, C(8)H_A, C(1')H_A, C(2')H_A), 1.49–1.69 (3H, m, C(7)H_B, C(8)H_B, C(2')H_B), 1.69–1.91 (6H, m, C(1)H_B, C(2)H₂, C(6)H_B, C(9)H_A, C(1')H_B), 1.91–2.02 (1H, m, C(6a)H), 2.82–2.91 (3H, m, C(3)H_A, C(5)H, C(9)H_B), 3.21 (1H, app q, *J* 7.7, C(3)H_B), 3.48, (1H, app t, *J* 2.3, C(10a)H); δ_{H} (400 MHz, CDCl₃) 0.90 (3H, t, *J* 7.3, C(3')H₃), 1.24–1.43 (3H, m, C(7)H_A, C(2')H₂), 1.43–1.96 (12H, m, C(1)H₂, C(2)H₂, C(6)H₂, C(7)H_B, C(8)H₂, C(9)H_A, C(1')H₂), 2.00–2.10 (1H, m, C(6a)H), 2.82 (1H, td, *J* 7.9, 3.0, C(3)H_A), 2.91–3.01 (2H, m, C(5)H, C(9)H_B), 3.08 (1H, app q, *J* 7.9, C(3)H_B), 3.39 (1H, app d, *J* 3.7, C(10a)H); δ_{C} (100 MHz, C₆D₆) 14.5 (C(3')), 20.4 (C(8)), 20.8 (C(2')), 22.0 (C(2)), 30.1, 30.1 (C(1), C(6)), 31.6 (C(7)), 35.4 (C(1')), 50.0 (C(9)), 51.0 (C(3)), 53.7 (C(5)), 58.3 (C(6a)), 76.5 (C(10a)); δ_{C} (100 MHz, CDCl₃) 14.3 (C(3')), 20.1 (C(8)), 20.6 (C(2')), 21.3 (C(2)), 29.4 (C(6)), 29.7 (C(1)), 31.0 (C(7)), 34.0 (C(1')), 50.1 (C(9)), 50.3 (C(3)), 53.6 (C(5)), 58.3 (C(6a)), 76.9 (C(10a));²⁷ *m/z* (ESI⁺) 209 ([M + H]⁺, 100%); HRMS (ESI⁺) *m/z*: [M + H]⁺ Calcd for C₁₃H₂₅N₂⁺ 209.2012; Found 209.2013.

(5*R*,6*aR*,10*aS*)-5-Pentyldecahydrodipyrrolo[1,2-*a*:1',2'-*c*]pyrimidine [(+)-tetraponerine 5] **56.**

Palladium black (31 mg, 20% w/w) was added to a stirred, degassed solution of **52** (154 mg, 0.330 mmol) in MeOH/H₂O/AcOH (10:1:1, 2 mL). The resultant mixture was stirred under an atmosphere of H₂ (5 atm) at rt for 48 h. The reaction mixture was then filtered through a short plug of Celite[®] (eluent MeOH/H₂O/AcOH, 10:1:1) and concentrated in vacuo. The residue was dissolved in CH₂Cl₂ (2 mL), then K₂CO₃ (137 mg, 0.99 mmol) and 4-bromobutanal (99 mg, 0.66 mmol) were added to the resultant solution. The reaction mixture was stirred at rt for 2 h, then H₂O (2 mL) and satd aq NaHCO₃ (10 mL) were added. The reaction mixture was then extracted with CH₂Cl₂ (3 × 5 mL) and the combined organic extracts were dried and concentrated in vacuo. Purification via flash column chromatography (eluent 30–40 °C petrol/Me₂CO/NH₄OH, 100:10:1) gave (+)-tetraponerine 5 (**56**) as a colourless oil (11 mg, 14% from **52**, >99:1 dr); [α]_D²⁵ +12.4 (*c* 0.13 in CHCl₃); {lit.⁴ for a sample isolated from the natural source [α]_D²⁰ +10 (*c* 0.2 in CHCl₃); lit.⁵ [α]_D²⁰ +10 (*c* 0.24 in CHCl₃); lit.⁶ [α]_D²⁰ +14 (*c* 1.6 in CHCl₃)}; $\nu_{\max}$ (ATR) 2980, 2930, 2794 (C–H);²⁶ δ_{H} (400 MHz, C₆D₆) 0.93 (3H, t, *J* 6.8, C(5')H₃), 1.28–1.46 (10H, m, C(1)H_A, C(6)H_A, C(7)H_A, C(8)H_A, C(1')H_A, C(2')H_A, C(3')H₂, C(4')H₂), 1.50–1.69 (3H, m, C(7)H_B, C(8)H_B, C(2')H_B), 1.69–1.92 (6H, m, C(1)H_B, C(2)H₂,

C(6)*H*_B, C(9)*H*_A, C(1')*H*_B), 1.94–2.01 (1H, m, C(6a)*H*), 2.83–2.92 (3H, m, C(3)*H*_A, C(5)*H*, C(9)*H*_B), 3.23 (1H, app q, *J* 7.8, C(3)*H*_B), 3.51 (1H, app t, *J* 2.1, C(10a)*H*); δ_H (400 MHz, CDCl₃) 0.88 (3H, t, *J* 6.8, C(5')*H*₃), 1.22–1.42 (7H, m, C(7)*H*_A, C(2')*H*₂, C(3')*H*₂, C(4')*H*₂), 1.43–1.96 (12H, m, C(1)*H*₂, C(2)*H*₂, C(6)*H*₂, C(7)*H*_B, C(8)*H*₂, C(9)*H*_A, C(1')*H*₂), 2.01–2.10 (1H, m, C(6a)*H*), 2.82 (1H, app td, *J* 8.0, 2.8, C(3)*H*_A), 2.89–3.01 (2H, m, C(5)*H*, C(9)*H*_B), 3.08 (1H, app q, *J* 8.0, C(3)*H*_B), 3.39 (1H, app d, *J* 3.5, C(10a)*H*); δ_C (100 MHz, C₆D₆) 14.4 (C(5')), 20.4 (C(8)), 22.0 (C(2)), 23.2 (C(4')), 27.5 (C(2')), 30.1, 30.2 (C(1), C(6)), 31.6 (C(7)), 32.5 (C(3')), 33.2 (C(1')), 50.0 (C(9)), 51.1 (C(3)), 54.1 (C(5)), 58.3 (C(6a)), 76.5 (C(10a)); δ_C (100 MHz, CDCl₃) 14.3 (C(5')), 20.2 (C(8)), 21.3 (C(2)), 22.8 (C(4')), 27.2 (C(2')), 29.4 (C(6)), 29.7 (C(1)), 31.0 (C(7)), 31.7 (C(1')), 32.1 (C(3')), 50.0 (C(9)), 50.3 (C(3)), 53.9 (C(5)), 58.3 (C(6a)), 76.8 (C(10a));²⁷ *m/z* (ESI⁺) 237 ([M + H]⁺, 100%); HRMS (ESI⁺) *m/z*: [M + H]⁺ Calcd for C₁₅H₂₉N₂⁺ 237.2325; Found 237.2324.

ASSOCIATED CONTENT

Supporting Information Available: Copies of ¹H and ¹³C NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E-mail: steve.davies@chem.ox.ac.uk.

Notes

The authors declare no competing financial interest.

References and Notes

- ¹ Brakman, J. C.; Daloze, D.; Pasteels, J. M.; Van Hecke, P.; Declercq, J. M.; Stinwell, V.; Francke W. Z. *Naturforsch.* **1987**, 42c, 627.
- ² Kem, W. R.; Wildeboer, K.; LeFrancois, S.; Raja, M.; Marszalec, W.; Braekman, J.-C. *Cell. Mol. Neurobiol.* **2004**, 24, 535.
- ³ Yue, C.; Royer, J.; Husson, H.-P. *J. Org. Chem.* **1990**, 55, 1140.
- ⁴ Macours, P.; Braekman, J.-C.; Daloze, D. *Tetrahedron* **1995**, 51, 1415.
- ⁵ Yue, C.; Gauthier, I.; Royer, J.; Husson, H.-P. *J. Org. Chem.* **1996**, 61, 4949.

- ⁶ Bosque, I.; Gonzalez-Gomez, J. C.; Loza, M. I.; Brea, J. *J. Org. Chem.* **2014**, *79*, 3982.
- ⁷ Merlin, P.; Braekman, J.-C.; Daloze, D.; Pasteels, J. M. *J. Chem. Ecol.* **1988**, *14*, 517.
- ⁸ (a) Merlin, P.; Braekman, J.-C.; Daloze, D. *Tetrahedron Lett.* **1988**, *29*, 1691. (b) Jones, T. H. *Tetrahedron Lett.* **1990**, *31*, 1535. (c) Jones, T. H. *Tetrahedron Lett.* **1990**, *31*, 4543. (d) Merlin, P.; Braekman, J.-C.; Daloze, D. *Tetrahedron* **1991**, *47*, 3805. (e) Barluenga, J.; Tomás, M.; Kouznetsov, V.; Rubio, E. *J. Org. Chem.* **1994**, *59*, 3699. (f) Devijver, C.; Macours, P.; Braekman, J.-C.; Daloze, D.; Pasteels, J. M. *Tetrahedron* **1995**, *51*, 10913. (g) Kim, J. T.; Gevorgyan, V. *Org. Lett.* **2002**, *4*, 4697. (h) Kim, J. T.; Butt, J.; Gevorgyan, V. *J. Org. Chem.* **2004**, *69*, 5638. (i) Charette, A. B.; Mathieu, S.; Martel, J. *Org. Lett.* **2005**, *7*, 5401.
- ⁹ (a) Takahata, H.; Kubota, M.; Ikota, N. *J. Org. Chem.* **1999**, *64*, 8594. (b) Stragies, R.; Blechert, S. *J. Am. Chem. Soc.* **2000**, *122*, 9584. (c) Airiau, E.; Girard, N.; Pizzeti, M.; Salvadori, J.; Taddei, M.; Mann, A. *J. Org. Chem.* **2010**, *75*, 8670. (d) Bosque, I.; González-Gómez, J. C.; Guijarro, A.; Foubelo, F.; Yus, M. *J. Org. Chem.* **2012**, *77*, 10340. (e) Strassnig, I.; Körber, K.; Hüniger, U.; Kunz, H. *Synthesis* **2015**, *47*, 2299. (f) Pacheco, J. C. O.; Lipp, A.; Nauth, A. M.; Acke, F.; Dietz, J.-P.; Opatz, T. *Chem. Eur. J.* **2016**, *22*, 5409.
- ¹⁰ Several formal syntheses of the tetraponerine alkaloids have also been reported; see: (a) Plehiers, M.; Heilporn, S.; Ekelmans, D.; Leclercq, S.; Sangermano, M.; Braekman, J.-C.; Daloze, D. *Can. J. Chem.* **2000**, *78*, 1030. (b) Bisai, A.; Singh, V. K. *Tetrahedron Lett.* **2007**, *48*, 1907. (c) Liu, J.-D.; Chen, Y.-C.; Zhang, G.-B.; Li, Z.-Q.; Chen, P.; Du, J.-Y.; Tu, Y.-Q.; Fan, C.-A. *Adv. Synth. Catal.* **2011**, *353*, 2721.
- ¹¹ Various structural analogues of the tetraponerine alkaloids have also been prepared; see: (a) Zhu, Z.; Quirion, J.-C.; Husson, H.-P. *Tetrahedron Lett.* **1989**, *30*, 5137. (b) Rouchaud, A.; Braekman, J.-C. *Eur. J. Org. Chem.* **2009**, 2666. See also: Ref 8h.
- ¹² For a recent review of this methodology, see: Davies, S. G.; Fletcher, A. M.; Roberts, P. M.; Thomson, J. E. *Tetrahedron: Asymmetry* **2012**, *23*, 1111.
- ¹³ Davies, S. G.; Fletcher, A. M.; Roberts, P. M.; Smith, A. D. *Tetrahedron* **2008**, *65*, 10192.
- ¹⁴ Csatayová, K.; Davies, S. G.; Fletcher, A. M.; Ford, J. G.; Klauber, D. J.; Roberts, P. M.; Thomson, J. E. *J. Org. Chem.* **2014**, *79*, 10932.
- ¹⁵ (a) Davies, S. G.; Iwamoto, K.; Smethurst, C. A. P.; Smith, A. D.; Rodríguez-Solla, H. *Synlett* **2002**, 1146. (b) Chippindale, A. M.; Davies, S. G.; Iwamoto, K.; Parkin, R. M.; Smethurst, C. A. P.; Smith, A. D.; Rodríguez-Solla, H. *Tetrahedron* **2003**, *59*, 3253. (c) Csatayová, K.; Davies, S. G.; Figuccia, A. L. A.; Fletcher, A. M.; Ford, J. G.; Lee, J. A.; Roberts, P. M.; Seward, B. G.; Song, H.; Thomson, J. E. *Tetrahedron* **2015**, *71*, 9131.

- ¹⁶ *tert*-Butyl sorbate **3** was prepared in 91% yield upon esterification of commercially available sorbic acid with isobutene; see: Ref 14.
- ¹⁷ For example, see: (a) Davies, S. G.; Durbin, M. J.; Goddard, E. C.; Kelly, P. M.; Kurosawa, W.; Lee, J. A.; Nicholson, R. L.; Price, P. D.; Roberts, P. M.; Russell, A. J.; Scott, P. M.; Smith, A. D. *Org. Biomol. Chem.* **2009**, 7, 761. (b) Davies, S. G.; Lee, J. A.; Roberts, P. M.; Thomson, J. E.; Yin, J. *Tetrahedron* **2011**, 67, 6382.
- ¹⁸ For example, see: (a) Magano, J.; Bowles, D.; Conway, B.; Nanninga, T. N.; Winkle, D. D. *Tetrahedron Lett.* **2009**, 50, 6325. (b) Davies, S. G.; Ichihara, O.; Roberts, P. M.; Thomson, J. E. *Tetrahedron* **2011**, 67, 216. (c) Davies, S. G.; Lee, J. A.; Roberts, P. M.; Stonehouse, J. P.; Thomson, J. E. *J. Org. Chem.* **2012**, 77, 9724.
- ¹⁹ The application of an alternative olefination procedure gave a 75:25 dr [(*Z*):(*E*)] mixture of olefins, which was analysed by ¹³C NMR spectroscopy. The configuration of the double bond within (*Z*)-**24** was then assigned from the diagnostic differences in chemical shift for the allylic carbon atoms within the two diastereoisomeric olefin products: for the (*E*)-diastereoisomer $\delta_{\text{C}} = 35.4$ (C(3)), 25.8 (C(6)) and for the (*Z*)-diastereoisomer $\delta_{\text{C}} = 29.6$ (C(3)), 20.9 (C(6)). (*Z*)-Olefins typically appear ~6 ppm lower in their ¹³C NMR spectra than the corresponding (*E*)-diastereoisomers; for a discussion of this “ γ -steric effect” in the ¹³C NMR spectra of *cis*- and *trans*-olefins, see: Jacobsen, N. E. In *NMR Data Interpretation Explained: Understanding 1D and 2D NMR Spectra of Organic Compounds and Natural Products*; Wiley, Hoboken, New Jersey, USA, 2017; pp 212.
- ²⁰ The (*Z*)-configurations of the C=C double bonds within **30**, **46** and **52** were assigned by analogy to that within (*Z*)-**24** in the absence of authentic samples of the corresponding (*E*)-diastereoisomers.
- ²¹ Oxidation of *ent*-**36** to give the corresponding pyrrole was previously noted upon attempted Pd-mediated tandem hydrogenation/hydrogenolysis; see: Ref 15a.
- ²² As (+)-tetraonerine 2 (**49**) and (+)-tetraonerine 6 (**50**) were derived from the common precursor (*R,R,R,R*)-**42**, the configurations of **46**, **48** and **50** remain secure.
- ²³ Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometallics* **1996**, 15, 1518.
- ²⁴ The solution of Ph₃P=CH₂ was prepared on the requisite scale by the addition of KHMDS (0.5 M in PhMe, 1.0 equiv) to stirred suspension of methyltriphenylphosphonium iodide (1.0 M in THF, 1.0 equiv) at rt and the resultant suspension was stirred at rt for 1 h.

²⁵ The solution of $\text{Ph}_3\text{P}=\text{CHCH}_2\text{CH}_3$ was prepared on the requisite scale by the addition of KHMDS (0.5 M in PhMe, 1.0 equiv) to stirred suspension of propyltriphenylphosponium bromide (1.0 M in THF, 1.0 equiv) at rt and the resultant suspension was stirred at rt for 1 h.

²⁶ The Bolmann band frequencies and intensities in the IR spectra of the tetraponerine alkaloids have previously been used to assign the configuration at the C(5)-position. The even numbered alkaloids (i.e., T2, T4, T6 and T8) typically have more intense Bohlmann bands at $\sim 2790\text{ cm}^{-1}$, whereas the odd numbered alkaloids (i.e., T1, T3, T5 and T7) display weaker Bohlmann bands at $\sim 2805\text{ cm}^{-1}$; for further discussion of this analysis, see: Garraffo, H. M.; Spande, T. F.; Jain, P.; Kaneko, T.; Jones, T. H.; Blum, M. S.; Ali, T. M. M.; Snelling, R. R.; Isbell, L. A.; Robertson, H. G.; Daly, J. W. *Rapid Commun. Mass Spectrom.* **2001**, *15*, 1409.

²⁷ For both (+)-tetraponerine 1 (**55**) and (+)-tetraponerine 5 (**56**), the resonance corresponding to C(10a) in their ^{13}C NMR spectra (in CDCl_3) was obscured by the solvent peak.