

## **Abstract**

**Objective:** To examine job stress and health behaviours, including their co-occurrence, in Australians aged 31–41 year assessed in 2009-11.

**Methods:** Cross-sectional analyses using multivariable regression models of the association between the Effort Reward Imbalance (ERI) scale and health behaviours (smoking, alcohol consumption, diet, physical activity and BMI) both individually and co-occurring (0-3 versus 4-5 behaviours) were undertaken. Covariates included sociodemographics, personality and life events.

**Results:** Greater ERI was associated with a significantly lower prevalence of having co-occurring healthy behaviours and poorer diets in both sexes. Higher ERI was also associated greater physical inactivity and sedentary behaviour in men and smoking, high alcohol consumption and more pedometer-measured physical activity in women.

**Conclusion:** Job stress at work was associated with a range of unhealthy behaviours, which may explain the higher chronic disease associated with job stress.

**Key terms** effort-reward imbalance, job stress, health behaviours; cross-sectional

## Introduction

The adverse effects of job stress on physical and mental diseases have been well investigated, but the role of health risk behaviours in these associations remains unclear. There is evidence that health risk behaviours account for a large proportion of the adverse effect of job stress on some cardiovascular diseases like coronary artery disease.<sup>1</sup> As such, health risk behaviours may be a target for reducing the effects of job stress on health.

Health risk behaviours differ according to levels of job stress but existing studies have some limitations. Authors for a recent individual participant data meta-analysis that included over 100,000 people<sup>2</sup> reported that people with higher levels of job stress were more likely to be current smokers, consume heavy amounts of alcohol and be less physically active than people with lower levels of job stress,<sup>3</sup> with similar findings reported by others.<sup>4,5</sup> There has been little investigation of the association between diet and job stress, despite its importance for physical<sup>6-8</sup> and mental<sup>9</sup> health. Most studies of health risk behaviours and job stress have been conducted in Europe, especially in Finland,<sup>3-5</sup> with very little is known outside this region. Contextual differences in the distributions of risk factors, characteristics of the workforce and society could result in different associations between these factors, suggesting that region specific analyses are warranted.

Researchers have tended to examine the association between individual risk factors and job stress. While this is important for understanding potential mechanisms, it disregards that health behaviours commonly co-occur. When they have examined the association between job stress and co-occurring risk behaviours the results have been inconsistent.<sup>3,4,5</sup> In the IPD-Work Consortium, investigators found similar associations between job stress and individual or co-occurring health risk behaviours.<sup>3</sup> In contrast, in the Finnish Public Sector Study, job stress was more strongly associated with co-occurring than individual health risk behaviours.

<sup>4 5</sup>. Many studies have lacked information on potentially important factors that might confound associations between job stress and health risk behaviours including major life events (e.g., divorce or illness) <sup>10 11</sup> and personality, <sup>12 13</sup> as noted by others. <sup>3-5</sup> A better understanding of the extent to which job stress is associated with both individual and co-occurring health behaviours with account for these important confounding factors is important for understanding its links with chronic disease but also for devising interventions to improve workers' health.

The aims of this study were to examine the association between job stress and a suite of health risk behaviours, including their co-occurrence, with adjustment for a wide range of potential confounding factors.

## **Method**

### **Sample**

The Childhood Determinants of Adult Health (CDAH) study is a cohort study that follows up the participants of the 1985 Australian Schools Health and Fitness Survey (ASHFS). The details of sampling and data collection is described elsewhere. <sup>14</sup> This study uses data from the second adult follow-up of ASHFS participants conducted in 2009-11 (CDAH-2) that comprised questionnaire assessments of health risk behaviours and job stress, as specified below. As specified in the participation flow chart (Figure 1), after excluding participants with incomplete job stress information, 1,481 participants were eligible for inclusion although the number in some analyses is less than this due to missing data for some health risk behaviours and covariates.

## **Measures**

### **Job stress**

Job stress in this study was measured with the ERI ratio, using a validated questionnaire that included 17 items (6 effort items and 11 reward items).<sup>15</sup> The effort scale covers workplace interruptions, job responsibility, pressure to work overtime and job demand, while the reward scale covers adequate support, being treated unfairly and adequate gains and reflects employee's responses of esteem, job security and job promotion. Standard scoring was applied to devise scores for effort, reward and the ERI ratio and higher scores indicate higher job stress.

### ***Health risk behaviours***

#### **Smoking**

Smoking status was measured by standard self-reported questions regarding current and past smoking history with people defined as 'never smoker' (smoked <100 cigarettes in lifetime), 'ex-smoker' (current non-smoker that was a daily smoker) or 'current smoker' (current weekly or daily smoker).

#### **Body mass index (BMI)**

BMI was calculated from self-reported height and weight. A correction factor was applied based on self-report and measured height and weight from an earlier follow-up to give estimates of measured BMI from self-reported values, as previously.<sup>16</sup> BMI was reported into healthy weight (BMI <25 kg/m<sup>2</sup>), overweight (25-29.9 kg/m<sup>2</sup>) and obese ( $\geq$  30 kg/m<sup>2</sup>).

#### **Diet**

Participants completed a 127 item food frequency and habits questionnaire. The Dietary Guideline Index (DGI) used for assessing participants' diet quality, based on the Dietary Guidelines for Australian Adults<sup>17</sup> and the Australian Guide to Healthy Eating (AGHE).<sup>18</sup>

The DGI reflects the consumption of the 5 core food groups (vegetables, fruits, cereals, meat and alternatives, and dairy) and includes 15 components including whole-grain cereals, fruit and vegetables, lean meat, reduced or low fat dairy, and dietary variety.<sup>19</sup> The DGI score with a potential range of 0-150, a higher score indicates greater compliance with the Dietary Guidelines. This score has been shown to be a valid measure of diet quality<sup>19</sup> and was associated with cardio-metabolic risk factors in Australian adults.<sup>20</sup> We also examined consumption of ‘extra foods’ that are not essential to provide nutrition and typically contain too much fat, sugar and salt.<sup>19 21</sup> Frequency of takeaway food consumption was measured from participants’ response to the question, ‘how many times per week would you usually eat hot takeaway meals?’ with responses ranging from ‘I don’t eat takeaway’ to ‘6–7 meals per week’ and presented as dichotomous variable (‘less than twice per week’ or ‘twice a week or more’).<sup>8</sup>

#### Alcohol consumption

We used two items for measuring alcohol consumption. One measured standard drinks consumed per day from responses to the question: ‘On a day, when you drink alcohol, how many standard drinks do you usually have?’ with responses being ‘none’, ‘1 or 2 drinks per day’, ‘3 or 4 drinks per day’ or ‘more than 5 drinks per day’. The other estimated the total volume of alcohol (grams/day) consumed in the previous 12 months from the usual frequency of consumption of 10 common types of alcoholic beverages multiplied by the average alcohol concentration of each beverage from the Food Frequency Questionnaire.<sup>22</sup>

#### Physical activity

The long version of the International Physical Activity Questionnaire (IPAQ-L) was used to assess levels of physical activity. Participants self-reported duration and frequency of leisure time (LTPA), domestic and gardening (DGPA), work-related and transport-related physical

activity. We summed the total minutes of LTPA, work related physical activity, DGPA and transport-related physical activity were by multiplying frequency by duration, and then multiplied by the resting metabolic rate to get the MET-minutes/week. Additionally, we summed the total sitting time on weekdays and weekends, respectively. Steps per day were also measured using a Yamax Dighwalker SW-200 pedometer worn for seven consecutive days, as described elsewhere.<sup>23</sup>

### Co-occurrence of health risk behaviours

We calculated a Healthy Lifestyle Score that is associated with biomedical cardiovascular risk factors<sup>8</sup> and mental health in this cohort<sup>24</sup> and is similar to other scores.<sup>25 26</sup> Our score comprised five 'healthy' items assigned one point each: body mass index (BMI)  $<25\text{kg/m}^2$ , never smoker or ex-smoker  $\geq 12$  months,  $\geq 3$  hours of moderate to vigorous leisure time physical activity (LTPA) per week,  $\leq 20$  grams of alcohol per day and for indicating a 'healthy' diet, scoring in the 75<sup>th</sup> percentile of a validated Dietary Guideline Index that assessed adherence to Australian dietary guidelines from a Food Frequency Questionnaire.<sup>27</sup> The total score ranged from zero (no healthy behaviours) to five (all healthy behaviours).

### Covariates

Sociodemographic covariates were age, marital status (not married/de facto, currently married/de facto or divorced/widowed/separated), parental status (children or no children), highest level of education attained (secondary schooling only, vocational training or tertiary degree). Work covariates were occupation type (manual, non-manual or professional/managerial), working hours and work schedule (regular daytime, shift work or other). Individual covariates were the number of adverse life events in the previous five years (e.g. financial difficulties or physical illness) reported by participants,<sup>28</sup> personality assessed by the 60-item NEO Personality Inventory that measures neuroticism, extroversion,

agreeableness, openness and conscientiousness,<sup>29</sup> DSM-IV diagnoses of major depression/dysthymia or anxiety from the lifetime version of the Computerised International Diagnostic Interview administered over the telephone<sup>30</sup>.

## **Data analysis**

We used sex-specific log multinomial regression (for variables with three or more categories), log binomial regression (for dichotomous variables) and linear regression (for continuous variables) to estimate associations between the total ERI scale as well as the individual effort and reward scales and the health risk behaviours. We multiplied the ERI ratio by 10 (corresponding to a 0.1 unit change on the ERI scale), and divided effort and reward by 5 (corresponding to a 5 unit change in these scales) to aid interpretation.

Extra food consumption, domains of physical activity and total alcohol consumption were log transformed. For domains of physical activity there were a large number of zeros, so two step analyses were used. First, a binary variable was created for doing any activity versus none with log binomial regression used to investigate associations with ERI. Then, linear regression was performed on the amount of activity within the any activity group. In addition to looking at the HLS as a continuous variable, we made categories of low (scores 0 to 2), and high (scores 3 to 5). For daily steps, we categorised people into sedentary (<5000 steps/day), low active (5000-7499 steps/day), somewhat active (7500-9999 steps/day), active ( $\geq$  10000 steps/day) and high active (>12500 steps/day).<sup>31</sup>

We present models that are adjusted for sociodemographic factors, sociodemographic plus work factors; and age, work factors plus individual factors. Purposeful model building was used to select the specific covariates included in each model with factors that satisfied these criteria for any outcome included across all analyses. Previous studies of job stress and health

have tended to include all work and SEP variables in a single model. However, there is a risk that such models are over-adjusted<sup>5</sup> so we did sensitivity analyses testing the effect of each SEP and work characteristic separately.

We used multiple imputation with chained equations and with 30 estimations to impute missing data on covariates. The following variables were used in the imputation model: childhood school academic attainment and smoking status, education level, marital status, sex, age, and state, height, weight and self-rated health from an earlier adult follow-up.

All analyses were conducted with Stata version 12.1 (Statacorp, 2012).

### **Ethical considerations**

The CDAH study was approved by Tasmanian Health and Medical Human Research Ethics Committee and all participants provided written informed consent.

### **Results**

The included participants are shown in Figure 1 and their characteristics in Table 1.

In men, after adjusting for sociodemographic factors, higher ERI was significantly associated with a lower mean Healthy Lifestyle Score, lower DGI, greater consumption of extra foods, less MET minutes of leisure-time physical activity per week and more minutes of sitting during the weekend (Table 2). After additionally adjusting for work-related and individual factors of personality and life events, the associations persisted between ERI and greater extra food consumption, less MET minutes of leisure-time physical activity per week, more minutes of sitting time during the weekend. In women, after adjustment for sociodemographic factors, higher ERI was associated with lower mean Healthy Lifestyle Scores and higher pedometer steps counts (Table 3). The magnitude of these associations was



similar after further adjustment for work and individual covariates. Associations between weekday sitting time and ERI were completely attenuated by adjusted for work and individual factors.

In men, after adjusting for sociodemographic factors, higher ERI was significantly associated with a lower prevalence of ‘high’ Healthy Lifestyle Scores, which persisted after additionally adjusting for work-related and individual factors of personality and life events (Table 4). In women, after adjustment for sociodemographic factors, higher ERI was associated with a lower prevalence of a ‘high’ Healthy Lifestyle Score, a higher prevalence of ‘high active’ number of steps/day, a higher probability of doing any physical activity in the workplace; being a current smoker, consuming takeaway food twice a week or more and consuming five or more alcohol drinks per day (Table 5). The magnitude of these associations was similar after further adjustment for work and individual covariates but associations for alcohol consumption and any physical activity in the workplace were no longer significant.

Higher levels of effort in men were associated with greater consumption of extra foods and more minutes of sitting during the weekend across all models (Table 1, Supplementary Digital Content). Higher effort was also associated with lower DGI in men when adjusted for age, but not after adjusting for work or individual factors (Table 1, Supplementary Digital Content). In women, higher levels of effort were associated with a lower mean Healthy Lifestyle Score, doing more minutes of physical activity in the workplace, and less minutes of transport related physical activity even after adjusting for age, work and individual factors (Table 1, Supplementary Digital Content). In women, higher levels of effort were also associated with a lower prevalence of a high Healthy Lifestyle Score, a higher prevalence of current smoking and consumption of takeaway foods twice a week or more even after adjusting for age, work and individual factors (Table 2, Supplementary Digital Content).

Higher levels of reward in men were associated with higher scores on the dietary guideline index, less consumption of extra foods and less time sitting on the weekends independent of sociodemographic factors (Table 3, Supplementary Digital Content). In women, there was a trend toward a significant association between higher reward and lower body mass index but this was attenuated with adjustment for individual factors (Table 3, Supplementary Digital Content). Higher levels of reward in men were associated with a trend toward a greater prevalence of a high healthy lifestyle score and a lower likelihood of being a current smoker but did not reach statistical significance in the fully adjusted models (Table 4, Supplementary Digital Content). In women, higher levels of reward were associated with a lesser likelihood of being a current smoker and consuming takeaway food more than twice per week across all models (Table 4, Supplementary Digital Content).

Sensitivity analyses looking at the effect of individual markers of SEP and work characteristics showed that working hours were particularly influential on the associations between job stress and behavioural risk factors, particularly physical activity, in women (Tables 5 to 10, Supplementary Digital Content).

## **Discussion**

Higher job stress was associated with a range of health behaviours, including less adherence to dietary guidelines, lower levels of leisure time physical activity, more sitting time during weekends, being a current smoker and consuming greater amounts of alcohol. There was also evidence that higher job stress was associated with co-occurring health behaviours. These effects were largely independent of confounding factors, including life events and personality that have not been examined in previous studies. These findings contribute to our understanding of how job stress might lead to poorer health outcomes and suggest targets to improve the health and wellbeing of workers.

Higher levels of job stress were associated with a greater co-occurrence of health behaviours. The score we used to measure co-occurring health behaviours not only included behaviours previously examined in relation to job stress (e.g. smoking, alcohol, overweight and physical inactivity) but also items about diet which have not been considered by other investigators.<sup>4 5</sup> In support of our finding, Heikkila and colleagues reported that people with high job strain were 25% more likely to have four unhealthy behaviours than those without job strain in cross-sectional analyses including over 100,000 people.<sup>3</sup> Kouvonen et al. also found similar findings, with people with high ERI ratios being 40% more likely to have three or more unhealthy behaviours independent of sociodemographic factors.<sup>5</sup> Both authors noted they could not account for other important factors, such as personality. Our results confirm that associations between job stress and co-occurring health behaviours remain even after accounting for personality factors.

Job stress may be associated with a greater burden of unhealthy behaviours as people may engage in these behaviours to relieve stress related to work.<sup>4</sup> According to the self-medication theory, people engage in unhealthy behaviours to relieve the unfavourable symptoms associated with stress. For example, they might use tobacco, consume 'comfort foods' or drink alcohol to diminish stress from work.<sup>32-36</sup> While engaging in these behaviours might make people feel less stress at the time, they are also associated with higher risk of chronic diseases<sup>37 38</sup> and likely explain at least some of the link between job stress and such diseases, as noted by others.<sup>1</sup> Ideally, we would be able to address the causes of job stress to alleviate its effects on health behaviours and outcomes. However, the causes of job stress are complex and influenced by individual and workplace characteristics.<sup>39</sup> There is a potential role for workplace health promotion programs,<sup>40</sup> with there being some evidence that sustained, well designed and implemented programs have benefits on health behaviours<sup>41</sup> and some aspects of job stress.<sup>42</sup>

Higher job stress was also associated with several individual health behaviours. People with higher levels of job stress had a range of poorer dietary behaviours including consuming more extra foods each day, meeting less of the dietary guidelines (in men) and eating more takeaway foods (in women). There are no comparable studies of job stress and diet but studies of general stress support our findings. The investigators for the longitudinal Midlife in the US II study found evidence for an association between higher stress and greater consumption of energy-dense foods,<sup>43</sup> which include takeaway and ‘extra foods’ measured in our study. In experimental studies, craving for fat, sugar and energy increased during stressful situations and parts of the brain that control mood, like the meso-limbic dopaminergic brain “reward” system, are activated by eating these types of food.<sup>34</sup> Therefore, people with higher job stress might eat more of these types of food as it reduces their negative emotional responses to stress at work. An alternative hypothesis is that these foods are readily available with this of benefit to ‘time poor’ workers.<sup>44 45</sup> In support of this, women with higher ERI ratios had a higher probability of eating takeaway food twice or more times per week. Long working hours are associated with greater job stress<sup>45</sup> but adjusting for working hours in our models did not change these associations. Therefore, the association between poorer dietary behaviours and job stress are possibly related to the effects of these foods on mood regulation.

Higher ERI was related to lower levels of leisure-time physical activity and more time spent sitting during weekend, with these associations independent of SEP, work factors, personality and life events. The negative association between ERI and less leisure-time physical activity is consistent with findings from a previous study.<sup>46</sup> One mechanism might be that higher job stress is related to excessive fatigue manifesting as more sitting time during the weekends and lower levels of leisure-time physical activity.<sup>47</sup> For women, higher ERI was associated with doing more pedometer-measured steps per day, which is likely linked to the association between greater job stress and greater work-related physical activity. Related to this is the

apparent role of long working hours, with sensitivity analyses showing that work hours were particularly important for the association between ERI and physical activity. Long hours at work potentially increase job stress by increasing work-life conflicts<sup>48</sup> and others have noted the need to consider the role of long working hours in the association between job stress and health risk behaviours.<sup>2</sup> There are also possible bi-directional associations between work-related factors and health outcomes, with previous analyses in this cohort showing that people with greater levels of work-related physical activity had a higher risk of depression.<sup>49</sup>

We did not find any association between ERI and BMI in this study. Our results are in agreement with some previous studies that also found no association between job stress and BMI,<sup>50 51</sup> but in contrast to another that reported a U-shaped relationship between job stress and BMI.<sup>52</sup> One reason for the lack of an association in our study is that the levels of job stress were low with only 1.6% of the participants having a ERI ratio higher than the cut-off for high levels of job stress.<sup>53</sup> Similarly, although our cohort had reasonable distribution of different body sizes, most participants were healthy weight or overweight. There were less people that were underweight or obese, where the greatest associations with job stress have been demonstrated.<sup>52</sup> We note the longitudinal studies showing that people with higher job stress appear to gain more weight<sup>54</sup> and also that the preponderance of behaviours associated with weight gain in those with job stress in this study (e.g. consumption of energy dense foods and low levels of physical activity) may expose this group to weight gain overtime.

There are some limitations of this study. This study is cross-sectional and therefore we cannot draw any conclusions about causality. Our measures were mostly self-reported, with the exception of pedometers. This may contribute to misclassification of some health risk behaviours; however, all measures we used were validated. Although this study was a national study, the sample was not fully representative of the working population and this

may limit the generalizability of these findings. In line with this, the levels of jobs stress were low meaning and our findings might not be transferable to other contexts where there are higher levels of job stress.

The present study also has several strengths. Ours is the first study to examine the full ERI model in relation to co-occurrence of personal health risk behaviours and a range of dietary behaviours. We controlled for often missed confounders (reference to the paper that said people should account for these things), such as life events and personality. We concluded that while these were associated with both job stress and health risk behaviours, they did not appear to be confounders or mediators of the association between these two factors. Finally, the present study focuses on a national cohort of Australians, whereas most previous studies have been focused on European populations.

In conclusion, higher ERI was associated with poorer health risk behaviours independent of SEP, work factors, life events and personality. Our findings support those of others that suggest job stress may lead to poorer health outcomes via such health behaviours.

## Reference

1. Kivimaki M, Nyberg ST, Fransson EI et al. Associations of job strain and lifestyle risk factors with risk of coronary artery disease: a meta-analysis of individual participant data. *CMAJ* 2013;185:763-9.
2. Kivimaki M, Singh-Manoux A, Virtanen M, Ferrie JE, Batty GD, Rugulies R. IPD-Work consortium: pre-defined meta-analyses of individual-participant data strengthen evidence base for a link between psychosocial factors and health. *Scand J Work Environ Health* 2015;41:312-21.
3. Heikkila K, Fransson EI, Nyberg ST et al. Job strain and health-related lifestyle: findings from an individual-participant meta-analysis of 118,000 working adults. *Am J Public Health* 2013;103:2090-7.
4. Kouvonen A, Kivimaki M, Vaananen A et al. Job strain and adverse health behaviors: the Finnish Public Sector Study. *J Occup Environ Med* 2007;49:68-74.
5. Kouvonen A, Kivimaki M, Virtanen M et al. Effort-reward imbalance at work and the co-occurrence of lifestyle risk factors: cross-sectional survey in a sample of 36,127 public sector employees. *BMC Public Health* 2006;6:24.
6. Epel E, Lapidus R, McEwen B, Brownell K. Stress may add bite to appetite in women: a laboratory study of stress-induced cortisol and eating behavior. *Psychoneuroendocrinology* 2001;26:37-49.
7. Nishitani N, Sakakibara H. Relationship of obesity to job stress and eating behavior in male Japanese workers. *Int J Obes* 2006;30:528-33.
8. Smith KJ, McNaughton SA, Gall SL, Blizzard L, Dwyer T, Venn AJ. Takeaway food consumption and its associations with diet quality and abdominal obesity: a cross-sectional study of young adults. *Int J Behav Nutr Phys Act* 2009;6:29.

9. Smith KJ, Sanderson K, McNaughton SA, Gall SL, Dwyer T, Venn AJ. Longitudinal associations between fish consumption and depression in young adults. *Am J Epidemiol* 2014;179:1228-35.
10. Keyes KM, Hatzenbuehler ML, Hasin DS. Stressful life experiences, alcohol consumption, and alcohol use disorders: the epidemiologic evidence for four main types of stressors. *Psychopharmacology* 2011;218:1-17.
11. Bhagat RS. Effects of Stressful Life Events on Individual-Performance Effectiveness and Work Adjustment Processes within Organizational Settings - a Research Model. *Academy of Management Review* 1983;8:660-671.
12. Vollrath; M, Knoch; D, Cassano L. Personality, Risky Health Behaviour, and Perceived Susceptibility to Health Risks. *European Journal of Personality* 1999;13:39-50.
13. Tornroos M, Hintsanen M, Hintsala T et al. Personality traits of the five-factor model are associated with effort-reward imbalance at work: a population-based study. *Journal of Occupational and Environmental Medicine* 2012;54:875-80.
14. Dwyer T, Gibbons LE. The Australian Schools Health and Fitness Survey. Physical fitness related to blood pressure but not lipoproteins. *Circulation* 1994;89:1539-44.
15. Siegrist J, Starke D, Chandola T et al. The measurement of effort-reward imbalance at work: European comparisons. *Soc Sci Med* 2004;58:1483-99.
16. Venn AJ, Thomson RJ, Schmidt MD et al. Overweight and obesity from childhood to adulthood: a follow-up of participants in the 1985 Australian Schools Health and Fitness Survey. *Med J Aust* 2007;186:458-60.
17. Pomerleau J, Gilmore A, McKee M, Rose R, Haerpfer C. Determinants of smoking in eight countries of the former Soviet Union: results from the Living Condition, Lifestyles and Health Study. *Addiction* 2004;99:1577-1585.



18. Drazin D, Rosner J, Nuno M et al. Type of admission is associated with outcome of spontaneous subarachnoid hemorrhage. *Int J Stroke* 2013.
19. McNaughton SA, Ball K, Crawford D, Mishra GD. An index of diet and eating patterns is a valid measure of diet quality in an Australian population. *J Nutr* 2008;138:86-93.
20. McNaughton SA, Dunstan DW, Ball K, Shaw J, Crawford D. Dietary quality is associated with diabetes and cardio-metabolic risk factors. *J Nutr* 2009;139:734-42.
21. Smith KJ, Blizzard L, McNaughton SA, Gall SL, Dwyer T, Venn AJ. Daily eating frequency and cardiometabolic risk factors in young Australian adults: cross-sectional analyses. *Br J Nutr* 2012;108:1086-94.
22. Gall SL, Abbott-Chapman J, Patton GC, Dwyer T, Venn A. Intergenerational educational mobility is associated with cardiovascular disease risk behaviours in a cohort of young Australian adults: The Childhood Determinants of Adult Health (CDAH) Study. *BMC Public Health* 2010;10:55.
23. Cleland VJ, Schmidt MD, Salmon J, Dwyer T, Venn A. Correlates of pedometer-measured and self-reported physical activity among young Australian adults. *J Sci Med Sport* 2011;14:496-503.
24. Gall SL, Sanderson K, Smith KJ, Patton G, Dwyer T, Venn A. Bi-directional associations between healthy lifestyles and mood disorders in young adults: The Childhood Determinants of Adult Health Study. *Psychol Med* 2016;46:2535-48.
25. Lloyd-Jones DM, Hong Y, Labarthe D et al. Defining and setting national goals for cardiovascular health promotion and disease reduction: the American Heart Association's strategic Impact Goal through 2020 and beyond. *Circulation* 2010;121:586-613.
26. Khaw K-T, Wareham N, Bingham S, Welch A, Luben R, Day N. Combined Impact of Health Behaviours and Mortality in Men and Women: The EPIC-Norfolk Prospective Population Study. *PLoS Med* 2008;5:e12.

27. Sanjoti P, David K, Neville O, Konrad J. Spousal concordance and reliability of the 'Prudence Score' as a summary of diet and lifestyle. *Australian and New Zealand Journal of Public Health* 2009;33:320-324.
28. Brugha T, Bebbington P, Tennant C, Hurry J. The List of Threatening Experiences: a subset of 12 life event categories with considerable long-term contextual threat. *Psychol Med* 1985;15:189-94.
29. Paul T. Costa J, McCrae RR. Normal Personality Assessment in Clinical Practice: The NEO Personality Inventory. *Psychological Assessment* 1992;4:5-13.
30. Andrews G, Peters L. The psychometric properties of the Composite International Diagnostic Interview. *Soc Psychiatry Psychiatr Epidemiol* 1998;33:80-8.
31. Tudor-Locke C, Bassett DR, Jr. How many steps/day are enough? Preliminary pedometer indices for public health. *Sports Med* 2004;34:1-8.
32. Perdikaris P, Kletsiou E, Gymnopoulos E, Matziou V. The relationship between workplace, job stress and nurses' tobacco use: a review of the literature. *Int J Environ Res Public Health* 2010;7:2362-75.
33. Leeies M, Pagura J, Sareen J, Bolton JM. The use of alcohol and drugs to self-medicate symptoms of posttraumatic stress disorder. *Depress Anxiety* 2010;27:731-6.
34. Dallman MF, Pecoraro NC, la Fleur SE. Chronic stress and comfort foods: self-medication and abdominal obesity. *Brain Behav Immun* 2005;19:275-80.
35. Morris MJ, Beilharz JE, Maniam J, Reichelt AC, Westbrook RF. Why is obesity such a problem in the 21st century? The intersection of palatable food, cues and reward pathways, stress, and cognition. *Neurosci Biobehav Rev* 2014.
36. Vasse RM, Nijhuis FJ, Kok G. Associations between work stress, alcohol consumption and sickness absence. *Addiction* 1998;93:231-41.

37. Martin J O'Donnell, Denis Xavier, Lisheng Liu et al. Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): a case-control study. *Lancet* 2010;376:113-123.
38. Yusuf S, Hawken S, Ounpuu S et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004;364:937-52.
39. Michie S. Causes and management of stress at work. *Occupational and Environmental Medicine* 2002;59:67.
40. Kilpatrick M, Blizzard L, Sanderson K et al. Investigating Employee-Reported Benefits of Participation in a Comprehensive Australian Workplace Health Promotion Program. *J Occup Environ Med* 2016;58:505-13.
41. Rongen A, Robroek SJ, van Lenthe FJ, Burdorf A. Workplace health promotion: a meta-analysis of effectiveness. *Am J Prev Med* 2013;44:406-15.
42. Jarman L, Martin A, Venn A, Otahal P, Sanderson K. Does workplace health promotion contribute to job stress reduction? Three-year findings from Partnering Healthy@Work. *BMC Public Health* 2015;15:1293.
43. Tsenkova V, Boylan JM, Ryff C. Stress eating and health. Findings from MIDUS, a national study of US adults. *Appetite* 2013;69:151-5.
44. Alan Warde, Martens L. *Eating Out: Social Differentiation, Consumption and Pleasure*. United Kingdom: Cambridge University Press, 2000.
45. Welfare AIoHa. Australia's health 2014. In: Welfare AIoHa (ed.), Canberra: Australian Institute of Health and Welfare 2014.
46. Fransson EI, Heikkila K, Nyberg ST et al. Job strain as a risk factor for leisure-time physical inactivity: an individual-participant meta-analysis of up to 170,000 men and women: the IPD-Work Consortium. *Am J Epidemiol* 2012;176:1078-89.

47. Stults-Kolehmainen MA, Sinha R. The effects of stress on physical activity and exercise. *Sports Med* 2014;44:81-121.
48. Tytherleigh \* MY, Webb C, Cooper CL, Ricketts C. Occupational stress in UK higher education institutions: a comparative study of all staff categories. *Higher Education Research & Development* 2005;24:41-61.
49. McKercher CM, Schmidt MD, Sanderson KA, Patton GC, Dwyer T, Venn AJ. Physical activity and depression in young adults. *Am J Prev Med* 2009;36:161-4.
50. Siegrist J, Rodel A. Work stress and health risk behavior. *Scand J Work Environ Health* 2006;32:473-81.
51. Ishizaki M, Nakagawa H, Morikawa Y et al. Influence of job strain on changes in body mass index and waist circumference--6-year longitudinal study. *Scand J Work Environ Health* 2008;34:288-96.
52. Nyberg ST, Heikkila K, Fransson EI et al. Job strain in relation to body mass index: pooled analysis of 160 000 adults from 13 cohort studies. *J Intern Med* 2012;272:65-73.
53. Siegrist J, Starke D, Chandola T et al. The measurement of effort–reward imbalance at work: European comparisons. *Social Science & Medicine* 2004;58:1483-1499.
54. Fujishiro K, Lividoti Hibert E, Schernhammer E, Rich-Edwards JW. Shift work, job strain and changes in the body mass index among women: a prospective study. *Occupational and Environmental Medicine* 2017;74:410.

## **Figure legends**

Figure 1. Participants in the CDAH study

