



# Incidence, Demographics, and Clinical Characteristics of Diabetes of the Exocrine Pancreas (Type 3c): A Retrospective Cohort Study

Diabetes Care 2017;40:1–8 | <https://doi.org/10.2337/dc17-0542>

Chris Woodmansey,<sup>1</sup>  
 Andrew P. McGovern,<sup>1</sup>  
 Katherine A. McCullough,<sup>1,2</sup>  
 Martin B. Whyte,<sup>1,2</sup> Neil M. Munro,<sup>1</sup>  
 Ana C. Correa,<sup>1</sup> Piers A.C. Gatenby,<sup>1,3</sup>  
 Simon A. Jones,<sup>1,4</sup> and  
 Simon de Lusignan<sup>1,5</sup>

## OBJECTIVE

This study was conducted to describe the incidence of diabetes after pancreatic disease, assess how these patients are classified by clinicians, and compare clinical characteristics with type 1 and type 2 diabetes.

## RESEARCH DESIGN AND METHODS

Primary care records in England ( $n = 2,360,631$ ) were searched for incident cases of adult-onset diabetes between 1 January 2005 and 31 March 2016. We examined demographics, diabetes classification, glycemic control, and insulin use in those with and without pancreatic disease (subcategorized into acute pancreatitis or chronic pancreatic disease) before diabetes diagnosis. Regression analysis was used to control for baseline potential risk factors for poor glycemic control ( $\text{HbA}_{1c} \geq 7\%$  [ $53 \text{ mmol/mol}$ ]) and insulin requirement.

## RESULTS

We identified 31,789 new diagnoses of adult-onset diabetes. Diabetes after pancreatic disease (2.59 [95% CI 2.38–2.81] per 100,000 person-years) was more common than type 1 diabetes (1.64 [1.47–1.82];  $P < 0.001$ ). The 559 cases of diabetes after pancreatic disease were mostly classified by clinicians as type 2 diabetes (87.8%) and uncommonly as diabetes of the exocrine pancreas (2.7%). Diabetes after pancreatic disease was diagnosed at a median age of 59 years and BMI of  $29.2 \text{ kg/m}^2$ . Diabetes after pancreatic disease was associated with poor glycemic control (adjusted odds ratio, 1.7 [1.3–2.2];  $P < 0.001$ ) compared with type 2 diabetes. Insulin use within 5 years was 4.1% (3.8–4.4) with type 2 diabetes, 20.9% (14.6–28.9) with diabetes after acute pancreatitis, and 45.8% (34.2–57.9) with diabetes after chronic pancreatic disease.

## CONCLUSIONS

Diabetes of the exocrine pancreas is frequently labeled type 2 diabetes but has worse glycemic control and a markedly greater requirement for insulin.

Diabetes of the exocrine pancreas is thought to constitute 9% of diabetes in hospitalized patients (1). This form of diabetes results when a process, such as inflammation, neoplasia, or surgical resection, disrupts the global architecture or physiology of the pancreas, often resulting in both exocrine and endocrine dysfunction (2,3). Diabetes of the exocrine pancreas is now the suggested universal nomenclature (4); terms such as type 3c diabetes and secondary pancreatic diabetes have previously been used. To our

<sup>1</sup>Department of Clinical and Experimental Medicine, University of Surrey, Guildford, Surrey, U.K.

<sup>2</sup>The Centre for Diabetes, Endocrinology and Research (CEDAR), Royal Surrey County Hospital, Guildford, Surrey, U.K.

<sup>3</sup>Regional Oesophagogastric Unit/Gastrointestinal Surgery, Royal Surrey County Hospital, Guildford, Surrey, U.K.

<sup>4</sup>Division of Healthcare Delivery Science/Center for Healthcare Innovation and Delivery Science (CHIDS), Department of Population Health, New York University, Langone Medical Centre, New York, NY

<sup>5</sup>Royal College of General Practitioners Research and Surveillance Centre, London, U.K.

Corresponding author: Simon de Lusignan, [s.lusignan@surrey.ac.uk](mailto:s.lusignan@surrey.ac.uk).

Received 21 March 2017 and accepted 5 August 2017.

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc17-0542/-/DC1>.

© 2017 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetesjournals.org/content/license>.

knowledge, preceding pancreatic disease has never been systematically identified in a cohort of people with diabetes in primary care, and thus, the comparative incidence and clinical characteristics of diabetes of the exocrine pancreas remain uncertain.

Recognizing the role of pancreatic damage in the development of a patient's diabetes is vital to inform appropriate management plans. In diabetes of the exocrine pancreas, the production of insulin is usually reduced. This is caused by  $\beta$ -cell dysfunction after pancreatic inflammation or by absolute  $\beta$ -cell loss (5,6). In addition, pancreatic polypeptide upregulates insulin receptor expression in the liver, and its loss can lead to hepatic insulin resistance (7), an important physiological difference between diabetes of the exocrine pancreas and type 1 diabetes. Glucagon production from pancreatic  $\alpha$ -cells is also diminished, which may explain the episodes of severe hypoglycemia that are reported in some patients (8,9). Failure to recognize this altered physiology may result in suboptimal treatment. Newer incretin-based agents, such as glucagon-like peptide 1 receptor agonists ("incretins") and dipeptidyl peptidase 4 inhibitors ("gliptins"), would usually be considered contraindicated because of the presence of pancreatic damage (10). Insulinopenic patients with diabetes of the exocrine pancreas could be unfairly labeled as poorly compliant if they are misclassified as type 2 diabetes and show resistance to oral antihyperglycemic agents. Malabsorption secondary to pancreatic exocrine dysfunction is common, and pancreatic enzyme and vitamin D replacement is required to prevent malnutrition and osteoporotic bone disease (11–13).

Few studies have investigated how the distinct pathophysiology of diabetes of the exocrine pancreas manifests clinically. Limited evidence on the likelihood of significant hyperglycemia in diabetes of the exocrine pancreas makes deciding when to escalate antihyperglycemic therapy difficult. Retinopathy screening of patients with diabetes after chronic pancreatitis has suggested that when poor glycemic control is present in diabetes of the exocrine pancreas, the risk of microvascular disease is as high as with type 1 diabetes (14).

That 40% of hospital inpatients with diabetes of the exocrine pancreas are

misdiagnosed as type 2 diabetes has been suggested (1). Whether all diabetes that occurs after pancreatic disease should be considered diabetes of the exocrine pancreas is not clear, and there are currently no validated diagnostic criteria for diabetes of the exocrine pancreas.  $\beta$ -Cell failure does occur in adults with type 2 diabetes, but usually after a period of sustained insulin resistance (15), and only 1.7% of people with newly diagnosed type 2 diabetes required insulin as a first line therapy in a recent large U.K. retrospective cohort (16). If insulin is commonly required early in the course of diabetes in adults with prior pancreatic disease, it would imply that type 2 diabetes is not the usual explanation for diabetes in this patient group.

There is scarce evidence on how the clinical characteristics of diabetes after acute pancreatitis compare with those of diabetes after chronic pancreatic disease. Hypothetically, acute pancreatitis may have a lower propensity toward diabetes if an attack results in less significant organ damage than occurs with chronic pancreatic disease. However, a recent systematic review of cohorts of patients with acute pancreatitis found a prevalence of diabetes of 23%, and that 27 of 108 patients with diabetes after severe acute pancreatitis were using insulin if monitored beyond 5 years (17). Furthermore, a study of the prevalence of diabetes after pancreatitis or pancreatic cancer in New Zealand identified the largest cohort of cases of diabetes of the exocrine pancreas described in the literature thus far and found acute pancreatitis was the most frequent preceding disease (18). Comparison of diabetes after acute pancreatitis with both diabetes after chronic pancreatic disease and diabetes in the absence of preceding pancreatic disease would help confirm that diabetes of the exocrine pancreas routinely follows acute pancreatitis and ascertain to what extent diabetes of the exocrine pancreas is a homogenous group.

We aimed to describe the incidence of diabetes after pancreatic disease in the general population and to assess whether this patient group is more commonly classified by doctors as type 1 diabetes, as type 2 diabetes, or as diabetes of the exocrine pancreas. We also assessed whether the clinical course of diabetes after pancreatic disease differs from type 1 diabetes and type 2 diabetes by comparing

trends in glycemic control and insulin use over time.

## RESEARCH DESIGN AND METHODS

### Study Design

We identified new diagnoses (incident cases) of diabetes and then performed a retrospective cohort study comparing those with preceding pancreatic disease to those without. Our comparative groups were adults with a new diagnosis of 1) type 1 diabetes with no prior pancreatic disease, 2) type 2 diabetes with no prior pancreatic disease, and 3) diabetes after pancreatic disease, which we subdivided into 1) diabetes after acute pancreatitis and 2) diabetes after chronic pancreatic disease. We report the baseline characteristics and population incidence for each group along with the diabetes classification, assigned by treating physicians, of those with diabetes after pancreatic disease. Clinical outcome measures were glycemic control and insulin use.

### Setting and Data Source

We searched routinely collected electronic records from patients in England who were registered at primary care practices participating in the Royal College of General Practitioners Research and Surveillance Centre (RCGP RSC) (19). The study period was 1 January 2005 to 31 March 2016, inclusive.

The RCGP RSC extracts data once weekly from the primary care records of more than 2 million patients. All patients in the U.K. are registered with a primary care practice and have a unique National Health Service number that prevents "double-counting" in research. U.K. primary care is fully computerized. Consultations with secondary care specialists (in the large majority via the public National Health Service, but on occasion in the private sector) occurs by referral from the primary care doctor, and the specialist will write back to the primary care doctor with the findings of each consultation. The events of any acute hospital admissions are also fed back to patients' primary care practices. This information is then coded into the patient's primary care record. Data in U.K. primary care systems are encoded using Read codes. Read codes can be considered as equivalent to ICD codes but provide a more comprehensive coding system that allows the recording of diagnoses, problems, clinical findings, and treatments (20). The RCGP

RSC was initially established in 1964 for surveillance of influenza and respiratory infections (21) but has since expanded its scope to allow more detailed research.

## Participants and Study Groups

### Diabetes

We used Read codes to identify adults with diabetes. Adulthood was defined as age 18 years or older in accordance with national U.K. diabetes guidelines (22,23). We excluded all those whose diagnosis of diabetes preceded our study period (Read code of diagnosis before the study period, oral antihyperglycemic medication or insulin use before the study period, first diabetes Read code less than 1 year after registration at a practice). We excluded from our primary analysis those who had a conflicted coding of both type 1 and type 2 diabetes but no history of pancreatic disease but reanalyzed them as part of a secondary sensitivity analysis.

### Pancreatic Disease

We also used Read codes to identify all those with a diagnosis of pancreatic disease preceding their diagnosis of diabetes. This group was subdivided into those with a history of acute pancreatitis only and those with any history of chronic pancreatic disease. The chronic conditions we identified were chronic pancreatitis, pancreatic cancer, hemochromatosis, cystic fibrosis, and surgical pancreatic resection. We included single and repeated episodes of acute pancreatitis in the acute pancreatitis group. However, a patient who was ever diagnosed with a chronic pancreatic condition before developing diabetes was included in the chronic pancreatic disease group and excluded from the acute pancreatitis group.

## Baseline Characteristics and Outcome Measures

### Baseline Characteristics

The baseline characteristics of age, sex, ethnicity, index of multiple deprivation score, smoking status, and BMI were extracted from the time of diagnosis of diabetes. These baseline characteristics were later adjusted for as possible confounders of glycemic control and insulin use (for which we also adjusted for HbA<sub>1c</sub> at diagnosis) in the statistical models as described below.

### Classification of Diabetes After Pancreatic Disease

We examined whether those with diabetes after pancreatic disease were coded by their treating physicians as type 1

diabetes, type 2 diabetes, or diabetes of the exocrine pancreas. Any coding containing any possible synonymous term such as “secondary pancreatic diabetes mellitus” or “secondary diabetes mellitus” we defined as consistent with a diagnosis of diabetes of the exocrine pancreas.

### Glycemic Control

We extracted HbA<sub>1c</sub> values at diagnosis, at 1 year, and at 5 years for each group. We used a 12-month window of 6 months before to 6 months after the 1-year and 5-year times, during which if a measurement of HbA<sub>1c</sub> was made, we extracted it for analysis. We used the HbA<sub>1c</sub> value closest to the desired time point when more than one value was measured in the window. We determined the number with poor glycemic control at 1 year and at 5 years. We defined poor glycemic control as HbA<sub>1c</sub>  $\geq 7\%$  (53 mmol/mol), according to the generic target of the American Diabetes Association and the European Association for the Study of Diabetes hyperglycemia guidelines (24). We calculated the likelihood of poor glycemic control at both 1 and 5 years for diabetes after pancreatic disease and for type 1 diabetes, with type 2 diabetes as the reference group.

### Insulin Use

We assessed the number of patients who started insulin within 1 year and within 5 years of diabetes diagnosis and the proportion of those with sufficient follow-up who had started insulin at each time point. We calculated the likelihood of progression to insulin therapy for diabetes after pancreatic disease at both 1 and 5 years, with type 2 diabetes as the reference group.

### Statistical Methods

We calculated incidence rates as the number of new diagnoses per the total study group person-years. We used the Pearson  $\chi^2$  test with Yates correction for continuity to test for significant differences between incidence rates, the proportion of people taking insulin, and the proportion of people with poor glycemic control (25). We created logistic regression models for odds of poor glycemic control at 1 year and at 5 years (26). To analyze time to insulin use, we produced Kaplan-Meier curves comparing type 2 diabetes without preceding pancreatic disease, diabetes after acute pancreatitis, and diabetes after chronic pancreatic

disease (27). We initially created Cox proportional hazards models to assess the hazard of progression to insulin; however, these failed proportional hazards testing (28,29). We therefore created logistic regression models for odds of progression to insulin at 1 year and 5 years. For all the statistical models we report unadjusted odds ratios and odds ratios after adjustment for baseline characteristics. Our null hypothesis for all statistical tests was that there was no difference between the compared groups, and we set a confidence level of 95% such that a *P* value of  $<0.05$  would lead us to reject our null hypothesis. All statistical testing was performed using R 3.2.5 statistical software.

## RESULTS

We identified 31,789 new diagnoses (incident cases) of adult-onset diabetes with a median follow-up time of 4.5 years (interquartile range [IQR] 2.0–7.4) from the date of diabetes diagnosis (Fig. 1); of these, 559 cases were of diabetes after pancreatic disease. We subdivided this group into 361 cases of diabetes after acute pancreatitis and 198 after chronic pancreatic disease.

### Incidence


The incidence of adult-onset diabetes after pancreatic disease was higher than the incidence of adult-onset type 1 diabetes (2.59 [95% CI 2.38–2.81] per 100,000 person-years vs. 1.64 [1.47–1.82] per 100,000 person-years; *P* < 0.001) in our population. Adult-onset type 2 diabetes had the highest incidence (142.89 [41.31–144.50] per 100,000 person-years). The proportion of diabetes after pancreatic disease among adult-onset diabetes was 1.8% compared with a proportion of 1.1% for type 1 diabetes.

### Demographics

Diabetes after pancreatic disease was diagnosed at a median age of 59 years (IQR 49–70) in patients with a median BMI of 29.2 kg/m<sup>2</sup> (IQR 25.7–34.3). Full characteristics for each of the groups at baseline (diagnosis of diabetes) are provided in Table 1.

### Classification of Diabetes After Pancreatic Disease

Diabetes after pancreatic disease was rarely classified as diabetes of the exocrine pancreas (2.7% [95% CI 1.6–4.5]). A proportion was classified as type 1



**Figure 1**—Identification of study groups and exclusions.

diabetes (7.7% [5.7–10.3]), but diabetes after pancreatic disease was most commonly diagnosed as type 2 diabetes (87.8% [84.8–90.4]) (Table 2).

#### Glycemic Control

Mean HbA<sub>1c</sub> levels (Supplementary Table 1) were higher in people with diabetes after pancreatic disease than in people

with type 2 diabetes. HbA<sub>1c</sub> levels were higher at presentation ( $P = 0.002$ ) and remained higher at 1 year ( $P < 0.001$ ) and 5 years ( $P < 0.001$ ) after diagnosis. Mean HbA<sub>1c</sub> levels in people with diabetes after pancreatic disease were 8.3% (67 mmol/mol) at diagnosis, 7.1% (54 mmol/mol) at 1 year, and 7.6% (60 mmol/mol) at 5 years. Mean HbA<sub>1c</sub> levels in people with type 2 diabetes were 7.9% (63 mmol/mol) at diagnosis, 6.8% (51 mmol/mol) at 1 year, and 7.2% (55 mmol/mol) at 5 years. HbA<sub>1c</sub> levels were also higher in type 1 diabetes than in type 2 diabetes, with mean HbA<sub>1c</sub> levels of 12.1% (109 mmol/mol) at diagnosis, 7.6% (60 mmol/mol) at 1 year, and 8.6% (70 mmol/mol) at 5 years. Mean HbA<sub>1c</sub> levels in diabetes after acute pancreatitis were no different from the mean HbA<sub>1c</sub> levels in diabetes after chronic pancreatic disease at presentation ( $P = 0.214$ ), at 1 year ( $P = 0.642$ ), or at 5 years ( $P = 0.779$ ).

People with diabetes after pancreatic disease had a greater likelihood of poor glycemic control (Supplementary Table 2) than those with type 2 diabetes. Diabetes after pancreatic disease had poor control in 40.3% (95% CI 35.7–45.1) of cases compared with 32.5% (31.9–33.1) of cases of type 2 diabetes at 1 year after diagnosis ( $P < 0.001$ ). Diabetes after pancreatic disease had an unadjusted odds ratio for poor control at 1 year of 1.4 (1.1–1.7) compared with type 2 diabetes ( $P = 0.001$ ). After adjustment for baseline characteristics, the odds ratio for poor control was 1.3 (1.1–1.6;  $P = 0.005$ ). At 5 years after diagnosis, diabetes after pancreatic disease had poor control in 61.9% (54.8–68.5) of cases compared with 46.3% (45.5–47.2) of cases of type 2 diabetes ( $P < 0.001$ ). Diabetes after pancreatic disease had an unadjusted odds ratio for poor control at 5 years of 1.7 (1.3–2.2) compared with type 2 diabetes ( $P < 0.001$ ). After adjustment for baseline characteristics, the odds ratio for poor control was 1.7 (1.3–2.2;  $P < 0.001$ ). People with type 1 diabetes also had worse control than those with type 2 diabetes, with adjusted odds ratios for poor glycemic control of 1.4 (1.1–1.8) at 1 year ( $P = 0.007$ ) and 2.5 (1.7–3.7) at 5 years ( $P < 0.001$ ).

#### Insulin Use

Diabetes after pancreatic disease was associated with early initiation of insulin therapy (Supplementary Table 3) compared with



**Table 1—Baseline characteristics at the time of diagnosis of diabetes**

	Diabetes with no prior pancreatic disease		Diabetes after pancreatic disease		
	Type 1 diabetes <i>n</i> = 354	Type 2 diabetes <i>n</i> = 30,876	All <i>N</i> = 559	After acute pancreatitis <i>n</i> = 361	After chronic pancreatic disease <i>n</i> = 198
Age (years)					
<20	29 (8.2)	17 (0.1)	2 (0.4)	1 (0.3)	1 (0.5)
20–29	96 (27.1)	226 (0.7)	17 (3.0)	8 (2.2)	9 (4.5)
30–39	103 (29.1)	1,284 (4.2)	38 (6.8)	22 (6.1)	16 (8.1)
40–49	65 (18.4)	4,517 (14.6)	85 (15.2)	52 (14.4)	33 (16.7)
50–59	36 (10.2)	7,546 (24.4)	147 (26.3)	90 (24.9)	57 (28.8)
60–69	18 (5.1)	8,727 (28.3)	123 (22.0)	80 (22.2)	43 (21.7)
70–79	5 (1.4)	6,141 (19.9)	94 (16.8)	64 (17.7)	30 (15.2)
≥80	2 (0.6)	2,418 (7.8)	53 (9.5)	44 (12.2)	9 (4.5)
Sex					
Female	129 (36.4)	13,490 (43.7)	230 (41.1)	157 (43.5)	73 (36.9)
Male	225 (63.6)	17,386 (56.3)	329 (58.9)	204 (56.5)	125 (63.1)
Ethnicity					
Asian	9 (2.5)	2,148 (7.0)	21 (3.8)	16 (4.4)	5 (2.5)
Black	7 (2.0)	921 (3.0)	8 (1.4)	4 (1.1)	4 (2.0)
Mixed	1 (0.3)	229 (0.7)	2 (0.4)	2 (0.6)	0 (0.0)
Other	1 (0.3)	230 (0.7)	3 (0.5)	2 (0.6)	1 (0.5)
White	244 (68.9)	22,248 (72.1)	430 (76.9)	276 (76.5)	154 (77.8)
Unknown	92 (26.0)	5,100 (16.5)	95 (17.0)	61 (16.9)	34 (17.2)
IMD score					
First quintile	82 (23.2)	7,298 (23.6)	126 (22.5)	88 (24.4)	38 (19.2)
Second quintile	86 (24.3)	6,778 (22.0)	109 (19.5)	70 (19.4)	39 (19.7)
Third quintile	62 (17.5)	5,382 (17.4)	87 (15.6)	52 (14.4)	35 (17.7)
Fourth quintile	51 (14.4)	5,636 (18.3)	109 (19.5)	71 (19.7)	38 (19.2)
Fifth quintile	69 (19.5)	5,595 (18.1)	125 (22.4)	78 (21.6)	47 (23.7)
Unknown	4 (1.1)	187 (0.6)	3 (0.5)	2 (0.6)	1 (0.5)
Alcohol consumption*					
None	61 (17.2)	8,634 (28.0)	145 (25.9)	105 (29.1)	40 (20.2)
Safe	105 (29.7)	6,755 (21.9)	107 (19.1)	71 (19.7)	36 (18.2)
Hazardous	112 (31.6)	12,510 (40.5)	177 (31.7)	121 (33.5)	56 (28.3)
Alcoholic	21 (5.9)	1,425 (4.6)	103 (18.4)	49 (13.6)	54 (27.3)
Unknown	55 (15.5)	1,552 (5.0)	27 (4.8)	15 (4.2)	12 (6.1)
Smoking status					
Never smoked	122 (34.5)	7,280 (23.6)	100 (17.9)	60 (16.6)	40 (20.2)
Former smoker	138 (39.0)	19,310 (62.5)	318 (56.9)	217 (60.1)	101 (51.0)
Current smoker	94 (26.6)	4,224 (13.7)	140 (25.0)	84 (23.3)	56 (28.3)
Unknown	0 (0.0)	62 (0.2)	1 (0.2)	0 (0.0)	1 (0.5)
BMI (kg/m <sup>2</sup> )					
<18.5	11 (3.1)	59 (0.2)	9 (1.6)	0 (0.0)	9 (4.5)
≥18.5–24.9	107 (30.2)	2,242 (7.3)	84 (15.0)	40 (11.1)	44 (22.2)
≥25–29.9	47 (13.3)	7,451 (24.1)	130 (23.3)	89 (24.7)	41 (20.7)
≥30.0	35 (9.9)	14,886 (48.2)	196 (35.1)	151 (41.8)	45 (22.7)
Unknown	154 (43.5)	6,238 (20.2)	140 (25.0)	81 (22.4)	59 (29.8)
HbA <sub>1c</sub> (%) [mmol/mol]					
<7.0 [ $<53$ ]	3 (0.8)	10,500 (34.0)	170 (30.4)	114 (31.6)	56 (28.3)
≥7.0–7.9 [ $\geq 53$ –63]	8 (2.3)	5,740 (18.6)	94 (16.8)	64 (17.7)	30 (15.2)
≥8.0–8.9 [ $\geq 64$ –74]	4 (1.1)	2,030 (6.6)	32 (5.7)	26 (7.2)	6 (3.0)
≥9.0–9.9 [ $\geq 75$ –85]	15 (4.2)	1,344 (4.4)	31 (5.5)	17 (4.7)	14 (7.1)
≥10.0–10.9 [ $\geq 86$ –96]	20 (5.6)	1,186 (3.8)	23 (4.1)	17 (4.7)	6 (3.0)
≥11.0–11.9 [ $\geq 97$ –107]	33 (9.3)	1,018 (3.3)	24 (4.3)	12 (3.3)	12 (6.1)
≥12.0 [ $\geq 108$ ]	85 (24.0)	1,381 (4.5)	43 (7.7)	23 (6.4)	20 (10.1)
Unknown	186 (52.5)	7,677 (24.9)	142 (25.4)	88 (24.4)	54 (27.3)

Data are presented as *n* (%). Groups are split into type 1 diabetes with no prior history of pancreatic disease, type 2 diabetes with no prior history of pancreatic disease, and diabetes that developed after pancreatic disease. Diabetes after pancreatic disease is subdivided into diabetes after acute pancreatitis and diabetes after any chronic pancreatic disease. IMD, index of multiple deprivation. \*Alcohol consumption was divided into four categories based on Read code data: “none” (current nondrinker), “safe” (<14 units per week or only occasional consumption), “hazardous” (>14 units per week or excess consumption), or “alcoholic” (alcohol-related disease complications or alcoholism treatment).

**Table 2—Classification of diabetes by clinicians, obtained from Read code data, for diabetes that developed after pancreatic disease**

Classification of diabetes	Diabetes after pancreatic disease		
	All	Diabetes after acute pancreatitis	Diabetes after chronic pancreatic disease
Coding consistent with type 1 diabetes	43 (7.7)	14 (3.9)	29 (14.6)
Coding consistent with type 2 diabetes	491 (87.8)	336 (93.1)	155 (78.3)
Conflicted type 1 and type 2 diabetes codes	10 (1.8)	2 (0.6)	8 (4.0)
Coding consistent with diabetes of the exocrine pancreas	15 (2.7)	9 (2.5)	6 (3.0)
Total	559 (100)	361 (100)	198 (100)

Data are presented as *n* (%).

type 2 diabetes. At 1 year after diagnosis, 1.4% (95% CI 1.3–1.6) of those with type 2 diabetes required insulin, rising to 4.1% (3.8–4.4) at 5 years. In comparison, insulin use in diabetes after pancreatic disease was 16.3% (13.1–20.0) at 1 year ( $P < 0.001$ ), rising to 29.6% (23.6–36.4) at 5 years ( $P < 0.001$ ). Diabetes after pancreatic disease had an unadjusted odds ratio for insulin use at 1 year of 13.5 (10.3–17.5) compared with type 2 diabetes ( $P < 0.001$ ). After adjustment for baseline characteristics, the odds ratio for insulin use was 9.6 (7.0–13.2;  $P < 0.001$ ). Diabetes after pancreatic disease had an unadjusted odds ratio for insulin use at 5 years of 9.9 (7.2–13.4) compared with type 2 diabetes ( $P < 0.001$ ). After adjustment for baseline characteristics, the odds ratio for insulin use was 7.4 (5.2–10.4;  $P < 0.001$ ).

Diabetes after chronic pancreatic disease had a higher rate of insulin use than diabetes after acute pancreatitis (Fig. 2); however, patients in both of these subgroups had a greater requirement for insulin than those with type 2 diabetes (Supplementary Table 3). At 1 year after diagnosis, 9.7% (95% CI 6.8–13.7) of those with diabetes after acute pancreatitis were using insulin, rising to 20.9% (14.6–28.9) at 5 years. At 1 year after diagnosis, 28.9% (22.2–36.7) of those with diabetes after chronic pancreatic disease were using insulin, rising to 45.8% (34.2–57.9) at 5 years (all  $P < 0.001$  compared with type 2 diabetes).

### Supplementary Analyses

A summary of the patient characteristics and outcomes further subdivided by individual pancreatic diseases is provided in Supplementary Table 4. We also performed a sensitivity analysis that demonstrated that inclusion of patients with


conflicted type 1 and type 2 diabetes diagnoses did not affect our findings (data not shown).

### CONCLUSIONS

Diabetes after pancreatic disease is frequently labeled type 2 diabetes but follows a different clinical course, with worse glycemic control and a markedly greater requirement for insulin. This is the largest study to include all forms of diabetes of the exocrine pancreas to date, and to our knowledge, the only study to systematically identify pancreatic disease

in a cohort of people with newly diagnosed diabetes.

We show that diabetes after pancreatic disease has a higher incidence in adults than type 1 diabetes. We were unable to find any other primary evidence comparing the incidence, prevalence, or demographics of diabetes of the exocrine pancreas with type 1 or type 2 diabetes in the general population. This finding, however, is in keeping with a large epidemiological study of the prevalence of diabetes of the exocrine pancreas in New Zealand, in which the researchers hypothesized that



**Figure 2—Kaplan-Meier curves of freedom from insulin use over time for type 2 diabetes, diabetes after acute pancreatitis, and diabetes after chronic pancreatic disease. The shaded areas represent the 95% CIs. Log-rank  $P$  for difference:  $P < 0.001$ . The table is the number patients at risk over time.**

at a rate of 1.13 per 1,000 general population, the prevalence of diabetes of the exocrine pancreas may be not dissimilar to type 1 diabetes (18).

Important limitations when interpreting our findings are the retrospective collection of data and the use of Read codes to identify pancreatic disease and to classify diabetes. This method can result in misclassification and selection bias (30). The RCGP RSC has been shown to be a source of high-quality data and is representative of the population of England (19). However generalizability of our data to other countries may be limited, especially in those with a different burden of pancreatic disease. This is of particular significance when considering ethnic groups of low prevalence in the U.K. For example, Pendharkar et al. (31) found people of Māori and Pacific ethnicity had an incidence of diabetes after pancreatitis that was three times that of people with New Zealand European ethnicity. It is likely that our findings regarding the glycemic control and insulin requirements of those with diabetes after pancreatic disease relate to the pathophysiology of diabetes of the exocrine pancreas and so can be generalized. A limitation of these findings is the varied availability of HbA<sub>1c</sub> data for patients. Data were missing for 13.6% of cases of diabetes after pancreatic disease at 1 year and for 13.7% of cases at 5 years (Supplementary Table 1). The absence of HbA<sub>1c</sub> data were adjusted for as a potential confounder in the logistic regression models.

Our study indicates that many people with diabetes of the exocrine pancreas are currently misdiagnosed. We found that only 2.7% of people with diabetes after pancreatic disease are diagnosed with diabetes of the exocrine pancreas. Most patients were labeled type 2 diabetes, despite a sevenfold increased insulin requirement within 5 years, by which time 45.8% of patients with diabetes after chronic pancreatic disease are using insulin. Ewald et al. (1) used diagnostic criteria, including the presence of both exocrine pancreatic dysfunction and pathological pancreatic imaging, to retrospectively identify diabetes of the exocrine pancreas in their hospital's inpatients and found 49% were initially misclassified. Whether all diabetes that develops after pancreatic disease should be managed according to recommendations for diabetes of the exocrine pancreas (32) or

whether some patients would be appropriate for standard type 1 or type 2 diabetes management pathways is an area that requires further study. In our population, people with type 2 diabetes and people with diabetes after pancreatic disease were both most frequently diagnosed with diabetes between the ages of 50 and 69 years and at a BMI ( $\geq 30.0$  kg/m<sup>2</sup>) compatible with obesity. This may partially explain why diabetes of the exocrine pancreas is often mistaken for type 2 diabetes. However, a higher percentage of people with diabetes after pancreatic disease were diagnosed when younger than 50 years old and at a BMI compatible with being underweight ( $< 18.5$  kg/m<sup>2</sup>) or at a healthy weight ( $\geq 18.5$ – $24.9$  kg/m<sup>2</sup>), and it is not possible to tell from our retrospective study whether diagnosis was more common at higher age and BMI because this was the context in which diabetes first developed or because of another factor such as more frequent screening for diabetes in that demographic.

Acute pancreatitis was the most common pancreatic disease preceding diabetes in our population. This may be the result of a higher background prevalence of acute pancreatitis than the other pancreatic diseases. Considering the variety of pathological processes that cause pancreatic injury, diabetes of the exocrine pancreas is likely to be heterogeneous in nature. Patients may lie along a spectrum ranging from those with an underlying predisposition toward type 2 diabetes, in whom an additional pancreatic insult becomes diabetogenic, to those in whom a total pancreatectomy removes all endogenous insulin production. There is some evidence that the greater the severity of pancreatic damage the greater the failure of insulin production (33,34). In our study, diabetes after acute pancreatitis displayed HbA<sub>1c</sub> levels equivalent to those of diabetes after chronic pancreatic disease but was associated with a lower rate of insulin use. A possible explanation for this could be that although acute pancreatitis alters insulin production (33), diabetes after acute pancreatitis involves comparatively less failure of insulin production and more insulin resistance than diabetes after chronic pancreatic disease. One study after acute pancreatitis found HOMA-insulin resistance levels (a calculation of insulin resistance) had a stronger association with the presence

of prediabetes or diabetes than did the HOMA-% $\beta$  reduction (a calculation of  $\beta$ -cell function) (35). We note that the pathways that result in type 1 diabetes, type 2 diabetes, and diabetes of the exocrine pancreas need not be mutually exclusive. When adipocytokines implicated in the development of type 2 diabetes were studied as a possible mechanism for chronic hyperglycemia after acute pancreatitis, a suggested role was found for interleukin-6 (36). Clinical differences should not be disregarded, however, because we found diabetes after acute pancreatitis had significantly worse glycemic control (at 1 year and 5 years) and higher rates of insulin prescription than type 2 diabetes.

The combination of increased likelihood of poor glycemic control and accelerated requirement for insulin suggests that patients with diabetes after pancreatic disease may benefit from more frequent review than is normally necessitated in type 2 diabetes. Given the complexity of both classifying and managing diabetes after pancreatic disease, these patients may benefit from management in a specialist setting or with multidisciplinary dietitian and gastroenterology input. However, it is not possible from our observational study to draw inferences about whether the increased likelihood of poor glycemic control could be prevented with more frequent recognition of diabetes of the exocrine pancreas or whether comparatively poor glycemic control is unavoidable as a result of the underlying pathophysiology. Interventional studies may help to elucidate the risk-benefit profile of various antihyperglycemic agents and whether more frequent review reduces complications or hospitalizations.

In summary, diabetes that develops after pancreatic disease is significantly associated with poor glycemic control and early insulin therapy but is rarely identified as diabetes of the exocrine pancreas. Clinicians should elicit whether a patient has any history of pancreatic disease when they first present with diabetes and consider the diagnosis of diabetes of the exocrine pancreas. Diabetes of the exocrine pancreas must be appropriately recognized to tailor management, including choice of antihyperglycemic therapy, and consideration of malabsorption requiring pancreatic enzyme and vitamin D prescription. Greater awareness

of diabetes of the exocrine pancreas is required to appropriately manage this diabetes subgroup.

**Acknowledgments.** The authors acknowledge the patients in the RCGP RSC who formed the basis of this analysis.

Q:2

**Duality of Interest.** A.P.M. reports grants from Eli Lilly and AstraZeneca unrelated to this study. M.B.W. reports personal fees from AstraZeneca (speaker fee) unrelated to this study. S.d.L. heads a diabetes Real World Evidence Centre funded by Eli Lilly and reports a research grant from AstraZeneca; neither are directly related to this study. No other potential conflicts of interest relevant to this article were reported.

**Author Contributions.** C.W. contributed to the literature search and to data analysis. C.W., A.P.M., K.A.M., M.B.W., N.M.M., and P.A.C.G. contributed to background research and data interpretation. C.W., A.P.M., K.A.M., and S.d.L. contributed to conception of the study. C.W., A.P.M., and S.A.J. contributed to statistical analysis. C.W., A.P.M., S.A.J., and S.d.L. contributed to study design and planning of data analysis. A.C.C. contributed to data collection. S.A.J. contributed to validity of statistical models. All authors contributed substantially to writing of the manuscript. S.d.L. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.


## References

- Ewald N, Kaufmann C, Raspe A, Kloer HU, Bretzel RG, Hardt PD. Prevalence of diabetes mellitus secondary to pancreatic diseases (type 3c). *Diabetes Metab Res Rev* 2012;28:338–342
- Hart PA, Bellin MD, Andersen DK, et al.; Consortium for the Study of Chronic Pancreatitis, Diabetes, and Pancreatic Cancer (CPDPC). Type 3c (pancreatogenic) diabetes mellitus secondary to chronic pancreatitis and pancreatic cancer. *Lancet Gastroenterol Hepatol* 2016;1:226–237
- Lankisch PG, Löhner-Happe A, Otto J, Creutzfeldt W. Natural course in chronic pancreatitis. Pain, exocrine and endocrine pancreatic insufficiency and prognosis of the disease. *Digestion* 1993;54:148–155
- Petrov MS. Diabetes of the exocrine pancreas: American Diabetes Association-compliant lexicon. *Pancreatol* 2017;17:523–526
- Sasikala M, Talukdar R, Pavan Kumar P, et al.  $\beta$ -Cell dysfunction in chronic pancreatitis. *Dig Dis Sci* 2012;57:1764–1772
- Jethwa P, Sodergren M, Lala A, et al. Diabetic control after total pancreatectomy. *Dig Liver Dis* 2006;38:415–419
- Brunicaudi FC, Chaiken RL, Ryan AS, et al. Pancreatic polypeptide administration improves abnormal glucose metabolism in patients with chronic pancreatitis. *J Clin Endocrinol Metab* 1996;81:3566–3572
- Cui Y, Andersen DK. Pancreatogenic diabetes: special considerations for management. *Pancreatol* 2011;11:279–294
- Knop FK, Vilsbøll T, Larsen S, Madsbad S, Holst JJ, Krarup T. Glucagon suppression during OGTT worsens while suppression during IVGTT sustains alongside development of glucose intolerance in patients with chronic pancreatitis. *Regul Pept* 2010;164:144–150
- Tibaldi JM. Incorporating incretin-based therapies into clinical practice for patients with type 2 diabetes. *Adv Ther* 2014;31:289–317
- Duggan SN, Ewald N, Kelleher L, Griffin O, Gibney J, Conlon KC. The nutritional management of type 3c (pancreatogenic) diabetes in chronic pancreatitis. *Eur J Clin Nutr* 2017;71:3–8
- Trang T, Chan J, Graham DY. Pancreatic enzyme replacement therapy for pancreatic exocrine insufficiency in the 21(st) century. *World J Gastroenterol* 2014;20:11467–11485
- Klapdor S, Richter E, Klapdor R. Vitamin D status and per-oral vitamin D supplementation in patients suffering from chronic pancreatitis and pancreatic cancer disease. *Anticancer Res* 2012;32:1991–1998
- Gullo L, Parenti M, Monti L, Pezzilli R, Barbara L. Diabetic retinopathy in chronic pancreatitis. *Gastroenterology* 1990;98:1577–1581
- Chang-Chen KJ, Muller R, Bernal-Mizrachi E. Beta-cell failure as a complication of diabetes. *Rev Endocr Metab Disord* 2008;9:329–343
- Sharma M, Nazareth I, Petersen I. Trends in incidence, prevalence and prescribing in type 2 diabetes mellitus between 2000 and 2013 in primary care: a retrospective cohort study. *BMJ Open* 2016;6:e010210
- Das SL, Singh PP, Phillips ARJ, Murphy R, Windsor JA, Petrov MS. Newly diagnosed diabetes mellitus after acute pancreatitis: a systematic review and meta-analysis. *Gut* 2014;63:818–831
- Pendharkar SA, Mathew J, Petrov MS. Age- and sex-specific prevalence of diabetes associated with diseases of the exocrine pancreas: a population-based study. *Dig Liver Dis* 2017;49:540–544
- Correa A, Hinton W, McGovern A, et al. Royal College of General Practitioners Research and Surveillance Centre (RCGP RSC) sentinel network: a cohort profile. *BMJ Open* 2016;6:e011092
- de Lusignan S. Codes, classifications, terminologies and nomenclatures: definition, development and application in practice. *Inform Prim Care* 2005;13:65–70
- Elliot AJ, Fleming DM. Surveillance of influenza-like illness in England and Wales during 1966–2006. *Euro Surveill* 2006;11:249–250
- National Institute for Health and Clinical Excellence. Type 1 diabetes in adults: diagnosis and management. NICE guideline [NG17] [Internet], updated July 2016. Available from <https://www.nice.org.uk/guidance/ng17>. Accessed 7 July 2017
- National Institute for Health and Clinical Excellence. Type 2 diabetes in adults: management. NICE guideline [NG28] [Internet], updated May 2017. Available from <https://www.nice.org.uk/guidance/ng28>. Accessed 7 July 2017
- Inzucchi SE, Bergenstal RM, Buse JB, et al. Management of hyperglycemia in type 2 diabetes, 2015: a patient-centered approach: update to a position statement of the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care* 2015;38:140–149
- Yates F. Tests of significance for 2 x 2 contingency tables. *J R Stat Soc [Ser A]* 1984;147:426–463
- Cox DR. The regression analysis of binary sequences. *J R Stat Soc B* 1958;20:215–242
- Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. *J Am Stat Assoc* 1958;53:457–481
- Cox DR. Regression models and life-tables. *J R Stat Soc B* 1972;34:187–220
- Grambsch PM, Therneau TM. Proportional hazards tests and diagnostics based on weighted residuals. *Biometrika* 1994;81:515–526
- Kleinbaum DG, Morgenstern H, Kupper LL. Selection bias in epidemiologic studies. *Am J Epidemiol* 1981;113:452–463
- Pendharkar SA, Mathew J, Zhao J, Windsor JA, Exeter DJ, Petrov MS. Ethnic and geographic variations in the incidence of pancreatitis and post-pancreatitis diabetes mellitus in New Zealand: a nationwide population-based study. *N Z Med J* 2017;130:55–68
- Rickels MR, Bellin M, Toledo FG, et al.; PancreasFest Recommendation Conference Participants. Detection, evaluation and treatment of diabetes mellitus in chronic pancreatitis: recommendations from PancreasFest 2012. *Pancreatol* 2013;13:336–342
- Malecka-Panas E, Gasiorowska A, Kropiwnicka A, Zlobinska A, Drzewoski J. Endocrine pancreatic function in patients after acute pancreatitis. *Hepatogastroenterology* 2002;49:1707–1712
- Perusicová J. Glucose tolerance and secretion of insulin in chronic pancreatitis. *Acta Univ Carol Med Monogr* 1989;131:1–92
- Pendharkar SA, Asrani VM, Xiao AY, et al. Relationship between pancreatic hormones and glucose metabolism: a cross-sectional study in patients after acute pancreatitis. *Am J Physiol Gastrointest Liver Physiol* 2016;311:G50–G58
- Gillies N, Pendharkar SA, Asrani VM, Mathew J, Windsor JA, Petrov MS. Interleukin-6 is associated with chronic hyperglycemia and insulin resistance in patients after acute pancreatitis. *Pancreatol* 2016;16:748–755



# AUTHOR QUERIES

## PLEASE ANSWER ALL QUERIES

Q1: For consistency, the text is edited to reflect that 95% CI may be omitted in subsequent OR (and other values) within the same paragraph. Please confirm the changes are okay in the abstract and text or adjust as needed for consistency 

Q2: Check that the conflict of interest information for each author is presented in full in the Duality of Interest section. 