

Critical Questions in Ovarian Cancer Research and Treatment: Report of an AACR Special Conference

Robert C. Bast, Jr.^{1†}, Ursula A. Matulonis^{2†}, Anil K. Sood^{1†}, Ahmed A. Ahmed³, Adaobi E. Amobi⁴, Frances R. Balkwill⁵, Monicka Wielgos-Bonvallet⁶, David D. L. Bowtell⁷, James D. Brenton⁸, Joan S. Brugge⁹, Robert L. Coleman¹, Giulio F. Draetta¹, Kai Doberstein¹⁰, Ronny I. Drapkin¹⁰, Mark A. Eckert¹¹, Robert P. Edwards¹², Kevin M. Elias¹³, Darren Ennis¹⁴, Andrew Futreal¹, David M. Gershenson¹, Roger A. Greenberg¹¹, David G. Huntsman¹⁵, Jennifer Xiao Ye Ji¹⁵, Elise C. Kohn¹⁶, Claudia Iavarone⁹, Ernst R. Lengyel¹¹, Douglas A. Levine⁶, Christopher J. Lord¹⁷, Zhen Lu¹, Gordon B. Mills¹, Francesmary Modugno¹², Brad H. Nelson^{15,18}, Kunle Odunsi⁴, Jessica A. Pilsworth¹⁵, Robert K. Rottapel¹⁹, Daniel J. Powell, Jr.¹⁰, Li Shen⁴, le-Ming Shih²⁰, David R. Spriggs²¹, Josephine Walton¹⁴, Kaiyang Zhang²², Rugang Zhang²³, and Lee Zou²¹

¹ University of Texas MD Anderson Cancer Center, Houston, TX, ² Dana-Farber Cancer Institute, Boston, MA, ³ University of Oxford, United Kingdom, ⁴ Roswell Park Cancer Institute, Buffalo, NY, ⁵ Barts Cancer Institute, United Kingdom, ⁶ NYU Langone Medical Center, New York, NY, ⁷ Peter MacCallum Cancer Center, Australia, ⁸ University of Cambridge, United Kingdom, ⁹ Harvard Medical School, Boston, MA, ¹⁰ University of Pennsylvania, Philadelphia, PA, ¹¹ University of Chicago, Chicago, IL, ¹² University of Pittsburgh, Pittsburgh, PA, ¹³ Brigham and Women's Hospital, Boston, MA, ¹⁴ University of Glasgow, United Kingdom, ¹⁵ University of British Columbia, Canada, ¹⁶ CTEP NCI, Rockville, MD, ¹⁷ Harvard Medical School, Boston, MA, ¹⁷ Institute of Cancer Research, London, United Kingdom, ¹⁸ BC Cancer Agency, Canada, ¹⁹ University of Toronto, Canada, ²⁰ Johns Hopkins University School of Medicine, Baltimore, MD, ²¹ Massachusetts General Hospital, Boston, MD, ²² University of Helsinki, Finland, ²³ The Wistar Institute, Philadelphia, PA,

*Address correspondence to: Robert C. Bast, University of Texas MD Anderson Cancer Center, 1400 Pressler Street, Unit 1439, Houston Texas 77030, USA. Phone: (713)792-7743; Email:rbast@mdanderson.org

†Co-corresponding authors.

Funding: RCB: NIH/NCI P50 CA083639, P50 CA217685, U01 CA200462; RZ: R01 CA163377, R01 CA202919; DGH: R01 CA19567001; GBM: P50 CA217685, Ovarian Cancer Research Foundation, Prospect Creek Foundation, Adelson Medical Research Foundation; AKS: NIH/NCI P50 CA217685, ACS Research Professor

Award; RLC: CPRIT RP210214; RP160145; UAM: Breast Cancer Research Foundation; ECK: CTEP/CDTD/NCI; FM: University School of Medicine Dean's Faculty Advancement Fund

Conflicts of Interest: RCB: Royalties for CA125 Fujirebio Diagnostics, Inc.; AKS: SAB for Kiyatec, research funding from M-Trap and stockholder in Biopath; KME: Editorial Board *Current Problems in Cancer*, Elsevier; GBM: Consultant/SAB: AstraZeneca, Catena Pharmaceuticals, Critical Outcome Technologies, ImmunoMET, Ionis, Medimmune, Nuevolution, Pfizer, Precision Medicine, Signalchem Lifesciences, Symphogen, Takeda/Millennium Pharmaceuticals, Tarveda, Stock/Options: Catena Pharmaceuticals, ImmunoMet, SignalChem, Spindletop Ventures, Tarveda, Licensed Technology: HRD assay to Myriad Genetics, Sponsored Research: Abbvie, Adelson Medical Research Foundation, AstraZeneca, Breast Cancer Research Foundation, Critical Outcome Technologies, Horizon Diagnostics, Illumina, Immunomet, Ionis, Karus Therapeutics, Komen Research Foundation, Nanostring, Ovarian Cancer Research Foundation, Pfizer, Prospect Creek Foundation, Takeda/Millennium Pharmaceuticals, Tesaro; RID: SAB for Repare Therapeutics and Siamab Therapeutics. Authors: UAM, AAA, AEA, FRB, MWB, DDB, JDB, JSB, RLC, GFD, KD, MAE, RPE, DE, AF, DMG, RAG, DGH, JXJ, ECK, CI, ERL, DAL, CJL, ZL, FM, BHN, KO, JAP, RKR, DJPJ, LS, LMS, DRS, JW, KZ, RZ and LZ declare that they have no conflict of interest.

Substantial progress has been made in understanding ovarian cancer at the molecular and cellular level. Significant improvement in 5-year survival has been achieved through cytoreductive surgery, combination platinum-based chemotherapy, and more effective treatment of recurrent cancer. There are now more than 280,000 ovarian cancer survivors in the United States. Despite these advances, long term survival in late stage disease has improved little over the last four decades. Poor outcomes relate, in part, to late stage at initial diagnosis, intrinsic drug resistance and the persistence of dormant drug resistant cancer cells after primary surgery and chemotherapy. Our ability to accelerate progress in the clinic will depend on the ability to answer several critical questions regarding this disease. To assess current answers, an AACR Special Conference regarding “Critical Questions in Ovarian Cancer Research and Treatment” was held in Pittsburgh, PA, from October 1-3, 2017. While clinical, translational and basic investigators conducted much of the discussion, advocates participated in the meeting, and many presentations were directly relevant to patient care, including treatment with PARP inhibitors, attempts to improve immunotherapy by overcoming the immune suppressive effects of the microenvironment and a better understanding of the heterogeneity of the disease.

Can we detect ovarian cancer earlier? Disease limited to the ovary (Stage I) can be cured with currently available surgery and chemotherapy in up to 90% of cases and disease limited to the pelvis (Stage II) can be cured in 70%, but currently only 20-25% of patients are diagnosed in these early stages. Computer simulations suggest that detection of a greater fraction of ovarian cancers in early stage could reduce mortality by 15-43%.^{1,2} The relatively low prevalence of ovarian cancer (1:2500), requires a screening strategy that has high sensitivity (>75%) and extremely high specificity (>99.6%) to achieve a positive predictive value of 10%, i.e., 10 operations for each case of ovarian cancer detected. Neither the serum biomarker CA125 nor transvaginal sonography (TVS) used alone can achieve this sensitivity or specificity. A Risk of Ovarian Cancer Algorithm (ROCA) has, however, been developed that measures the trend of CA125 from year to year. Rising CA125 has triggered TVS in 1-3% of women screened and abnormal TVS has prompted laparotomy. Both the Normal Risk Ovarian Cancer Screening Study (NROSS)³ and the United Kingdom Collaborative Trial of Ovarian Cancer Screening (UKCTOCS)⁴ conducted in postmenopausal women at average risk demonstrated that, used in this way, CA125 and TVS achieved >99.6% specificity with 3-4 operations for each case detected. Both studies detected early stage disease in 40-70% of cases. With >200,000 participants, the UKCTOCS was powered to detect a survival advantage. While, overall, the study did not attain statistical significance, a pre-specified subset

of patients with prevalent disease demonstrated a 20% reduction in mortality ($P < 0.021$). With wide confidence limit around this estimate, additional follow-up will be required, but clearly there is room for improvement in serum biomarkers and in imaging.

For more effective detection, greater sensitivity is required in the initial phase of two stage strategies, while maintaining very high specificity. HE4 and CA72-4 antigens can detect approximately 16% of early stage ovarian cancers missed by CA125, but do not provide lead time. Small amounts of ovarian or fallopian tube cancer can evoke the production of autoantibodies. Virtually all high grade serous ovarian cancers have mutations of *TP53*, and autoantibodies against TP53 can be detected in 20% of patients, rising a median of 8 months prior to CA125 and 22 months prior to diagnosis in patients with normal CA125.⁵ HE4 antigen-autoantibody complexes are found in sera from 39% of early stage ovarian cancer patients, CA125 in 62% and the combination in 80%.

New data indicate that circulating tumor DNA (ctDNA) can be detected in blood or in cervical secretions in 5% of cases with early stage ovarian cancer complementing CA125 and promising to improve detection, which may be particularly relevant for surveillance of women with BRCA1/2 germ line mutations who are delaying preventive bilateral salpingo-oophorectomy to complete their families.⁶ In studies of *TP53* ctDNA, pre-selection of DNA fragments from plasma prior to assay substantially enhanced sensitivity and this might also prove useful for detecting amplified or mutant DNA in cervical secretions or uterine washings. As *TP53* is mutated in a wide spectrum of cancers determining tissue of origin for ctDNA assays could prove problematic. Promising data have also been obtained by neural-network analysis of a 9 miRNA panel that can distinguish malignant from benign pelvic masses that include early stage disease.⁷

With regard to prevention, use of oral contraceptives (OC) and pregnancy reduce risk by about 30% each, with greater protection conferred with longer OC duration and increasing parity. Recent data suggest that breast feeding can also decrease the risk of ovarian cancer by about 30%.⁸ Longer total duration, increasing number of offspring nursed, and earlier age at first breastfeeding increase the protective effect.

How does the microenvironment influence cancer growth and resistance to treatment? Ovarian cancer growth and resistance to treatment are affected by angiogenesis, extra-cellular matrix and cancer-associated fibroblasts in the tumor microenvironment. Anti-angiogenic therapy has targeted vascular endothelial

growth factor (VEGF) and its receptors (VEGFRs). Most clinical trials with VEGF/VEGF-R targeted drugs have shown improved progression-free survival, but not overall survival. Recently identified, resistance to anti-angiogenic therapy has been associated with the accumulation of tumor-associated macrophages (TAMs). Targeting TAMs using CSF1R-targeted drugs in combination with anti-VEGF drugs has improved outcomes in preclinical models.⁹ Another potential target is EGFL6, one of the most highly expressed genes in tumor endothelial cells.¹⁰

Matrix proteins are dysregulated in ovarian cancer. A comprehensive profile of the ovarian cancer “matrisome” has been obtained by measuring and integrating multiple components including gene expression, proteomics, cytokine and chemokine levels, cellularity, and extracellular matrix organization in clinical biopsies. An expression pattern for 22 matrisome genes distinguished patients with a shorter overall survival in high-grade serous ovarian cancer (HGSOC) and in twelve other primary solid cancers, suggesting that there may be a common matrix response to human cancer.¹¹ Networks of cytokines and chemokines appear to regulate the influx of leukocytes into ovarian cancer metastases and an index of matrisome proteins correlates with infiltration of CD4+ and FOXP3+ T cells.

Cancer-associated fibroblasts (CAFs) from metastatic sites have high expression of nicotinamide N-methyltransferase (NNMT). Functionally, NNMT regulates the methylation of repressive histone marks and expression of genes involved in CAF differentiation by depleting S-adenosyl methionine levels. Knockdown of NNMT in CAFs was sufficient to attenuate their ability to promote the proliferation, migration, and metastasis of ovarian cancer cells.¹²

Do ovarian cancers have distinctive metabolic vulnerabilities? Unlike many other cancers, epithelial ovarian cancer metastases often remain within the abdominal cavity. Microscopic residual disease (MRD) can persist following primary chemotherapy, grow progressively and give rise to recurrence, intestinal obstruction and death. Isolating MRD and measuring genotypic and phenotypic changes in tiny specimens can be challenging. Techniques have been developed to produce whole genome sequencing of picogram quantities of DNA and to measure the prevalence of mutations using phase information.

Within the peritoneal cavity, certain metabolic pathways are key drivers of ovarian cancer cell growth and survival. Ovarian cancer cells can condition CAFs, which, in turn, regulate important cancer cell activities in a

paracrine fashion. Adipocytes in the omentum provide fatty acids to adjacent ovarian cancer cells to generate much needed energy. The fatty acid receptor, CD36, is upregulated when ovarian cancer cells are co-cultivated with adipocytes.¹³

Metabolism can affect drug resistance. Most ovarian cancer patients receive a combination of carboplatin and paclitaxel, but less than half of patients respond to paclitaxel.¹⁴ Knockdown of the glycolytic enzyme 6-phosphofructo-2-kinase/fructose-2, 6-biphosphatase 2 (PFKFB2) enhances paclitaxel response in ovarian cancer cells with wild type *TP53*. Silencing PFKFB2 increases the rate of glycolysis, but decreases the flow of intermediates through the pentose-phosphate pathway in cancer cell lines with wtTP53, decreasing NADPH. Reactive oxygen species (ROS) accumulate and stimulate phosphorylation of Janus kinase (JNK), induce G1 cell cycle arrest, and initiate apoptosis that depends upon upregulation of p21Cip1 and Puma. Targeting PFKFB2 is a promising strategy for sensitizing ovarian cancers with wild type TP53 to paclitaxel.

Salt-induced kinase 2 (SIK2) is upregulated in 30% of primary ovarian cancers and overexpressed in omental metastases to a greater extent than in primary cancers.¹⁵ Adipocytes activate SIK2 in ovarian cancer cells leading to downstream phosphorylation of p85 and activation of the PI3K pathway. In addition, SIK2 augments AMPK in regulating fatty acid oxidation and energy production. Targeting SIK2, either genetically or by specific small molecule inhibitors, significantly reduces metastasis in vivo and adipocyte-induced cancer cell proliferation in culture. Knockdown of SIK2 also enhances sensitivity to paclitaxel by inhibiting centrosome splitting and PI3Kinase activity and downregulating survivin.¹⁶ Novel small molecule inhibitors of SIK2-ARN-3236 and ARN3261 enhance paclitaxel response in culture and in xenografts.¹⁷ ARN-3261 will enter first-in-human trials later this year.

Argininosuccinate synthase 1 (ASS1), a crucial enzyme for synthesis of arginine, is lost in a fraction of clear cell ovarian cancers. Since cells with no expression of ASS1 become dependent on external arginine, deprivation of arginine may provide a promising strategy to enhance chemotherapy efficacy.¹⁸

How do we measure and target the genetic, epigenetic and transcriptional heterogeneity of different types of ovarian cancer, common and rare? A significant fraction of HGSOCs are thought to arise from serous tubal intraepithelial carcinomas (STICs) in the fimbriae of fallopian tubes. While mutation of *TP53* is an early event, loss of heterozygosity for *TP53*, mutations of *BRCA1*, *BRCA2*, loss of *PTEN* and copy number

abnormalities are also found in STICS.¹⁹ Evolutionary analyses reveal that *TP53* signatures and STICs are precursors of metastatic ovarian carcinoma and several years can elapse between development of a STIC and initiation of ovarian carcinoma with metastases, providing an important window of opportunity for early detection and prevention of this disease by surgically removing the fallopian tubes.

HGSOC is typified by frequent copy number alterations across the entire genome with loss of homologous recombination DNA repair machinery. Aside from universal mutation of *TP53* and mutation of *BRCA1* or *BRCA2* in 15-20% of cases, HGSOC has a paucity of dominant acting mutations, making targeted therapy directed against driver mutations a difficult strategy to deploy clinically. A recent study has begun to decode the complexity of copy number changes, identifying seven distinct copy number signatures that predict both overall survival and the probability of platinum-resistant relapse. Copy number signatures can also be used to combinations of agents that are likely to be more effective.²⁰

HGSOC cell lines appear to depend upon “quality control” pathways which sense and destroy damaged transcripts and proteins. Nonsense mediated decay (NMD) is highly active in HGSOC cell lines.²¹ Interference with enzymes that control NMD either genetically or with small molecule inhibitors is deleterious to HGSOC cell lines. NMD inhibition triggers cell death through activation of the unfolded protein response. Recent studies of HGSOC with multi-parametric mass spectrometry (CyTOF) have identified rare cell phenotypes within ovarian tumors in addition to the dominant cell subset.²² Rare populations included ovarian cancer cells that co-expressed vimentin and E-cadherin which may play a role in epithelial mesenchymal transition, as well as populations that co-expressed vimentin, HE4 and c-myc that were associated with poor patient outcome.

Many widely used human and murine ovarian cancer cell lines do not resemble the genotype of HGSOC.²³ Genetically engineered mouse models of HGSOC have been developed in which *TP53*, *BRCA1*, *BRCA2*, *PTEN* and *NF1* have been deleted in fallopian tube epithelial cells. While these models have great potential to expand our understanding of HGSOC biology, their use requires large-scale breeding programs and primary cancers can take many months to develop. Thus, transplantable models remain valuable research tools. ID8, a widely-used syngeneic model of ovarian cancer, lacks the frequent mutations observed in human HGSOC. Using CRISPR/Cas9 gene editing²⁴, novel ID8 sublines have been developed with deletion of *TP53*, *PTEN* and *NF-1*.²⁵ Loss of *PTEN* and *NF1* significantly increased the rate of intraperitoneal

growth when compared to loss of *TP53*. By contrast, *BRCA1* loss had no effect on intraperitoneal growth, while loss of *BRCA2* actually decreased growth.

A recent review underlines the possible strengths of 1) orthotopic mouse models where cancer cells are injected into the ovarian bursa, 2) patient derived xenografts (PDXs) where human ovarian cancer cells are grown subcutaneously or intraperitoneally in immune-incompetent nu/nu (T-cell deficient), SCID (T and B cell deficient) or NSG (T, B and NK deficient) mice, and humanized mice reconstituted with human immunocytes.²⁶ A majority of PDX models have been shown to correlate histologically, genotypically and in response to platinum-based chemotherapy, but with repeated passage copy number abnormalities have diverged, consistent with genetic instability. Use of humanized mice has permitted evaluation of immunotherapy that cannot be evaluated with PDX's in immune-deficient mice, although response to check point inhibitors has been much greater than those encountered in the clinic. An important principle to recall is that xenograft models are derived from a single patient and large numbers of different models must be tested to encompass the heterogeneity of clinical cancer.

Low-grade serous carcinomas (LGSC) account for approximately 10% of all serous cancers and can arise de novo or from serous borderline tumors. LGSC occur at a younger age than HGSOC and exhibit relative chemoresistance, but are associated with prolonged overall survival. While virtually all LGSC have wtTP53, up to 40% contain a KRAS mutation and 5-10% contain a BRAF mutation. At least 80% of LGSCs are ER+, and approximately 50% are PR+. In addition, the IGF-1 pathway and angiogenesis appear to be potential therapeutic targets. Clinical benefit of conventional chemotherapy in LGSC is limited,²⁷ but bevacizumab,²⁸ hormonal therapies,^{Error! Bookmark not defined., 29} and targeted agents, such as MEK or BRAF inhibitors have exhibited greater activity in LGSOC than in HGSOC. Currently, second-generation trials of MEK inhibitors³⁰ are nearing completion and biomarker studies within these trials should provide important information on the relationship of mutational status to anti-tumor activity.

Small cell carcinomas of the ovary hypercalcemic type (SCCOHT) are rare but highly aggressive cancers that exhibit truncating or splice site mutations in the SMARCA4 gene that encodes BRG1, 1 of 2 potential ATPases within the SWI/SNF complex. SCCOHT are quite distinctive in that these SMARCA4 mutations are likely initiating events and occur within the context of an extremely quiescent genome. Treatment with EZH2 inhibitors has been proposed and is being tested. While immune modulation with check point inhibitors have

been associated with high mutational burden in other types of cancer, four cases of SCCOHT responded impressively to immune modulation therapy, despite a very low mutational burden.³¹

Adult-type granulosa cell tumors (AGCT) account for only 5% of all ovarian cancer and are characterized by a C402G somatic missense mutation in the transcription factor *FOXL2*.³² Across three cohorts, approximately 20% of the cancers thought to be AGCT did not have the pathognomonic mutation and these cancers accounted for 70% of all deaths from disease within the first five years. ~~Recently, a *TERT* C228T promoter mutation was identified in 50% of AGCT, but not in other histotypes, and was found more frequently in recurrent tumors ($p = 0.003334$).~~ Recently, a *TERT* C228T promoter mutation was identified in 22% of primary AGCT and 41% of recurrent AGCT, but not in other histotypes. *TERT* mutation was significantly more frequent in recurrent tumors ($p = 0.003334$).

Mature cystic teratomas (MCT) are the most common germ cell tumor of the ovary. Less than one percent of MCTs transform, usually into a squamous cell carcinoma (SCC), but the mean overall survival of patients with transformed MCT is less than 2 years. In a series of 25 cases, the SCC components had *TP53* abnormalities in 80% of cases, *PIK3CA* mutations in 52% and *CDKN2A* abnormalities (deletion or loss-of-function mutation) in 44%.³³ Cases with mutant *TP53* had a better prognosis than wild type cases.

Can we predict, detect and reverse drug resistance to conventional and targeted therapy?

Targeted therapy can evoke adaptive responses in tissue culture, animal models, and patient samples. Measurement of nodes and pathways that are upregulated by drug treatment can identify effective combinations of targets and drugs. Using reverse phase protein arrays to identify adaptive responses after treatment with targeted therapy, mutant *RAS* has been found to be a potent mediator of resistance to PARP inhibitors that can be overcome by combinations of PARP and MEK or ERK inhibitors, which could be relevant to treatment of low-grade serous cancers.³⁴ PARP inhibitors induced a STING response that sensitizes syngeneic tumors to PARP inhibitors and immune checkpoint inhibitors including anti-PD1 or anti-PDL1. BRD4 inhibitors induce marked HR defects that synergize with PARP inhibitors primarily through the downregulation of CTIP.³⁵

MEK inhibition with cobimetinib (GDC-0973) alone had minimal effect on 14 HGSOC PDX models, but produced strong upregulation of the pro-apoptotic protein BIM, which undergoes degradation following ERK activation. Combining targeting of the MEK pathway with inhibition of the anti-apoptotic proteins BCL-2/XL

navitoclax (ABT-263) was more effective in reducing cell number and increasing cell death than single agents in the majority of PDX models assessed in vitro and in vivo. Moreover, high pre-treatment protein levels of BIM predicted response to combination therapy.³⁶

Assay of plasma ctDNA promises to identify relevant targets for therapy. A panel of 508 cancer genes has been used to identify actionable mutations and copy-number alterations in patients with HGSOC. Tumor variants could be detected at a low frequency (0.01%) in ctDNA. Altered variants correlated with improved response to treatment. Interestingly, the fraction of tumor-derived variants increased during treatment even in patients with a complete clinical response, potentially detecting subclones that remain refractory to treatment. Some of these variants contain potentially actionable mutations. Interestingly, mutations in chromatin modifiers were significantly enriched among patients with poor response.³⁷

How can we optimize the ability of PARP inhibitors to exploit defects in DNA repair? HGSCs can respond to PARP inhibitors, particularly in the presence of *BRCA 1* or *2* mutations or other abnormalities that compromise homologous recombination (HR) DNA repair. HR dysfunction occurs in nearly 50% of HGSCs.³⁸ Biallelic loss of *BRCA1*, *BRCA2*, or many of its interacting partners renders cells up to several hundred-fold more sensitive to PARPi. Three PARPi are currently approved for treatment of recurrent ovarian cancer including Olaparib, Rucaparib, and Niraparib.^{39,40,41,42,43} Resistance to these PARP inhibitors is a growing problem. A number of mechanisms have been identified, including reversion mutations in *BRCA1* or *BRCA2*, defects in DNA repair caused by loss of REV7 or 53BP1 function⁴⁴, alterations in proteins that stabilize replication forks and upregulation of PgP transporters. PARP inhibitor resistance can also be caused by mutations in the *PARP1* gene itself. Despite PARP inhibitor resistance, *PARP1* mutant tumor cells retain the platinum salt sensitivity seen in *BRCA1* mutant cells, suggesting that cross resistance does not occur.

One rationale for developing combinations of other targeted therapy with PARPi is to convert an HR proficient cancer into one that is HR deficient (HRD) by adding another agent that inhibits HR or replication fork protection such as anti-angiogenic agents, VEGFR inhibitors, CDK inhibitors, PI3K inhibitors, PD1 blockade, ATR inhibitors and CHK1 inhibitors. Trials combining PARPi and these various agents are currently in phase I through III. The oral VEGFR inhibitor Cediranib has been combined with Olaparib in Phase I-II trials showing enhanced efficacy in ovarian cancers that were *BRCA* wildtype suggesting that the addition of an anti-angiogenic agent rendered the cancer cell more HR-like and enhanced the efficacy of the PARPi.⁴⁵ Phase III prospective

trials of this combination are now underway. A phase I clinical trial of the PI3K inhibitor BKM120 and Olaparib shows clinical activity in BRCA wildtype platinum resistant ovarian cancer with a 64% rate of benefit, largely from stable disease.⁴⁶

ATR (Ataxia telangiectasia and Rad3 related) kinase inhibition can overcome acquired resistance to PARPi in *BRCA* mutant cancer cells.⁴⁷ BRCA1 and BRCA2 control genome integrity through protection of stalled replication forks,⁴⁸ in addition to their previously established roles in HR.⁴⁹ Interestingly, genetic perturbations that mitigate replication fork stress in *BRCA* mutant cells have been implicated in resistance to PARPi and platinum compounds.⁵⁰ ATR is a primary sensor and transducer of replication stress signals, where its kinase activity is induced by events on extended single stranded DNA segments. Subsequent ATR dependent phosphorylation and activation of the checkpoint kinase 1 (Chk1) stabilizes replication forks and prevents catastrophic replication origin firing that leads to genome fragmentation and cell death.⁵¹ Combinations of PARPi and ATRi have synergistic activity in *BRCA* mutant ovarian cancer cells that had acquired resistance to PARPi.⁵²

Chk1 activation is a potential determinant of response to PARPi. Chk1 phosphorylation can be elevated specifically in response to small molecule PARPi that trap PARP1 on chromatin. In contrast, PARPi agents that produce minimal PARP1 trapping do not activate Chk1. Chk1 inhibition in combination with a non-trapping PARPi resulted in improved tumor regression in PDX models of HGSOC. Chk1 phosphorylation may be a convenient biomarker of response to PARPi.⁵³

Approximately 20% of HGSCs with amplification of *CCNE1* (encoding cyclin E2), exhibit HR proficiency and de novo resistance to PARPi or platinum compounds. Synthetic lethality has been observed between the presence of high levels of cyclin E and *BRCA* mutation,⁵⁴ explaining the mutually exclusive nature of *CCNE1* amplifications and *BRCA* mutations in HGSC. Rad51 homolog overexpression enhanced transformation of cyclin E expressing cells. The Rad51 homologs (Rad51 B, C, D, XRCC2 and XRCC3) are thought to promote canonical Rad51 filament formation and enhance HR and replication fork protection activities.⁵⁵ XRCC2 is upregulated in cyclin E overexpressing cells and is also synthetic lethal in this setting.⁵⁶

DNA damage can induce inflammatory cytokine signaling that might augment effects of immunotherapy.⁵⁷ Inflammatory cytokines modify the tumor microenvironment by recruiting immune cells that are critical for both local and systemic responses to immunotherapy and radiotherapy in preclinical murine cancer models.⁵⁸ Cell

cycle progression through mitosis following ionizing radiation or PARPi is essential to activate type I interferon responses. Mitotic progression dependent inflammatory signaling involves micronuclei formation. Micronuclei frequently rupture in the subsequent interphase,⁵⁹ thus exposing genomic DNA to the pattern recognition receptor cGAS. Activation of cGAS within micronuclei signals through the STING protein to promote inflammatory cytokine dependent gene expression. Interestingly, inhibiting progression through mitosis or loss of pattern recognition by cGAS-STING also impairs systemic anti-tumor immune responses in the context of therapy combining ionizing radiation and immune checkpoint blockade. DNA damage dependent inflammation could be used to harness immune responses that eradicate both chemotherapy-sensitive and -resistant populations.

How can we enhance the immune response in ovarian cancer? In contrast to some dramatic results report in other cancer types, monotherapy with checkpoint inhibitors that bind CTLA4, PD1 or PDL1 has produced response rates of <15% in unselected ovarian cancer patients. Use of checkpoint inhibitors individually or in combination can induce substantial toxicity from autoimmune disease. There is clearly a need to identify biomarkers for response or lack of response to these agents. Based on infiltration of CD8, Treg and B cells into the epithelial and stromal compartments, ovarian cancers can be immunologically “cold”, “warm” or “hot”. High levels of CD8 TIL are associated with a favorable prognosis, but have not yet been shown to predict a response to PD1/PDL1 targeted immunotherapy, consistent with the presence of additional immunosuppressive factors in the tumor microenvironment.

Better understanding of immunologically inert or “cold” tumors may represent an attractive therapeutic opportunity, as they can express high levels of tumor-specific antigens with corresponding systemic T-cell and antibody responses. Following neoadjuvant chemotherapy, TIL^{high} tumors showed increases in multiple immune markers after chemotherapy; (ii) TIL^{low} tumors underwent similar increases, achieving patterns indistinguishable from the first group; and (iii) TIL^{negative} cases generally remained negative.⁶⁰ Tumor deposits with a high degree of clonal heterogeneity generally have low densities of immune infiltrates, suggesting a means by which tumor evolutionary processes are insulated from immunologic attack.⁶¹ T-cell clones track with individual tumor clones across space, suggesting that the immune system contends with intratumoral heterogeneity by battling each tumor clone individually. In particular, observations of clonally diverse primary foci present in conjunction with distal clonally pure sites could indicate local immune privilege at sites with divergent clones and active immuno-

selection at more clonally pure sites. This work is providing novel insights into the relationship between the clonal architecture of tumors and antitumor immunity. One route to deliver T-cell immunotherapy to ovarian cancer patients is to enrich naturally-occurring tumor-reactive T cells from “hot” tumors and expand them to large numbers for autologous infusion. Methods have now been developed to enrich and expand tumor-reactive tumor infiltrating (TIL) lymphocytes and to eliminate non-reactive bystander cell subsets. The subset of ovarian cancer TILs with potent antitumor activity expresses the CD137 molecule and can be enriched through magnetic sorting. This tumor-reactive TIL fraction expands in an HLA-dependent manner in presence of IL-7 and IL-15, but not IL-2. Antigen presenting cells that are genetically-modified to express co-stimulatory ligands for T-cells, such as CD137L, can expand these tumor-reactive ovarian cancer TILs to levels greater than those seen in standard IL-2 culture conditions. Based upon these and other findings, several clinical trials of TIL therapy for ovarian cancer are ongoing.

Multiple approaches are being tested to overcome the immunosuppression observed in ovarian cancers. Indoleamine-2,3-dioxygenase 1 (IDO1) expression depletes tryptophan (TRP) and enhances synthesis of immunosuppressive metabolites that can decrease the activity of checkpoint inhibitors in ovarian and other solid cancers.⁶² Administration of the IDO inhibitor INCB024360 before surgical resection of ovarian cancer reduced IDO enzyme activity, increased CD8⁺ T cell infiltration and reduced suppressive Treg cells. A second strategy to reprogram the immunosuppressive ovarian cancer microenvironment utilizes the repeated intraperitoneal administration of a human IL-12 DNA expression vector within a synthetic polyethyleneglycol-polyethyleneimine-cholesterol delivery system during neoadjuvant chemotherapy, prior to interval cytoreduction.⁶³ IP IL12 gene therapy led to a preferential increase in IL-12 and IFN- γ levels in peritoneal cavity of patients, a decrease in Tregs, an increase in CD8⁺/ratio in 60-80% patients, and a shift from naïve CD8⁺ cells to effector memory cells. A third approach to overcoming immunosuppression is to engineer T cells that not only express the T cell receptor specific for NY-ESO-1, but also a decoy receptor that renders the T cells resistant to immunosuppression by TGF β . A phase I/II clinical trial testing this approach is currently open and accruing.

For patients with immunologically “cold” ovarian cancers, lysates generated from the patient’s own tumor have been used to vaccinate against shared antigens as well as patient-specific mutated antigens.⁶⁴ In this prime-and-boost approach, a patient can first be vaccinated against patient-specific antigens to induce antitumor immunity, and the vaccine-primed T cells then harvested and expanded to high numbers outside of the body

before reinfusion into the patient. The feasibility and safety of this approach has now been established in ovarian cancer with evidence of biologic activity. Particularly promising results have been obtained using personalized vaccines generated by pulsing autologous dendritic cells (DCs) with oxidized autologous whole-tumor cell lysate (OCDL), which was injected intranodally in platinum-treated, immunotherapy-naïve, recurrent ovarian cancer patients alone or with bevacizumab with or without low dose cyclophosphamide.⁶⁵ Vaccination induced T cell responses to autologous tumor antigen, which were associated with significantly prolonged survival. Vaccination also amplified T cell responses against mutated neo-epitopes derived from non-synonymous somatic tumor mutations, and this included priming of T cells against previously unrecognized neoepitopes, as well as novel T cell clones of markedly higher avidity against previously recognized neoepitopes.

Inhibitors of DNA methyl transferase (DNMTI) can enhance expression and presentation of tumor antigens that can be recognized by the adaptive immune system, including “cancer testis antigens” such as NY-ESO-1. Adoptive transfer of HLA-A*02 restricted clones of NY-ESO-1 specific CD8 TCR gene-engineered T cells in combination with the demethylation agents decitabine and SGI-110 elicited synergistic inhibition of tumor growth, curing a fraction of mice. In the NY-ESO-1 negative OVCAR3 model, demethylation agents not only induced expression of NY-ESO-1 tumor antigen and MHC I and II, rendering the tumor visible for recognition by CD8 T cells, but also dramatically promoted persistence and accumulation of adoptively transferred T cells at tumor site, as well as reduction of suppressive myeloid cells in the tumor.⁶⁶ DNMTI rendered the tumor visible for recognition by NY-ESO-1 specific CD4 T cells leading to significant tumor inhibition, and improved the persistence of CD4 T cells at peripheral and tumor sites.

MUC16, the glycoprotein encoding the CA125 antigen, can function as an oncogene. The carboxy-terminal portion of the MUC16/CA125 protein transforms NIH/3T3 cells, increases invasive tumor properties, activates the AKT and ERK pathways, and contributes to the biologic properties of ovarian cancer. The MUC16 oncogenic effects are mediated through N-glycosylation of asparagine sites within the 58-amino-acid domain between the putative cleavage site and the cell membrane. Oncogenic signaling requires the presence of Galectin-3 and growth factor receptors co-localized on lipid rafts. With sufficient N-glycosylation and Galectin-3, MUC16 stabilizes pro-growth receptors on the cancer cell surface and enhances signaling through decreased receptor turnover. Monoclonal antibodies that block Galectin-3-mediated MUC16 interactions with cell surface signaling molecules inhibit invasion of ovarian cancer cells, directly blocking the in vivo growth of MUC16-bearing

ovarian cancer xenografts, providing a new therapeutic approach. MUC16-targeted CAR-T cells directed at the most proximal portions of MUC16 have been developed and are currently being evaluated in clinical trials.⁶⁷

Conclusions. In answering these “Critical Questions”, we have learned that screening algorithms measuring the trend of CA125 values over time can achieve adequate specificity, but we must improve the sensitivity of panels of biomarkers for early detection of ovarian cancer, possibly utilizing autoantibodies, antigen-autoantibody complexes and nucleic acids. Macrophages in the tumor microenvironment can increase resistance to anti-angiogenic therapy and cytokines, chemokines and matrix proteins can influence the influx of immunoregulatory cells. Ovarian cancer cells exhibit distinctive metabolic changes that can be targeted including a dependence on fatty acids from adipocytes, aberrant glycolytic pathways, overexpression of SIK2 and dependence on arginine in different histotypes. Heterogeneity is observed between and within ovarian cancers of the same histotype. New targets have been identified in rare histotypes and new technologies have identified diverse and potentially important subpopulations within high grade serous ovarian cancers. There is still a critical need to develop more predictive animal models, as well as to test new agents and approaches in multiple models as each may reflect the genotype and phenotype of only a single patient. Targeting pathways upregulated by individual drugs can overcome adaptive resistance. Three different PARP inhibitors have been approved for treatment of women with ovarian cancer and combinations of PARP inhibitors with PI3K inhibitors, MEK inhibitors, ATM inhibitors and CHK1 inhibitors are being evaluated to overcome PARP inhibitor resistance. Finally, immunotherapy with immune check point inhibitors has produced only a 10-15% response rate in ovarian cancers, related, in part, to the heterogeneity of immune infiltrates in tumor tissue. Novel approaches are being developed to overcome the immunosuppressive microenvironment ovarian cancers, to present autologous tumor associated antigens more effectively and to administer genetically engineered CAR –T cells. These advances promise further improvement in patient outcomes over the next years.

REFERENCES

¹ Havrilesky LJ, Sanders GD, Kulasingam S, Myers ER. Reducing ovarian cancer mortality through screening: is it possible, and can we afford it? *Gynecol Onco.* 2008; 111:179-87.

-
- ² Moss Ha, Berchuck A, Neely ML, Myers ER, Havrilesky LJ. Estimating cost-effectiveness of a multimodal ovarian cancer screening program in the United States: secondary analysis of the UK Collaborative Trial of Ovarian Cancer Screening (UKCTOCS). *JAMA Oncol* 2018; 4:190-195.
- ³ Lu, KH, Skates S, Hernandez MA, et al. A 2-stage ovarian cancer screening strategy using the Risk of Ovarian Cancer Algorithm (ROCA) identifies early-stage incident cancers and demonstrates high positive predictive value. *Cancer*. 2013; 119:3454-3461.
- ⁴ Jacobs IJ, Menon U, Ryan A, et al. Ovarian cancer screening and mortality in the UK Collaborative Trial of Ovarian Cancer Screening (UKCTOCS): a randomised controlled trial. *Lancet*. 2016; 387:945-956.
- ⁵ Yang WL, Gentry-Maharaj A, Simmons A, et al. Elevation of TP53 Autoantibody Prior to CA125 in Preclinical Invasive Epithelial Ovarian Cancer. *Clin Cancer Res*. 2017; 23:5912-5922.
- ⁶ Wang Y, Li L, Douville C, et al. Evaluation of liquid from the papanicolaou test and other liquid biopsies for the detection of endometrial and ovarian cancers. *Sci Transl Med*. 2018; 10:433.
- ⁷ Elias KM, Fendler W, Stawiski K, et al. Diagnostic potential for a serum miRNA neural network for detection of ovarian cancer. *Elife* 2017; 31:6
- ⁸ Sung HK, Ma SH, Choi JY, et al. The effect of breastfeeding duration and parity on the risk of epithelial ovarian cancer: A systematic review and meta-analysis. *J Prev Med Public Health*. 2016; 49:349-66.
- ⁹ Lyons YA, Pradeep S, Wu SY, et al. Macrophage depletion through colony stimulating factor 1 receptor pathway blockade overcomes adaptive resistance to anti-VEGF therapy. *Oncotarget*. 2017; 8:96496-96505.
- ¹⁰ Noh K, Mangala LW, Han HD, et al. Differential effects of EGFL6 on tumor versus wound angiogenesis. *Cell Rep*. 2017; 21:2785-2795.
- ¹¹ Pearce OMT, Delaine-Smith RM, Maniati E, et al. Deconstruction of a metastatic tumor microenvironment reveals a common matrix response in human cancers. *Cancer Discovery*. 2018; 8:304-319.
- ¹² Eckert MA, Coscia F, Chryplewicz A, et al. Nicotinamide N-methyltransferase metabolically reprograms the stroma to promote ovarian cancer metastasis. *Proceedings of an AACR Special Conference on Addressing Critical Questions in Ovarian Cancer Research and Treatment*. 2017; PR05
- ¹³ Ladanyi A, Mukherjee A, Kenny HA, et al. Adipocyte-induced CD36 expression drives ovarian cancer progression and metastasis. *Oncogene*. 2018; 37:2285-2301.

-
- ¹⁴ Muggia FM, Braly Ps, Brady MF, et al. Phase III randomized study of cisplatin versus paclitaxel versus cisplatin and paclitaxel in patients with suboptimal stage III or IV ovarian cancer: a gynecologic oncology group study. J Clin Oncol 2000; 18:106-15.
- ¹⁵ Miranda F, Mannion D, Liu S, et al. Salt-Inducible Kinase 2 Couples Ovarian Cancer Cell Metabolism with Survival at the Adipocyte-Rich Metastatic Niche. Cancer Cell 2016; 30:273-289.
- ¹⁶ Ahmed AA, Lu Z, Jennings NB, et al. SIK2 is a centrosome kinase required for bipolar mitotic spindle formation that provides a potential target for therapy in ovarian cancer. Cancer Cell. 2010; 18:109-121.
- ¹⁷ Zhou J, Alfraidi A, Zhang S, et al. A novel compound ARN-3236 inhibits Salt Inducible Kinase 2 and sensitizes ovarian cancer cell lines and xenografts to paclitaxel. Clin Cancer Res. 2017; 23:1945-1954.
- ¹⁸ Cheon DJ, Walts AE, Beach JA, et al. Differential expression of argininosuccinate synthetase in serous and non-serous ovarian carcinomas. J Pathol Clin Res. 2014; 1:41-53.
- ¹⁹ Labidi-Galy SI, Papp E, Hallberg D, et al. High grade serous ovarian carcinomas originate in the fallopian tube. Nat Commun. 2017; 8:1093.
- ²⁰ Macintyre G, Goranova TE, De Silva D, et al. Copy number signatures and mutational processes in ovarian carcinomas. Nature Genetics 2018; 50: 1262-1270.
- ²¹ Francis K, Burston H, Mokady D, et al. Functional genetic architecture of serous ovarian cancer. Proceedings of an AACR Special Conference on Addressing Critical Questions in Ovarian Cancer Research and Treatment. 2017; IA27.
- ²² Gonzalez VD, Smusik N, Chen TJ, et al Commonly occurring cell subsets in high grade serous ovarian tumors identified by single cell mass spectrometry. Cell Reports 2018; 22:1875-1888.
- ²³ Domcke S, Sinha R, Levine DA, Sander C, Schultz N. Evaluating cell lines as tumour models by comparison of genomic profiles. Nat Commun. 2013; 4:2126.
- ²⁴ Walton JB, Blagih J, Ennis D, et al. CRISPR/Cas9-Mediated Trp53 and Brca2 Knockout to Generate Improved Murine Models of Ovarian High-Grade Serous Carcinoma. Cancer Res. 2016; 76:6118-6129.
- ²⁵ Walton JB, Farquharson M, Mason S, et al. CRISPR/Cas9-derived models of ovarian high grade serous carcinoma targeting Brca1, Pten and Nf1, and correlation with platinum sensitivity. Sci Rep. 2017; 7:16827.

²⁶ Magnotti E, Marasco WA. The latest animal models of ovarian cancer for novel drug discovery. *Expert Opin Drug Discov* 2018; 13:249-257.

1	
---	--

²⁷ Gershenson DM, Bodurka DC, Lu KH, et al. Impact of Age and Primary Disease Site on Outcome in Women with Low-Grade Serous Carcinoma of the Ovary or Peritoneum: Results of a Large Single-Institution Registry of a Rare Tumor. *J Clin Oncol*. 2015; 33:2675-82.

²⁸ Dalton HJ, Fleming ND, Sun CC, Bhosale P, Schmeler KM, Gershenson DM. Activity of bevacizumab-containing regimens in recurrent low-grade serous ovarian or peritoneal cancer: A single institution experience. *Gynecol Oncol*. 2017; 145:37-40.

²⁹ Gershenson DM, Bodurka DC, Coleman RL, Lu KH, Malpica A, Sun CC. Hormonal Maintenance Therapy for Women With Low-Grade Serous Cancer of the Ovary or Peritoneum. *J Clin Oncol*. 2017; 35:1103-1111.

³⁰ Farley J, Brady WE, Vathipadiekal V, et al. Selumetinib in women with recurrent low-grade serous carcinoma of the ovary or peritoneum: an open-label, single-arm, phase 2 study. *Lancet Oncol*. 2013; 14:134-40.

³¹ Jelinic P, Ricca J, Van Oudenhove E, et al, Immune-active microenvironment in small cell carcinoma of the ovary, hypercalcemic type: rationale for immune check point blockade. *J Natl Cancer Inst* 2018; 110: 787-790.

³² Yanagida S, Anglesio MS, Nazeran TM, et al. Clinical and genetic analysis of recurrent adult-type granulosa cell tumor of the ovary: Persistent preservation of heterozygous c.402C>G FOXL2 mutation. *PLoS One*. 2017; 12: e0178989.

³³ Ennis D, Cooke SL, Evers L, et al. The driver mutational landscape of ovarian squamous cell carcinomas arising in mature cystic teratoma. *Proceedings of an AACR Special Conference on Addressing Critical Questions in Ovarian Cancer Research and Treatment*. 2017; PR11.

³⁴ Sun C, Fang Y, Yin J, et al. Rational combination therapy with PARP and MEK inhibitors capitalizes on therapeutic liabilities in RAS mutant cancers. *Sci Transl Med*. 2017; 9:392.

³⁵ Sun C, Yin J, Yin J, et al. BRD4 inhibition is synthetic lethal with PARP inhibitors through the induction of homologous recombination deficiency. *Cancer Cell*. 2018; 33:401-416.

³⁶ Zervantonakis IK, Iavarone C, Chen HY, et al. Systems analysis of apoptotic priming in ovarian cancer identifies vulnerabilities and predictors of drug response. *Nat Commun*. 2017; 8:365.

³⁷ Phallen J, Sausen M, Adleff V, et al. Direct detection of early-stage cancers using circulating tumor DNA. *Sci Transl Med*. 2017; 9:403.

³⁸ Jiang Q, Greenberg RA. Deciphering the BRCA1 Tumor Suppressor Network. *J Biol Chem.* 2015; 290:17724-32.

³⁹ Ledermann J, Harter P, Gourley C, et al. Olaparib maintenance therapy in platinum-sensitive relapsed ovarian cancer. *N Engl J Med.* 2012; 366:1382-92.

⁴⁰ Kaufman B, Shapira-Frommer R, Schmutzler RK, et al. Olaparib monotherapy in patients with advanced cancer and a germline BRCA1/2 mutation. *J Clin Oncol.* 2015; 33:244-50.

⁴¹ Pujade-Lauraine E, Ledermann JA, Selle F, et al. Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): a double-blind, randomised, placebo-controlled, phase 3 trial. *Lancet Oncol.* 2017; 18:1274-1284.

⁴² Mirza MR, Monk BJ, Herrstedt J, et al. Niraparib Maintenance Therapy in Platinum-Sensitive, Recurrent Ovarian Cancer. *N Engl J Med.* 2016; 375:2154-2164.

⁴³ Swisher EM, Lin KK, Oza AM, et al. Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): an international, multicentre, open-label, phase 2 trial. *Lancet Oncol.* 2017; 18:75-87.

⁴⁴ Lord CJ, Ashworth A. Mechanisms of resistance to therapies targeting BRCA-mutant cancers. *Nat Med.* 2013; 19:1381-8.

⁴⁵ Liu JF, Barry WT, Birrer M, et al. Combination cediranib and olaparib versus olaparib alone for women with recurrent platinum-sensitive ovarian cancer: a randomised phase 2 study. *Lancet Oncol.* 2014; 15:1207-14.

⁴⁶ Matulonis UA, Wulf GM, Barry WT, et al. Phase I dose escalation study of the PI3kinase pathway inhibitor BKM120 and the oral poly (ADP ribose) polymerase (PARP) inhibitor olaparib for the treatment of high-grade serous ovarian and breast cancer. *Ann Oncol.* 2017; 28:512-518.

⁴⁷ Yazinski SA, Comaills V, Buisson R, et al. ATR inhibition disrupts rewired homologous recombination and fork protection pathways in PARP inhibitor-resistant BRCA-deficient cancer cells. *Genes Dev.* 2017; 31:318-332.

⁴⁸ Schlacher K, Christ N, Siaud N, Egashira A, Wu H, Jasin M. Double-strand break repair-independent role for BRCA2 in blocking stalled replication fork degradation by MRE11. *Cell.* 2011; 145:529-42.

⁴⁹ Scully R, Livingston D. In search of the tumour-suppressor functions of BRCA1 and BRCA2. *Nature.* 2000; 408:429-432.

-
- ⁵⁰ Ray Chaudhuri A, Callen E, Ding X, et al. Replication fork stability confers chemoresistance in BRCA-deficient cells. *Nature*. 2016; 535:382-7.
- ⁵¹ Toledo LI, Altmeyer M, Rask MB, et al. ATR prohibits replication catastrophe by preventing global exhaustion of RPA. *Cell*. 2013; 155:1088-103.
- ⁵² Kim H, George E, Ragland R, et al. Targeting the ATR/CHK1 Axis with PARP Inhibition Results in Tumor Regression in BRCA-Mutant Ovarian Cancer Models. *Clin Cancer Res*. 2017; 23:3097-3108.
- ⁵³ Wielgos-Bonvallet M, Cybulska P, Oudenhove EV, et al. Exploring the effects of PARP inhibition on CHK1 activation as a potential determinant of synergy with CHK1 inhibition. *Proceedings of an AACR Special Conference on Addressing Critical Questions in Ovarian Cancer Research and Treatment*. 2017; PR02.
- ⁵⁴ Etemadmoghadam D, Weir BA, Au-Yeung G, et al. Synthetic lethality between CCNE1 amplification and loss of BRCA1. *Proc Natl Acad Sci U S A*. 2013; 110:19489-94.
- ⁵⁵ Godin SK, Sullivan MR, Bernstein KA. Novel insights into RAD51 activity and regulation during homologous recombination and DNA replication. *Biochem Cell Biol*. 2016; 94:407-418.
- ⁵⁶ Nielsen FC, van Overeem Hansen T, Sorensen CS. Hereditary breast and ovarian cancer: new genes in confined pathways. *Nat Rev Cancer*. 2016; 16:599-612.
- ⁵⁷ Harding SM, Benci JL, Irianto J, Discher DE, Minn AJ, Greenberg RA. Mitotic progression following DNA damage enables pattern recognition within micronuclei. *Nature*. 2017; 548:466-470.
- ⁵⁸ Minn AJ, Wherry EJ. Combination Cancer Therapies with Immune Checkpoint Blockade: Convergence on Interferon Signaling. *Cell*. 2016; 165:272-5 (2016).
- ⁵⁹ Hatch EM, Fischer AH, Deerinck TJ, Hetzer MW. Catastrophic nuclear envelope collapse in cancer cell micronuclei. *Cell*. 2013; 154:47-60.
- ⁶⁰ Lo CS, McPherson A, Milne K, et al. Neoadjuvant chemotherapy of ovarian cancer results in three patterns of tumor-infiltrating lymphocyte response with distinct implications for immunotherapy. *Clin Cancer Res*. 2017 Feb 15; 23(4):925-934.
- ⁶¹ Zhang AW, McPherson A, Milne K, et al. Interfaces of Malignant and Immunologic Clonal Dynamics in Ovarian Cancer. *Cell*. 2018; 173:1755-1769.
- ⁶² Amobi A, Qian F, Lugade AA, Odunsi K. Tryptophan Catabolism and Cancer Immunotherapy Targeting IDO Mediated Immune Suppression. *Adv Exp Med Biol*. 2017; 1036:129-144.

⁶³ Odunsi K, Immunotherapy in ovarian cancer. 1 Ann Oncol. 2017;28(suppl_8):viii1-viii7

⁶⁴ Rodriguez Garcia A, Minutolo NG, Robinson JM, Powell DJ. T-cell target antigens across major gynecologic cancers. Gynecol Oncol. 2017 Jun; 145(3):426-435.

⁶⁵ Tanyi JL, Bobisse S, Ophir E, et al. Personalized cancer vaccine effective mobilizes T cell antitumor immunity in ovarian cancer. Science Trans Med 2018; 10:1-14.

⁶⁶ Shen L, McGray B, Miliotto A, et al. Epigenetic reprogramming promotes an immunogenic ovarian cancer tumor microenvironment and synergizes with adoptive transfer of engineered T cells expressing NY-ESO-1 specific T cell receptors. Proceedings of an AACR Special Conference on Addressing Critical Questions in Ovarian Cancer Research and Treatment. 2017; PR12

⁶⁷ Yeku OO, Purdoin TJ, Koneru M, et al. Armored CAR T cells enhance antitumor efficacy and overcome the tumor microenvironment. Sci Rep. 2017 ;7(1):10541.