

Presumed filter-feeding in a deep-sea benthic shrimp (Decapoda, Caridea, Stylodactylidae), with records of the deepest occurrence of carideans

Mary Wicksten¹, Sammy De Grave², Scott France³, Christopher Kelley⁴

1 Department of Biology, Texas A&M University, College Station Texas 77843-3258, USA **2** Oxford University Museum of Natural History, Parks Road, Oxford, OX1 3PW, United Kingdom **3** Department of Biology, University of Louisiana at Lafayette, P.O. Box 43602, Lafayette, Louisiana, 70504 USA **4** Hawaii Undersea Research Laboratory, University of Hawaii, Honolulu, Hawaii 96822 USA

Corresponding author: Mary Wicksten (Wicksten@bio.tamu.edu)

Academic editor: I. Wehrtmann | Received 29 October 2016 | Accepted 3 January 2017 | Published 17 January 2017

<http://zoobank.org/D217CEED-CCE5-4BB8-A3C9-D3A173643E82>

Citation: Wicksten M, De Grave S, France S, Kelley C (2017) Presumed filter-feeding in a deep-sea benthic shrimp (Decapoda, Caridea, Stylodactylidae), with records of the deepest occurrence of carideans. ZooKeys 646: 17–23. <https://doi.org/10.3897/zookeys.646.10969>

Abstract

Using the remotely operated vehicle *Deep Discoverer*, we observed a large stylodactylid shrimp resting on a sedimented sea floor at 4826 m in the Marianas Trench Marine National Monument. The shrimp was not collected but most closely resembled *Bathystylodactylus bathyalis*, known previously only from a single broken specimen. Video footage shows the shrimp facing into the current and extending its upraised and fringed first and second pereopods, presumably capturing passing particles. The video footage is the first ever to show a living deep-sea stylodactylid and constitutes the deepest record for the family. We provide a list of the deepest reports of caridean shrimps world-wide.

Keywords

Deep-sea shrimp, Stylodactylidae, feeding, carideans, Marianas Trench

Introduction

Benthic caridean shrimps living at 1000 m or more are poorly known, for the most part represented by specimens taken by trawls. To the best of our knowledge, none of them has been brought to the surface alive. Often, the fragile pereopods and antennae are broken or torn off. The feeding modes and form of the appendages of vent shrimps (family Alvinocarididae) have been studied in detail (Komai and Segonzac 2008). Other carideans living on mud or hard substrates away from vents are poorly studied. It is difficult to maneuver collecting equipment among rocks without entanglement. Shrimps living on mud may be difficult to see because they burrow into the sediment or dart away if disturbed. In situ photographs often do not show the shrimp in sufficient detail for identification.

There are three species of *Bathystylodactylus*, the deepest known members of the family Stylodactylidae. These are known from six specimens: one each of *B. inflatus* from off Taiwan and *B. bathyalis* from the Coral Sea and four of *B. echinus* from the eastern Pacific (Hanamura and Takeda 1996; Cleve 1994; Wicksten and Martin 2004). All known specimens are damaged, missing at least some of the appendages. We report here on a living shrimp photographed in situ, provide the deepest report of a stylodactylid shrimp to date, and compare the depth records of the deepest known carideans.

Methods

We obtained photographs and video of the shrimp from the U.S. National Oceanographic and Atmospheric Administration (NOAA) Ship *Okeanos Explorer* (*OkEx*), administered through the Office of Exploration and Research. The ship is equipped with a 6,000 m-rated dual-body system comprising the remotely-operated vehicle (ROV) *Deep Discoverer* (*D2*) and *Seirios* camera sled. During operations, *Seirios* is tethered to the ship with a standard oceanographic armored, fiber-optic cable (1.73 cm diameter) and *D2* is linked to *Seirios* with a neutrally buoyant tether, thus isolating the ROV from the *OkEx* surface motion to allow precise maneuvering and steady imaging of deep-sea communities.

The ROV *D2* is outfitted with two maneuverable and four fixed video cameras; scientific observations are made primarily using two high-definition video cameras. Light is supplied by 26 LED lamps (195000 lumens total), with eight of these on four hydraulically positioned booms. Paired lasers (10 cm apart) mounted on the fixed, high-definition video camera provide size scale in the imagery. The ROV traversed the seafloor at a speed of approximately ~ 0.1–0.3 knots (1 knot = 0.514 m s⁻¹) with the cameras generally set on wide-angle view, and zooms were initiated to obtain detailed imagery when objects of scientific or aesthetic interest were encountered. High-definition video was transmitted from the *D2* in HD-SDI 1080i format. *OkEx* is equipped with high-speed communication capabilities to enable scientists on the shore to participate in ship operations in time via telepresence (see <http://oceanexplorer.noaa.gov/>

okeanos/collaboration-tools/welcome.html). Scientists (and anyone with an Internet connection) are able to observe live video feeds from the ROV *D2* and participate in real-time via a private Internet chat room and satellite teleconference line.

Results

From 20 April to 10 July, 2016, the *OkEx* was engaged in the “2016 Exploration of the Marianas” Expedition, a baseline study of deep-water environments in and around the Commonwealth of the Northern Mariana Islands and the Mariana Trench Marine National Monument (MTMNM). *D2* Dive 13 of Leg 3 (30 June 2016) of this expedition explored a site informally named “Twin Peaks” on a seamount in the MTMNM. The *D2* spent 4 hours 48 min traversing the bottom, from 4840 meters depth upslope to 4787 m. The seafloor was mostly thickly sedimented with clay-like particles that were easily disturbed into clouds by the ROV thrusters; small outcroppings of sedimentary rocks were seen frequently as well as occasional large boulders.

Approximately 2 hours 17 min after the ROV reached the bottom and the benthic exploration began, an observation was made of a single individual stylodactylid shrimp (estimated total length 120 mm) at 4826 m depth (21.41774° N, 145.89294° E). No other stylodactylids were observed during the dive or on other dives during this expedition. Other benthic shrimps (Superfamily Penaeoidea) and thread-leg shrimps (Caridea: Nematocarcinidae) were observed on soft substrates, as well as other carideans associated with soft corals (Order Gorgonacea) and sponges (Hexactinellida). When initially observed, the shrimp was facing away from the ROV camera. Participating scientists noted this was an atypical shrimp species for the expedition, so the ROV settled to capture detailed images. After a couple of minutes the ROV was repositioned to get a lateral view of the shrimp. During the approximately 4 minutes of observation (including the ROV maneuver) the shrimp did not move from its initial position on the bottom, facing into the weak boundary layer current. The following environmental data were recorded during the observation: temperature 1.47236°C, salinity 34.69294 PSU, dissolved oxygen 4.96223 mg/L.

With anterior legs upraised, the shrimp faced into the current, presumably using its legs as a net to capture passing particles. The shrimp had first and second pereopods fringed with setae and with extremely slender chelae, characteristic of the family Stylodactylidae (Cleva 1990). The deepest species, belonging to the genus *Bathystylodactylus*, have a long toothed rostrum, eyes without pigment, and a carapace that is pubescent or studded with minute spinules (Hanamura and Takeda 1996, Wicksten and Martin 2004). The shrimp in the video appeared to most closely resemble *Bathystylodactylus bathyalis*, previously only known from a single broken specimen collected at 3515–3502 m in the Coral Sea (Cleva 1994). The present video footage is the first for any of the deep-water dwelling Stylodactylidae. Other invertebrates in the area included a large long-legged isopod (family Munnopsidae), an enteropneust (phylum Hemichordata), a swimming holothurian (Echinodermata), and a hermit crab

(*Parapagurus* sp.) with an associated commensal sea anemone (Actiniaria). The ROV was not equipped with collecting equipment that could capture any of these animals and so their species identification remains uncertain.

As in the euphausiaceans, the long, setose legs of the stylodactylid seem to form a “filter basket” that captures particles. The shrimp in the video was not seen to open and expel water by means of pumping, as can euphausiaceans, but instead relied on passive transfer of food particles by the boundary layer current. No other group of carideans is known to feed in this manner (Wicksten 2010). Other decapods filter-feed by means of long feathery antennae, as in the hermit crabs of the genus *Paguritta* and the mole crabs, family Hippidae; or setose third maxillipeds, as in crabs of the family Porcellanidae (Riisgard and Larsen 2010).

Discussion

Relatively few caridean shrimps live at bathyal depths, i.e. greater than 1000 m, with far fewer recorded at abyssal depths of 3000 m and deeper. As well as species of the Stylodactylidae, caridean shrimps reported from abyssal depths include species of the families Bythocarididae, Crangonidae, Disciadidae, Nematocarcinidae, Oplophoridae, Pandalidae and Pasiphaeidae (Table 1). The present observation of *Bathystylodactylus* sp. is the deepest report of a member of this family. The oplophorid *Heterogenys microphthalmus* has been collected in trawls as deep as 5060 m (Crosnier 1987). Jamieson et al. (2009) included photographs of “*Acantheephyra* sp.” from baited traps as deep as



Figure 1. *Bathystylodactylus* cf. *bathyalis*, 4826 m, in situ. Photo extracted from high-definition video captured using ROV *Deep Discoverer*. (Image courtesy of the NOAA Office of Ocean Exploration and Research 2016 Deepwater Exploration of the Marianas).

Table 1. Deepest family level records for caridean shrimp species living at depths bathyal and abyssal depths. Systematics follows De Grave & Fransen (2011), with modifications in De Grave et al. (2014, 2015). Depths as well as sampling or observation method are given.

Acanthephyridae
<i>Acanthephyra quadrispinosa</i> Kemp, 1939: Crosnier (1987), 5040–5060 m, trawl.
<i>Heterogenys microphthalma</i> (Smith, 1885): Crosnier (1987), 5040–5060 m, trawl.
Alvinocarididae
<i>Rimicaris hybisae</i> Nye, Copley & Plouviez, 2012: 4960 m, TV grab and slurp gun.
Bathypalaemonellidae
<i>Bathypalaemonella serratipalma</i> Pequegnat, 1970: Cleva (2001), 2660–2750 m, beam trawl.
Bresiliidae
<i>Bresilia pacifica</i> Hendrickx, 2014: 2010–2046 m, benthic sledge.
Bythocarididae
<i>Bythocaris cryonesus</i> Bowman & Manning, 1973: 3803 m, minnow trap.
Crangonidae
<i>Placopsicrangan formosa</i> Komai & Chan, 2009: 4807–4824, beam trawl.
Disciadiidae
<i>Lucaya bigelowi</i> Chace, 1939: 4773 m, open net.
Glyphocrangonidae
<i>Glyphocrangon atlantica</i> Chace, 1939: Holthuis (1971), 6364–6373 m, collecting method not specified.
Hippolytidae
<i>Leontocaris amplexipes</i> Bruce, 1990: Ahyong (2010), 2182–2119 m, sled.
Nematocarcinidae
<i>Nematocarcinus challengerii</i> Burukovsky, 2006: 5477 m, trawl.
Oplophoridae
<i>Systellaspis debilis</i> (A. Milne Edwards, 1881): Crosnier (1987), 4987–5025 m, trawl.
Palaemonidae
<i>Periclimenes pholeter</i> Holthuis, 1973: Bruce (2011), 2148 m, collecting method not specified.
Pandalidae
<i>Stylopandalus richardi</i> (Coutière, 1905): Hayashi & Miyake (1969), 3600 m, midwater trawl.
Pasiphaeidae
<i>Parapasiphae compta</i> Smith, 1884: Crosnier (1988), 4990 m, trawl.
Physetocarididae
<i>Physetocaris microphthalma</i> Chace, 1940: Wasmer (1985), 1200–2200 m, non-closing trawl.
Psalidopodidae
<i>Psalidopus tosaensis</i> Toriyama & Horikawa, 1993: 2765–2881 m, beam trawl.
Stylodactylidae
<i>Bathystylodactylus</i> sp.: present report, 4820 m, video observation.
Thoridae
<i>Lebbeus laurentae</i> Wicksten, 2010: Komai et al. (2012), 2640 m, slurp gun.

6890 m in the Kermadec Trench, but their photographs (Fig. 1B, C) show a shrimp with a short upturned rostrum, characteristic of *H. microphthalma*. If confirmed this would be the deepest recorded caridean shrimp record, a record currently held by *Glyphocrangon atlantica* at 6364–6373 m (Holthuis 1971).

Video link: <http://oceanexplorer.noaa.gov/oceanos/explorations/ex1605/dailyupdates/media/video/0630-mudmonsters/0630-mudmonsters.html>

References

- Ahyong ST (2010) New species and new records of Caridea (Hippolytidae: Pasiphaeidae) from New Zealand. In: De Grave S, Fransen CHJM (Eds) Contributions to shrimp taxonomy. Zootaxa 2372: 341–357.
- Bowman TE, Manning RB (1973) Two Arctic bathyal crustaceans: the shrimp *Bythocaris cryonensis* new species, and the amphipod *Eurythenes gryllus*, with in situ photographs from Ice Island T-3. Crustaceana 23: 187–201. <https://doi.org/10.1163/156854072X00363>
- Bruce AJ (2011) A new record of *Periclimenes pholeter* Holthuis, 1973 (Crustacea: Decapoda: Pontoniinae) from the Red Sea. Cahiers de Biologie Marine 52: 119–120.
- Burukovsky RN (2006) Systematics of shrimps of the genus *Nematocarcinus* (Decapoda, Nematocarcinidae). Description of the type specimen, *N. paucidentatus* and some unidentified species of the family Nematocarcinidae from the collection of the British Museum (samples from the “Challenger”). Zoologicheskii Zhurnal 85: 896–900. [In Russian]
- Chace FA Jr (1939) Reports on the scientific results of the first Atlantis expedition to the West Indies, under the joint auspices of the University of Havana and Harvard University. Preliminary descriptions of one new genus and seventeen new species of decapod and stomatopod Crustacea. Memorias de la Sociedad Cubana de Historia Natural 13: 31–54.
- Cleva R (1990) Crustacea Decapoda: Les genres et les espèces indo-ouest pacifiques de Styrodactylidae. In: Crosnier A (Ed.) Résultats des Campagnes MUSORSTOM. Vol. 6. Memoires du Muséum National d’Histoire Naturelle (A) 145: 71–136.
- Cleva R (1994) Some Australian Styrodactylidae (Crustacea: Decapoda), with descriptions of two new species. The Beagle 11: 53–64.
- Cleva R (2001) Les Bathypalaemonellidae de Saint-Laurent, 1985 (Crustacea, Decapoda, Caridea) avec description d’une espèce nouvelle et définition d’un genre nouveau. Zoosystema 23: 757–782.
- Crosnier A (1987) Oplophoridae (Crustacea Decapoda) récoltés de 1971 à 1982 par les navires françaises dans l’océan Indien occidental sud. Bulletin du Muséum national d’Histoire naturelle, Paris, 4ème série, section A, Zoologie, Biologie et Écologie animales 9: 695–726.
- Crosnier A (1988) Sur la présence de *Parapasiphae compta* Smith, 1884, dans l’Atlantique oriental (Crustacea Decapoda Pasiphaeidae). Bulletin du Muséum national d’Histoire naturelle, Paris, 1988; series 4, 799–803.
- De Grave S, Fransen CHJM (2011) Carideorum Catalogus: The recent species of the dendrobranchiate, stenopodidean, procarididean and caridean shrimps (Crustacea: Decapoda). Zoologische Mededelingen 85: 195–589.
- De Grave S, Fransen CHJM, Page TJ (2015) Let’s be pals again: major systematic changes in Palaemonidae (Crustacea: Decapoda). PeerJ 3: e1167. <https://doi.org/10.7717/peerj.1167>
- De Grave S, Li CP, Tsang LM, Chu KH, Chan TY (2014) Unweaving hippolytoid systematics (Crustacea, Decapoda, Hippolytidae): resurrection of several families. Zoologica Scripta 43: 496–507. <https://doi.org/10.1111/zsc.12067>
- Hanamura Y, Takeda M (1996) Establishment of a new genus *Bathystylodactylus* (Crustacea: Decapoda: Stylodactylidae) with description of a new species from northwestern Pacific. Zoological Science 13: 929–934. <https://doi.org/10.2108/zsj.13.929>

- Hayashi KI, Miyake S (1969) Bathypelagic caridean shrimps collected by “Koyo Maru” during the International Indian Ocean Expedition. Ohmu. Occasional Papers of Zoological Laboratory, Faculty of Agriculture, Kyushu University, Fukuoka, Japan 2: 59–77.
- Hendrickx ME (2014) First record of the caridean shrimp genus *Bresilia* Calman, 1896 (Crustacea: Decapoda: Bresiliidae) from the East Pacific and description of a new species. *Zootaxa* 3878: 185–195. <https://doi.org/10.11646/zootaxa.3878.2.5>
- Holthuis LB (1971) Biological results of the University of Miami deep-sea Expeditions. 75. The Atlantic shrimps of the deep-sea genus *Glyphocrangon* A. Milne Edwards, 1881. *Bulletin of Marine Science* 21: 267–373.
- Jamieson AJ, Fujii T, Solan M, Matsumoto AK, Bagley PM, Priede IG (2009) First findings of decapod crustacean in the hadal zone. *Deep-Sea Research I* 56: 641–647. <https://doi.org/10.1016/j.dsr.2008.11.003>
- Komai T, Chan TY (2009) New genus and species of Crangonidae (Decapoda: Caridea) with a large plate-like eye from the abyssal zone off Taiwan, northwestern Pacific. *Journal of Crustacean Biology* 29: 254–265. <https://doi.org/10.1651/08-3080R.1>
- Komai T, Segonzac M (2008) Taxonomic review of the hydrothermal vent shrimps genera *Rimicaris* Williams & Rona and *Chorocaris* Martin & Hessler (Crustacea: Decapoda: Caridea: Alvinocarididae). *Journal of Shellfish Research* 27: 21–41. [https://doi.org/10.2983/0730-8000\(2008\)27\[21:TROTHV\]2.0.CO;2](https://doi.org/10.2983/0730-8000(2008)27[21:TROTHV]2.0.CO;2)
- Komai T, Tsuchida S, Segonzac M (2012) Records of species of the hippolytid genus *Lebbeus* White, 1847 (Crustacea: Decapoda: Caridea) from hydrothermal vents in the Pacific Ocean, with descriptions of three new species. *Zootaxa* 3241: 35–63.
- Nye V, Copley J, Plouviez S (2012) A new species of *Rimicaris* (Crustacea: Decapoda: Caridea: Alvinocarididae) from hydrothermal vent fields on the Mid-Cayman spreading centre, Caribbean. *Journal of the Marine Biological Association of the United Kingdom* 92: 1057–1072. <https://doi.org/10.1017/S0025315411002001>
- Riisgard HU, Larsen PS (2010) Particle capture mechanisms in suspension-feeding invertebrates. *Marine Ecology Progress Series* 418: 255–293. <https://doi.org/10.3354/meps08755>
- Toriyama M, Horikawa H (1993) A new caridean shrimp, *Psolidopus tosaensis*, from Tosa Bay, Japan (Decapoda: Caridea: Psalidopodidae). *Bulletin of the Nansei National Fishery Research Institute* 26: 1–8.
- Wasmer RA (1985) New records for *Physetocaris microphthalma* Chace (Decapoda, Caridea, Physetocarididae) from the South Pacific. *Crustaceana* 49: 315–318. <https://doi.org/10.1163/156854085X00648>
- Wicksten MK (2010) Infraorder Caridea Dana, 1852. Chapter 64 In: Schram FR, Von Vaupel Klein JC (Eds) *Treatise on Zoology: the Crustacea*. Vol. 9 Part A. Eucarida: Euphausiacea, Amphionidacea, and Decapoda (partim): 163–206.
- Wicksten MK, Martin JW (2004) A new species of caridean shrimp of the family Stylodactylidae from the eastern Pacific Ocean. *Proceedings of the Biological Society of Washington* 117: 377–384.