

Odour hedonics and the ubiquitous appeal of vanilla

Charles Spence

Crossmodal Research Laboratory, Oxford University, Oxford OX2 6GG, UK;

charles.spence@psy.ox.ac.uk

ABSTRACT

Our food choices and consumption behaviours are often influenced by odour hedonics, especially in the case of those orthonasally-experienced aromas (i.e., those odours that are food-related). The origin(s) of odour hedonics remains one of the most intriguing in olfactory science and, over the years, several fundamentally different accounts have been put forward to try and explain the varying hedonic responses that people have to a wide range of odorants. Associative learning, innate, and molecular accounts of odour pleasantness have all been suggested. Here, the origins of the hedonic response to vanilla which is one of the most liked smells cross-culturally is explored. The history of vanilla's use in food and medicine is overviewed, with a focus on its neurocognitive appeal. While vanilla is one of the most widely liked aromas, it is also rated as smelling sweet to most people. Food scientists are becoming increasingly interested in the possibility that such 'sweet smells' could be used to help maintain the sweetness of commercial food products while, at the same time, reducing the use of calorific sweeteners. Such an approach is likely to be facilitated by the low cost of artificial vanilla flavouring (when compared to the high and fluctuation price of natural vanilla pods).

Introduction

Although the importance of the sense of smell has been downplayed by cognitive neuroscientists over the last 150 years or so¹, olfaction undoubtedly plays an important role in well-being^{2,3} and interpersonal interaction/attraction^{4,5}. It is our sense of smell that helps us to detect danger and off-taints in food and drink^{6,7} and orthonasal olfactory cues also play a crucial role in food choice as well^{8,9}. At the same time, however, olfactory stimuli are rarely experienced in isolation and are typically experienced in combination with other senses (e.g., as when eating and drinking). That said, the role of olfaction in humans is probably supportive rather than leading, especially given the ubiquity of visual dominance¹⁰. It should be noted here that visual cues have also been shown to dominate over our perception of the flavour of food and drink^{11,12}.

It has been suggested that all smell sensations carry an element of pleasure or pain and, indeed, hedonics may be one of the primary, or dominant, responses that people have to odorants¹³⁻¹⁵. In fact, the close link between emotion and olfaction is likely related to the shared neural substrates involved, including brain structures such as the amygdala, hippocampus, insula, anterior cingulate cortex and orbitofrontal cortex¹⁶. However, researchers would appear to disagree with regards to the question of where such differences in odour hedonics originate. One argument is that our hedonic responses to odours are mostly acquired as a result of associative learning (i.e., via prior experience with odours; e.g., in the context of foods¹⁷; i.e., we learn to like the smell of vanilla because of the sweet taste with which it is so often paired in food and drink (even though the fermented vanilla pod itself tastes very bitter). Similarly, we learn to like the smell of coffee because of what it is normally associated with (i.e., caffeine); And we learn to like the ‘new car’ smell¹⁸⁻²⁰, not because of anything unique about the molecular structure of the volatile organic chemicals *per se*, but rather because this olfactory stimulus happens to be associated with a highly-rewarding experience³. Another account for our response to certain evolutionarily important odorants, such as those found in blood, is that they are innate, and likely preserved across species^{21,cf.22}. One might at least consider the possibility that the sweetness that humans associate with floral scents represents such a food-related preserved response (see ²³ for other suggestions regarding our innate response to certain food odorants). However, according to an increasingly-popular alternative account, olfactory pleasantness (at least in the case of unfamiliar monomolecular odorants) can sometimes be predicted on the basis of the structure

of the molecular odorants themselves. According to Joussain and colleagues (2011, p. 563)²⁴, this link between the physicochemical structure of an odorant and hedonics “may already be engraved” at the level of the olfactory receptor neurons in the olfactory epithelium. The question of the relationships between odour hedonics (and their origin) and food behaviour is of widespread interest given its applicability to food scientists/industry etc. Furthermore, gaining a better understanding of the mechanisms behind the positive odour hedonics associated with vanilla and its strong connection with sweetness can potentially help bias people’s food choice, food characteristics associated with preferential intake, and possibly also help to tackle the growing global obesity crisis²⁵.

In this narrative historical review, my particular interest is in trying to understand the origins of our hedonic response to the odour of vanilla (vanillin) which, at least according to the latest research, is currently one of the most liked smells cross-culturally²⁶. Vanilla flavouring is very widely used by the food industry globally, and the link between odour and food preference as it relates to vanilla is an intriguing one.— Later in this review, I will take a look at how the emerging knowledge and insights can be applied in the future in a food science/commercial context²⁷.

The world’s most liked smell

Arshamian et al. recently conducted a study of 225 individuals from nine diverse non-western cultures (including in several remote groups of hunter-gatherers and horticulturalists) in which the various groups of participants had to rank order ten odorants in terms of their hedonic ratings. The odorants, delivered from scented pens (thus minimizing any non-olfactory cues), were chosen on the basis of prior research from an urban sample in New York City²⁶ to cover a wide range of the hedonic spectrum. The odorants included vanillin, linalool, eugenol, ethyl butyrate, 2-Phenylethanol, 1-Octen-3-ol, 2-Isobutyl-3-methoxypyrazine, octanoic acid, diethyl sulphide, and isovaleric acid. Regardless of an individual’s cultural background, vanilla was the most liked of the ten odorants tested. Odour valence ratings correlated strongly and positively across all cultures. In fact, only 6% of the between-group variance was explained by culture. This result is certainly consistent with the title of Rain’s book claiming that vanilla is ‘the world’s favorite flavor and fragrance’,²⁷ though, of course, Arshamian et al.’s findings fall far short of rigorously supporting such a

claim empirically. Indeed, it is interesting to note how other researchers stress the chemical, biological, and cultural considerations, underpinning odour hedonics when considered in a global context²⁸. Vanilla was found elsewhere to be a very popular aroma cross-culturally, coming in just behind banana, peppermint, and lemon in a study of 22 aromas evaluated by 30 participants in each of 16 countries²⁹. On the basis of their findings, Pangborn and her colleagues “concluded that degree of liking of food-related aromas varies across regions, probably due to differences in traditional food habits and availability of regional flavour sources.” (29, p. 111).

Cross-cultural differences in odour hedonics have also been reported by other researchers. For instance, a German-Japanese study of odour hedonics, which did not include vanilla amongst the 18 olfactory stimuli tested, highlighted a number of cross-cultural differences between these populations in the case of specific odorants such as, for example, dried fish, soybean, and church incense³⁰. The former two odorants were rated as much more liked amongst the Japanese participants, whereas the latter smell was liked more by the German participants³¹. As well as any cultural differences in people’s odour preferences, some modest variation in the liking for seasonally-appropriate odours have been reported in both Japan and Germany^{32,33}. Furthermore, people’s responses to isovaleric acid, one of the less pleasant odorants presented to participants in Arshamian et al.’s study²⁷, changes markedly depending on how it is described (i.e., ‘ripe cheese’ vs. ‘sweaty socks’)^{34,35}, see also 36,37. Such results can presumably be taken to suggest that odour hedonics, at least for many real-world odorants, cannot simply be reduced to molecular structure^{24,38-40}. Furthermore, a range of other factors have also been shown to exert an influence over odour hedonics (in general, i.e., not specific to vanilla odour⁴¹): These include physiological factors (e.g., gestation, menstrual cycle, satiety; ⁴²⁻⁴³, gender ^{45,46}, aging⁴⁷, certain pathological states⁴⁸, the chemical environment (drugs, smoking, and pollution).

The ambient aroma of vanilla has also been documented to exert an ~~un~~ significant influence over people’s mood and arousal⁴⁹. More speculatively, it has been suggested that those companies wishing to reduce their customer’s shock when sending out large bills should consider scenting them with vanilla as this might help to relieve stress and anxiety⁵⁰. Elsewhere it was found that those individuals scented with vanilla, but not with, camphor, were more likely to be successful in ‘foot-in-the-door’-type procedures⁵¹. The scent of vanilla is apparently particularly popular amongst North American real-estate brokers⁵². Elsewhere, Hultén⁵³

documented a 65% increase in sales when a pleasant vanilla fragrance was released in the glassware section of an IKEA store over two weekends^{cf.54,55}. Due to its relaxing effect, vanilla has also been suggested as a potentially beneficial scent in a healthcare setting (see ³, 2021, p. 195). According to Hoppough⁵⁶, Sony stores also pump in a scent that incorporates notes of vanilla⁵⁷. Vanilla's sweet properties/associations can potentially be used to influence perceived sweetness and thereby bias consumer food behaviour in a healthy direction. At the same time, however, and as we will see later, it has also been suggested that ambient odour of vanilla may increase appetite (thus potentially biasing people to consume more).

The history of vanilla

The phenomenal global popularity of vanilla, both as a flavouring and as an ambient fragrance, is surprising given the origins of this aroma in nature^{58,59}. The vanilla orchid originated in Mexico, growing naturally in the Veracruz region of Papantla, where the exclusive pollinator of this orchid is the Melipona bee⁶⁰. In 1836, Charles Francois Antoine Morren travelled to Mexico in the hope of discovering the secret of how to produce vanilla beans⁶¹. While studying a vanilla orchid one day, Morren apparently observed the Mexican Melipona bee land on a vanilla orchid. He watched the bee lift a protective hood-like membrane covering the throat of the orchid before disappearing inside. The bee collected its pollen and left the orchid before flying on to the next flower and repeating the process. Surprisingly, unlike European orchids, the flower did not wither away, but produced a pod instead.

Vanilla flavour and functions

Originally, the flavour of vanilla was obtained from vanilla beans, the seed capsules of *Vanilla* spp. plants, which are now known to be native to tropical rainforests located from Mexico to Brazil^{62,63}. In recent years, vanilla orchids have been cultivated extensively in tropical areas such as Madagascar, Indonesia, and starting on Reunion Island in around 1819^{64,65}. Crucially, vanilla beans have no vanilla flavour upon harvesting. They only acquire their characteristic flavour after curing, which involves repeated drying and fermenting of the beans⁶⁶. Vanilla flavour, in other words, only develops as a result of fermentation of the

vanilla pod^{64,67}. The orchid genus *Vanilla* (Family: Orchidaceae) consists of 110 species, three of which are cultivated for their flavor-related commercial value (*Vanilla planifolia* Jacks. ex Andrews, *Vanilla tahitensis* J. W. Moore, and *Vanilla pompona* Schiede, Family: Orchidaceae⁶⁸). Originating in Central America, vanilla was also used as a flavouring agent. The Spanish first brought it to Europe from Mexico in around 1520⁶⁴; The absence of the appropriate insect pollinator, meant, however, that no vanilla pods were grown in Europe⁶⁴.

Originally, vanilla was used medicinally in Mesoamerica⁶⁹, and only later did it come to be used as a sweet flavouring. Indeed, it was the stomach-settling properties of vanilla that led to its incorporation into Coca-Cola during the 1920s^{27,69}. Contemporary research has proposed a number of pharmaceutical benefits of vanilla (e.g., in terms of fighting cancer⁷⁰⁻⁷²). Furthermore, like many herbs and spices, vanillin has been shown to help preserve food by inhibiting the activity of food spoilage yeasts, thus functioning as a highly-effective antimicrobial agent^{73,74}. (Note that vanilla is considered a spice⁷⁵.) In the 13th Century, Aztecs mixed cured vanilla pods with copal to scent their temples during sacred rituals, while the green bean was used medicinally (e.g., to treat venomous insect bites and heal wounds²⁷). Bythrow (69, p. 131) writes that: “The Aztecs and Totonacs knew that they had discovered a sacred plant in the vanilla orchid. Keeping their cultivation strategies a secret, early Americans utilized vanilla in their medicinal and culinary practices and introduced the Spanish conquerors to the sweet bean.” Given that fermented vanilla pods actually taste very bitter, it is an intriguing question as to when the aroma of vanilla first became associated with sweetness. Interestingly, vanillin is also unusual in that it is one of the very few odorants that has been shown to activate human bitter taste receptors TAS2R14, TAS2R20, and TAS2R39⁷⁶. According to Bythrow⁶⁹, Europeans historically valued vanilla for its flavour, its lore as a love potion, and also for its medicinal uses. It would certainly be interesting to know whether, in the ancient world, cured vanilla pods also smelled sweet to people, or rather just pleasant/relaxing⁷⁷⁻⁷⁹.

Nowadays, the aroma of cured vanilla beans includes descriptors such as sweet, vanillin, floral, prune/raisin, spicy, woody, and tobacco-like⁸⁰. That said, I would be tempted to question Bythrow’s (69, p. 131) assertion, quoted earlier, that the vanilla bean was perceived as sweet when it was first introduced to the Spanish conquistadors by the native Aztecs and Totonacs; my intuition is that this bitter-tasting bean may only have become sweet to the majority of the population once Coca-Cola introduced vanilla to its medicinal tonic in the

1920s. According to Rain²⁷, vanilla remained part of the American pharmacopoeia until 1916 for stomach distress, hysteria, and as a nerve stimulant. Here, though, it's interesting to note that Thomas Jefferson is credited with first adding a vanilla pod to ice-cream⁸¹. Jefferson apparently returned from France in 1789 with his chef—newly trained in making frozen desserts. In Philadelphia in 1791, Jefferson sent to France for 50 vanilla bean pods, implying that this was already popular flavouring.

The Aztecs are most often cited as the first culture to use and domesticate vanilla to flavour their drink of choice, *chocolat*⁶⁹. Chocolate was originally served as a bitter liquid, mixed with spices or corn puree. While it is uncertain whether it was Cortes who brought vanilla back to Spain, the flavour began to appear in Spain shortly after his conquests²⁷. After its arrival to Europe in the Sixteenth Century, sugar was added and it became popular throughout society, first among the ruling classes and then among the common people^{82,83}. Sugar was presumably only widely combined with vanilla in chocolate drinks once the price of sugar had declined⁸⁴⁻⁸⁷. Similarly, one only sees the formerly very expensive vanilla appearing in French perfume recipes and store inventories around the start of the 19th Century (see 88, pp. 155-156).

Contemporary production of vanilla flavouring

Vanilla beans containing the flavour and their extracts using solvents such as ethanol and glycerol are referred to as 'natural vanilla flavour', whereas synthesised aroma compounds such as vanillin contained in vanilla beans and its analogue ethyl vanillin are referred to as 'synthetic vanilla flavour'⁸⁹. The key structural feature of vanillin, the main compound responsible for the characteristic flavour of vanilla, is the vanillyl (3-methoxy-4-hydroxyphenyl) group. Synthetic pure vanillin lacks complex flavour notes, and thus cannot replace vanilla in high quality products. Although customers typically prefer nonsynthetic flavours, non-synthetic vanilla extract is approximately 200 times more expensive than its synthetic counterpart^{64,90}, in part because it is one of the most labour-intensive crops to grow worldwide^{66,91}. As a result, vanilla pods are expensive with the price fluctuating sometimes unpredictably according to current political and climatic conditions. This price instability is one of the reasons for the shift to artificial vanilla flavour or pure vanillin by many in the food and beverage industry. Demand for cured vanilla pods, over 16,000 metric tons in

2015⁹², exceeds supply by as much as ten to one. With the high prices and the inability of non-synthetic vanilla production to meet the global demand for the vanilla flavour, the focus for many in the flavour industry has turned to pure vanillin.

Contemporary popularity of vanilla

Nowadays, vanillin, the key compound that gives rise to the distinctive scent of vanilla is used widely in foods such as cola beverages and ice-cream⁹⁰, sugar cookies, puff pastries, and butter creams⁶⁹. The vanilla flavour from pods tends to be used for high quality products such as expensive chocolates and luxury ice creams (see ⁹³, pp. 61-62), which account for approximately 75% of the total vanilla plant flavour used. Long associated with dessert, vanilla is the second most expensive spice, just behind saffron, according to Bythrow⁶⁹, due primarily to the extensive processing required. Other major uses include cola-type beverages and baked foods⁹⁰. Indeed, there is plenty of evidence to suggest that the aroma of vanilla tends to enhance the sweetness of whatever foodstuff it is added to^{94,95}. Recently, it has been proposed that the addition of the sweet aroma of vanilla might prove to be an effective strategy in terms of sugar reduction⁹⁶. In recent years, numerous different research groups have become interested in crossmodal approaches to sugar reduction⁹⁷⁻¹⁰⁰. The results of one study revealed that a 25% sugar reduction in yoghurt with 0.2% vanilla (or strawberry) flavour added did not affect the sweetness, as shown in temporal profiles, and hedonic ratings⁹⁶. That said, the study reported that vanilla flavours, e.g., such as vanillin, appear to have appetite-enhancing effects⁸⁹, suggesting that the presence of this odorant might actually lead to an increase in people's consumption instead¹⁰¹⁻¹⁰².

Ultimately, though, the question has to be why the flavour of the fermented bean of this obscure (not to mention, hard to pollinate) orchid originating in Mexico should have become such a highly-liked and dominant flavour and fragrance globally in recent centuries. A secondary question is when exactly did this bitter-tasting cured vanilla pod come to be associated with sweetness^{cf.76}.

Explaining odour hedonics

Having documented the global popularity of vanilla (both as a flavouring, and as an ambient fragrance), the question becomes one of how to explain its ubiquity in terms of the acquired, innate, and molecular accounts of odour hedonics that were mentioned earlier^{cf.24}.

Predicting odour hedonics for unfamiliar monomolecular odorants

Our response to a variety of unfamiliar volatile aromatic chemicals may be predicted on the basis of the structure of the molecules themselves. When the number of distinct olfactory notes ascribed to a range of 411 monomolecular odorants was assessed¹⁰³, both expert and non-expert participants perceived structurally more complex monomolecular odorants as having a larger number of olfactory notes. Functional magnetic resonance imaging has revealed that more structurally-complex odorant molecules tend to give rise to a larger number of descriptions, and evoke greater activity in the dorsal anterior cingulate gyrus than do simpler odorants¹⁰⁴. The latter brain area is responsible for odour-related verbal and lexical processing, and for the resolving of cognitive conflict, perhaps associated with competing odour descriptors.

The pleasantness of an unfamiliar odour tracks, or at least correlates weakly with, the size of the molecule. In a study reported by Kahn and colleagues,¹⁴ 185 participants took part in one of seven studies involving some subset of 144 monomolecular odorants and 16 mixtures which were rated a range of (perceptual) dimensions. Elsewhere, hedonic judgments of chemical compounds were positively correlated with molecular size/weight, meaning that larger molecules tended to be rated as more pleasant¹⁰⁵. The correlation between molecular weight and olfactory pleasantness for novel monomolecular odorants has been documented elsewhere^{106,cf.24}).

It would appear that more complex monomolecular odorants give rise to perceptual experiences that involve a larger number of describable olfactory elements (or notes) and are also rated as more pleasant. Although the correlation between physicochemical properties of individual volatile odour molecules and odour hedonics in the case of unfamiliar monomolecular compounds is significant, it is not particularly strong^{14,105}. As such, the associative learning (or acquired/hedonic) account of why we like the smell of particular volatiles that we happen to be very familiar with (i.e., such as vanilla) may, in most everyday circumstances at least, explain more of the variance than a purely molecular account^{cf.107-109}.

The crowd-sourced DREAM ([Dialogue for Reverse Engineering And Methods](#)) Olfaction Prediction Challenge utilised a large olfactory psychophysical data set to develop machine-learning algorithms to predict the sensory attributes of molecules based on their chemoinformatic features¹¹⁰. Intriguingly, the resulting models accurately predicted odour intensity and pleasantness, as well as successfully predicting 8 among 19 rated semantic descriptors (“garlic,” “burnt,” “spices,” “fish,” “sweet,” “fruit,” “flower,” and “sour”). That said, the psychophysical data was collected from just one cultural group, and none of the models did a particularly good job of accounting for differences at the level of the individual observer. Relevant to the theme of the present review, the ‘bakery’ olfactory descriptor could be predicted on the basis of the similarity of a given molecule to vanillin or ethyl vanillin.

Here, though, it should be noted that the majority of scents that we come across in daily life actually consist of several tens or even hundreds of different volatiles. In fact, some of the most desirable flavours (e.g., think of the flavour of quality wine or coffee) may contain well over a thousand distinct volatile compounds¹¹¹⁻¹¹⁴. Even though only a small proportion of these are likely perceptible consciously, and hence contribute to the unified aroma/flavour percept^{23,115}, the majority of natural odours are composed of a multitude of volatiles unlike the monomolecular olfactory stimuli used in the majority of studies. According to the research, somewhere between 40 and 80 volatiles contribute to the complex natural flavour of vanilla^{64,116-118}. Perhaps unsurprisingly, natural vanilla is considered a complex flavour¹¹⁹ (though see ¹¹³). It is relevant here to note that attempts to predict the pleasantness of odour mixtures have thus far been restricted to binary combinations of one pleasant and one unpleasant odorant. In particular, researchers have reported that they were able to predict the perceived pleasantness of binary mixtures of two odorants (three of which were pleasant and three unpleasant), with pleasantness ratings falling in-between, and dependent on relative concentration¹²⁰. In this case, the prediction was based on the pleasantness of the elements weighted by their respective perceived intensities.

Innate responses to evolutionarily-important odours

In a few cases, it has been suggested that our response to evolutionarily-important odorants may be innate/hard-wired. This suggestion has been made, for example, with regard to volatile chemical *trans*-4,5-epoxy-(*E*)-2-decenal (E2D) that is found in mammalian blood. Its

presence tends to induce postural avoidance, increases physiological arousal, and enhances visual perception of affective stimuli in humans. The presence of this odorant would appear to serve as an alarm signal across species²¹. That said, there would seem to be little reason to suspect that vanilla, or vanillin, would be associated with such an innate response.

Associative learning and the response to vanillin

Familiarity breeds liking (as in the ‘mere exposure’ effect¹²¹⁻¹²⁴). Our early preference for umami and sweet taste might thus be the result of exposure to amniotic fluid and breast milk which both apparently have these as dominant tastes (¹²⁵; see also ¹²⁶⁻¹²⁸). On the challenge of knowing whether the detection of olfactory stimuli in water should be described as tasting or smelling see Parker and Stabler¹²⁹. It is important to note that after birth, classical conditioning can modify olfactory preferences^{130,131}. That said, the preference for the taste/flavour of breast milk declines prior to adulthood for most of us (though there was a short-lived trend for breast-milk ice-cream¹³²⁻¹³⁴).

It would certainly seem relevant here to consider the influence of early exposure to vanilla in flavouring of foods on later liking¹³⁵. ~~One of the dominant notes (or volatile chemicals) giving rise to the aroma of vanilla, v~~Vanillin, ~~(a simple phenolic compound). In fact, vanillin, one of~~ the odorants used in Arshamian et al.’s study²⁶ also happens to be present in human breast milk depending on the mother’s diet (and is also present in artificial breast formulations¹³⁶⁻¹³⁹). Various beneficial effects on infant health and immune response have been associated with the consumption of breast milk¹⁴⁰, including the transfer of various taste/flavour preferences from the mother to her offspring. The flavours of alcohol and garlic have, for instance, been shown to pass to the breast milk from the mother’s diet¹⁴¹⁻¹⁴³. Though here, it is also important to note that changes in the odour of the mother’s breath or sweat may also impact on a baby’s weaning behaviour, and both may exert an influence a baby’s subsequent feeding behaviour.

Vanilla was also apparently added to ~~bottle~~-milk formula for new-borns and small babies for decades in Germany (¹⁴⁴; see also ¹³⁵). Intriguingly, Mennella and Beauchamp (¹³⁸, p. 14) note, in passing, how “some authorities have recommended that vanilla be added to formulas intended for infants beginning the weaning process, reportedly for both its flavoring

properties and its ability to provide variety to the bottle-fed infant's otherwise "bland" diet (Food Advisory Committee, 1992). And finally, vanilla-flavoured pacifiers (Ross Laboratories, Inc.) are currently being distributed in some hospitals."¹³⁸ These authors also write that: "These data support the hypothesis that flavors, either consumed by the mother and transmitted to her milk or added to formula, are detected by the infant and serve to modulate feeding. They also suggest that experience with a flavor in milk alters the infant's responsiveness to that flavor during subsequent feedings. It is hypothesized that under the natural condition of breast-feeding, infants become familiar with the flavors consumed by their mothers, and such experiences may impact on later food and flavor acceptability and choice." (¹³⁸, p. 13; see also ¹⁴⁵⁻¹⁴⁸). Indeed, long ago, Steiner¹⁴⁹ observed that newborns exhibited a "smiling" expression accompanied by sucking and licking movements when exposed to the smell of vanilla. On the flip-side, however, Soussignan and colleagues addressed the same question using butyric acid (rancid butter smell) and vanillin odours, they found that while butyric acid induced more facial expressions of disgust than smiles in newborns (65% vs. 33%), the smell of vanillin induced as many facial expressions of disgust as smiles (33% vs. 33%).¹⁵⁰ Bear in mind here, as stated earlier, that vanillin turns out to be one of the very few odorants capable of activating the human bitter taste receptors⁷⁶.

Scenting a breastfed infant's toy with vanilla leads to increased behavioural interaction (that is, increased looking and reduced vocalization) as compared to an unscented toy or one that had been scented with ethanol¹⁵¹. Haller and colleagues conducted an intriguing questionnaire study of 133 people (aged between 12 and 59 years of age) visiting an exhibition in Germany¹³⁵. The researchers asked whether the participants had been breast or bottle fed as a newborn. The participants were then given tomato ketchup, and tomato ketchup adulterated with a very low level of vanillin, and asked which they preferred. (Note that the food chosen was one in which vanillin would normally not have been experienced, hence implying that any preference changes that were observed must have been attributable to the presence of vanillin rather than to a previously-familiar combination of ingredients.) Intriguingly, the results revealed that those who reported having been bottle-fed as a baby (N = 30) preferred the ketchup that had been adulterated with a low-level of vanillin (66.7% to 33.3% for original), whereas those who had been breastfed as a small child (N = 103) preferred the unadulterated ketchup instead (70.9% vs. 29.1% for the adulterated version).

A growing body of research now shows that we come to acquire a liking for those flavours in breast milk that our mothers consumed during pregnancy^{136,152}. Schaal et al. conducted an intriguing of 24 neonates born to mothers who either had or had not consumed anise flavour during pregnancy. The babies born to anise consuming mothers showed a stable preference for anise odour immediately after birth and on day four. By contrast, those infants born to anise non-consuming mothers displayed aversion or neutral responses instead. In recent years, a number of programs have started to encourage pregnant mothers to consume carrot-flavoured milk, etc., in order to increase the likelihood that their offspring will consume these vegetables after birth¹⁵³⁻¹⁵⁵. What such findings suggest is that we acquire a liking for that which we are exposed to during our early taste/flavour experiences¹⁵⁶. Vanilla flavouring is ubiquitous, and crucially, the flavour passes to the breast milk as well as often being added to formula milk, and even to pacifiers.

Molecular vs. perceptual accounts of olfactory perception

It is also important to consider whether the key volatile, vanillin, ~~that gives rise to the smell of vanilla~~ is also present, as one of the volatiles, in other sweet-flavoured foods²³. Indeed, it is currently unclear whether it is the physical presence of specific volatile or its perception that is key to driving the odour-induced taste enhancement, and odour hedonics. Given that the prediction of perceptual response to combinations of odorants is still challenging (to say the least), it is unclear whether it is the physical presence of a specific volatile, or the perceptual properties of the volatile mixture, that are key to determining learning/behaviour/hedonics. Note that, in this regard, humans are very different from mice, say, who are able to decompose odour mixtures into their component parts¹⁵⁷⁻¹⁵⁹. Much the same conclusion, i.e., supporting a perceptual account, emerges from those studies of flavour pairing, where shared compounds have proven to be a poor predictor of the pleasantness of the combination¹⁶⁰⁻¹⁶¹. Rather, it is matching (or possibly contrasting) the perceptual qualities of the component stimuli which may well be made up of several tens or hundreds of different volatile molecules that are key to creating great-tasting flavour combinations.

One of the peculiar features of vanilla is that its psychophysical curve is particularly flat. Indeed, when Moskowitz and his colleagues evaluated the hedonic character of 32 odorous molecules as a function of the concentration of the stimulus, vanillin was found to be one of

the few molecules that showed a psychophysical response pattern with almost no relationship between the intensity of the stimulus and the pleasant nature of the odorant.¹⁶² That is, vanillin was rated as pleasant regardless of the concentration (i.e., low, medium, or high) at which it was delivered, unlike what is typically seen for the majority of odorants. It is interesting to speculate as to whether this peculiarity might be linked to the fact that vanillin is one of the few pure olfactants (i.e., with no trigeminal component¹⁶³⁻¹⁶⁶). That said, according to research contemporary research, it would appear that ethyl vanillin is a TRPA1 agonist¹⁶⁷. At the same time, however, it is worth noting that little is currently known about the neural processing of the reward value of vanilla odour¹⁶⁸⁻¹⁶⁹.

Genetic determinants of response to olfactory stimuli

Before closing this section, it is important to note that genetically-determined individual differences play an important role in determining our liking for certain odorants/flavours¹⁷⁰. In the case of coriander/cilantro, for example, one and the same molecule smells pleasantly citrusy/herbal to one section of the population, but unpleasantly soapy to another section of the population¹⁷¹⁻¹⁷³. Androstenone is another chemosensory stimulus that delivers very different perceptual and hedonic responses as a function of an individual's genetic make-up¹⁷⁴. One should probably also consider the incidence of anosmia to vanilla that affects roughly 1% of the population¹⁷⁵⁻¹⁷⁶. Such specific anosmias¹⁷⁷⁻¹⁸⁰ are likely to influence odour hedonics too¹⁸¹.

Given that we live in different taste worlds, it is clearly important to identify those flavours that are most widely liked in order to bias food choice. While the smell of vanilla has been identified as one of the most widely liked aromas²⁶, it is also described as smelling sweet. Food scientists are becoming increasingly interested in the possibility that sweet smells such as vanilla could be used to maintain the sweetness of commercial food products while, at the same time, reducing the use of calorific sweeteners²⁶. At the same time, however, it has also been suggested that vanilla aroma may also encourage consumption⁸⁹. Such an approach is likely to be facilitated by the low cost of artificial vanilla flavouring (when compared to the high and fluctuation price of vanilla pods).

Conclusions

Recently, Arshamian et al. documented the cultural independence of hedonic responses to a range of familiar odorants, such as vanilla²⁶. It is tempting to think that such observations reflect a molecular basis for odour hedonics^{14,182}. It is, however, important to note that the majority of such claims (i.e., for the molecular basis of olfactory pleasantness) appearing in the literature, have been based on predicting people's responses to unlabelled monomolecular odorants that are often novel/unfamiliar^{103,105,110}, thus effectively eliminating the marked effects of learning/semantics that have frequently been documented so frequently in the literature. The recently-discovered existence of olfactory 'metamers' that have a different molecular structure, but which smell indistinguishable also argues against an unmediated molecular account of odour pleasantness¹⁸³.

Arshamian and colleagues (²⁶, p. 4) conclude that: "Our data do not, however, adjudicate between learned versus innate explanations of odor pleasantness perception. Global regularities in odor perception could indicate common and shared experiences across all human groups. Infant data from diverse cultural contexts could adjudicate between these possibilities, although even here there are challenges since the fetus is already being enculturated into a specific chemical environment."^{26,cf.184}. There is little reason to consider an innate account as providing a plausible explanation for the ubiquitous preference that has been demonstrated for vanilla. When it comes to familiar odorants, the associative learning/conditioning more likely to play a key role in determining valence^{108,185}. As such, one would expect cultural differences depending on how/when a given odorant is experienced during development¹⁸⁶. However, it is important to stop short of inferring from this that molecular account can explain odour hedonics. Clearly, as we have seen, the molecular structure of these volatile compounds is important in helping to understand people's response to unfamiliar monomolecular odorants. However, the globally positive hedonic response to vanillin, ~~the key compound giving rise to what we recognize as odorants~~ is more likely associative/familiarity-based. Meanwhile, the explanation for why trigeminal odorants are typically rated as unpleasant is presumably physiological in origin.

Of the more than 25,000 varieties of *Orchidaceae* (orchids are the most prolific member of the angiosperm family¹⁸⁷), it is the pod from the vanilla orchid, that nowadays flavours so many of our foods and drinks, not to mention scenting our homes, and many home and personal care (HPC) products as well¹⁸⁸. According to the research that has been reviewed

here, vanilla is likely so universally liked, not because of its molecular structure *per se*, but because of ubiquitous early exposure (it is one of the odorants that passes to breast milk), not to mention in a historical context its anti-microbial, medicinal, and anxiety-relieving properties^{69,189,190}. Furthermore, the suggestion is made that the bitter-tasting vanilla pod may only have taken on sweet properties when Europeans started to drink sweetened chocolate after vanilla's arrival in Europe in the 16th Century (see ¹⁹¹⁻¹⁹³ on the biological and cultural role of sweeteners).

REFERENCES

1. McGann, J. P. Poor human olfaction is a 19th-century myth. *Science*, **356**, eaam7263 (2017).
2. Spence, C. Using ambient scent to enhance well-being in the multisensory built environment. *Frontiers in Psychology* (SI: Smells, Well-being, and the Built Environment), **11**:598859 (2020). doi: 10.3389/fpsyg.2020.598859.
3. Spence, C. *Sensehacking: How to use the power of your senses for happier, healthier living*. London, UK: Viking Penguin (2021).
4. Croy, I., Bojanowski, V., & Hummel, T. Men without a sense of smell exhibit a strongly reduced number of sexual relationships, women exhibit reduced partnership security - a reanalysis of previously published data. *Biological Psychology*, **92**(2), 292-294 (2013). <https://doi.org/10.1016/j.biopsycho.2012.11.008>
5. Spence, C. The scent of attraction & the smell of success: Crossmodal influences on person perception. *Cognitive Research: Principles & Implications*, **6**(1):46 (2021). DOI: [10.1186/s41235-021-00311-3](https://doi.org/10.1186/s41235-021-00311-3). <https://rdcu.be/cngR9>.
6. Croy, I., Negioas, S., Novakova, L., Landin, B. N., & Hummel, T. Learning about the functions of the olfactory system from people without a sense of smell. *PLoS One*, **7**:e33365 (2012).
7. Sharma, A., Kumar, R., Aier, I., Semwal, R., Tyagi, P., & Varadwaj, P. Sense of smell: Structural, functional, mechanistic advancements and challenges in human olfactory research. *Current Neuropharmacology*, **17**(9), 891-911 (2019). <https://doi.org/10.2174/1570159X17666181206095626>.
8. Spence, C. Leading the consumer by the nose: On the commercialization of olfactory-design for the food and beverage sector. *Flavour*, **4**:31 (2015). DOI 10.1186/s13411-015-0041-1
9. Spence, C. Enhancing the experience of food and drink via neuroscience-inspired olfactory design. *The Senses and Society*, **12**, 209-221 (2017). [http://dx.doi.org/10.1080/17458927.2017.1270800](https://dx.doi.org/10.1080/17458927.2017.1270800).
10. Hutmacher, F. Why is there so much more research on vision than on any other sensory modality? *Frontiers in Psychology*, **10**:2246 (2019). doi: 10.3389/fpsyg.2019.02246.

11. Spence, C. On the psychological impact of food colour. *Flavour*, **4**:21 (2015). DOI 10.1186/s13411-015-0031-3.
12. Spence, C. On the manipulation, and meaning(s), of colour in food: A historical perspective. *Journal of Food Science* (submitted).
13. Herz, R. *The scent of desire: Discovering our enigmatic sense of smell*. New York, NY: William Morrow (2007).
14. Khan, R. M., Luk, C.-H., Flinker, A., Aggarwal, A., Lapid, H., Haddad, R., & Sobel, N. Predicting odor pleasantness from odorant structure: Pleasantness as a reflection of the physical world. *Journal of Neuroscience*, **27**, 10015-10023 (2007).
15. Yeshurun, Y., & Sobel, N. An odor is not worth a thousand words: From multidimensional odors to unidimensional odor objects. *Annual Review of Psychology*, **61**, 219-241 (2010).
16. Soudry, Y., Lemogne, C., Malinvaud, D., Consoli, S. M., & Bonfils, P. Olfactory system and emotion: Common substrates. *European Annals of Otorhinolaryngology, Head and Neck Diseases*, **128**(1), 18-23 (2011).
17. Stevenson, R. J., & Boakes, R. A. Sweet and sour smells: Learned synaesthesia between the senses of taste and smell. In G. A. Calvert, C. Spence, & B. E. Stein (Eds.), *The handbook of multisensory processing* (pp. 69-83). Cambridge, MA: MIT Press (2004).
18. Aikman, L. Perfume, the business of illusion. *National Geographic*, **99**, 531-550 (1951).
19. Moran, T. Ah, the aroma of a just-baked sedan. *New York Times*, **May 14th**, 40 (2000).
20. Moran, T. Sniffing car parts: Yes, the job stinks. *New York Times*, **May 14th**, 40 (2000).
21. Arshamian, A., Laska, M., Gordon, A. R., Norberg, M., Lahger, C., Porada, D. K., Jelvez Serra, N., Johansson, E., Schaefer, M., Amundin, M., Melin, H., Olsson, A., Olsson, M. J., Stensmyr, M., & Lundström, J. N. A mammalian blood odor component serves as an approach-avoidance cue across phylum border - from flies to humans. *Scientific Reports*, **7**:13635 (2017). DOI:10.1038/s41598-017-13361-9.
22. Mandaïron, N., Poncelet, J., Bensafi, M., & Didier, A. Humans and mice express similar olfactory preferences. *PLoS One*, **4**:4209 (2009).
23. Spence, C. Factors affecting odour-induced taste enhancement. *Food Quality & Preference*, **96**:104393 (2022). <https://doi.org/10.1016/j.foodqual.2021.104393>.
24. Jossain, P., Chakirian, A., Kermen, F., Rouby, C., & Bensafi, M. Physicochemical influence on odor hedonics: Where does it occur first? *Communicative & Integrative Biology*, **4**, 563-565 (2011).
25. Caballero, B. The global epidemic of obesity: An overview. *Epidemiologic Reviews*, **29**, 1-5 (2007).
26. Arshamian, A., Gerkin, R. C., Kruspe, N., Wnuk, E., Floyd, S., O'Meara, C., Rodriguez, G. G., Lundström, J. N., Mainland, J. D., & Majid, A. The perception of odor pleasantness is shared across cultures. *Current Biology*, **32**(9), [P2001-P2006.E3](https://doi.org/10.1016/j.cub.2022.02.062). (2022). <https://doi.org/10.1016/j.cub.2022.02.062>.
27. Rain, P. *Vanilla: The cultural history of the world's favorite flavor and fragrance*. USA: Jeremy P. Tarcher (2004).
28. Kaiser, R. *Meaningful scents around the world: Olfactory, chemical, biological, and cultural considerations*. Weinheim: Wiley (2006).

29. Pangborn, R. M., Guinard, J.-X., & Davis, R. G. Regional aroma preferences. *Food Quality and Preference*, **1**(1), 11-19 (1988).
30. Ayabe-Kanamura, S., Schicker, I., Laska, M., Hudson, R., Distel, H., Kobayakawa, T., & Saito, S. Differences in perception of everyday odors: A Japanese-German cross-cultural study. *Chemical Senses*, **23**, 31-38 (1998).
31. Schleidt, M., Neumann, P., & Morishita, H. Pleasure and disgust: Memories and associations of pleasant and unpleasant odours in Germany and Japan. *Chemical Senses*, **13**, 279-293 (1988).
32. Seo, H.-S., Buschhüter, D., & Hummel, T. Odor attributes change in relation to the time of the year. Cinnamon odor is more familiar and pleasant during Christmas season than summertime. *Appetite*, **53**(2), 222-225 (2009). <http://doi.org/10.1016/j.appet.2009.06.011>.
33. Wada, Y., Inada, Y., Yang, J., Kunieda, S., Masuda, T., Kimura, A., Kanazawa, S., & Yamaguchi, M. K. Infant visual preference for fruit enhanced by congruent in-season odor. *Appetite*, **58**(3), 1070-1075 (2012). <https://doi.org/10.1016/j.appet.2012.02.002>.
34. De Araujo, I. E., Rolls, E. T., Velazco, M. I., Margot, C., & Cayeux, I. Cognitive modulation of olfactory processing. *Neuron*, **46**, 671-679 (2005).
35. Herz, R. S., & von Clef, J. The influence of verbal labelling on the perception of odors: Evidence for olfactory illusions? *Perception*, **30**, 381-391 (2001).
36. Bensafi, M., Rinck, F., Schaal, B., & Rouby, C. Verbal cues modulate hedonic perception of odors in 5-year-old children as well as in adults. *Chemical Senses*, **32**, 855-862 (2007).
37. Djordjevic, J., Lundstrom, J. N., Clément, F., Boyle, J. A., Pouliot, S., & Jones-Gotman, M. A rose by any other name: Would it smell as sweet? *Journal of Neurophysiology*, **99**, 386-393(2008).
38. Cain, W. S. *Lability of odor pleasantness*. In J. H. A. Kroeze (Ed.), *Preference behavior and chemoreception* (pp. 303-315). London, UK: IRL Press (1979).
39. Poncelet, J., Rinck, F., Ziesel, A., Joussain, P., Thevenet, M., Rouby, C., & Bensafi, M. Semantic knowledge influences prewired hedonic responses to odors. *PLoS One*, **5**:13878 (2010). DOI: 10.1371/journal.pone.0013878.
40. Rouby, C., Pouliot, S., & Bensafi, M. Odor hedonics and their modulators. *Food Quality and Preference*, **8**, 545-549 (2009).
41. Distel, H., Ayabe-Kanamura, S., Martinez-Gomez, M., Schicker, I., Kobayakawa, T., Saito, S., & Hudson, R. Perception of everyday odors – correlation between intensity, familiarity and strength of hedonic judgment. *Chemical Senses*, **24**, 191-199 (1999).
42. Faas, M. M., Melgert, B. N., & de Vos, P. A brief review on how pregnancy and sex hormones interfere with taste and food intake. *Chemosensory Perception*, **3**, 51-56 (2010)..
43. Kringelbach, M. L., O'Doherty, J., Rolls, E. T., & Andrews, C. Sensory-specific satiety for the flavour of food is represented in the orbitofrontal cortex. *NeuroImage*, **11**, S767 (2000).
44. Rolls, E. T., & Rolls, J. H. Olfactory sensory-specific satiety in humans. *Physiology & Behavior*, **61**, 461-473 (1997).
45. Brand, G., & Millot, J.-L. Sex differences in human olfaction: Between evidence and enigma. *Quarterly Journal of Experimental Psychology*, **54B**, 259-270 (2001).

46. Spence, C. Do men and women really live in different taste worlds? *Food Quality & Preference*, **73**, 38-45 (2019). <https://doi.org/10.1016/j.foodqual.2018.12.002>.
47. Joussain, P., Thevenet, M., Rouby, C., & Bensafi, M. Effect of aging on hedonic appreciation of pleasant and unpleasant odors. *PLoS ONE*, **8(4)**:e61376 (2013). <https://doi.org/10.1371/journal.pone.0061376>.
48. Zhao, J. B., Wang, Y. L., Ma, Q. W., Zhao, J. B., Zhang, X. Y., & Zou, L. Q. The chemosensory pleasure scale: A new assessment for measuring hedonic smell and taste capacities. *Chemical Senses*, **44(7)**, 457-464 (2019). DOI: [10.1093/chemse/bjz040](https://doi.org/10.1093/chemse/bjz040).
49. De Wijk, R. A., & Zijlstra, S. M. Differential effects of exposure to ambient vanilla and citrus aromas on mood, arousal and food choice. *Flavour*, **1**:24 (2012). <http://www.flavourjournal.com/content/1/1/24>.
50. Harris, E., Sengupta, C., & Gray, A. DOPE: The Digiscent-operated persuasive environment (2000). <http://www.stanford.edu/~demian23/captology/DOPE/>.
51. Saint-Bauzel, R., & Fointiat, V. The sweet smell of the requester: Vanilla, camphor, and foot-in-the-door. *Social Behavior and Personality*, **40(3)**, 369-374 (2012). <http://dx.doi.org/10.2224/sbp.2012.40.3.369>.
52. Demetros, V. A. M. The sweet smell of success. *The Crafts Report*, **April** (1997). <http://www.craftsreport.com/april97/aroma.html>.
53. Hultén, B. Sensory cues and shoppers' touching behaviour: The case of IKEA. *International Journal of Retail and Distribution Management*, **40**, 273-289 (2012).
54. Fetterman, J., & O'Donnell, J. Just browsing the mall? That's what you think. *USA Today*, **September 1st** (2006). http://usatoday30.usatoday.com/money/industries/retail/2006-09-01-retail-cover-usat_x.htm.
55. Trivedi, B. Recruiting smell for the hard sell. *New Scientist*, **2582**, 36-39 (2006).
56. Hoppough, S.. What's that smell? *Forbes*, **September 15th** (2006). <https://www.forbes.com/forbes/2006/1002/076.html#3ef667ec5377>.
57. Herz, R. S. The emotional, cognitive, and biological basics of olfaction: Implications and considerations for scent marketing. In A. Krishna (Ed.), *Sensory marketing: Research on the sensuality of products* (pp. 87-107). New York, NY: Routledge (2010).
58. Abreu-Runkel, R. *Vanilla: A global history*. London: Reaktion Books (2020).
59. Ecott, T. *Vanilla: Travels in search of a luscious substance*. London, UK: Michael Joseph (2004).
60. Youssef, J., & Spence, C. Introducing diners to the range of experiences in creative Mexican cuisine, including the consumption of insects. *International Journal of Gastronomy & Food Science*, **25**:100371 (2021). <https://doi.org/10.1016/j.ijgfs.2021.100371>.
61. Sethi, S. The bittersweet story of vanilla. *Smithsonian Magazine*, **April 3rd** (2017). <https://www.smithsonianmag.com/science-nature/bittersweet-story-vanilla-180962757/>.
62. Gamboa-Gaitán, M. A., & Otero-Ospina, J. T. Colombian vanilla and its microbiota. III. Diversity and structure of the endophytic community. *Acta Botanica Hungaria*, **58**, 241-256 (2016).
63. Hotta, M., Ogata, K., Nitta, A., Hoshikawa, K., Yanagi, M., & Yamazaki, K. *Useful plants of the world*. Tokyo: Heibonsha (1989).

64. Khoyratty, S., Kodjac, H., & Verpoorte, R. Vanilla flavor production methods: A review. *Industrial Crops & Products*, **125**, 433-442 (2018). doi: 10.1016/j.indcrop.2018.09.028.
65. Nishi, T. Shokubutsu-no hanashi arekore 7. *Puranta*, **64**, 49-53 (1999).
66. Dignum, M. J. W., Kerler, J., & Verpoorte, R. Vanilla curing under laboratory conditions. *Food Chemistry*, **79**, 165-171 (2002).
67. Walton, N. J., Mayer, M. J., & Narbad, A. Vanillin. *Phytochemistry*, **63**, 505-515 (2003). doi: 10.1016/S0031-9422(03)00149-3.
68. Soto-Arenas, M. A., & Cribb, P. A new infrageneric classification and synopsis of the genus *Vanilla* Plum. ex Mil. (Orchidaceae: Vanillinae). *Lankesteriana*, **9**, 355-398 (2010).
69. Bythrow, J. D. Vanilla as a medicinal plant. *Seminars in Integrative Medicine*, **3**, 129-131 (2005). doi: 10.1016/j.sigm.2006.03.001.
70. Bezerra, D. P., Soares, A. K., & de Sousa, D. P. Overview of the role of vanillin on redox status and cancer development. *Oxidative Medicine and Cell Longevity*, **2016**:9734816 (2016). doi: 10.1155/2016/9734816.
71. Bezerra, D. P., Soares, A. K., & de Sousa, D. P. Therapeutic potential of vanillin and its main metabolites to regulate the inflammatory response and oxidative stress. *Mini-Reviews in Medicinal Chemistry*, **19**, 1681-1693 (2019). doi: 10.2174/1389557519666190312164355.
72. Cheng, H. M., Chen, F. Y., Li, C. C., Lo, H. Y., Liao, Y. F., Ho, T. Y., et al. Oral administration of vanillin improves imiquimod-induced psoriatic skin inflammation in mice. *Journal of Agriculture Food Chemistry*, **65**, 10233-10242 (2017). doi: 10.1021/acs.jafc.7b04259
73. Fitzgerald, D. J., Stratford, M., Gasson, M. J., Ueckert, J., Bos, A., & Narbad, A. Mode of antimicrobial action of vanillin against *Escherichia coli*, *Lactobacillus plantarum* and *Listeria innocua*. *Journal of Applied Microbiology*, **97**, 104-113 (2004). doi: 10.1111/j.1365-2672.2004.02275.x
74. Fitzgerald, D. J., Stratford, M., & Narbad, A. Analysis of the inhibition of food spoilage yeasts by vanillin. *International Journal of Food Microbiology*, **86**, 113-122 (2003).
75. Cameron, K. M. Vanilla phylogeny and classification. In D. Havkin-Frenkel & F. C. Belanger (Eds.), *Handbook of vanilla science and technology* (p. 243). Oxford, UK: Wiley-Blackwell (2011).
76. Morini, G., Winnig, M., Vennegeerts, T., Borgonovo, G., & Bassoli, A. Vanillin activates human bitter taste receptors TAS2R14, TAS2R20, and TAS2R39. *Frontiers in Nutrition*, **8**:683627 (2021). doi: 10.3389/fnut.2021.683627.
77. Draycott, J. Smelling trees, flowers and herbs in the ancient world. In M. Bradley (Ed.), *Smell and the ancient senses* (pp. 60-73). London, UK: Routledge (2015).
78. Huber, B., Larsen, T., Spengler, R. N., & Boivin, N. How to use modern science to reconstruct ancient scents. *Nature Human Behaviour*, s41562-022-01325-7 (2022). doi:10.1038/s41562-022-01325-7.
79. Classen, C., Howes, D., & Synnott, A. *Aroma: The cultural history of smell*. London, UK: Routledge (1994).

80. Ranadive, A. S. Quality control of vanilla beans and extracts. In D. Havkin-Frenkel & F. C. Belanger (Eds.), *Handbook of vanilla science and technology* (pp. 141-160). Oxford, UK: Wiley-Blackwell (2011).
81. Diamond, A. Make Thomas Jefferson's recipe for ice cream. The co-author of the Declaration of Independence also drafted a radical recipe. *The Smithsonian Magazine*, **July** (2020). <https://www.smithsonianmag.com/history/thomas-jefferson-ice-cream-recipe-180975200/>.
82. Kerr, J. *History of chocolate*. Field Museum (2007). <http://archive.fieldmuseum.org/chocolate/history.html>.
83. Newquist, H. P. (2017). *The book of chocolate: The amazing story of the world's favorite candy*. Viking Press Inc. New York, NY (2017).
84. Macinnis, P. *Bittersweet: The story of sugar*. Sydney: Allen & Unwin (2002).
85. Mintz, S. *Sweetness and power: The place of sugar in modern history*. New York, NY: Penguin (1985).
86. Mintz, S. Sweetness and meaning. In C. Korsmeyer (Ed.), *The taste culture reader: Experiencing food and drink* (pp. 110-122). Oxford, UK: Berg (2005).
87. Spence, C., & Youssef, J. Personalized rock: A nostalgic fairground revival confection. *International Journal of Gastronomy & Food Science*, **17**:100150 (2020). <https://doi.org/10.1016/j.ijgfs.2019.100150>.
88. Muchembled, R. *Smells: A cultural history of odours in early modern times*. Cambridge, UK: Polity Press (2021).
89. Ogawa, K., Tashima, A., Sadakata, M., & Morinaga, O. Appetite-enhancing effects of vanilla flavours such as vanillin. *Journal of Natural Medicine*, **72**(3), 798-802 (2018). doi: 10.1007/s11418-018-1206-x.
90. Royal Society of Chemistry. Chemistry in its element – Vanillin (2016). <http://www.rsc.org/chemistryworld/podcast/CIEcompounds/transcripts/vanillin.asp>.
91. Dignum, M., Kerler, J., & Verpoorte, R. Vanilla production: Technological, chemical and biosynthetic aspects. *Food Research International*, **17**, 199-219 (2001).
92. Givaudan. Anything but plain vanilla. Givaudan taste essentials vanilla (2016). <https://www.givaudan.com/flavours/world-flavours/taste-essentials/vanilla>.
93. Derval, D. *The right sensory mix: Targeting consumer product development scientifically*. Heidelberg: Springer (2010).
94. Kuo, Y.-L., Pangborn, R. M., & Noble, A. C. Temporal patterns of nasal, oral, and retronasal perception of citral and vanillin and interaction of these odourants with selected tastants. *International Journal of Food Science and Technology*, **28**, 127-137 (1993).
95. Sakai, N., Kobayakawa, T., Gotow, N., Saito, S., & Imada, S. Enhancement of sweetness ratings of aspartame by a vanilla odor presented either by orthonasal or retronasal routes. *Perceptual and Motor Skills*, **92**, 1002-1008 (2001).
96. Oliveira, A. A. A., Andrade, A. C., Bastos, S. C., Condino, J. P. F., Júnior, A. C., & Pinheiro, A. C. M. Use of strawberry and vanilla natural flavors for sugar reduction: A dynamic sensory study with yogurt. *Food Research International*, **139**:109972 (2021). <https://doi.org/10.1016/j.foodres.2020.109972>.

97. Alcaire, F., Antúnez, L., Vidal, L., Giménez, A., & Ares, G. Aroma-related cross-modal interactions for sugar reduction in milk desserts: Influence on consumer perception. *Food Research International*, **97**, 45-50 (2017). <https://doi.org/10.1016/j.foodres.2017.02.019>.
98. Hoppert, K., Zahn, S., Jänecke, L., Mai, R., Hoffmann, S., & Rohm, H. Consumer acceptance of regular and reduced-sugar yogurt enriched with different types of dietary fiber. *International Dairy Journal*, **28**(1), 1-7 (2013). <https://doi.org/10.1016/j.idairyj.2012.08.005>.
99. Velázquez, A. L., Vidal, L., Varela, P., & Ares, G. Cross-modal interactions as a strategy for sugar reduction in products targeted at children: Case study with vanilla milk desserts. *Food Research International*, **130**:108920 (2020). <https://doi.org/10.1016/j.foodres.2019.108920>.
100. Wang, G., Hayes, J. E., Ziegler, G. R., Roberts, R. F., & Hopfer, H. Dose-response relationships for vanilla flavor and sucrose in skim milk: Evidence of synergy. *Beverages*, **4**:73 (2018). doi: 10.3390/beverages4040073.
101. Guo, J., Han, X., Zhan, J., You, Y., & Huang, W. Vanillin alleviates high fat diet-induced obesity and improves the gut microbiota composition. *Frontiers in Microbiology*, **9**:2733 (2018). <https://doi.org/10.3389/fmicb.2018.02733>.
102. Takai, E., Aoyagi, T., Ichikawa, K., Matsuura, Y., Kinoshita, F., & Takada, H. Effect of olfactory stimulation with vanilla odor on degree of electrical activity to control gastrointestinal motility. In *Universal Access in Human-Computer Interaction. Access to Media, Learning and Assistive Environments: 15th International Conference, UAHCI 2021, Held as Part of the 23rd HCI International Conference, HCII 2021, Virtual Event, July 24-29, 2021, Proceedings, Part II*. Springer-Verlag, Berlin, Heidelberg, 519-530 (2021). https://doi.org/10.1007/978-3-030-78095-1_38.
103. Kermen, F., Chakirian, A., Sezille, C., Joussain, P., Le Goff, G., Ziessel, A., Chastrette, M., Mandairon, N., Didier, A., Rouby, C., & Bensafi, M. Molecular complexity determines the number odour notes and the pleasantness of smells. *Scientific Reports*, **1**:206 (2011).
104. Sezille, C., Fournel, A., Rouby, C., Rinck, F., & Bensafi, M. Hedonic appreciation and verbal description of pleasant and unpleasant odors in untrained, trainee cooks, flavorists, and perfumers. *Frontiers in Psychology*, **5**:12 (2014). doi: 10.3389/fpsyg.2014.00012.
105. Zarzo, M. Hedonic judgments of chemical compounds are correlated with molecular size. *Sensors*, **11**, 3667-3686 (2011). doi:10.3390/s110403667.
106. Schiffman, S. S. Physiochemical correlates of olfactory quality. *Science*, **185**, 112-117 (1974).
107. Engen, T. *The acquisition of odour hedonics*. In S. Van Toller & G. H. Dodd (Eds.), *Perfumery: The psychology and biology of fragrance* (pp. 79-90). London, UK: Chapman & Hall (1988).
108. Hermans, D., & Baeyens, F. *Acquisition and activation of odor hedonics in everyday situations: Conditioning and priming studies*. In C. Rouby, B. Schaal, D. Dubois, R. Gervais, & A. Holley (Eds.), *Olfaction, taste, and cognition* (pp. 119-139). Cambridge, UK: Cambridge University Press (2002).
109. Sell, C. Structure-odor relationships: On the unpredictability of odor. *Angewandte Chemie International Edition*, **45**, 6254-6261 (2006).

110. Keller, A., Gerkin, R. C., Guan, Y., Dhurandhar, A., Turu, G., Szalai, B., Mainland, J. D., Ihara, Y., Yu, C. W., Wolfinger, R., et al. Predicting human olfactory perception from chemical features of odor molecules. *Science*, **355**, 820-826 (2017).
111. Clarke, R. J. *Coffee volume 1 chemistry*. New York, NY: Springer (2013).
112. Rapp, A. Natural flavours of wine: Correlation between instrumental analysis and sensory perception. *Fresenius' Journal of Analytical Chemistry*, **337**, 777-785 (1990).
113. Spence, C., & Wang, Q. J. On the meaning(s) of complexity in the chemical senses. *Chemical Senses*, **43**, 451-461 (2018).
114. Tao, Y. S., & Li, H. Active volatiles of cabernet sauvignon wine from Changli county. *Natural Science*, **1**, 176-182 (2009).
115. Dunkel, A., Steinhaus, M., Kotthoff, M., Nowak, B., Krautwurst, D., Schieberle, P., & Hofmann, T. Nature's chemical signatures in human olfaction: A foodborne perspective for future biotechnology. *Angewandte Chemie International Edition*, **53**, 7124-7143 (2014).
116. Gu, F., Chen, Y., Fang, Y., Wu, G., & Tan, L. Contribution of *Bacillus* isolates to the flavor profiles of vanilla beans assessed through aroma analysis and chemometrics. *Molecules*, **20**, 18422-18436 (2015).
117. Toth, S., Lee, K. J., Havkin-Frenkel, D., Belanger, F. C., & Hartman, T. G. Volatile compounds in vanilla. In D. Havkin-Frenkel & F. C. Belanger (Eds.), *Handbook of vanilla science and technology* (pp. 183-218). Oxford, UK: Wiley-Blackwell (2011).
118. Zhang, S., & Mueller, C. Comparative analysis of volatiles in traditionally cured bourbon and Ugandan Vanilla bean (*Vanilla planifolia*) extracts. *Journal of Agriculture and Food Chemistry*, **60**, 10433-10444 (2012).
119. Gleason-Allured, J. Vanilla: Anything but plain. *Perfumer and Flavorist* (2011). <http://www.perfumerflavorist.com/flavor/application/vanilla/132347233.html>.
120. Lapid, H., Harel, D., & Sobel, N. Prediction models for the pleasantness of binary mixtures in olfaction. *Chemical Senses*, **33**, 599-609 (2008).
121. Balogh, R. D., & Porter, R. H. Olfactory preferences resulting from mere exposure in human neonates. *Infant Behavior and Development*, **9**, 395-401 (1986).
122. Ishii, K. Does mere exposure enhance positive evaluation, independent of stimulus recognition? A replication study in Japan and the USA. *Japanese Psychological Research*, **47**, 280-285 (2005).
123. Monahan, J. L., Murphy, S. T., & Zajonc, R. B. Subliminal mere exposure: Specific, general and affective effects. *Psychological Science*, **11**, 462-466 (2000).
124. Pliner, P. The effects of mere exposure on liking for edible substances. *Appetite*, **3**, 283-290 (1982).
125. Varendi, H., Porter, R. H., & Winberg, J. Attractiveness for amniotic fluid odor: Evidence for prenatal learning? *Acta Paediatrica*, **85**, 1223-1227 (1996).
126. DeSnoo, K. Das trinkende kind im uterus [The drinking child in the uterus]. *Monatschrift für Geburtshilfe und Gynäkologie*, **105**, 88 (1937).
127. Marlier, L., Schaal, B., & Soussignan, R. Neonatal responsiveness to the odor of amniotic and lacteal fluids: A test of perinatal chemosensory continuity. *Child Development*, **69**, 611-623 (1998).

128. Schaal, B., Marlier, L., & Soussignan, R. Neonatal responsiveness to the odour of amniotic fluid. *Biology of Neonate*, **67**, 397-406 (1995). <https://doi.org/10.1159/000244192>.
129. Parker, G. H., & Stabler, E. M. On certain distinctions between taste and smell. *American Journal of Physiology*, **32**, 230-240 (1913).
130. Sullivan, R. M., Taborskybarba, S., Mendoza, R., Itano, A., Leon, M., Cotman, C. W., et al. Olfactory classical conditioningconditioning in. Nneonates. *Pediatrics*, **87**, 511-518 (1991).
131. Doucet, S., Soussignan, R., Sagot, P., & Schaal, B. The secretion of areolar (Montgomery's) glands from lactating women elicits selective, unconditional responses in neonates. *PLoS ONE*, **4(10)**: e7579 (2009). <https://doi.org/10.1371/journal.pone.0007579>.
132. Allen, J. New Yorkers sample cheese made from human breast milk. *Reuters*. **May 2nd** (2011). <http://www.reuters.com/article/us-cheese-breastmilk-idUSTRE7413X020110502>.
133. Chappell, B. Breast milk ice cream a hit at London store. *NPR*, **February 25th** (2011). <https://www.npr.org/sections/thetwo-way/2011/02/25/134056923/breast-milk-ice-cream-a-hit-at-london-store>.
134. Shah, K. Oh great, breast milk ice cream is back. "When it comes to ice cream this summer, breast is best." *Eater*, **April 24th** (2015). <https://www.eater.com/2015/4/24/8491303/breast-milk-ice-cream-london-the-lickators-royal-baby>.
135. Haller, R., Rummel, C., Henneberg, S., Pollmer, U., & Köster, E. The influence of early experience with vanillin in food preference later in life. *Chemical Senses*, **24**, 465-467 (1999). doi: 10.1093/chemse/24.4.465.
136. Mennella, J. A., & Beauchamp, G. The infant's responses to flavored milk. *Infant Behavior and Development*, **17**, 819 (1994).
137. Mennella, J. A., & Beauchamp, G. K. Early flavor experiences: When do they start? *Nutrition Today*, **29**, 25-31 (1994).
138. Mennella, J. A., & Beauchamp, G. K. The human infants' response to vanilla flavors in mother's milk and formula. *Infant Behavior and Development*, **19**, 13-19 (1996).
139. Mennella, J. A., & Beauchamp, G. K. The role of early life experiences in flavor perception and delight. In L. Dubé, A. Bechara, A. Dagher, A. Drewnowski, J. Lebel, P. James, Yada, R. Y., & M.-C. LaFlamme-Sanders (Eds.), *Obesity prevention: The role of brain and society on individual behavior* (pp. 203-217). Amsterdam, NL: Elsevier (2010).
140. Newman, J. How breast milk protects newborns. *Scientific American*, **December**, 76-79 (1995).
141. Hepper, P. G. Adaptive fetal learning. Prenatal exposure to garlic affects postnatal preferences. *Animal Behaviour*, **36**, 935-936 (1988).
142. Mennella, J. A., & Beauchamp, G. K. Maternal diet alters the sensory qualities of human milk and the nursing's behavior. *Pediatrics*, **88**, 737-744 (1991).
143. Mennella, J. A., & Beauchamp, G. K. The transfer of alcohol to human milk: Effects on flavor and the infant's behavior. *New England Journal of Medicine*, **325**, 981-985 (1991).
144. Pollmer, U. Novel foods: Flavour design and malnutrition. *Agro-Industrial High-Technology*, **2**, 43-45 (1991).

145. Capretta, P. J., Petersik, J. T., & Steward, D. J. Acceptance of novel flavours is increased after early experience of diverse taste. *Nature*, **254**, 689-691 (1975).
146. Forestell, C. A. Flavor perception and preference development in human infants. *Annals of Nutrition and Metabolism*, **70**(Suppl. 3), 17-25 (2017).
147. Galef, B. G., & Sherry, D. F. Mother's milk: A medium for transmission of cues reflecting the flavour of mother's diet. *Journal of Comparative Physiology and Psychology*, **83**, 374-378 (1973).
148. Jerome, N. W. Taste experience and the development of a dietary preference for sweet in humans: Ethnic and cultural variations in early taste experience. In J. M. Weiffenbach (Ed.), *Taste and development: The genesis of sweet preference* (pp. 235-248) Bethesda, MD: U.S. Department of Health, Education, and Welfare (1977).
149. Steiner, J. E. Human facial expressions in response to taste and smell stimulation. In H. W. Reese (Ed.), *Advances in child development and behaviour*, **13**, 257-295. Orlando, FL: Academic (1979).
150. Soussignan, R., Schaal, B., Marlier, L., & Jiang, T. Facial and autonomic responses to biological and artificial olfactory stimuli in human neonates: Re-examining early hedonic discrimination of odors. *Physiology & Behavior*, **62**(4), 745-758. doi: 10.1016/s0031-9384(97)00187-x (1997).
151. Mennella, J. A., & Beauchamp, G. K. Infants' exploration of scented toys: Effects of prior experience. *Chemical Senses*, **23**, 11-17 (1998).
152. Schaal, B., Marlier, L., & Soussignan, R. Human foetuses learn odours from their pregnant mother's diet. *Chemical Senses*, **25**, 729-737 (2000). <https://doi.org/10.1093/chemse/25.6.729>.
153. Mennella, J. A., Jagnow, C. P., & Beauchamp, G. K. Prenatal and postnatal flavor learning by human infants. *Pediatrics*, **107**:e88, 1-6 (2001).
154. Hausner, H., Bredie, W. L. P., Mølgaard, C., Petersen, M. A., & Møller, P. Differential transfer of dietary flavour compounds into human breast milk. *Physiology & Behavior*, **95**, 118-124 (2008).
155. Loos, H. M., Reger, D., & Schaal, B. The odour of human milk: Its chemical variability and detection by newborns. *Physiology & Behavior*, **199**, 88-99 (2019). doi:10.1016/j.physbeh.2018.11.008.
156. Ventura, A. K., & Mennella, J. A. Innate and learned preferences for sweet taste during childhood. *Current Opinion in Clinical Nutrition and Metabolic Care*, **14**(4), 379-384 (2011). <https://doi.org/10.1097/MCO.0b013e328346df65>.
157. Holy, T. E. Scenting Waldo: Analyzing olfactory scenes. *Nature Neuroscience*, **17**, 1144-1145 (2014).
158. Rokni, D., Hemmelder, V., Kapoor, V., & Murthy, V. N. An olfactory cocktail party: Figure-ground segregation of odorants in rodents. *Nature Neuroscience*, **17**, 1225-1232 (2014).
159. Rokni, D., & Murthy, V. N. Analysis and synthesis in olfaction. *ACS Chemical Neuroscience*, **5**, 870-872 (2014).
160. Spence, C. Flavour pairing: A critical review of the literature on food and beverage pairing. *Food Research International*, **133**:109124 (2020). <https://doi.org/10.1016/j.foodres.2020.109124>.

161. Spence, C. Gastrophysics: Getting creative with pairing flavours. *International Journal of Gastronomy & Food Science*, **27**:100433 (2022). <https://doi.org/10.1016/j.ijgfs.2021.100433>.
162. Moskowitz, H. R., Dravnieks, A., & Klarman, L. A. Odor intensity and pleasantness for a diverse set of odorants. *Perception & Psychophysics*, **19**, 122-128 (1976). <https://doi.org/10.3758/BF03204218>.
163. Doty, R., Brugger, W. E., Jurs, P. C., Orndorff, M. A., Snyder, P. J., & Lowry, L. D. Intranasal trigeminal stimulation from odorous volatiles: Psychometric responses from anosmic and normal humans. *Physiology & Behavior*, **20**, 175-185 (1978).
164. Frasnelli, J., Hummel, T., Berg, J., Huang, G., & Doty, R. L. Intranasal localizability of odorants: Influence of stimulus volume. *Chemical Senses*, **36**(4), 405-410 (2011). doi: 10.1093/chemse/bjr001.
165. Kobal, G., & Hummel, T. Olfactory and intranasal trigeminal event-related potentials in anosmic patients. *Laryngoscope*, **108**(7), 1033-1035 (1998). doi: 10.1097/00005537-199807000-00015.
166. Savic, I., Gulyás, B., & Berglund, H. Odorant differentiated pattern of cerebral activation: Comparison of acetone and vanillin. *Human Brain Mapping*, **17**(1), 17-27 (2002). doi: 10.1002/hbm.10045.
167. Wu, S. W., Fowler, D. K., Shaffer, F. J., Lindberg, J. E. M., & Peters, J. H. Ethyl vanillin activates TRPA1. *Journal of Pharmacology and Experimental Therapeutics*, **362**(3), 368-377 (2017). doi: 10.1124/jpet.116.239384.
168. Midroit, M., Chalençon, L., Renier, N., Milton, A., Thevenet, M., Sacquet, J., Breton, M., Forest, J., Noury, N., Richard, M., Raineteau, O., Ferdenzi, C., Fournel, A., Wesson, D. W., Bensafi, M., Didier, A., & Mandairon, N. Neural processing of the reward value of pleasant odorants. *Current Biology*, **31**(8), 1592-1605.e9 (2021). doi: 10.1016/j.cub.2021.01.066.
169. Wilson, D. A., & East, B. S. Good scents: A short road from olfaction to satisfaction. *Current Biology*, **31**(8), R374-R376 (2021). <https://doi.org/10.1016/j.cub.2021.01.099>.
170. Mennella, J. A., Pepino, M. Y., & Reed, D. R. Genetic and environmental determinants of bitter perception and sweet preferences. *Pediatrics*, **115**, e216-e222 (2005).
171. Eriksson, N., Wu, S., Do, C. B., Kiefer, A. K., Tung, J. Y., Mountain, J. L., Hinds, D. A., & Francke, U. A genetic variant near olfactory receptor genes influences cilantro preference. *Flavour*, **1**:22 (2012).
172. Mauer, L., & El-Sohemy, A. Prevalence of cilantro (*Coriandrum sativum*) disliking among different ethnocultural groups. *Flavour*, **1**:8 (2012).
173. McGee, H. Cilantro haters, it's not your fault. *The New York Times*, **April 13th** (2010). http://www.nytimes.com/2010/04/14/dining/14curious.html?_r=0.
174. Wysocki, C. J., & Beauchamp, G. K. Ability to smell androstenone is genetically determined. *Proceedings of the National Academy of Sciences USA*, **81**, 4899-4902 (1984).
175. Howgego, J. Sense for scents traced down to genes: Genetic factors for ability to smell violets and certain foods point to a future of 'personalized' scents and flavours. *Nature Neuroscience*, **August 1st** (2013). <http://www.nature.com/news/sense-for-scents-traced-down-to-genes-1.13493>.

176. Reed, D. R., & Knaapila, A. Genetics of taste and smell: Poisons and pleasures. *Progress in Molecular Biology Translational Science*, **94**, 213-240 (2010).
177. Jaeger, S. R., McRae, J. F., Bava, C. M., Beresford, M. K., Hunter, D., Jia, Y., Chheang, S. L., Jin, D., Peng, M., Gamble, J. C., Atkinson, K. R., Axten, L. G., Paisley, A. G., Tooman, L., Pineau, B., Rouse, S. A., & Newcomb, R. D. A Mendelian trait for olfactory sensitivity affects odor experience and food selection. *Current Biology*, **23**, 1601-1605 (2013).
178. McRae, J. F., Jaeger, S. R., Bava, C. M., Beresford, M. K., Hunter, D., Jia, Y., Chheang, S. L., Jin, D., Peng, M., Gamble, J. C., Atkinson, K. R., Axten, L. G., Paisley, A. G., Williams, L., Tooman, L., Pineau, B., Rouse, S. A., & Newcomb, R. D. Identification of regions associated with variation in sensitivity to food-related odors in the human genome. *Current Biology*, **23**, 1596-1600 (2013).
179. Pelchat, M. L., Bykowski, C., Dule, F. F., & Reed, D. R. Excretion and perception of a characteristic odor in urine after asparagus ingestion: A psychophysical and genetic study. *Chemical Senses*, **36**, 9-17 (2011).
180. Plotto, A., Barnes, K. W., & Goodner, K. L. Specific anosmia observed for β -ionone, but not for α -ionone: Significance for flavour research. *Journal of Food Science*, **71**, 401-406 (2006).
181. Simmen, D., & Briner, H. R. Olfaction in rhinology – methods of assessing the sense of smell. *Rhinology*, **44**, 98-101 (2006).
182. Axel, R. The molecular logic of smell. *Scientific American*, **273(October)**, 154-159 (1995).
183. Ravia, A., Snitz, K., Honigstein, D., Finkel, M., Zirler, R., Perl, O., Secundo, L., Laudamiel, C., Harel, D., & Sobel, N. A measure of smell enables the creation of olfactory metamers. *Nature*, **588**, 118-123 (2020). <https://doi.org/10.1038/s41586-020-2891-7>.
184. Schaal, B., Saxton, T. K., Loos, H., Soussignan, R., & Durand, K. Olfaction scaffolds the developing human from neonate to adolescent and beyond. *Philosophical Transactions of the Royal Society London B Biological Sciences*, **375**:20190261 (2020).
185. Wilson, D. A., & Stevenson, R. J. *Learning to smell*. Baltimore: Johns Hopkins University Press (2006).
186. Blank, D. M., & Mattes, R. D. Sugar and spice: Similarities and sensory attributes. *Nursing Research*, **39**, 290-293 (1990).
187. Hsiao, Y.-Y., Pan, Z.-J., Hsu, C.-C., Yang, Y.-P., Hsu, Y.-C., Chuang, Y.-C., Shih, H.-H., Chen, W.-H., Tsai, W.-C., & Chen, H.-H. Research on orchid biology and biotechnology. *Plant and Cell Physiology*, **52(9)**, 1467-1486 (2011). <https://doi.org/10.1093/pcp/pcr100>.
188. Herz, R. S., Larsson, M., Trujillo, R., Casola, M. C., Ahmed, F. K., Lipe, S., & Brashear, M. E. A three-factor benefits framework for understanding consumer preference for scented household products: psychological interactions and implications for future development. *Cognitive Research: Principles and Implications*, **7**:28 (2022). <https://doi.org/10.1186/s41235-022-00378-6>.
189. Wang, X., Chen, Y., Wang, Q., Sun, L., Li, G., Zhang, C., et al. Support for natural small-molecule phenols as anxiolytics. *Molecules*, **22**:E2138 (2017). doi: 10.3390/molecules22122138.

190. Hunt, S. R. Sugar and spice. *Pharmaceutical Journal*, **191**, 632-635 (1963).
191. Beauchamp, G. K. Why do we like sweet taste: A bitter tale? *Physiology and Behavior*, **164**, 432-437 (2016). <https://doi.org/10.1016/j.physbeh.2016.05.007>.
192. Beidler, L. M. The biological and cultural role of sweeteners. In *Sweeteners: Issues and uncertainties* (pp. 11-18). National Academy of Sciences Academy Forum, Fourth of a Series. Washington, DC: National Academy of Sciences (2016).
193. Moskowitz, H. The psychology of sweetness. In H. L. Sipple & K. W. McNutt (Eds.), *Sugars in nutrition* (pp. 37-64). New York, NY: Academic Press (1974).

CS (charles.spence@psy.ox.ac.uk) is the author to whom correspondence and requests for materials should be addressed.

Competing financial interests: The author declares no competing financial interests.