

**Age and sex-specific associations of carotid pulsatility with small vessel disease burden  
in TIA and ischaemic stroke**

#Kui Kai Lau, MRCP;<sup>1</sup> #Petra Pego, MD;<sup>1</sup> Sara Mazzucco, MD PhD;<sup>1</sup> Linxin Li, MD DPhil;<sup>1</sup>  
Dominic PJ Howard, MA, MRCS, DPhil;<sup>1,2</sup> Wilhelm Küker, MD FRCR;<sup>1</sup>  
\*Peter M Rothwell, MD PhD FRCP FMedSci<sup>1</sup>

<sup>1</sup>Centre for Prevention of Stroke and Dementia, Nuffield Department of Clinical  
Neurosciences University of Oxford

<sup>2</sup>Department of Vascular Surgery, Oxford University Hospitals NHS Trust, Oxford, United  
Kingdom

# These authors contributed equally

\* Corresponding author contact details:

Professor Peter M Rothwell

Tel 0044 1865 231603

Fax 0044 1865 234639

Email: [peter.rothwell@ndcn.ox.ac.uk](mailto:peter.rothwell@ndcn.ox.ac.uk)

Number of tables: 4

Number of figures: 0

Number of supplementary tables: 2

Word count abstract: 250

Word count text: 2782

**Cover title:** carotid pulsatility and small vessel disease

**List of Tables:**

Table 1. Clinical and imaging characteristics of the study population

Table 2. Clinical predictors of common carotid and internal carotid artery pulsatility index

Table 3. Relationships of common carotid and internal carotid artery pulsatility index with individual neuroimaging markers and global burden of small vessel disease, stratified by age

Table 4. Relationships of common carotid and internal carotid artery pulsatility index with individual neuroimaging markers and global burden of small vessel disease, stratified by sex

**List of Supplementary Tables:**

Table I. Image sequence parameters of the study population

Table II. Aetiology of TIA/ischaemic strokes according to TOAST classification

**Indexing terms:** carotid pulsatility, small vessel disease, lacunes, stroke, transient ischaemic attack

**Contributors:**

KKL collected data, did the statistical analysis and interpretation, wrote and revised the manuscript. PP, SM, LL and DPJH collected data. WK provided study supervision and acquired imaging data. PMR conceived and designed the overall study, provided study supervision and funding, acquired, analysed and interpreted data, and wrote and revised the manuscript.

## **Abstract**

**Background and Purpose:** Although large artery stiffness has been implicated in the pathogenesis of cerebral small vessel disease (SVD), whether carotid pulsatility, a convenient surrogate marker of arterial stiffness, is similarly associated with global burden of SVD is unknown.

**Methods:** We studied consecutive patients with TIA or non-disabling ischaemic stroke from the Oxford Vascular Study who had a brain MRI and carotid duplex ultrasound during 2002-2014. We determined clinical correlates of common carotid (CCA) and internal carotid artery (ICA) pulsatility index (PI) and their associations with the Total SVD Score on MRI, stratified by age (median=72).

**Results:** In 587 patients, CCA and ICA-PI were both independently associated with age, diabetes and premorbid mean pulse pressure after adjustment for age, sex and cardiovascular risk factors (all  $p < 0.05$ ). ICA-PI was strongly associated with SVD markers and burden, particularly lacunes, in patients aged  $< 70$  (age and sex-adjusted OR of top vs. bottom PI quartile: 5.35, 1.95-14.70,  $p = 0.001$ ; increasing SVD Score: 2.30, 1.01-5.25,  $p = 0.048$ ), but not in patients aged  $\geq 70$  ( $p > 0.05$ ). No associations between CCA-PI with SVD Score were noted at any age. In 94 consecutive patients who also received transcranial Doppler ultrasound, strong associations between middle cerebral artery (MCA)-PI and an increasing SVD Score was noted (unadjusted OR - MCA: 4.26, 1.45-12.55,  $p = 0.009$ ; ICA: 2.37, 0.81-6.87,  $p = 0.11$ ; CCA: 1.33, 0.45-3.96,  $p = 0.61$ ).

**Conclusions:** ICA and MCA-PI are associated with global SVD burden, especially in individuals aged  $< 70$  and may be causally related.

## Introduction

Cerebral small vessel disease (SVD) accounts for approximately 20% of all strokes and up to half of all dementias.<sup>1</sup> However, although hypertension is one of the main risk factors leading to SVD, the pathogenesis of SVD is not fully understood.<sup>1</sup>

Stiffening of the large vessels has been associated with hypertension and with the pathogenesis of cerebral SVD.<sup>2-10</sup> Previous studies have demonstrated that large artery stiffness is associated with silent cerebral infarcts,<sup>4, 5, 7</sup> cerebral microbleeds,<sup>6</sup> white matter hyperintensity (WMH)<sup>3-5, 7, 8</sup> and strokes of lacunar subtype.<sup>10</sup> Furthermore, stiffness of the aorta is associated with an increased arterial pulsatility measured at the common carotid artery (CCA)<sup>3, 10</sup> and middle cerebral artery (MCA).<sup>9, 11</sup> Studies have also shown that MCA pulsatility is correlated with leukoaraiosis,<sup>9</sup> suggesting that leukoaraiosis and lacunar infarcts may result partly from increased arterial pulsatility transmitted to the cerebral vessels secondary to large artery stiffness.<sup>2-5, 8, 9</sup>

However, causation is uncertain. The burden of leukoaraiosis varies with age and sex, being more severe in women than in men at older ages,<sup>12</sup> and could be due to differences in pulsatility. Sex differences are also seen in the carotid bifurcation and internal carotid artery (ICA) anatomy with the ICA/CCA ratio and ICA/external carotid artery ratio being greater in women compared with men.<sup>13</sup> Moreover, large inter- and intraindividual variations in carotid bifurcation anatomy exist<sup>14</sup> and could account for some unexplained susceptibility to developing SVD at an individual level. These individual and sex differences in carotid anatomy could influence associations between carotid pulsatility and global SVD burden. If so, one might expect to see stronger associations between ICA pulsatility with SVD than with CCA pulsatility.

We therefore investigated the age and sex-specific associations of ICA versus CCA pulsatility with individual neuroimaging markers and global burden of SVD (Total SVD

Score) in patients with transient ischaemic attack (TIA) or ischaemic stroke, nested in the population-based Oxford Vascular Study (OXVASC). In a subset of patients, we also compared the relationships of carotid artery and MCA pulsatility with the Total SVD Score.

## Methods

We prospectively studied patients with TIA or non-disabling ischaemic stroke from OXVASC. In brief, OXVASC is an on-going population-based study of all acute vascular events occurring within a population of 92728 individuals, irrespective of age, who are registered with 100 general practitioners in nine general practices of Oxfordshire, UK.<sup>15</sup> The analysis herein includes 606 cases of TIA/ischaemic stroke recruited from November 1, 2004 to September 30, 2014 who had a cerebral magnetic resonance imaging (MRI) and also a carotid duplex ultrasound. The imaging protocol of OXVASC has been described elsewhere.<sup>16, 17</sup> Briefly, from April 1, 2002 - March 31, 2010 (phase 1), non-contrast CT brain and carotid duplex ultrasound were the first-line investigations and MRI and MR angiography was performed in selected patients when clinically indicated. From April 1, 2010 onwards (phase 2), brain MRI and MR angiography became the first-line imaging methods. However, carotid duplex ultrasound was also performed in selected patients in instances where MRI was contraindicated, MR angiography was not available or if carotid stenosis was identified on MR angiography and required more accurate quantification of stenosis via ultrasonography. At the end of the study period, 94 consecutive patients also received a transcranial Doppler (TCD) ultrasound upon ascertainment.

We collected demographic data, atherosclerotic risk factors, details of hospitalisation of index event during face-to-face interview and cross-referenced these with primary care records and hospital records. Cause of TIA/ischaemic stroke was classified according to the modified Trial of Org 10172 in Acute Stroke Treatment (TOAST) criteria.<sup>18</sup> Premorbid blood pressure readings for all 606 patients during the 20 years prior to ascertainment (13404 readings in total, median number of readings per patient: 16, interquartile range [IQR]: 7-33) was also retrieved from the primary care practices and the mean systolic blood pressure, diastolic blood pressure and pulse pressure used for analysis.

Patients were scanned predominantly (488/606) with 2 scanners – Achieva, Philips Healthcare (369/443 patients who received a 1.5T MRI), and Magnetom Verio, Siemens Healthcare (119/163 patients who received a 3T MRI). Details of scan parameters are documented in Supplementary Table I. PVSs were defined as small (<3mm) punctate (if perpendicular to the plane of scan) or linear (if longitudinal to the plane of scan) hyperintensities on T2 images in the basal ganglia (BG) and centrum-semiovale (CS) based on a previously validated scale.<sup>19</sup> Burden of PVSs were then stratified into 3 groups: <11, 11-20 and >20. The severity of subcortical and periventricular WMH was determined for each patient according to the Fazekas scale.<sup>20</sup> Cerebral microbleeds were defined as rounded, hypodense foci up to 10mm in size and were differentiated from microbleed mimics based on current guidelines.<sup>21</sup> The location and number of microbleeds were scored according to the Microbleed Anatomical Rating Scale.<sup>22</sup> Lacunes were defined as rounded or ovoid lesions, >3 and <20mm in diameter, in the BG, internal capsule, CS or brainstem, of cerebrospinal fluid signal density on T2 and fluid-attenuated inversion recovery and no increased signal on diffusion weighted imaging.<sup>23</sup> Burden of SVD was estimated by calculating the Total SVD Score where one point is allocated to each of the following: 1) presence of lacunes, 2) presence of microbleeds, 3) moderate-severe (>10) BG-PVSs and 4) severe periventricular and/or moderate-severe deep WMH.<sup>24</sup>

One neuroradiologist (WK) provided ongoing supervision of interpretation of the MRI images during the study period. Definitions of neuroimaging biomarkers were based on STRIVE.<sup>23</sup> The intra-rater  $\kappa$  for 50 randomly selected scans was: lacunes - 0.85, microbleed burden (0, 1, 2-4,  $\geq 5$ ) – 0.88, periventricular WMH burden (Fazekas grade 0, 1, 2, 3) – 0.75; subcortical WMH burden (Fazekas grade 0, 1, 2, 3) WMH burden – 0.80 and PVS burden (<11, 11-20, >20) - 0.86 (BG) and 0.84 (CS).

Carotid duplex ultrasound scans were performed at the Vascular Laboratory, Oxford Regional Vascular Unit of John Radcliffe Hospital, Oxford, according to current guidelines.<sup>25</sup>

The ultrasound protocol included B-mode imaging (transverse and longitudinal plane) of the proximal, mid and distal CCA, carotid bifurcation and proximal, mid and distal extracranial ICA. Presence of carotid plaques and plaque morphology were documented. Peak systolic velocity (PSV) and end-diastolic velocity (EDV) were measured by pulsed-wave Doppler analysis on the CCA and extracranial ICA bilaterally. In a subgroup of patients, intracranial vessels were examined by means of TCD sonography (DopplerBox, Compumedics DWL, Singen, Germany). MCA blood flow velocities were recorded with a handheld 2MHz probe through temporal bone window at the depth that provided the best signal. Each session was stored in the hard disc of the TCD device for subsequent off-line analysis.<sup>26</sup> PI was calculated according to Gosling's pulsatility index as the (peak systolic velocity – end-diastolic velocity)/mean flow velocity for the explored extracranial and intracranial arteries (CCA, ICA and MCA).<sup>27</sup>

Patients gave written informed consent after an event or assent was obtained from relatives for patients who were unable to provide consent. OXVASC was approved by the local research ethics committee.

### **Statistical analysis**

The mean CCA, ICA and MCA-PI of the left and right carotid or cerebral arteries were used for analysis if neither artery had >50% stenosis. In patients who had unilateral >50% stenosis of the CCA, ICA or MCA, the contralateral CCA, ICA or MCA-PI were used for analysis. Patients were excluded from the main analysis if they had bilateral >50% stenosis of the CCA, ICA and/or MCA.

We determined the clinical predictors of CCA and ICA-PI by linear regression in a univariate model, model adjusted for age and sex as well as a multi-variate model adjusted for all co-variates (age, sex, vascular risk factors, glomerular filtration rate, and premorbid mean systolic blood pressure, diastolic blood pressure and pulse pressure). We also determined, by



ordinal regression, the odds ratios (OR) of lacunes, an increasing burden of subcortical and periventricular WMH (Fazekas grade 0, 1, 2, 3), cerebral microbleeds (0, 1, 2-4,  $\geq 5$ ), BG and CS-PVSs (<11, 11-20,  $\geq 20$ ) and Total SVD Score (0, 1, 2, 3, 4) in patients with the top CCA or ICA-PI quartile, compared with those in the bottom CCA or ICA-PI quartile as reference, in an analysis stratified by median age (<70 vs.  $\geq 70$ ) and by sex. In the 94/606 patients who also had TCD ultrasound performed and with satisfactory bone window, we similarly determined by ordinal regression the ORs of an increasing Total SVD Score in patients with the top MCA-PI quartile, compared with those in the bottom quartile as reference.

All analyses were done with SPSS version 22.

### **Role of the funding source**

The funding source had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had the final responsibility for the decision to submit for publication.

## Results

A total of 606 patients with TIA or non-disabling stroke had a MRI and carotid Doppler ultrasound during the study period. After excluding 29 patients who had bilateral CCA or ICA stenosis >50%, 587 patients (306 TIA, 281 ischemic stroke) were included in the final analysis. Carotid Doppler ultrasound was performed on a median of 2 days (IQR 0-6) after TIA/ischaemic stroke presentation. Baseline clinical and imaging characteristics of patients are shown in Table 1. The mean (SD) age of the population was 70 (14) years and 51% were males. Lacunes were present in 20.4% of the study population, 15.7% had microbleeds, 11.2% with Fazekas grade 3 subcortical WMH, 10.9% with Fazekas grade 3 periventricular WMH, 20.8% with >20 BG-PVSs and 42.9% with >20 CS-PVSs (Table 1). The mean (SD) Total SVD Score was 1.16 (1.17). The mean (SD) CCA-PI was 1.57 (0.29), mean ICA-PI was 1.28 (0.27) and mean MCA-PI was 1.08 (0.24).

After adjusting for age, sex, vascular risk factors and premorbid blood pressure, age, diabetes and premorbid mean pulse pressure were identified as independent predictors of CCA-PI (Table 2). Male sex was associated with an increasing CCA-PI (0.119, 0.074-0.164,  $p<0.0001$ ). However, this relationship was lost for ICA-PI ( $p=0.40$ ) (Table 2).

We studied the associations between CCA and ICA-PI with an increasing burden of individual neuroimaging markers of SVD and global burden of SVD stratified by age (Table 3). In the 261/587 patients aged <70, significant associations between ICA-PI with lacunes (5.35, 1.95-14.70,  $p=0.001$ ), increasing burden of periventricular WMH (2.86, 1.26-6.46,  $p=0.012$ ), BG-PVSs (2.83, 1.07-7.46,  $p=0.035$ ) and Total SVD Score (2.30, 1.01-5.25,  $p=0.048$ ) was noted, even after adjusting for age and sex. However, there were no relationships between ICA-PI with neuroimaging markers of SVD in the 326/587 patients aged  $\geq 70$  (Total SVD Score – ICA: 1.20, 0.59-2.42,  $p=0.62$ ) (Table 3). No relationships between CCA-PI with Total SVD Score were noted in patients aged <70 (1.08, 0.52-2.25,

p=0.84) or  $\geq 70$  (1.20, 0.66-2.21, p=0.55) (Table 3). We also noted no sex-differences in between CCA-PI and ICA-PI with neuroimaging markers of SVD in men (Total SVD Score – CCA: 1.84, 1.01-3.35, p=0.046; ICA: 3.72, 2.01-6.88, p<0.0001) and women (CCA: 3.12, 1.67-5.82, p=0.0004,  $p_{het}=0.23$ ; ICA: 6.06, 3.20-11.47, p<0.0001;  $p_{het}=0.28$ ) (Table 4).

In 94 consecutive patients who also had a TCD ultrasound performed, we noted stronger univariate associations between MCA-PI and ICA-PI (B 0.499, 0.341-0.657, p<0.0001) than between MCA-PI and CCA-PI (0.353, 0.193-0.512, p<0.0001). Risk associations with Total SVD Score increased progressively when arteries in closer proximity to the brain parenchyma were imaged (unadjusted OR of top vs. bottom quartile of PI – CCA: 1.33, 0.45-3.96, p=0.61; ICA: 2.37, 0.81-6.87, p=0.11; MCA: 4.26, 1.45-12.55, p=0.008). However, due to the small number of patients who had TCD ultrasound performed, this analysis was underpowered and we did not adjust for confounding factors.

## Discussion

In patients with TIA or non-disabling ischaemic stroke, we demonstrated significant age-specific associations between ICA-PI with neuroimaging markers of SVD. In patients aged <70, ICA-PI was most significantly associated with lacunes, but associations were also noted between ICA-PI with periventricular WMH and BG-PVS burden, as well as the overall burden of SVD. However, no associations between neuroimaging markers of SVD and ICA-PI were noted in patients aged  $\geq 70$ . We also demonstrated that the overall associations between PI and SVD burden were in general stronger with more distal arterial beds, and that in a subset of individuals, SVD burden appeared most strongly associated with MCA-PI, followed by ICA-PI and then CCA-PI.

A number of previous studies have demonstrated that PI, measured at the CCA or MCA is a reflection of underlying large artery stiffness.<sup>9-11</sup> Our results support these findings, as age, diabetes and premorbid pulse pressure, all of which have previously been associated with arterial stiffness,<sup>28-30</sup> were noted to be independent predictors of CCA and ICA-PI in our study. Male sex was also a strong independent predictor of CCA-PI, but we did not note any associations between ICA-PI and sex on univariate analysis ( $p=0.40$ ). These observations may be due to the underlying sex differences in carotid bifurcation and ICA anatomy, as although men may have larger CCA and ICA lumen sizes compared with women, the ICA/CCA ratio and ICA/external carotid artery ratio have been noted to be greater in women compared with men.<sup>13</sup> However, despite known sex-differences in carotid anatomy, we were not able to delineate any significant sex differences in risk associations between CCA and ICA-PI with SVD burden.

In common with previous studies that have noted associations between large artery stiffness or arterial pulsatility with leukoaraiosis,<sup>3-5, 7-9</sup> there were significant associations between ICA-PI with periventricular WMH burden in all patients. However, we also found significant

age-specific associations between carotid PI and SVD, such that in younger individuals age <70, additional associations between ICA-PI with lacunes, BG-PVS and global SVD burden were present, but these associations were attenuated in patients aged  $\geq 70$ . Age is perhaps the most important independent predictor of arterial stiffness<sup>28</sup> and cerebral SVD,<sup>16, 31</sup> and hence any associations between carotid PI and SVD burden would be confounded by age.

In a small subgroup of consecutive patients who also had TCD ultrasound performed, we noted that the association between PI and SVD burden appeared to increase progressively with more distal arterial beds (MCA>ICA>CCA). Together with findings from previous studies,<sup>3-5, 8, 9</sup> our results support the hypothesis that arterial stiffening results in an increased pulsatile flow that is propagated distally, along the large arterial beds, and is subsequently transmitted to the cerebral small vessels.<sup>2-5, 8</sup> This process may then result in the various parenchymal lesions as a consequence of cerebral SVD (e.g. leukoaraiosis, lacunes and BG-PVSs), possibly due to cerebrovascular endothelial failure, blood brain barrier dysfunction, alterations in perfusion during diastole, increased endothelial shear stress and/or impaired cerebral autoregulation.<sup>1, 2</sup> However, it is uncertain how much MCA-PI is also a measure of the resistance characteristics of the distal small vessel vascular bed, and hence how much the MCA-PI risk association is due to reverse causation is unknown.

Our study has a number of limitations. Firstly, the series of patients who received a TCD ultrasound was small and although our findings have been consistent with those from other studies,<sup>6, 7, 9-11</sup> our results would need to be confirmed in larger cohorts. Second, in contrast to previous studies,<sup>6</sup> we were not able to demonstrate significant associations between carotid pulsatility and cerebral microbleed burden. Cerebral microbleeds could be secondary to hypertensive or cerebral amyloid angiopathy. Our cohort however, lacked power to determine whether carotid PI was a stronger predictor of strictly deep-seated microbleeds (i.e. suggestive of hypertensive angiopathy, only 7/587 patients with strictly deep microbleeds in our cohort) or strictly lobar microbleeds (suggestive of cerebral amyloid angiopathy, 44/587

patients with strictly lobar microbleeds in our cohort). Third, although we demonstrated that ICA and MCA-PI may be surrogate markers of underlying SVD burden, their prognostic value over and above traditional vascular risk factors or neuroimaging findings in patients with TIA/ischaemic stroke remains uncertain. Similarly, whether ICA or MCA-PI are able to predict progression of SVD burden was not studied in our cohort. Finally, our cohort spanned a 10-year period, during which the neuroimaging protocol of OXVASC has changed. However, when we stratified our analyses by MRI scanner, we noted similar associations between CCA and ICA-PI with Total SVD Score with no heterogeneity (CCA:  $p_{\text{het}}=0.16$ , ICA:  $p_{\text{het}}=0.090$ ). Similarly, we have recently shown in our cohort that the prognostic value for recurrent stroke of individual neuroimaging markers of SVD, such as PVSs,<sup>16</sup> as well as the Total SVD Score<sup>17</sup> were robust to the variations in scanner type and sequences used in our study.

In conclusion, our findings suggest that ICA-PI and MCA-PI are surrogate markers of large artery stiffness and underlying SVD burden and support the hypothesis that stiffening of the large arteries results in increased transmission of pulsatile flow along the carotid and cerebral arteries and may play a role in the pathophysiology of cerebral SVD. More research is required to determine how much MCA-PI reflects input pressure or flow characteristics as compared with distal resistance.

## **Acknowledgements**

We are grateful to all the staff in the general practices that collaborated in the Oxford Vascular Study: Abingdon Surgery, Stert St, Abingdon; Malthouse Surgery, Abingdon; Marcham Road Family Health Centre, Abingdon; The Health Centre, Berinsfield; Key Medical Practice; Kidlington; 19 Beaumont St, Oxford; East Oxford Health Centre, Oxford; Church Street Practice, Wantage. We also acknowledge the use of the facilities of the Acute Vascular Imaging Centre, Oxford.

## **Sources of Funding**

The Oxford Vascular Study is funded by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC), Wellcome Trust, Wolfson Foundation, British Heart Foundation and the European Union's Horizon 2020 programme (grant 666881, SVDs@target). Professor Rothwell is in receipt of a NIHR Senior Investigator award. KKL is funded by a University of Oxford Croucher Scholarship. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

## **Declaration of interests**

We declare no competing interests.

## References

1. Wardlaw JM, Smith C, Dichgans M. Mechanisms of sporadic cerebral small vessel disease: Insights from neuroimaging. *The Lancet. Neurology*. 2013;12:483-497
2. O'Rourke MF, Safar ME. Relationship between aortic stiffening and microvascular disease in brain and kidney: Cause and logic of therapy. *Hypertension*. 2005;46:200-204
3. Mitchell GF, van Buchem MA, Sigurdsson S, Gotal JD, Jonsdottir MK, Kjartansson O, et al. Arterial stiffness, pressure and flow pulsatility and brain structure and function: The age, gene/environment susceptibility--reykjavik study. *Brain*. 2011;134:3398-3407
4. Brandts A, van Elderen SG, Westenberg JJ, van der Grond J, van Buchem MA, Huisman MV, et al. Association of aortic arch pulse wave velocity with left ventricular mass and lacunar brain infarcts in hypertensive patients: Assessment with mr imaging. *Radiology*. 2009;253:681-688
5. Henskens LH, Kroon AA, van Oostenbrugge RJ, Gronenschild EH, Fuss-Lejeune MM, Hofman PA, et al. Increased aortic pulse wave velocity is associated with silent cerebral small-vessel disease in hypertensive patients. *Hypertension*. 2008;52:1120-1126
6. Seo WK, Lee JM, Park MH, Park KW, Lee DH. Cerebral microbleeds are independently associated with arterial stiffness in stroke patients. *Cerebrovasc Dis*. 2008;26:618-623
7. Tsao CW, Seshadri S, Beiser AS, Westwood AJ, Decarli C, Au R, et al. Relations of arterial stiffness and endothelial function to brain aging in the community. *Neurology*. 2013;81:984-991
8. van Elderen SG, Brandts A, Westenberg JJ, van der Grond J, Tamsma JT, van Buchem MA, et al. Aortic stiffness is associated with cardiac function and cerebral



- small vessel disease in patients with type 1 diabetes mellitus: Assessment by magnetic resonance imaging. *Eur Radiol*. 2010;20:1132-1138
9. Webb AJ, Simoni M, Mazzucco S, Kuker W, Schulz U, Rothwell PM. Increased cerebral arterial pulsatility in patients with leukoaraiosis: Arterial stiffness enhances transmission of aortic pulsatility. *Stroke; a journal of cerebral circulation*. 2012;43:2631-2636
  10. Wohlfahrt P, Krajcoviechova A, Jozifova M, Mayer O, Vanek J, Filipovsky J, et al. Large artery stiffness and carotid flow pulsatility in stroke survivors. *J Hypertens*. 2014;32:1097-1103; discussion 1103
  11. Kim JY, Bushnell CD, Park JH, Han SM, Im JH, Han SW, et al. Central aortic pressure and pulsatility index in acute ischemic stroke. *J Neuroimaging*. 2015;25:438-442
  12. Simoni M, Li L, Paul NL, Gruter BE, Schulz UG, Kuker W, et al. Age- and sex-specific rates of leukoaraiosis in tia and stroke patients: Population-based study. *Neurology*. 2012;79:1215-1222
  13. Schulz UG, Rothwell PM. Sex differences in carotid bifurcation anatomy and the distribution of atherosclerotic plaque. *Stroke*. 2001;32:1525-1531
  14. Schulz UG, Rothwell PM. Major variation in carotid bifurcation anatomy: A possible risk factor for plaque development? *Stroke*. 2001;32:2522-2529
  15. Rothwell PM, Coull AJ, Giles MF, Howard SC, Silver LE, Bull LM, et al. Change in stroke incidence, mortality, case-fatality, severity, and risk factors in oxfordshire, uk from 1981 to 2004 (oxford vascular study). *Lancet*. 2004;363:1925-1933
  16. Lau KK, Li L, Lovelock CE, Zamboni G, Chan TT, Chiang MF, et al. Clinical correlates, ethnic differences, and prognostic implications of perivascular spaces in transient ischemic attack and ischemic stroke. *Stroke; a journal of cerebral circulation*. 2017;48:1470-1477

17. Lau KK, Li L, Schulz U, Simoni M, Chan KH, Ho SL, et al. Total small vessel disease score and risk of recurrent stroke: Validation in 2 large cohorts. *Neurology*. 2017;88:2260-2267
18. Adams HP, Jr., Bendixen BH, Kappelle LJ, Biller J, Love BB, Gordon DL, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. Toast. Trial of org 10172 in acute stroke treatment. *Stroke; a journal of cerebral circulation*. 1993;24:35-41
19. Potter GM, Chappell FM, Morris Z, Wardlaw JM. Cerebral perivascular spaces visible on magnetic resonance imaging: Development of a qualitative rating scale and its observer reliability. *Cerebrovascular diseases*. 2015;39:224-231
20. Fazekas F, Chawluk JB, Alavi A, Hurtig HI, Zimmerman RA. Mr signal abnormalities at 1.5 t in alzheimer's dementia and normal aging. *AJR. American journal of roentgenology*. 1987;149:351-356
21. Greenberg SM, Vernooij MW, Cordonnier C, Viswanathan A, Al-Shahi Salman R, Warach S, et al. Cerebral microbleeds: A guide to detection and interpretation. *The Lancet. Neurology*. 2009;8:165-174
22. Gregoire SM, Chaudhary UJ, Brown MM, Yousry TA, Kallis C, Jager HR, et al. The microbleed anatomical rating scale (mars): Reliability of a tool to map brain microbleeds. *Neurology*. 2009;73:1759-1766
23. Wardlaw JM, Smith EE, Biessels GJ, Cordonnier C, Fazekas F, Frayne R, et al. Neuroimaging standards for research into small vessel disease and its contribution to ageing and neurodegeneration. *The Lancet. Neurology*. 2013;12:822-838
24. Staals J, Makin SD, Doubal FN, Dennis MS, Wardlaw JM. Stroke subtype, vascular risk factors, and total mri brain small-vessel disease burden. *Neurology*. 2014;83:1228-1234

25. Oates CP, Naylor AR, Hartshorne T, Charles SM, Fail T, Humphries K, et al. Joint recommendations for reporting carotid ultrasound investigations in the united kingdom. *Eur J Vasc Endovasc Surg*. 2009;37:251-261
26. Mazzucco S, Li L, Tuna MA, Pendlebury ST, Wharton R, Rothwell PM, et al. Hemodynamic correlates of transient cognitive impairment after transient ischemic attack and minor stroke: A transcranial doppler study. *Int J Stroke*. 2016;11:978-986
27. Oughton JA, Rose S, Galloway G, Khoo SK, O'Neill S, Coulthard A. Carotid ultrasound pulsatility indices and cardiovascular risk in australian women. *J Med Imaging Radiat Oncol*. 2015;59:20-25
28. Lee HY, Oh BH. Aging and arterial stiffness. *Circ J*. 2010;74:2257-2262
29. Prenner SB, Chirinos JA. Arterial stiffness in diabetes mellitus. *Atherosclerosis*. 2015;238:370-379
30. Van Bortel LM, Struijker-Boudier HA, Safar ME. Pulse pressure, arterial stiffness, and drug treatment of hypertension. *Hypertension*. 2001;38:914-921
31. van Dijk EJ, Prins ND, Vrooman HA, Hofman A, Koudstaal PJ, Breteler MM. Progression of cerebral small vessel disease in relation to risk factors and cognitive consequences: Rotterdam scan study. *Stroke*. 2008;39:2712-2719

**Table 1. Clinical and imaging characteristics of the study population**

	<b>N=587 (TIA N=306, ischaemic stroke N=281)</b>
<b>Clinical characteristics</b>	
Mean age, yr (SD)	69.5 (13.6)
Males (%)	300 (51.1)
Hypertension (%)	342 (58.3)
Hyperlipidaemia (%)	245 (41.7)
Diabetes (%)	85 (14.5)
Ever-smokers (%)	333 (56.7)
Atrial fibrillation (%)	81 (13.8)
Prior transient ischaemic attack / stroke (%)	133 (22.7)
Ischaemic heart disease (%)	103 (16.7)
Glomerular filtration rate, ml/min/1.73m <sup>2</sup>	71.1 (21.4)
<b>Imaging characteristics</b>	
Common carotid artery pulsatility index (SD)	1.57 (0.29)
Internal carotid artery pulsatility index (SD)	1.28 (0.27)
Middle cerebral artery pulsatility index (SD)*	1.08 (0.24)
N with diffusion weighted imaging positive lesion (%)	125 (21.3)
N with basal ganglia perivascular spaces (%)	
1-10 (%)	316 (53.8)
11-20 (%)	149 (25.4)
>20(%)	122 (20.8)
N with centrum semi-ovale perivascular spaces (%)	
1-10 (%)	147 (25.0)
11-20 (%)	188 (32.0)
>20(%)	252 (42.9)
N with lacunes (%)	120 (20.4)
N with microbleeds (%)	92 (15.7)
1 microbleed (%)	42 (7.2)
2-4 microbleeds (%)	27 (4.6)
≥5 microbleeds (%)	27 (4.6)
Strictly deep microbleeds (%)	7 (1.2)
Strictly lobar microbleeds (%)	44 (7.5)
N with periventricular white matter hyperintensity (%)	
Grade 1 (%)	215 (36.6)
Grade 2 (%)	121 (20.6)
Grade 3 (%)	64 (10.9)
N with subcortical white matter hyperintensity (%)	
Grade 1 (%)	198 (33.7)
Grade 2 (%)	116 (19.8)
Grade 3 (%)	66 (11.2)
Mean Total Small Vessel Disease score	1.16 (1.17)

\*Middle cerebral artery pulsatility index was measured in a subset of 94/587 patients

**Table 2. Clinical predictors of common carotid and internal carotid artery pulsatility index**

	<b>Univariate B (95% CI)</b>	<b>p</b>	<b>Age and sex adjusted B (95% CI)</b>	<b>p</b>	<b>Multivariate adjusted B (95% CI)</b>	<b>p</b>
<b>Common carotid artery pulsatility index</b>						
Age	0.005 (0.004, 0.007)	<0.0001	0.006 (0.004, 0.007)	<0.0001	0.002 (0.000, 0.004)	0.016
Male sex	0.106 (0.061, 0.152)	<0.0001	0.107 (0.063, 0.151)	<0.0001	0.119 (0.074, 0.164)	<0.0001
History of hypertension	0.021 (-0.026, 0.068)	0.38	-0.012 (-0.058, 0.034)	0.60	-0.050 (-0.105, 0.005)	0.075
Hyperlipidaemia	0.032 (-0.015, 0.079)	0.18	0.014 (-0.031, 0.059)	0.54	-0.006 (-0.054, 0.041)	0.80
Diabetes	0.095 (0.030, 0.161)	0.004	0.075 (0.013, 0.138)	0.018	0.079 (0.016, 0.143)	0.014
Ever-smoking	0.006 (-0.041, 0.053)	0.80	-0.008 (-0.054, 0.038)	0.74	-0.022 (-0.067, 0.022)	0.33
Atrial fibrillation	0.097 (0.031, 0.164)	0.004	0.064 (-0.001, 0.128)	0.055	0.063 (0.000, 0.127)	0.051
Glomerular filtration rate	-0.002 (-0.003, -0.001)	0.0004	-0.001 (-0.002, 0.000)	0.16	0.000 (-0.002, 0.001)	0.49
Premorbid mean SBP (per SD increase)	0.049 (0.027, 0.070)	<0.0001	0.023 (0.001, 0.046)	0.045	-0.003 (-0.061, 0.054)	0.91
Premorbid mean DBP (per SD increase)	-0.025 (-0.046, -0.003)	0.028	-0.028 (-0.049, -0.007)	0.009	-0.019 (-0.056, 0.017)	0.30
Premorbid mean PP (per SD increase)	0.081 (0.060, 0.102)	<0.0001	0.063 (0.039, 0.087)	<0.0001	0.075 (0.026, 0.124)	0.003
<b>Internal carotid artery pulsatility index</b>						
Age	0.008 (0.007, 0.010)	<0.0001	0.008 (0.007, 0.010)	<0.0001	0.005 (0.004, 0.007)	<0.0001
Male sex	0.019 (-0.025, 0.062)	0.40	0.020 (-0.020, 0.060)	0.32	0.025 (-0.015, 0.065)	0.22
History of hypertension	0.103 (0.059, 0.146)	<0.0001	0.051 (0.009, 0.092)	0.017	0.012 (-0.037, 0.062)	0.62
Hyperlipidaemia	0.027 (-0.017, 0.072)	0.22	0.005 (-0.035, 0.046)	0.80	-0.032 (-0.074, 0.010)	0.14
Diabetes	0.110 (0.049, 0.171)	0.0005	0.091 (0.034, 0.147)	0.002	0.098 (0.042, 0.155)	0.001
Ever-smoking	-0.008 (-0.052, 0.036)	0.72	0.004 (-0.037, 0.045)	0.86	-0.014 (-0.054, 0.026)	0.49
Atrial fibrillation	0.081 (0.018, 0.144)	0.012	0.018 (-0.041, 0.077)	0.55	0.013 (-0.044, 0.069)	0.66
Glomerular filtration rate	-0.002 (-0.003, -0.001)	<0.0001	0.000 (-0.001, 0.001)	0.72	0.001 (0.000, 0.002)	0.11
Premorbid mean SBP (per SD increase)	0.081 (0.062, 0.101)	<0.0001	0.046 (0.026, 0.066)	<0.0001	0.020 (-0.031, 0.071)	0.44
Premorbid mean DBP (per SD increase)	-0.012 (-0.032, 0.009)	0.28	-0.011 (-0.030, 0.008)	0.26	-0.024 (-0.056, 0.009)	0.16
Premorbid mean PP (per SD increase)	0.110 (0.091, 0.128)	<0.0001	0.076 (0.055, 0.097)	<0.0001	0.063 (0.020, 0.107)	0.004

CI=confidence interval; SBP=systolic blood pressure; DBP=diastolic blood pressure; PP=pulse pressure

**Table 3. Relationships of common carotid and internal carotid artery pulsatility index with individual neuroimaging markers and global burden of small vessel disease, stratified by age**

	Age <70 (n=261)				Age ≥70 (n=326)			
	Univariate OR (95% CI)*	p	Age and sex adjusted OR (95% CI)*	p	Univariate OR (95% CI)*	p	Age and sex adjusted OR (95% CI)*	p
<b>Lacunae</b>								
CCA-PI	1.68 (0.70-4.04)	0.25	1.43 (0.57-3.55)	0.45	1.03 (0.47-2.28)	0.94	1.00 (0.45-2.25)	1.00
ICA-PI	6.17 (2.29-16.65)	0.0003	5.35 (1.95-14.70)	0.001	1.24 (0.49-3.16)	0.65	1.04 (0.40-2.73)	0.93
<b>Subcortical WMH burden</b>								
CCA-PI	1.55 (0.80-3.00)	0.20	1.23 (0.60-2.49)	0.58	1.30 (0.72-2.36)	0.39	1.19 (0.65-2.18)	0.58
ICA-PI	2.15 (0.99-4.65)	0.052	1.49 (0.66-3.35)	0.34	1.90 (0.96-3.78)	0.067	1.45 (0.72-2.95)	0.30
<b>Periventricular WMH burden</b>								
CCA-PI	1.80 (0.92-3.50)	0.084	1.40 (0.69-2.87)	0.35	1.46 (0.80-2.67)	0.22	1.30 (0.70-2.40)	0.40
ICA-PI	3.87 (1.78-8.42)	0.001	2.86 (1.26-6.46)	0.012	1.36 (0.68-2.71)	0.38	0.99 (0.49-2.02)	0.99
<b>Cerebral microbleed burden</b>								
CCA-PI	1.55 (0.46-5.20)	0.48	1.22 (0.35-4.25)	0.75	1.26 (0.54-2.95)	0.59	1.05 (0.44-2.49)	0.91
ICA-PI	1.50 (0.37-6.01)	0.57	1.14 (0.27-4.71)	0.86	1.99 (0.72-5.53)	0.19	1.57 (0.55-4.46)	0.40
<b>BG-PVS burden</b>								
CCA-PI	1.00 (0.45-2.23)	1.00	0.55 (0.23-1.32)	0.18	1.40 (0.76-2.58)	0.27	1.16 (0.62-2.16)	0.65
ICA-PI	4.25 (1.72-10.52)	0.002	2.83 (1.07-7.46)	0.035	1.57 (0.78-3.19)	0.21	0.99 (0.48-2.07)	0.98
<b>CS-PVS burden</b>								
CCA-PI	1.49 (0.77-2.88)	0.23	1.12 (0.56-2.23)	0.76	1.23 (0.65-2.32)	0.53	1.15 (0.60-2.19)	0.67
ICA-PI	2.71 (1.26-5.81)	0.01	1.96 (0.89-4.32)	0.095	1.67 (0.82-3.40)	0.16	1.48 (0.72-3.08)	0.29
<b>Total Small Vessel Disease Score</b>								
CCA-PI	1.50 (0.75-2.97)	0.25	1.08 (0.52-2.25)	0.84	1.48 (0.82-2.68)	0.19	1.20 (0.66-2.21)	0.55
ICA-PI	3.37 (1.54-7.38)	0.002	2.30 (1.01-5.25)	0.048	1.83 (0.93-3.63)	0.082	1.20 (0.59-2.42)	0.62

OR=odds ratio; CI=confidence interval; CCA=common carotid artery; ICA=internal carotid artery; PI=pulsatility index; WMH=white matter hyperintensity; BG=basal ganglia; CS=centrum semi-ovale; PVS=perivascular space

\*Odds of increasing small vessel disease burden in patients in top quartile of common carotid or internal carotid artery pulsatility index compared with patients in bottom quartile as referent

**Table 4. Relationships of common carotid and internal carotid artery pulsatility index with individual neuroimaging markers and global burden of small vessel disease, stratified by sex**

	Univariate OR (95% CI)*			
	Men	p	Women	p
<b>Lacunae</b>				
CCA-PI	1.58 (0.64-3.91)	0.32	1.51 (0.70-3.27)	0.30
ICA-PI	2.99 (1.17-7.62)	0.022	2.61 (1.12-6.08)	0.026
<b>Subcortical WMH burden</b>				
CCA-PI	1.78 (0.97-3.23)	0.061	2.62 (1.41-4.89)	0.002
ICA-PI	3.20 (1.74-5.91)	0.0002	3.97 (2.14-7.39)	<0.0001
<b>Periventricular WMH burden</b>				
CCA-PI	2.00 (1.10-3.65)	0.024	3.62 (1.92-6.83)	<0.0001
ICA-PI	4.44 (2.39-8.22)	<0.0001	5.30 (2.81-9.98)	<0.0001
<b>Cerebral microbleed burden</b>				
CCA-PI	2.22 (0.77-6.38)	0.14	1.59 (0.64-4.01)	0.33
ICA-PI	2.21 (0.94-5.21)	0.070	4.43 (1.38-14.28)	0.013
<b>BG-PVS burden</b>				
CCA-PI	1.42 (0.76-2.65)	0.27	3.12 (1.62-6.01)	0.001
ICA-PI	2.89 (1.48-5.65)	0.002	7.49 (3.61-15.53)	<0.0001
<b>CS-PVS burden</b>				
CCA-PI	1.41 (0.77-2.59)	0.27	2.82 (1.48-5.37)	0.002
ICA-PI	3.28 (1.78-6.04)	0.0001	4.03 (2.14-7.58)	<0.0001
<b>Total Small Vessel Disease Score</b>				
CCA-PI	1.84 (1.01-3.35)	0.046	3.12 (1.67-5.82)	0.0004
ICA-PI	3.72 (2.01-6.88)	<0.0001	6.06 (3.20-11.47)	<0.0001

OR=odds ratio; CI=confidence interval; CCA=common carotid artery; ICA=internal carotid artery; PI=pulsatility index; WMH=white matter hyperintensity; BG=basal ganglia; CS=centrum semi-ovale; PVS=perivascular space

\*Odds of increasing small vessel disease burden in patients in top quartile of common carotid or internal carotid artery pulsatility index compared with patients in bottom quartile as referent