

## **Advances in Hypoxia-Inducible Factor Biology**

**Hani Choudhry<sup>1</sup>, Adrian L. Harris<sup>2</sup>**

<sup>1</sup>Department of Biochemistry, Cancer Metabolism and Epigenetic Unit, Faculty of Science, Cancer and Mutagenesis Unit, King Fahd Center for Medical Research, King Abdulaziz University, Jeddah, Saudi Arabia. <sup>2</sup>Molecular Oncology Laboratories, Department of Oncology, University of Oxford, Weatherall Institute of Molecular Medicine, Oxford, OX3 9DS, UK

### **Corresponding Author & Lead Contact:**

Prof. Adrian L. Harris –Molecular Oncology Laboratories,  
Department of Oncology, University of Oxford, Weatherall  
Institute of Molecular Medicine, Oxford, OX3 9DS, UK:  
Tel: +44 (0)1865 222457;  
email: [aharris.lab@imm.ox.ac.uk](mailto:aharris.lab@imm.ox.ac.uk).

## **Summary**

Hypoxia inducible factor (HIF), a central regulator for detecting and adapting to cellular oxygen levels, transcriptionally activates genes modulating oxygen homeostasis and metabolic activation. Beyond this, HIF influences many other processes in the complex response to hypoxic stress. Hypoxia, in part through HIF-dependent mechanisms, influences epigenetic factors, including DNA methylation and histone acetylation, which modulate hypoxia-responsive gene expression in cells. Hypoxia profoundly affects expression of many noncoding RNAs classes that have clinicopathological implications in cancer. HIF can regulate noncoding RNAs production while, conversely, noncoding RNAs can modulate HIF expression. There is recent evidence for crosstalk between circadian rhythms and hypoxia-induced signalling, suggesting involvement of molecular clocks in adaptation to fluxes in nutrient and oxygen sensing. HIF induces increased production of cellular vesicles facilitating intercellular communication at a distance, for example promoting angiogenesis in hypoxic tumors. Overall, understanding the complex networks underlying cellular and genomic regulation in response to hypoxia via HIF may identify novel and specific therapeutic targets for many diseases.

**Keywords:** Hypoxia, HIF, exosomes, non-coding RNAs, epigenetics, circadian rhythms, molecular clocks, cancer

## Introduction

Oxygen is essential for mammalian metabolism and physiological functions because of its use in cellular energy production and cofactor/substrate for many enzymes. The central regulator of oxygen detection and adaptation at the cellular level is hypoxia inducible factor (HIF), a heterodimeric transcription factor that consists of either HIF-1 $\alpha$  or HIF-2 $\alpha$  and HIF-1 $\beta$ /ARNT subunits. In the presence of oxygen (normoxia), HIF $\alpha$  interacts and binds to the von Hippel–Lindau (VHL) protein, which consequently activates the ubiquitin ligase system, leading to proteasomal degradation of HIF $\alpha$ . Hydroxylation of proline residues in HIF $\alpha$  is vital for VHL binding and depends on  $\alpha$ -ketoglutarate-dependent dioxygenases, prolyl hydroxylases (PHD), and the asparaginyl hydroxylase, factor-inhibiting HIF (FIH). During hypoxia, PHDs are inactive, leading to HIF- $\alpha$  stabilization and dimerization with HIF-1 $\beta$ . Upon dimerization, HIF translocates to the nucleus to bind to E-box-like hypoxia response elements (HREs) within the promoter region that contain the sequence 5'-[A/G]CGTG-3' (Kaelin and Ratcliffe, 2008). HIF activates genes that control cellular oxygen homeostasis, including genes involved in oxygen consumption, erythrocyte production, angiogenesis, and mitochondrial metabolism. Hypoxic cells counter stress by transcriptional and post-transcriptional mechanisms, mainly regulated by HIF. These molecular changes allow cells to adapt to stress by lowering oxygen consumption through a shift to glycolysis rather than oxidative metabolism and by reducing energy demand for cellular processes such as cell division (Ratcliffe, 2013). Most solid tumors have some degree of hypoxia and is associated with poor clinical outcome. Induction of HIF activity upregulates genes involved in many hallmarks of cancer, including metabolic reprogramming, cell proliferation, invasion and metastasis, apoptosis, and resistance to therapies (Rankin and Giaccia, 2016).

In contrast to these, other well-studied pathways, in this review, we discuss recent advances in the biology of hypoxia and HIF regulating epigenetics, non-coding RNAs, biological clocks and cellular vesicles. These findings provide new insights on the complex networks underlying cellular and genomic regulation in response to hypoxia and may provide novel targets for future therapies.

## Hypoxia, HIFs and epigenetics

DNA and histone methylation are two key mechanisms of epigenetic regulation of gene expression. The histone methylation is modulated via histone methyltransferases and histone demethylases. Histone acetylation is a common marker of gene activation, whereas histone deacetylation is a marker of repression. Methylation status of specific amino acids in histones lead to gene activation or repression. For example, histone 3 lysine 4 methylations (H3K4me, me2 and me3) are markers of gene activation, while histone 3 lysine 9 methylations (H3K9me2 and me3) and H3K27me3 are markers of transcription suppression (Dawson, 2017).

Several histone lysine demethylases (KDM) are induced under hypoxia upon HIF-1 $\alpha$  stabilization

**Table 1.** All Jumonji-type KDMs are members of the 2-oxoglutarate-dependent dioxygenase family (2-OGDO), thus dependent on oxygen and a Krebs cycle intermediate “2-oxoglutarate” to catalyse their enzymatic reactions. Most of KDM enzymes are structurally similar to the HIF hydroxylase Factor Inhibiting HIF-1 (FIH), suggesting that KDM enzymes may act as molecular oxygen sensors in the cell. KDMs can either enhance or suppress gene expression. For instance, KDMs can demethylate histone 3 di- or trimethylated lysine 4 (H3K4me2 and 3) and H3K36me2 and 3 (activation markers) or H3K9me1 and 2 and H3K27me2 and 3 (repression markers). Several Jumonji domain containing histone lysine demethylases (KDM2-7) present different protein binding domains, allowing their interaction with other chromatin associated proteins (Estaras et al., 2013).

Pollard *et al.* and Beyer *et al.* (2008) reported KDM3A and KDM4B induction under hypoxia in HIF-1 $\alpha$  dependent manner (Beyer et al., 2008; Pollard et al., 2008). Krieg *et al.* (2010) identified KDM3A dependent genes, many of which are hypoxia inducible genes such as adrenomedullin (ADM), heme oxygenase 1 (HMOX1), and SERPINE1. KDM3A loss significantly reduces expression of these genes and tumor growth *in vivo* models (Krieg et al., 2010). Therefore, hypoxic regulation of KDM3A acts as a signal amplifier to facilitate hypoxic gene expression, ultimately enhancing tumor growth.

KDM4B is also induced by hypoxia in a HIF-1 $\alpha$ -dependent manner in colorectal cancer cells, which decreases H3K9me3 level at the promoter of hypoxia-inducible genes (Salminen et al., 2016). KDM4B

is involved in cell proliferation, apoptosis, and cell cycle arrest. Hence, the expression of KDM4B in CRC correlates positively with the hypoxia marker, carbonic anhydrase 9 (CA9) expression (Salminen et al., 2016).

Genome-wide analysis of HIF-1 $\alpha$  binding sites reported HRE sites on the promoter region of two other histone demethylases, KDM4C (JMJD2C) and KDM5B (JARID1B). Both of KDMs found significantly upregulated under hypoxia (Xia et al., 2009). KDM5B catalyzes the removal of methyl groups from tri-, di-, and monomethylated lysine 4 of histone H3 (H3K4me<sub>3/2/1</sub>), while KDM4C converts specific trimethylated histone residues to dimethylated residues. At protein level, KDM4C selectively interacts with HIF-1 $\alpha$  and that enhances HIF-1 $\alpha$  binding to HRE to activate transcription of several genes which encode proteins for metabolic reprogramming and metastasis such as BNIP3, LDHA, LOX, PDK1, and SLC2A1 (Luo et al., 2012). KDM4C is overexpressed in several malignancies, including prostate, bladder, and lung cancers. In contrast, KDM4C is reported downregulated in melanoma and triple receptor negative breast cancer, where it inhibits cell proliferation (Klein et al., 2014).

Protein expression of KDM5B and KDM3A (JMJD1A) are significantly induced in hypoxic cells and in renal cancer cells that have lost VHL expression. However, the demethylase activity of both enzymes is reduced but remains active under low oxygen stress (Beyer et al., 2008), suggesting that HIF mediates upregulation of their expression which might provide a secondary mechanism to retain a dynamic regulation of H3K9 methylation in hypoxia. Recently, KDM3A was reported to enhance glycolysis through interaction with HIF-1 $\alpha$  in cancer cells. KDM3A facilitate HIF1 $\alpha$ -mediated Phosphoglycerate Kinase 1 (PGK1) transcription by reducing the levels of H3K9me<sub>2</sub> at the HRE site on the PGK1 promoter in hypoxia (Wan et al., 2017). Mutated JMJD1A (H1120Y) failed to interact with HIF-1 $\alpha$  and was unable to reduce the level of H3K9me<sub>2</sub> at the HRE of PGK1 promoter, thus suggesting that the demethylase activity of KDM3A is essential for cooperation with HIF-1 $\alpha$  and for the modulation of levels of H3K9me<sub>2</sub> to enhance glycolysis. Generally, hypoxia globally induces levels of H3K4me<sub>2</sub>, H3K4me<sub>3</sub>, H3K79me<sub>3</sub>, H3K9me<sub>2</sub>, H3K9me<sub>3</sub> and H3K36me<sub>3</sub> in different cell line models. These changes indicate hypoxia inhibits or significant reduces the enzymatic activity of different histone demethylase such as KDM5A, KMD6B (Johnson et al., 2008; Prickaerts et al., 2016; Tausendschon et

al., 2011; Zhou et al., 2010). There are limited data on biochemical and kinetics of KDMs activity with respect to oxygen availability (Cascella and Mirica, 2012), however, it is clear that enzymatic activity of KDMs is altered under hypoxia with potential consequences on gene regulation.

Hypoxia also influences the remodeling of the nucleosome architecture for transcriptional regulation. Johnson *et al.* (2008) found an increase in total histone H3 occupancy at the promoters of hypoxia-suppressed genes such as *ALB* and *AFP* and decreased occupancy at hypoxia-upregulated genes such as *EGR1* and vascular endothelial growth factor (*VEGF*), suggesting an inverse correlation between nucleosome occupancy and transcriptional activation in response to hypoxia. Hypoxia-induced *VEGF* and *EGR1* promoters were marked with increased H3K9 acetylation and a significant decrease in H3K9me2 and H3K27me2 methylation (Johnson et al., 2008). In contrast, hypoxia-repressed *AFP* and *ALB* promoters showed low acetylation of H3K9 and high H3K9 and H3K27 methylation.

Several chromatin-modifying complexes control hypoxia-inducible gene expression. The SWI/SNF chromatin remodeling complex is associated with HIF-1 $\alpha$  to regulate hypoxia inducible genes (Kenneth et al., 2009). A nucleosome remodeling deacetylase complex, Metastasis-associated 1 (MTA1) is significantly induced in hypoxic breast cancer cells. MTA1 physically associates with HIF-1 $\alpha$  to increase HIF-1 $\alpha$  transcriptional activity and VEGF expression (Yoo et al., 2006). The imitation switch (ISWI) is another chromatin remodeling complex linked to the hypoxic response. Unlike the SWI/SNF complex, ISWI downregulation promotes HIF-1 $\alpha$  activity without affecting its levels. Moreover, ISWI depletion alters a subset of HIF-1 $\alpha$  target genes and, subsequently, reduces autophagy and increases cell death during hypoxia. The ISWI is vital for the regulation of gene and protein expression level of FIH by altering RNA polymerase II loading onto the FIH promoter (Melvin and Rocha, 2012). These findings suggest a dynamic remodeling of the chromatin structure under hypoxia, which influences the downstream regulation of transcription.

Pan-genomic analysis of DNaseI hypersensitivity sites and nucleosome occupancy revealed that, under hypoxia, a large number of HIF binding sites overlap with open chromatin sites defined in normoxic cells (Schodel et al., 2011; Tanimoto et al., 2010). Thus, HIF may bind at pre-existing open chromatin

sites as described for glucocorticoid receptor binding at the genome wide level (John et al., 2011). The open chromatin conformation is associated with HIF binding at the enhancer site in cells with VHL loss, but not in cells with intact VHL (Platt et al., 2016), suggesting cell type specificity in the formation of enhancer for HIF transcriptional response.

Several studies shown that hypoxia modulate chromatin remodelling and histone modifications to facilitate transcription of HIF dependent genes. Integrated genomic analysis using massively parallel sequencing revealed that, under hypoxia, RNAPol2 is already bound at the promoter of most hypoxia-inducible genes in normoxic cells. However, during hypoxia, there was a slight increase in loading of RNAPol2 at the TSS with significant increase in release of pre-bound promoter-paused RNAPol2 across the body of the gene. Similarly, little change in H3K4me3 signal was observed at the promoter under hypoxia, but the signal increased downstream of the TSS (Choudhry et al., 2014) . These findings support the hypothesis that RNAPol2 recruits histone methyl transferases that trimethylate H3K4 to facilitate transcriptional regulation during hypoxia.

Large numbers of HIF-binding sites are located at long chromosomal distances from hypoxia inducible transcripts. Therefore, it is challenging to define the specific targets of HIF binding sites and to determine how HIF affects the chromatin conformation over the distance to interact with functional elements. Recently, Platt *et al.* (2016) employed a high-resolution chromosome conformation method to defined specific high attitude interactions of HIF-binding regions with single or multiple promoters of hypoxia-inducible genes. They reported the presence of multiple chromosomal loops, which generate interaction between promoter-enhancer, and enhancer-enhancer during hypoxia (Platt et al., 2016). They found that promoter distant HIF-binding sites establish cis-interactions with HIF inducible genes to maintain the transcriptional repertoire, supporting the notion that hypoxia causes dynamic changes in the chromosome structure for HIF to bind at primed enhancer–promoter complexes, allowing rapid activation of its transcriptional cascade through the release of promoter paused RNA polymerase II.

Hypoxia-induced histone demethylases regulate not only gene expression, but also modulate chromatin structures such as heterochromatin and polycomb complexes. For instance, hypoxia modulates chromatin conformation to induce expression of SLC2A3 (GLUT3), through interaction between

KDM3A and HIF-1 $\alpha$  in endothelial cells (Mimura et al., 2012). KDM3A is recruited at the promoter of SLC2A3 gene in a HIF-1 $\alpha$ -dependent manner, inducing SLC2A3 expression and subsequently promoting glycolysis.

Reduction of heterochromatin because of increase expression of specific histone demethylases (KDM3, KDM4, and KDM6) is commonly associated with oncogenesis (Slee et al., 2012; Young and Hendzel, 2013). In contrast, KDM2 and KDM5 overexpression promotes the maintenance of heterochromatin and genome stability (Li et al., 2014). Hypoxia upregulates euchromatic histone-lysine N-methyltransferase 2 (G9a) to increase the H3K9me2 level and, consequently, controls the expression of several hypoxia-regulated genes, including breast cancer 1, early onset (*BRCA1*), colon cancer, nonpolyposis type 2 (*Mlh1*) and mutL homolog 1 (Lu et al., 2011). Hypoxia also modulates the deSUMOylation of transcriptional repressor CTCF (CCCTC binding factor) (Wang et al., 2012).

Histone deacetylases (HDACs) are other epigenetic modifications, which are altered in hypoxic conditions. Hypoxia regulate the expression and activity of several HDACs. Histone Deacetylase 1 (HDAC1) overexpression enhances angiogenesis of human endothelial cells by downregulating hypoxia-responsive tumor suppressor genes, including *p53* and *VHL* (Kim et al., 2001). Several HDACs (Class I and Class IIa) found to promote HIF-1 $\alpha$  stability and accumulation by regulating HIF-1 $\alpha$ -PHD2 interaction. HDAC4, HDAC5, HDAC6 enhance HIF-1 $\alpha$  stability and activity through modulation of acetylation level and activity of key HIF-1 $\alpha$  cofactors such as HSP90 and p300 (Chen et al., 2015; Geng et al., 2011; Kong et al., 2006; Schoepflin et al., 2016). Inhibition of either HDAC4 or HDAC5 reduces HIF-1 $\alpha$  protein levels and suppresses HIF-1 $\alpha$  activity (Chen et al., 2015). Therefore, HDACs may be involved in the regulation of posttranslational processing of HIF-1 $\alpha$  by modulating the Hsp70/Hsp90. On the other hand, HDAC inhibition (Vorinostat) was reported to increase HIF-2 $\alpha$  accumulation, which suppresses sarcomas tumor growth in a HIF-2 $\alpha$  dependent manner (Nakazawa et al., 2016). These findings provide a rationale for targeting HIF with HDAC inhibitors. Hypoxia upregulates HDAC3 to regulate the expression of EMT markers in cancer cells (Wu et al., 2011). HDAC3 deacetylates the histone 3 lysine 4 (H3K4Ac) in the promoter regions of EMT marker genes and elevates the levels of H3K4me2 and H3K4me3. Hypoxia-induced EMT is abolished upon HDAC3 suppression.

Under hypoxia, HDAC3 interacts with the WD repeat domain 5 (WDR5), which increases the activity of histone methyltransferases (HMT) (Wu et al., 2011). HDAC3 enhances HMT activity by increasing the stability of the HMT complex or through conformational remodeling. However, the molecular mechanisms remain to be explored.

Histone modifications in collaboration with DNA methylation play a vital role in controlling gene expression and modulating chromatin structure. The HIF-1 $\alpha$  promoter contains an HRE, which harbors a CpG island. In colon cancer, this site is aberrantly demethylated, allowing binding of HIF-1 $\alpha$  to its own promoter (Koslowski et al., 2011). This leads to auto-transactivation of HIF-1 $\alpha$  expression and transactivation of HIF-1 $\alpha$  target genes. Treating tumor cells with DNMT inhibitor, 5-aza-deoxycytidine (5-aza-dC) elevates the expression of hypoxia-induced genes. DNA methylation can also affect HIF-1 $\alpha$  stability. For example, *VHL* hypermethylation results in HIF-1 $\alpha$  constitutive activation. Epigenetic silencing of *VHL* has been associated with increased nuclear translocation of HIF-1 $\alpha$  and upregulation of HIF-1 $\alpha$  target genes such as CA9 and GLUT1 (Schmitt et al., 2009). *VHL* hypermethylation has been observed in many solid epithelial tumors and cancer cell lines (Sanchez-Vega et al., 2013; Stewart et al., 2016). In gastric cancer, CA9 expression is associated with tumor progression and metastasis. CA9 overexpression has been linked with promoter DNA hypomethylation (Nakamura et al., 2011). In addition, studies reported *PHD3* silencing because of DNA methylation in multiple myeloma, prostate cancer, B cell lymphoma, and breast cancer cell lines (Hatzimichael et al., 2010; Place et al., 2011). However, epigenetic silencing of *PHD3* showed no significant effect on HIF-1 $\alpha$  or HIF-2 $\alpha$  protein levels. The above studies reported that PHD3 and VHL, two vital enzymes involved in HIF-1 $\alpha$  protein destabilization and regulators of the HIF pathway, are commonly hypermethylated under hypoxia.

HRE methylation status can have a profound impact on HIF transactivation of its target genes. HRE hypermethylation blocks the HIF binding at erythropoietin (EPO) enhancer and represses its expression (Wenger et al., 1998). Mammalian stanniocalcin-2 (STC2) is a secreted glycoprotein, which is overexpressed in cancer and involved in EMT, drug resistance, and cell proliferation (Chen et al., 2016). HRE hypermethylation at *STC2* promoter has been reported in different cancer cell lines. Treating cancer cells with 5-aza-2'-deoxycytidine (5-aza-CdR) under hypoxia significantly increased STC2

expression (Law et al., 2008). Another example is BCL2/Adenovirus E1B 19kDa Interacting Protein 3 (BNIP3) is induced by hypoxia in a HIF-dependent manner. BNIP3 is a mitochondrial protein that functions as a pro-apoptotic factor (Feng et al., 2016; Vasagiri and Kutala, 2014). In contrast to many other cancers, some colorectal and pancreatic cancer cell lines showed little or no hypoxic induction of BNIP3 despite an intact HIF signaling pathway (Bacon et al., 2007; Okami et al., 2004). Analysis revealed that *BNIP3* promoter is located within a CpG island and is hypermethylated under hypoxia, thus blocking *BNIP3* induction. Hypermethylation silencing of BNIP3 is detected in 66% of primary colorectal and 49% of primary gastric cancers and associated with worse clinical outcome (Murai et al., 2005).

The ten-eleven translocation (TET) family proteins promote key epigenetic changes and regulate crucial cancer development processes. In mammals, TET enzymes catalyze DNA demethylation by hydroxylating 5-methylcytosine (5mC) to 5-hydroxymethylcytosine (5hmC). 5hmC is further oxidized to 5-formylcytosine (5fC) and 5-carboxylcytosine (5caC) and subsequently substituted with an unmodified cytosine by base-excision repair (BER) to complete demethylation (Ito et al., 2011; Tahiliani et al., 2009). Similar to PHDs, TET enzymes are dependent on Fe<sup>2+</sup> and are  $\alpha$ -ketoglutarate dioxygenases. Decrease in 5mC oxidation due to reduce TET activity causes an increase in DNA methylation levels. Shahrzad *et al.* reported global DNA hypomethylation in hypoxic colorectal and melanoma cancers measured by the amount of 5mC. They found an inverse correlation between the magnitude of tumor hypoxia and the incidence of methylation both in cells and in xenograft models (Shahrzad et al., 2007). In addition, prolong hypoxic stress causes hypomethylation of retrotransposable Alu or short interspersed nuclear elements (SINEs) and reverses transcriptase coding long interspersed nuclear element (LINE) transcripts in glial tumor and osteosarcoma (Pal et al., 2010). These repetitive elements play a vital role increasing genomic instability during hypoxia. In hepatocellular carcinoma, hypoxia induces global DNA demethylation by decreasing S-adenosylmethionine (SAM) levels both *in vitro* and *in vivo*. Methionine adenosyltransferase 2A (MAT2A) catalyzes the production of SAM. Hypoxia induces MAT2A expression in a HIF-1 $\alpha$

dependent manner and recruits p300 and HDAC1 (Liu et al., 2011b). This suggests that hypoxia promotes DNA demethylation via HIF-1 $\alpha$  activation and transcriptional upregulation of MAT2A.

In contrary to these observations, other studies reported that hypoxia causes DNA hypermethylation. Watson *et al.* (2009) found an increase in global DNA methylation and H3K9 histone acetylation in hypoxia-adapted PwR-1E benign prostate epithelial cells. They found upregulation of DNA methyltransferase DNMT3b and gene-specific changes in DNA methylation (Watson et al., 2009). In another study, prolonged hypoxia exposure of human cardiac fibroblast cells caused a pro-fibrotic phenotype. The hypoxia-induced pro-fibrotic phenotype was associated with global DNA hypermethylation and increased levels of DNMT1 and DNMT3B (Watson et al., 2014).

Recently, Thienpont and colleagues (2016) reported that hypoxia promotes hypermethylation through 5hmC loss in a number of human cancer and murine cell lines. Hypoxia-induced hypermethylation is due to reduced activity of TETs, but not their expression. The hypoxic reduction of TET activity was independent of TET expression, metabolism, HIF, and reactive oxygen species. Using DNA-immunoprecipitation sequencing (DIP-seq), they found that global loss of 5hmC in hypoxic conditions and primarily at promoter region of genes associated with gain of 5mC. The Cancer Genome Atlas (TCGA) data revealed an inverse association between TET2 and TET3 expression and DNA hypermethylation, while a positive association with TET1 and TET3 mutations and hypermethylation (Thienpont et al., 2016), indicating that reduced TET activity limits 5hmC generation and promotes DNA hypermethylation. Interestingly, hypoxia-induced DNA hypermethylation in breast tumors occurred at the promoter region of tumor suppressor genes, but oncogenes were not affected (Thienpont et al., 2016). These findings reveal that hypoxia reduces TET activity, leading to downregulation of tumor suppressor genes, and establishes a link between hypoxia and DNA methylation. In contrast, other studies reported a positive regulation and activity of the TET proteins and induction of global DNA hypomethylation during hypoxia. Wu et al. (2015) reported that HIF-1 $\alpha$ -dependent regulation of TET1 and TET3 promotes global DNA hydroxymethylation during hypoxia. Hypoxia-induced overexpression of TET1/TET3 proteins activates the TNF $\alpha$ - p38-MAPK pathway to enhance breast cancer stemness. Expression of 5hmC, TET1, and TET3 was significantly correlated with tumor

hypoxia, tumor progression, and poor clinical outcome in breast cancer (Wu et al., 2015b). Moreover, HIF-independent TET1 overexpression, leading to global DNA hypomethylation, was also reported in scleroderma fibroblasts (Hattori et al., 2015). Tsai et al. (2014) demonstrated that hypoxia regulates the expression of TET1 that interacts with HIF-1 $\alpha$  and HIF-2 $\alpha$  to enhance their transactivation activity and promote hypoxia-induced EMT. Furthermore, several groups reported that TET proteins are linked with glucose and lipids metabolism (Chen et al., 2013; Tsai et al., 2014). Accumulation of fumarate and succinate metabolites can increase HIF-1 $\alpha$  levels and expression of HIF target genes in part by inhibiting the PHD enzymes in cancer cells (Isaacs et al., 2005; Selak et al., 2005). Fumarate and succinate also inhibit TET1 and TET2 enzymes, resulting in DNA hypermethylation and downregulation of HIF dependent genes (Laukka et al., 2016). There are several contradictory data in the literature on the regulation of TETs and their impact on DNA methylation in hypoxic conditions. Further investigations are needed to understand the mechanism underlying TET protein function and its up/downstream regulators in hypoxia.

Many studies demonstrated that hypoxia provokes epigenomic alterations in the chromatin and DNA methylation landscape, which consequently modulates the transcriptional output of tissues. However, hypoxic epigenetic change is a complex mechanism, which is variable in response because of various factors, including level of oxygen, exposure time to hypoxia, type of hypoxic treatments, tissue and cell line specific responses, and hypoxia dependent and independent protein synthesis. These factors could explain inconclusive results from many global histone and DNA methylation studies on hypoxia. For example, KDM3A and KDM4B activity does not change at 1% O<sub>2</sub> tension, whereas the reduction of O<sub>2</sub> to 0.2% significantly abrogates histone demethylase activity (Beyer et al., 2008).

Overall, it is clear that different epigenetic factors including transcriptional co-regulators, DNA methylation, histone and chromatin modifications mediate regulation of hypoxia-responsive gene expression through HIF dependent and non-dependent mechanisms in cancer. Further epigenetic mechanisms to control hypoxia response will be the major focus of upcoming research endeavors.

### **Hypoxia, HIFs and non-coding RNAs**

Recent large-scale genomic sequencing projects revealed that less than 2% of human transcriptional output encodes for proteins, while the remaining genome encrypts different classes of non-coding RNAs. Emerging evidence has been revealed that hypoxia regulates expression of different non-coding RNAs classes and they in turn influence on HIF expression and stability. These hypoxia/HIF associated non-coding RNAs modulate range physiological and pathological pathways including cellular growth, metabolism and angiogenesis.

#### *HIF and microRNA*

MicroRNAs (miRNAs) are a class of small RNAs (~22 nucleotide duplexes) that regulate RNA stability and mRNA translation. The extensive literature on the regulation of individual and genome-wide miRNA expression, biogenesis miRNA pathway, and the regulation of miRNA target genes in hypoxia has been recently reviewed (Choudhry et al., 2016). Hypoxia has a profound impact on miRNAs expression, which is controlled by a complex network of gene regulation. For instance, hypoxia directly regulates miRNA expression through HIF or indirectly by other hypoxia-regulated factors such as Oct-4. miRNA also regulates expression or stabilization of HIF-1 $\alpha$  and/or HIF-2 $\alpha$  by direct binding to their mRNAs or through controlling the expression of a regulatory unit of the HIFs such as VHL or a PHD. In addition, miRNA biogenesis machinery is tightly regulated by hypoxia.

Hypoxic miRNA expression signatures have been develop from primary tumors and cancer cell lines, including brain, breast, colorectal, and bladder cancers (Agrawal et al., 2014; Blick et al., 2015; Choudhry et al., 2016). Most miRNAs identified in the hypoxia signature such as miR -23, miR-24, miR-26, miR-27, miR-103, miR-107, miR-181, miR-210, and miR-213 are commonly overexpressed in different tumor types.

Several differentially expressed miRNAs were identified in human endothelial cells grown under hypoxia (1% O<sub>2</sub> for 24 hours), including miR-210, which is overexpressed in most hypoxic tumors, and miR-150, which is downregulated in hypoxic tumor studies (Voellenkle et al., 2012). Similarly, RNA-seq analysis of the breast cancer cell line (MCF-7) grown under hypoxia indicated the upregulation of 41 miRNAs and downregulation of 28 miRNA by hypoxia (Camps et al., 2014). Analysis of hypoxic

responses at different time points (16, 32, and 48 hours) indicated an increase in the number of differentially expressed miRNAs with time. Several hypoxia-regulated miRNAs are located within intronic regions of protein coding genes. However, there was no significant association between the miRNA and host gene expression during hypoxia (Camps et al., 2014). Some of the upregulated hypoxia-induced miRNAs (miR-210-3p, miR-27a-3p, and miR-24-3p) were associated with a hypoxic signature in breast cancer.

Various miRNAs upregulated by hypoxia are direct targets of HIF-1 $\alpha$  and/or HIF-2 $\alpha$ . HIF binds to their promoter region to induce their expression and target both upstream and downstream signaling molecules that function as oncogenes and/or tumor suppressors. Changes in hypoxia-regulated miRNAs have been associated with clinico-pathological features and clinical outcome in various cancer types (Favaro et al., 2011; Rupaimoole et al., 2016a). miR-34 expression is decrease in hypoxic conditions to modulate the tumor microenvironment and EMT by targeting *IL6*, *NOTCH1*, and *JAG1* (Rokavec et al., 2014). Another study reported that miR-34 targets *CD44*, hence influencing cancer stem cell signaling (Liu et al., 2011a). miR-15 and miR-16 function as tumor suppressor by downregulating anti-apoptotic oncogenes. miR-155 is upregulated in hypoxia and has an HRE in its promoter in colorectal cancer cell lines (Bruning et al., 2011). Hypoxic downregulation of miRNAs, Let-7a, miR-135a, miR-146a, and miR-30, causes the upregulation of pro-metastatic genes, including *RHOB1*, *TAGLN*, *SRTAD1*, *TXNIP*, *JAG1*, *CTGF*, and *JUN*.

miR-210 is a well-established target of HIF-1 $\alpha$  and is strongly induced in most cancer types in response to hypoxia. miR-210 reduces the level of glycerol-3-phosphate dehydrogenase 1-like (GPD1L) to stabilize HIF-1 $\alpha$  protein by downregulating HIF-1 $\alpha$  hyperhydroxylation, suggesting a positive feedback loop of HIF-1 $\alpha$  dependent regulation via miR-210 (Kelly et al., 2011). In addition, miR-210 feedback loop functions through targeting HIF-1 $\alpha$  inhibitor such as Succinate Dehydrogenase Complex Subunit D (SDHD). Thus, miR-210 downregulates SDHD functions to promote HIF-1 $\alpha$  stabilization (Gorospe et al., 2011).

Many miR-210 targets regulate several key cellular processes, including cell cycle, cell morphology, polarization, differentiation, apoptosis, metabolism, tumor migration, and metastasis (Qin et al., 2014). The mitochondrial iron sulfur scaffold protein (ISCU) is among the critical targets of miR-210 (Chan et al., 2009). ISCU is a vital component of iron sulfur clusters, which are cofactors for many enzymes, including those that modulate metabolism, iron homeostasis, and oxidative stress. In addition, ISCU is a key part of TCA cycle components, including succinate dehydrogenase and mitochondrial complex components. miR-210 reduces the ISCU expression, which results in decreased tricarboxylic acid “TCA” cycle and mitochondria electron transport chain function (Favaro et al., 2010). In breast cancer, miR-210 and ISCU levels were inversely correlated and low ISCU expression was associated with poor recurrence-free survival. miR-210 enhances angiogenesis and metastasis by promoting glucose transporters such as GLUT-1. Induction of GLUT-1 generates an extracellular microenvironment that supports angiogenesis through upregulation of VEGF and Platelet-Derived Growth Factor (PDGF)(Bailey et al., 2012). In addition, miR-210 can regulate VEGF cellular expression by regulating levels of *EFNA3* and *PTP1B*, which are adverse regulators of VEGF. Downregulation of *EFNA3* or *PTP1B* by miR-210 causes increased VEGF production and leads to angiogenesis (Fasanaro et al., 2008; Kim et al., 2013). miR-210 is also involved in controlling apoptosis by negatively regulating the expression of key molecules of apoptosis-related cell signaling pathways, including apoptosis-Inducing Factor Mitochondrion-associated 3 (*AIFM3*) (Yang et al., 2012), *CASP8AP2* (Kim et al., 2009), and *SIN3A*. miR-210 promotes metastasis by downregulating the vacuole membrane protein 1 (*VMP1*), which inhibits cell migration and invasion. Increased level of miR-210 results in a decrease in *VMP1* gene and protein expression in various cancer types, thus promoting cell migration and invasion (Ying et al., 2011).

In addition to HIF, several other factors play a role in modulating miRNA expression in hypoxia. For example, hypoxia activates the AKT signaling, resulting in an increase of miR-21 expression in an NF- $\kappa$ B- and CREB- dependent manner (Polytarchou et al., 2011). HIFs control the expression of several transcription factors. The promoter region of miR-210 contains several conserved transcription factor sites, including Oct-4, which is regulated by hypoxia (Ivan et al., 2008). HIFs also control the expression

of the transcription factor, TWIST1, which regulates oncogenic miR-10b that mediates metastasis (Haque et al., 2011). In endothelial cells, C/EBP- $\alpha$ /RUNX-1 transcription factor induces the expression of miR-424 during hypoxia to modulate vascular remodelling (Ghosh et al., 2010). These findings indicate the role of HIF-independent regulation of miRNA expression under hypoxia.

Studies reported that many miRNAs modulate the expression of HIFs and/or regulatory units of the HIF pathway. These miRNAs are commonly downregulated in hypoxia, resulting in increased *HIF* gene and protein expression. miR-18a is suppressed by hypoxia in different cancer types and directly targets the 3'-UTR of HIF-1 $\alpha$  to inhibit metastasis (Wu et al., 2015a). Moreover, hypoxia reduces the expression of miR-199a, which targets 3'-UTR of HIF-1 $\alpha$ , and HIF-2 $\alpha$ . Changes in miR-199a affect HIF levels and the expression of genes that control cell migration and metastasis, including the matrix-remodeling enzyme, lysyl oxidase (LOX) (Joshi et al., 2014). miR-138 directly targeted *SOX4* and *HIF-1 $\alpha$*  in ovarian cancer, modulating tumor migration and invasion phenotypes (Yeh et al., 2013). miR338-3p also binds the 3'-UTR of HIF-1 $\alpha$ , resulting in decrease expression of HIF-1 $\alpha$  and its regulated genes such as *VEGF* and *GLUT-1* (Xu et al., 2014).

Recent studies uncovered a novel oncogenic role of hypoxia in regulating the miRNA biogenesis pathway (Bandara et al., 2017). Hypoxia activates EGFR signaling to enhance cell growth and tumorigenesis. This activation increases the phosphorylated form of AGO2 at Tyr 393, hence obstructing the interaction between DICER1 and AGO2 and blocking miRNA accumulation and maturation. Hypoxia-induced EGFR dependent AGO2-Tyr 393 phosphorylation is vital for cellular adaptation under hypoxia and is associated with poor clinical survival outcome of patients with breast cancer (Shen et al., 2013). Global miRNA downregulation in the hypoxic tumor microenvironment results from alteration and suppression of key members of the miRNA biogenesis pathway. Hypoxia downregulates Dicer and Drosha in an ETS1/ELK1-dependent manner. Moreover, deep RNA sequencing revealed aberrant miRNA maturation under hypoxia that was linked with increased ovarian cancer progression. Similarly, in breast cancer, enzymes involved in microRNA biogenesis (Dicer, Drosha, TARPB2, and DCGR8) were significantly repressed under hypoxic conditions (Bandara et al., 2014). Recently, Rupaimoole *et al.* (2016b) demonstrated that Dicer downregulation during hypoxia

results from hypoxia-induced miR-630, which targets the 3'-UTR of Dicer. They reported that mice treated with a combination of anti-miR-630 and anti-VEGF therapy presented smaller tumors with less metastatic nodules compared with mice treated with anti-VEGF therapy alone (Rupaimoole et al., 2016b). These findings not only provide novel insights on miRNA biogenesis downregulation detected under hypoxia, but also identify potential therapeutic targets that are deregulated in cancer. In another study, DICER suppression during hypoxia was due to epigenetic alteration. This involves inhibition of oxygen-dependent H3K27me3 demethylases, KDM6A/B, which consequently hypermethylate the *DICER* promoter, leading to DICER downregulation of in hypoxic breast cancer cells (van den Beucken et al., 2014).

### *HIF and lncRNAs*

Several long noncoding RNAs (lncRNAs) are also aberrantly expressed in hypoxic tumor microenvironments. LncRNAs are a heterogeneous class of non-coding RNAs, which are more than 200 nucleotides in length. They include antisense RNAs, transcribed ultraconserved regions (T-UCR), intergenic RNAs, and pseudogenes. Growing evidence has established that lncRNAs play vital roles in regulating genome at several levels, including genomic imprinting, transcription activation or inactivation, RNA splicing, translation control and RNA interference

Recent studies demonstrated the regulation of lncRNAs in response to hypoxia modulate the HIF pathway, metabolism, tumor growth, and metastasis (Choudhry et al., 2016). However, the exact mechanisms underlying the interaction of lncRNAs with the hypoxic tumor microenvironment require further elucidation. Current findings indicate two mechanisms of interaction of lncRNAs and hypoxia in cancer, **Figure 1**. First, hypoxia regulates hypoxia-responsive lncRNA expression via HIF. Additionally, lncRNAs modulate HIF expression and pathways **Table 2**.

HIF directly regulates lncRNAs transcription in hypoxia. *In silico* and experimental analyses identified the presence of HREs at the promoter region of most hypoxia-responsive lncRNAs. HIF ChIP-seq analysis identified that HIF-2 $\alpha$  bound slightly more at lncRNA promoters than HIF-1 $\alpha$  in hypoxic MCF-7 breast cancer cells. NEAT1 was identified among the most highly upregulated lncRNAs during

hypoxia and is dependent on HIF-2 $\alpha$  **Figure 1a** (Choudhry et al., 2015). On the other hand, UCA1 (Xue et al., 2014), EFNA3 (Gomez-Maldonado et al., 2015), H19 (Matouk et al., 2016), MALAT1 (Choudhry et al., 2014; Michalik et al., 2014), and HOTAIR (Bhan et al., 2017; Zhou et al., 2015), FALEC (Zhao et al., 2017), HAS2-AS1 (Zhu et al., 2017) are example of HIF-1 $\alpha$  dependent lncRNAs that contains HREs on their promoters. These studies provide evidence that the hypoxia-responsive lncRNAs can be directly regulated by HIF in different types of cancer.

Through different mechanisms, some lncRNAs can also be indirectly influenced by HIF. For instance, HIF upregulates the expression of WT1 lncRNA in a TET2- and TET3- (DNA demethylating enzymes) dependent manner in leukemia cells (McCarty and Loeb, 2015). Moreover, HIF-1 $\alpha$  dependent HDAC3 reduces lncRNA-LET expression by decreasing acetylation levels in the lncRNA-LET promoter region, suggesting that lncRNA-LET is indirectly regulated by HIF-1 $\alpha$  **Figure 1b** (Yang et al., 2013). These examples indicate that HIF indirectly regulates lncRNA expression via epigenetic modulation.

LncRNAs have the ability to control gene expression through diverse mechanisms as a guide, scaffold, decoy, or sponge. Recent studies demonstrated the role of lncRNAs in the direct and indirect regulation of HIF expression and pathway. An antisense lncRNA named HIF-2 $\alpha$  promoter upstream transcript (HIF2PUT) is a cis-regulatory lncRNA, which co-regulates *HIF-2 $\alpha$*  mRNA in osteosarcoma (Wang et al., 2015). Similarly, HIF-1 $\alpha$  locus contains two antisense transcripts, one transcribed from the 3'-UTR "3'aHIF-1 $\alpha$ " and the other from the 5'-promoter region "5'aHIF-1 $\alpha$ " of the sense *HIF-1 $\alpha$*  mRNA (Bertozi et al., 2011). These aHIF-1 $\alpha$  lncRNAs negatively regulate *HIF-1 $\alpha$*  mRNA level *in cis* by modulating chromatin or mRNA stability (Uchida et al., 2004) and associated with clinical outcome (Tasharrofi et al., 2016). In mesenchymal glioblastoma stem-like cells (GSCs), aHIF-1 $\alpha$  interacts with proteins including insulin-like growth factor 2 mRNA-binding protein 2 (IGF2BP2) and ATP-dependent RNA helicase A (DHX9) to regulate expression of high mobility group AT-hook 1 (HMGA1) and HIF-2 $\alpha$  in hypoxic stress (Mineo et al., 2016).. The above studies indicate a new role of lncRNAs directly regulating HIF levels.

Other lncRNAs indirectly regulate HIF levels. For instance, in pancreatic ductal adenocarcinoma (PDAC), low expression of lncRNA ENST00000480739 inhibits HIF-1 $\alpha$  expression by upregulating osteosarcoma amplified-9 (OS-9) (Sun et al., 2014). OS-9 induces an interaction between HIF-1 $\alpha$  and proline hydroxylase domain (PHD2/3) protein, leading to HIF-1 $\alpha$  degradation (Baek et al., 2005). OS-9 inhibition in hypoxia abrogates ENST00000480739-induced downregulation of HIF-1 $\alpha$ . In addition, ENST00000480739 upregulation significantly reduces the expression of HIF-1 $\alpha$ -dependent genes, including *MXI-1*, *PDGFC* and *MMP28* (Sun et al., 2014), indicating that ENST00000480739 may negatively control HIF-1 $\alpha$  activity by inducing OS-9 expression. Similarly, RERT-lncRNA transcriptionally upregulates PHD1 (EGLN2) expression and consequently suppress HIF-1 $\alpha$  activity (Zhu et al., 2012). Long intergenic non-coding RNA, regulator of reprogramming “Linc-RoR,” is also induced under hypoxia and modulates HIF-1 $\alpha$  expression. Suppression of hypoxia-induced linc-RoR reduces the expression of HIF-1 $\alpha$  and pyruvate dehydrogenase kinase isozyme 1 (PDK1), indicating that linc-RoR promotes hypoxia responses by controlling HIF-1 $\alpha$  levels and its target genes (Takahashi et al., 2014). Studies indicated that linc-RoR might function as a miRNA sponge under hypoxia to modulate and control the expression of miRNAs such as miR-145 that regulates the expression of HIF-1 $\alpha$  and HIF-dependent genes (Takahashi et al., 2014; Zhou et al., 2014). Similar, PVT1 and UCA1 act as sponges for miR-186 and miR-18a, respectively, hence regulate HIF-1 $\alpha$  expression (Huang et al., 2017; Li et al., 2016c). Low expression of Maternally expressed gene 3 (MEG3) lncRNA has been reported in multiple cancer types. Recently, downregulation of MEG3 promoter due to hypermethylation leads to c-Jun inhibition of PHLPP1 transcription, which enhance protein translation of HIF-1 $\alpha$  and activation of the Akt/p70S6K/S6 axis (Zhou et al., 2017). The lncRNA-SARCC (Suppressing Androgen Receptor in Renal Cell Carcinoma) found differentially respond to hypoxia in a VHL-dependent manner in renal cancer. LncRNA-SARCC physically interacts with androgen receptor (AR) to suppress the AR/HIF-2 $\alpha$ /C-MYC axis (Zhai et al., 2016). HIF-2 $\alpha$  suppress LncRNA-SARCC expression via binding to its hypoxia-responsive elements in the LncRNA promoter, suggesting a negative feedback loop between LncRNA-SARCC and HIF-2 $\alpha$  that modulates tumorigenesis for RCC in hypoxia.

Under hypoxia, HIF-1 $\alpha$  epigenetically decreases lncRNA-LET expression, while lncRNA-LET destabilizes *HIF-1 $\alpha$*  mRNA level via degrading nuclear factor 90 (NF90). Low lncRNA-LET expression promotes HIF-1 $\alpha$  accumulation (Yang et al., 2013). On the other hand, HIF-1 $\alpha$  directly binds at the promoter region for the transcriptional activation of lincRNA-p21, while lincRNA-p21 enhances HIF-1 $\alpha$  accumulation by disrupting the HIF-1 $\alpha$ -VHL interaction during hypoxia **Figure 1c** (Yang et al., 2014). Both lincRNA-p21 and lncRNA-LET form feedback loops that reciprocally regulate HIF levels under hypoxia.

LncRNAs can also modulate HIF-1 $\alpha$  accumulation and activation in normoxia. Recently, a ~1.5 kb long intergenic non-coding RNA for kinase activation (LINK-A) was found to be important for the growth factor-induced normoxic HIF-1 $\alpha$  signalling in triple receptor-negative breast cancer (TNBC) (Lin et al., 2016). LINK-A is essential for the recruitment and enzymatic activation of Tyrosine protein kinase 6, also known as breast tumour kinase (BRK). This activation is triggered by Heparin-binding EGF-like growth factor (HB-EGF), which mediates epidermal growth factor receptor (EGFR) and transmembrane glycoprotein NMB (GPNMB) heterodimerization 'EGFR:GPNMB'. LINK-A recruits activated BRK along with leucine-rich repeat kinase 2 (LRRK2) which leads to HIF-1 $\alpha$  phosphorylation at Tyr 565 and Ser 797, respectively. The phosphorylation at Tyr 565 suppresses hydroxylation at the Pro 564, which averts HIF1 $\alpha$  protosomal degradation in normoxia. Whereas, the Ser 797 phosphorylation enhances HIF1 $\alpha$ -p300 interaction resulting in HIF-1 $\alpha$  target genes activation in the presence of HB-EGF **Figure 1d** (Lin et al., 2016). Interestingly, LINK-A expression and LINK-A induced normoxic HIF1 $\alpha$  signalling activation were associated with TNBC and induced glycolysis reprogramming in breast cancer. Another example of a lncRNA regulating HIF-1 $\alpha$  activity is lncRNA cancer susceptibility candidate 9 (CASC9) which is found highly expressed in nasopharyngeal carcinoma (NPC) (Su et al., 2017). CASC9 binds HIF-1 $\alpha$  and promotes the stabilization and activation of HIF-1 $\alpha$ , thus inducing glycolysis metabolism and tumorigenesis. Both LINK-A and CASC9 need to be further investigated for their expression and function in hypoxia.

The host gene of microRNA-31 (MIR31HG), also known as a long noncoding HIF-1 $\alpha$  co-activating RNA (LncHIFCAR), was reported to activate the pseudohypoxia signature required for hypoxia-

induced metabolic reprogramming (Shih et al., 2017). Suppression of lncHIFCAR but not mir-31 significantly downregulated hypoxia-induced glucose uptake and lactate production. Mechanistically, lncHIFCAR physically interacts with HIF-1 $\alpha$  and binds to HIF-1 target genes. Consequently, this results in the recruitment of co-activator p300 which enabled the activation of HIF-1 transcriptional network (Shih et al., 2017). Moreover, hypoxia-responsive lncRNA uc.475 regulates polypeptide glycosylation (Ferdin et al., 2013). These are examples of lncRNAs that have an effect on the regulation or stabilization of HIF, hence they potentially modulate tumor metabolism.

Increasing evidence suggests that hypoxia-responsive lncRNAs play important roles in regulating all the hallmarks of cancer. Several hypoxia-responsive lncRNAs are involved in regulating apoptosis via different pathways. Suppression of HIF-1 $\alpha$ -dependent UCA1 during hypoxia induces apoptosis in bladder cancer cells through upregulation of Bax and cell cycle arrest at G1 phase and downregulation of Bcl-2 (Zhen et al., 2017). Hypoxic exosomal lncRNA-UCA1 reported to stimulate tumor growth and progression via EMT (Xue et al., 2017). H19 is overexpressed in various types of tumors with anti-apoptotic function and dependent on the tumor suppressor p53 (Matouk et al., 2016). NEAT1, HIF2PUT, MALAT1, and H19 regulate the proliferation of cancer cells during hypoxia. Hypoxia-induced NEAT1 is essential for the formation of nuclear structures called paraspeckles during hypoxia. NEAT1 knockdown remarkably inhibits cell proliferation and survival in both normoxia and hypoxia. However, the inhibition was more significant in hypoxic conditions (Choudhry et al., 2015). Similar regulation of cell proliferation was observed for UCA1 and hypoxia-induced noncoding ultraconserved transcripts (HINCUTs) (Ferdin et al., 2013; Xue et al., 2014). aHIF-1 $\alpha$  shown to promote mesenchymal glioblastoma stem-like cells (GSCs) growth, self-renewal, and hypoxia-dependent molecular reprogramming (Mineo et al., 2016). Recently, histone methyl-transferase MLL1 modulation with HIF-1 $\alpha$  and p300 found to control induction of HOTAIR under hypoxia hence promote tumorigenesis (Bhan et al., 2017).

The hypoxic microenvironment promotes angiogenesis and EMT via a series of signaling pathways, including HIF pathways. Examples of lncRNAs that probably regulate cancer angiogenesis include linc-ROR, which regulates HIF-1 activity (Takahashi et al., 2014) and hypoxia-induced MALAT1

that promotes vascular growth *in vivo* (Michalik et al., 2014). HIF-1 $\alpha$  dependent induction of HAS2-AS1 promotes hypoxia-regulated EMT and invasiveness via stabilizing HAS2 (Zhu et al., 2017).

In addition to miRNAs and lncRNAs, in-depth investigation of hypoxic non-coding transcriptome identified other classes of hypoxia-responsive non-coding RNAs such as snRNAs, piwiRNAs, tRNAs, and circular RNAs (Boeckel et al., 2015; Choudhry et al., 2016). However, their functional implication in hypoxia biology and metabolism remains to be determined.

### **Hypoxia, HIFs and the circadian clock**

In living organisms, the circadian clock characterizes the 24-h oscillations that control daily rhythmicity of molecular, physiological, and behavioral processes. A master clock modulating circadian rhythms is located in the suprachiasmatic nucleus (SCN) of the hypothalamus (Takahashi, 2017). The SCN is a central oscillator induced by environmental signals, thus coordinating the SCN-driven output rhythms. A set of genes coordinate the biological rhythms including clock genes, CLOCK/NPAS2, BMAL1/2(ARNTL/2), ARNT3, Period (PER1, PER2, and PER3), cryptochrome (CRY1 and CRY2), and MOP3. In mammals, the molecular clock is composed of activators (CLOCK/BMAL) that promote the transcription of repressors (PER/CRY) that, in turn, suppress the forward limb in a cycle, which repeats itself every 24 hours, with an additional supporting loop involving the nuclear receptor REV-ERB and the retinoic acid receptor-related orphan receptor (ROR) transcription factors (Takahashi, 2017). A growing number of studies demonstrated that alteration of the circadian clock is linked with the initiation and progression of several types of cancers, including HCC, colorectal, breast, blood and lung cancers (Fu and Kettner, 2013).

Recent studies reported interrelationships between hypoxic response pathways and circadian pathways. The fundamental components of circadian clockwork and oxygen homeostasis are the PAS protein family members (PER and CLOCK) and HIF-1 $\alpha$ . Numerous genes are involved in the circadian clock circuitry, and response to hypoxia is deregulated in diseases and stress conditions, hence influencing the physiological process as well as disease progression and outcome.

The underlying mechanism could involve the transcriptional control of circadian gene expression by HIF or through HIF protein–protein interactions, leading to decreased proteolytic degradation of circadian proteins. *In vivo* experiments revealed that hypoxia upregulates levels of circadian proteins, PER1 and CLOCK, possibly through protein interaction between PER1 and HIF-1 $\alpha$  (Chilov et al., 2001). In addition, HIF-1 $\alpha$  interacts with BMAL1/MOP3 and CLOCK to regulate gene expression (Ghorbel et al., 2003; Hogenesch et al., 1998), suggesting that HIF-1 $\alpha$  may regulate the circadian clockwork, and the crosstalk between hypoxia and circadian pathways modulates the expression of target genes. Hypoxia-induced HCC with CoCl<sub>2</sub> revealed an increase in the expression of Clock, BMAL1, and CRY2, and decreased levels of PER1, PER2, PER3, CRY1, and CKI $\epsilon$  in hypoxia. Both HIF-1 $\alpha$  and HIF-2 $\alpha$  are involved in regulating the expression of circadian genes in HCC. However, they transcriptionally regulate different set of circadian genes (Yu et al., 2015).

The post-translational modifications of core clock proteins significantly contribute in the circadian system flux. These modifications control the timing between the activation and the repression of circadian transcription to regulate distinct functions. For example, BMAL1, a clock protein, is subject to various post-translational modifications, including phosphorylation through casein kinase (CK) 1  $\delta/\epsilon$  and CK 2 (Agostino et al., 2009; Sahar et al., 2010). Under hypoxia, CK1 $\delta$  plays a vital role, phosphorylating the N-terminal heterodimerization (PAS) domain of HIF-1 $\alpha$ . This obstructs the association with ARNT, thereby controlling the activity of HIF-1 $\alpha$  during hypoxia (Kalouisi et al., 2010).

The circadian organization of the molecular clockwork affects VEGF levels in hypoxic cells. For instance, PER2 and CRY1 inhibit hypoxia-induced VEGF promoter activity, indicating a negative feedback loop, which periodically suppresses the transcriptional upregulation of *VEGF* during hypoxia, resulting in the circadian fluctuation of its gene expression (Koyanagi et al., 2003). Moreover, the basic helix-loop-helix (bHLH) transcription factors, differentiated embryonic chondrocyte-expressed gene 1 (DEC1) and DEC2, are part of an important crosstalk between hypoxia responses and circadian pathways (Sato et al., 2016).

Recently, number of studies have showed the crosstalk and feedback loops between circadian clock and oxygen sensing pathways **Figure 2**. Peek *et al.* (2017) reported a bidirectional crosstalk between circadian and HIF pathways that modulates the metabolic adaptation in a tissue-specific manner. Genetic disruption of the core clock component, BMAL1, in mouse skeletal muscle reduces anaerobic glycolysis, HIF-1 $\alpha$  activity, and the expression of its target genes, including *PHD3*, *VEGFA*, *MCT4*, *PK-M*, and *LDHA*. HIF-1 $\alpha$  binds directly on the promoters of several core clock genes. In response to strenuous exercise, expression level of CLOCK and HIF-1 $\alpha$  target genes varied depending on the time of day in mice. They concluded that circadian clock cooperate with HIF-1 $\alpha$  to mediate anaerobic glycolysis in Muscle (Peek *et al.*, 2017). Another study showed that the hypoxia response is gated by the circadian clock through HIF-1 $\alpha$  that act as a regulatory node that connects both pathways (Wu *et al.*, 2017). Genome-wide analysis of HIF-1 $\alpha$  and BMAL1 binding revealed hypoxia-clock reciprocal regulation at the genome level. An *in vivo* heart attack model described that the circadian clock protects the heart from hypoxia-induced cell death (Wu *et al.*, 2017), suggesting that the circadian clock can be utilized therapeutically to reduce the severity of hypoxia-associated diseases. Adamovich *et al.* (2017) measured continuous oxygen levels in the blood and tissue of rodents and identified daily rhythms in tissue oxygenation. Interestingly, the physiological oxygen rhythms synchronize clocks in cultured cells in a HIF-1 $\alpha$ -dependent manner. Moreover, modulation of oxygen levels hasten the recovery of wild-type mice from a jet lag protocol, but not that of HIF-1 $\alpha$ -deficient mice (Adamovich *et al.*, 2017). These recent studies provide strong evidence of important crosstalk between circadian and hypoxia signaling pathways in HIF-1 $\alpha$  dependent fashion (Adamovich *et al.*, 2017; Peek *et al.*, 2017; Wu *et al.*, 2017) **Figure 2**.

Currently, circadian rhythm and hypoxia-signaling are active areas of research aiming at identifying the molecular mechanisms and regulators of these pathways in cells. The absence of models in which these pathways can be investigated under controlled conditions delayed our progress in understanding the crosstalk between hypoxia and the circadian clock in physiological and pathological states.

Interestingly, circadian clocks are capable of not only predict changes in the external light cycle, but also regulate adaptation to flux in nutrient and oxygen sensing.

### **Hypoxia, HIFs and cellular vesicles**

The discovery of cellular vesicles including exosomes and microvesicles led to a better understanding of cell-cell interaction and communication. Exosomes originate from endocytic multivesicular bodies that follow the endosomal pathway and end up with exocytosis. Exosomes range from 30–100 nm in size. Recent studies suggest that cancer-associated exosomes, through their content, induce metastasis and create a pre-metastatic niche. For instance, low cellular pH is a hallmark of tumor malignancy. In cancer cells, the acidic microenvironment leads to increased exosomal release and uptake by recipient cells (Parolini et al., 2009). Riches *et al.* measured the release of exosomes from normal human mammary and breast cancer cells. They reported a significant increase in release of exosomes from cancer cells compared to that from normal parent cells within 24 hours (Riches et al., 2014). Recent findings clearly suggest that cellular vesicles serve as a significant mediator of cell-to-cell communication within the tumor microenvironment to facilitate tumorigenesis and modulate metabolism. For example, breast cancer cells can suppress glucose uptake of stromal cells in the pre-metastatic niche, by releasing cellular vesicles that contain high levels of the miR-122 which downregulate the glycolytic enzyme pyruvate kinase (Fong et al., 2015). *In vivo* suppression of miR-122 restores glucose uptake in distant organs and significantly reduce metastasis. These data suggest breast cancer cells derived vesicles can reprogram energy metabolism and create a niche with high glucose availability for their own consumption. Proteomic analysis of exosomes released from HCC revealed enrichment of glucose metabolism regulatory proteins (Zhang et al., 2017). Exosomal miR-126 is an emerging novel regulator of different aspects of cell metabolism including glucose homeostasis, mitochondrial respiration and regulation of genes involved in gluconeogenesis and oxidative stress (Tao et al., 2016; Tomasetti et al., 2014). Recently, cancer-associated fibroblasts exosomes shown to promote metabolic reprogramming in cancer cells through suppressing mitochondrial function and inducing glucose metabolism of other cancer cells (Zhao et al., 2016).

Increasing evidence suggests that hypoxia secreted cellular vesicles in a tumor microenvironment are involved in a number of functions, such as prompting intratumoral heterogeneity, responding to immunological reactions, inducing cancer-associated fibroblasts, metabolic reprogramming and promoting angiogenesis and metastasis.

#### *HIF and exosomes*

Many studies reported that tumor hypoxia increases the release of exosomes from malignant cells. HIF regulates the expression of many plasma membrane receptors, including glucose transporter (GLUT-1), transferrin receptor, and Epidermal Growth Factor Receptor (EGFR), and it is thought that increased expression of receptors causes their activation and internalization, which consequently induces endocytosis and promotes exosome release. Hypoxia-induced exosomes content varies depending on cell origin including signal transducers, transcription factors, enzyme, lipids, mRNAs, and non-coding RNAs **Table 3**. Aberrant exosomal secretion is associated with the adaptation to a different microenvironment that promotes cancer survival. King *et al.* (2012) found that exosome secretion increased as which contained high levels of HIF-regulated miR-210 (King *et al.*, 2012).

Several studies indicated that hypoxia-induced exosomes contribute to tumor angiogenesis and metastasis. Aga *et al.* (2013) detected HIF-1 $\alpha$  in exosomes secreted from invasive Epstein–Barr virus (EBV) malignancy, nasopharyngeal carcinoma (NPC). The oncoprotein, latent membrane protein 1 (LMP1), upregulates HIF-1 $\alpha$  expression in exosomes. Interestingly, transcriptionally active *HIF-1 $\alpha$*  was detected in recipient cells upon exosome uptake. In addition, HIF-1 $\alpha$  regulates exosome-mediated pro-metastatic effects through epithelial–mesenchymal transition (EMT) changes in E- and N-cadherin expression in the recipient cells (Aga *et al.*, 2014). Exosomes secreted from prostate cancer under hypoxia enhance stemness, invasiveness, and EMT by modulating the cancer-associated fibroblast phenotype in prostate stromal cells (Ramteke *et al.*, 2015). Hypoxic prostate cancer exosomes are unique and express high levels of proteins such as heat shock proteins (HSP90 and HSP70), matrix metalloproteinases, AKT, annexin II, IL6, TGF $\beta$ 2,  $\beta$ -catenin, TNF1 $\alpha$ , CD63, and CD81 compared to those observed in normoxic secreted exosomes (Ramteke *et al.*, 2015). Similarly, Rong *et al.* (2016)

showed that hypoxia induces the secretion of exosomes that express immunosuppressive cytokines such as TGF- $\beta$  and IL-10 from breast cancer cells (Rong et al., 2016).

Co-culturing of K562 leukemic cells with human umbilical vein endothelial cells (HUVECs) significantly promoted tube formation by HUVECs during hypoxia (Tadokoro et al., 2013). Exosomes released during hypoxia were similar in size as those secreted during normoxia. Interestingly, exosomal miR-210 released from leukemic cells during hypoxia downregulates the expression of the anti-angiogenic factor, Ephrin-A3 (EFNA3), in HUVECs, indicating the interaction between cancer and endothelial cells via exosomes to modulate angiogenesis during hypoxia (Tadokoro et al., 2013). Similarly, Umezu *et al.* (2014) developed an *in vitro* chronic hypoxia model by continuous growth of multiple myeloma (MM) cell lines (more than 6 months) under hypoxic conditions, named hypoxia-resistant MM cells. This model mimics the *in vivo* microenvironment of bone marrow myeloma cells. The number of exosomes secreted from hypoxia-resistant MM cells is twice that secreted by parental cells grown in normoxia or hypoxia (Umezu et al., 2014). Exosomes released from hypoxia-resistant MM cells promote tube formation of HUVECs during normoxia, but to a greater degree under hypoxia. Interestingly, secreted miR-135b by hypoxia-resistant MM cells induces the endothelial tube formation in hypoxia and enhances HIF-1 transcriptional activity by inhibiting factor-inhibiting hypoxia-inducible factor 1 (FIH-1) (Umezu et al., 2014). Further studies reported that exosomes secreted from pericytes upon HIF activation promote the angiogenic activity of endothelial cells (Mayo and Bearden, 2015). Moreover, exosomes derived from hypoxic endothelial cells remodel the extracellular matrix via exosome-associated lysyl oxidase-like 2 (LOXL2) for fibrosis and wound healing (de Jong et al., 2016). Recently, Ling *et al.* demonstrated that hypoxia increases oral squamous cell carcinoma (OSCC)-derived exosomes, which promotes cell migration and invasion in a HIF-1 $\alpha$ - and HIF-2 $\alpha$ -dependent manner (Li et al., 2016a). miR-21 was the most significantly upregulated miRNA in exosomes released from hypoxic OSCC cells. miR-21 suppression in hypoxic OSCC cells reduced miR-21 expression in exosomes and significantly impaired cancer cell migration and invasion. Exosomal miR-21 promotes the expression of snail and vimentin, while decreasing E-cadherin expression. Moreover, exosomal miR-21 levels and HIF-1 $\alpha$ /HIF-2 $\alpha$  expression were associated with lymph node metastasis in patients

with OSCC (Li et al., 2016a). Moreover, miR-24-3p is found enriched in hypoxic cells and exosomes from neural progenitor cells (NPC) cells and serum. miR-24-3p regulates T-cell proliferation and differentiation and may be a prognostic marker for NPC (Ye et al., 2016).

Both *in vitro* and clinical studies using specimens from patients with glioblastoma multiforme (GBM) showed enrichment of hypoxia-regulated mRNAs and proteins such as platelet-derived growth factors (PDGF), matrix metalloproteinases, IL-8, caveolin 1, and lysyl oxidase in exosomes (Kucharzewska et al., 2013). Many of these molecules modulate angiogenesis and are associated with poor clinical outcome in patients with glioma. In GBM, hypoxia-induced exosomes stimulate endothelial cells through several growth factors and cytokines, leading to the activation of PI3K/AKT signaling and migration. Hypoxia-induced GBM exosomes significantly increase cell proliferation, tumor vascularization, and pericyte vessel coverage (Kucharzewska et al., 2013). This study indicates that exosomal protein and RNA content recapitulates the oxygenation status of exosome releasing cells and patient tumors.

Hypoxic regions are also detected in adipose tissues because of adipocyte hypertrophy. Hypoxia induces release of exosomes from differentiated adipocytes. Hypoxia-derived exosomes are enriched in proteins responsible for *de novo* conversion of acetyl-CoA to fatty acids, including fatty acid synthase, acetyl-CoA carboxylase, and glucose-6-phosphate dehydrogenase, suggesting a role of hypoxia-induced release of exosomes by adipocytes in regulating lipogenesis. Given that lipid metabolism is one of the hallmarks of hypoxic cancer cells, increased accumulation of lipids during hypoxia may stimulate and induce the biogenesis and secretion of extracellular vesicles. Hypoxic prostate cancer exosomes are significantly enriched in triglycerides, owing to the activation of lipogenic enzymes and signaling molecules, which play important roles in prostate cancer invasiveness (Schlaepfer et al., 2015). Moreover, fatostatin and silibinin (lipogenesis inhibitors) reduce the concentration of extracellular vesicles and VEGF level in exosomes, suggesting the important role of lipid metabolism in hypoxic prostate cancer exosomes supporting cancer invasiveness and metastasis.

### *HIF and microvesicles*

HIF also regulates other extracellular vesicles such as microvesicles. These vesicles are budded directly from the membranes and shed from almost all cell types. Microvesicles have a wide range of sizes from 100 nm to 1  $\mu$ m and play an important role in intercellular communication. Wang *et al.* (2014) reported that HIF mediates microvesicle production by breast cancer cells during hypoxia. HIF increases the activity of a member of the Rab family of small GTPases, RAB22A, which co-localizes with budding microvesicles at the cell surface. RAB22A is HIF-dependent and essential for the formation of vesicle, trafficking, and membrane fusion. Thus, microvesicle biogenesis depends on HIF-1 $\alpha$  during hypoxia. RAB22A is required to promote invasion and metastasis of recipient breast cancer cells (Wang *et al.*, 2014). Similarly, Berchem *et al.* (2016) reported that hypoxia-derived microvesicles from GR-Heu and K562 tumor cells suppress natural killer (NK) cell function and decrease cytotoxicity compared to macrovesicles released during normoxia. Impaired cytotoxicity of NK cells by hypoxia-derived microvesicles results from decreased expression of NKG2D in a tumor growth factor (TGF- $\beta$ 1)-dependent manner. In addition, miR-23a is significantly overexpressed during hypoxia when compared to that in normoxic tumor microvesicles. miR-23a directly regulates the expression of Lysosomal-associated membrane protein 1 (LAMP-1)/CD107a in NK cells (Berchem *et al.*, 2016), suggesting that suppressive signals of immune cells such as TGF- $\beta$ 1 and miR23a in the hypoxia-derived microvesicles impair the antitumor immune response during hypoxia.

The above studies suggest that hypoxia/HIF affects the synthesis of extracellular vesicles and their biological effects in the tumor microenvironment, contributing to the major hallmarks of cancer and supplementing many of the pathways induced directly in cells by hypoxia e.g. metabolism, angiogenesis and EMT. It is also reasonable to think that cellular vesicles are heterogeneous and this may contribute to tumour heterogeneity adding new layer of complexity in our understanding of cancer.

Hypoxia/HIF regulation on exosome and microvesicles function is a new and exciting area of research, however, it is in early stages. The direct mechanism of extracellular vesicles induction by HIF and

impact of HIF in modulating processes including cellular vesicles formation, content selection, loading, trafficking, and release remains to be determined. How these processes are orchestrated during hypoxia should be further investigated.

### **Future Perspectives**

Decades of research on hypoxia and HIF biology significantly improved our understanding of oxygen homeostasis in health and diseases. However, advancement in analytical tools and genomic technologies opened new important areas for investigation. Recently, pan-genomic analysis several cancer risk polymorphisms overlap with HIF-binding sites hence modulate HIF signalling and cancer development (Grampp et al., 2016; Grampp et al., 2017).

The role of HIF in hypoxia responses has been the topic of many investigations. Some of these studies provide inconsistent findings because of the experimental tools used and design biases. For instance, researchers utilized either cell-based *in vitro* models or animal models maintained under hypoxia to infer their conclusions. However, hypoxia responses vary between cell lines from the same cancer type and between models and within subpopulations of the same cell line (Ledaki et al., 2015). The other major barrier to investigating the effects of hypoxia on tumor cells *in vitro* is that cells can only grow for a few days under hypoxic conditions. In most studies, cells are commonly exposed to hypoxia for 24–72 h. In such cases, tumor cells might respond to acute hypoxic stress, which could be different from long-term intratumoral hypoxia.

The new mechanisms described above point to many ways of individualising responses in each cell, influencing neighbouring cells and those at a great distance, diversifying the response well beyond the recognised core metabolism genes. This provides the possibility of substantial advances in biomarkers of hypoxia in specific tissues, but also far more specific modulation of the hypoxia response than previously recognised e.g. use of JQ1 a BRD4 modulator that specifically inhibited induction of a subset of genes in hypoxia (da Motta et al., 2017).

Deciphering the role of epigenetics, non-coding RNAs, circadian rhythms and extracellular vesicles in hypoxia will advance our understanding of cellular modulation under low oxygen stress, and should lead to the development of more specific therapeutic agents.

### **Acknowledgements**

The authors would like to acknowledge the support from the Deanship of Scientific Research (DSR), King Abdulaziz University (KAU), Jeddah the Ministry of Education for Saudi Arabia as well as the KAU Research Endowment Fund (WAQF) (HC). In addition, acknowledge the support from the Cancer Research UK (ALH) and Breast Cancer Research Foundation (ALH).

**Conflict of interest disclosure:** None

## References

- Adamovich, Y., Ladeuix, B., Golik, M., Koeners, M.P., and Asher, G. (2017). Rhythmic Oxygen Levels Reset Circadian Clocks through HIF1alpha. *Cell Metab* 25, 93-101.
- Aga, M., Bentz, G.L., Raffa, S., Torrisi, M.R., Kondo, S., Wakisaka, N., Yoshizaki, T., Pagano, J.S., and Shackelford, J. (2014). Exosomal HIF1alpha supports invasive potential of nasopharyngeal carcinoma-associated LMP1-positive exosomes. *Oncogene* 33, 4613-4622.
- Agostino, P.V., Harrington, M.E., Ralph, M.R., and Golombek, D.A. (2009). Casein kinase-1-epsilon (CK1epsilon) and circadian photic responses in hamsters. *Chronobiol Int* 26, 126-133.
- Agrawal, R., Pandey, P., Jha, P., Dwivedi, V., Sarkar, C., and Kulshreshtha, R. (2014). Hypoxic signature of microRNAs in glioblastoma: insights from small RNA deep sequencing. *BMC Genomics* 15, 686.
- Bacon, A.L., Fox, S., Turley, H., and Harris, A.L. (2007). Selective silencing of the hypoxia-inducible factor 1 target gene BNIP3 by histone deacetylation and methylation in colorectal cancer. *Oncogene* 26, 132-141.
- Baek, J.H., Mahon, P.C., Oh, J., Kelly, B., Krishnamachary, B., Pearson, M., Chan, D.A., Giaccia, A.J., and Semenza, G.L. (2005). OS-9 interacts with hypoxia-inducible factor 1alpha and prolyl hydroxylases to promote oxygen-dependent degradation of HIF-1alpha. *Mol Cell* 17, 503-512.
- Bailey, K.M., Wojtkowiak, J.W., Hashim, A.I., and Gillies, R.J. (2012). Targeting the metabolic microenvironment of tumors. *Adv Pharmacol* 65, 63-107.
- Bandara, V., Michael, M.Z., and Gleadle, J.M. (2014). Hypoxia represses microRNA biogenesis proteins in breast cancer cells. *BMC Cancer* 14, 533.
- Bandara, V., Michael, M.Z., and Gleadle, J.M. (2017). MicroRNA biogenesis in hypoxia. *Microna*.
- Berchem, G., Noman, M.Z., Bosseler, M., Paggetti, J., Baconnais, S., Le Cam, E., Nanbakhsh, A., Moussay, E., Mami-Chouaib, F., Janji, B., et al. (2016). Hypoxic tumor-derived microvesicles negatively regulate NK cell function by a mechanism involving TGF-beta and miR23a transfer. *Oncoimmunology* 5, e1062968.
- Bertozzi, D., Iurlaro, R., Sordet, O., Marinello, J., Zaffaroni, N., and Capranico, G. (2011). Characterization of novel antisense HIF-1alpha transcripts in human cancers. *Cell Cycle* 10, 3189-3197.
- Beyer, S., Kristensen, M.M., Jensen, K.S., Johansen, J.V., and Staller, P. (2008). The histone demethylases JMJD1A and JMJD2B are transcriptional targets of hypoxia-inducible factor HIF. *J Biol Chem* 283, 36542-36552.
- Bhan, A., Deb, P., Shihabeddin, N., Ansari, K.I., Brotto, M., and Mandal, S.S. (2017). Histone methylase MLL1 coordinates with HIF and regulate lncRNA HOTAIR expression under hypoxia. *Gene* 629, 16-28.
- Blick, C., Ramachandran, A., McCormick, R., Wigfield, S., Cranston, D., Catto, J., and Harris, A.L. (2015). Identification of a hypoxia-regulated miRNA signature in bladder cancer and a role for miR-145 in hypoxia-dependent apoptosis. *Br J Cancer* 113, 634-644.
- Boeckel, J.N., Jae, N., Heumuller, A.W., Chen, W., Boon, R.A., Stellos, K., Zeiher, A.M., John, D., Uchida, S., and Dimmeler, S. (2015). Identification and Characterization of Hypoxia-Regulated Endothelial Circular RNA. *Circ Res* 117, 884-890.
- Bruning, U., Cerone, L., Neufeld, Z., Fitzpatrick, S.F., Cheong, A., Scholz, C.C., Simpson, D.A., Leonard, M.O., Tambuwala, M.M., Cummins, E.P., et al. (2011). MicroRNA-155 promotes resolution of hypoxia-inducible factor 1alpha activity during prolonged hypoxia. *Mol Cell Biol* 31, 4087-4096.
- Camps, C., Saini, H.K., Mole, D.R., Choudhry, H., Reczko, M., Guerra-Assuncao, J.A., Tian, Y.M., Buffa, F.M., Harris, A.L., Hatzigeorgiou, A.G., et al. (2014). Integrated analysis of microRNA and mRNA expression and association with HIF binding reveals the complexity of microRNA expression regulation under hypoxia. *Mol Cancer* 13, 28.

Cascella, B., and Mirica, L.M. (2012). Kinetic analysis of iron-dependent histone demethylases: alpha-ketoglutarate substrate inhibition and potential relevance to the regulation of histone demethylation in cancer cells. *Biochemistry* 51, 8699-8701.

Chan, S.Y., Zhang, Y.Y., Hemann, C., Mahoney, C.E., Zweier, J.L., and Loscalzo, J. (2009). MicroRNA-210 controls mitochondrial metabolism during hypoxia by repressing the iron-sulfur cluster assembly proteins ISCU1/2. *Cell Metab* 10, 273-284.

Chen, B., Zeng, X., He, Y., Wang, X., Liang, Z., Liu, J., Zhang, P., Zhu, H., Xu, N., and Liang, S. (2016). STC2 promotes the epithelial-mesenchymal transition of colorectal cancer cells through AKT-ERK signaling pathways. *Oncotarget* 7, 71400-71416.

Chen, Q., Chen, Y., Bian, C., Fujiki, R., and Yu, X. (2013). TET2 promotes histone O-GlcNAcylation during gene transcription. *Nature* 493, 561-564.

Chen, S., Yin, C., Lao, T., Liang, D., He, D., Wang, C., and Sang, N. (2015). AMPK-HDAC5 pathway facilitates nuclear accumulation of HIF-1alpha and functional activation of HIF-1 by deacetylating Hsp70 in the cytosol. *Cell Cycle* 14, 2520-2536.

Chilov, D., Hofer, T., Bauer, C., Wenger, R.H., and Gassmann, M. (2001). Hypoxia affects expression of circadian genes PER1 and CLOCK in mouse brain. *FASEB J* 15, 2613-2622.

Choudhry, H., Albukhari, A., Morotti, M., Haider, S., Moralli, D., Smythies, J., Schodel, J., Green, C.M., Camps, C., Buffa, F., et al. (2015). Tumor hypoxia induces nuclear paraspeckle formation through HIF-2alpha dependent transcriptional activation of NEAT1 leading to cancer cell survival. *Oncogene* 34, 4482-4490.

Choudhry, H., Harris, A.L., and McIntyre, A. (2016). The tumour hypoxia induced non-coding transcriptome. *Mol Aspects Med* 47-48, 35-53.

Choudhry, H., Schodel, J., Oikonomopoulos, S., Camps, C., Grampp, S., Harris, A.L., Ratcliffe, P.J., Ragoussis, J., and Mole, D.R. (2014). Extensive regulation of the non-coding transcriptome by hypoxia: role of HIF in releasing paused RNAPol2. *EMBO Rep* 15, 70-76.

da Motta, L.L., Ledaki, I., Purshouse, K., Haider, S., De Bastiani, M.A., Baban, D., Morotti, M., Steers, G., Wigfield, S., Bridges, E., et al. (2017). The BET inhibitor JQ1 selectively impairs tumour response to hypoxia and downregulates CA9 and angiogenesis in triple negative breast cancer. *Oncogene* 36, 122-132.

Dawson, M.A. (2017). The cancer epigenome: Concepts, challenges, and therapeutic opportunities. *Science* 355, 1147-1152.

de Jong, O.G., van Balkom, B.W., Gremmels, H., and Verhaar, M.C. (2016). Exosomes from hypoxic endothelial cells have increased collagen crosslinking activity through up-regulation of lysyl oxidase-like 2. *J Cell Mol Med* 20, 342-350.

Estaras, C., Fueyo, R., Akizu, N., Beltran, S., and Martinez-Balbas, M.A. (2013). RNA polymerase II progression through H3K27me3-enriched gene bodies requires JMJD3 histone demethylase. *Mol Biol Cell* 24, 351-360.

Fasanaro, P., D'Alessandra, Y., Di Stefano, V., Melchionna, R., Romani, S., Pompilio, G., Capogrossi, M.C., and Martelli, F. (2008). MicroRNA-210 modulates endothelial cell response to hypoxia and inhibits the receptor tyrosine kinase ligand Ephrin-A3. *J Biol Chem* 283, 15878-15883.

Favaro, E., Lord, S., Harris, A.L., and Buffa, F.M. (2011). Gene expression and hypoxia in breast cancer. *Genome Med* 3, 55.

Favaro, E., Ramachandran, A., McCormick, R., Gee, H., Blancher, C., Crosby, M., Devlin, C., Blick, C., Buffa, F., Li, J.L., et al. (2010). MicroRNA-210 regulates mitochondrial free radical response to hypoxia and krebs cycle in cancer cells by targeting iron sulfur cluster protein ISCU. *PLoS One* 5, e10345.

Feng, C.C., Lin, C.C., Lai, Y.P., Chen, T.S., Marthandam Asokan, S., Lin, J.Y., Lin, K.H., Viswanadha, V.P., Kuo, W.W., and Huang, C.Y. (2016). Hypoxia suppresses myocardial survival pathway through HIF-1alpha-IGFBP-3-dependent signaling and enhances cardiomyocyte autophagic and apoptotic effects mainly via FoxO3a-induced BNIP3 expression. *Growth Factors* 34, 73-86.

Ferdin, J., Nishida, N., Wu, X., Nicoloso, M.S., Shah, M.Y., Devlin, C., Ling, H., Shimizu, M., Kumar, K., Cortez, M.A., et al. (2013). HINCUTs in cancer: hypoxia-induced noncoding ultraconserved transcripts. *Cell Death Differ* 20, 1675-1687.

Fong, M.Y., Zhou, W., Liu, L., Alontaga, A.Y., Chandra, M., Ashby, J., Chow, A., O'Connor, S.T., Li, S., Chin, A.R., et al. (2015). Breast-cancer-secreted miR-122 reprograms glucose metabolism in premetastatic niche to promote metastasis. *Nat Cell Biol* 17, 183-194.

Fu, L., Chen, L., Yang, J., Ye, T., Chen, Y., and Fang, J. (2012). HIF-1alpha-induced histone demethylase JMJD2B contributes to the malignant phenotype of colorectal cancer cells via an epigenetic mechanism. *Carcinogenesis* 33, 1664-1673.

Fu, L., and Kettner, N.M. (2013). The circadian clock in cancer development and therapy. *Prog Mol Biol Transl Sci* 119, 221-282.

Geng, H., Harvey, C.T., Pittsenbarger, J., Liu, Q., Beer, T.M., Xue, C., and Qian, D.Z. (2011). HDAC4 protein regulates HIF1alpha protein lysine acetylation and cancer cell response to hypoxia. *J Biol Chem* 286, 38095-38102.

Ghorbel, M.T., Coulson, J.M., and Murphy, D. (2003). Cross-talk between hypoxic and circadian pathways: cooperative roles for hypoxia-inducible factor 1alpha and CLOCK in transcriptional activation of the vasopressin gene. *Mol Cell Neurosci* 22, 396-404.

Ghosh, G., Subramanian, I.V., Adhikari, N., Zhang, X., Joshi, H.P., Basi, D., Chandrashekhar, Y.S., Hall, J.L., Roy, S., Zeng, Y., et al. (2010). Hypoxia-induced microRNA-424 expression in human endothelial cells regulates HIF-alpha isoforms and promotes angiogenesis. *J Clin Invest* 120, 4141-4154.

Gomez-Maldonado, L., Tiana, M., Roche, O., Prado-Cabrero, A., Jensen, L., Fernandez-Barral, A., Guijarro-Munoz, I., Favaro, E., Moreno-Bueno, G., Sanz, L., et al. (2015). EFNA3 long noncoding RNAs induced by hypoxia promote metastatic dissemination. *Oncogene* 34, 2609-2620.

Gorospe, M., Tominaga, K., Wu, X., Fahling, M., and Ivan, M. (2011). Post-Transcriptional Control of the Hypoxic Response by RNA-Binding Proteins and MicroRNAs. *Front Mol Neurosci* 4, 7.

Grampp, S., Platt, J.L., Lauer, V., Salama, R., Kranz, F., Neumann, V.K., Wach, S., Stohr, C., Hartmann, A., Eckardt, K.U., et al. (2016). Genetic variation at the 8q24.21 renal cancer susceptibility locus affects HIF binding to a MYC enhancer. *Nat Commun* 7, 13183.

Grampp, S., Schmid, V., Salama, R., Lauer, V., Kranz, F., Platt, J.L., Smythies, J., Choudhry, H., Goppelt-Struebe, M., Ratcliffe, P.J., et al. (2017). Multiple renal cancer susceptibility polymorphisms modulate the HIF pathway. *PLoS Genet* 13, e1006872.

Haque, I., Banerjee, S., Mehta, S., De, A., Majumder, M., Mayo, M.S., Kambhampati, S., Campbell, D.R., and Banerjee, S.K. (2011). Cysteine-rich 61-connective tissue growth factor-nephroblastoma-overexpressed 5 (CCN5)/Wnt-1-induced signaling protein-2 (WISP-2) regulates microRNA-10b via hypoxia-inducible factor-1alpha-TWIST signaling networks in human breast cancer cells. *J Biol Chem* 286, 43475-43485.

Hattori, M., Yokoyama, Y., Hattori, T., Motegi, S., Amano, H., Hatada, I., and Ishikawa, O. (2015). Global DNA hypomethylation and hypoxia-induced expression of the ten eleven translocation (TET) family, TET1, in scleroderma fibroblasts. *Exp Dermatol* 24, 841-846.

Hatzimichael, E., Dasoula, A., Shah, R., Syed, N., Papoudou-Bai, A., Coley, H.M., Dranitsaris, G., Bourantas, K.L., Stebbing, J., and Crook, T. (2010). The prolyl-hydroxylase EGLN3 and not EGLN1 is inactivated by methylation in plasma cell neoplasia. *Eur J Haematol* 84, 47-51.

Hogenesch, J.B., Gu, Y.Z., Jain, S., and Bradfield, C.A. (1998). The basic-helix-loop-helix-PAS orphan MOP3 forms transcriptionally active complexes with circadian and hypoxia factors. *Proc Natl Acad Sci U S A* 95, 5474-5479.

Huang, T., Liu, H.W., Chen, J.Q., Wang, S.H., Hao, L.Q., Liu, M., and Wang, B. (2017). The long noncoding RNA PVT1 functions as a competing endogenous RNA by sponging miR-186 in gastric cancer. *Biomed Pharmacother* 88, 302-308.

Huang, Z., and Feng, Y. (2016). Exosomes derived from hypoxic colorectal cancer cells promotes angiogenesis through Wnt4 induced beta-catenin signaling in endothelial cells. *Oncol Res*.

Isaacs, J.S., Jung, Y.J., Mole, D.R., Lee, S., Torres-Cabala, C., Chung, Y.L., Merino, M., Trepel, J., Zbar, B., Toro, J., et al. (2005). HIF overexpression correlates with biallelic loss of fumarate hydratase in renal cancer: novel role of fumarate in regulation of HIF stability. *Cancer Cell* 8, 143-153.

Ito, S., Shen, L., Dai, Q., Wu, S.C., Collins, L.B., Swenberg, J.A., He, C., and Zhang, Y. (2011). Tet proteins can convert 5-methylcytosine to 5-formylcytosine and 5-carboxylcytosine. *Science* 333, 1300-1303.

Ivan, M., Harris, A.L., Martelli, F., and Kulshreshtha, R. (2008). Hypoxia response and microRNAs: no longer two separate worlds. *J Cell Mol Med* 12, 1426-1431.

John, S., Sabo, P.J., Thurman, R.E., Sung, M.H., Biddie, S.C., Johnson, T.A., Hager, G.L., and Stamatoyannopoulos, J.A. (2011). Chromatin accessibility pre-determines glucocorticoid receptor binding patterns. *Nat Genet* 43, 264-268.

Johnson, A.B., Denko, N., and Barton, M.C. (2008). Hypoxia induces a novel signature of chromatin modifications and global repression of transcription. *Mutat Res* 640, 174-179.

Joshi, H.P., Subramanian, I.V., Schnettler, E.K., Ghosh, G., Rupaimoole, R., Evans, C., Saluja, M., Jing, Y., Cristina, I., Roy, S., et al. (2014). Dynamin 2 along with microRNA-199a reciprocally regulate hypoxia-inducible factors and ovarian cancer metastasis. *Proc Natl Acad Sci U S A* 111, 5331-5336.

Jung, K.O., Youn, H., Lee, C.H., Kang, K.W., and Chung, J.K. (2017). Visualization of exosome-mediated miR-210 transfer from hypoxic tumor cells. *Oncotarget* 8, 9899-9910.

Kaelin, W.G., Jr., and Ratcliffe, P.J. (2008). Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Mol Cell* 30, 393-402.

Kalousi, A., Mylonis, I., Politou, A.S., Chachami, G., Paraskeva, E., and Simos, G. (2010). Casein kinase 1 regulates human hypoxia-inducible factor HIF-1. *J Cell Sci* 123, 2976-2986.

Kelly, T.J., Souza, A.L., Clish, C.B., and Puigserver, P. (2011). A hypoxia-induced positive feedback loop promotes hypoxia-inducible factor 1 $\alpha$  stability through miR-210 suppression of glycerol-3-phosphate dehydrogenase 1-like. *Mol Cell Biol* 31, 2696-2706.

Kenneth, N.S., Mudie, S., van Uden, P., and Rocha, S. (2009). SWI/SNF regulates the cellular response to hypoxia. *J Biol Chem* 284, 4123-4131.

Kim, H.W., Haider, H.K., Jiang, S., and Ashraf, M. (2009). Ischemic preconditioning augments survival of stem cells via miR-210 expression by targeting caspase-8-associated protein 2. *J Biol Chem* 284, 33161-33168.

Kim, J.H., Park, S.G., Song, S.Y., Kim, J.K., and Sung, J.H. (2013). Reactive oxygen species-responsive miR-210 regulates proliferation and migration of adipose-derived stem cells via PTPN2. *Cell Death Dis* 4, e588.

Kim, M.S., Kwon, H.J., Lee, Y.M., Baek, J.H., Jang, J.E., Lee, S.W., Moon, E.J., Kim, H.S., Lee, S.K., Chung, H.Y., et al. (2001). Histone deacetylases induce angiogenesis by negative regulation of tumor suppressor genes. *Nat Med* 7, 437-443.

King, H.W., Michael, M.Z., and Gleadle, J.M. (2012). Hypoxic enhancement of exosome release by breast cancer cells. *BMC Cancer* 12, 421.

Klein, B.J., Piao, L., Xi, Y., Rincon-Arango, H., Rothbart, S.B., Peng, D., Wen, H., Larson, C., Zhang, X., Zheng, X., et al. (2014). The histone-H3K4-specific demethylase KDM5B binds to its substrate and product through distinct PHD fingers. *Cell Rep* 6, 325-335.

Kong, X., Lin, Z., Liang, D., Fath, D., Sang, N., and Caro, J. (2006). Histone deacetylase inhibitors induce VHL and ubiquitin-independent proteasomal degradation of hypoxia-inducible factor 1 $\alpha$ . *Mol Cell Biol* 26, 2019-2028.

Koslowski, M., Luxemburger, U., Tureci, O., and Sahin, U. (2011). Tumor-associated CpG demethylation augments hypoxia-induced effects by positive autoregulation of HIF-1 $\alpha$ . *Oncogene* 30, 876-882.

Koyanagi, S., Kuramoto, Y., Nakagawa, H., Aramaki, H., Ohdo, S., Soeda, S., and Shimeno, H. (2003). A molecular mechanism regulating circadian expression of vascular endothelial growth factor in tumor cells. *Cancer Res* 63, 7277-7283.

Krieg, A.J., Rankin, E.B., Chan, D., Razorenova, O., Fernandez, S., and Giaccia, A.J. (2010). Regulation of the histone demethylase JMJD1A by hypoxia-inducible factor 1 alpha enhances hypoxic gene expression and tumor growth. *Mol Cell Biol* *30*, 344-353.

Kucharzewska, P., Christianson, H.C., Welch, J.E., Svensson, K.J., Fredlund, E., Ringner, M., Morgelin, M., Bourseau-Guilmain, E., Bengzon, J., and Belting, M. (2013). Exosomes reflect the hypoxic status of glioma cells and mediate hypoxia-dependent activation of vascular cells during tumor development. *Proc Natl Acad Sci U S A* *110*, 7312-7317.

Laukka, T., Mariani, C.J., Ithantola, T., Cao, J.Z., Hokkanen, J., Kaelin, W.G., Jr., Godley, L.A., and Koivunen, P. (2016). Fumarate and Succinate Regulate Expression of Hypoxia-inducible Genes via TET Enzymes. *J Biol Chem* *291*, 4256-4265.

Law, A.Y., Lai, K.P., Ip, C.K., Wong, A.S., Wagner, G.F., and Wong, C.K. (2008). Epigenetic and HIF-1 regulation of stanniocalcin-2 expression in human cancer cells. *Exp Cell Res* *314*, 1823-1830.

Ledaki, I., McIntyre, A., Wigfield, S., Buffa, F., McGowan, S., Baban, D., Li, J.L., and Harris, A.L. (2015). Carbonic anhydrase IX induction defines a heterogeneous cancer cell response to hypoxia and mediates stem cell-like properties and sensitivity to HDAC inhibition. *Oncotarget* *6*, 19413-19427.

Lee, H.Y., Choi, K., Oh, H., Park, Y.K., and Park, H. (2014). HIF-1-dependent induction of Jumonji domain-containing protein (JMJD) 3 under hypoxic conditions. *Mol Cells* *37*, 43-50.

Li, L., Li, C., Wang, S., Wang, Z., Jiang, J., Wang, W., Li, X., Chen, J., Liu, K., Li, C., et al. (2016a). Exosomes Derived from Hypoxic Oral Squamous Cell Carcinoma Cells Deliver miR-21 to Normoxic Cells to Elicit a Prometastatic Phenotype. *Cancer Res* *76*, 1770-1780.

Li, W., He, X., Xue, R., Zhang, Y., Zhang, X., Lu, J., Zhang, Z., and Xue, L. (2016b). Combined over-expression of the hypoxia-inducible factor 2alpha gene and its long non-coding RNA predicts unfavorable prognosis of patients with osteosarcoma. *Pathol Res Pract* *212*, 861-866.

Li, X., Liu, L., Yang, S., Song, N., Zhou, X., Gao, J., Yu, N., Shan, L., Wang, Q., Liang, J., et al. (2014). Histone demethylase KDM5B is a key regulator of genome stability. *Proc Natl Acad Sci U S A* *111*, 7096-7101.

Li, X., Wu, Y., Liu, A., and Tang, X. (2016c). Long non-coding RNA UCA1 enhances tamoxifen resistance in breast cancer cells through a miR-18a-HIF1alpha feedback regulatory loop. *Tumour Biol* *37*, 14733-14743.

Lin, A., Li, C., Xing, Z., Hu, Q., Liang, K., Han, L., Wang, C., Hawke, D.H., Wang, S., Zhang, Y., et al. (2016). The LINK-A lncRNA activates normoxic HIF1alpha signalling in triple-negative breast cancer. *Nat Cell Biol* *18*, 213-224.

Liu, C., Kelnar, K., Liu, B., Chen, X., Calhoun-Davis, T., Li, H., Patrawala, L., Yan, H., Jeter, C., Honorio, S., et al. (2011a). The microRNA miR-34a inhibits prostate cancer stem cells and metastasis by directly repressing CD44. *Nat Med* *17*, 211-215.

Liu, Q., Liu, L., Zhao, Y., Zhang, J., Wang, D., Chen, J., He, Y., Wu, J., Zhang, Z., and Liu, Z. (2011b). Hypoxia induces genomic DNA demethylation through the activation of HIF-1alpha and transcriptional upregulation of MAT2A in hepatoma cells. *Mol Cancer Ther* *10*, 1113-1123.

Lu, Y., Chu, A., Turker, M.S., and Glazer, P.M. (2011). Hypoxia-induced epigenetic regulation and silencing of the BRCA1 promoter. *Mol Cell Biol* *31*, 3339-3350.

Luo, W., Chang, R., Zhong, J., Pandey, A., and Semenza, G.L. (2012). Histone demethylase JMJD2C is a coactivator for hypoxia-inducible factor 1 that is required for breast cancer progression. *Proc Natl Acad Sci U S A* *109*, E3367-3376.

Matouk, I.J., Halle, D., Raveh, E., Gilon, M., Sorin, V., and Hochberg, A. (2016). The role of the oncofetal H19 lncRNA in tumor metastasis: orchestrating the EMT-MET decision. *Oncotarget* *7*, 3748-3765.

Mayo, J.N., and Bearden, S.E. (2015). Driving the Hypoxia-Inducible Pathway in Human Pericytes Promotes Vascular Density in an Exosome-Dependent Manner. *Microcirculation* *22*, 711-723.

McCarty, G., and Loeb, D.M. (2015). Hypoxia-sensitive epigenetic regulation of an antisense-oriented lncRNA controls WT1 expression in myeloid leukemia cells. *PLoS One* *10*, e0119837.

Melvin, A., and Rocha, S. (2012). Chromatin as an oxygen sensor and active player in the hypoxia response. *Cell Signal* *24*, 35-43.

Michalik, K.M., You, X., Manavski, Y., Doddaballapur, A., Zornig, M., Braun, T., John, D., Ponomareva, Y., Chen, W., Uchida, S., et al. (2014). Long noncoding RNA MALAT1 regulates endothelial cell function and vessel growth. *Circ Res* *114*, 1389-1397.

Mimura, I., Nangaku, M., Kanki, Y., Tsutsumi, S., Inoue, T., Kohro, T., Yamamoto, S., Fujita, T., Shimamura, T., Suehiro, J., et al. (2012). Dynamic change of chromatin conformation in response to hypoxia enhances the expression of GLUT3 (SLC2A3) by cooperative interaction of hypoxia-inducible factor 1 and KDM3A. *Mol Cell Biol* *32*, 3018-3032.

Mineo, M., Ricklefs, F., Rooj, A.K., Lyons, S.M., Ivanov, P., Ansari, K.I., Nakano, I., Chiocca, E.A., Godlewski, J., and Bronisz, A. (2016). The Long Non-coding RNA HIF1A-AS2 Facilitates the Maintenance of Mesenchymal Glioblastoma Stem-like Cells in Hypoxic Niches. *Cell Rep* *15*, 2500-2509.

Murai, M., Toyota, M., Suzuki, H., Satoh, A., Sasaki, Y., Akino, K., Ueno, M., Takahashi, F., Kusano, M., Mita, H., et al. (2005). Aberrant methylation and silencing of the BNIP3 gene in colorectal and gastric cancer. *Clin Cancer Res* *11*, 1021-1027.

Nakamura, J., Kitajima, Y., Kai, K., Hashiguchi, K., Hiraki, M., Noshiro, H., and Miyazaki, K. (2011). Expression of hypoxic marker CA IX is regulated by site-specific DNA methylation and is associated with the histology of gastric cancer. *Am J Pathol* *178*, 515-524.

Nakazawa, M.S., Eisinger-Mathason, T.S., Sadri, N., Ochocki, J.D., Gade, T.P., Amin, R.K., and Simon, M.C. (2016). Epigenetic re-expression of HIF-2alpha suppresses soft tissue sarcoma growth. *Nat Commun* *7*, 10539.

Okami, J., Simeone, D.M., and Logsdon, C.D. (2004). Silencing of the hypoxia-inducible cell death protein BNIP3 in pancreatic cancer. *Cancer Res* *64*, 5338-5346.

Pal, A., Srivastava, T., Sharma, M.K., Mehndiratta, M., Das, P., Sinha, S., and Chattopadhyay, P. (2010). Aberrant methylation and associated transcriptional mobilization of Alu elements contributes to genomic instability in hypoxia. *J Cell Mol Med* *14*, 2646-2654.

Parolini, I., Federici, C., Raggi, C., Lugini, L., Palleschi, S., De Milito, A., Coscia, C., Iessi, E., Logozzi, M., Molinari, A., et al. (2009). Microenvironmental pH is a key factor for exosome traffic in tumor cells. *J Biol Chem* *284*, 34211-34222.

Peek, C.B., Levine, D.C., Cedernaes, J., Taguchi, A., Kobayashi, Y., Tsai, S.J., Bonar, N.A., McNulty, M.R., Ramsey, K.M., and Bass, J. (2017). Circadian Clock Interaction with HIF1alpha Mediates Oxygenic Metabolism and Anaerobic Glycolysis in Skeletal Muscle. *Cell Metab* *25*, 86-92.

Place, T.L., Fitzgerald, M.P., Venkataraman, S., Vorrink, S.U., Case, A.J., Teoh, M.L., and Domann, F.E. (2011). Aberrant promoter CpG methylation is a mechanism for impaired PHD3 expression in a diverse set of malignant cells. *PLoS One* *6*, e14617.

Platt, J.L., Salama, R., Smythies, J., Choudhry, H., Davies, J.O., Hughes, J.R., Ratcliffe, P.J., and Mole, D.R. (2016). Capture-C reveals preformed chromatin interactions between HIF-binding sites and distant promoters. *EMBO Rep* *17*, 1410-1421.

Pollard, P.J., Loenarz, C., Mole, D.R., McDonough, M.A., Gleadle, J.M., Schofield, C.J., and Ratcliffe, P.J. (2008). Regulation of Jumonji-domain-containing histone demethylases by hypoxia-inducible factor (HIF)-1alpha. *Biochem J* *416*, 387-394.

Polytarchou, C., Iliopoulos, D., Hatzia Apostolou, M., Kottakis, F., Maroulakou, I., Struhl, K., and Tschlis, P.N. (2011). Akt2 regulates all Akt isoforms and promotes resistance to hypoxia through induction of miR-21 upon oxygen deprivation. *Cancer Res* *71*, 4720-4731.

Prickaerts, P., Adriaens, M.E., Beucken, T.V., Koch, E., Dubois, L., Dahlmans, V.E., Gits, C., Evelo, C.T., Chan-Seng-Yue, M., Wouters, B.G., et al. (2016). Hypoxia increases genome-wide bivalent epigenetic marking by specific gain of H3K27me3. *Epigenetics Chromatin* *9*, 46.

Qin, Q., Furong, W., and Baosheng, L. (2014). Multiple functions of hypoxia-regulated miR-210 in cancer. *J Exp Clin Cancer Res* *33*, 50.

Ramteke, A., Ting, H., Agarwal, C., Mateen, S., Somasagara, R., Hussain, A., Graner, M., Frederick, B., Agarwal, R., and Deep, G. (2015). Exosomes secreted under hypoxia enhance invasiveness and stemness of prostate cancer cells by targeting adherens junction molecules. *Mol Carcinog* 54, 554-565.

Rankin, E.B., and Giaccia, A.J. (2016). Hypoxic control of metastasis. *Science* 352, 175-180.

Ratcliffe, P.J. (2013). Oxygen sensing and hypoxia signalling pathways in animals: the implications of physiology for cancer. *J Physiol* 591, 2027-2042.

Riches, A., Campbell, E., Borger, E., and Powis, S. (2014). Regulation of exosome release from mammary epithelial and breast cancer cells - a new regulatory pathway. *Eur J Cancer* 50, 1025-1034.

Rokavec, M., Oner, M.G., Li, H., Jackstadt, R., Jiang, L., Lodygin, D., Kaller, M., Horst, D., Ziegler, P.K., Schwitalla, S., et al. (2014). IL-6R/STAT3/miR-34a feedback loop promotes EMT-mediated colorectal cancer invasion and metastasis. *J Clin Invest* 124, 1853-1867.

Rong, L., Li, R., Li, S., and Luo, R. (2016). Immunosuppression of breast cancer cells mediated by transforming growth factor-beta in exosomes from cancer cells. *Oncol Lett* 11, 500-504.

Rupaimoole, R., Calin, G.A., Lopez-Berestein, G., and Sood, A.K. (2016a). miRNA Deregulation in Cancer Cells and the Tumor Microenvironment. *Cancer Discov* 6, 235-246.

Rupaimoole, R., Ivan, C., Yang, D., Gharpure, K.M., Wu, S.Y., Pecot, C.V., Previs, R.A., Nagaraja, A.S., Armaiz-Pena, G.N., McGuire, M., et al. (2016b). Hypoxia-upregulated microRNA-630 targets Dicer, leading to increased tumor progression. *Oncogene* 35, 4312-4320.

Sahar, S., Zocchi, L., Kinoshita, C., Borrelli, E., and Sassone-Corsi, P. (2010). Regulation of BMAL1 protein stability and circadian function by GSK3beta-mediated phosphorylation. *PLoS One* 5, e8561.

Salminen, A., Kaarniranta, K., and Kauppinen, A. (2016). Hypoxia-Inducible Histone Lysine Demethylases: Impact on the Aging Process and Age-Related Diseases. *Aging Dis* 7, 180-200.

Sanchez-Vega, F., Gotea, V., Petrykowska, H.M., Margolin, G., Krivak, T.C., DeLoia, J.A., Bell, D.W., and Elnitski, L. (2013). Recurrent patterns of DNA methylation in the ZNF154, CASP8, and VHL promoters across a wide spectrum of human solid epithelial tumors and cancer cell lines. *Epigenetics* 8, 1355-1372.

Sato, F., Bhawal, U.K., Yoshimura, T., and Muragaki, Y. (2016). DEC1 and DEC2 Crosstalk between Circadian Rhythm and Tumor Progression. *J Cancer* 7, 153-159.

Schlaepfer, I.R., Nambiar, D.K., Ramteke, A., Kumar, R., Dhar, D., Agarwal, C., Bergman, B., Graner, M., Maroni, P., Singh, R.P., et al. (2015). Hypoxia induces triglycerides accumulation in prostate cancer cells and extracellular vesicles supporting growth and invasiveness following reoxygenation. *Oncotarget* 6, 22836-22856.

Schmitt, A.M., Schmid, S., Rudolph, T., Anlauf, M., Prinz, C., Kloppel, G., Moch, H., Heitz, P.U., Komminoth, P., and Perren, A. (2009). VHL inactivation is an important pathway for the development of malignant sporadic pancreatic endocrine tumors. *Endocr Relat Cancer* 16, 1219-1227.

Schodel, J., Oikonomopoulos, S., Ragoussis, J., Pugh, C.W., Ratcliffe, P.J., and Mole, D.R. (2011). High-resolution genome-wide mapping of HIF-binding sites by ChIP-seq. *Blood* 117, e207-217.

Schoepflin, Z.R., Shapiro, I.M., and Risbud, M.V. (2016). Class I and IIa HDACs Mediate HIF-1alpha Stability Through PHD2-Dependent Mechanism, While HDAC6, a Class IIb Member, Promotes HIF-1alpha Transcriptional Activity in Nucleus Pulposus Cells of the Intervertebral Disc. *J Bone Miner Res* 31, 1287-1299.

Selak, M.A., Armour, S.M., MacKenzie, E.D., Boulahbel, H., Watson, D.G., Mansfield, K.D., Pan, Y., Simon, M.C., Thompson, C.B., and Gottlieb, E. (2005). Succinate links TCA cycle dysfunction to oncogenesis by inhibiting HIF-alpha prolyl hydroxylase. *Cancer Cell* 7, 77-85.

Shahzad, S., Bertrand, K., Minhas, K., and Coomber, B.L. (2007). Induction of DNA hypomethylation by tumor hypoxia. *Epigenetics* 2, 119-125.

Shen, J., Xia, W., Khotskaya, Y.B., Huo, L., Nakanishi, K., Lim, S.O., Du, Y., Wang, Y., Chang, W.C., Chen, C.H., et al. (2013). EGFR modulates microRNA maturation in response to hypoxia through phosphorylation of AGO2. *Nature* 497, 383-387.

Shi, X.F., Wang, H., Kong, F.X., Xu, Q.Q., Xiao, F.J., Yang, Y.F., Ge, R.L., and Wang, L.S. (2017). Exosomal miR-486 regulates hypoxia-induced erythroid differentiation of erythroleukemia cells through targeting Sirt1. *Exp Cell Res* 351, 74-81.

Shih, J.W., Chiang, W.F., Wu, A.T.H., Wu, M.H., Wang, L.Y., Yu, Y.L., Hung, Y.W., Wang, W.C., Chu, C.Y., Hung, C.L., et al. (2017). Long noncoding RNA LncHIFCAR/MIR31HG is a HIF-1 $\alpha$  co-activator driving oral cancer progression. *Nat Commun* 8, 15874.

Slee, R.B., Steiner, C.M., Herbert, B.S., Vance, G.H., Hickey, R.J., Schwarz, T., Christan, S., Radovich, M., Schneider, B.P., Schindelbauer, D., et al. (2012). Cancer-associated alteration of pericentromeric heterochromatin may contribute to chromosome instability. *Oncogene* 31, 3244-3253.

Stewart, G.D., Powles, T., Van Neste, C., Meynert, A., O'Mahony, F., Laird, A., Deforce, D., Van Nieuwerburgh, F., Trooskens, G., Van Criekinge, W., et al. (2016). Dynamic epigenetic changes to VHL occur with sunitinib in metastatic clear cell renal cancer. *Oncotarget* 7, 25241-25250.

Su, X., Li, G., and Liu, W. (2017). The Long Noncoding RNA Cancer Susceptibility Candidate 9 Promotes Nasopharyngeal Carcinogenesis via Stabilizing HIF1 $\alpha$ . *DNA Cell Biol* 36, 394-400.

Sun, Y.W., Chen, Y.F., Li, J., Huo, Y.M., Liu, D.J., Hua, R., Zhang, J.F., Liu, W., Yang, J.Y., Fu, X.L., et al. (2014). A novel long non-coding RNA ENST00000480739 suppresses tumour cell invasion by regulating OS-9 and HIF-1 $\alpha$  in pancreatic ductal adenocarcinoma. *Br J Cancer* 111, 2131-2141.

Tadokoro, H., Umezumi, T., Ohyashiki, K., Hirano, T., and Ohyashiki, J.H. (2013). Exosomes derived from hypoxic leukemia cells enhance tube formation in endothelial cells. *J Biol Chem* 288, 34343-34351.

Tahiliani, M., Koh, K.P., Shen, Y., Pastor, W.A., Bandukwala, H., Brudno, Y., Agarwal, S., Iyer, L.M., Liu, D.R., Aravind, L., et al. (2009). Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* 324, 930-935.

Takahashi, J.S. (2017). Transcriptional architecture of the mammalian circadian clock. *Nat Rev Genet* 18, 164-179.

Takahashi, K., Yan, I.K., Haga, H., and Patel, T. (2014). Modulation of hypoxia-signaling pathways by extracellular linc-RoR. *J Cell Sci* 127, 1585-1594.

Tanimoto, K., Tsuchihara, K., Kanai, A., Arauchi, T., Esumi, H., Suzuki, Y., and Sugano, S. (2010). Genome-wide identification and annotation of HIF-1 $\alpha$  binding sites in two cell lines using massively parallel sequencing. *Hugo J* 4, 35-48.

Tao, H., Wang, M.M., Zhang, M., Zhang, S.P., Wang, C.H., Yuan, W.J., Sun, T., He, L.J., and Hu, Q.K. (2016). MiR-126 Suppresses the Glucose-Stimulated Proliferation via IRS-2 in INS-1 beta Cells. *PLoS One* 11, e0149954.

Tasharofi, B., Soudyab, M., Nikpayam, E., Iranpour, M., Mirfakhraie, R., Sarrafzadeh, S., Geranpayeh, L., Azargashb, E., Sayad, A., and Ghafouri-Fard, S. (2016). Comparative expression analysis of hypoxia-inducible factor- $\alpha$  and its natural occurring antisense in breast cancer tissues and adjacent noncancerous tissues. *Cell Biochem Funct* 34, 572-578.

Tausendschon, M., Dehne, N., and Brune, B. (2011). Hypoxia causes epigenetic gene regulation in macrophages by attenuating Jumonji histone demethylase activity. *Cytokine* 53, 256-262.

Thienpont, B., Steinbacher, J., Zhao, H., D'Anna, F., Kuchnio, A., Ploumaki, A., Ghesquiere, B., Van Dyck, L., Boeckx, B., Schoonjans, L., et al. (2016). Tumour hypoxia causes DNA hypermethylation by reducing TET activity. *Nature* 537, 63-68.

Tomasetti, M., Nocchi, L., Staffolani, S., Manzella, N., Amati, M., Goodwin, J., Kluckova, K., Nguyen, M., Strafella, E., Bajzikova, M., et al. (2014). MicroRNA-126 suppresses mesothelioma malignancy by targeting IRS1 and interfering with the mitochondrial function. *Antioxid Redox Signal* 21, 2109-2125.

Tsai, Y.P., Chen, H.F., Chen, S.Y., Cheng, W.C., Wang, H.W., Shen, Z.J., Song, C., Teng, S.C., He, C., and Wu, K.J. (2014). TET1 regulates hypoxia-induced epithelial-mesenchymal transition by acting as a co-activator. *Genome Biol* 15, 513.

Uchida, T., Rossignol, F., Matthay, M.A., Mounier, R., Couette, S., Clottes, E., and Clerici, C. (2004). Prolonged hypoxia differentially regulates hypoxia-inducible factor (HIF)-1 $\alpha$  and HIF-2 $\alpha$  expression in lung epithelial cells: implication of natural antisense HIF-1 $\alpha$ . *J Biol Chem* 279, 14871-14878.

Umezū, T., Tadokoro, H., Azuma, K., Yoshizawa, S., Ohyashiki, K., and Ohyashiki, J.H. (2014). Exosomal miR-135b shed from hypoxic multiple myeloma cells enhances angiogenesis by targeting factor-inhibiting HIF-1. *Blood* *124*, 3748-3757.

van den Beucken, T., Koch, E., Chu, K., Rupaimoole, R., Prickaerts, P., Adriaens, M., Voncken, J.W., Harris, A.L., Buffa, F.M., Haider, S., et al. (2014). Hypoxia promotes stem cell phenotypes and poor prognosis through epigenetic regulation of DICER. *Nat Commun* *5*, 5203.

Vasagiri, N., and Kutala, V.K. (2014). Structure, function, and epigenetic regulation of BNIP3: a pathophysiological relevance. *Mol Biol Rep* *41*, 7705-7714.

Vinas, J.L., Burger, D., Zimpelmann, J., Haneef, R., Knoll, W., Campbell, P., Gutsol, A., Carter, A., Allan, D.S., and Burns, K.D. (2016). Transfer of microRNA-486-5p from human endothelial colony forming cell-derived exosomes reduces ischemic kidney injury. *Kidney Int* *90*, 1238-1250.

Voellenkle, C., Rooij, J., Guffanti, A., Brini, E., Fasanaro, P., Isaia, E., Croft, L., David, M., Capogrossi, M.C., Moles, A., et al. (2012). Deep-sequencing of endothelial cells exposed to hypoxia reveals the complexity of known and novel microRNAs. *RNA* *18*, 472-484.

Wan, W., Peng, K., Li, M., Qin, L., Tong, Z., Yan, J., Shen, B., and Yu, C. (2017). Histone demethylase JMJD1A promotes urinary bladder cancer progression by enhancing glycolysis through coactivation of hypoxia inducible factor 1 $\alpha$ . *Oncogene*.

Wang, J., Wang, Y., and Lu, L. (2012). De-SUMOylation of CCCTC binding factor (CTCF) in hypoxic stress-induced human corneal epithelial cells. *J Biol Chem* *287*, 12469-12479.

Wang, T., Gilkes, D.M., Takano, N., Xiang, L., Luo, W., Bishop, C.J., Chaturvedi, P., Green, J.J., and Semenza, G.L. (2014). Hypoxia-inducible factors and RAB22A mediate formation of microvesicles that stimulate breast cancer invasion and metastasis. *Proc Natl Acad Sci U S A* *111*, E3234-3242.

Wang, Y., Yao, J., Meng, H., Yu, Z., Wang, Z., Yuan, X., Chen, H., and Wang, A. (2015). A novel long non-coding RNA, hypoxia-inducible factor-2 $\alpha$  promoter upstream transcript, functions as an inhibitor of osteosarcoma stem cells in vitro. *Mol Med Rep* *11*, 2534-2540.

Watson, C.J., Collier, P., Tea, I., Neary, R., Watson, J.A., Robinson, C., Phelan, D., Ledwidge, M.T., McDonald, K.M., McCann, A., et al. (2014). Hypoxia-induced epigenetic modifications are associated with cardiac tissue fibrosis and the development of a myofibroblast-like phenotype. *Hum Mol Genet* *23*, 2176-2188.

Watson, J.A., Watson, C.J., McCrohan, A.M., Woodfine, K., Tosetto, M., McDaid, J., Gallagher, E., Betts, D., Baugh, J., O'Sullivan, J., et al. (2009). Generation of an epigenetic signature by chronic hypoxia in prostate cells. *Hum Mol Genet* *18*, 3594-3604.

Wenger, R.H., Kvietikova, I., Rolfs, A., Camenisch, G., and Gassmann, M. (1998). Oxygen-regulated erythropoietin gene expression is dependent on a CpG methylation-free hypoxia-inducible factor-1 DNA-binding site. *Eur J Biochem* *253*, 771-777.

Wu, F., Huang, W., and Wang, X. (2015a). microRNA-18a regulates gastric carcinoma cell apoptosis and invasion by suppressing hypoxia-inducible factor-1 $\alpha$  expression. *Exp Ther Med* *10*, 717-722.

Wu, M.Z., Chen, S.F., Nieh, S., Benner, C., Ger, L.P., Jan, C.I., Ma, L., Chen, C.H., Hishida, T., Chang, H.T., et al. (2015b). Hypoxia Drives Breast Tumor Malignancy through a TET-TNF $\alpha$ -p38-MAPK Signaling Axis. *Cancer Res* *75*, 3912-3924.

Wu, M.Z., Tsai, Y.P., Yang, M.H., Huang, C.H., Chang, S.Y., Chang, C.C., Teng, S.C., and Wu, K.J. (2011). Interplay between HDAC3 and WDR5 is essential for hypoxia-induced epithelial-mesenchymal transition. *Mol Cell* *43*, 811-822.

Wu, Y., Tang, D., Liu, N., Xiong, W., Huang, H., Li, Y., Ma, Z., Zhao, H., Chen, P., Qi, X., et al. (2017). Reciprocal Regulation between the Circadian Clock and Hypoxia Signaling at the Genome Level in Mammals. *Cell Metab* *25*, 73-85.

Xia, X., Lemieux, M.E., Li, W., Carroll, J.S., Brown, M., Liu, X.S., and Kung, A.L. (2009). Integrative analysis of HIF binding and transactivation reveals its role in maintaining histone methylation homeostasis. *Proc Natl Acad Sci U S A* *106*, 4260-4265.

Xu, H., Zhao, L., Fang, Q., Sun, J., Zhang, S., Zhan, C., Liu, S., and Zhang, Y. (2014). MiR-338-3p inhibits hepatocarcinoma cells and sensitizes these cells to sorafenib by targeting hypoxia-induced factor 1alpha. *PLoS One* 9, e115565.

Xue, M., Chen, W., Xiang, A., Wang, R., Chen, H., Pan, J., Pang, H., An, H., Wang, X., Hou, H., et al. (2017). Hypoxic exosomes facilitate bladder tumor growth and development through transferring long non-coding RNA-UCA1. *Mol Cancer* 16, 143.

Xue, M., Li, X., Li, Z., and Chen, W. (2014). Urothelial carcinoma associated 1 is a hypoxia-inducible factor-1alpha-targeted long noncoding RNA that enhances hypoxic bladder cancer cell proliferation, migration, and invasion. *Tumour Biol* 35, 6901-6912.

Yang, F., Huo, X.S., Yuan, S.X., Zhang, L., Zhou, W.P., Wang, F., and Sun, S.H. (2013). Repression of the long noncoding RNA-LET by histone deacetylase 3 contributes to hypoxia-mediated metastasis. *Mol Cell* 49, 1083-1096.

Yang, F., Zhang, H., Mei, Y., and Wu, M. (2014). Reciprocal regulation of HIF-1alpha and lincRNA-p21 modulates the Warburg effect. *Mol Cell* 53, 88-100.

Yang, W., Sun, T., Cao, J., Liu, F., Tian, Y., and Zhu, W. (2012). Downregulation of miR-210 expression inhibits proliferation, induces apoptosis and enhances radiosensitivity in hypoxic human hepatoma cells in vitro. *Exp Cell Res* 318, 944-954.

Ye, S.B., Zhang, H., Cai, T.T., Liu, Y.N., Ni, J.J., He, J., Peng, J.Y., Chen, Q.Y., Mo, H.Y., Jun, C., et al. (2016). Exosomal miR-24-3p impedes T-cell function by targeting FGF11 and serves as a potential prognostic biomarker for nasopharyngeal carcinoma. *J Pathol* 240, 329-340.

Yeh, Y.M., Chuang, C.M., Chao, K.C., and Wang, L.H. (2013). MicroRNA-138 suppresses ovarian cancer cell invasion and metastasis by targeting SOX4 and HIF-1alpha. *Int J Cancer* 133, 867-878.

Ying, Q., Liang, L., Guo, W., Zha, R., Tian, Q., Huang, S., Yao, J., Ding, J., Bao, M., Ge, C., et al. (2011). Hypoxia-inducible microRNA-210 augments the metastatic potential of tumor cells by targeting vacuole membrane protein 1 in hepatocellular carcinoma. *Hepatology* 54, 2064-2075.

Yoo, Y.G., Kong, G., and Lee, M.O. (2006). Metastasis-associated protein 1 enhances stability of hypoxia-inducible factor-1alpha protein by recruiting histone deacetylase 1. *EMBO J* 25, 1231-1241.

Young, L.C., and Hendzel, M.J. (2013). The oncogenic potential of Jumonji D2 (JMJD2/KDM4) histone demethylase overexpression. *Biochem Cell Biol* 91, 369-377.

Yu, C., Yang, S.L., Fang, X., Jiang, J.X., Sun, C.Y., and Huang, T. (2015). Hypoxia disrupts the expression levels of circadian rhythm genes in hepatocellular carcinoma. *Mol Med Rep* 11, 4002-4008.

Zhai, W., Sun, Y., Jiang, M., Wang, M., Gasiewicz, T.A., Zheng, J., and Chang, C. (2016). Differential regulation of LncRNA-SARCC suppresses VHL-mutant RCC cell proliferation yet promotes VHL-normal RCC cell proliferation via modulating androgen receptor/HIF-2alpha/C-MYC axis under hypoxia. *Oncogene* 35, 4866-4880.

Zhang, J., Lu, S., Zhou, Y., Meng, K., Chen, Z., Cui, Y., Shi, Y., Wang, T., and He, Q.Y. (2017). Motile hepatocellular carcinoma cells preferentially secrete sugar metabolism regulatory proteins via exosomes. *Proteomics*.

Zhao, H., Yang, L., Baddour, J., Achreja, A., Bernard, V., Moss, T., Marini, J.C., Tudawe, T., Seviour, E.G., San Lucas, F.A., et al. (2016). Tumor microenvironment derived exosomes pleiotropically modulate cancer cell metabolism. *Elife* 5, e10250.

Zhao, R., Sun, F., Bei, X., Wang, X., Zhu, Y., Jiang, C., Zhao, F., Han, B., and Xia, S. (2017). Upregulation of the long non-coding RNA FALEC promotes proliferation and migration of prostate cancer cell lines and predicts prognosis of PCa patients. *Prostate* 77, 1107-1117.

Zhen, S., Hua, L., Liu, Y.H., Sun, X.M., Jiang, M.M., Chen, W., Zhao, L., and Li, X. (2017). Inhibition of long non-coding RNA UCA1 by CRISPR/Cas9 attenuated malignant phenotypes of bladder cancer. *Oncotarget* 8, 9634-9646.

Zhou, C., Huang, C., Wang, J., Huang, H., Li, J., Xie, Q., Liu, Y., Zhu, J., Li, Y., Zhang, D., et al. (2017). LncRNA MEG3 downregulation mediated by DNMT3b contributes to nickel malignant transformation of human bronchial epithelial cells via modulating PHLPP1 transcription and HIF-1alpha translation. *Oncogene*.

Zhou, C., Ye, L., Jiang, C., Bai, J., Chi, Y., and Zhang, H. (2015). Long noncoding RNA HOTAIR, a hypoxia-inducible factor-1alpha activated driver of malignancy, enhances hypoxic cancer cell proliferation, migration, and invasion in non-small cell lung cancer. *Tumour Biol* 36, 9179-9188.

Zhou, X., Gao, Q., Wang, J., Zhang, X., Liu, K., and Duan, Z. (2014). Linc-RNA-RoR acts as a "sponge" against mediation of the differentiation of endometrial cancer stem cells by microRNA-145. *Gynecol Oncol* 133, 333-339.

Zhou, X., Sun, H., Chen, H., Zavadil, J., Kluz, T., Arita, A., and Costa, M. (2010). Hypoxia induces trimethylated H3 lysine 4 by inhibition of JARID1A demethylase. *Cancer Res* 70, 4214-4221.

Zhu, G., Wang, S., Chen, J., Wang, Z., Liang, X., Wang, X., Jiang, J., Lang, J., and Li, L. (2017). Long noncoding RNA HAS2-AS1 mediates hypoxia-induced invasiveness of oral squamous cell carcinoma. *Mol Carcinog*.

Zhu, Z., Gao, X., He, Y., Zhao, H., Yu, Q., Jiang, D., Zhang, P., Ma, X., Huang, H., Dong, D., et al. (2012). An insertion/deletion polymorphism within RERT-lncRNA modulates hepatocellular carcinoma risk. *Cancer Res* 72, 6163-6172.

**Table 1. HIF dependent histone lysine demethylases:**

Histone lysine Demethylases (KDMs)	Other symbols	Target	Types cells	References
KDM3A	JMJD1A JHDM2A	H3K9me1 H3K9me2	liver cancer, glioblastoma, clear-cell renal cell carcinoma, osteosarcoma, breast cancer and cervical carcinoma	(Krieg et al., 2010; Pollard et al., 2008; Xia et al., 2009)
KDM2B	JHDM1	H3K4me3 H3K36me1 H3K26me2	liver cancer, glioblastoma	(Xia et al., 2009)
KDM4B	JMJD2B	H3K9me2 H3K9me3 H3K36me2 H3K36me3	liver cancer, glioblastoma, clear-cell renal cell carcinoma, osteosarcoma, breast cancer and cervical carcinoma, colon cancer, epithelial cells	(Beyer et al., 2008; Fu et al., 2012; Krieg et al., 2010; Pollard et al., 2008; Xia et al., 2009)
KDM5B	JAR1D1B	H3K4me2 H3K4me3	liver cancer, glioblastoma, clear-cell renal cell carcinoma	(Krieg et al., 2010; Xia et al., 2009)
KDM6B	JMJD3	H3K27me2 H3K27me3	liver cancer, glioblastoma, adipose tissue	(Lee et al., 2014; Xia et al., 2009)
KDM4C	JMJD2C	H3K9me2 H3K9me3 H3K36me2 H3K36me3	liver cancer, glioblastoma, osteosarcoma, breast cancer and cervical carcinoma, epithelial cells	(Beyer et al., 2008; Pollard et al., 2008; Xia et al., 2009)

**Table 2. LncRNAs regulate HIF pathways in cancer:**





lncRNA	Regulation in hypoxia	HIF dependence	Cancer types	Impact on HIF pathway	Reference
lncRNA-LET	Downregulated		Gallbladder cancer, squamous-cell lung cancer, hepatocellular carcinoma and colorectal cancers	<ul style="list-style-type: none"> <li>Regulates HIF1<math>\alpha</math> level</li> </ul>	(Yang et al., 2013)
HIF2PUT			Osteosarcoma	<ul style="list-style-type: none"> <li>Upregulates HIF2<math>\alpha</math></li> </ul>	(Li et al., 2016b)
aHIF-1 $\alpha$	Upregulated	HIF-1 $\alpha$	Multiple cancer types	<ul style="list-style-type: none"> <li>Downregulates HIF1<math>\alpha</math></li> </ul>	(Uchida et al., 2004)
Linc-ROR	Upregulated		Hepatocellular carcinoma	<ul style="list-style-type: none"> <li>Upregulates HIF1<math>\alpha</math></li> </ul>	(Takahashi et al., 2014)
lincRNA-p21	Upregulated	HIF-1 $\alpha$	Cervical cancer, breast cancer	<ul style="list-style-type: none"> <li>Upregulates HIF1<math>\alpha</math></li> <li>Enhances glycolysis</li> </ul>	(Yang et al., 2014)
RERT			Hepatocellular carcinoma	<ul style="list-style-type: none"> <li>Downregulates HIF</li> </ul>	(Sun et al., 2014)
ENST00000480739			Pancreatic cancer	<ul style="list-style-type: none"> <li>Downregulates HIF-1<math>\alpha</math></li> </ul>	(Sun et al., 2014)
PVT1			Gastric cancer	<ul style="list-style-type: none"> <li>Downregulates HIF-1<math>\alpha</math></li> </ul>	(Huang et al., 2017)
MEG3			Lung cancer	<ul style="list-style-type: none"> <li>Upregulate HIF-1<math>\alpha</math> protein translation</li> </ul>	(Zhou et al., 2017)
UCA1	Upregulated	HIF-1 $\alpha$	Breast cancer	<ul style="list-style-type: none"> <li>Upregulate HIF-1<math>\alpha</math></li> </ul>	(Li et al., 2016c)
LncRNA-SARCC	Downregulated in RCC VHL wild-type	HIF-2 $\alpha$	Renal cell carcinoma	<ul style="list-style-type: none"> <li>Upregulate HIF-2<math>\alpha</math></li> </ul>	(Zhai et al., 2016)
CASC9			Nasopharyngeal carcinoma	<ul style="list-style-type: none"> <li>Upregulate HIF-1<math>\alpha</math></li> </ul>	(Su et al., 2017)
LncHIFCAR	Upregulated		Oral carcinoma	<ul style="list-style-type: none"> <li>Upregulate HIF-1<math>\alpha</math></li> </ul>	(Shih et al., 2017)
LINK-A			Triple-negative breast cancer	<ul style="list-style-type: none"> <li>Upregulate HIF-1<math>\alpha</math></li> </ul>	(Lin et al., 2016)


**Table 3. Secreted molecules in exosomes under hypoxia:**

Molecules	Type	Cancer	HIF dependent	Reference
HIF-1 $\alpha$	Transcription Factor	Nasopharyngeal carcinoma		(Aga et al., 2014)
miR-210	Non-coding RNA	Breast cancer leukaemia		(Jung et al., 2017; King et al., 2012)
miR-486	Non-coding RNA	Erythroleukemia, Endothelial cells		(Shi et al., 2017; Vinas et al., 2016)
miR-24-3p	Non-coding RNA	Nasopharyngeal carcinoma.		(Ye et al., 2016)
miR-135b	Non-coding RNA	Multiple myeloma		(Umezu et al., 2014)
miR-21	Non-coding RNA	Oral squamous cell carcinoma	HIF-1 $\alpha$ and HIF-2 $\alpha$	(Li et al., 2016a)
Myristic, Palmitic, Palmitoleic, Stearic, Oleic, Linoleic, Arachidonic	Fatty acid	Prostate cancer		(Schlaepfer et al., 2015)
TGF- $\beta$ , IL10		Breast cancer		(Rong et al., 2016)
Wnt4	mRNA and Protein	Colorectal cancer	HIF-1 $\alpha$	(Huang and Feng, 2016)
<b>Proteins:</b> MMP9, PTX3, IL8, PDGF-AB/AA, CD26, PAI1,CAV1  <b>mRNAs:</b> IGFBP3, NDRG1, LOX, ADM, IGFBP3, BNIP3, ID2, NDRG1, PLOD2, PAI1	mRNA and Proteins	Glioblastoma		(Kucharzewska et al., 2013)
Tetraspanins (CD81 and CD63), HSP70, HSP90 , MMP-2, MMP-9, TGF- $\beta$ 2, TNF1 $\alpha$ , IL6	Protein, cytokines, growth factors	Prostate cancer		(Ramteke et al., 2015)

**Figure 1. Examples of regulatory mechanisms between lncRNAs and HIFs:** **a)** HIF-2 $\alpha$  binds HRE at the promoter region of NEAT1 to induce its expression under hypoxia. Upregulated NEAT1 increases the formation of paraspeckle bodies. On the other hand, upon hypoxia the editing of RNA is also increased. Hyper-edited RNA molecules (in particular A to I editing) bind to paraspeckle bodies and retained in the nucleus, thus inhibit their translation. **b)** Hypoxia downregulates lncRNA-LET expression through hypoxia-induced HDAC3, a HIF-1 $\alpha$  dependent gene. Hypoxia-induced HDAC3 reduces the histone H3 and H4 acetylation-mediated modulation at the promoter region of lncRNA-LET which consequently suppress its expression. Decrease lncRNA-LET expression increases the abundance of NF90 protein hence upregulates HIF-1 $\alpha$  level. **c)** Hypoxia induces lincRNA-P21 expression in a HIF-1 $\alpha$  dependent manner. Hypoxia induced lincRNA-P21 binds to VHL and HIF-1 $\alpha$  that causes disassociation of VHL and HIF-1 $\alpha$ , therefore inhibiting proteasome-dependent degradation of HIF-1 $\alpha$  under hypoxia. lincRNA-p21 is modulate the Warburg effect through regulating expression of a number of glycolysis related genes. **d)** Signal from Heparin-binding EGF (HB-EGF) triggers 'EGFR:GPNMB' heterodimerization. This lead to the recruitment of LINK-A which interacts with both BRK and LRRK2 to facilitate phosphorylation of HIF1 $\alpha$  and promotes glycolysis reprogramming in TNBC.

**Figure 2. Bidirectional interactions between circadian clock and HIF pathways:** Under hypoxia, HIF-1 $\alpha$  heterodimerize with HIF-1 $\beta$  in nucleus and binds to HREs in the promoters of hypoxia-induced genes. Upon recruitment of CBP/P300 (co-activator), HIF-1 $\alpha$ :HIF-1 $\beta$  complex drive the transcription of target genes such as glycolysis related genes and some circadian clock genes such PERs, CRYs and DECs. On the other hand, BMAL1 heterodimerize with CLOCK to form a BMAL1:CLOCK complex which binds at E-box containing motifs and transactivate the clock-controlled genes such as PERs, CRYs, REV-ERB, RORs, including HIF-1 $\alpha$ . The CBP/P300 act as co-activator to induce the activation. There are number of interconnected feedback loops between HIF and circadian clock pathways that regulate the circadian transcription. The key feedback is a native loop of PER and CRY (transcriptionally activated by HIF and by BMAL1:CLOCK) lead to the repression of CLOCK:BMAL1 mediated transcription. The other feedback loop is by ROR which activates BMAL1:CLOCK by binding the ROR-elements (RORE). In addition, ROR physically interact with HIF-1 $\alpha$  hence, inducing the stabilization and transcriptional activation of HIF-1 $\alpha$ . The REV-ERB act as a repressor for BMAL1 transcription by competing at the same ROR sequence. HIF-1 $\alpha$  dependent DECs also negatively feedback by suppressing BMAL1:CLOCK transactivation. These interactions and feedback loops provide strong evidence that HIF play an important role in modulating the circadian rhythm.

**(A) Hypoxia****(B) Hypoxia****(C) Hypoxia****(D) Normoxia**



## References

- Adamovich, Y., Ladeuix, B., Golik, M., Koeners, M.P., and Asher, G. (2017). Rhythmic Oxygen Levels Reset Circadian Clocks through HIF1alpha. *Cell Metab* 25, 93-101.
- Aga, M., Bentz, G.L., Raffa, S., Torrisi, M.R., Kondo, S., Wakisaka, N., Yoshizaki, T., Pagano, J.S., and Shackelford, J. (2014). Exosomal HIF1alpha supports invasive potential of nasopharyngeal carcinoma-associated LMP1-positive exosomes. *Oncogene* 33, 4613-4622.
- Agostino, P.V., Harrington, M.E., Ralph, M.R., and Golombek, D.A. (2009). Casein kinase-1-epsilon (CK1epsilon) and circadian photic responses in hamsters. *Chronobiol Int* 26, 126-133.
- Agrawal, R., Pandey, P., Jha, P., Dwivedi, V., Sarkar, C., and Kulshreshtha, R. (2014). Hypoxic signature of microRNAs in glioblastoma: insights from small RNA deep sequencing. *BMC Genomics* 15, 686.
- Bacon, A.L., Fox, S., Turley, H., and Harris, A.L. (2007). Selective silencing of the hypoxia-inducible factor 1 target gene BNIP3 by histone deacetylation and methylation in colorectal cancer. *Oncogene* 26, 132-141.
- Baek, J.H., Mahon, P.C., Oh, J., Kelly, B., Krishnamachary, B., Pearson, M., Chan, D.A., Giaccia, A.J., and Semenza, G.L. (2005). OS-9 interacts with hypoxia-inducible factor 1alpha and prolyl hydroxylases to promote oxygen-dependent degradation of HIF-1alpha. *Mol Cell* 17, 503-512.
- Bailey, K.M., Wojtkowiak, J.W., Hashim, A.I., and Gillies, R.J. (2012). Targeting the metabolic microenvironment of tumors. *Adv Pharmacol* 65, 63-107.
- Bandara, V., Michael, M.Z., and Gleadle, J.M. (2014). Hypoxia represses microRNA biogenesis proteins in breast cancer cells. *BMC Cancer* 14, 533.
- Bandara, V., Michael, M.Z., and Gleadle, J.M. (2017). MicroRNA biogenesis in hypoxia. *Microna*.
- Berchem, G., Noman, M.Z., Bosseler, M., Paggetti, J., Baconnais, S., Le Cam, E., Nanbakhsh, A., Moussay, E., Mami-Chouaib, F., Janji, B., et al. (2016). Hypoxic tumor-derived microvesicles negatively regulate NK cell function by a mechanism involving TGF-beta and miR23a transfer. *Oncoimmunology* 5, e1062968.
- Bertozzi, D., Iurlaro, R., Sordet, O., Marinello, J., Zaffaroni, N., and Capranico, G. (2011). Characterization of novel antisense HIF-1alpha transcripts in human cancers. *Cell Cycle* 10, 3189-3197.
- Beyer, S., Kristensen, M.M., Jensen, K.S., Johansen, J.V., and Staller, P. (2008). The histone demethylases JMJD1A and JMJD2B are transcriptional targets of hypoxia-inducible factor HIF. *J Biol Chem* 283, 36542-36552.
- Bhan, A., Deb, P., Shihabeddin, N., Ansari, K.I., Brotto, M., and Mandal, S.S. (2017). Histone methylase MLL1 coordinates with HIF and regulate lncRNA HOTAIR expression under hypoxia. *Gene* 629, 16-28.
- Blick, C., Ramachandran, A., McCormick, R., Wigfield, S., Cranston, D., Catto, J., and Harris, A.L. (2015). Identification of a hypoxia-regulated miRNA signature in bladder cancer and a role for miR-145 in hypoxia-dependent apoptosis. *Br J Cancer* 113, 634-644.
- Boeckel, J.N., Jae, N., Heumuller, A.W., Chen, W., Boon, R.A., Stellos, K., Zeiher, A.M., John, D., Uchida, S., and Dimmeler, S. (2015). Identification and Characterization of Hypoxia-Regulated Endothelial Circular RNA. *Circ Res* 117, 884-890.
- Bruning, U., Cerone, L., Neufeld, Z., Fitzpatrick, S.F., Cheong, A., Scholz, C.C., Simpson, D.A., Leonard, M.O., Tambuwala, M.M., Cummins, E.P., et al. (2011). MicroRNA-155 promotes resolution of hypoxia-inducible factor 1alpha activity during prolonged hypoxia. *Mol Cell Biol* 31, 4087-4096.
- Camps, C., Saini, H.K., Mole, D.R., Choudhry, H., Reczko, M., Guerra-Assuncao, J.A., Tian, Y.M., Buffa, F.M., Harris, A.L., Hatzigeorgiou, A.G., et al. (2014). Integrated analysis of microRNA and mRNA expression and association with HIF binding reveals the complexity of microRNA expression regulation under hypoxia. *Mol Cancer* 13, 28.

Cascella, B., and Mirica, L.M. (2012). Kinetic analysis of iron-dependent histone demethylases: alpha-ketoglutarate substrate inhibition and potential relevance to the regulation of histone demethylation in cancer cells. *Biochemistry* 51, 8699-8701.

Chan, S.Y., Zhang, Y.Y., Hemann, C., Mahoney, C.E., Zweier, J.L., and Loscalzo, J. (2009). MicroRNA-210 controls mitochondrial metabolism during hypoxia by repressing the iron-sulfur cluster assembly proteins ISCU1/2. *Cell Metab* 10, 273-284.

Chen, B., Zeng, X., He, Y., Wang, X., Liang, Z., Liu, J., Zhang, P., Zhu, H., Xu, N., and Liang, S. (2016). STC2 promotes the epithelial-mesenchymal transition of colorectal cancer cells through AKT-ERK signaling pathways. *Oncotarget* 7, 71400-71416.

Chen, Q., Chen, Y., Bian, C., Fujiki, R., and Yu, X. (2013). TET2 promotes histone O-GlcNAcylation during gene transcription. *Nature* 493, 561-564.

Chen, S., Yin, C., Lao, T., Liang, D., He, D., Wang, C., and Sang, N. (2015). AMPK-HDAC5 pathway facilitates nuclear accumulation of HIF-1alpha and functional activation of HIF-1 by deacetylating Hsp70 in the cytosol. *Cell Cycle* 14, 2520-2536.

Chilov, D., Hofer, T., Bauer, C., Wenger, R.H., and Gassmann, M. (2001). Hypoxia affects expression of circadian genes PER1 and CLOCK in mouse brain. *FASEB J* 15, 2613-2622.

Choudhry, H., Albukhari, A., Morotti, M., Haider, S., Moralli, D., Smythies, J., Schodel, J., Green, C.M., Camps, C., Buffa, F., et al. (2015). Tumor hypoxia induces nuclear paraspeckle formation through HIF-2alpha dependent transcriptional activation of NEAT1 leading to cancer cell survival. *Oncogene* 34, 4482-4490.

Choudhry, H., Harris, A.L., and McIntyre, A. (2016). The tumour hypoxia induced non-coding transcriptome. *Mol Aspects Med* 47-48, 35-53.

Choudhry, H., Schodel, J., Oikonomopoulos, S., Camps, C., Grampp, S., Harris, A.L., Ratcliffe, P.J., Ragoussis, J., and Mole, D.R. (2014). Extensive regulation of the non-coding transcriptome by hypoxia: role of HIF in releasing paused RNAPol2. *EMBO Rep* 15, 70-76.

da Motta, L.L., Ledaki, I., Purshouse, K., Haider, S., De Bastiani, M.A., Baban, D., Morotti, M., Steers, G., Wigfield, S., Bridges, E., et al. (2017). The BET inhibitor JQ1 selectively impairs tumour response to hypoxia and downregulates CA9 and angiogenesis in triple negative breast cancer. *Oncogene* 36, 122-132.

Dawson, M.A. (2017). The cancer epigenome: Concepts, challenges, and therapeutic opportunities. *Science* 355, 1147-1152.

de Jong, O.G., van Balkom, B.W., Gremmels, H., and Verhaar, M.C. (2016). Exosomes from hypoxic endothelial cells have increased collagen crosslinking activity through up-regulation of lysyl oxidase-like 2. *J Cell Mol Med* 20, 342-350.

Estaras, C., Fueyo, R., Akizu, N., Beltran, S., and Martinez-Balbas, M.A. (2013). RNA polymerase II progression through H3K27me3-enriched gene bodies requires JMJD3 histone demethylase. *Mol Biol Cell* 24, 351-360.

Fasanaro, P., D'Alessandra, Y., Di Stefano, V., Melchionna, R., Romani, S., Pompilio, G., Capogrossi, M.C., and Martelli, F. (2008). MicroRNA-210 modulates endothelial cell response to hypoxia and inhibits the receptor tyrosine kinase ligand Ephrin-A3. *J Biol Chem* 283, 15878-15883.

Favaro, E., Lord, S., Harris, A.L., and Buffa, F.M. (2011). Gene expression and hypoxia in breast cancer. *Genome Med* 3, 55.

Favaro, E., Ramachandran, A., McCormick, R., Gee, H., Blancher, C., Crosby, M., Devlin, C., Blick, C., Buffa, F., Li, J.L., et al. (2010). MicroRNA-210 regulates mitochondrial free radical response to hypoxia and krebs cycle in cancer cells by targeting iron sulfur cluster protein ISCU. *PLoS One* 5, e10345.

Feng, C.C., Lin, C.C., Lai, Y.P., Chen, T.S., Marthandam Asokan, S., Lin, J.Y., Lin, K.H., Viswanadha, V.P., Kuo, W.W., and Huang, C.Y. (2016). Hypoxia suppresses myocardial survival pathway through HIF-1alpha-IGFBP-3-dependent signaling and enhances cardiomyocyte autophagic and apoptotic effects mainly via FoxO3a-induced BNIP3 expression. *Growth Factors* 34, 73-86.

Ferdin, J., Nishida, N., Wu, X., Nicoloso, M.S., Shah, M.Y., Devlin, C., Ling, H., Shimizu, M., Kumar, K., Cortez, M.A., et al. (2013). HINCUTs in cancer: hypoxia-induced noncoding ultraconserved transcripts. *Cell Death Differ* 20, 1675-1687.

Fong, M.Y., Zhou, W., Liu, L., Alontaga, A.Y., Chandra, M., Ashby, J., Chow, A., O'Connor, S.T., Li, S., Chin, A.R., et al. (2015). Breast-cancer-secreted miR-122 reprograms glucose metabolism in premetastatic niche to promote metastasis. *Nat Cell Biol* 17, 183-194.

Fu, L., Chen, L., Yang, J., Ye, T., Chen, Y., and Fang, J. (2012). HIF-1 $\alpha$ -induced histone demethylase JMJD2B contributes to the malignant phenotype of colorectal cancer cells via an epigenetic mechanism. *Carcinogenesis* 33, 1664-1673.

Fu, L., and Kettner, N.M. (2013). The circadian clock in cancer development and therapy. *Prog Mol Biol Transl Sci* 119, 221-282.

Geng, H., Harvey, C.T., Pittsenbarger, J., Liu, Q., Beer, T.M., Xue, C., and Qian, D.Z. (2011). HDAC4 protein regulates HIF1 $\alpha$  protein lysine acetylation and cancer cell response to hypoxia. *J Biol Chem* 286, 38095-38102.

Ghorbel, M.T., Coulson, J.M., and Murphy, D. (2003). Cross-talk between hypoxic and circadian pathways: cooperative roles for hypoxia-inducible factor 1 $\alpha$  and CLOCK in transcriptional activation of the vasopressin gene. *Mol Cell Neurosci* 22, 396-404.

Ghosh, G., Subramanian, I.V., Adhikari, N., Zhang, X., Joshi, H.P., Basi, D., Chandrashekar, Y.S., Hall, J.L., Roy, S., Zeng, Y., et al. (2010). Hypoxia-induced microRNA-424 expression in human endothelial cells regulates HIF- $\alpha$  isoforms and promotes angiogenesis. *J Clin Invest* 120, 4141-4154.

Gomez-Maldonado, L., Tiana, M., Roche, O., Prado-Cabrero, A., Jensen, L., Fernandez-Barral, A., Guijarro-Munoz, I., Favaro, E., Moreno-Bueno, G., Sanz, L., et al. (2015). EFNA3 long noncoding RNAs induced by hypoxia promote metastatic dissemination. *Oncogene* 34, 2609-2620.

Gorospe, M., Tominaga, K., Wu, X., Fahling, M., and Ivan, M. (2011). Post-Transcriptional Control of the Hypoxic Response by RNA-Binding Proteins and MicroRNAs. *Front Mol Neurosci* 4, 7.

Grampp, S., Platt, J.L., Lauer, V., Salama, R., Kranz, F., Neumann, V.K., Wach, S., Stohr, C., Hartmann, A., Eckardt, K.U., et al. (2016). Genetic variation at the 8q24.21 renal cancer susceptibility locus affects HIF binding to a MYC enhancer. *Nat Commun* 7, 13183.

Grampp, S., Schmid, V., Salama, R., Lauer, V., Kranz, F., Platt, J.L., Smythies, J., Choudhry, H., Goppelt-Struebe, M., Ratcliffe, P.J., et al. (2017). Multiple renal cancer susceptibility polymorphisms modulate the HIF pathway. *PLoS Genet* 13, e1006872.

Haque, I., Banerjee, S., Mehta, S., De, A., Majumder, M., Mayo, M.S., Kambhampati, S., Campbell, D.R., and Banerjee, S.K. (2011). Cysteine-rich 61-connective tissue growth factor-nephroblastoma-overexpressed 5 (CCN5)/Wnt-1-induced signaling protein-2 (WISP-2) regulates microRNA-10b via hypoxia-inducible factor-1 $\alpha$ -TWIST signaling networks in human breast cancer cells. *J Biol Chem* 286, 43475-43485.

Hattori, M., Yokoyama, Y., Hattori, T., Motegi, S., Amano, H., Hatada, I., and Ishikawa, O. (2015). Global DNA hypomethylation and hypoxia-induced expression of the ten eleven translocation (TET) family, TET1, in scleroderma fibroblasts. *Exp Dermatol* 24, 841-846.

Hatzimichael, E., Dasoula, A., Shah, R., Syed, N., Papoudou-Bai, A., Coley, H.M., Dranitsaris, G., Bourantas, K.L., Stebbing, J., and Crook, T. (2010). The prolyl-hydroxylase EGLN3 and not EGLN1 is inactivated by methylation in plasma cell neoplasia. *Eur J Haematol* 84, 47-51.

Hogenesch, J.B., Gu, Y.Z., Jain, S., and Bradfield, C.A. (1998). The basic-helix-loop-helix-PAS orphan MOP3 forms transcriptionally active complexes with circadian and hypoxia factors. *Proc Natl Acad Sci U S A* 95, 5474-5479.

Huang, T., Liu, H.W., Chen, J.Q., Wang, S.H., Hao, L.Q., Liu, M., and Wang, B. (2017). The long noncoding RNA PVT1 functions as a competing endogenous RNA by sponging miR-186 in gastric cancer. *Biomed Pharmacother* 88, 302-308.

Huang, Z., and Feng, Y. (2016). Exosomes derived from hypoxic colorectal cancer cells promotes angiogenesis through Wnt4 induced beta-catenin signaling in endothelial cells. *Oncol Res*.

Isaacs, J.S., Jung, Y.J., Mole, D.R., Lee, S., Torres-Cabala, C., Chung, Y.L., Merino, M., Trepel, J., Zbar, B., Toro, J., et al. (2005). HIF overexpression correlates with biallelic loss of fumarate hydratase in renal cancer: novel role of fumarate in regulation of HIF stability. *Cancer Cell* 8, 143-153.

Ito, S., Shen, L., Dai, Q., Wu, S.C., Collins, L.B., Swenberg, J.A., He, C., and Zhang, Y. (2011). Tet proteins can convert 5-methylcytosine to 5-formylcytosine and 5-carboxylcytosine. *Science* 333, 1300-1303.

Ivan, M., Harris, A.L., Martelli, F., and Kulshreshtha, R. (2008). Hypoxia response and microRNAs: no longer two separate worlds. *J Cell Mol Med* 12, 1426-1431.

John, S., Sabo, P.J., Thurman, R.E., Sung, M.H., Biddie, S.C., Johnson, T.A., Hager, G.L., and Stamatoyannopoulos, J.A. (2011). Chromatin accessibility pre-determines glucocorticoid receptor binding patterns. *Nat Genet* 43, 264-268.

Johnson, A.B., Denko, N., and Barton, M.C. (2008). Hypoxia induces a novel signature of chromatin modifications and global repression of transcription. *Mutat Res* 640, 174-179.

Joshi, H.P., Subramanian, I.V., Schnettler, E.K., Ghosh, G., Rupaimoole, R., Evans, C., Saluja, M., Jing, Y., Cristina, I., Roy, S., et al. (2014). Dynamin 2 along with microRNA-199a reciprocally regulate hypoxia-inducible factors and ovarian cancer metastasis. *Proc Natl Acad Sci U S A* 111, 5331-5336.

Jung, K.O., Youn, H., Lee, C.H., Kang, K.W., and Chung, J.K. (2017). Visualization of exosome-mediated miR-210 transfer from hypoxic tumor cells. *Oncotarget* 8, 9899-9910.

Kaelin, W.G., Jr., and Ratcliffe, P.J. (2008). Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Mol Cell* 30, 393-402.

Kalousi, A., Mylonis, I., Politou, A.S., Chachami, G., Paraskeva, E., and Simos, G. (2010). Casein kinase 1 regulates human hypoxia-inducible factor HIF-1. *J Cell Sci* 123, 2976-2986.

Kelly, T.J., Souza, A.L., Clish, C.B., and Puigserver, P. (2011). A hypoxia-induced positive feedback loop promotes hypoxia-inducible factor 1 $\alpha$  stability through miR-210 suppression of glycerol-3-phosphate dehydrogenase 1-like. *Mol Cell Biol* 31, 2696-2706.

Kenneth, N.S., Mudie, S., van Uden, P., and Rocha, S. (2009). SWI/SNF regulates the cellular response to hypoxia. *J Biol Chem* 284, 4123-4131.

Kim, H.W., Haider, H.K., Jiang, S., and Ashraf, M. (2009). Ischemic preconditioning augments survival of stem cells via miR-210 expression by targeting caspase-8-associated protein 2. *J Biol Chem* 284, 33161-33168.

Kim, J.H., Park, S.G., Song, S.Y., Kim, J.K., and Sung, J.H. (2013). Reactive oxygen species-responsive miR-210 regulates proliferation and migration of adipose-derived stem cells via PTPN2. *Cell Death Dis* 4, e588.

Kim, M.S., Kwon, H.J., Lee, Y.M., Baek, J.H., Jang, J.E., Lee, S.W., Moon, E.J., Kim, H.S., Lee, S.K., Chung, H.Y., et al. (2001). Histone deacetylases induce angiogenesis by negative regulation of tumor suppressor genes. *Nat Med* 7, 437-443.

King, H.W., Michael, M.Z., and Gleagle, J.M. (2012). Hypoxic enhancement of exosome release by breast cancer cells. *BMC Cancer* 12, 421.

Klein, B.J., Piao, L., Xi, Y., Rincon-Arango, H., Rothbart, S.B., Peng, D., Wen, H., Larson, C., Zhang, X., Zheng, X., et al. (2014). The histone-H3K4-specific demethylase KDM5B binds to its substrate and product through distinct PHD fingers. *Cell Rep* 6, 325-335.

Kong, X., Lin, Z., Liang, D., Fath, D., Sang, N., and Caro, J. (2006). Histone deacetylase inhibitors induce VHL and ubiquitin-independent proteasomal degradation of hypoxia-inducible factor 1 $\alpha$ . *Mol Cell Biol* 26, 2019-2028.

Koslowski, M., Luxemburger, U., Tureci, O., and Sahin, U. (2011). Tumor-associated CpG demethylation augments hypoxia-induced effects by positive autoregulation of HIF-1 $\alpha$ . *Oncogene* 30, 876-882.

Koyanagi, S., Kuramoto, Y., Nakagawa, H., Aramaki, H., Ohdo, S., Soeda, S., and Shimeno, H. (2003). A molecular mechanism regulating circadian expression of vascular endothelial growth factor in tumor cells. *Cancer Res* 63, 7277-7283.

Krieg, A.J., Rankin, E.B., Chan, D., Razorenova, O., Fernandez, S., and Giaccia, A.J. (2010). Regulation of the histone demethylase JMJD1A by hypoxia-inducible factor 1 alpha enhances hypoxic gene expression and tumor growth. *Mol Cell Biol* *30*, 344-353.

Kucharzewska, P., Christianson, H.C., Welch, J.E., Svensson, K.J., Fredlund, E., Ringner, M., Morgelin, M., Bourseau-Guilmain, E., Bengzon, J., and Belting, M. (2013). Exosomes reflect the hypoxic status of glioma cells and mediate hypoxia-dependent activation of vascular cells during tumor development. *Proc Natl Acad Sci U S A* *110*, 7312-7317.

Laukka, T., Mariani, C.J., Ithantola, T., Cao, J.Z., Hokkanen, J., Kaelin, W.G., Jr., Godley, L.A., and Koivunen, P. (2016). Fumarate and Succinate Regulate Expression of Hypoxia-inducible Genes via TET Enzymes. *J Biol Chem* *291*, 4256-4265.

Law, A.Y., Lai, K.P., Ip, C.K., Wong, A.S., Wagner, G.F., and Wong, C.K. (2008). Epigenetic and HIF-1 regulation of stanniocalcin-2 expression in human cancer cells. *Exp Cell Res* *314*, 1823-1830.

Ledaki, I., McIntyre, A., Wigfield, S., Buffa, F., McGowan, S., Baban, D., Li, J.L., and Harris, A.L. (2015). Carbonic anhydrase IX induction defines a heterogeneous cancer cell response to hypoxia and mediates stem cell-like properties and sensitivity to HDAC inhibition. *Oncotarget* *6*, 19413-19427.

Lee, H.Y., Choi, K., Oh, H., Park, Y.K., and Park, H. (2014). HIF-1-dependent induction of Jumonji domain-containing protein (JMJD) 3 under hypoxic conditions. *Mol Cells* *37*, 43-50.

Li, L., Li, C., Wang, S., Wang, Z., Jiang, J., Wang, W., Li, X., Chen, J., Liu, K., Li, C., et al. (2016a). Exosomes Derived from Hypoxic Oral Squamous Cell Carcinoma Cells Deliver miR-21 to Normoxic Cells to Elicit a Prometastatic Phenotype. *Cancer Res* *76*, 1770-1780.

Li, W., He, X., Xue, R., Zhang, Y., Zhang, X., Lu, J., Zhang, Z., and Xue, L. (2016b). Combined over-expression of the hypoxia-inducible factor 2alpha gene and its long non-coding RNA predicts unfavorable prognosis of patients with osteosarcoma. *Pathol Res Pract* *212*, 861-866.

Li, X., Liu, L., Yang, S., Song, N., Zhou, X., Gao, J., Yu, N., Shan, L., Wang, Q., Liang, J., et al. (2014). Histone demethylase KDM5B is a key regulator of genome stability. *Proc Natl Acad Sci U S A* *111*, 7096-7101.

Li, X., Wu, Y., Liu, A., and Tang, X. (2016c). Long non-coding RNA UCA1 enhances tamoxifen resistance in breast cancer cells through a miR-18a-HIF1alpha feedback regulatory loop. *Tumour Biol* *37*, 14733-14743.

Lin, A., Li, C., Xing, Z., Hu, Q., Liang, K., Han, L., Wang, C., Hawke, D.H., Wang, S., Zhang, Y., et al. (2016). The LINK-A lncRNA activates normoxic HIF1alpha signalling in triple-negative breast cancer. *Nat Cell Biol* *18*, 213-224.

Liu, C., Kelnar, K., Liu, B., Chen, X., Calhoun-Davis, T., Li, H., Patrawala, L., Yan, H., Jeter, C., Honorio, S., et al. (2011a). The microRNA miR-34a inhibits prostate cancer stem cells and metastasis by directly repressing CD44. *Nat Med* *17*, 211-215.

Liu, Q., Liu, L., Zhao, Y., Zhang, J., Wang, D., Chen, J., He, Y., Wu, J., Zhang, Z., and Liu, Z. (2011b). Hypoxia induces genomic DNA demethylation through the activation of HIF-1alpha and transcriptional upregulation of MAT2A in hepatoma cells. *Mol Cancer Ther* *10*, 1113-1123.

Lu, Y., Chu, A., Turker, M.S., and Glazer, P.M. (2011). Hypoxia-induced epigenetic regulation and silencing of the BRCA1 promoter. *Mol Cell Biol* *31*, 3339-3350.

Luo, W., Chang, R., Zhong, J., Pandey, A., and Semenza, G.L. (2012). Histone demethylase JMJD2C is a coactivator for hypoxia-inducible factor 1 that is required for breast cancer progression. *Proc Natl Acad Sci U S A* *109*, E3367-3376.

Matouk, I.J., Halle, D., Raveh, E., Gilon, M., Sorin, V., and Hochberg, A. (2016). The role of the oncofetal H19 lncRNA in tumor metastasis: orchestrating the EMT-MET decision. *Oncotarget* *7*, 3748-3765.

Mayo, J.N., and Bearden, S.E. (2015). Driving the Hypoxia-Inducible Pathway in Human Pericytes Promotes Vascular Density in an Exosome-Dependent Manner. *Microcirculation* *22*, 711-723.

McCarty, G., and Loeb, D.M. (2015). Hypoxia-sensitive epigenetic regulation of an antisense-oriented lncRNA controls WT1 expression in myeloid leukemia cells. *PLoS One* *10*, e0119837.

Melvin, A., and Rocha, S. (2012). Chromatin as an oxygen sensor and active player in the hypoxia response. *Cell Signal* *24*, 35-43.

Michalik, K.M., You, X., Manavski, Y., Doddaballapur, A., Zornig, M., Braun, T., John, D., Ponomareva, Y., Chen, W., Uchida, S., et al. (2014). Long noncoding RNA MALAT1 regulates endothelial cell function and vessel growth. *Circ Res* *114*, 1389-1397.

Mimura, I., Nangaku, M., Kanki, Y., Tsutsumi, S., Inoue, T., Kohro, T., Yamamoto, S., Fujita, T., Shimamura, T., Suehiro, J., et al. (2012). Dynamic change of chromatin conformation in response to hypoxia enhances the expression of GLUT3 (SLC2A3) by cooperative interaction of hypoxia-inducible factor 1 and KDM3A. *Mol Cell Biol* *32*, 3018-3032.

Mineo, M., Ricklefs, F., Rooj, A.K., Lyons, S.M., Ivanov, P., Ansari, K.I., Nakano, I., Chiocca, E.A., Godlewski, J., and Bronisz, A. (2016). The Long Non-coding RNA HIF1A-AS2 Facilitates the Maintenance of Mesenchymal Glioblastoma Stem-like Cells in Hypoxic Niches. *Cell Rep* *15*, 2500-2509.

Murai, M., Toyota, M., Suzuki, H., Satoh, A., Sasaki, Y., Akino, K., Ueno, M., Takahashi, F., Kusano, M., Mita, H., et al. (2005). Aberrant methylation and silencing of the BNIP3 gene in colorectal and gastric cancer. *Clin Cancer Res* *11*, 1021-1027.

Nakamura, J., Kitajima, Y., Kai, K., Hashiguchi, K., Hiraki, M., Noshiro, H., and Miyazaki, K. (2011). Expression of hypoxic marker CA IX is regulated by site-specific DNA methylation and is associated with the histology of gastric cancer. *Am J Pathol* *178*, 515-524.

Nakazawa, M.S., Eisinger-Mathason, T.S., Sadri, N., Ochocki, J.D., Gade, T.P., Amin, R.K., and Simon, M.C. (2016). Epigenetic re-expression of HIF-2alpha suppresses soft tissue sarcoma growth. *Nat Commun* *7*, 10539.

Okami, J., Simeone, D.M., and Logsdon, C.D. (2004). Silencing of the hypoxia-inducible cell death protein BNIP3 in pancreatic cancer. *Cancer Res* *64*, 5338-5346.

Pal, A., Srivastava, T., Sharma, M.K., Mehndiratta, M., Das, P., Sinha, S., and Chattopadhyay, P. (2010). Aberrant methylation and associated transcriptional mobilization of Alu elements contributes to genomic instability in hypoxia. *J Cell Mol Med* *14*, 2646-2654.

Parolini, I., Federici, C., Raggi, C., Lugini, L., Palleschi, S., De Milito, A., Coscia, C., Iessi, E., Logozzi, M., Molinari, A., et al. (2009). Microenvironmental pH is a key factor for exosome traffic in tumor cells. *J Biol Chem* *284*, 34211-34222.

Peek, C.B., Levine, D.C., Cedernaes, J., Taguchi, A., Kobayashi, Y., Tsai, S.J., Bonar, N.A., McNulty, M.R., Ramsey, K.M., and Bass, J. (2017). Circadian Clock Interaction with HIF1alpha Mediates Oxygenic Metabolism and Anaerobic Glycolysis in Skeletal Muscle. *Cell Metab* *25*, 86-92.

Place, T.L., Fitzgerald, M.P., Venkataraman, S., Vorrink, S.U., Case, A.J., Teoh, M.L., and Domann, F.E. (2011). Aberrant promoter CpG methylation is a mechanism for impaired PHD3 expression in a diverse set of malignant cells. *PLoS One* *6*, e14617.

Platt, J.L., Salama, R., Smythies, J., Choudhry, H., Davies, J.O., Hughes, J.R., Ratcliffe, P.J., and Mole, D.R. (2016). Capture-C reveals preformed chromatin interactions between HIF-binding sites and distant promoters. *EMBO Rep* *17*, 1410-1421.

Pollard, P.J., Loenarz, C., Mole, D.R., McDonough, M.A., Gleadle, J.M., Schofield, C.J., and Ratcliffe, P.J. (2008). Regulation of Jumonji-domain-containing histone demethylases by hypoxia-inducible factor (HIF)-1alpha. *Biochem J* *416*, 387-394.

Polytarchou, C., Iliopoulos, D., Hatzia Apostolou, M., Kottakis, F., Maroulakou, I., Struhl, K., and Tschlis, P.N. (2011). Akt2 regulates all Akt isoforms and promotes resistance to hypoxia through induction of miR-21 upon oxygen deprivation. *Cancer Res* *71*, 4720-4731.

Prickaerts, P., Adriaens, M.E., Beucken, T.V., Koch, E., Dubois, L., Dahlmans, V.E., Gits, C., Evelo, C.T., Chan-Seng-Yue, M., Wouters, B.G., et al. (2016). Hypoxia increases genome-wide bivalent epigenetic marking by specific gain of H3K27me3. *Epigenetics Chromatin* *9*, 46.

Qin, Q., Furong, W., and Baosheng, L. (2014). Multiple functions of hypoxia-regulated miR-210 in cancer. *J Exp Clin Cancer Res* *33*, 50.

Ramteke, A., Ting, H., Agarwal, C., Mateen, S., Somasagara, R., Hussain, A., Graner, M., Frederick, B., Agarwal, R., and Deep, G. (2015). Exosomes secreted under hypoxia enhance invasiveness and stemness of prostate cancer cells by targeting adherens junction molecules. *Mol Carcinog* 54, 554-565.

Rankin, E.B., and Giaccia, A.J. (2016). Hypoxic control of metastasis. *Science* 352, 175-180.

Ratcliffe, P.J. (2013). Oxygen sensing and hypoxia signalling pathways in animals: the implications of physiology for cancer. *J Physiol* 591, 2027-2042.

Riches, A., Campbell, E., Borger, E., and Powis, S. (2014). Regulation of exosome release from mammary epithelial and breast cancer cells - a new regulatory pathway. *Eur J Cancer* 50, 1025-1034.

Rokavec, M., Oner, M.G., Li, H., Jackstadt, R., Jiang, L., Lodygin, D., Kaller, M., Horst, D., Ziegler, P.K., Schwitalla, S., et al. (2014). IL-6R/STAT3/miR-34a feedback loop promotes EMT-mediated colorectal cancer invasion and metastasis. *J Clin Invest* 124, 1853-1867.

Rong, L., Li, R., Li, S., and Luo, R. (2016). Immunosuppression of breast cancer cells mediated by transforming growth factor-beta in exosomes from cancer cells. *Oncol Lett* 11, 500-504.

Rupaimoole, R., Calin, G.A., Lopez-Berestein, G., and Sood, A.K. (2016a). miRNA Deregulation in Cancer Cells and the Tumor Microenvironment. *Cancer Discov* 6, 235-246.

Rupaimoole, R., Ivan, C., Yang, D., Gharpure, K.M., Wu, S.Y., Pecot, C.V., Previs, R.A., Nagaraja, A.S., Armaiz-Pena, G.N., McGuire, M., et al. (2016b). Hypoxia-upregulated microRNA-630 targets Dicer, leading to increased tumor progression. *Oncogene* 35, 4312-4320.

Sahar, S., Zocchi, L., Kinoshita, C., Borrelli, E., and Sassone-Corsi, P. (2010). Regulation of BMAL1 protein stability and circadian function by GSK3beta-mediated phosphorylation. *PLoS One* 5, e8561.

Salminen, A., Kaarniranta, K., and Kauppinen, A. (2016). Hypoxia-Inducible Histone Lysine Demethylases: Impact on the Aging Process and Age-Related Diseases. *Aging Dis* 7, 180-200.

Sanchez-Vega, F., Gotea, V., Petrykowska, H.M., Margolin, G., Krivak, T.C., DeLoia, J.A., Bell, D.W., and Elnitski, L. (2013). Recurrent patterns of DNA methylation in the ZNF154, CASP8, and VHL promoters across a wide spectrum of human solid epithelial tumors and cancer cell lines. *Epigenetics* 8, 1355-1372.

Sato, F., Bhawal, U.K., Yoshimura, T., and Muragaki, Y. (2016). DEC1 and DEC2 Crosstalk between Circadian Rhythm and Tumor Progression. *J Cancer* 7, 153-159.

Schlaepfer, I.R., Nambiar, D.K., Ramteke, A., Kumar, R., Dhar, D., Agarwal, C., Bergman, B., Graner, M., Maroni, P., Singh, R.P., et al. (2015). Hypoxia induces triglycerides accumulation in prostate cancer cells and extracellular vesicles supporting growth and invasiveness following reoxygenation. *Oncotarget* 6, 22836-22856.

Schmitt, A.M., Schmid, S., Rudolph, T., Anlauf, M., Prinz, C., Kloppel, G., Moch, H., Heitz, P.U., Komminoth, P., and Perren, A. (2009). VHL inactivation is an important pathway for the development of malignant sporadic pancreatic endocrine tumors. *Endocr Relat Cancer* 16, 1219-1227.

Schodel, J., Oikonomopoulos, S., Ragoussis, J., Pugh, C.W., Ratcliffe, P.J., and Mole, D.R. (2011). High-resolution genome-wide mapping of HIF-binding sites by ChIP-seq. *Blood* 117, e207-217.

Schoepflin, Z.R., Shapiro, I.M., and Risbud, M.V. (2016). Class I and IIa HDACs Mediate HIF-1alpha Stability Through PHD2-Dependent Mechanism, While HDAC6, a Class IIb Member, Promotes HIF-1alpha Transcriptional Activity in Nucleus Pulposus Cells of the Intervertebral Disc. *J Bone Miner Res* 31, 1287-1299.

Selak, M.A., Armour, S.M., MacKenzie, E.D., Boulahbel, H., Watson, D.G., Mansfield, K.D., Pan, Y., Simon, M.C., Thompson, C.B., and Gottlieb, E. (2005). Succinate links TCA cycle dysfunction to oncogenesis by inhibiting HIF-alpha prolyl hydroxylase. *Cancer Cell* 7, 77-85.

Shahzad, S., Bertrand, K., Minhas, K., and Coomber, B.L. (2007). Induction of DNA hypomethylation by tumor hypoxia. *Epigenetics* 2, 119-125.

Shen, J., Xia, W., Khotskaya, Y.B., Huo, L., Nakanishi, K., Lim, S.O., Du, Y., Wang, Y., Chang, W.C., Chen, C.H., et al. (2013). EGFR modulates microRNA maturation in response to hypoxia through phosphorylation of AGO2. *Nature* 497, 383-387.

Shi, X.F., Wang, H., Kong, F.X., Xu, Q.Q., Xiao, F.J., Yang, Y.F., Ge, R.L., and Wang, L.S. (2017). Exosomal miR-486 regulates hypoxia-induced erythroid differentiation of erythroleukemia cells through targeting Sirt1. *Exp Cell Res* 351, 74-81.

Shih, J.W., Chiang, W.F., Wu, A.T.H., Wu, M.H., Wang, L.Y., Yu, Y.L., Hung, Y.W., Wang, W.C., Chu, C.Y., Hung, C.L., et al. (2017). Long noncoding RNA LncHIFCAR/MIR31HG is a HIF-1 $\alpha$  co-activator driving oral cancer progression. *Nat Commun* 8, 15874.

Slee, R.B., Steiner, C.M., Herbert, B.S., Vance, G.H., Hickey, R.J., Schwarz, T., Christan, S., Radovich, M., Schneider, B.P., Schindelhauer, D., et al. (2012). Cancer-associated alteration of pericentromeric heterochromatin may contribute to chromosome instability. *Oncogene* 31, 3244-3253.

Stewart, G.D., Powles, T., Van Neste, C., Meynert, A., O'Mahony, F., Laird, A., Deforce, D., Van Nieuwerburgh, F., Trooskens, G., Van Criekinge, W., et al. (2016). Dynamic epigenetic changes to VHL occur with sunitinib in metastatic clear cell renal cancer. *Oncotarget* 7, 25241-25250.

Su, X., Li, G., and Liu, W. (2017). The Long Noncoding RNA Cancer Susceptibility Candidate 9 Promotes Nasopharyngeal Carcinogenesis via Stabilizing HIF1 $\alpha$ . *DNA Cell Biol* 36, 394-400.

Sun, Y.W., Chen, Y.F., Li, J., Huo, Y.M., Liu, D.J., Hua, R., Zhang, J.F., Liu, W., Yang, J.Y., Fu, X.L., et al. (2014). A novel long non-coding RNA ENST00000480739 suppresses tumour cell invasion by regulating OS-9 and HIF-1 $\alpha$  in pancreatic ductal adenocarcinoma. *Br J Cancer* 111, 2131-2141.

Tadokoro, H., Umezumi, T., Ohyashiki, K., Hirano, T., and Ohyashiki, J.H. (2013). Exosomes derived from hypoxic leukemia cells enhance tube formation in endothelial cells. *J Biol Chem* 288, 34343-34351.

Tahiliani, M., Koh, K.P., Shen, Y., Pastor, W.A., Bandukwala, H., Brudno, Y., Agarwal, S., Iyer, L.M., Liu, D.R., Aravind, L., et al. (2009). Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* 324, 930-935.

Takahashi, J.S. (2017). Transcriptional architecture of the mammalian circadian clock. *Nat Rev Genet* 18, 164-179.

Takahashi, K., Yan, I.K., Haga, H., and Patel, T. (2014). Modulation of hypoxia-signaling pathways by extracellular linc-RoR. *J Cell Sci* 127, 1585-1594.

Tanimoto, K., Tsuchihara, K., Kanai, A., Arauchi, T., Esumi, H., Suzuki, Y., and Sugano, S. (2010). Genome-wide identification and annotation of HIF-1 $\alpha$  binding sites in two cell lines using massively parallel sequencing. *Hugo J* 4, 35-48.

Tao, H., Wang, M.M., Zhang, M., Zhang, S.P., Wang, C.H., Yuan, W.J., Sun, T., He, L.J., and Hu, Q.K. (2016). MiR-126 Suppresses the Glucose-Stimulated Proliferation via IRS-2 in INS-1 beta Cells. *PLoS One* 11, e0149954.

Tasharofi, B., Soudyab, M., Nikpayam, E., Iranpour, M., Mirfakhraie, R., Sarrafzadeh, S., Geranpayeh, L., Azargashb, E., Sayad, A., and Ghafouri-Fard, S. (2016). Comparative expression analysis of hypoxia-inducible factor- $\alpha$  and its natural occurring antisense in breast cancer tissues and adjacent noncancerous tissues. *Cell Biochem Funct* 34, 572-578.

Tausendschon, M., Dehne, N., and Brune, B. (2011). Hypoxia causes epigenetic gene regulation in macrophages by attenuating Jumonji histone demethylase activity. *Cytokine* 53, 256-262.

Thienpont, B., Steinbacher, J., Zhao, H., D'Anna, F., Kuchnio, A., Ploumaki, A., Ghesquiere, B., Van Dyck, L., Boeckx, B., Schoonjans, L., et al. (2016). Tumour hypoxia causes DNA hypermethylation by reducing TET activity. *Nature* 537, 63-68.

Tomasetti, M., Nocchi, L., Staffolani, S., Manzella, N., Amati, M., Goodwin, J., Kluckova, K., Nguyen, M., Strafella, E., Bajzikova, M., et al. (2014). MicroRNA-126 suppresses mesothelioma malignancy by targeting IRS1 and interfering with the mitochondrial function. *Antioxid Redox Signal* 21, 2109-2125.

Tsai, Y.P., Chen, H.F., Chen, S.Y., Cheng, W.C., Wang, H.W., Shen, Z.J., Song, C., Teng, S.C., He, C., and Wu, K.J. (2014). TET1 regulates hypoxia-induced epithelial-mesenchymal transition by acting as a co-activator. *Genome Biol* 15, 513.

Uchida, T., Rossignol, F., Matthay, M.A., Mounier, R., Couette, S., Clottes, E., and Clerici, C. (2004). Prolonged hypoxia differentially regulates hypoxia-inducible factor (HIF)-1 $\alpha$  and HIF-2 $\alpha$  expression in lung epithelial cells: implication of natural antisense HIF-1 $\alpha$ . *J Biol Chem* 279, 14871-14878.

Umezū, T., Tadokoro, H., Azuma, K., Yoshizawa, S., Ohyashiki, K., and Ohyashiki, J.H. (2014). Exosomal miR-135b shed from hypoxic multiple myeloma cells enhances angiogenesis by targeting factor-inhibiting HIF-1. *Blood* *124*, 3748-3757.

van den Beucken, T., Koch, E., Chu, K., Rupaimoole, R., Prickaerts, P., Adriaens, M., Voncken, J.W., Harris, A.L., Buffa, F.M., Haider, S., et al. (2014). Hypoxia promotes stem cell phenotypes and poor prognosis through epigenetic regulation of DICER. *Nat Commun* *5*, 5203.

Vasagiri, N., and Kutala, V.K. (2014). Structure, function, and epigenetic regulation of BNIP3: a pathophysiological relevance. *Mol Biol Rep* *41*, 7705-7714.

Vinas, J.L., Burger, D., Zimpelmann, J., Haneef, R., Knoll, W., Campbell, P., Gutsol, A., Carter, A., Allan, D.S., and Burns, K.D. (2016). Transfer of microRNA-486-5p from human endothelial colony forming cell-derived exosomes reduces ischemic kidney injury. *Kidney Int* *90*, 1238-1250.

Voellenkle, C., Rooij, J., Guffanti, A., Brini, E., Fasanaro, P., Isaia, E., Croft, L., David, M., Capogrossi, M.C., Moles, A., et al. (2012). Deep-sequencing of endothelial cells exposed to hypoxia reveals the complexity of known and novel microRNAs. *RNA* *18*, 472-484.

Wan, W., Peng, K., Li, M., Qin, L., Tong, Z., Yan, J., Shen, B., and Yu, C. (2017). Histone demethylase JMJD1A promotes urinary bladder cancer progression by enhancing glycolysis through coactivation of hypoxia inducible factor 1 $\alpha$ . *Oncogene*.

Wang, J., Wang, Y., and Lu, L. (2012). De-SUMOylation of CCCTC binding factor (CTCF) in hypoxic stress-induced human corneal epithelial cells. *J Biol Chem* *287*, 12469-12479.

Wang, T., Gilkes, D.M., Takano, N., Xiang, L., Luo, W., Bishop, C.J., Chaturvedi, P., Green, J.J., and Semenza, G.L. (2014). Hypoxia-inducible factors and RAB22A mediate formation of microvesicles that stimulate breast cancer invasion and metastasis. *Proc Natl Acad Sci U S A* *111*, E3234-3242.

Wang, Y., Yao, J., Meng, H., Yu, Z., Wang, Z., Yuan, X., Chen, H., and Wang, A. (2015). A novel long non-coding RNA, hypoxia-inducible factor-2 $\alpha$  promoter upstream transcript, functions as an inhibitor of osteosarcoma stem cells in vitro. *Mol Med Rep* *11*, 2534-2540.

Watson, C.J., Collier, P., Tea, I., Neary, R., Watson, J.A., Robinson, C., Phelan, D., Ledwidge, M.T., McDonald, K.M., McCann, A., et al. (2014). Hypoxia-induced epigenetic modifications are associated with cardiac tissue fibrosis and the development of a myofibroblast-like phenotype. *Hum Mol Genet* *23*, 2176-2188.

Watson, J.A., Watson, C.J., McCrohan, A.M., Woodfine, K., Tosetto, M., McDaid, J., Gallagher, E., Betts, D., Baugh, J., O'Sullivan, J., et al. (2009). Generation of an epigenetic signature by chronic hypoxia in prostate cells. *Hum Mol Genet* *18*, 3594-3604.

Wenger, R.H., Kvietikova, I., Rolfs, A., Camenisch, G., and Gassmann, M. (1998). Oxygen-regulated erythropoietin gene expression is dependent on a CpG methylation-free hypoxia-inducible factor-1 DNA-binding site. *Eur J Biochem* *253*, 771-777.

Wu, F., Huang, W., and Wang, X. (2015a). microRNA-18a regulates gastric carcinoma cell apoptosis and invasion by suppressing hypoxia-inducible factor-1 $\alpha$  expression. *Exp Ther Med* *10*, 717-722.

Wu, M.Z., Chen, S.F., Nieh, S., Benner, C., Ger, L.P., Jan, C.I., Ma, L., Chen, C.H., Hishida, T., Chang, H.T., et al. (2015b). Hypoxia Drives Breast Tumor Malignancy through a TET-TNF $\alpha$ -p38-MAPK Signaling Axis. *Cancer Res* *75*, 3912-3924.

Wu, M.Z., Tsai, Y.P., Yang, M.H., Huang, C.H., Chang, S.Y., Chang, C.C., Teng, S.C., and Wu, K.J. (2011). Interplay between HDAC3 and WDR5 is essential for hypoxia-induced epithelial-mesenchymal transition. *Mol Cell* *43*, 811-822.

Wu, Y., Tang, D., Liu, N., Xiong, W., Huang, H., Li, Y., Ma, Z., Zhao, H., Chen, P., Qi, X., et al. (2017). Reciprocal Regulation between the Circadian Clock and Hypoxia Signaling at the Genome Level in Mammals. *Cell Metab* *25*, 73-85.

Xia, X., Lemieux, M.E., Li, W., Carroll, J.S., Brown, M., Liu, X.S., and Kung, A.L. (2009). Integrative analysis of HIF binding and transactivation reveals its role in maintaining histone methylation homeostasis. *Proc Natl Acad Sci U S A* *106*, 4260-4265.

Xu, H., Zhao, L., Fang, Q., Sun, J., Zhang, S., Zhan, C., Liu, S., and Zhang, Y. (2014). MiR-338-3p inhibits hepatocarcinoma cells and sensitizes these cells to sorafenib by targeting hypoxia-induced factor 1alpha. *PLoS One* 9, e115565.

Xue, M., Chen, W., Xiang, A., Wang, R., Chen, H., Pan, J., Pang, H., An, H., Wang, X., Hou, H., et al. (2017). Hypoxic exosomes facilitate bladder tumor growth and development through transferring long non-coding RNA-UCA1. *Mol Cancer* 16, 143.

Xue, M., Li, X., Li, Z., and Chen, W. (2014). Urothelial carcinoma associated 1 is a hypoxia-inducible factor-1alpha-targeted long noncoding RNA that enhances hypoxic bladder cancer cell proliferation, migration, and invasion. *Tumour Biol* 35, 6901-6912.

Yang, F., Huo, X.S., Yuan, S.X., Zhang, L., Zhou, W.P., Wang, F., and Sun, S.H. (2013). Repression of the long noncoding RNA-LET by histone deacetylase 3 contributes to hypoxia-mediated metastasis. *Mol Cell* 49, 1083-1096.

Yang, F., Zhang, H., Mei, Y., and Wu, M. (2014). Reciprocal regulation of HIF-1alpha and lincRNA-p21 modulates the Warburg effect. *Mol Cell* 53, 88-100.

Yang, W., Sun, T., Cao, J., Liu, F., Tian, Y., and Zhu, W. (2012). Downregulation of miR-210 expression inhibits proliferation, induces apoptosis and enhances radiosensitivity in hypoxic human hepatoma cells in vitro. *Exp Cell Res* 318, 944-954.

Ye, S.B., Zhang, H., Cai, T.T., Liu, Y.N., Ni, J.J., He, J., Peng, J.Y., Chen, Q.Y., Mo, H.Y., Jun, C., et al. (2016). Exosomal miR-24-3p impedes T-cell function by targeting FGF11 and serves as a potential prognostic biomarker for nasopharyngeal carcinoma. *J Pathol* 240, 329-340.

Yeh, Y.M., Chuang, C.M., Chao, K.C., and Wang, L.H. (2013). MicroRNA-138 suppresses ovarian cancer cell invasion and metastasis by targeting SOX4 and HIF-1alpha. *Int J Cancer* 133, 867-878.

Ying, Q., Liang, L., Guo, W., Zha, R., Tian, Q., Huang, S., Yao, J., Ding, J., Bao, M., Ge, C., et al. (2011). Hypoxia-inducible microRNA-210 augments the metastatic potential of tumor cells by targeting vacuole membrane protein 1 in hepatocellular carcinoma. *Hepatology* 54, 2064-2075.

Yoo, Y.G., Kong, G., and Lee, M.O. (2006). Metastasis-associated protein 1 enhances stability of hypoxia-inducible factor-1alpha protein by recruiting histone deacetylase 1. *EMBO J* 25, 1231-1241.

Young, L.C., and Hendzel, M.J. (2013). The oncogenic potential of Jumonji D2 (JMJD2/KDM4) histone demethylase overexpression. *Biochem Cell Biol* 91, 369-377.

Yu, C., Yang, S.L., Fang, X., Jiang, J.X., Sun, C.Y., and Huang, T. (2015). Hypoxia disrupts the expression levels of circadian rhythm genes in hepatocellular carcinoma. *Mol Med Rep* 11, 4002-4008.

Zhai, W., Sun, Y., Jiang, M., Wang, M., Gasiewicz, T.A., Zheng, J., and Chang, C. (2016). Differential regulation of LncRNA-SARCC suppresses VHL-mutant RCC cell proliferation yet promotes VHL-normal RCC cell proliferation via modulating androgen receptor/HIF-2alpha/C-MYC axis under hypoxia. *Oncogene* 35, 4866-4880.

Zhang, J., Lu, S., Zhou, Y., Meng, K., Chen, Z., Cui, Y., Shi, Y., Wang, T., and He, Q.Y. (2017). Motile hepatocellular carcinoma cells preferentially secrete sugar metabolism regulatory proteins via exosomes. *Proteomics*.

Zhao, H., Yang, L., Baddour, J., Achreja, A., Bernard, V., Moss, T., Marini, J.C., Tudawe, T., Seviour, E.G., San Lucas, F.A., et al. (2016). Tumor microenvironment derived exosomes pleiotropically modulate cancer cell metabolism. *Elife* 5, e10250.

Zhao, R., Sun, F., Bei, X., Wang, X., Zhu, Y., Jiang, C., Zhao, F., Han, B., and Xia, S. (2017). Upregulation of the long non-coding RNA FALEC promotes proliferation and migration of prostate cancer cell lines and predicts prognosis of PCa patients. *Prostate* 77, 1107-1117.

Zhen, S., Hua, L., Liu, Y.H., Sun, X.M., Jiang, M.M., Chen, W., Zhao, L., and Li, X. (2017). Inhibition of long non-coding RNA UCA1 by CRISPR/Cas9 attenuated malignant phenotypes of bladder cancer. *Oncotarget* 8, 9634-9646.

Zhou, C., Huang, C., Wang, J., Huang, H., Li, J., Xie, Q., Liu, Y., Zhu, J., Li, Y., Zhang, D., et al. (2017). LncRNA MEG3 downregulation mediated by DNMT3b contributes to nickel malignant transformation of human bronchial epithelial cells via modulating PHLPP1 transcription and HIF-1alpha translation. *Oncogene*.

Zhou, C., Ye, L., Jiang, C., Bai, J., Chi, Y., and Zhang, H. (2015). Long noncoding RNA HOTAIR, a hypoxia-inducible factor-1alpha activated driver of malignancy, enhances hypoxic cancer cell proliferation, migration, and invasion in non-small cell lung cancer. *Tumour Biol* 36, 9179-9188.

Zhou, X., Gao, Q., Wang, J., Zhang, X., Liu, K., and Duan, Z. (2014). Linc-RNA-RoR acts as a "sponge" against mediation of the differentiation of endometrial cancer stem cells by microRNA-145. *Gynecol Oncol* 133, 333-339.

Zhou, X., Sun, H., Chen, H., Zavadil, J., Kluz, T., Arita, A., and Costa, M. (2010). Hypoxia induces trimethylated H3 lysine 4 by inhibition of JARID1A demethylase. *Cancer Res* 70, 4214-4221.

Zhu, G., Wang, S., Chen, J., Wang, Z., Liang, X., Wang, X., Jiang, J., Lang, J., and Li, L. (2017). Long noncoding RNA HAS2-AS1 mediates hypoxia-induced invasiveness of oral squamous cell carcinoma. *Mol Carcinog*.

Zhu, Z., Gao, X., He, Y., Zhao, H., Yu, Q., Jiang, D., Zhang, P., Ma, X., Huang, H., Dong, D., et al. (2012). An insertion/deletion polymorphism within RERT-lncRNA modulates hepatocellular carcinoma risk. *Cancer Res* 72, 6163-6172.