

Report Number 11/68

Comment on “Frequency-dependent dispersion in porous media”

by

Yohan Davit and Michel Quintard



Oxford Centre for Collaborative Applied Mathematics  
Mathematical Institute  
24 - 29 St Giles'  
Oxford  
OX1 3LB  
England



# Comment on “Frequency-dependent dispersion in porous media”

Yohan Davit

Mathematical Institute, University of Oxford, 24-29 St Giles', Oxford, OX1 3LB, UK

Michel Quintard

Université de Toulouse; INPT, UPS; IMFT (Institut de Mécanique des Fluides de Toulouse); Allée Camille Soula, F-31400 Toulouse, France and CNRS; IMFT; F-31400 Toulouse, France

In a recent paper, Valdès-Parada and Alvarez-Ramirez [Phys. Rev. E 84 031201 (2011)] used the technique of volume averaging to derive a “frequency-dependent” dispersion tensor,  $\mathbf{D}_\gamma^*$ , the goal of which is to describe solute transport in porous media undergoing periodic processes. We describe two issues with this dispersion tensor. Firstly, we demonstrate that the definition of  $\mathbf{D}_\gamma^*$  is erroneous and derive a corrected version,  $\mathbf{D}_\gamma^{*c}$ . With this modification, the approach of Valdès-Parada and Alvarez-Ramirez becomes strictly equivalent to the one devised by Moyne [Adv. Water Res. 20 63 (1997)]. Secondly, we show that the term “frequency-dependent dispersion” is misleading because  $\mathbf{D}_\gamma^*$  and  $\mathbf{D}_\gamma^{*c}$  do not depend on the process operating frequency,  $\chi$ . The study carried out by Valdès-Parada and Alvarez-Ramirez represents a spectral analysis of the relaxation of  $\mathbf{D}_\gamma^*$  towards its steady-state, independently of any periodic operation or excitation.

## I. INTRODUCTION

The investigation of dynamic dispersion in porous media has a long history. It was first explored by addressing issues with the early time regime and unsteady flows in Taylor’s tube problem [1]. One of the simplest models that was proposed, e.g., in [2], is an advection-dispersion equation with a time-dependent longitudinal dispersion coefficient,  $D_\gamma^{*s}(t)$ ,

$$\partial_t \langle c_{A\gamma} \rangle^\gamma + \langle v_\gamma \rangle^\gamma \partial_x \langle c_{A\gamma} \rangle^\gamma = D_\gamma^{*s}(t) \partial_x^2 \langle c_{A\gamma} \rangle^\gamma, \quad (1)$$

where  $\langle \cdot \rangle^\gamma$  is the cross-sectional average over the fluid phase ( $\gamma$ ). Here, and throughout this publication, volume fractions have been assumed constant and disappear from the analysis.

This approach was criticized by a number of authors, including Sir Geoffrey Taylor himself. As he pointed out in [3], and as discussed later on in [4], “*it seems therefore that no physical meaning can be attached to the use of equations in which the coefficient of diffusion varies with the time of diffusion, even though the formulae produced by their use do represent adequately the concentrations in particular cases*”. To circumvent this problem, a delay-diffusion description was developed (see, for instance, [4] and [5]),

$$\begin{aligned} & \partial_t \langle c_{A\gamma} \rangle^\gamma + \langle v_\gamma \rangle^\gamma \partial_x \langle c_{A\gamma} \rangle^\gamma \\ &= \int_0^t \partial_\tau D_\gamma^{*c}(\tau) \partial_x^2 \langle c_{A\gamma} \rangle^\gamma(t - \tau) d\tau \\ &= \partial_t D_\gamma^{*c} \star \partial_x^2 \langle c_{A\gamma} \rangle^\gamma(t), \end{aligned} \quad (2)$$

based on the introduction of a memory function, i.e., a time-convolution, denoted here by  $\star$ . In Eq (2), the time derivative used in the integrand is fundamental because it ensures that  $D_\gamma^{*c}(t)$  conforms with essential requirements associated with a notion of dynamic dispersion. One of these requirements is that Eq (2) should degenerate into Taylor’s dispersion equation in the limit of

long times, i.e., when macroscopic times are significantly larger than characteristic times associated with the relaxation of  $D_\gamma^{*c}(t)$  towards Taylor’s dispersion coefficient,  $D_\gamma^{*c}(\infty)$ . Roughly speaking, this asymptotic behaviour corresponds to passing  $D_\gamma^{*c}(t)$  to the limit  $u(t)D_\gamma^{*c}(\infty)$  in the convolution product, where  $u(t)$  is the unit step function. The result of this operation leads to the following expression for the right-hand side of Eq (2):

$$\begin{aligned} & \int_0^t D_\gamma^{*c}(\infty) \partial_\tau u(\tau) \partial_x^2 \langle c_{A\gamma} \rangle^\gamma(t - \tau) d\tau \\ &= D_\gamma^{*c}(\infty) \int_0^t \delta(\tau) \partial_x^2 \langle c_{A\gamma} \rangle^\gamma(t - \tau) d\tau \\ &= D_\gamma^{*c}(\infty) \partial_x^2 \langle c_{A\gamma} \rangle^\gamma(t), \end{aligned} \quad (3)$$

where  $\delta(t)$  is the Dirac distribution. Here, we see that the time derivative is indispensable because it yields the correct asymptotic result, Eq (3), obtained by Taylor in [1]. Note that a rigorous analysis of convergence, although an interesting problem, is beyond the scope of this comment.

By formulating their model in the Laplace and frequency domains, rather than the time domain, Valdès-Parada and Alvarez-Ramirez [6] obscure physical interpretation and comparison with previous works. The term dynamic dispersion, as used in [6], suggests a notion similar to  $D_\gamma^{*s}$ . However, applying an inverse Laplace transform to Eq (14) in [6] yields

$$\begin{aligned} & \partial_t \langle c_{A\gamma} \rangle^\gamma + \nabla \cdot (\langle \mathbf{v}_\gamma \rangle^\gamma \langle c_{A\gamma} \rangle^\gamma) \\ &= \nabla \cdot \left[ \int_0^t \mathbf{D}_\gamma^*(\tau) \cdot \nabla \langle c_{A\gamma} \rangle^\gamma(t - \tau) d\tau \right] \\ &= \nabla \cdot [\mathbf{D}_\gamma^* \star \nabla \langle c_{A\gamma} \rangle^\gamma(t)], \end{aligned} \quad (4)$$

which hints at a connection with the convolution formulation, Eq (2), and  $D_\gamma^{*c}(t)$ . In this expression, the tensor,  $\mathbf{D}_\gamma^*(t)$ , corresponding to Eq (15) in [6], can be written in

the time domain as

$$\mathbf{D}_\gamma^*(t) = \mathcal{D}_\gamma \left( \mathbf{l}\delta(t) + \frac{1}{\mathcal{V}_\gamma} \int_{A_{\gamma\kappa}} \mathbf{n}_{\gamma\kappa} \mathbf{b}_\gamma(\mathbf{r}, t) dA \right) - \langle \tilde{\mathbf{v}}_\gamma \mathbf{b}_\gamma(\mathbf{r}, t) \rangle^\gamma, \quad (5)$$

where  $\mathbf{b}_\gamma$  solves

$$\begin{aligned} \partial_t \mathbf{b}_\gamma(\mathbf{r}, t) + \mathbf{v}_\gamma \cdot \nabla \mathbf{b}_\gamma(\mathbf{r}, t) + \tilde{\mathbf{v}}_\gamma \delta(t) \\ = \mathcal{D}_\gamma \nabla^2 \mathbf{b}_\gamma(\mathbf{r}, t), \text{ in the phase } (\gamma), \end{aligned} \quad (6)$$

with

$$-\mathbf{n}_{\gamma\kappa} \cdot \nabla \mathbf{b}_\gamma(\mathbf{r}, t) = \mathbf{n}_{\gamma\kappa} \delta(t), \text{ on } A_{\gamma\kappa}. \quad (7)$$

To ensure uniqueness of the solution (see details in [6]), we prescribe the zero initial condition,  $\mathbf{b}_\gamma(\mathbf{r}, t < 0) = \mathbf{0}$ , the average constraint,  $\langle \mathbf{b}_\gamma(\mathbf{r}, t) \rangle^\gamma = \mathbf{0}$ , and local periodicity.

Eqs (4) and (5) are not wrong and can be used to describe solute transport. However,  $\mathbf{D}_\gamma^*(t)$ , Eq (5), cannot be interpreted as a dispersion tensor because Eq (4) neglects the time derivative used with the convolution in Eq (2). Therefore, it exhibits features that are inconsistent with the notion of dispersion:

1.  $\mathbf{D}_{\gamma,ij}^*$  have dimensions of  $[(\text{Length})^2 \cdot (\text{Time})^{-2}]$ , instead of  $[(\text{Length})^2 \cdot (\text{Time})^{-1}]$ .
2.  $\mathbf{D}_\gamma^*$  does not yield the correct asymptotic behaviour. Considering Eq (5), we remark that  $\mathbf{D}_\gamma^*$  may be approximated by  $\mathbf{D}_\gamma^{*c}(\infty) \delta(t)$  in the long-time limit, but not by  $\mathbf{D}_\gamma^{*c}(\infty) u(t)$ .
3.  $\mathbf{D}_{\gamma,ij}^*$  are distributions, not functions, and  $\mathbf{D}_\gamma^*$  is not necessarily positive-semidefinite. Indeed, the identity tensor on the right-hand side of Eq (5) is weighted by the Dirac distribution,  $\delta(t)$ . Therefore, diagonal components  $\mathbf{D}_{\gamma,ii}^*$  will be *negative* in the early times of the diffusive regime, when the tensor describing tortuosity effects,  $\frac{1}{\mathcal{V}_\gamma} \int_{A_{\gamma\kappa}} \mathbf{n}_{\gamma\kappa} \mathbf{b}_\gamma(\mathbf{r}, t) dA$ , is dominant.

It follows that a three-dimensional analogue of Eq (2) should be used instead of Eq (4), and reads

$$\begin{aligned} \partial_t \langle c_{A\gamma} \rangle^\gamma + \nabla \cdot (\langle \mathbf{v}_\gamma \rangle^\gamma \langle c_{A\gamma} \rangle^\gamma) \\ = \nabla \cdot [\partial_t \mathbf{D}_\gamma^{*c} \cdot \star \nabla \langle c_{A\gamma} \rangle^\gamma(t)], \end{aligned} \quad (8)$$

where  $\mathbf{D}_\gamma^{*c}(t)$  is the correct transient dispersion tensor. This idea was used by Moyne, in [7], to develop a two-equation model of transient dispersion. Note that all these models are based on the assumption that spatial memory effects can be neglected, which may be inaccurate if the non-local spatial and temporal effects are strongly coupled. To model such effects, a variety of techniques may be used, including: non-local formulations involving spatial convolutions (see [8]); direct numerical

computation of the transport equations at the pore-scale; and higher order theories.

In addition to this problem in the definition of the dispersion tensor, there is an issue with the notion of frequency used in [6]. In theory,  $\mathbf{D}_\gamma^{*c}(t)$  and  $\mathbf{D}_\gamma^*(t)$  may depend on the frequency,  $\chi$ , of external periodic operations. For example, Smith in [5], considered the case of an oscillatory flow with a frequency-dependent velocity,  $\mathbf{v}_\gamma(t, \chi)$ . In this case, the dispersion coefficient,  $\mathbf{D}_\gamma^{*c}(t, \chi)$ , depends on the frequency of the excitation signal,  $\mathbf{v}_\gamma(t, \chi)$ . More generally,  $\mathbf{D}_\gamma^{*c}(t)$  will exhibit frequency-dependence if at least one of the parameters oscillates in the initial boundary value problem given by Eqs (13) in [6] and Eqs (6) and (7) herein. In such cases, either the velocity field,  $\mathbf{v}_\gamma$ , the molecular diffusion,  $\mathcal{D}_\gamma$ , or the position of the interface,  $A_{\gamma\kappa}$ , would need to vary in time following a periodic pattern.

*Such frequency-dependence is incompatible with the hypotheses made in [6].* Indeed,  $\mathcal{D}_\gamma$  is a constant scalar;  $A_{\gamma\kappa}$  is static; and the flow is implicitly steady. This is obvious in Eq (3) in [6] where the Laplace transform does not apply to the velocity field. Consequently, the model is not as general as suggested in the introduction of [6] and is in fact limited to the description of periodic discharges of solute at the macroscopic level. In this case,  $\mathbf{D}_\gamma^{*c}(t)$  has been implicitly decoupled from macroscale source terms during upscaling and is not frequency-dependent.

In our opinion, this is the result of a misinterpretation in the frequency-analysis. In Eq (8),  $\mathbf{D}_\gamma^{*c}(t)$  is a transient parameter and its relaxation can be decomposed into a spectrum of frequencies,  $\omega$ , via Fourier or Laplace transforms. However, the result of this analysis is independent of the frequency of the solute discharge, which will affect the behaviour of the average concentration  $\langle c_{A\gamma} \rangle^\gamma$  only through macroscale boundary conditions. Hence, Figs 4 and 6 in [6] must be understood as the spectral decomposition of  $\mathbf{D}_\gamma^*(t)$  relaxing towards its non-periodic steady-state, not as the frequency-analysis of the response of the dispersion tensor to periodic excitations. In addition, Eqs (6) and (14) in [6] need to be treated with great care because they describe spatial variations for a *fixed value of Fourier's frequency*,  $\omega$ , as illustrated in [6]-§VI. Many problems would involve time-dependent boundary conditions characterized by a spectrum of frequencies for which the impact of the convolution would be emphasized.

## II. CORRECTION TO THE DISPERSION TENSOR IN THE CASE OF PERIODIC SOLUTE DISCHARGE

In this section, we show that, in the case of periodic macroscale solute discharge, Eq (8) and  $\mathbf{D}_\gamma^{*c}(t)$  can be readily obtained using results in [6]. Our analysis is based on the following relationship between  $\mathbf{D}_\gamma^{*c}(t)$  and  $\mathbf{D}_\gamma^*(t)$  (see Eqs (4) and (8)):

$$\mathbf{D}_\gamma^*(t) = \partial_t \mathbf{D}_\gamma^{*c}(t). \quad (9)$$

Further, consider the unilateral Laplace transform defined by  $\bar{c}_{A\gamma}(s) = \int_0^\infty e^{-st} c_{A\gamma} dt$ . Following [6], we simplify expressions in the Laplace domain by imposing zero initial conditions. With this constraint, we have

$$\bar{\mathbf{D}}_\gamma^{*c}(s) = \frac{1}{s} \bar{\mathbf{D}}_\gamma^*(s). \quad (10)$$

Applying the Laplace transform to Eq (5) and then using Eq (10) yields

$$\begin{aligned} \bar{\mathbf{D}}_\gamma^{*c}(s) = \mathcal{D}_\gamma \left( \mathbf{I} \frac{1}{s} + \frac{1}{\mathcal{V}_\gamma} \int_{A_{\gamma\kappa}} \mathbf{n}_{\gamma\kappa} \frac{\bar{\mathbf{b}}_\gamma(\mathbf{r}, s)}{s} dA \right) \\ - \left\langle \tilde{\mathbf{v}}_\gamma \frac{\bar{\mathbf{b}}_\gamma(\mathbf{r}, s)}{s} \right\rangle^\gamma. \end{aligned} \quad (11)$$

Further, we define  $\bar{\mathbf{B}}_\gamma$  by

$$\bar{\mathbf{B}}_\gamma(\mathbf{r}, s) = \frac{\bar{\mathbf{b}}_\gamma(\mathbf{r}, s)}{s}. \quad (12)$$

Using Eqs (13) in [6] leads to the following differential equation for  $\bar{\mathbf{B}}_\gamma(\mathbf{r}, s)$ :

$$\begin{aligned} s \bar{\mathbf{B}}_\gamma(\mathbf{r}, s) + \mathbf{v}_\gamma \cdot \nabla \bar{\mathbf{B}}_\gamma(\mathbf{r}, s) + \frac{1}{s} \tilde{\mathbf{v}}_\gamma \\ = \mathcal{D}_\gamma \nabla^2 \bar{\mathbf{B}}_\gamma(\mathbf{r}, s), \text{ in the phase } (\gamma), \end{aligned} \quad (13)$$

with the boundary condition

$$-\mathbf{n}_{\gamma\kappa} \cdot \nabla \bar{\mathbf{B}}_\gamma(\mathbf{r}, s) = \frac{1}{s} \mathbf{n}_{\gamma\kappa}, \text{ on } A_{\gamma\kappa}, \quad (14)$$

and local periodicity. We also have the average constraint  $\langle \bar{\mathbf{B}}_\gamma \rangle^\gamma = \mathbf{0}$ .

Returning to the time domain, the so-called closure given by Eq (12) in [6] becomes

$$\tilde{c}_{A\gamma} = \partial_t \mathbf{B}_\gamma(\mathbf{r}, t) \cdot \star \nabla \langle c_{A\gamma} \rangle^\gamma = \partial_t B_{\gamma,i} * \partial_i \langle c_{A\gamma} \rangle^\gamma, \quad (15)$$

which is reminiscent of the closure proposed in [7] for the two-equation heat transfer problem. Here,  $\mathbf{B}_\gamma(\mathbf{r}, t)$  solves the following initial boundary value problem:

$$\begin{aligned} \partial_t \mathbf{B}_\gamma(\mathbf{r}, t) + \mathbf{v}_\gamma \cdot \nabla \mathbf{B}_\gamma(\mathbf{r}, t) + \tilde{\mathbf{v}}_\gamma u(t) \\ = \mathcal{D}_\gamma \nabla^2 \mathbf{B}_\gamma(\mathbf{r}, t), \text{ in the phase } (\gamma), \end{aligned} \quad (16)$$

with the boundary condition

$$-\mathbf{n}_{\gamma\kappa} \cdot \nabla \mathbf{B}_\gamma(\mathbf{r}, t) = \mathbf{n}_{\gamma\kappa} u(t), \text{ on } A_{\gamma\kappa}, \quad (17)$$

and local periodicity. We also have the initial condition,  $\mathbf{B}_\gamma(\mathbf{r}, t < 0) = \mathbf{0}$ , and the average constraint,  $\langle \mathbf{B}_\gamma \rangle^\gamma = \mathbf{0}$ . Again, we emphasize that the solution of this problem *does not depend on the frequency* of any source term associated with  $\langle c_{A\gamma} \rangle^\gamma$ , and that  $\mathbf{B}_\gamma(\mathbf{r}, t)$ , solution of an inhomogeneous linear advection-diffusion equation, will relax towards a non-periodic steady-state field  $\mathbf{B}_\gamma(\mathbf{r}, \infty)$  in the asymptotic regime.




Figure 1. Logarithmic plots illustrating the temporal behaviour of  $\mathbf{D}_{\gamma,xx}^{*c}/\mathcal{D}_\gamma$  (implicitly weighted by  $u(\tau)$ ) for the unit cell presented in [6] and  $\varepsilon_\gamma = 0.56$ . Results were computed using COMSOL Multiphysics 4.2. In the short time regime,  $\mathbf{D}_{\gamma,xx}^{*c}/\mathcal{D}_\gamma$  tends towards unity because we have assumed zero initial condition for  $\mathbf{B}'_\gamma(\mathbf{r}, t)$ . In the long-time regime,  $\mathbf{D}_{\gamma,xx}^{*c}/\mathcal{D}_\gamma < 1$  when tortuosity effects dominate dispersion effects and  $\mathbf{D}_{\gamma,xx}^{*c}/\mathcal{D}_\gamma > 1$  when dispersion effects dominate tortuosity effects.

In the time domain, the dispersion tensor,  $\mathbf{D}_\gamma^{*c}(t)$ , reads

$$\begin{aligned} \mathbf{D}_\gamma^{*c}(t) = \mathcal{D}_\gamma \left( \mathbf{I} u(t) + \frac{1}{\mathcal{V}_\gamma} \int_{A_{\gamma\kappa}} \mathbf{n}_{\gamma\kappa} \mathbf{B}_\gamma(\mathbf{r}, t) dA \right) \\ - \langle \tilde{\mathbf{v}}_\gamma \mathbf{B}_\gamma(\mathbf{r}, t) \rangle^\gamma. \end{aligned} \quad (18)$$

The traditional steady-state dispersion tensor can be expressed as

$$\begin{aligned} \mathbf{D}_\gamma^{*c}(\infty) = \mathcal{D}_\gamma \left( \mathbf{I} + \frac{1}{\mathcal{V}_\gamma} \int_{A_{\gamma\kappa}} \mathbf{n}_{\gamma\kappa} \mathbf{B}_\gamma(\mathbf{r}, \infty) dA \right) \\ - \langle \tilde{\mathbf{v}}_\gamma \mathbf{B}_\gamma(\mathbf{r}, \infty) \rangle^\gamma. \end{aligned} \quad (19)$$

### III. NUMERICAL RESULTS

To illustrate the temporal behaviour of this transient dispersion tensor, we use COMSOL Multiphysics 4.2 to solve numerically the following nondimensionalized closure problem for the periodic unit cell described in [6] with  $\varepsilon_\gamma = 0.56$ :

$$\partial_\tau \mathbf{B}'_\gamma + \mathbf{v}'_\gamma \cdot \nabla_{\mathbf{r}'} \mathbf{B}'_\gamma + \tilde{\mathbf{v}}'_\gamma = \frac{1}{Pe} \nabla_{\mathbf{r}'}^2 \mathbf{B}'_\gamma, \text{ in the phase } (\gamma), \quad (20)$$

with  $\mathbf{B}'_\gamma = \mathbf{B}_\gamma/(l_\gamma - l_\kappa)$ ,  $\mathbf{r}' = \mathbf{r}/(l_\gamma - l_\kappa)$ ,  $\tau = \sqrt{\langle \mathbf{v}_\gamma \rangle^\gamma \cdot \langle \mathbf{v}_\gamma \rangle^\gamma} t / (l_\gamma - l_\kappa)$ ,  $Pe = \sqrt{\langle \mathbf{v}_\gamma \rangle^\gamma \cdot \langle \mathbf{v}_\gamma \rangle^\gamma} (l_\gamma - l_\kappa) / \mathcal{D}_\gamma$  and  $\mathbf{v}'_\gamma = \mathbf{v}_\gamma / \sqrt{\langle \mathbf{v}_\gamma \rangle^\gamma \cdot \langle \mathbf{v}_\gamma \rangle^\gamma}$ . Boundary conditions are

$$-\mathbf{n}_{\gamma\kappa} \cdot \nabla_{\mathbf{r}'} \mathbf{B}'_\gamma = \mathbf{n}_{\gamma\kappa} u(\tau), \text{ on } A_{\gamma\kappa}, \quad (21)$$

and local periodicity. We also have the initial condition,  $\mathbf{B}'_\gamma(\mathbf{r}', \tau < 0) = \mathbf{0}$ , and the average constraint,  $\langle \mathbf{B}'_\gamma \rangle^\gamma = \mathbf{0}$ .

Similarly, the flow was obtained by imposing a pressure gradient and solving periodic Stokes equations. The longitudinal dispersion coefficient,  $D_{\gamma,xx}^{*c}(t)/\mathcal{D}_\gamma$ , was computed using Eq (18). Results for different values of the Péclet number,  $Pe$ , are plotted in Fig 1. The dispersion coefficient clearly relaxes towards its steady-state,  $D_{\gamma,xx}^{*c}(\infty)$ . Fig 6a in [6] shows a very similar behaviour for the longitudinal dispersion with a reflection symmetry that indicates the time/frequency relationship  $\omega \sim t^{-1}$ . This shows that the analysis carried out in [6] describes the relaxation of  $D_\gamma^*(t)$  *in the frequency domain*.

This application also highlights that time-dependent problems must be treated with care. In general, formulations similar to Eq (8) should be used, especially

when boundary conditions are characterized by a single relatively large frequency, a spectrum involving large frequencies, or when small-time relaxation phenomena are of interest. On the other hand, if relevant time-scale constraints are satisfied, the asymptotic version of Eq (8) may be used.

## ACKNOWLEDGMENTS

This publication was based on work supported in part by Award No KUK-C1-013-04, made by King Abdullah University of Science and Technology (KAUST). The authors would like to acknowledge the many valuable suggestions made by Helen Byrne.

- 
- [1] G. Taylor, Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences (1934-1990) **219**, 186 (1953).
  - [2] W. Gill and R. Sankarasubramanian, Proceedings of the Royal Society A: Mathematical and Physical Sciences **316**, 341 (1970).
  - [3] G. Taylor, Advances in Geophysics **6**, 101 (1959).
  - [4] R. Smith, Journal of Fluid Mechanics **105**, 469 (1981).
  - [5] R. Smith, Journal of Fluid Mechanics **114**, 379 (1982).
  - [6] F.J. Valdés-Parada and J. Alvarez-Ramirez, Physical Review E **84**, 031201 (2011).
  - [7] C. Moyne, Advances in Water Resources **20**, 63 (1997).
  - [8] D. Koch and J. Brady, Journal of Fluid Mechanics **180**, 387 (1987).



## RECENT REPORTS

45/11	A fibrocontractive mechanochemical model of dermal wound closure incorporating realistic growth factor	Murphy Hall Maini McCue McElwain
46/11	A two-compartment mechanochemical model of the roles of transforming growth factor $\beta$ and tissue tension in dermal wound healing	Murphy Hall Maini McCue McElwain
47/11	Effects of demographic noise on the synchronization of a metapopulation in a fluctuating environment	Lai Newby Bressloff
48/11	High order weak methods for stochastic differential equations based on modified equations	Abdulle Cohen Vilmart Zygalakis
49/11	The kinetics of ice-lens growth in porous media	Style Peppin
50/11	Wound healing angiogenesis: the clinical implications of a simple mathematical model	Flegg Byrne Flegg McElwain
51/11	Wound healing angiogenesis: the clinical implications of a simple mathematical model	Du Gunzburger Lehoucq Zhou
52/11	Image Inpainting based on coherence transport with Adapted distance functions	März
53/11	Surface growth kinematics via local curve evolution	Moulton Goriely
54/11	A multiple scales approach to evaporation induced Marangoni convection	Hennessey Münch
55/11	The dynamics of bistable liquid crystal wells	Luo Majumdar Erban
56/11	Real-Time Fluid Effects on Surfaces using the Closest Point Method	Auer Macdonald Treib Schneider Westermann
57/11	Isolating intrinsic noise sources in a stochastic genetic switch	Newby
58/11	Riemann-Cartan Geometry of Nonlinear Dislocation Mechanics	Yavari Goriely
59/11	Helices through 3 or 4 points?	Goriely

60/11	Bayesian data assimilation in shape registration	Cotter Cotter Vialard
61/11	Asymptotic solution of a model for bilayer organic diodes and solar cells	Richardson Please Kirkpatrick
62/11	Neural field model of binocular rivalry waves	Bressloff Webber
63/11	Front propagation in stochastic neural fields	Bressloff Webber
64/11	Stability estimates for a twisted rod under terminal loads: a three-dimensional study	Majumdar Prior Goriely
65/11	Adaptive Finite Element Method Assisted by Stochastic Simulation of Chemical Systems	Cotter Vejchodsky Erban
66/11	On the shape of force-free field lines in the solar corona	Prior Berger
67/11	Tear film thickness variations and the role of the tear meniscus	Please Fulford Fulford Collins

**Copies of these, and any other OCCAM reports can be obtained from:**

**Oxford Centre for Collaborative Applied Mathematics  
Mathematical Institute  
24 - 29 St Giles'  
Oxford  
OX1 3LB  
England  
[www.maths.ox.ac.uk/occam](http://www.maths.ox.ac.uk/occam)**