

Bacterial defences: mechanisms, evolution and antimicrobial resistance

William P. J. Smith ^{1,2,3}*, Benjamin R. Wucher ⁴, Carey D. Nadell ⁴ and Kevin R. Foster ^{2,3}*

1. *Division of Genomics, Infection and Evolution, University of Manchester, UK*

2. *Department of Biology, University of Oxford, Oxford, UK*

3. *Department of Biochemistry, University of Oxford, Oxford, UK*

4. *Department of Biological sciences, Dartmouth College, Hanover, NH, USA*

*E-mails: william.smith-4@manchester.ac.uk; kevin.foster@biology.ox.ac.uk

11 **Abstract**

12 Throughout their evolutionary history, bacteria have faced diverse threats from other
13 microorganisms, including competing bacteria, bacteriophages and predators. In response to
14 these threats, they have evolved sophisticated defence mechanisms that today also protect
15 bacteria against antibiotics and other therapies. In this Review, we explore the protective
16 strategies of bacteria, including the mechanisms, evolution and clinical implications of these
17 ancient defences. We also review the countermeasures that attackers have evolved to
18 overcome bacterial defences. We argue that understanding how bacteria defend themselves
19 in nature is important for the development of new therapies, and for minimising resistance
20 evolution.

[H1] Introduction

Bacteria are amongst the most ancient organisms on Earth¹, but across virtually every ecosystem, they are threatened by **competitor** [G] bacteria^{2–5}, **bacteriophages**⁶ [G] and **predators**⁷ [G], which are all equipped with a broad range of means to attack them. Whereas the widespread human use of antibiotics dates back a mere century, these three biotic threats have been shaping the evolution and physiology of bacteria for billions of years.

Bacteria have evolved a panoply of **defence mechanisms** [G] to avoid or mitigate harm from biotic threats. Understanding these defences is important for several reasons. They offer insights into bacterial biology, illustrating ecological challenges that bacteria faced in the past and the mechanisms that evolved to overcome them. These mechanisms are phylogenetically widespread and influence the physiology of diverse bacterial species; some components of animal innate immune systems even trace their origins to bacterial defence mechanisms⁸. Ancient defences are also central to how modern bacteria respond to antimicrobial therapies. Many defences offer broad protection against various threats, which means that bacteria often have **preadaptations** [G] that potentiate resistance to antimicrobials in the clinic. Moreover, as we search for new **biotherapeutic** [G] alternatives to antibiotics, including probiotic bacteria and phage therapy, we face many of the same challenges from these preadaptations that render bacteria hard to kill⁹.

In this Review, we explore bacterial defence mechanisms through an evolutionary lens and discuss their relevance for treating bacterial infections. We discuss the threats that bacteria face from microbial predators, competitors and viruses, and then identify common principles of defence that protect against these threats. The set of known bacterial defences is large and ever-growing, such that exhaustively cataloguing every mechanism is beyond the scope of this article. Instead, we select examples that illustrate different categories of defence, and discuss their regulation and their evolution. We close by examining how attackers have evolved to overcome bacterial defences, and discuss how the study of defences can inform on the treatment of bacterial disease (Box 1).

[H1] Bacteria face myriad threats

In a given environment, abiotic factors (for example, light, salinity or heat) produce **stressors** [G], and for host-associated bacteria, immune cells and responses may contribute others (for example, antimicrobial peptides). In this Review, however, our focus centres on the biotic challenges presented by bacterial competitors, phages, and predation by eukaryotes and specialised bacteria (Figure 1).

[H2] Bacterial competitors.

Most bacteria live in dense, multi-species communities, where competition for space and nutrient resources is severe²⁻⁴. Commensurately, bacteria have evolved diverse strategies for inhibiting and killing their competitors, many of which involve the use of specialised **weaponry** [G] (recently reviewed in Ref. ¹⁰). Antibacterial weapons are extraordinarily diverse, encompassing molecular toxins¹¹, antimicrobial peptides¹² and proteins^{13,14}, toxin-injecting¹⁵ and membrane-puncturing¹⁶ nanomachines, and even weaponized phages¹⁷. These myriad weapons harm a target bacterium by attacking its key cellular structures and processes, which results in growth inhibition or cell death. For example, diffusible peptide-based toxins (bacteriocins) often damage DNA and RNA¹⁸, or compromise cell envelopes via pore-forming¹⁹ or wall-degrading activity²⁰. Protein toxins injected via the type VI secretion system (T6SS) frequently attack the bacterial cell wall or membrane(s)²¹, lysing intoxicated cells quickly and thereby clearing a path to new targets²². Antibiotics, a diverse group of secondary metabolite toxins, have broad but overlapping activities, and common targets include gene transcription and protein translation, DNA synthesis and replication, and the cell envelope^{23,24}.

[H2] Bacteriophages.

Phages are the most numerous biological entities in the biosphere²⁵, and are a leading cause of bacterial mortality in many environments²⁶ (for a recent review, see Ref. ⁶). Phages differ widely in their evolutionary relationships with hosts, spanning a continuum from **parasitism** [G] to **mutualism** [G]²⁷. To replicate all phages must inject their genetic material into bacterial hosts. For lytic phages, the injected genetic material is immediately copied and transcribed to assemble progeny phage particles, which kill and burst the host cell to disperse. Temperate phages (for example, λ -coliphages) also reproduce via host lysis under certain conditions, but have the additional ability to lysogenize host bacteria²⁸, whereby the phage inserts its genome into the bacterial chromosome, which enables it to replicate vertically alongside its host as it grows and divides. A third class of phages (for example, filamentous phages) exhibit a chronic replicative cycle, whereby new phages are continuously extruded from the host²⁹. Lysis of cells infected with lytic phages is triggered by envelope-degrading endolysins and holins³⁰; temperate phages kill via similar mechanisms but may lie dormant for long periods before those mechanisms are induced. Cells with chronic phage infections are generally not killed²⁹, but still suffer from reduced fitness owing to the diversion of cellular resources towards phage assembly³¹.

[H2] Eukaryotic and bacterial predators.

As well as viral infection, bacteria have long faced the threat of predation, particularly from free-living protozoa that feed via phagocytosis in soil and aquatic environments⁷. Some

bacteria are also facultative or obligate bacterial predators: the soil bacterium *Myxococcus xanthus* moves rapidly in large groups, digesting encountered prey with secreted hydrolytic enzymes³². *Bdellovibrio* and like organisms (BALOs) are small bacteria that burrow inside Gram-negative bacteria: once inside the periplasm, a BALO cell grows by digesting the cytosolic contents of the host with hydrolytic enzymes, fueling rapid growth³³. Once the resources of the host are exhausted, the BALO cell divides to form multiple progeny cells, which are released via host-cell lysis³⁴. Meanwhile, the Candidate Phyla Radiation, a diverse group of small-celled bacteria representing approximately 15% of all bacterial diversity³⁵, may incorporate other new types of predatory or parasitic bacteria. Although the biology of this group remains poorly understood, members often have reduced genomes, and seem to rely on other bacteria to survive and reproduce³⁶.

[H1] Classes of bacterial defence

Bacteria have a wide range of defensive mechanisms against competitors, phages and predators. These mechanisms operate at a range of spatial scales, from molecular and cellular defences, to those that require bacteria to work as a group (Figure 2).

[H2] Molecular-scale defences.

[H3] *Target modification and protection.* To kill a bacterium, attackers deploy harmful **agents** **[G]** that interact with specific molecular targets to disrupt vital cellular processes of the target cell. Modification³⁷ or protection³⁸ of a target structure can attenuate these interactions and prevent or lessen harm. β -lactam antibiotics, such as penicillin, kill bacteria by inhibiting cell-wall cross-linking enzymes. In methicillin-resistant *Staphylococcus aureus* (MRSA), the genes *mecA* and *mecC* encode modified cross-linking enzymes that are insensitive to almost all β -lactam drugs³⁹. Modification can be post-translational as well as genetic; for instance, the enzymatic methylation of bacterial ribosomes can prevent multiple classes of antibiotics from binding with this target³⁷.

[H3] *Target repair and compensation.* Cells can compensate for the presence of a harmful agent via generalized physiological responses that repair damaged targets. Exposure of bacteria to antibiotics, other competitor toxins, or phages, often results in oxidative DNA damage^{40,41}. Subsequently, repair of oxidised DNA occurs via the base excision repair (BER) and nucleotide excision repair (NER) systems, which are both highly conserved and ancient pathways^{42,43}. Apart from chromosomal repair, some species possess RNA ligases that can mend 16S rRNA damage caused by ribotoxic bacteriocins⁴⁴. Similarly, the extrusion of filamentous phages can compromise the inner membrane of *Escherichia coli*, but the expression of membrane-binding phage-shock proteins suppresses proton leakage and

maintains the proton-motive force⁴⁵. Sometimes it suffices to simply replace lost targets: when intoxicated with cell-wall-degrading T6SS toxins, *Vibrio cholerae* responds by increasing peptidoglycan synthesis to compensate⁴⁶.

[H3] Agent modification, binding and degradation. Harmful agents can be neutralised before they inflict damage. Multiple classes of antibiotics are neutralized through modification, via the enzymatic addition of acetyl, phosphoryl or adenyl groups⁴⁷. Toxic agents can also be inactivated via binding to other molecules: the expression of cognate immunity proteins confers resistance to many bacteriocins¹⁹, T6SS⁴⁸ and Cdi⁴⁹ effectors, and enables cells to safely use these toxic proteins as weapons¹⁰. In the same way, expression of orphan immunity proteins (that is, those for which a bacterium does not produce a cognate toxin) enables bacteria to survive attacks from non-kin cells^{50,51}.

Bacteria also have diverse systems to degrade harmful agents. β -lactamases are ancient proteins that hydrolyse the ring structure of beta-lactam antibiotics, such as penicillin⁵². Restriction-modification (RM) systems encode restriction endonucleases, which bind to and cleave phage and other foreign DNA at specific recognition sites. Target modification also has a role here, but is directed at host DNA: recognition sequences on host DNA are modified (e.g. via methylation) to protect them from degradation, while unmodified phage DNA is destroyed by the endonuclease. Multiple classes of RM systems have been characterised across both bacteria and archaea^{53,54}, providing innate immunity against a subset of phages. Recently-discovered antiviral defences, such as DISARM⁵⁴ (defence island system associated with restriction–modification) and Dnd⁵⁵ (DNA phosphorothioation) systems, function in a similar manner, respectively attacking foreign DNA that lacks methyl- or sulphur modification.

The degradation of harmful agents reaches astonishing complexity in CRISPR–Cas systems, which provide bacteria with adaptive immunity against phages whose genomic signatures have previously been encountered. These systems store fragments of foreign DNA in the bacterial genome, which then guide Cas restriction enzymes to degrade DNA in the cell that resembles that of past phage infections⁵⁶ or other mobile genetic elements⁵⁷. The recently-discovered prokaryotic Argonaute (pAgo) proteins operate on a similar principle, providing guided DNA interference against harmful genetic elements including plasmids, transposons and phages⁵⁸.

[H2] Cellular defences

[H3] Membranes, capsules and extracellular vesicles. Most harmful agents must enter a cell before they can cause harm, and bacterial membranes are often pivotal in restricting this entry.

Indeed, the outer membrane of Gram-negative bacteria may have evolved in part to better protect cells from antimicrobial compounds⁵⁹. The structures decorating a membrane are also crucial to barrier function: some structures (for example, transporters or surface polysaccharides) function as binding sites or entry points for phages and protein toxins, and bacteria that lack such structures, or have modified those structures, benefit from resistance. Other surface structures (for example, lipopolysaccharides⁶⁰ and curli fibres⁶¹) confer protection by occluding phage- or toxin-binding sites, or by armouring the cell against mechanical insult. For example, bacterial capsules, which are protective sheaths of exopolysaccharides, can armour cells against penetration by the T6SS^{62,63}. Similarly, a layer of interlocking surface proteins, known as the S-layer⁶⁴, can protect bacteria from entry by *Bdellovibrio* bacteria⁶⁵, as can certain lipopolysaccharides⁶⁶. Beyond their barrier role, membranes can perform additional defensive functions when shed as bubble-like extracellular vesicles⁶⁷. As well as enhancing envelope stability (by removing misfolded or mislocalised envelope components)⁶⁷, vesicles can function as extracellular 'decoys', absorbing antibiotics, peptide toxins and phages, and carrying toxin-degrading enzymes⁶⁸. Vesicle release is actively upregulated in response to envelope stress, and is thought to have intersecting roles in anti-phage and anti-toxin defence⁶⁸.

[H3] Efflux pumps. When the cell envelope fails to stop harmful molecules from entering, bacteria can instead force them back out. Efflux pumps are a diverse group of membrane transport proteins universal to bacteria, with a broad range of substrate specificities⁶⁹ and physiological functions⁷⁰. In particular, they are an effective and fast-acting antibiotic resistance mechanism⁷¹, sufficient in some cases to protect antibiotic-producing bacteria against their own toxins⁷².

[H3] Motility. Using flagellae, type IV pili or other motility systems⁷³, bacteria can evade threats that would otherwise kill them. In planktonic environments, bacteria with sufficiently high swimming speeds ($>30 \mu\text{m s}^{-1}$) can avoid capture by protozoan predators, despite meeting them more often at high speeds⁷⁴. Indeed, motility can be beneficial even if a bacterium cannot 'outrun' a threat: *Bdellovibrio* predators swim approximately twice as fast as *V. cholerae* prey cells⁷⁵, but the drag forces generated by prey motility impede predator attachment⁶⁶. However, motility is not always a good defence: many phages bind to motility systems as part of their infection process⁷⁶, and movement can also spread phage within bacterial groups⁷⁷.

[H2] Multicellular defences

[H3] Biofilms. Clonal groups of bacteria often work together, collectively enduring threats which would kill single cells⁷⁸. The most ubiquitous example of a multicellular defence in

bacteria is the formation of **biofilms** [G]. Biofilms underlie a range of chronic infections, and often form in response to antibiotics and competition from other strains^{79–81}. They can render bacteria extremely hard to kill, for multiple reasons. Diffusion limitation of solutes, such as oxygen or nutrients, means that many biofilms contain large numbers of slow-growing or dormant cells, which are more tolerant of toxins that target cell growth and division machinery than their fast-growing counterparts⁸². The outer regions of a biofilm can also protect cells deeper inside, collectively absorbing⁸³ and degrading⁸⁴ toxins and limiting their penetration into the community. Cells in biofilms also produce a slimy matrix of polysaccharides, proteins, DNA and other compounds: these surround cells and create an additional physical barrier that can inhibit the passage of antibiotics⁸⁵, block T6SS attacks^{62,78} and screen cells from phages⁶¹ and predators⁸⁶. Matrix production can also function as an offensive strategy, which enables bacteria within the biofilm to spread out and smother competitors⁸⁷. Matrix-trapped phage can even become weapons, protecting a biofilm from invasion by competing bacteria⁸⁸.

[H3] Phenotypic heterogeneity. Another **collective defence** [G] strategy displayed by bacteria is to maintain standing population variability in phenotype (for example, growth phase), such that not all individuals fare equally badly when conditions deteriorate. Such phenotypic heterogeneity is associated with clinical antibiotic tolerance⁸⁹, and is also a route through which bacteria resist toxins from competitors⁹⁰. Sources of this variability include the gradients in nutrients and other solutes discussed above, which commonly occur in biofilms, and can drive differences in cell physiology across space⁹¹. However, phenotypic variation also emerges in the absence of environmental gradients, via stochastic mechanisms. A key example of this is the ability of bacteria to switch epigenetically to slow-growing antibiotic-tolerant ‘persister’ states⁹², or to rapid growth modes that avoid antibiotic accumulation⁹⁰. An evolutionary experiment showed that antibiotic treatment can select for *E. coli* point mutations that increase the rate of this switching, which results in high levels of multi-drug tolerance⁹³. This result suggests that production of persister cells represents an evolved defence mechanism.

[H3] Counterattacks. Sometimes offence is the best defence — true to this maxim, many bacterial species launch en-masse **counterattacks** [G] to eliminate perceived threats¹⁰. Of course, counterattack strategies can be protective at the individual level too: environmental *V. cholerae* cells use the T6SS as an anti-grazer defence⁹⁴, whereas *Pseudomonas aeruginosa* cells respond to T6SS-mediated attacks by competitors with spatially coordinated T6SS firing^{95,96}. However, for many secreted toxins, lethality is strongly dependent on producer cell density^{97,98}, making counterattacks more effective when undertaken collectively⁹⁹. Some bacteria regulate toxin counterattacks via autoinduction: when toxin concentration and

production are connected in a positive feedback loop, a minor aggression may be met with disproportionate retaliation^{83,100}. In some cases, mass-counterattacks lead to runaway conflict escalation and even mutual destruction^{100,101}.

[H3] Suicide. Saving nearby clonemates via self-sacrifice is another striking form of defence shown by bacteria. Active cell suicide is both collective and cooperative by definition, as it kills the individual while benefiting neighbouring cells. Many bacteria protect their kin from the spread of a phage infection using a strategy called abortive infection¹⁰², whereby an infected cell pre-emptively triggers its own lysis, or growth arrest, before phage particle assembly is completed, thereby sparing kin from subsequent infection. Multiple anti-phage defences, including bacterial gasdermins¹⁰³, the CBASS system¹⁰⁴, and certain toxin–antitoxin¹⁰⁵ and CRISPR systems¹⁰⁶, function in this way; other recent discoveries (for example, RADAR¹⁰⁷, Theoris¹⁰⁸ and Zorya¹⁰⁹ systems) may behave likewise. Interestingly, cell suicide is also at the heart of some striking examples of counterattack: colicin toxins produced by *E. coli* are too large to pass through standard secretion apparatus, necessitating destructive cell lysis for their release. Other large protein weapons, such as eCISs (extracellular contractile injection systems)¹¹⁰ and R tailocins¹⁴, are similarly constrained. However, it was shown that only *E. coli* cells that have already sustained lethal damage undergo the lytic toxin release pathway, which reduces the effective costs of suicide¹¹¹. The result can be a massive counterattack by the doomed cells, paralleling suicidal stinging by honeybees.

[H1] Competition sensing and defence regulation

Bacteria use some defensive structures by default; for example, the outer membrane is a permanent protective feature of Gram-negative bacteria⁵⁹. However, many defences are not fixed and are instead **plastic responses [G]** to perceived threats. These responses are distinct from evolutionary responses (population changes in genotype), which we discuss in the next section. Critical to using plastic defences is the ability to infer that a threat is present or likely to occur, and bacteria use a range of information sources (cues) to achieve this¹¹² when acclimating to new and hostile environments (Figure 3).

[H2] Bacteria sense attacks through direct and indirect means.

First, many bacteria regulate defences by sensing attack signatures; that is, cues that result directly from a biotic threat. Physiological stress is a primary indicator that a focal bacterium could be under attack, and bacteria detect stress using a wide range of **stress responses [G]**^{79,113}. These regulatory networks respond to diverse forms of stress, of both biotic and abiotic origins. However, there is strong evidence that bacteria differentiate between different stress cues, deploying anti-competitor defences only in response to stressors that are likely to stem

from a biotic threat (Figure 3). This behaviour is known as **competition sensing** [G]⁷⁹, and is thought to regulate a wide range of defences. The clearest evidence for competition sensing comes from the upregulation of antibacterial toxins, because in that case one can infer that the likely function of the response is to cope with competitors. For other defences, such as DNA repair systems, it is more challenging to tell if the response evolved primarily due to biotic or abiotic stressors. However, multiple of the major stress responses are known to be activated by biotic threats, which is consistent with their use in competition sensing^{41,81}.

Antibacterial weapons often target vital structures such as the cell envelope or chromosome (Figure 1). Damage to these components, sensed via specific stress response pathways¹¹³, is frequently used to regulate counter-attacks and structure-specific repair pathways⁷⁹. As cellular damage often results in the production of reactive oxygen species⁴¹, many bacteria also use oxidative stress as a cue to produce toxins^{79,114}. General stress responses can also be used to regulate defences: when attacked by T6SS-armed competitors, *Salmonella enterica* serovar *Typhimurium* activates various damage responses, including the general stress response, to induce biofilm formation and efflux pump expression⁸¹. In some cases, cellular perturbation is sensed without a canonical stress response: *P. aeruginosa* bacteria directly sense oncoming T6SS attacks through the resulting perturbations to its membranes, likely via the TagQRST pathway⁹⁵. By sensing the specific location of these strikes, defenders gain valuable information on the position of the attacker cells, helping them to more effectively counter-attack with their own T6SS weaponry¹¹⁵. There is also evidence that competition sensing by *P. aeruginosa* is induced by the cytotoxins of *Staphylococcus aureus*, which is a key ecological competitor during infections¹¹⁶.

Competition sensing, therefore, enables bacteria to infer the presence of competitors, and the efficient activation of defences and counter-attacks. There is growing evidence that stress responses can play analogous roles in sensing and responding to cell damage stemming from other biotic threats. Envelope stress responses are frequently triggered during phage infection: filamentous phages compromise *E. coli* membrane integrity during chronic infection, triggering the so-called 'phage shock' cascade, and activating membrane repair pathways¹¹⁷. Likewise, lytic phages stimulate phage shock proteins in *Lactococcus lactis*, which responds by altering its metabolism to restore loss of proton-motive force¹¹⁸. Certain toxin–antitoxin systems sense phage infection via canonical stress responses, or via transcriptional changes caused by infection¹¹⁹. In a similar vein, cellular damage can warn of predator activity. *Tetrahymena* ciliates engulf bacteria to feed on them, but this can activate the bacterial SOS response. When *Tetrahymena* eat enterohemorrhagic *E. coli*, the result is that the engulfed bacteria retaliate by suicidally releasing shiga toxins, killing the predator from within, and

protecting kin cells from the predator¹²⁰. Shiga toxins are the causative agents of enterohemorrhagic diarrhoea¹²¹, underscoring that anti-predator defences can be linked to human disease.

Cell damage is a reliable indicator of an urgent threat^{79,112}, but by the time a cell detects injury, it may already be too late for defensive action. For instance, *E. coli* cell invasion by *Bdellovibrio* predators prompts host upregulation of genes associated with osmotic, envelope and general stress responses, but these do not seem to confer any resistance to the predator¹²². In such cases, detecting alternative attack signatures, such as chemical cues that precede an attack, may provide an important alternative to damage sensing⁷⁹. Through 'danger sensing [G]'¹²³, bacteria intercept chemical signatures of the attacker: for example, peptidoglycan sheddings¹²⁴, or signal molecules (Figure 3). Some bacteria express receptors for quorum sensing molecules that they themselves do not produce¹²⁵, which enables them to 'eavesdrop' on the communications among competitor strains and thereby monitor their density^{79,123}. Similarly, the perception of predator-associated chemical cues is widespread in planktonic microorganisms¹²⁶; for instance, *Pseudomonas fluorescens* responds to diffusible cues produced by protozoan predators by producing membrane-disrupting biosurfactants that are toxic to protozoa¹²⁷. Intriguingly, some bacteria are even capable of directly sensing attackers' toxins (for example, antimicrobial peptides¹²³ and β -lactam antibiotics¹²⁸), and responding before the toxin takes its effect. In the sense that genetic material injected by phages is itself a harmful agent, anti-phage systems that detect foreign DNA (for example, CRISPR, restriction–modification and DISARM systems) fall into this sensing category.

When attacked, bacteria can also forewarn their kin of danger, priming defences in advance of physiological harm. When attacked by phage or antibiotics, *P. aeruginosa* cells produce a quinolone signal that repels other clonemates from the affected area¹²⁹. Similarly: in response to neighbour infection, non-infected *Bacillus subtilis* cells can modify phage binding sites (cell wall teichoic acid polymers) on their surface, adding anlyl groups that hinder phage binding¹³⁰. Cell lysate factors, such as DNA and other mislocalised cytoplasmic molecules¹³¹, often serve as danger cues for bacteria, eliciting toxin and exopolysaccharide production in kin cells. These cues are sensed via transduction pathways (for example, the Gac–Rsm and PhoPQ pathways in *P. aeruginosa* and other Gammaproteobacteria) that are often independent from classic response pathways¹³¹. As discussed, these enable cells to raise defences and launch counterattacks before they enter stress states^{123,132}.

[H2] Bacteria associate nutrient depletion with competition.

Short of direct threat, certain environmental changes can also imply the presence of competing organisms. Nutrient starvation may indicate **exploitation competition** [G], driven by high numbers of clonemates, competitors or both¹¹² (Figure 3). Consistent with their use in competition sensing, bacteria use starvation stress pathways to regulate the production of anti-competitor toxins⁷⁹. For example, the stringent response is a ubiquitous signalling cascade that is triggered by limitations to key resources, such as amino-acids, fatty acids, inorganic phosphate or iron¹³³. As well as triggering cell cycle arrest and the cessation of growth, the stringent response upregulates the production of toxins across diverse bacterial species^{134–136}.

[H2] Bacteria use kin density to forecast threats.

A third important information source for defence regulation is **quorum sensing** [G]^{112,137,138}. By monitoring the concentration of density cues (both canonical quorum sensing autoinducers and other ‘quorum-related’ cues⁷⁹; for example, peptidoglycan fragments¹²⁴) bacteria can sense high kin densities and prepare for an expected attack (Figure 3). Recent work demonstrated that CRISPR–Cas activity and adaptation is regulated via quorum sensing, such that antiviral defences are primed when bacteria are at high density, and most vulnerable to virulent phage¹³⁹. Density sensing also informs whether bacterial groups have sufficient members for collective defences to be effective. Biofilm defences are frequently regulated using quorum sensing^{137,140}; various bacterial species also use quorum sensing to control collective counterattacks, using antibiotics¹⁴¹, bacteriocins¹⁴² or T6SSs¹⁴³. For instance, when at high cell density, *P. aeruginosa* produces the phenazine pyocyanin in a quorum sensing-dependent manner. Among a wealth of other potential functions, pyocyanin production was recently found to stimulate upregulation of multiple efflux pump systems, which means cells are better defended against a range of antibiotics¹⁴⁴.

[H1] Evolution of defences

How did bacteria acquire their impressive defensive functions? At a fundamental level, the evolution of biological functions (‘adaptation’ in evolutionary biology) is driven by natural selection acting on variation¹⁴⁵. In bacteria, two key processes generate the variation upon which natural selection depends. Mutation, stemming from DNA replication error or chromosomal rearrangements¹⁴⁶, generates raw genetic sequence variation, and **horizontal gene transfer** (HGT) [G] adds further variation by mixing alleles and genes among different cells¹⁴⁷. Phages, competitors and predators can then generate natural selection and favour bacterial variants with improved defences. In this section, we discuss how these processes enable the evolution of defensive traits, before examining how this impacts bacterial genomes (Figure 4).

[H2] Evolutionary processes.

[H3] *Mutations and other genetic changes.* Compared with larger organisms, mutational variation often arises quickly in bacteria because of their short generation times and large population sizes¹⁴⁸, which can enable the rapid emergence of protective phenotypes. Simple point mutations can drastically reduce toxin-binding affinities of their targets, generating resistance to antibiotics¹⁴⁹, bacteriocins¹⁵⁰ and phages¹⁵¹ (Figure 4a). Minor changes in regulatory genes can also provide protection against harmful agents. For example, inactivation of a repressor gene (*ramR*) in *S. Typhimurium* results in over-expression of the AcrAB efflux pump, conferring resistance to diverse quinolones, phenicol, and tetracycline antibiotics¹⁵². Likewise, alterations to regulators of lipopolysaccharide¹⁵³ and cell wall synthesis¹⁵⁴ have been shown to generate resistance to bacteriocins, antibiotics and phages. Mutation rates can also increase in times of stress¹⁵⁵, or at low cell density¹⁵⁶, potentially accelerating defensive adaptation¹⁵⁷.

[H3] *Horizontal gene transfer.* Bacteria can also acquire new defensive genes from other microorganisms via conjugation, natural transformation and transduction¹⁵⁸ (Figure 4a). These HGT events have a central role in bacterial evolution¹⁵⁹, and seem to be particularly important for defence evolution¹⁶⁰. Importantly, HGT can provide a suite of new genes to a recipient cell in a single step¹⁵⁹, which confers a complex protective phenotype much faster than would be possible through mutation alone. In parallel, HGT can rapidly generate novel and beneficial combinations of alleles via recombination¹⁶¹. HGT has facilitated the spread of defences against bacterial, viral and eukaryotic threats. Resistance to antibiotics is often conferred by plasmids¹⁶² and integrative conjugative elements¹⁶³. Other antibacterial weapons and their cognate defences, including bacteriocins¹⁹, T6SS⁵¹ and Cdi¹⁶⁴ systems, are frequently encoded on mobile elements, such that bacteria can gain both resistance and potentially counterattack capability through HGT. Many phage protection systems are also extensively shared via HGT^{165–167}. Though less well-documented, anti-predator toxins can be acquired in the same manner: the biosynthetic operon for the toxin pyrrolnitrin seems to be mobile¹⁶⁸, and confers protection against protozoa to various Gram-negative bacteria¹⁶⁹.

[H3] *Natural selection and genetic drift.* Natural selection can act on the genetic variation generated by mutation and HGT whenever a threat affects survival and reproduction, and so bacterial fitness. In some situations, low population sizes can introduce stochastic changes in the frequency of a given genotype, which can limit defence evolution via genetic drift and related processes¹⁷⁰. Nevertheless, the potential strength of natural selection for bacterial defences is made clear by evolutionary experiments with competitors, phage and predators,

where the rapid evolution of defences has been observed^{154,171–173}. This potential is further underlined by the current antimicrobial resistance crisis: the widespread use of antimicrobials by humans has created concerted selection for drug-resistant bacteria, making previously treatable infections deadly.

However, even when a particular defence is under strong natural selection, it may not lead to the fixation of a given genotype. The utility of some defensive genes can diminish as they become more common (frequency-dependent selection¹⁷²). For example, variability in O-antigen composition of a pathogen is thought to be driven by frequency-dependent selection for evasion of host immune cells¹⁷⁴, intestinal protozoa¹⁷⁵ or phages¹⁷⁶, as rarer genotypes can have an advantage if they are less likely to be recognised. In other cases, **pleiotropy [G]** can limit, or enhance, selection for defensive attributes¹⁷⁷. Many defensive adaptations have secondary phenotypic effects that are subject to evolutionary trade-offs (antagonistic pleiotropy). For instance, bacteria that gain resistance to a lytic phage might suffer enhanced susceptibility to another¹⁷². Alternatively, resistance to one threat might also enhance protection to another (synergistic pleiotropy, also referred to as a ‘trade-up’)^{177,178}. Moreover, even strong trade-offs can be insufficient to drive the loss of a defensive adaptation. Compensatory mutations can substantially reduce the fitness costs of defensive genes, which enables them to persist even in the absence of a threat¹⁵⁸. This has worrying consequences for the long-term maintenance of antibiotic resistance genes: once a bacterium gains resistance, it may not easily lose it¹⁷⁹.

[H2] Evolutionary consequences.

[H3] Genomic organisation of defences. The evolution of defences can have major impacts on bacterial genomes. Across diverse environments and lifestyles, genomes are replete with genes that encode defensive functions¹⁸⁰. These genes are often clustered together in specialised repositories (Figure 4b–d), each encoding protection against a particular class of threat. Perhaps best-known are bacterial ‘defence islands’: these mosaic-like chromosomal regions are enriched in diverse anti-viral defences, and have been the source of multiple recent defence system discoveries^{54,58,109}. In addition to antiviral genes, bacteria retain clusters of toxin immunity and detoxification genes for use during anti-competitor warfare. Examples include the recently-discovered antagonism resistance (*arc1-3*) clusters in *P. aeruginosa*¹³², and the orphan immunity gene libraries (dubbed ‘acquired interbacterial defence’ (AID) arrays) widely found among human gut *Bacteroides* species^{50,51} (Figure 4b).

Some clusters acquire new defensive genes in a highly ordered manner. Many of the AID immunity genes seem to be actively captured via recombinases, which enables gut bacteria

to expand into niches occupied by aggressive competitors⁵¹. CRISPR spacer libraries can likewise be regarded as gene capture systems, which generate arrays of phage DNA templates that guard against future infections⁵⁶ (Figure 4c). Integrons, which are ancient DNA-scavenging machines that capture mobile gene cassettes¹⁸¹, commonly confer antibiotic resistance, and are another example of active defence acquisition (Figure 4d). Multi-resistance integrons (MRIs) that contain up to eight resistance cassettes have been reported¹⁸², and super-integrons with >200 cassettes are also known¹⁸³. Integron gene expression is triggered by cellular stress, and bacteria also seem to alter the expression of different integron genes by shuffling their order¹⁸⁴.

Some defences are always found in a given species (that is; they form part of its core genome). Core defences include the outer membrane of Gram-negative bacteria (thought to be an adaptation to ancient antibiotic warfare⁵⁹), some restriction–modification systems¹⁸⁵, and multi-drug efflux pumps¹⁸⁶. However, many defence genes are found in the accessory genome, and are a major contributor to intraspecific variation among bacteria^{187–189}. Indeed, the content of the accessory genome can be overwhelming defensive¹⁶⁰: in certain marine bacteria, anti-phage systems represent >90% of all accessory genes¹⁹⁰.

[H3] The impact of selfish genes. The beneficial acquisition of new defensive capacities through HGT can occur as a by-product of the infectious actions of mobile genetic elements¹⁵⁹. This can blur the lines of what can be considered a ‘bacterial’ defensive adaptation: a mobile element may be the primary recipient of the benefit of the defensive system¹⁶⁷. Consider superinfection exclusion, whereby phage infection of a bacterium prevents similar phages from infecting the same cell. While this may benefit the bacterium, superinfection exclusion presumably evolved due to benefits to the infecting phage, which then avoids competing with other phages for the hosts’ resources¹⁹¹. In a similar vein: some anti-phage or anti-plasmid systems may have first evolved not in bacterial chromosomes, but in mobile genetic elements, either as adaptations to fend off competing genetic parasites (using, for example, CRISPR and restriction-modification systems¹⁶⁷), or as systems to ensure their own maintenance during host replication (for instance, some toxin–antitoxin modules¹⁹²). Nevertheless, even if defence genes did not originate as bacterial adaptations, bacteria may still benefit from inter-parasite conflict, or come to integrate and exploit selfish genes for their own ends. For example, CRISPR–Cas systems are often now part of the bacterial chromosome, and are no longer under the direct control of mobile elements¹⁹³.

[H1] Overcoming bacterial defences

Bacterial defences have the potential to coevolve with the offensive strategies of their aggressors. A new defence mechanism can generate natural selection on attackers for countermeasures [G], examples of which are shown in Figure 5. Countermeasures may precipitate an evolutionary arms race, whereby attackers and defenders become progressively better-adapted to defeat each other¹⁹⁴. However, such escalation is only one possibility; coevolutionary dynamics can also be cyclical, which may facilitate the coexistence of many different types of attack and defence strategy¹⁹⁵. Coevolution can also be short-lived if antagonists diverge to the point of non-interaction: for instance, if a phage switches host preference away from a focal bacterium¹⁹⁶. Alternatively, a defender might simply develop such a strong defence that an attacker is tolerated³⁶ or driven to extinction¹⁹⁷. Whichever the trajectory it takes, the coevolution of attack and defence, measure and countermeasure, seems to be a major driver of bacterial diversity¹⁹⁸.

[H2] Bacterial competitor countermeasures.

Consistent with the prevalence of inter-bacterial warfare^{2,4,5}, bacteria have numerous adaptations for thwarting the defences of competitors. One solution to the evolution of resistance is for an attacker to innovate new toxins; this selects for attackers with novel toxins, driving diversification of bacterial weapons^{199,200}. Resistant targets may simply select for attackers that produce more toxin¹⁷³, or for those that secrete cocktails of multiple toxins (Figure 5a). Of 102 bacteriocin-producing faecal *E. coli* isolates surveyed in a study in 2006, the majority (58%) produced two or more different bacteriocins²⁰¹; similarly, *P. aeruginosa* releases multiple tailocins and other bacteriocins simultaneously²⁰². A diverse cocktail of toxins may also maintain lethal function over a wider range of environmental conditions, and can benefit from synergistic interactions between toxins²¹. Mirroring antibiotic combination therapy, toxin cocktails may also make resistance less likely to evolve in the first place²⁰³ (Box 1).

A more sophisticated countermeasure is to directly inhibit a defensive mechanism, thereby negating resistance to a particular attack (Figure 5a). The adjuvant Clavulanic acid, which inhibits β -lactamase enzymes, functions in this way: the soil bacterium *Streptomyces clavuligerus* co-regulates clavulanic acid production with the synthesis of the antibiotic cephamycin C, to destroy β -lactamase-protected competitors²⁰⁴. A related approach is to deploy efflux pump inhibitors²⁰⁵ that limit the ability of target bacteria to remove toxins from the cell – another adjuvant countermeasure used in combination with antibiotic therapy²⁰⁶.

Attackers have also evolved ways of surmounting barriers to cell entry (Figure 5b). For example, some bacteria produce 'Trojan Horse' toxins called sideromycins²⁰⁷, which comprise

an antibiotic covalently attached to a siderophore molecule. Siderophores are used by cells to scavenge iron and are imported via dedicated receptors, which enables sideromycins to enter the cell and deliver their antibiotic cargo via the same route²⁰⁸. Some bacterial weapons take a more direct route to toxin translocation: the bacterial T6SS physically punctures target cells, conveying toxins into the target cell without the need to rely upon specific surface receptors or transporter machinery. This direct approach to toxin delivery affords the T6SS a very broad range of target organisms, spanning both Gram-negative and Gram-positive bacteria, fungal cells and other eukaryotes²⁰⁹. Finally, attackers can thwart collective defences (Figure 5c), using proteases and surfactants to disperse biofilm-dwelling bacteria²¹⁰, and quorum-quenching molecules to disrupt intercellular signalling and collective responses, including biofilm formation²¹¹. Additionally, attackers can avoid mass retaliation by deploying 'silent' toxins that are poorly detected by stress responses, thereby suppressing alarm signalling¹⁰¹.

[H2] Phage and predator countermeasures.

Phages have a well-described set of counter-adaptions that enable them to bypass bacterial defences²¹². These adaptations include counter-modification of phage tail fibres, enabling binding of modified cell surface receptors²¹³, and epigenetic modification of phage DNA to mimic the host DNA, thereby escaping degradation via restriction–modification systems²¹⁴. Similarly, defence against restriction (Dar) proteins, injected into hosts by coliphage P1, mask the recognition sites used by restriction enzymes²¹⁵. Some phages encode anti-CRISPR proteins that bind to and inhibit CRISPR–Cas complexes²¹⁶; others boast tail sections with hydrolytic domains, which enables them to penetrate the thick polysaccharide capsules of host cells²¹⁷. Phages also have evolved ways of bypassing bacterial abortive infection mechanisms, thus preventing hosts from interrupting construction of progeny phage¹⁰². For example, coliphage T4 encodes Dmd, an antitoxin 'mimic' that disarms suicide toxins during infection²¹⁸. Finally, paralleling bacterial quorum sensing, some phages use their own 'arbitrium' peptide signal to assess local phage density, transitioning from lytic to lysogenic lifestyles when uninfected hosts become scarce²¹⁹. While not a counter-measure *per se*, this example underlines the sophistication of the responses of phages to their hosts. Meanwhile: though less well-studied, predator adaptations to bacterial defences are also known²²⁰. These include countermeasures to overcome toxin production by prey: mirroring *P. aeruginosa*, the free-living amoeba *Acanthamoeba castellanii* has modified cytochrome oxidases, which enable it to tolerate prey-produced cyanide¹⁶⁹. Some eukaryote predators may also be able to suppress toxin production by prey¹⁶⁹, including via quorum quenching mechanisms²²¹.

[H1] Conclusion

Bacteria have evolved a wide range of defensive adaptations that can make them difficult to kill. Knowledge of these defences has already driven technological revolutions in microbiology and beyond, providing researchers with new tools (restriction enzymes⁵³, CRISPR gene-editing^{56,222} and DNAi/RNAi silencing⁵⁸) and therapeutic approaches (novel antivirals²²³, antimicrobials²²⁴ and biotherapeutics²²⁵). In addition to these applications, defence systems are also central to understanding bacterial biology: they are deeply integrated into their core regulatory networks^{79,81,123}, and can determine which species will persist in a given environment^{4,51,94,172}. Some defences protect only against a particular threat, but many are general and protect against a range of attacks^{144,154,226}. Still others alter bacterial virulence^{120,121,186}, with the potential to exacerbate disease transmission and severity.

These are indeed exciting times for the study of bacterial defences. Spearheaded by bioinformatic¹⁶⁵ and high-throughput²²⁷ approaches, the staggering diversity of bacteria has become clear and with this, the myriad ways they can defend themselves. The past 5 years alone have seen an explosion in the number of novel anti-phage systems identified in bacterial defence islands^{104,109,228} (>50 since 2018), with many more likely awaiting discovery. The diversity and spread of these anti-phage systems highlights how little, in comparison, we know of anti-competitor and anti-predator defences. What might these same approaches teach us about bacterial adaptations against ever-present predator or competitor threats? Early signs are promising: as with the bountiful phage defence islands, anti-competitor defence genes also form clusters in bacterial genomes^{51,132,184}; mining these might therefore reveal novel routes through which bacteria evade rivals' attacks.

As well as examining survival mechanisms, we must understand their broader impact within microbial communities, and the conditions and pathways that trigger them. A major current goal is to control bacteria and their communities, both ecologically and evolutionarily^{3,229,230}. Replacing a pathogen in a community with a biotherapeutic strain²³¹, for example, will require us to understand both the attack and defence strategies of bacteria^{5,10}. And whenever we attempt to eliminate bacteria, whether via antibiotics or one of the emerging alternatives, there is the potential for evolution⁹. As for antibiotic resistance evolution, therefore, the study of how bacterial defences evolve in nature and in the clinic is an important topic for the future.

References

1. Xavier, J. C. *et al.* The metabolic network of the last bacterial common ancestor. *Commun. Biol.* 2021 41 **4**, 1–10 (2021).
2. Foster, K. R. & Bell, T. Competition, not cooperation, dominates interactions among culturable microbial species. *Curr. Biol.* **22**, 1845–1850 (2012).
3. Nadell, C. D., Drescher, K. & Foster, K. R. Spatial structure, cooperation and competition in biofilms. *Nat. Rev. Microbiol.* 2016 149 **14**, 589–600 (2016).
4. Peterson, S. B., Bertolli, S. K. & Mougous, J. D. The Central Role of Interbacterial Antagonism in Bacterial Life. *Curr. Biol.* **30**, R1203–R1214 (2020).
5. Palmer, J. D. & Foster, K. R. Bacterial species rarely work together. *Science* (80-.). **376**, 581–582 (2022).
6. Chevallereau, A., Pons, B. J., van Houte, S. & Westra, E. R. Interactions between bacterial and phage communities in natural environments. *Nat. Rev. Microbiol.* 2021 201 **20**, 49–62 (2021).
7. Pérez, J., Moraleda-Muñoz, A., Marcos-Torres, F. J. & Muñoz-Dorado, J. Bacterial predation: 75 years and counting! *Environmental Microbiology* **18**, 766–779 (2016).
8. Wein, T. & Sorek, R. Bacterial origins of human cell-autonomous innate immune mechanisms. *Nat. Rev. Immunol.* **22**, (2022).
9. Brockhurst, M. A. *et al.* Assessing evolutionary risks of resistance for new antimicrobial therapies. *nature.com*
10. Granato, E. T., Meiller-Legrand, T. A. & Foster, K. R. The Evolution and Ecology of Bacterial Warfare. *Curr. Biol.* **29**, R521–R537 (2019).
11. Clardy, J., Fischbach, M. A. & Currie, C. R. The natural history of antibiotics. *Current Biology* **19**, R437 (2009).
12. M Hassan, M. K. I. N. D. D. F. L. Natural antimicrobial peptides from bacteria: characteristics and potential applications to fight against antibiotic resistance. *J. Appl. Microbiol.* **113**, 723–736 (2012).
13. Riley, M. A. & Wertz, J. E. Bacteriocin diversity: ecological and evolutionary perspectives. *Biochimie* **84**, 357–364 (2002).
14. Vacheron, J., Heiman, C. M. & Keel, C. Live cell dynamics of production, explosive release and killing activity of phage tail-like weapons for *Pseudomonas* kin exclusion. *Commun. Biol.* 2021 41 **4**, 1–14 (2021).
15. Ho, B. T., Dong, T. G. & Mekalanos, J. J. A View to a Kill: The Bacterial Type VI Secretion System. *Cell Host Microbe* **15**, 9–21 (2014).
16. Nakayama, K. *et al.* The R-type pyocin of *Pseudomonas aeruginosa* is related to P2 phage, and the F-type is related to lambda phage. *Mol. Microbiol.* **38**, 213–231

641 (2000).

642 17. Davies, E. V. *et al.* Temperate phages both mediate and drive adaptive evolution in
643 pathogen biofilms. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 8266–8271 (2016).

644 18. Sharp, C., Bray, J., Housden, N. G., Maiden, M. C. J. & Kleanthous, C. Diversity and
645 distribution of nuclease bacteriocins in bacterial genomes revealed using Hidden
646 Markov Models. *PLOS Comput. Biol.* **13**, e1005652 (2017).

647 19. Cascales, E. *et al.* Colicin biology. *Microbiol. Mol. Biol. Rev.* **71**, 158–229 (2007).

648 20. Chikindas, M. L., Weeks, R., Drider, D., Chistyakov, V. A. & Dicks, L. M. Functions
649 and emerging applications of bacteriocins. *Current Opinion in Biotechnology* **49**, 23–
650 28 (2018).

651 21. LaCourse, K. D. *et al.* Conditional toxicity and synergy drive diversity among
652 antibacterial effectors. *Nat. Microbiol.* **3**, 440–446 (2018).

653 22. Smith, W. P. J. *et al.* The evolution of the type VI secretion system as a disintegration
654 weapon. *PLoS Biol.* **18**, e3000720 (2020).

655 23. Kohanski, M. A., Dwyer, D. J. & Collins, J. J. How antibiotics kill bacteria: From
656 targets to networks. *Nature Reviews Microbiology* **8**, 423–435 (2010).

657 24. Boolchandani, M., D’Souza, A. W. & Dantas, G. Sequencing-based methods and
658 resources to study antimicrobial resistance. *Nature Reviews Genetics* **20**, 356–370
659 (2019).

660 25. Rohwer, F. Global phage diversity. *Cell* **113**, 141 (2003).

661 26. Suttle, C. A. The significance of viruses to mortality in aquatic microbial communities.
662 *Microb. Ecol. 1994 282* **28**, 237–243 (1994).

663 27. Secor, P. R. & Dandekar, A. A. More than simple parasites: The sociobiology of
664 bacteriophages and their bacterial hosts. *mBio* **11**, (2020).

665 28. Howard-Varona, C., Hargreaves, K. R., Abedon, S. T. & Sullivan, M. B. Lysogeny in
666 nature: mechanisms, impact and ecology of temperate phages. *ISME J.* **11**, 1511–
667 1520 (2017).

668 29. Clokie, M. R. J., Millard, A. D., Letarov, A. V. & Heaphy, S. Phages in nature.
669 *Bacteriophage* **1**, 31–45 (2011).

670 30. Fernandes, S. & São-José, C. Enzymes and mechanisms employed by tailed
671 bacteriophages to breach the bacterial cell barriers. *Viruses* **10**, 396 (2018).

672 31. Sausset, R., Petit, M. A., Gaboriau-Routhiau, V. & De Paepe, M. New insights into
673 intestinal phages. *Mucosal Immunol.* **13**, 205–215 (2020).

674 32. Muñoz-Dorado, J., Marcos-Torres, F. J., García-Bravo, E., Moraleda-Muñoz, A. &
675 Pérez, J. Myxobacteria: Moving, killing, feeding, and surviving together. *Frontiers in*
676 *Microbiology* **7**, (2016).

677 33. Laloux, G. Shedding Light on the Cell Biology of the Predatory Bacterium *Bdellovibrio*

- bacteriovorus. *Front. Microbiol.* **10**, 3136 (2020).
34. Bratanis, E., Andersson, T., Lood, R. & Bukowska-Faniband, E. Biotechnological Potential of Bdellovibrio and Like Organisms and Their Secreted Enzymes. *Front. Microbiol.* **11**, 662 (2020).
35. Danczak, R. E. *et al.* Members of the Candidate Phyla Radiation are functionally differentiated by carbon-and nitrogen-cycling capabilities. doi:10.1186/s40168-017-0331-1
36. Bor, B. *et al.* Rapid evolution of decreased host susceptibility drives a stable relationship between ultrasmall parasite TM7x and its bacterial host. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 12277–12282 (2018).
37. Blair, J. M. A., Webber, M. A., Baylay, A. J., Ogbolu, D. O. & Piddock, L. J. V. Molecular mechanisms of antibiotic resistance. *Nature Reviews Microbiology* **13**, 42–51 (2015).
38. Wilson, D. N., Haurlyiuk, V., Atkinson, G. C. & O'Neill, A. J. Target protection as a key antibiotic resistance mechanism. *Nature Reviews Microbiology* 1–12 (2020). doi:10.1038/s41579-020-0386-z
39. Larsen, J. *et al.* Emergence of methicillin resistance predates the clinical use of antibiotics. *Nat. 2022 6027895* **602**, 135–141 (2022).
40. Fan, X. Y. *et al.* Oxidation of dCTP contributes to antibiotic lethality in stationary-phase mycobacteria. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 2210–2215 (2018).
41. Dong, T. G. *et al.* Generation of reactive oxygen species by lethal attacks from competing microbes. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 2181–2186 (2015).
42. Wozniak, K. J. & Simmons, L. A. Bacterial DNA excision repair pathways. *Nat. Rev. Microbiol.* **20**, 465–477 (2022).
43. Kisker, C., Kuper, J. & Van Houten, B. Prokaryotic Nucleotide Excision Repair. *Cold Spring Harb. Perspect. Biol.* **5**, (2013).
44. TP, M. *et al.* RtcB2-PrfH Operon Protects E. coli ATCC25922 Strain from Colicin E3 Toxin. *Int. J. Mol. Sci.* **23**, (2022).
45. Joly, N. *et al.* Managing membrane stress: The phage shock protein (Psp) response, from molecular mechanisms to physiology. *FEMS Microbiology Reviews* **34**, 797–827 (2010).
46. Hersch, S. J. *et al.* Envelope stress responses defend against type six secretion system attacks independently of immunity proteins. *Nat. Microbiol.* **5**, 706–714 (2020).
47. Munita, J. M. & Arias, C. A. Mechanisms of Antibiotic Resistance. doi:10.1128/microbiolspec.VMBF-0016-2015
48. Yang, X., Long, M. & Shen, X. Effector–Immunity Pairs Provide the T6SS Nanomachine its Offensive and Defensive Capabilities. *Molecules* **23**, 1009 (2018).

- 715 49. Aoki, S. K. *et al.* A widespread family of polymorphic contact-dependent toxin delivery
716 systems in bacteria. *Nature* **468**, 439–442 (2010).
- 717 50. Coyne, M. J., Zitomersky, N. L., McGuire, A. M., Earl, A. M. & Comstock, L. E.
718 Evidence of extensive DNA transfer between bacteroidales species within the human
719 gut. *MBio* **5**, (2014).
- 720 51. Ross, B. D. *et al.* Human gut bacteria contain acquired interbacterial defence
721 systems. *224 | Nat. |* **575**, (2019).
- 722 52. Bush, K. Past and present perspectives on β -lactamases. *Antimicrobial Agents and*
723 *Chemotherapy* **62**, (2018).
- 724 53. Tock, M. R. & Dryden, D. T. F. The biology of restriction and anti-restriction. *Current*
725 *Opinion in Microbiology* **8**, 466–472 (2005).
- 726 54. Ofir, G. *et al.* DISARM is a widespread bacterial defence system with broad anti-
727 phage activities. *Nat. Microbiol.* **3**, 90–98 (2018).
- 728 55. Wang, L., Jiang, S., Deng, Z., Dedon, P. C. & Chen, S. DNA phosphorothioate
729 modification-a new multi-functional epigenetic system in bacteria. *FEMS Microbiol.*
730 *Rev.* **43**, 109–122 (2019).
- 731 56. Barrangou, R. *et al.* CRISPR provides acquired resistance against viruses in
732 prokaryotes. *Science (80-.).* **315**, 1709–1712 (2007).
- 733 57. Westra, E. R. & Levin, B. R. It is unclear how important CRISPR-Cas systems are for
734 protecting natural populations of bacteria against infections by mobile genetic
735 elements. *Proceedings of the National Academy of Sciences of the United States of*
736 *America* **117**, 27777–27785 (2020).
- 737 58. Lisitskaya, L., Aravin, A. A. & Kulbachinskiy, A. DNA interference and beyond:
738 structure and functions of prokaryotic Argonaute proteins. *Nature Communications* **9**,
739 1–12 (2018).
- 740 59. Gupta, R. S. Origin of diderm (Gram-negative) bacteria: Antibiotic selection pressure
741 rather than endosymbiosis likely led to the evolution of bacterial cells with two
742 membranes. *Antonie van Leeuwenhoek, International Journal of General and*
743 *Molecular Microbiology* **100**, 171–182 (2011).
- 744 60. Dy, R. L., Richter, C., Salmond, G. P. C. & Fineran, P. C. Remarkable Mechanisms in
745 Microbes to Resist Phage Infections. *Annu. Rev. Virol.* **1**, 307–331 (2014).
- 746 61. Vidakovic, L., Singh, P. K., Hartmann, R., Nadell, C. D. & Drescher, K. Dynamic
747 biofilm architecture confers individual and collective mechanisms of viral protection.
748 *Nat. Microbiol.* **3**, 26–31 (2017).
- 749 62. Toska, J., Ho, B. T. & Mekalanos, J. J. Exopolysaccharide protects *Vibrio cholerae*
750 from exogenous attacks by the type 6 secretion system. *Proc. Natl. Acad. Sci.* **115**,
751 7997–8002 (2018).

63. Flaugnatti, N. *et al.* Human commensal gut Proteobacteria withstand type VI secretion attacks through immunity protein-independent mechanisms. *Nat. Commun.* **2021** 121 **12**, 1–13 (2021).
64. Bharat, T. A. M. *et al.* Structure of the hexagonal surface layer on *Caulobacter crescentus* cells. *Nat. Microbiol.* **2**, 17059 (2017).
65. Koval, S. F. & Hynes, S. H. Effect of paracrystalline protein surface layers on predation by *Bdellovibrio bacteriovorus*. *J. Bacteriol.* **173**, 2244–2249 (1991).
66. Duncan, M. C. *et al.* *Vibrio cholerae* motility exerts drag force to impede attack by the bacterial predator *Bdellovibrio bacteriovorus*. *Nat. Commun.* **9**, (2018).
67. Gao, L. & van der Veen, S. Role of Outer Membrane Vesicles in Bacterial Physiology and Host Cell Interactions. *Infect. Microbes Dis.* **2**, 3–9 (2020).
68. Manning, A. J. & Kuehn, M. J. Contribution of bacterial outer membrane vesicles to innate bacterial defense. *BMC Microbiol.* **11**, (2011).
69. Webber, M. A. The importance of efflux pumps in bacterial antibiotic resistance. *J. Antimicrob. Chemother.* **51**, 9–11 (2003).
70. Saier, M. H. *et al.* Evolutionary origins of multidrug and drug-specific efflux pumps in bacteria. *FASEB J.* **12**, 265–274 (1998).
71. Du, D. *et al.* Multidrug efflux pumps: structure, function and regulation. *Nat. Rev. Microbiol.* **16**, 523–539 (2018).
72. Tahlan, K. *et al.* Initiation of actinorhodin export in *Streptomyces coelicolor*. *Mol. Microbiol.* **63**, 951–961 (2007).
73. Wadhwa, N. & Berg, H. C. Bacterial motility: machinery and mechanisms. *Nat. Rev. Microbiol.* **2021** 203 **20**, 161–173 (2021).
74. Matz, C. & Jürgens, K. High Motility Reduces Grazing Mortality of Planktonic Bacteria Downloaded from. *Appl. Environ. Microbiol.* **71**, 921–929 (2005).
75. Lambert, C. *et al.* Characterizing the flagellar filament and the role of motility in bacterial prey-penetration by *Bdellovibrio bacteriovorus*. *Mol. Microbiol.* **60**, 274–286 (2006).
76. Bertozzi Silva, J., Storms, Z. & Sauvageau, D. Host receptors for bacteriophage adsorption. *FEMS Microbiol. Lett.* **363**, 2 (2016).
77. Li, X., Gonzalez, F., Esteves, N., Scharf, B. E. & Chen, J. Formation of phage lysis patterns and implications on co-propagation of phages and motile host bacteria. *PLOS Comput. Biol.* **16**, e1007236 (2020).
78. Granato, E. T., Smith, W. P. J. & Foster, K. R. Collective protection against the type VI secretion system in bacteria. *bioRxiv* 2022.09.12.507624 (2022). doi:10.1101/2022.09.12.507624
79. Cornforth, D. M. & Foster, K. R. *Competition sensing: the social side of bacterial*

stress responses. (2013). doi:10.1038/nrmicro2977

80. Oliveira, N. M. *et al.* Biofilm formation as a response to ecological competition. *PLoS Biol.* **13**, (2015).

81. Lories, B. *et al.* Biofilm Bacteria Use Stress Responses to Detect and Respond to Competitors. *Curr. Biol.* **30**, 1231-1244.e4 (2020).

82. Sadiq, F. A. *et al.* Phenotypic and genetic heterogeneity within biofilms with particular emphasis on persistence and antimicrobial tolerance. *Future Microbiol.* **12**, 1087–1107 (2017).

83. Krishna Kumar, R. *et al.* Droplet printing reveals the importance of micron-scale structure for bacterial ecology. *Nat. Commun.* **12**, 1–12 (2021).

84. Bagge, N. *et al.* Dynamics and Spatial Distribution of β -Lactamase Expression in *Pseudomonas aeruginosa* Biofilms. *Antimicrob. Agents Chemother.* **48**, 1168–1174 (2004).

85. Billings, N. *et al.* The Extracellular Matrix Component Psl Provides Fast-Acting Antibiotic Defense in *Pseudomonas aeruginosa* Biofilms. *PLOS Pathog.* **9**, e1003526 (2013).

86. Wucher, B. R., Elsayed, M., Adelman, J. S., Kadouri, D. E. & Nadell, C. D. Bacterial predation transforms the landscape and community assembly of biofilms. *Curr. Biol.* (2021). doi:10.1016/j.cub.2021.03.036

87. Xavier, J. B. & Foster, K. R. Cooperation and conflict in microbial biofilms. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 876–881 (2007).

88. Bond, M. C., Vidakovic, L., Singh, P. K., Drescher, K. & Nadell, C. D. Matrix-trapped viruses can prevent invasion of bacterial biofilms by colonizing cells. *Elife* **10**, (2021).

89. Arnoldini, M. *et al.* Bistable Expression of Virulence Genes in *Salmonella* Leads to the Formation of an Antibiotic-Tolerant Subpopulation. *PLOS Biol.* **12**, e1001928 (2014).

90. Łapińska, U. *et al.* Fast bacterial growth reduces antibiotic accumulation and efficacy. *Elife* **11**, (2022).

91. Williamson, K. S. *et al.* Heterogeneity in *Pseudomonas aeruginosa* biofilms includes expression of ribosome hibernation factors in the antibiotic-tolerant subpopulation and hypoxia-induced stress response in the metabolically active population. *J. Bacteriol.* **194**, 2062–2073 (2012).

92. Balaban, N. Q., Merrin, J., Chait, R., Kowalik, L. & Leibler, S. Bacterial persistence as a phenotypic switch. *Science (80-.).* **305**, 1622–1625 (2004).

93. Van Den Bergh, B. *et al.* Frequency of antibiotic application drives rapid evolutionary adaptation of *Escherichia coli* persistence. *Nat. Microbiol.* **1**, (2016).

94. Drebes Dörr, N. C. & Blokesch, M. Interbacterial competition and anti-predatory behaviour of environmental *Vibrio cholerae* strains. *Environ. Microbiol.* **22**, 4485–4504

(2020).

95. Basler, M., Ho, B. T. & Mekalanos, J. J. Tit-for-Tat: Type VI Secretion System Counterattack during Bacterial Cell-Cell Interactions. *Cell* **152**, 884–894 (2013).
96. Kamal, F. *et al.* Differential Cellular Response to Translocated Toxic Effectors and Physical Penetration by the Type VI Secretion System. *Cell Rep.* **31**, 107766 (2020).
97. Rendueles, O., Amherd, M. & Velicer, G. J. Positively Frequency-Dependent Interference Competition Maintains Diversity and Pervades a Natural Population of Cooperative Microbes. *Curr. Biol.* **25**, 1673–1681 (2015).
98. Gordon, D. M. & Riley, M. A. A theoretical and empirical investigation of the invasion dynamics of colicinogeny. *Microbiology* **145**, 655–661 (1999).
99. Booth, S. C., Smith, W. P. J. & Foster, K. R. Bows and swords: why bacteria carry short and long-range weapons. *bioRxiv* 2022.10.13.512033 (2022).
doi:10.1101/2022.10.13.512033
100. Mavridou, D. A. I., Gonzalez, D., Kim, W., West, S. A. & Correspondence, K. R. F. Bacteria Use Collective Behavior to Generate Diverse Combat Strategies. *Curr. Biol.* **28**, 345-355.e4 (2018).
101. Gonzalez, D., Sabnis, A., Foster, K. R. & Mavridou, D. A. I. Costs and benefits of provocation in bacterial warfare. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 7593–7598 (2018).
102. Lopatina, A., Tal, N. & Sorek, R. Abortive Infection: Bacterial Suicide as an Antiviral Immune Strategy. *Annual Review of Virology* **7**, 371–384 (2020).
103. Johnson, A. G. *et al.* Bacterial gasdermins reveal an ancient mechanism of cell death. *Science* **375**, 221–225 (2022).
104. Cohen, D. *et al.* Cyclic GMP–AMP signalling protects bacteria against viral infection. *Nat.* 2019 5747780 **574**, 691–695 (2019).
105. Koga, M., Otsuka, Y., Lemire, S. & Yonesaki, T. Escherichia coli rnlA and rnlB Compose a Novel Toxin–Antitoxin System. *Genetics* **187**, 123–130 (2011).
106. Watson, B. N. J. *et al.* Type I-F CRISPR-Cas resistance against virulent phages results in abortive infection and provides population-level immunity. *Nat. Commun.* 2019 101 **10**, 1–8 (2019).
107. Gao, L. *et al.* Diverse Enzymatic Activities Mediate Antiviral Immunity in Prokaryotes. *Science* **369**, 1077 (2020).
108. Ka, D., Oh, H., Park, E., Kim, J. H. & Bae, E. Structural and functional evidence of bacterial antiphage protection by Thoeris defense system via NAD⁺ degradation. *Nat. Commun.* **11**, (2020).
109. Doron, S. *et al.* Systematic discovery of antiphage defense systems in the microbial pangenome. *Science (80-.).* **359**, eaar4120 (2018).

- 863 110. Geller, A. M. *et al.* The extracellular contractile injection system is enriched in
864 environmental microbes and associates with numerous toxins. *Nat. Commun.* 2021
865 121 **12**, 1–15 (2021).
- 866 111. Granato, E. T. & Foster, K. R. The Evolution of Mass Cell Suicide in Bacterial
867 Warfare. *Curr. Biol.* **30**, 2836-2843.e3 (2020).
- 868 112. Niehus, R., Oliveira, N. M., Li, A., Fletcher, A. G. & Foster, K. R. The evolution of
869 strategy in bacterial warfare via the regulation of bacteriocins and antibiotics.
870 doi:10.7554/eLife.69756
- 871 113. Boor, K. J. Bacterial stress responses: What doesn't kill them can make them
872 stronger. *PLoS Biology* **4**, 0018–0020 (2006).
- 873 114. Shin, J. H., Singh, A. K., Cheon, D. J. & Roe, J. H. Activation of the SoxR regulon in
874 *Streptomyces coelicolor* by the extracellular form of the pigmented antibiotic
875 actinorhodin. *J. Bacteriol.* **193**, 75–81 (2011).
- 876 115. Smith, W. P. J. *et al.* The evolution of tit-for-tat in bacteria via the type VI secretion
877 system. *Nat. Commun.* **11**, (2020).
- 878 116. Wang, G. Z. *et al.* Staphylococcal secreted cytotoxins are competition sensing signals
879 for *Pseudomonas aeruginosa*. *bioRxiv* 2023.01.29.526047 (2023).
880 doi:10.1101/2023.01.29.526047
- 881 117. Flores-Kim, J. & Darwin, A. J. The Phage Shock Protein Response. *Annu. Rev.*
882 *Microbiol.* **70**, 83–101 (2016).
- 883 118. Fallico, V., Ross, R. P., Fitzgerald, G. F. & McAuliffe, O. Genetic response to
884 bacteriophage infection in *Lactococcus lactis* reveals a four-strand approach involving
885 induction of membrane stress proteins, D-alanylation of the cell wall, maintenance of
886 proton motive force, and energy conservation. *J. Virol.* **85**, 12032–12042 (2011).
- 887 119. Song, S. & Wood, T. K. A Primary Physiological Role of Toxin/Antitoxin Systems Is
888 Phage Inhibition. *Front. Microbiol.* **11**, 1895 (2020).
- 889 120. Aijaz, I. & Koudelka, G. B. Cheating, facilitation and cooperation regulate the
890 effectiveness of phage-encoded exotoxins as antipredator molecules.
891 *Microbiologyopen* **8**, e00636 (2019).
- 892 121. Pacheco, A. R. & Sperandio, V. Shiga toxin in enterohemorrhagic *E. coli*: regulation
893 and novel anti-virulence strategies. *Front. Cell. Infect. Microbiol.* **2**, 81 (2012).
- 894 122. Lambert, C., Ivanov, P. & Sockett, R. E. A transcriptional 'scream' early response of
895 *E. coli* prey to predatory invasion by *Bdellovibrio*. *Curr. Microbiol.* **60**, 419–427 (2010).
- 896 123. LeRoux, M., Peterson, S. B. & Mougous, J. D. Bacterial danger sensing. *Journal of*
897 *Molecular Biology* **427**, 3744–3753 (2015).
- 898 124. Bertsche, U., Mayer, C., Götz, F. & Gust, A. A. Peptidoglycan perception—Sensing
899 bacteria by their common envelope structure. *Int. J. Med. Microbiol.* **305**, 217–223

900 (2015).

901 125. Diggle, S. P., Gardner, A., West, S. A. & Griffin, A. S. Evolutionary theory of bacterial
 902 quorum sensing: when is a signal not a signal? *Philos. Trans. R. Soc. B Biol. Sci.* **362**,
 903 1241–1249 (2007).

904 126. Pohnert, G., Steinke, M. & Tollrian, R. Chemical cues, defence metabolites and the
 905 shaping of pelagic interspecific interactions. *Trends Ecol. Evol.* **22**, 198–204 (2007).

906 127. Mazzola, M., De Bruijn, I., Cohen, M. F. & Raaijmakers, J. M. Protozoan-induced
 907 regulation of cyclic lipopeptide biosynthesis is an effective predation defense
 908 mechanism for *Pseudomonas fluorescens*. *Appl. Environ. Microbiol.* **75**, 6804–6811
 909 (2009).

910 128. Li, L. *et al.* Sensor histidine kinase is a β -lactam receptor and induces resistance to β -
 911 lactam antibiotics. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 1648–1653 (2016).

912 129. Bru, J. L. *et al.* PQS produced by the *pseudomonas aeruginosa* stress response
 913 repels swarms away from bacteriophage and antibiotics. *J. Bacteriol.* **201**, (2019).

914 130. Tzipilevich, E., Pollak-Fiyaksel, O., Shraiteh, B. & Ben-Yehuda, S. Bacteria elicit a
 915 phage tolerance response subsequent to infection of their neighbors. *EMBO J.* **41**,
 916 (2022).

917 131. LeRoux, M. *et al.* Kin cell lysis is a danger signal that activates antibacterial pathways
 918 of *pseudomonas aeruginosa*. *Elife* **2015**, 1–65 (2015).

919 132. Ting, S. Y. *et al.* Discovery of coordinately regulated pathways that provide innate
 920 protection against interbacterial antagonism. *Elife* **11**, (2022).

921 133. Irving, S. E., Choudhury, N. R. & Corrigan, R. M. The stringent response and
 922 physiological roles of (pp)pGpp in bacteria. *Nat. Rev. Microbiol.* **2020** *194* **19**, 256–
 923 271 (2020).

924 134. Inaoka, T., Takahashi, K., Ohnishi-Kameyama, M., Yoshida, M. & Ochi, K. Guanine
 925 Nucleotides Guanosine 5'-Diphosphate 3'-Diphosphate and GTP Co-operatively
 926 Regulate the Production of an Antibiotic Bacilysin in *Bacillus subtilis* *. *J. Biol. Chem.*
 927 **278**, 2169–2176 (2003).

928 135. Kuhar, I. & Žgur-Bertok, D. Transcription regulation of the colicin K cka gene reveals
 929 induction of colicin synthesis by differential responses to environmental signals. *J.*
 930 *Bacteriol.* **181**, 7373–7380 (1999).

931 136. Ochi, K. Occurrence of the stringent response in *Streptomyces* sp. and its significance
 932 for the initiation of morphological and physiological differentiation. *J. Gen. Microbiol.*
 933 **132**, 2621–2631 (1986).

934 137. Hammer, B. K. & Bassler, B. L. Quorum sensing controls biofilm formation in *Vibrio*
 935 *cholerae*. *Mol. Microbiol.* **50**, 101–104 (2003).

936 138. Rutherford, S. T. & Bassler, B. L. Bacterial Quorum Sensing: Its Role in Virulence and

- Possibilities for Its Control. *Cold Spring Harb. Perspect. Med.* **2**, (2012).
139. Høyland-Kroghsbo, N. M. *et al.* Quorum sensing controls the pseudomonas aeruginosa CRISPR-Cas adaptive immune system. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 131–135 (2017).
140. Zhou, L., Zhang, Y., Ge, Y., Zhu, X. & Pan, J. Regulatory Mechanisms and Promising Applications of Quorum Sensing-Inhibiting Agents in Control of Bacterial Biofilm Formation. *Front. Microbiol.* **11**, 2558 (2020).
141. Takano, E. Gamma-butyrolactones: Streptomyces signalling molecules regulating antibiotic production and differentiation. *Curr. Opin. Microbiol.* **9**, 287–294 (2006).
142. Fontaine, L. *et al.* Quorum-Sensing Regulation of the Production of Blp Bacteriocins in Streptococcus thermophilus. *J. Bacteriol.* **189**, 7195 (2007).
143. Silverman, J. M., Brunet, Y. R., Cascales, E. & Mougous, J. D. Structure and Regulation of the Type VI Secretion System. *Annu. Rev. Microbiol.* **66**, 453 (2012).
144. Meirelles, L. A., Perry, E. K., Bergkessel, M. & Newman, D. K. Bacterial defenses against a natural antibiotic promote collateral resilience to clinical antibiotics. *PLoS Biol.* **19**, e3001093 (2021).
145. Charles Darwin. *On the Origin of Species: A Facsimile of the First Edition*. (Harvard University Press, 2003).
146. Raeside, C. *et al.* Large Chromosomal Rearrangements during a Long-Term Evolution Experiment with Escherichia coli. *MBio* **5**, 1377–1391 (2014).
147. Frazão, N., Sousa, A., Lässig, M. & Gordo, I. Horizontal gene transfer overrides mutation in Escherichia coli colonizing the mammalian gut. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 17906–17915 (2019).
148. Woodford, N. & Ellington, M. J. The emergence of antibiotic resistance by mutation. *Clin. Microbiol. Infect.* **13**, 5–18 (2007).
149. Macwana, S. & Muriana, P. M. Spontaneous bacteriocin resistance in Listeria monocytogenes as a susceptibility screen for identifying different mechanisms of resistance and modes of action by bacteriocins of lactic acid bacteria. *J. Microbiol. Methods* **88**, 7–13 (2012).
150. Riley, M. A. & Gordon, D. M. The ecology and evolution of bacteriocins. *J. Ind. Microbiol. Biotechnol.* **17**, 151–158 (1996).
151. Wright, R. C. T., Friman, V.-P., Smith, M. C. M. & Brockhurst, M. A. Cross-resistance is modular in bacteria–phage interactions. *PLOS Biol.* **16**, e2006057 (2018).
152. Abouzeed, Y. M., Baucheron, S. & Cloeckaert, A. ramR mutations involved in efflux-mediated multidrug resistance in Salmonella enterica serovar Typhimurium. *Antimicrob. Agents Chemother.* **52**, 2428–2434 (2008).
153. Cannatelli, A. *et al.* MgrB inactivation is a common mechanism of colistin resistance in

974 KPC-producing *klebsiella pneumoniae* of clinical origin. *Antimicrob. Agents*
975 *Chemother.* **58**, 5696–5703 (2014).

976 154. Koch, G. *et al.* Evolution of resistance to a last-resort antibiotic in *staphylococcus*
977 *aureus* via bacterial competition. *Cell* **158**, 1060–1071 (2014).

978 155. Bjedov, I. *et al.* Stress-induced mutagenesis in bacteria. *Science (80-.)*. **300**, 1404–
979 1409 (2003).

980 156. Krašovec, R. *et al.* Opposing effects of final population density and stress on
981 *Escherichia coli* mutation rate. *ISME J. 2018 12 12* **12**, 2981–2987 (2018).

982 157. Ram, Y. & Hadany, L. Evolution of stress-induced mutagenesis in the presence of
983 horizontal gene transfer. *Am. Nat.* **194**, 73–89 (2019).

984 158. MacLean, R. C. & San Millan, A. The evolution of antibiotic resistance. *Science (80-.)*. **365**, 1082–1083 (2019).

985
986 159. Hall, J. P. J., Brockhurst, M. A. & Harrison, E. Sampling the mobile gene pool:
987 innovation via horizontal gene transfer in bacteria. *Philos. Trans. R. Soc. B Biol. Sci.*
988 **372**, (2017).

989 160. Meaden, S. & Fineran, P. C. Bacterial defense islands limit viral attack. *Science (80-.)*. **374**, 399–400 (2021).

990
991 161. Didelot, X. & Maiden, M. C. J. Impact of recombination on bacterial evolution. *Trends*
992 *Microbiol.* **18**, 315 (2010).

993 162. San Millan, A. Evolution of Plasmid-Mediated Antibiotic Resistance in the Clinical
994 Context. *Trends Microbiol.* **26**, 978–985 (2018).

995 163. Botelho, J. & Schulenburg, H. The Role of Integrative and Conjugative Elements in
996 Antibiotic Resistance Evolution. *Trends in Microbiology* **29**, 8–18 (2021).

997 164. Ruhe, Z. C. *et al.* CDI Systems Are Stably Maintained by a Cell-Contact Mediated
998 Surveillance Mechanism. *PLoS Genet.* **12**, e1006145 (2016).

999 165. Makarova, K. S., Wolf, Y. I., Snir, S. & Koonin, E. V. Defense Islands in Bacterial and
1000 Archaeal Genomes and Prediction of Novel Defense Systems. *J. Bacteriol.* **193**,
1001 6039–6056 (2011).

1002 166. Bernheim, A. & Sorek, R. The pan-immune system of bacteria: antiviral defence as a
1003 community resource. *Nat. Rev. Microbiol. 2019 182* **18**, 113–119 (2019).

1004 167. Rocha, E. P. C. & Bikard, D. Microbial defenses against mobile genetic elements and
1005 viruses: Who defends whom from what? *PLOS Biol.* **20**, e3001514 (2022).

1006 168. Costa, R., Van Aarle, I. M., Mendes, R. & Van Elsas, J. D. Genomics of pyrrolnitrin
1007 biosynthetic loci: Evidence for conservation and whole-operon mobility within Gram-
1008 negative bacteria. *Environ. Microbiol.* **11**, 159–175 (2009).

1009 169. Jousset, A., Rochat, L., Scheu, S., Bonkowski, M. & Keel, C. Predator-prey chemical
1010 warfare determines the expression of biocontrol genes by rhizosphere-associated

1011 *pseudomonas fluorescens*. *Appl. Environ. Microbiol.* **76**, 5263–5268 (2010).

1012 170. Alexander, H. K. & Craig MacLean, R. Stochastic bacterial population dynamics
 1013 restrict the establishment of antibiotic resistance from single cells. *Proc. Natl. Acad.*
 1014 *Sci. U. S. A.* **117**, 19455–19464 (2020).

1015 171. Herron, M. D. *et al.* De novo origins of multicellularity in response to predation. *Sci.*
 1016 *Reports* 2019 91 **9**, 1–9 (2019).

1017 172. Koskella, B. & Brockhurst, M. A. Bacteria–phage coevolution as a driver of ecological
 1018 and evolutionary processes in microbial communities. *Fems Microbiol. Rev.* **38**, 916
 1019 (2014).

1020 173. Libberton, B., Horsburgh, M. J. & Brockhurst, M. A. The effects of spatial structure,
 1021 frequency dependence and resistance evolution on the dynamics of toxin-mediated
 1022 microbial invasions. *Evol. Appl.* **8**, 738–750 (2015).

1023 174. Butela, K. & Lawrence, J. Population Genetics of Salmonella: Selection for Antigenic
 1024 Diversity. in *Bacterial Population Genetics in Infectious Disease* (eds. D. Ashley
 1025 Robinson, Daniel Falush & Edward J. Feil) 287–319 (John Wiley & Sons, Ltd, 2010).
 1026 doi:10.1002/9780470600122.CH15

1027 175. Wildschutte, H., Wolfe, D. M., Tamewitz, A. & Lawrence, J. G. Protozoan predation,
 1028 diversifying selection, and the evolution of antigenic diversity in Salmonella. *Proc.*
 1029 *Natl. Acad. Sci. U. S. A.* **101**, 10644–10649 (2004).

1030 176. Mostowy, R. J. & Holt, K. E. Diversity-Generating Machines: Genetics of Bacterial
 1031 Sugar-Coating. *Trends Microbiol.* **26**, 1008–1021 (2018).

1032 177. Burmeister, A. R. *et al.* Pleiotropy complicates a trade-off between phage resistance
 1033 and antibiotic resistance. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 11207–11216 (2020).

1034 178. German, G. J. & Misra, R. The TolC protein of Escherichia coli serves as a cell-
 1035 surface receptor for the newly characterized TLS bacteriophage. *J. Mol. Biol.* **308**,
 1036 579–585 (2001).

1037 179. Andersson, D. I. & Hughes, D. Antibiotic resistance and its cost: Is it possible to
 1038 reverse resistance? *Nature Reviews Microbiology* **8**, 260–271 (2010).

1039 180. Puigbò, P., Makarova, K. S., Kristensen, D. M., Wolf, Y. I. & Koonin, E. V.
 1040 Reconstruction of the evolution of microbial defense systems. doi:10.1186/s12862-
 1041 017-0942-y

1042 181. Rowe-Magnus, D. A. & Mazel, D. The role of integrons in antibiotic resistance gene
 1043 capture. *Int. J. Med. Microbiol.* **292**, 115–125 (2002).

1044 182. Akrami, F., Rajabnia, M. & Pournajaf, A. Resistance integrons; A mini review. *Caspian*
 1045 *Journal of Internal Medicine* **10**, 370–376 (2019).

1046 183. Hocquet, D. *et al.* Evidence for induction of integron-based antibiotic resistance by the
 1047 SOS response in a clinical setting. *PLoS Pathog.* **8**, 1002778 (2012).

1048 184. Souque, C., Escudero, J. A. & Maclean, R. C. Integron activity accelerates the
1049 evolution of antibiotic resistance. *Elife* **10**, 1–47 (2021).

1050 185. Oliveira, P. H., Touchon, M. & Rocha, E. P. C. The interplay of restriction-modification
1051 systems with mobile genetic elements and their prokaryotic hosts. *Nucleic Acids Res.*
1052 **42**, 10618–10631 (2014).

1053 186. Alcalde-Rico, M., Hernando-Amado, S., Blanco, P. & Martínez, J. Multidrug efflux
1054 pumps at the crossroad between antibiotic resistance and bacterial virulence. *Front.*
1055 *Microbiol.* **7**, 1483 (2016).

1056 187. van Houte, S., Buckling, A. & Westra, E. R. Evolutionary Ecology of Prokaryotic
1057 Immune Mechanisms. *Microbiol. Mol. Biol. Rev.* **80**, 745–763 (2016).

1058 188. Avrani, S., Wurtzel, O., Sharon, I., Sorek, R. & Lindell, D. Genomic island variability
1059 facilitates *Prochlorococcus*–virus coexistence. *Nat. 2011 4747353* **474**, 604–608
1060 (2011).

1061 189. Cordero, O. X. *et al.* Ecological populations of bacteria act as socially cohesive units
1062 of antibiotic production and resistance. *Science (80-.)*. **337**, 1228–1231 (2012).

1063 190. Hussain, F. A. *et al.* Rapid evolutionary turnover of mobile genetic elements drives
1064 bacterial resistance to phages. *Science* **374**, 488–492 (2021).

1065 191. Hunter, M. & Fusco, D. Superinfection exclusion: A viral strategy with short-term
1066 benefits and long-term drawbacks. *PLoS Comput. Biol.* **18**, (2022).

1067 192. Van Melder, L. & De Bast, M. S. Bacterial Toxin–Antitoxin Systems: More Than
1068 Selfish Entities? *PLOS Genet.* **5**, e1000437 (2009).

1069 193. Koonin, E. V. & Makarova, K. S. Origins and evolution of CRISPR-Cas systems.
1070 *Philos. Trans. R. Soc. B* **374**, (2019).

1071 194. Dawkins, R. & Krebs, J. R. Arms races between and within species. *Proc. R. Soc.*
1072 *London. Ser. B. Biol. Sci.* **205**, 489–511 (1979).

1073 195. Kerr, B., Riley, M. A., Feldman, M. W. & Bohannan, B. J. M. Local dispersal promotes
1074 biodiversity in a real-life game of rock–paper–scissors. *Nature* **418**, 171–174 (2002).

1075 196. Longdon, B., Brockhurst, M. A., Russell, C. A., Welch, J. J. & Jiggins, F. M. The
1076 Evolution and Genetics of Virus Host Shifts. *PLoS Pathog.* **10**, e1004395 (2014).

1077 197. Friman, V.-P. & Buckling, A. Effects of predation on real-time host-parasite
1078 coevolutionary dynamics. *Ecol. Lett.* **16**, 39–46 (2013).

1079 198. Paterson, S. *et al.* Antagonistic coevolution accelerates molecular evolution. *Nature*
1080 **464**, 275–278 (2010).

1081 199. Czárán, T. L., Hoekstra, R. F. & Pagie, L. Chemical warfare between microbes
1082 promotes biodiversity. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 786 (2002).

1083 200. Riley, M. A. Positive selection for colicin diversity in bacteria. *Mol. Biol. Evol.* **10**,
1084 1048–1059 (1993).

1085 201. Gordon, D. M. & O'Brien, C. L. Bacteriocin diversity and the frequency of multiple
1086 bacteriocin production in *Escherichia coli*. *Microbiology* **152**, 3239–3244 (2006).

1087 202. Michel-Briand, Y. & Baysse, C. The pyocins of *Pseudomonas aeruginosa*. *Biochimie*
1088 **84**, 499–510 (2002).

1089 203. Merker, M. *et al.* Evolutionary Approaches to Combat Antibiotic Resistance:
1090 Opportunities and Challenges for Precision Medicine. *Front. Immunol.* **11**, 1938
1091 (2020).

1092 204. Paradkar, A. Clavulanic acid production by *Streptomyces clavuligerus*: Biogenesis,
1093 regulation and strain improvement. *Journal of Antibiotics* **66**, 411–420 (2013).

1094 205. Lee, M. D. *et al.* Microbial fermentation-derived inhibitors of efflux-pump-mediated
1095 drug resistance. *Farmaco* **56**, 81–85 (2001).

1096 206. Bambeke, F., Pages, J.-M. & Lee, V. Inhibitors of Bacterial Efflux Pumps as Adjuvants
1097 in Antibiotic Treatments and Diagnostic Tools for Detection of Resistance by Efflux.
1098 *Recent Pat. Antiinfect. Drug Discov.* **1**, 157–175 (2008).

1099 207. Braun, V., Pramanik, A., Gwinner, T., Köberle, M. & Bohn, E. Sideromycins: Tools
1100 and antibiotics. in *BioMetals* **22**, 3–13 (Biomaterials, 2009).

1101 208. Tillotson, G. Trojan horse antibiotics—a novel way to circumvent Gram-negative
1102 bacterial resistance? *Infect. Dis.* **9**, 45–52 (2016).

1103 209. Coulthurst, S. The Type VI secretion system: A versatile bacterial weapon. *Microbiol.*
1104 *(United Kingdom)* **165**, 503–515 (2019).

1105 210. Boles, B. R. & Horswill, A. R. agr-mediated dispersal of *Staphylococcus aureus*
1106 biofilms. *PLoS Pathog.* **4**, e1000052 (2008).

1107 211. Paluch, E., Rewak-Soroczyńska, J., Jędrusik, I., Mazurkiewicz, E. & Jermakow, K.
1108 Prevention of biofilm formation by quorum quenching. *Applied Microbiology and*
1109 *Biotechnology* **104**, 1871–1881 (2020).

1110 212. Samson, J. E., Magadán, A. H., Sabri, M. & Moineau, S. Revenge of the phages:
1111 defeating bacterial defences. *Nat. Rev. Microbiol.* **11**, 675–87 (2013).

1112 213. Paterson, S. *et al.* Antagonistic coevolution accelerates molecular evolution. *Nature*
1113 **464**, 275–278 (2010).

1114 214. Auer, B. & Schweiger, M. Evidence that *Escherichia coli* virus T1 induces a DNA
1115 methyltransferase. *J. Virol.* **49**, 588–590 (1984).

1116 215. Iida, S., Streiff, M. B., Bickle, T. A. & Arber, W. Two DNA antirestriction systems of
1117 bacteriophage P1, *darA*, and *darB*: characterization of *darA*- phages. *Virology* **157**,
1118 156–166 (1987).

1119 216. Pawluk, A., Davidson, A. R. & Maxwell, K. L. Anti-CRISPR: Discovery, mechanism
1120 and function. *Nat. Rev. Microbiol.* **16**, 12–17 (2018).

1121 217. Scholl, D., Adhya, S. & Merril, C. *Escherichia coli* K1's capsule is a barrier to

1122 bacteriophage T7. *Appl. Environ. Microbiol.* **71**, 4872–4874 (2005).

1123 218. Otsuka, Y. & Yonesaki, T. Dmd of bacteriophage T4 functions as an antitoxin against
1124 Escherichia coli LsoA and RnIA toxins. *Mol. Microbiol.* **83**, 669–681 (2012).

1125 219. Erez, Z. *et al.* Communication between viruses guides lysis–lysogeny decisions. *Nat.*
1126 *2017 5417638* **541**, 488–493 (2017).

1127 220. Jousset, A. Ecological and evolutive implications of bacterial defences against
1128 predators. *Environ. Microbiol.* **14**, 1830–1843 (2012).

1129 221. Kaplan, F. *et al.* Bacterial attraction and quorum sensing inhibition in caenorhabditis
1130 elegans exudates. *J. Chem. Ecol.* **35**, 878–892 (2009).

1131 222. Adli, M. The CRISPR tool kit for genome editing and beyond. *Nat. Commun.* **9**,
1132 (2018).

1133 223. Bernheim, A. *et al.* Prokaryotic viperins produce diverse antiviral molecules. *Nature*
1134 **589**, 120–124 (2021).

1135 224. Raffatellu, M. Learning from bacterial competition in the host to develop
1136 antimicrobials. *Nat. Med.* *2018 248* **24**, 1097–1103 (2018).

1137 225. Ghosh, C., Sarkar, P., Issa, R. & Haldar, J. Alternatives to Conventional Antibiotics in
1138 the Era of Antimicrobial Resistance. *Trends Microbiol.* **27**, 323–338 (2019).

1139 226. Hiltunen, T., Virta, M. & Anna-Liisa, L. Antibiotic resistance in the wild: an eco-
1140 evolutionary perspective. *Philos. Trans. R. Soc. B Biol. Sci.* **372**, (2017).

1141 227. Vassallo, C. N., Doering, C. R., Littlehale, M. L., Teodoro, G. I. C. & Laub, M. T. A
1142 functional selection reveals previously undetected anti-phage defence systems in the
1143 E. coli pangenome. *Nat. Microbiol.* *2022 710* **7**, 1568–1579 (2022).

1144 228. Millman, A. *et al.* An expanded arsenal of immune systems that protect bacteria from
1145 phages. *Cell Host Microbe* **30**, 1556-1569.e5 (2022).

1146 229. Coyte, K. Z., Schluter, J. & Foster, K. R. The ecology of the microbiome: Networks,
1147 competition, and stability. *Science* **350**, 663–6 (2015).

1148 230. Coyte, K. Z., Rao, C., Rakoff-Nahoum, S. & Foster, K. R. Ecological rules for the
1149 assembly of microbiome communities. *PLoS Biol.* **19**, (2021).

1150 231. García-Bayona, L. & Comstock, L. E. Bacterial antagonism in host-associated
1151 microbial communities. *Science* **361**, (2018).

1152 232. Ventola, C. The antibiotic resistance crisis: part 1: causes and threats. *P&T* **40**, 277–
1153 283 (2015).

1154 233. Aminov, R. I. & Mackie, R. I. Evolution and ecology of antibiotic resistance genes.
1155 *FEMS Microbiol. Lett.* **271**, 147–161 (2007).

1156 234. Abrudan, M. I. *et al.* Socially mediated induction and suppression of antibiosis during
1157 bacterial coexistence. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 11054–11059 (2015).

1158 235. González-Bello, C. Antibiotic adjuvants – A strategy to unlock bacterial resistance to

antibiotics. *Bioorganic and Medicinal Chemistry Letters* **27**, 4221–4228 (2017).

236. Zhang, Z. *et al.* Antibiotic production in *Streptomyces* is organized by a division of labor through terminal genomic differentiation. *Sci. Adv.* **6**, (2020).

237. Batra, A. *et al.* High potency of sequential therapy with only β -lactam antibiotics. *Elife* **10**, (2021).

238. Beardmore, R. E., Peña-Miller, R., Gori, F., Iredell, J. & Barlow, M. Antibiotic Cycling and Antibiotic Mixing: Which One Best Mitigates Antibiotic Resistance? *Mol. Biol. Evol.* **34**, 802–817 (2017).

239. Baym, M., Stone, L. K. & Kishony, R. Multidrug evolutionary strategies to reverse antibiotic resistance. *Science (80-.).* **351**, (2016).

240. Jamet, A. *et al.* A widespread family of polymorphic toxins encoded by temperate phages. *BMC Biol.* **15**, 1–12 (2017).

241. Wang, M. *et al.* Modular Design of Membrane-Active Antibiotics: From Macromolecular Antimicrobials to Small Scorpionlike Peptidomimetics. *J. Med. Chem.* **64**, 9894–9905 (2021).

242. Trejo-Hernández, A., Andrade-Domínguez, A., Hernández, M. & Encarnación, S. Interspecies competition triggers virulence and mutability in *Candida albicans*–*Pseudomonas aeruginosa* mixed biofilms. *ISME J. 2014 810* **8**, 1974–1988 (2014).

243. Flint, A., Stintzi, A. & Saraiva, L. M. Oxidative and nitrosative stress defences of *Helicobacter* and *Campylobacter* species that counteract mammalian immunity. *FEMS Microbiol. Rev.* **40**, 938–960 (2016).

244. Wille, J. & Coenye, T. Biofilm dispersion: The key to biofilm eradication or opening Pandora’s box? *Biofilm* **2**, 100027 (2020).

245. Dieltjens, L. *et al.* Inhibiting bacterial cooperation is an evolutionarily robust anti-biofilm strategy. *Nat. Commun. 2020 111* **11**, 1–11 (2020).

246. Brown, S. P., West, S. A., Diggle, S. P. & Griffin, A. S. Social evolution in micro-organisms and a Trojan horse approach to medical intervention strategies. *Philos. Trans. R. Soc. B Biol. Sci.* **364**, 3157–3168 (2009).

247. Gurney, J., Simonet, C., Wollein Waldetoft, K. & Brown, S. P. Challenges and opportunities for cheat therapy in the control of bacterial infections. *Nat. Prod. Rep.* **39**, 325–334 (2022).

248. Lin, D. M., Koskella, B. & Lin, H. C. Phage therapy: An alternative to antibiotics in the age of multi-drug resistance. *World J. Gastrointest. Pharmacol. Ther.* **8**, 162 (2017).

249. Atterbury, R. J. & Tyson, J. Predatory bacteria as living antibiotics – Where are we now? *Microbiol. (United Kingdom)* **167**, 1–8 (2021).

250. Torres-Barceló, C. The disparate effects of bacteriophages on antibiotic-resistant bacteria. <https://doi.org/10.1038/s41426-018-0169-z> **7**, (2018).

- 1196 251. Chan, B. K. *et al.* Phage selection restores antibiotic sensitivity in MDR *Pseudomonas*
1197 *aeruginosa*. *Sci. Reports 2016 6* **6**, 1–8 (2016).
1198 252. Roemhild, R. & Andersson, D. I. Mechanisms and therapeutic potential of collateral
1199 sensitivity to antibiotics. *PLOS Pathog.* **17**, e1009172 (2021).
1200

Acknowledgements

We are indebted to Elisa Granato, Rachel Wheatley, Connor Sharp and Michael Brockhurst for their helpful comments on the manuscript, and to Martin Jahn, Célia Souque, Erik Bakkeren, Frances Spragge, Jacob Palmer, Sean Booth, Olivier Cunrath, Craig Maclean and Laurie Comstock for their literature suggestions. C.D.N. is supported by the Simons Foundation award number 826672, NSF grant IOS 2017879, and grant RGY0077/2020 from the Human Frontier Science Program. B.R.W. received support from a Gillman Fellowship from the Department of Biological Sciences at Dartmouth. W.P.J.S. and K.R.F. are supported by the National Institutes of Health (project numbers R01AI093771 and R01AI120633), by European Research Council Grant 787932, and by Wellcome Trust Investigator award 209397/Z/17/Z. W.P.J.S. is also funded by a Sir Henry Wellcome Postdoctoral fellowship award, 222795/Z/21/Z.

Author contributions


W.P.J.S. researched data for article. W.P.J.S., K.R.F., B.R.W., and C.D.N. contributed substantially to the discussion of content. W.P.J.S. and K.R.F. wrote the article. W.P.J.S., B.R.W., C.D.N., and K.R.F. reviewed and edited the manuscript before submission.

Competing interests

The authors declare no competing interests.

Peer review information

Nature Reviews Microbiology thanks Jordan Vacheron, who co-reviewed with Clara Heiman, and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.



1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236

Figure 1. Bacteria face diverse threats from competitors, viruses and predators. Most attacks target select core cellular processes and functions of the bacterial target cell. Coloured squares indicate whether a given threat type typically acts on a particular target. Bacterial competitors antagonise a target bacterium via diverse mechanisms, including both contact-dependent weaponry (the type VI secretion system (T6SS); Cdi effectors) and diffusible weaponry (small molecules, peptide toxins, and tailocins). The majority of clinical antibiotics are also derived from bacteria and other microorganisms. Following infection of a bacterial cell, phages attack cell walls and membranes to release their progeny via cell lysis. Some bacterial predators, such as *Bdellovibrio* species and like organisms (BALOs), invade the host cell periplasm, injecting toxins that digest various cytoplasmic components. Many eukaryotic predators engulf and digest target bacteria whole in phagosome compartments.




Figure 2. Bacteria have evolved multiple lines of defence against biotic threats. At both the individual and collective level, bacteria draw upon a plethora of defensive adaptations to escape harm. Defences are arranged according to the spatial scale at which they operate. **(a)** *Molecular level*: attacks by competitors, phages and predators are mediated by harmful agents (for example, toxins, enzymes and genetic elements), which disrupt cellular functions by interacting with diverse targets. Bacteria can mitigate disruption at a molecular level, by altering the target or compensating for its disruption, or by destroying or binding the harmful agent. **(b)** *Cellular level*: macromolecular barriers, including cell membranes, S-layers, lipopolysaccharide (LPS) or capsules, prevent harmful agents from entering a bacterial cell. Efflux pumps remove harmful molecules that overcome barriers, and motile bacteria can escape harmful environments by repositioning themselves. Secreted membrane vesicles can bind and inactivate toxins and phages. Stress responses and other regulatory pathways enable these defences to be activated in response to specific or general threat cues. **(c)** *Multicellular level*: bacteria also create collective barriers (production of extracellular polymeric substances (EPS); biofilm formation) or resistant subpopulations (phenotypic heterogeneity), launch en-masse counterattacks, and, in some circumstances (e.g. abortive infection), commit suicide to protect kin cells.




Figure 3. Bacteria mount defences in response to diverse cues. Examples are ordered according to the proximity of potential harm, and grouped according to type. Some cues emanate from direct harm to a focal cell (harm from abiotic stressors; nutrient depletion or attacks by competitors); bacteria identify and distinguish these cues via competition sensing, and respond defensively. Bacteria can also respond to attacks before they themselves are harmed, activating defences in response to danger cues (kin lysate, non-kin toxins, signals and other molecular attacker signatures). Bacteria also use autoinducer-mediated quorum sensing, and other density-sensing mechanisms, to raise defences in anticipation of attacks.




Figure 4: Bacteria innovate, acquire and accumulate defences. (a) Random mutations alter bacterial susceptibility to threats (for example, via modification of target structures), occasionally conferring survival benefits. Bacteria may also acquire new defence genes via horizontal gene transfer: conjugation, natural transformation and phage transduction. (b) Bacteria accumulate toxin-immunity pairs and orphan immunity genes in their genomes, protecting them against the cognate toxins of both kin and competitor cells. (c) CRISPR–Cas systems remember past infections by storing phage and plasmid DNA samples in spacer libraries. (d) Gene cassettes encoding antibiotic resistance and other defensive functions are captured by integrons via site-specific recombination. Stress cues (lightning bolt) stimulate expression of captured genes; stress-induced integrases also shuffle cassettes, resulting in diverse gene expression profiles within a population.




Figure 5: Counter-adaptations to bacterial defences by competitors, phage and predators. (a) Attackers prevent degradation of their toxins (or DNA, in the case of phages) using adjuvants to inhibit defence enzyme function, or by modifying toxin structure. Toxin cocktails may offer toxin synergy and delay resistance evolution. (b) Competitors bypass the membranes of their target cells using toxin injection systems (the type VI secretion system (T6SS)) or by disguising toxins as useful substrates ('Trojan horses'). Some toxins kill without

triggering key stress responses, suppressing defensive behaviour ('Alarm suppression'). Efflux pump inhibitors prevent expulsion of absorbed molecular toxins. Phages penetrate cell capsules using tail-mounted hydrolases, and adapt to alterations in host receptor structure via counter-modification or stochastic expression of receptor-binding proteins. **(c)** Competitors degrade biofilms using dispersants and matrix hydrolases, and inhibit response coordination using quorum quenching. Phages override collective immunity by bypassing abortive infection mechanisms, using hijacked or surrogate immunity proteins to disarm suicide systems.

Box 1. Clinical implications of ancient bacterial defences

The study of bacterial defences can inform current and future antibacterial therapeutics.

[bH1] Origins of drug resistance. Understanding where resistance genes come from can help to predict and restrict antibiotic resistance proliferation²³². Environmental reservoirs harbour many old and diverse resistance genes²³³. For example, the methicillin-resistance genes found in methicillin-resistant *Staphylococcus aureus* (MRSA) seem to have first emerged in hedgehog-associated *Staphylococcus aureus*, as a protection against fungal β -lactam antibiotics³⁹. More generally, toxin-mediated competition among environmental bacteria is widespread²³⁴, and, along with phages and predators^{41,61}, can select for defences that increase virulence^{120,121,186} or protect bacteria against multiple different threats^{144,154,226}. Studying and surveying bacterial defences in environments with strong competition and conflict, therefore, may help to predict which resistance mechanisms are most likely to arise.

[bH1] New strategies against resistance. Many bacteria use antimicrobials to eliminate competitors¹⁰, which suggests they are often able to overcome the defences of their targets. We might look to bacteria, therefore, for strategies that help to overcome drug resistance. Evidence supporting this idea comes from the use of adjuvant therapy: *Streptomyces clavuligerus* produces clavulanic acid, which inhibits β -lactamase-based resistance mechanisms²⁰⁴. This strategy forms the basis for *Augmentin*, a therapeutic that uses both a β -lactam antibiotic and clavulanic acid to combat β -lactamase-based resistance²³⁵. Another feature of bacterial attack strategies is that they commonly use multiple different toxins against competitors^{10,201,236}. This contrasts with classic mono-therapy, which remains the clinical norm, but draws comparisons to a growing number of strategies that combine multiple antibiotics with the goal of limiting resistance evolution^{237–239}. In addition, many bacterial toxins are polymorphic, with a modular structure that enables new variants to be readily innovated as resistance emerges²⁴⁰. Adopting modular designs when developing new antimicrobials could enable us to exploit this adaptability²⁴¹.

[bH1] Targeting defences. The defensive responses of bacteria^{79,81,123} can increase virulence and protect against antimicrobial treatment, thereby exacerbating disease^{81,242,243}. Directly targeting defensive mechanisms, therefore, has the potential to greatly improve treatment efficacy when performed in combination with antibiotics or other bactericidal treatments. Diverse bacteria respond to antibiotic treatment by forming biofilms, which are notoriously difficult to treat⁸⁰. However, physical disruption of biofilm structures can increase bacterial exposure to antibiotics, sensitising recalcitrant infections²⁴⁴. Targeting defences also raises the possibility of treatments with a minimised risk of resistance evolution. Biofilm inhibitors can enhance antibiotic susceptibility while minimising resistance to the biofilm inhibitor, because resistant genotypes pay the fitness costs of EPS production²⁴⁵. A related defence-targeting strategy is to introduce strains of bacteria that do not contribute to collective defences ('cheat therapy')^{246,247}. Where cheater strains can outcompete the original strain, they have the potential to undermine defences and improve treatment outcomes without strong natural selection for resistance evolution.

[bH1] Exploiting novel antimicrobials. Phages²⁴⁸, predators²⁴⁹ and competing bacteria^{5,224} all have potential as alternative therapeutics for bacterial infections²²⁵. As we have discussed, however, bacteria have already evolved many defences against these threats. As with antibiotics, therefore, the rapid emergence of resistance in clinical settings seems to be likely^{9,250}. But these alternative antimicrobials share a potential major advantage over antibiotics: being biological, they have the potential to coevolve with their targets, such that resistance in a target is circumvented by countermeasures in the attacker. Although this outcome is far from guaranteed (it requires, amongst other things, that the survival of the therapeutic depends on defeating the target pathogen), it raises the possibility that evolution can be directed to overcome pathogen resistance as it emerges. Moreover, by combining therapies, one can exert contrasting selective pressures on pathogens, which may limit resistance evolution more than via antibiotic therapy alone^{177,251,252}.

1348 **Glossary of terms**

1349

1350 **Competitor**

1351 Another type of bacteria that competes with a focal bacterium for resources. Often this will be
1352 a genetically similar, but non-identical bacterium (for example, a different strain), as similar
1353 bacteria are most likely to have overlapping resource needs. Genetically identical organisms
1354 compete in an ecological sense, but not in an evolutionary sense (as they have the same
1355 evolutionary interests). In this Review, we use the term in the former sense.

1356

1357 **Bacteriophage** (phage)

1358 A virus that infects bacteria.

1359

1360 **Predator**

1361 An organism that consumes another for food, killing it in the process.

1362

1363 **Defence mechanisms**

1364 Traits that evolved, at least in part, to protect an organism against a threat. This term is often
1365 used in the context of bacterial defences against viral threats, but in this Review, we expand
1366 it to encompass protection against competitors and predators.

1367

1368 **Biotherapeutic**

1369 Medicine that is derived from (and often incorporating) biological entities. Phages are a
1370 potential biotherapeutic for treating bacterial infections.

1371

1372 **Preadaptations**

1373 Evolutionary adaptation which serves a different purpose from the one for which it first evolved.
1374 For instance, many modern efflux pumps function to remove antibiotics from bacterial cells,
1375 but homologous structures likely served different functions (e.g. metabolite export) in ancestral
1376 strains.

1377

1378 **Stressors**

1379 Changes in environmental or physiological conditions that perturb cell homeostasis.

1380

1381 **Weaponry**

1382 Cellular systems that evolved, at least in part, to harm other organisms.

1383

1384 **Parasitism**

1385 An evolutionary relationship between two organisms, in which one benefits at the expense of
1386 the other. In contrast to predators, parasites are generally smaller than and physically
1387 associated with the organisms they exploit.

1388

1389 **Mutualism**

1390 A mutually beneficial evolutionary relationship between two organisms; that is, one in which
1391 the fitness of the two parties are both improved by the presence of the other.

1392

1393 **Agents**

1394 Substances (particularly toxins and injected viral DNA) that, through interaction with targets,
1395 produce harm to a bacterial cell.

1396

1397 **Biofilms**

1398 Densely-packed cell groups that can contain billions or trillions of cells, enveloped by secreted
1399 extracellular matrix.

1400

1401 **Collective defence**

1402 Any defensive behaviour that becomes more effective when many individuals engage in it.
1403 Collective defences benefit the social partners of a focal bacterium, but do not always evolve
1404 for this reason.

1405

1406 **Counterattacks**

1407 Aggressions in response to aggression (apparent or actual).

1408

1409 **Plastic responses**

1410 'Programmed' alterations to bacterial phenotype in response to environmental change.
1411 Plasticity does *not* result from genetic change (though it may be genetically encoded).

1412

1413 **Stress responses**

1414 A set of regulatory pathways found in bacteria, which alter gene expression and cell physiology
1415 in response to harmful environmental changes and help the bacteria to survive stress.

1416

1417 **Competition sensing**

1418 The bacterial behaviour of discerning and responding to stress cues associated with
1419 competitor activity, often via stress responses. This is often used to regulate defences,
1420 especially counterattacks.

1421

1422 **Danger sensing**

1423 Conceptually similar to competition sensing, but pertaining to cues other than those resulting
1424 from direct harm to a focal cell.

1425

1426 **Exploitation competition**

1427 Mutually harmful interactions between bacteria, stemming from competition for contested
1428 resources (for example, space or nutrients). Contrasts with interference competition, where
1429 harm is inflicted directly via weaponry or other means.

1430

1431 **Quorum sensing**

1432 A widespread density-sensing mechanism found in bacteria and other microbes. Bacteria
1433 probe their effective density by secreting small molecules (autoinducers), which stimulate their
1434 own production. High autoinducer concentrations then become a proxy for high cell density or
1435 for restrictive spatial constraints that limit autoinducer diffusion. Quorum sensing is often used
1436 to regulate costly traits whose benefits depend on collective action.

1437

1438 **Horizontal gene transfer (HGT)**

1439 The flow of genetic information between two organisms, other than that which occurs via
1440 reproduction (vertical gene transfer).

1441

1442 **Pleiotropy**

1443 Phenomenon whereby one gene simultaneously affects multiple traits. Through pleiotropy, a
1444 defensive adaptation may affect the phenotype of a bacterium in unexpected ways (for
1445 example, reducing its fitness in the absence of a threat).

1446

1447 **Table of content:**

1448 In this Review, Smith, Foster and colleagues explore the protective strategies of bacteria,
1449 including the mechanisms, evolution and clinical implications of these ancient defences. They
1450 also review the countermeasures that attackers have evolved to overcome bacterial defences.