


Cell Reports

Oxytocin Modulates Nociception as an Agonist of Pain-Sensing TRPV1

Graphical Abstract



Authors

Yelena Nersesyan, Lusine Demirkhanyan, Deny Cabezas-Bratesco, ..., Carmen Domene, Sebastian Brauchi, Eleonora Zakharian

Correspondence

zakharel@uic.edu

In Brief

Oxytocin is known to suppress painful stimuli of inflammatory origin. Nersesyan et al. now find that oxytocin attenuates pain via the pain-sensing receptor TRPV1.

Highlights

- TRPV1 acts as an ionotropic oxytocin receptor in cells
- Oxytocin potentiates TRPV1 in cells and lipid bilayers
- Oxytocin interacts with TRPV1 at the extracellular pore loop region
- Oxytocin attenuates capsaicin-induced nociception via TRPV1 desensitization



Oxytocin Modulates Nociception as an Agonist of Pain-Sensing TRPV1

Yelena Nersesyan,¹ Lusine Demirkhanyan,¹ Deny Cabezas-Bratesco,² Victoria Oakes,³ Ricardo Kusuda,^{4,5} Tyler Dawson,¹ Xiaohui Sun,^{1,6} Chike Cao,⁷ Alejandro Martin Cohen,⁸ Bharath Chelluboina,¹ Krishna Kumar Veeravalli,¹ Katharina Zimmermann,⁴ Carmen Domene,^{3,9} Sebastian Brauchi,² and Eleonora Zakharian^{1,10,*}

¹Department of Cancer Biology and Pharmacology, University of Illinois College of Medicine, 1 Illini Drive, Peoria, IL 61605, USA

²Instituto de Fisiología, Facultad de Medicina, Universidad Austral de Chile, and Millennium Nucleus of Ion Channels-Associated Diseases (MiNICAD), Valdivia 5110566, Chile

³Department of Chemistry, King's College London, Britannia House, 7 Trinity Street, London, SE1 1DB, UK

⁴Department of Anesthesia, Friedrich-Alexander University Erlangen-Nürnberg, Krankenhausstrasse 12, 91054 Erlangen, Germany

⁵Department of Pharmacology, Ribeirão Preto Medical School, University of São Paulo, Av. Bandeirantes 3900, 14049-900 Ribeirão Preto, Brazil

⁶College of Pharmaceutical Sciences, Soochow University, Suzhou, Jiangsu 215123, China

⁷Ion Channel Research Unit, Duke University Medical Center, Durham, NC 27710, USA

⁸Department of Biochemistry and Molecular Biology, Faculty of Medicine, Dalhousie University, 5850 College Street, P.O. Box 15000, Halifax, NS B3H 4R2, Canada

⁹Chemistry Research Laboratory, Mansfield Road, University of Oxford, Oxford, OX1 3TA, UK

¹⁰Lead Contact

*Correspondence: zakharel@uic.edu

<https://doi.org/10.1016/j.celrep.2017.10.063>

SUMMARY

Oxytocin is a hormone with various actions. Oxytocin-containing parvocellular neurons project to the brainstem and spinal cord. Oxytocin release from these neurons suppresses nociception of inflammatory pain, the molecular mechanism of which remains unclear. Here, we report that the noxious stimulus receptor TRPV1 is an ionotropic oxytocin receptor. Oxytocin elicits TRPV1 activity in native and heterologous expression systems, regardless of the presence of the classical oxytocin receptor. In TRPV1 knockout mice, DRG neurons exhibit reduced oxytocin sensitivity relative to controls, and oxytocin injections significantly attenuate capsaicin-induced nociception in *in vivo* experiments. Furthermore, oxytocin potentiates TRPV1 in planar lipid bilayers, supporting a direct agonistic action. Molecular modeling and simulation experiments provide insight into oxytocin-TRPV1 interactions, which resemble DkTx. Together, our findings suggest the existence of endogenous regulatory pathways that modulate nociception via direct action of oxytocin on TRPV1, implying its analgesic effect via channel desensitization.

INTRODUCTION

Oxytocin is a versatile hormone with various physiological actions. The wide range of oxytocin-elicited effects include stimulation of uterine smooth muscle contraction to facilitate labor,

induction of lactation, and expression of complex social and bonding behaviors related to reproduction and care for offspring (Carter, 2014). Oxytocin is produced in the brain in the hypothalamic paraventricular (PVN), supraoptic (SON), and intermediate accessory nuclei (Sawchenko and Swanson, 1983). Magnocellular neurons release oxytocin into the bloodstream (Scharrer and Scharrer, 1940) and also innervate various regions of the brain, including the nucleus accumbens (Ross et al., 2009) and the central nucleus of the amygdala (Knobloch et al., 2012). The brainstem and various regions of the spinal cord (Swanson and Sawchenko, 1983) are innervated by oxytocin-containing parvocellular neurons, and the secreted neuropeptide modulates nociception via targeting C-type fibers in the dorsal root ganglion (DRG) (Juif and Poisbeau, 2013).

Many activities of oxytocin are accomplished through the canonical oxytocin receptor (OXTR), a G-protein-coupled receptor that is present in neural tissues, uterus, and breast (Gimpl and Fahrenholz, 2001). However, not all oxytocin-provoked actions have been linked to its cognate receptor. For example, oxytocin is present at distinctly high levels in brain regions that lack OXTR (Gimpl and Fahrenholz, 2001). Among oxytocin actions is the regulation of Ca^{2+} fluxes. These responses can involve (i) OXTR-mediated signaling via $\text{G}_{\alpha_q/11}$ that stimulates release of Ca^{2+} from internal stores and (ii) the influx of extracellular Ca^{2+} . The latter mechanism is dependent upon the extracellular Ca^{2+} concentrations, which suggested the participation of voltage-gated or ligand-coupled channels (Sanborn et al., 1998).

Here, we aimed to identify the receptor responsible for an ionotropic oxytocin effect. Our data indicate that oxytocin directly targets the pain receptor, TRPV1 (transient receptor potential vanilloid 1), a non-selective Ca^{2+} -permeable cation channel. TRPV1 is expressed in afferent somatosensory neurons and along the spinal cord. The original discovery of the TRPV1 channel as a heat and capsaicin receptor validated the existence of



nociceptive neural circuitry that mitigates awareness of noxious stimulus exposure (Caterina et al., 1997; Julius, 2013).

In this work, using native and heterologous expression systems, we find that oxytocin-elicited Ca^{2+} responses are mediated through TRPV1 channels. This activity is consistently observed in TRPV1-expressing cells, regardless of the presence of OXTR. Furthermore, DRG neurons isolated from the TRPV1-deficient mice displayed reduced oxytocin sensitivity in comparison to the wild-type and other controls. Planar lipid bilayer experiments confirmed that oxytocin can directly activate the TRPV1 channel. Molecular modeling indicated that, similarly to the tarantula double knot toxin (DkTx), oxytocin interacts with TRPV1 at the extracellular domain leading to channel gating. Together, our results provide solid evidence that oxytocin is a direct TRPV1 agonist. These results imply that oxytocin-induced suppression of nociception might be achieved directly through potentiation of the pain receptor TRPV1, causing analgesia upon the desensitization of the channel.

RESULTS

Oxytocin Elicits Ca^{2+} Influx through TRPV1 Channels

To assess the mechanism of oxytocin-induced anti-nociception, we tested whether it exerts any action on pain-sensing TRPV1. To test the effects of oxytocin on TRPV1 channels, initial experiments used a heterologous expression system, HEK293 cells, to stably express the channel. Oxytocin concentrations used were in the low micromolar range, which matches some physiological conditions. In the neurosecretory granules of the posterior pituitary, oxytocin is present at high concentrations ($>100\ \mu\text{M}$) and is complexed to its carrier protein neurophysin (Gimpl and Fahrenholz, 2001; Ludwig and Leng, 2006). Upon secretion, oxytocin concentrations can range from tens to hundreds of micromolar (Gimpl and Fahrenholz, 2001; Ludwig and Leng, 2006). In contrast, in cerebrospinal fluid, oxytocin is present in low picomolar concentrations, while plasma oxytocin can reach hundreds of picomolar (Kendrick et al., 1991; Winslow et al., 2003).

Under these experimental conditions, oxytocin induced rapid Ca^{2+} influx in cells stably expressing TRPV1 (Figures 1A and 1B), albeit modest in comparison to capsaicin. This effect was not observed in HEK293 cells, used as a control (Figures 1B and 1E). The observed TRPV1 responses displayed a rapid desensitization, which was noticeable during the individual and sequential oxytocin applications (Figure 1B). Co-application of capsaicin with oxytocin elicited a stronger Ca^{2+} response, indicating a cumulative effect of both compounds (Figures 1A and 1B). However, the co-application also exhibited a more rapid “run-down” in comparison to the control (Figures 1A and 1B). Importantly, the oxytocin-evoked activity was abolished (1) with the pre-application of TRPV1 antagonist capsazepine or (2) upon introduction of TRPV1-specific small interfering RNA (siRNA) (Figures 1C, 1D, and S1). The Ca^{2+} imaging performed on the HEK293 cells is summarized in Figure 1F.

Interesting, pre-activation of TRPV1 using capsaicin provoked stronger responses to oxytocin (Figure S2). Under these experimental conditions, oxytocin elicited TRPV1 responses beginning

at low picomolar concentrations (Figure S2). Similar channel behavior was observed when oxytocin was applied after potentiating TRPV1 with a low-pH pulse (pH 4.5 or 5.5; data not shown). These results indicated that oxytocin potency to TRPV1 increases when the channel is present in its active/open state. It is tempting to speculate that these conditions are physiologically relevant, reflecting increased oxytocin affinity to the channel following TRPV1 potentiation by inflammatory stimuli.

To determine whether oxytocin-evoked Ca^{2+} responses are mediated through the OXTR or vasopressin 1a receptors, we tested these responses using G-protein-coupled receptor inhibitors. Cells treated with the PLC inhibitor U73122, its negative control U73343 (following the protocol described previously [Horowitz et al., 2005; Thyagarajan et al., 2009]), or the oxytocin receptor (OXTR) inhibitor atosiban displayed no differences in the oxytocin-induced responses (Figure S3).

Next, we tested TRPV1 responses using neuronal F-11 cells. Transiently expressing TRPV1 F-11 cells promptly responded to oxytocin (Figure 1). Unlike HEK293, TRPV1-F-11 cells demonstrated more sustained activity and less desensitization (Figures 1G and 1H). Since the neuronal F-11 cells could also endogenously express the canonical oxytocin receptor, we again tested whether or not OXTR mediates the oxytocin-elicited Ca^{2+} signals using OXTR-specific siRNA. No difference occurred in the magnitude or kinetics of oxytocin-induced responses (Figures 1G, 1H, and S1), whereas F-11 cells transfected with control pcDNA vector were essentially unresponsive to all the stimuli (Figure 1I). The summary of Ca^{2+} influx in F-11 cells is presented in Figure 1J.

We next examined oxytocin-evoked activity using DRG neurons isolated from the wild-type (WT) and TRPV1 knockout (TRPV1-KO, referred to as TRPV1^{-/-}) mice (Figures 2A–2G). In the majority of experiments, DRG neurons that were sensitive to oxytocin were also sensitized by capsaicin. A few exceptions were noted when the initial oxytocin-elicited responses were of higher magnitude and were further desensitized to subsequent applications of capsaicin or oxytocin (Figure 2A). DRG neurons transiently expressing TRPV1 channels were sensitive to both agonists (Figure 2B).

On average, we found no essential differences between genders. In WT female mice, oxytocin-sensitive DRGs comprised ~22%, and male neurons comprised ~26% (Figure 2G). In contrast, only a few DRG neurons isolated from TRPV1-KO animals exhibited oxytocin sensitivity, comprising ~5% responsive neurons obtained from female mice and ~2.2% from the males (Figures 2C, 2D, and 2G). We hypothesized that these residual Ca^{2+} signals were mediated through the other oxytocin receptors present in DRG neurons, including OXTR, vasopressin 1a receptors (V1aR), or some other receptors. Notably, these residual Ca^{2+} responses were kinetically slower compared to the WT mice, which could reflect G-protein-mediated Ca^{2+} release from intracellular stores (Figures 2A and 2C). Rescuing TRPV1 expression in TRPV1-KO DRG neurons by transiently expressing the channel recovered their responses to both agonists (Figure 2E).

Alternatively, we tested whether other pain receptors such as TRPA1 or TRPM8 are implicated in oxytocin-elicited activity. However, DRG neurons from TRPA1-KO and TRPM8-KO mice displayed similar sensitivity to oxytocin as those obtained from the WT (Figure S4).




Figure 1. Oxytocin Evokes TRPV1 Activity in HEK293 Cells Stably Expressing the Channel

(A) Control experiment with 10 μM capsaicin-induced TRPV1 activation ($n_{\text{exp.}} = 6$; $n_{\text{cells}} = 81$). (B) Subsequent applications of 1 μM and 10 μM of oxytocin, followed by 10 μM capsaicin, elicited transient Ca^{2+} responses with oxytocin but a cumulative effect in the presence of both compounds ($n_{\text{exp.}} = 9$; $n_{\text{cells}} = 96$).

(C) Capsazepine significantly inhibited both the oxytocin- and oxytocin/capsaicin-induced responses ($n_{\text{exp.}} = 6$; $n_{\text{cells}} = 84$); p values in comparison to those of the control in (B) are the following: 1 μM oxytocin, $p = 1.6\text{E}-7$; 10 μM oxytocin, $p = 9.3\text{E}-7$; 10 μM oxytocin/capsaicin, $p = 4.25\text{E}-7$.

(D) Co-expression of TRPV1-specific siRNA (0.75 μg) and GFP (0.25 μg) significantly inhibited all the responses ($n_{\text{exp.}} = 6$; $n_{\text{cells}} = 66$); p values in comparison to those of the control in (B) are the following: 1 μM oxytocin, $p = 1.24\text{E}-4$; 10 μM oxytocin, $p = 3.58\text{E}-4$; 10 μM oxytocin/capsaicin, $p = 8.19\text{E}-4$.

(E) Control HEK293-cell non-expressing TRPV1 channels show no responses to any of the agonists ($n_{\text{exp.}} = 4$; $n_{\text{cells}} = 100$); p values in comparison to those of the control in (B) are the following: 1 μM oxytocin, $p = 1.04\text{E}-11$; 10 μM oxytocin, $p = 1.16\text{E}-9$; 10 μM oxytocin/capsaicin, $p = 1.88\text{E}-13$.

(F) The summary presents the means under all the conditions.

Oxytocin elicits TRPV1 responses in F-11 neuronal cells transiently expressing the channel.

(G) Oxytocin- and capsaicin-induced responses on F-11 cells with transiently expressed TRPV1 (0.5 μg) and GFP (0.15 μg) ($n_{\text{exp.}} = 6$; $n_{\text{cells}} = 71$). (H) Oxytocin- and capsaicin-induced responses on F-11 cells with transiently expressed TRPV1 (0.5 μg), GFP (0.15 μg), and OXR siRNA (0.4 μg) ($n_{\text{exp.}} = 6$; $n_{\text{cells}} = 60$).

(I) Oxytocin- and capsaicin-induced responses on F-11 cells with transiently expressed empty pcDNA vector (0.5 μg) and GFP (0.15 μg) ($n_{\text{exp.}} = 6$; $n_{\text{cells}} = 48$); p values in comparison to those of

TRPV1/GFP-expressing cells are the following: 1 μM oxytocin, $p = 0.009$; 10 μM oxytocin, $p = 2.1\text{E}-6$; 10 μM oxytocin/capsaicin, $p = 2.5\text{E}-9$.

(J) The summary presents the means of Ca^{2+} responses obtained under all the conditions.

All error bars represent $\pm\text{SEM}$. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, and **** $p < 0.0001$.

Together, these results suggest that the DRG sensitivity to oxytocin is, in large part, mediated through TRPV1 channels.

Oxytocin Evokes TRPV1 Channel Activation in Whole-Cell Patch Clamp

Next, we assessed the effect of oxytocin on TRPV1 activity using an electrophysiological approach. We found that oxytocin application elicited current potentiation in whole-cell patch-clamp recordings performed on HEK293 cells expressing TRPV1 (Figures 3A–3D). The channel current showed an outward rectification similar to that of capsaicin-evoked currents. However, the maximal response against oxytocin (10 μM) was modest compared to that of capsaicin (1 μM) (Figure 3A). The conductance versus voltage relation was similar under both conditions

(conductance/voltage [GV] curves), suggesting that differences in the current density are associated with differences in the desensitization induced by these agonists. In fact, oxytocin strongly desensitized the channel, while the observed desensitization for capsaicin was only partial (Figure 3B). Desensitization observed in the context of the intact cells impeded our ability to obtain the full dose-response curve, as concentrations over 30 μM rapidly inactivated the elicited current (Figure 3D).

Oxytocin Directly Activates TRPV1 in Planar Lipid Bilayers

The use of cellular systems to decipher direct agonistic actions of hormones can be complicated by the presence of various signaling pathways that give rise to intermediate effects. Thus,

Oxytocin-induced activity in DRG neurons of WT and TRPV1-KO mice




Figure 2. Oxytocin Sensitivity Is Diminished in DRG Neurons Isolated from TRPV1-Deficient Mice

(A) Oxytocin- and capsaicin-elicited Ca^{2+} responses in DRG neurons obtained from the WT mice (number of experiments [n_{exp}] = 33). (B) Oxytocin- and capsaicin-elicited Ca^{2+} responses in the WT DRG neurons transiently expressing TRPV1 (3 μ g) together with GFP (0.5 μ g), (n_{exp} = 6). (C and D) show oxytocin and capsaicin sensitivity of DRG neurons obtained from TRPV1-KO (C) female (n_{exp} = 21) and (D) male (n_{exp} = 21) mice, respectively. (E) Oxytocin- and capsaicin-induced responses obtained from the TRPV1-KO ($^{-/-}$) DRG neurons transiently expressing TRPV1 (3 μ g) along with GFP (0.5 μ g), (n_{exp} = 6). (F) Summary of oxytocin- and capsaicin-induced responses obtained from WT and WT with transiently expressed TRPV1 channels. (G) Pie graphs present distribution of the oxytocin-sensitive and oxytocin-insensitive DRG neurons isolated from the WT female (n_{cells} = 284), WT male (n_{cells} = 482), TRPV1-KO female (n_{cells} = 318), and TRPV1-KO male (n_{cells} = 318) mice. All error bars represent \pm SEM.

to validate direct actions of oxytocin on TRPV1, we used planar lipid bilayers to characterize oxytocin-induced channel activity.

For these experiments, the TRPV1 protein was purified and incorporated into bilayers as described previously (Lukacs et al., 2013; Sun and Zakharian, 2015). Protein purity was validated using liquid chromatography-mass spectrometry (LC-MS/MS). Distinct TRPV1 peptides were identified at high abundance (Figure S5; Table S1). After incorporating TRPV1 in the bilayers, we found that oxytocin activates the channel directly, similar to the other known agonists, heat (Sun and Zakharian, 2015) and capsaicin (Lukacs et al., 2013). Likewise,

TRPV1 opening required the presence of phosphatidylinositol-4,5-bisphosphate (PIP_2) (Figures 4 and S6).

Oxytocin-induced TRPV1 opening demonstrated distinct current rectification and profound voltage dependence. These features were evident in the single-channel conductance and the open probability (Figures 4A–4D). The channel exhibited the outward current with the mean slope conductance of ~ 81 pS, while the inward current was ~ 20 pS (Figures 4A and 4C). In addition, oxytocin-evoked TRPV1 openings showed strong voltage dependence, with marked low open probability obtained at negative voltages and an exponentially increasing open




Figure 3. Oxytocin Potentiates TRPV1 Currents in Whole-Cell Patch-Clamp Recordings

(A) Current-voltage (I-V) relations obtained from voltage ramps from -140 mV to $+140$ mV in the presence of vehicle (black), 10 μ M oxytocin (blue) or 1 μ M capsaicin (red). EM, membrane electric potential.

(B) Top: time course of representative 10 μ M oxytocin and 1 μ M capsaicin responses. Bottom: the overlay of normalized responses showing differences in the desensitization kinetics.

(C) Left: G-V curves were obtained by plotting peak tail currents at -100 mV in response to voltage-activated steady-state currents from -160 mV to $+160$ mV in 20 -mV increments: voltage for half-maximal activation of a channel (V_m) values for control, 121.66 ± 14.29 ($n = 12$); for oxytocin, 64.22 ± 8.96 ($n = 12$); and for capsaicin, 76.25 ± 21.03 ($n = 4$); charge z for control, 0.85 ± 0.07 ($n = 12$); for oxytocin, 0.75 ± 0.06 ($n = 12$); and for capsaicin, 0.67 ± 0.108 ($n = 4$). Right: representative traces obtained in transfected cells subjected to voltage steps in control conditions (black) or in the presence of 10 μ M oxytocin (blue) or 1 μ M capsaicin (red).

(D) Dose-response curve fitted to concentrations <30 μ M. $EC_{50} = 7.74 \pm 8.5$; $h = 1.5 \pm 0.6$. All error bars represent \pm SEM.

probability at positive potentials (Figures 4A, 4B, and 4D). In planar lipid bilayers, oxytocin activated TRPV1 with a half-maximal effective concentration (EC_{50}) of ~ 0.32 μ M (Figure 4E).

The comparison of oxytocin-evoked TRPV1 activity to that of capsaicin demonstrated that both conductance values of capsaicin-induced inward and outward currents were greater than that of oxytocin by about 20 and 22 pS, respectively (Figure 4F). The open probability of the channel was similar at negative voltages with both agonists, whereas the open probability was greater at positive voltages for capsaicin-evoked TRPV1 activity (Figure 4G).

In summary, the planar lipid bilayer experiments confirmed that oxytocin is a direct TRPV1 agonist, and this activity can occur independent of any other intermediate signaling pathway.

Identification of an Oxytocin-Binding Site

To investigate the interaction between oxytocin and the TRPV1 ion channel at the atomic level, computational studies were per-

formed on the TRPV1 homotetramer. To that end, oxytocin molecules were docked onto the pore loop region of structures representing the open and closed states of the channel, and MD simulations of the resulting complexes were performed (see Experimental Procedures). Negligible binding to the closed state was observed.

In contrast, the open state (denoted simulation Pep1) interacted with a single oxytocin molecule via its extracellular pore loop. The interaction persisted for approximately 200 ns, with the most prevalent binding pose (clustered using a 3 -Å root-mean-square deviation [RMSD] criterion) observed for 80 ns. In a further simulation (denoted simulation Pep2) with four oxytocin molecules initiated from this binding pose, interaction with a single oxytocin peptide persisted for the duration of the 200-ns trajectory, with a stable binding pose observed for 150 ns (Figures 5A–5E).

The identified binding site lies on the external surface of the transmembrane domain, in the interfacial region between two




Figure 4. Oxytocin Directly Activates TRPV1 in Planar Lipid Bilayers in the Presence of PIP₂

(A) Representative single-channel current traces of oxytocin-induced (0.5 μ M) TRPV1 obtained in the presence of PIP₂ (5 μ M); voltages are indicated on the left. (B) Representative ramp recording obtained at -150 to $+150$ mV. (C and D) Graphs demonstrate current voltage relationship (C) and voltage dependence (D) of oxytocin-induced TRPV1; $n = 12$. (E) Dose response of oxytocin-TRPV1 activation; $EC_{50} = 0.316 \mu$ M \pm 0.02 μ M. (F) Current voltage relationship of oxytocin- and capsaicin-induced TRPV1 channels. (G) Capsaicin induces higher open probability of TRPV1 outward channel activity, as indicated by open probability differences of oxytocin-elicited (2 μ M) and capsaicin-elicited (2 μ M) openings at -60 and $+60$ mV. All error bars represent \pm SEM.


adjacent subunits. The predicted binding site is illustrated in Figure 5E. The extension of this computational analysis and further discussion of the influence on the TRPV1 activation state are presented in the Supplemental Information (see also Figure S7).

Oxytocin Interaction with TRPV1: Similarity with Other Agonists

Next, we compared the TRPV1 activation of oxytocin to that of the tarantula toxin DkTx. The latter has been studied extensively using both experimental and computational approaches (Bae et al., 2016; Bohlen et al., 2010). Mutations I599A, F649A, A657P, and F659A inhibit channel activation by DkTx, whereas V595A and T695A attenuate channel activation to a lesser extent (Bohlen

et al., 2010). Conversely, Y631A enhanced activation by DkTx (Bohlen et al., 2010). The atomic model of TRPV1 in complex with DkTx, determined by Bae et al., advocated that some of the residues, Y631, F649, A657, and T650, were in direct contact with the toxin (Bae et al., 2016). Remarkably, such interactions are replicated in the model of oxytocin-bound TRPV1, thus suggesting that oxytocin may activate TRPV1 via a pathway similar to that of DkTx. Furthermore, our simulations revealed direct engagement between oxytocin and A649 in both *apo* and *holo* TRPV1 structures, identifying an explicit pathway by which oxytocin could modulate the behavior of this region and potentiate TRPV1.

To validate the TRPV1-oxytocin-binding site derived from computational analysis, we altered the most promising target



(legend on next page)

L635A and F649A mutants lose sensitivity to oxytocin




Figure 6. L635A and F649A Mutants Lose Sensitivity to Oxytocin

Patch-clamp recording show the current obtained from the wild-type TRPV1 and mutants. (A) demonstrates representative whole-cell currents recorded by using gap-free protocol with a holding potential at -70 mV. Traces are shown in current densities (picoamperes per picofarad; pA/pF). (B) presents the summary of the relative to control current of TRPV1 wt, L635A, D646A, and F649A constructs obtained upon application of $1 \mu\text{M}$ capsaicin and $10 \mu\text{M}$ oxytocin, $n = 5$ for each construct. (C) displays voltage ramp recordings of heat-induced potentiation of the wild-type TRPV1 and the mutants upon exposure to 40°C pulses. The summary of the heat-induced activities is presented in (D). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. All error bars represent $\pm\text{SEM}$.

residues by mutagenesis and tested their function. Patch-clamp recordings showed that L635A and F649A TRPV1 mutants completely lose their sensitivity to oxytocin. Surprisingly, they were also insensitive to capsaicin (Figures 6A and 6B) but displayed potentiation by heat (Figures 6C and 6D). Conversely, D646A mutation did not alter the sensitivity of TRPV1 to oxytocin (Figure 6A). Together, these experiments are consistent with the hypothesis that oxytocin interacts with TRPV1 via L635 and F649.

Oxytocin Reduces Capsaicin-Induced Aversive Behavior in Mice

In conclusion, our findings suggest that oxytocin suppresses nociception via activation and desensitization of the polymodal TRPV1 channels in nociceptors. To test this hypothesis, we performed *in vivo* experiments to quantify nociceptive behavior in mice subsequent to oxytocin and capsaicin application. Oxytocin alone did not provoke any noticeable aversive

response in mice. However, co-application of oxytocin with capsaicin significantly induced analgesia over the capsaicin-evoked pain stimuli (Figure 7). Together, these results support a direct link between oxytocin and antinociception via TRPV1 activity modulation.

DISCUSSION

Physiological Implication of Oxytocin in Pain Modulation via TRPV1

Oxytocin is a highly abundant neurohypophyseal peptide. Its prevalent site of expression is localized to the magnocellular neurons of the hypothalamic paraventricular and supraoptic nuclei (see, for review, Gimpl and Fahrenholz, 2001). Magnocellular oxytocin neurons of these nuclei innervate the forebrain and intensely release the hormone to the systemic circulation from the posterior pituitary in response to a variety of stimuli. The parvocellular oxytocin neurons further project to the brainstem and spinal cord, where the hormone was shown to modulate inflammatory pain processing (Eliava et al., 2016). Furthermore, this neuropeptide is also synthesized in peripheral tissues, such as uterus, placenta, amnion, corpus luteum, testis, and heart (Gimpl and Fahrenholz, 2001).

Physiologic roles of oxytocin range from various modalities of neuroendocrine reflexes to the complex social and bonding behaviors related to reproduction and care of offspring. It is now well established that oxytocin facilitates reproduction in all vertebrates (Gimpl and Fahrenholz, 2001). A well-known action of the hormone is the stimulation of milk ejection that was confirmed on

Figure 5. Modeling and Molecular Dynamics Simulations Display Oxytocin-TRPV1-Interacting Site

(A–E) Contact map displaying the number of interacting atom pairs (N_p) between TRPV1 and oxytocin in simulation Pep1 (A) and Pep2 (B), averaged over the simulation length, using a 4-\AA cutoff. The prominent interactions (defined as $N_p > 15$) are summarized in (C). The occurrence of hydrogen bonds is displayed in (D). In (E), a new cartoon representation of the TRPV1 system, in complex with oxytocin is illustrated; oxytocin and interacting TRPV1 residues are displayed in licorice representation.




Figure 7. Oxytocin Reduces Capsaicin-Induced Aversive Behavior in Mice

Mouse hindpaws were intracutaneously injected under light sevoflurane anesthesia with the vehicle ($n = 3$), oxytocin ($4 \mu\text{g}$ per paw; $n = 3$), capsaicin ($1.6 \mu\text{g}$ per paw; $n = 6$), or a combination of capsaicin and oxytocin ($4 \mu\text{g}$ per paw; $n = 6$), and nocifensive behavior was quantified as the time spent licking and flinching. Columns represent the amount of time (in seconds) animals spent licking and flinching the injected paw during the 30-min observation period following the injection (mean \pm SEM; p value according to Student's t test). nd, not determined. All error bars represent \pm SEM.

the model of oxytocin-deficient mice. The role of oxytocin in parturition is also established, although it involves more complex regulatory mechanisms. Some evidence indicates that, from the paraventricular nuclei of the hypothalamus, oxytocin can reach the central amygdala, where the hormone modulates emotion-related centers of the amygdala and brainstem (Stoop, 2012). Many oxytocin actions have been well characterized by the function of its classical receptor OXTR, a class I G-protein-coupled receptor that is differentially expressed in various tissues (Gimpl and Fahrenholz, 2001). The expression of the oxytocin receptor in the uterus and hypothalamus is strongly correlative with the presence of sex hormones, particularly estrogens. The estrogen receptors were shown to play the role of transcription factors of OXTR and are required for its expression mediated by estrogen binding (Young et al., 1998). Some oxytocin actions, however, imply OXTR-independent mechanisms.

Among its many activities, oxytocin is known to be involved in nociception and pain responses. These involve both peripheral components (Juif and Poisbeau, 2013) and central components (González-Hernández et al., 2014; Juif et al., 2013). Importantly, oxytocin neurons project directly into the C-type fibers of DRG (Juif and Poisbeau, 2013). In an animal model, oxytocin release suppresses nociception and induces analgesia by specifically affecting inflammatory pain pathways (Eliava et al., 2016). The oxytocin neuronal circuitry appears to occur via a dual process, by (1) releasing oxytocin from axons onto sensory spinal cord neurons and inhibiting their activity and (2) indirectly stimulating oxytocin release from supraoptic neurons into the periphery (Eliava et al., 2016). Thus, oxytocin modulates pain by reaching the spinal cord through fast neuronal projections and slower peripheral pathways (Eliava et al., 2016). Likewise, analgesic actions of

oxytocin have been documented for migraine patients treated with intranasal oxytocin administration (Wang et al., 2013). One aspect of oxytocin-promoted pain regulation involves endogenous analgesia (González-Hernández et al., 2014). However, it is still debatable as to whether this effect is mediated through the OXTR, V1aR, or some other receptor, as the majority of these studies have been based upon pharmacological approaches using either OXTR or V1aR antagonists (for a review, see González-Hernández et al., 2014).

Also important is the effect of oxytocin on thermoregulation. Similarly to capsaicin (Kobayashi et al., 1998), peripheral administration of oxytocin induces strong hypothermia (Hicks et al., 2014). Hence, in addition to analgesia, our finding suggests a possible molecular mechanism for oxytocin-induced hypothermia.

The results reported here suggest that several oxytocin effects may be mediated by interaction with the TRPV1 pain receptor. TRPV1 is a well-investigated receptor in the pain pathway. The original discovery of TRPV1's role in nociception inspired the development of novel classes of analgesic drugs (Caterina et al., 1997; Julius, 2013). Furthermore, TRPV1 serves as a molecular marker for the nociceptive neuronal circuitry that enables sensation of noxious heat, protons, neurogenic inflammation, and thermal hyperalgesia (Julius, 2013). Profound TRPV1 expression in C-fibers contributes to the signaling associated with acute and chronic pain conditions and could be implicated in various pathophysiological conditions such as arthritis, pancreatitis, migraine headache, and others.

Indeed, the expression of TRPV1 in trigeminal afferent neurons of *dura mater* suggested a plausible linkage of the channel actions to migraine (Huang et al., 2012; Shimizu et al., 2007). More evidence for this relationship was suggested by the effect of capsaicin on dilation of dural vessels (Akerman et al., 2003) and induction of trigeminal ganglion neuronal activity (Iwashita et al., 2013). However, the mechanisms by which TRPV1 contributes to initiating or propagating the neuronal signaling related to migraine are poorly understood (for a review, see Dussor et al., 2014). Our discovery of oxytocin as an endogenous agonist for the TRPV1 channel may shed more light onto the regulatory mechanisms that underlie physiology and pathology related to headaches and chronic migraine development.

The ability of oxytocin to modulate nociception is most likely achieved by activation of TRPV1, followed by its marked desensitization. Such mechanism is evident in nocifensive mice behavior, where oxytocin attenuates capsaicin-evoked nociceptive response (Figure 7). This is consistent with the fact that therapeutic targeting of TRPV1 has been primarily advanced via exogenous agonists that lead to prompt channel desensitization. These treatment options are traditionally based on capsaicin-containing remedies, such as capsaicin cream or high-concentration capsaicin patches (Dussor et al., 2014). Intranasal capsaicin administration has also been tested in treating migraine; similar to another TRPV1 agonist civamide, capsaicin reduced the frequency of cluster headache attacks (Diamond et al., 2000; Fusco et al., 2003). In contrast, attempts to identify effective and safe TRPV1 antagonists have been stymied by numerous complications, including dysregulation of internal body temperature. In this light, identifying the endogenous

mechanisms of TRPV1 potentiation and desensitization by oxytocin along with the patterns of expression and their functional relationship within the neuronal network offers a valuable asset to pain regulation.

EXPERIMENTAL PROCEDURES

For detailed methods, see the [Supplemental Information](#).

Cell Culture

HEK293 cells were maintained in minimal essential medium (MEM) as previously described (Zakharian et al., 2009). F-11 cells were cultured in DMEM/F12 medium as previously described (Zakharian et al., 2009). Mouse DRG neurons were cultured in primary neuron basal medium (PNBM), (Lonza, Allendale, NJ).

Whole-Cell Patch-Clamp Recordings

The whole-cell patch-clamp experiments were performed as previously described (Yudin et al., 2011; Zakharian et al., 2009). HEK293T cells were cotransfected with rTRPV1 and EGFP with Lipofectamine 2000 according to the manufacturer's protocol. Whole-cell patch-clamp recordings were performed 48–72 hr post-transfection at room temperature.

Intracellular Ca^{2+} Measurements

Ca^{2+} measurements were performed as previously described (Zakharian et al., 2009).

Preparation of the TRPV1 Protein from HEK Cells

TRPV1 protein isolation was performed as previously described (Sun and Zakharian, 2015; Zakharian et al., 2010). TRPV1 was purified by immunoprecipitation with anti-Myc-IgG conjugated to A/G protein magnetic beads (Pierce, Thermo Scientific, Milwaukee, WI). For the planar lipid bilayer experiments, the protein was eluted with Myc-peptide (150 $\mu\text{g}/\text{mL}$).

Planar Lipid Bilayer Measurements

Planar lipid bilayer measurements and temperature studies were performed as previously described (Sun and Zakharian, 2015; Zakharian, 2013; Zakharian et al., 2010).

Animal Studies

Age-matched male and female wild-type and TRPV1^{-/-}, TRPA^{-/-}, and TRPM8^{-/-} mice were purchased from Jackson Laboratory (Maine, USA).

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, seven figures, and one table and can be found with this article online at <https://doi.org/10.1016/j.celrep.2017.10.063>.

AUTHOR CONTRIBUTIONS

Y.N., L.D., D.C.-B., V.O., R.K., X.S., C.C., A.M.C., and E.Z. conducted experiments. Y.N., L.D., and E.Z. conducted imaging experiments. D.C.-B. conducted electrophysiological experiments, using whole-cell patch clamp. L.D., Y.N., T.D., and X.S. conducted electrophysiological experiments, using lipid bilayers. V.O. performed computational studies and analysis. L.D. conducted western blotting experiments. R.K. performed mouse behavioral experiments. E.Z. conducted experiments on protein isolation for mass spectrometry, and A.M.C. performed mass spectrometry experiments and their analysis. B.C. helped with the mouse tissue-isolation experiments. C.C. performed molecular biology studies. K.K.V. contributed reagents and material. Y.N., L.D., D.C.-B., V.O., R.K., X.S., A.M.C., K.Z., C.D., S.B., and E.Z. analyzed data. K.Z. designed and supervised mouse behavioral experiments. C.D. designed and supervised computational studies and simulation analysis. S.B. designed and supervised patch-clamp experiments. Y.N. and E.Z. designed the study, and E.Z. supervised the experiments. E.Z. wrote the manuscript,

with the contributions of Y.N., C.D., K.Z., and S.B. All authors critically read and edited the manuscript.

ACKNOWLEDGMENTS

We deeply appreciate the commitment of Dr. Liskin Swint-Kruse and her immense help in critical reading of the manuscript and fruitful suggestions. We are grateful to Angela Daniels for her exceptional help with the animal experiments.

This work was initially supported by the NIH through grant R01GM098052 to E.Z. C.D. and V.O. acknowledge the PRACE initiative for awarding the access to computational resources in ARCHER—the UK National Supercomputing Service (<http://www.archer.ac.uk>), the PDC Centre for High Performance Computing (PDC-HPC), CINECA, and the Jülich Supercomputing Center. V.O. was supported by BB/L015269/1 from the Biotechnology and Biological Sciences Research Council and Pfizer Neusentis. S.B. was supported by Anillo Científico ACT-1401. S.B. is a part of CISNe-UACH and UACH Program for Cell Biology.

Received: February 17, 2017

Revised: August 21, 2017

Accepted: October 16, 2017

Published: November 7, 2017

REFERENCES

- Akerman, S., Kaube, H., and Goadsby, P.J. (2003). Vanilloid type 1 receptors (VR1) on trigeminal sensory nerve fibres play a minor role in neurogenic dural vasodilatation, and are involved in capsaicin-induced dural dilation. *Br. J. Pharmacol.* 140, 718–724.
- Bae, C., Anselmi, C., Kalia, J., Jara-Oseguera, A., Schwieters, C.D., Krepiy, D., Won Lee, C., Kim, E.-H., Kim, J.I., Faraldo-Gómez, J.D., and Swartz, K.J. (2016). Structural insights into the mechanism of activation of the TRPV1 channel by a membrane-bound tarantula toxin. *eLife* 5, e11273.
- Bohlen, C.J., Priel, A., Zhou, S., King, D., Siemens, J., and Julius, D. (2010). A bivalent tarantula toxin activates the capsaicin receptor, TRPV1, by targeting the outer pore domain. *Cell* 141, 834–845.
- Carter, C.S. (2014). Oxytocin pathways and the evolution of human behavior. *Annu. Rev. Psychol.* 65, 17–39.
- Caterina, M.J., Schumacher, M.A., Tominaga, M., Rosen, T.A., Levine, J.D., and Julius, D. (1997). The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* 389, 816–824.
- Diamond, S., Freitag, F., Phillips, S.B., Bernstein, J.E., and Saper, J.R. (2000). Intranasal civamide for the acute treatment of migraine headache. *Cephalalgia* 20, 597–602.
- Dussor, G., Yan, J., Xie, J.Y., Ossipov, M.H., Dodick, D.W., and Porreca, F. (2014). Targeting TRP channels for novel migraine therapeutics. *ACS Chem. Neurosci.* 5, 1085–1096.
- Eliava, M., Melchior, M., Knobloch-Bollmann, H.S., Wahis, J., da Silva Gouveia, M., Tang, Y., Ciobanu, A.C., Triana Del Rio, R., Roth, L.C., Althammer, F., et al. (2016). A new population of parvocellular oxytocin neurons controlling magnocellular neuron activity and inflammatory pain processing. *Neuron* 89, 1291–1304.
- Fusco, B.M., Barzoi, G., and Agrò, F. (2003). Repeated intranasal capsaicin applications to treat chronic migraine. *Br. J. Anaesth.* 90, 812.
- Gimpl, G., and Fahrenholz, F. (2001). The oxytocin receptor system: structure, function, and regulation. *Physiol. Rev.* 81, 629–683.
- González-Hernández, A., Rojas-Piloni, G., and Condés-Lara, M. (2014). Oxytocin and analgesia: future trends. *Trends Pharmacol. Sci.* 35, 549–551.
- Hicks, C., Ramos, L., Reekie, T., Misagh, G.H., Narlawar, R., Kassiou, M., and McGregor, I.S. (2014). Body temperature and cardiac changes induced by peripherally administered oxytocin, vasopressin and the non-peptide oxytocin receptor agonist WAY 267,464: a biotelemetry study in rats. *Br. J. Pharmacol.* 171, 2868–2887.

- Horowitz, L.F., Hirdes, W., Suh, B.C., Hilgemann, D.W., Mackie, K., and Hille, B. (2005). Phospholipase C in living cells: activation, inhibition, Ca²⁺ requirement, and regulation of M current. *J. Gen. Physiol.* 126, 243–262.
- Huang, D., Li, S., Dhaka, A., Story, G.M., and Cao, Y.Q. (2012). Expression of the transient receptor potential channels TRPV1, TRPA1 and TRPM8 in mouse trigeminal primary afferent neurons innervating the dura. *Mol. Pain* 8, 66.
- Iwashita, T., Shimizu, T., Shibata, M., Toriumi, H., Ebine, T., Funakubo, M., and Suzuki, N. (2013). Activation of extracellular signal-regulated kinase in the trigeminal ganglion following both treatment of the dura mater with capsaicin and cortical spreading depression. *Neurosci. Res.* 77, 110–119.
- Juif, P.E., and Poisbeau, P. (2013). Neurohormonal effects of oxytocin and vasopressin receptor agonists on spinal pain processing in male rats. *Pain* 154, 1449–1456.
- Juif, P.E., Breton, J.D., Rajalu, M., Charlet, A., Goumon, Y., and Poisbeau, P. (2013). Long-lasting spinal oxytocin analgesia is ensured by the stimulation of allopregnanolone synthesis which potentiates GABA(A) receptor-mediated synaptic inhibition. *J. Neurosci.* 33, 16617–16626.
- Julius, D. (2013). TRP channels and pain. *Annu. Rev. Cell Dev. Biol.* 29, 355–384.
- Kendrick, K.M., Keverne, E.B., Hinton, M.R., and Goode, J.A. (1991). Cerebrospinal fluid and plasma concentrations of oxytocin and vasopressin during parturition and vaginocervical stimulation in the sheep. *Brain Res. Bull.* 26, 803–807.
- Kobayashi, A., Osaka, T., Namba, Y., Inoue, S., Lee, T.H., and Kimura, S. (1998). Capsaicin activates heat loss and heat production simultaneously and independently in rats. *Am. J. Physiol.* 275, R92–R98.
- Knobloch, H.S., Charlet, A., Hoffmann, L.C., Eliava, M., Khrulev, S., Cetin, A.H., Osten, P., Schwarz, M.K., Seeburg, P.H., Stoop, R., and Grinevich, V. (2012). Evoked axonal oxytocin release in the central amygdala attenuates fear response. *Neuron* 73, 553–566.
- Ludwig, M., and Leng, G. (2006). Dendritic peptide release and peptide-dependent behaviours. *Nat. Rev. Neurosci.* 7, 126–136.
- Lukacs, V., Rives, J.M., Sun, X., Zakharian, E., and Rohacs, T. (2013). Promiscuous activation of transient receptor potential vanilloid 1 (TRPV1) channels by negatively charged intracellular lipids: the key role of endogenous phosphoinositides in maintaining channel activity. *J. Biol. Chem.* 288, 35003–35013.
- Ross, H.E., Cole, C.D., Smith, Y., Neumann, I.D., Landgraf, R., Murphy, A.Z., and Young, L.J. (2009). Characterization of the oxytocin system regulating affiliative behavior in female prairie voles. *Neuroscience* 162, 892–903.
- Sanborn, B.M., Dodge, K., Monga, M., Qian, A., Wang, W., and Yue, C. (1998). Molecular mechanisms regulating the effects of oxytocin on myometrial intracellular calcium. *Adv. Exp. Med. Biol.* 449, 277–286.
- Sawchenko, P.E., and Swanson, L.W. (1983). The organization of forebrain afferents to the paraventricular and supraoptic nuclei of the rat. *J. Comp. Neurol.* 218, 121–144.
- Scharrer, E., and Scharrer, B. (1940). Secretory cells within the hypothalamus. *Res. Publ. Assoc. Res. Nerv. Ment. Dis.* 20, 170–194.
- Shimizu, T., Toriumi, H., Sato, H., Shibata, M., Nagata, E., Gotoh, K., and Suzuki, N. (2007). Distribution and origin of TRPV1 receptor-containing nerve fibers in the dura mater of rat. *Brain Res.* 1173, 84–91.
- Stoop, R. (2012). Neuromodulation by oxytocin and vasopressin. *Neuron* 76, 142–159.
- Sun, X., and Zakharian, E. (2015). Regulation of the temperature-dependent activation of transient receptor potential vanilloid 1 (TRPV1) by phospholipids in planar lipid bilayers. *J. Biol. Chem.* 290, 4741–4747.
- Swanson, L.W., and Sawchenko, P.E. (1983). Hypothalamic integration: organization of the paraventricular and supraoptic nuclei. *Annu. Rev. Neurosci.* 6, 269–324.
- Thyagarajan, B., Benn, B.S., Christakos, S., and Rohacs, T. (2009). Phospholipase C-mediated regulation of transient receptor potential vanilloid 6 channels: implications in active intestinal Ca²⁺ transport. *Mol. Pharmacol.* 75, 608–616.
- Wang, Y.L., Yuan, Y., Yang, J., Wang, C.H., Pan, Y.J., Lu, L., Wu, Y.Q., Wang, D.X., Lv, L.X., Li, R.R., et al. (2013). The interaction between the oxytocin and pain modulation in headache patients. *Neuropeptides* 47, 93–97.
- Winslow, J.T., Noble, P.L., Lyons, C.K., Sterk, S.M., and Insel, T.R. (2003). Rearing effects on cerebrospinal fluid oxytocin concentration and social buffering in rhesus monkeys. *Neuropsychopharmacology* 28, 910–918.
- Young, L.J., Wang, Z., Donaldson, R., and Rissman, E.F. (1998). Estrogen receptor alpha is essential for induction of oxytocin receptor by estrogen. *Neuroreport* 9, 933–936.
- Yudin, Y., Lukacs, V., Cao, C., and Rohacs, T. (2011). Decrease in phosphatidylinositol 4,5-bisphosphate levels mediates desensitization of the cold sensor TRPM8 channels. *J. Physiol.* 589, 6007–6027.
- Zakharian, E. (2013). Recording of ion channel activity in planar lipid bilayer experiments. *Methods Mol. Biol.* 998, 109–118.
- Zakharian, E., Thyagarajan, B., French, R.J., Pavlov, E., and Rohacs, T. (2009). Inorganic polyphosphate modulates TRPM8 channels. *PLoS ONE* 4, e5404.
- Zakharian, E., Cao, C., and Rohacs, T. (2010). Gating of transient receptor potential melastatin 8 (TRPM8) channels activated by cold and chemical agonists in planar lipid bilayers. *J. Neurosci.* 30, 12526–12534.