**Raising awareness of the plight of the critically endangered European mink in Spain is not
miscommunication – a response to Melero**

Lauren A Harrington¹, Madis Põdra², Asum Gómez³, Tiit Maran⁴

¹ Wildlife Conservation Research Unit, Zoology Department, University of Oxford, Recanati-Kaplan
Centre, Tubney House, Abingdon Road, Tubney, Oxfordshire OX13 5QL, UK

² European Mink Association, Av Diagonal 643, 2a planta, 08028 Barcelona, Spain

³ TRAGSATEC, Área de Vida Silvestre, C/Julián Camarillo 6A 4a planta, Sector B, 28037 Madrid

⁴ Institute of Veterinary Medicine and Animal Sciences, Estonian University of Life Sciences, Tartu,
Estonia; Species Conservation Lab, Tallinn Zoological Gardens, Ehitajate Road 150, Tallinn 13517,
Estonia

Corresponding author: Lauren A Harrington, lauren.harrington@zoo.ox.ac.uk, 01865 600126

We (TM, MP) estimate that fewer than 5,000 European mink persist, in less than 3% of their former range. The only remaining ‘large’ European populations are in the Danube Delta and northern Spain (Maran et al. 2016), and almost all populations are threatened by invasive American mink (Macdonald and Harrington 2003), established across Europe and Russia (Bouros et al. 2017). Effective control of American mink in Europe (and especially Spain) is crucial for European mink conservation and restoration. Whilst we applaud Melero’s (2017) attempt to address the success of American mink control in Spain, her suggestion that using the European mink as a native flagship species to raise awareness risks ‘miscommunication with the public’ is seriously misguided and potentially hugely damaging (in terms of public and political support) for on-going European mink conservation efforts.

Melero refers to speculation by Clavero (2014) that European mink in France and Spain may not be native (and thus not warrant conservation), based on their appearance in Spain in 1951, and France in 1839, and an incomplete interpretation of genetic analyses by Michaux et al. (2005) – an idea opposed by Díez-León et al. (2015) and Zuberogitia et al. (2016). French and Spanish European mink lack genetic variation, and likely arose from a small number of founders (Cabria et al. 2015). However, anthropogenic introduction is only one explanation – others (equally likely) are natural migration from Central Europe, and/or recent bottleneck events (Cabria et al. 2015). Additional information supports natural migration (Zuberogitia et al. 2016), and low genetic diversity at range margins is consistent with range expansion in other species (Excoffier et al. 2009).

Melero’s paper fuels a ‘controversy’ that does not exist, and lends weight to an idea that lacks scientific evidence, and her suggestion that the ‘details’ (of the apparent controversy) be explained to the general public would only create confusion, and provide excuse for inaction.

The relatively recent appearance of European mink in France/Spain is not fully understood but shifting distributional ranges are neither a unique, nor a new, phenomenon. The EU Habitats Directive protects listed species (including European mink) across their ‘natural range’, defined only as that to which the species could have ‘spread on its own...’ (Trouwborst et al. 2015). Where the mechanism

of ‘arrival’ (human-assisted or not) is ambiguous, Trouwborst suggests that ‘member states must opt for the course of action that is most likely to [ensure favourable conservation status]’. Indeed, Melero suggests that ‘any’ European mink population should be conserved for the ‘global benefit of the species’.

Melero draws comparisons with the Scottish mink removal project (Bryce et al. 2011), specifically the role of voluntary public participation, the success of which she attributes to the symbolism of water voles as Ratty in Kenneth Graham’s (1908) children’s classic, *The Wind in the Willows*. She fails to mention the earlier removal of American mink from the Scottish Hebridean Islands (www.snh.gov.uk), driven by popular desire to protect ground-nesting seabirds (e.g. Craik 1997). Both Scottish projects have been hugely successful.

We agree that there is much to learn from Scotland, but the lesson is not in downplaying the importance of the European mink. The lack of a charismatic European mink story character (to encourage participation in its conservation) is a pity, but hardly a basis on which to reject its worthiness as a flagship species. Further, there are other effective conservation models not based on voluntary labour, public participation, and childhood appeal. The Hebridean Mink Project used professional trappers and cost several million GBP (Munro 2014) in EU and government funding. (And in southern England, where *The Wind in the Willows* was written, there is no effective large-scale management of American mink despite the water-vole connection). We agree that the urgent need for action against American mink can be strengthened by raising awareness of impacts on other native species. However, there is strong scientific evidence (see Bouros et al. 2017) of the impact of American on European mink, the potential outcome (extinction of European mink) is clear, and awareness of this is crucial.

The success of the Scottish project is based on much more than the goodwill and involvement of water vole-loving volunteers – it is based on a large-scale, strategic, well-planned, well-monitored approach coordinated by a team of scientists, with the backing of national government (Scottish Mink Initiative, www.rafts.org.uk). Whether the workforce is voluntary (and there are potential problems

with this approach, Buesching et al. 2015) or paid (e.g through EU funding), this is what is needed in Spain, not the denigration of a critically endangered species.

Literature cited

Bouros G, Dekker J, Gómez A et al. (2016) *Neovison vison*. EU non-native organism risk assessment scheme. circabc.europa.eu/sd/a/a56cd4b4-4b2c-4b7f-979e-acda14ef2bfc/Neovison%20vison.pdf

Accessed 14 July 2017

Bryce R., Oliver MK, Davies L, Gray H, Urquhart J, Lambin X (2011). Turning back the tide of American mink invasion at an unprecedented scale through community participation and adaptive management. *Biol Conserv* 144:575-583.

Buesching CD, Slade E, Newman C, Riutta T, Riordan P, Macdonald DW (2015). Many hands make light work - but do they?: a critical evaluation of citizen science. In: Macdonald DW and Feber RE (eds) *Wildlife conservation on farmlands: conflict in the countryside* . Vol. 2. Oxford University Press, Oxford, pp293-317.

Cabria MT, González EG, Gómez-Moliner BJ et al (2015) Patterns of genetic variation in the endangered European mink (*Mustela lutreola* L., 1761). *BMC Evol Biol* 15:141

Clavero M (2014) Shifting baselines and the conservation of non-native species. *Conserv Biol* 28:1434–1436.

Craik C (1997) Long-term effects of North American mink *Mustela vison* on seabirds in western Scotland. *Bird Study* 44:303-309.

96 Díez-León M, Miranda R, Ariño AH, Galicia D (2015) Setting priorities for existing conservation
 97 needs of crayfish and mink. *Conserv Biol* 29:599–601.

98 Excoffier L, Foll M, Petit RJ (2009) Genetic consequences of range expansions. *Ann Rev Ecol Evol*
 99 *Syst* 40:481-501.

100 Macdonald DW, Harrington LA (2003) The American mink: the triumph and tragedy of adaptation
 101 out of context. *NZ J Zool* 30:421-441.

102 Maran T, Skumatov D, Gómez A, Põdra M, Abramov AV, Dinets V (2016) *Mustela lutreola*. IUCN
 103 Red List 2016: e.T14018A45199861. www.iucnredlist.org Accessed 13 July 2017.

104 Melero Y (2017) Communication of flagship species in conservation: lessons from invasive species
 105 management. *Biodivers Conserv*

106 Michaux JR, Hardy OJ, Justy F et al (2005) Conservation genetics and population history of the
 107 threatened European mink *Mustela lutreola*, with an emphasis on the west European population. *Mol*
 108 *Ecol* 14:2373–2388.

109 Munro A (2014) £6.4m cost of beating mink menace in Western Isles. *The Scotsman* 4 Feb
 110 2014. [http://www.scotsman.com/news/environment/6-4m-cost-of-beating-mink-menace-in-western-](http://www.scotsman.com/news/environment/6-4m-cost-of-beating-mink-menace-in-western-isles-1-3293703)
 111 [isles-1-3293703](http://www.scotsman.com/news/environment/6-4m-cost-of-beating-mink-menace-in-western-isles-1-3293703) Accessed 17 July 2017

112 Trouwborst A, Krofel M, Linnell JDC (2015) Legal implications of range expansions in a terrestrial
 113 carnivore: the case of the golden jackal (*Canis aureus*) in Europe. *Biodivers Conserv* 24:2593–2610.

114 Zuberogitia I, Podra M, Palazon S, Gomez A, Zabala N, Zabala J (2016) Misleading interpretation of
 115 shifting baseline síndrome in the conservation of European mink. *Biodivers Conserv* 25:1795-1800.

116
 117