

EGU2020-16003

<https://doi.org/10.5194/egusphere-egu2020-16003>

EGU General Assembly 2020

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



## Wave-resonance fingerprint in the 2010 summer: a modelling experiment

**Giorgia Di Capua**<sup>1,2</sup>, Kai Kornhuber<sup>3</sup>, Eftychia Rousi<sup>1</sup>, Sarah Sparrow<sup>4</sup>, David Wallom<sup>4</sup>, and Dim Coumou<sup>1,2</sup>

<sup>1</sup>Potsdam Institute for Climate Impact Research, Potsdam, Germany (dicapua@pik-potsdam.de)

<sup>2</sup>IVM-Institute for Environmental Studies, Free University of Amsterdam, Amsterdam, Netherlands

<sup>3</sup>Earth Institute, Columbia University, New York, United States

<sup>4</sup>Oxford e-Research Centre, University of Oxford, Oxford, United Kingdom

Summer 2010 was characterized by two contemporaneous extreme events: the Russian heat wave and the Pakistan flood. Several studies have shown a link between the two events, and Quasi-Resonant Amplification (QRA) has been suggested as an atmosphere-dynamic mechanism leading to the anomalous wavy circulation pattern which connected both extremes. Here, we aim at reproducing the 2010 circulation conditions in the Northern Hemisphere by obtaining a large ensemble of simulations from the Weather@home project within climateprediction.net (CPDN). We identify those ensemble members exhibiting a specific latitudinal temperature profile characterised by amplified high-latitude land warming (QRA - fingerprint) and investigate their surface temperature and upper level circulation properties. We show that when the QRA - fingerprint is present, the mid-latitude circulation bears similar characteristics to those observed in the 2010 summer: hot temperatures over European Russia and a wavy pattern in the upper-tropospheric meridional winds. As temperature profiles are projected to become increasingly similar to the QRA-fingerprint under future emission scenarios, these results provide further evidence that high latitude warming might favour persistent surface weather in the mid-latitudes.