

A NOTE ON THE RATIONAL HOMOLOGICAL DIMENSION OF LATTICES IN POSITIVE CHARACTERISTIC

SAM HUGHES

ABSTRACT. We show via ℓ^2 -homology that the rational homological dimension of a lattice in a product of simple simply connected Chevalley groups over global function fields is equal to the rational cohomological dimension and to the dimension of the associated Bruhat–Tits building.

1. INTRODUCTION

Let k be the function field of an irreducible projective smooth curve C defined over a finite field \mathbb{F}_q . Let S be a finite non-empty set of (closed) points of C . Let \mathcal{O}_S be the ring of rational functions whose poles lie in S . For each $p \in S$ there is a discrete valuation ν_x of k such that $\nu_p(f)$ is the order of vanishing of f at p . The valuation ring \mathcal{O}_p is the ring of functions that do not have a pole at p , that is

$$\mathcal{O}_S = \bigcap_{p \notin S} \mathcal{O}_p.$$

Let \bar{k} denote the algebraic closure of k . Let \mathbf{G} be an affine group scheme defined over \bar{k} such that $\mathbf{G}(\bar{k})$ is almost simple. For each $p \in S$ there is a completion k_p of k and the group $\mathbf{G}(k_p)$ acts on the Bruhat–Tits building X_p . Thus, we may embed $\mathbf{G}(\mathcal{O}_S)$ diagonally into the product $\prod_{p \in S} \mathbf{G}(k_p)$ as an arithmetic lattice.

The *rational cohomological dimension* of a group Γ is defined to be

$$\mathrm{cd}_{\mathbb{Q}}(\Gamma) := \sup\{n : H^n(\Gamma; M) \neq 0, \ M \text{ a } \mathbb{Q}\Gamma\text{-module}\},$$

the *rational homological dimension* is defined completely analogously as

$$\mathrm{hd}_{\mathbb{Q}}(\Gamma) := \sup\{n : H_n(\Gamma; M) \neq 0, \ M \text{ a } \mathbb{Q}\Gamma\text{-module}\}.$$

In [Gan12] it is shown that $\mathrm{cd}_{\mathbb{Q}}(\mathbf{G}(\mathcal{O}_S)) = \prod_{p \in S} \dim(X_p)$. In light of this Ian Leary asked the author what is $\mathrm{hd}_{\mathbb{Q}}(\mathbf{G}(\mathcal{O}_S))$?

MATHEMATICAL INSTITUTE, ANDREW WILES BUILDING, UNIVERSITY OF OXFORD, OXFORD OX2 6GG, UK

E-mail address: sam.hughes@maths.ox.ac.uk.

Date: 18th July, 2022.

2020 *Mathematics Subject Classification.* 20F67, 20J06.

Theorem A. *Let \mathbf{G} be a simple simply connected Chevalley group. Let k and \mathcal{O}_S be as above, then*

$$\mathrm{hd}_{\mathbb{Q}}(\mathbf{G}(\mathcal{O}_S)) = \mathrm{cd}_{\mathbb{Q}}(\mathbf{G}(\mathcal{O}_S)) = \prod_{p \in S} \dim(X_p).$$

More generally, we obtain the following.

Corollary B. *Let Γ be a lattice in a product of simple simply connected Chevalley groups over global function fields with associated Bruhat–Tits building X , then $\mathrm{hd}_{\mathbb{Q}}(\Gamma) = \mathrm{cd}_{\mathbb{Q}}(\Gamma) = \dim(X)$.*

The author expects these results are well-known, however, they do not appear in the literature so we take the opportunity to record them here.

Acknowledgements. The author would like to thank his PhD supervisor Professor Ian Leary for his guidance, support, and suggesting of the question. This note contains material from the author’s PhD thesis [Hug21a] and was originally part of [Hug21b], but was split off into a number of companion papers [Hug21c; Hug22] (see also [HV21]) at the request of the referee. This work was supported by the Engineering and Physical Sciences Research Council grant number 2127970. The author would like to thank the referee for a number of helpful comments.

Competing interests. The author declares none.

2. ℓ^2 -HOMOLOGY AND MEASURE EQUIVALENCE

Let Γ be a group. Both Γ and the complex group algebra $\mathbb{C}\Gamma$ act by left multiplication on the Hilbert space $\ell^2\Gamma$ of square-summable sequences. The *group von Neumann algebra* $\mathcal{N}\Gamma$ is the ring of Γ -equivariant bounded operators on $\ell^2\Gamma$. The non-zero divisors of $\mathcal{N}\Gamma$ form an Ore set and the Ore localization of $\mathcal{N}\Gamma$ can be identified with the *ring of affiliated operators* $\mathcal{U}\Gamma$.

There are inclusions $\mathbb{Q}\Gamma \subseteq \mathcal{N}\Gamma \subseteq \ell^2\Gamma \subseteq \mathcal{U}\Gamma$ and it is also known that $\mathcal{U}\Gamma$ is a self-injective ring which is flat over $\mathcal{N}\Gamma$. For more details concerning these constructions we refer the reader to [Lüc02] and especially to Theorem 8.22 of Section 8.2.3 therein. The *von Neumann dimension* and the basic properties we need can be found in [Lüc02, Section 8.3].

The ℓ^2 -Betti numbers of a group Γ , denoted $b_i^{(2)}(\Gamma)$, are then defined to be the von-Neumann dimensions of the homology groups $H_i(\Gamma; \mathcal{U}\Gamma)$. The following lemma is a triviality.

Lemma 2.1. *Let Γ be a discrete group and suppose that $b_i^{(2)}(\Gamma) > 0$, then the homology group $H_i(\Gamma; \mathcal{U}\Gamma)$ is non-trivial.*

Two countable groups Γ and Λ are said to be *measure equivalent* if there exist commuting, measure-preserving, free actions of Γ and Λ on some infinite Lebesgue measure space (Ω, m) , such that the action of each of the groups Γ and Λ admits a finite measure fundamental domain. The key examples of measure equivalent groups are lattices in the same locally-compact group [Gro93]. The relevance of this for us is the following deep theorem of Gaboriau.

Theorem 2.2 (Gaboriau’s Theorem [Gab02]). *Suppose a discrete group Γ is measure equivalent to a discrete group Λ , then $b_p(\Gamma) = 0$ if and only if $b_p(\Lambda) = 0$.*

3. PROOFS

Proof of Theorem A. We first note that the group $\Gamma := \mathbf{G}(\mathcal{O}_S)$ is measure equivalent to the product $\Lambda := \prod_{p \in S} \mathbf{G}(\mathbb{F}_q[t_p])$ for some suitably chosen $t_p \in \mathcal{O}_p$. By [PST18, Theorem 1.6] (see also [Dym04; Dym06; Dav+07]) the group $\mathbf{G}(\mathbb{F}_q[t_p])$ has one non-vanishing ℓ^2 -Betti number in dimension $\dim(X_p)$. Hence, by the Künneth formula Λ has one non-vanishing ℓ^2 -Betti number in dimension $d = \prod_{p \in S} \dim(X_p)$. Thus, by Gaboriau’s theorem, the group Γ has exactly one non-vanishing ℓ^2 -Betti number in dimension d . It follows from Lemma 2.1 that $\mathrm{hd}_{\mathbb{Q}}(\Gamma) \geq d$. The reverse inequality follows from the fact that Γ acts properly on the d -dimensional space $\prod_{p \in S} \dim(X_p)$. \square

Proof of Corollary B. The proof of the corollary is entirely analogous. First, we split \mathbf{G} into a product of simple groups $\prod_{i=1}^n \mathbf{G}_i$ corresponding to the decomposition of the Bruhat–Tits building $X = \prod_{i=1}^n X_i$. Let Λ_i be a lattice in \mathbf{G}_i and let $\Lambda = \prod_{i=1}^n \Lambda_i$. Each Λ_i has a non-vanishing ℓ^2 -Betti Number in dimension $\dim(X_i)$. In particular, Λ has a non-vanishing ℓ^2 -Betti Number in dimension $\dim(X) = \prod_{i=1}^n \dim(X_i)$. By Gaboriau’s Theorem Γ also has non-vanishing ℓ^2 -Betti Number in dimension $\dim(X)$. It follows from Lemma 2.1 that $\mathrm{hd}_{\mathbb{Q}}(\Gamma) \geq d$. The reverse inequality follows from the fact that Γ acts properly on the d -dimensional space $\prod_{p \in S} X_p$. \square

Remark 3.1. A similar argument can be applied to lattices in products of simple simply-connected algebraic groups over locally compact p -adic fields. One obtains the analogous result for such a lattice Γ that $\mathrm{cd}_{\mathbb{Q}}(\Gamma) = \mathrm{hd}_{\mathbb{Q}}(\Gamma) = \dim(X)$, where X is the associated Bruhat–Tits building.

REFERENCES

- [Dav+07] Michael W. Davis et al. “Weighted L^2 -cohomology of Coxeter groups”. In: *Geom. Topol.* 11 (2007), pp. 47–138. ISSN: 1465-3060. DOI: [10.2140/gt.2007.11.47](https://doi.org/10.2140/gt.2007.11.47).
- [Dym04] Jan Dymara. “ L^2 -cohomology of buildings with fundamental class”. In: *Proc. Amer. Math. Soc.* 132.6 (2004), pp. 1839–1843. ISSN: 0002-9939. DOI: [10.1090/S0002-9939-03-07234-4](https://doi.org/10.1090/S0002-9939-03-07234-4).

- [Dym06] Jan Dymara. “Thin buildings”. In: *Geom. Topol.* 10 (2006), pp. 667–694. ISSN: 1465-3060. DOI: [10.2140/gt.2006.10.667](https://doi.org/10.2140/gt.2006.10.667).
- [Gab02] Damien Gaboriau. “Invariants l^2 de relations d’équivalence et de groupes”. In: *Publ. Math. Inst. Hautes Études Sci.* 95 (2002), pp. 93–150. ISSN: 0073-8301. DOI: [10.1007/s102400200002](https://doi.org/10.1007/s102400200002).
- [Gan12] Giovanni Gandini. “Bounding the homological finiteness length”. In: *Bull. Lond. Math. Soc.* 44.6 (2012), pp. 1209–1214. ISSN: 0024-6093. DOI: [10.1112/blms/bds047](https://doi.org/10.1112/blms/bds047).
- [Gro93] M. Gromov. “Asymptotic invariants of infinite groups”. In: *Geometric group theory, Vol. 2 (Sussex, 1991)*. Vol. 182. London Math. Soc. Lecture Note Ser. Cambridge Univ. Press, Cambridge, 1993, pp. 1–295.
- [Hug21a] Sam Hughes. “Equivariant cohomology, lattices, and trees”. PhD thesis. School of Mathematical Sciences, University of Southampton, 2021.
- [Hug21b] Sam Hughes. *Graphs and complexes of lattices*. 2021. arXiv: [2104.13728](https://arxiv.org/abs/2104.13728) [math.GR].
- [Hug21c] Sam Hughes. “Lattices in a product of trees, hierarchically hyperbolic groups, and virtual torsion-freeness”. In: *Bull. London Math. Soc.* (2021). arXiv: [2105.02847](https://arxiv.org/abs/2105.02847) [math.GR]. To appear.
- [Hug22] Sam Hughes. *Irreducible lattices fibering over the circle*. 2022. arXiv: [2201.06525](https://arxiv.org/abs/2201.06525) [math.GR].
- [HV21] Sam Hughes and Motiejus Valiunas. *Commensurating HNN-extensions: hierarchical hyperbolicity and biautomaticity*. 2021. arXiv: [2203.11996](https://arxiv.org/abs/2203.11996) [math.GR].
- [Lüc02] Wolfgang Lück. *L^2 -invariants: theory and applications to geometry and K -theory*. Vol. 44. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, 2002, pp. xvi+595. ISBN: 3-540-43566-2. DOI: [10.1007/978-3-662-04687-6](https://doi.org/10.1007/978-3-662-04687-6).
- [PST18] Henrik Densing Petersen, Roman Sauer, and Andreas Thom. “ L^2 -Betti numbers of totally disconnected groups and their approximation by Betti numbers of lattices”. In: *J. Topol.* 11.1 (2018), pp. 257–282. ISSN: 1753-8416. DOI: [10.1112/topo.12056](https://doi.org/10.1112/topo.12056).