

UNIFORMLY SUPER MCDUFF II_1 FACTORS

ISAAC GOLDBRING, DAVID JEKEL, SRIVATSAV KUNNAWALKAM ELAYAVALLI, AND JENNIFER PI

ABSTRACT. We introduce and study the family of uniformly super McDuff II_1 factors. This family is shown to be closed under elementary equivalence and also coincides with the family of II_1 factors with the Brown property introduced in [1]. We show that a certain family of existentially closed factors, the so-called infinitely generic factors, are uniformly super McDuff, thereby improving a recent result of [6]. We also show that Popa's family of strongly McDuff II_1 factors are uniformly super McDuff. Lastly, we investigate when finitely generic II_1 factors are uniformly super McDuff.

1. INTRODUCTION

The study of central sequences in II_1 factors has been a central theme in operator algebras since the days of Murray and von Neumann, who introduced in [33] the notion of Property Gamma which states that nontrivial central sequences exist and is used to distinguish the hyperfinite II_1 factor $\mathcal{R}$ from the free group factor $L(\mathbb{F}_2)$. McDuff [32] leveraged the property of having abundantly many central sequences to construct continuum many non-isomorphic separable II_1 factors, settling a fundamental question open since the inception of the subject. In [31], McDuff showed that the central sequence algebra being noncommutative is equivalent to tensorially absorbing the hyperfinite II_1 factor; ever since then, this strengthening of property Gamma has been called the McDuff property and has been of great conceptual importance in the subject.

The model theory of II_1 factors, initiated in [12, 13, 14], has since been an active area of research in operator algebra theory (for a recent survey see [20] and the references therein). Among the various insights that model theory offers is in the analysis of ultrapowers; since the central sequences can be viewed as simply the commutant of the diagonal embedding of a II_1 factor M into its ultrapower $M^{\mathcal{U}}$, model theory is especially suited for studying central sequence algebras. One of the main avenues of research in the model theory of II_1 factors is to determine when two II_1 factors M and N elementarily equivalent, denoted $M \equiv N$, which, assuming the continuum hypothesis, is equivalent to asking if there exists an ultrafilter $\mathcal{U}$ such that the ultrapowers $M^{\mathcal{U}}$ and $N^{\mathcal{U}}$ are isomorphic. (In general, in this paper we assume the continuum hypothesis in order to be able to state such algebraic characterizations of model-theoretic properties. Additionally, throughout this introduction, all ultrafilters are assumed to be nonprincipal ultrafilters on $\mathbb{N}$.) The classification of II_1 factors up to elementary equivalence turns out to be much more challenging than the classification up to isomorphism (for instance, there are continuum many non-isomorphic separable II_1 factors that are elementarily equivalent to a given II_1 factor [14, Theorem 4.3]). The central sequences algebra has again played a crucial role in the elementary equivalence problem. Indeed, the first

I. Goldbring was partially supported by the National Science Foundation (NSF) grant DMS-2504477. D. Jekel was partially supported by NSF grant DMS-2002826. S. Kunawalkam Elayavalli was partially supported by NSF grant DMS-2350049.

handful of examples of non-elementarily equivalent II_1 factors (see [10, 11, 14, 21, 36]) and the first family of continuum many non-elementarily equivalent II_1 factors (interestingly coinciding with McDuff's factors; see [3]) pivotally use the structure of the central sequence algebra as the invariant (see also [7], which exhibits two non-elementarily equivalent II_1 factors without central sequences). Despite these results, it remains a challenging problem to further the classification up to elementary equivalence of II_1 factors with central sequences.

In this paper, we study this problem by identifying a proper subfamily of McDuff II_1 factors that are stable under elementary equivalence which we call *uniformly super McDuff* factors. The motivation for the terminology arises from [21], which introduced the super McDuff property, which is the property of admitting a II_1 factorial central sequence algebra and which has been of recent interest in the model theory of von Neumann algebras (see for instance [1, 6, 22, 27, 30]). To see the various examples and relationships between the super McDuff property and other subclasses of property Gamma factors, we direct the reader to [1, Section 6.4]. This paper introduces a uniform version of a finitary reformulation of the aforementioned super McDuff property, which we call the **uniform super McDuff property**. Throughout the paper, we write $(M)_1$ for the operator norm unit ball of a von Neumann algebra M .

Definition. The McDuff II_1 factor M is **uniformly super McDuff** if for all $n \geq 1$, there is $m \geq 1$ such that: for all $F \subseteq (M)_1$ with $|F| \leq n$, there is $G \subseteq (M)_1$ with $|G| \leq m$ for which, given any pair of projections $p, q \in M$ of the same trace with

$$\max_{x \in G} \max(\|[x, p]\|_2, \|[x, q]\|_2) < \frac{1}{m},$$

there is $u \in U(M)$ with $\|upu^* - q\|_2 < \frac{1}{n}$ and $\max_{x \in F} \|[x, u]\|_2 < \frac{1}{n}$.

As a first remark, we recall a notion first introduced in [1] and studied there in connection with proving a generalized version of a key result of Jung [29] (see also [2]).

Definition. A separable II_1 factor M has the **Brown property** if for every separable subfactor $N \subseteq M^{\mathcal{U}}$, there is a separable subfactor $N \subseteq P \subseteq M^{\mathcal{U}}$ with $P' \cap M^{\mathcal{U}}$ a II_1 factor.

Recalling that we are assuming the continuum hypothesis, any two nonprincipal ultrapowers of M are isomorphic, implying that the definition of the Brown property is independent of the choice of ultrafilter. Similarly, one sees that if M has the Brown property and N is elementarily equivalent to M , then N also has the Brown property. Our first main result shows that the uniform super McDuff property is closed under elementary equivalence and in fact coincides with the Brown property:

Theorem A. *For a II_1 factor M , the following are equivalent:*

- (1) M is uniformly super McDuff.
- (2) Every $N \equiv M$ is uniformly super McDuff.
- (3) Every $N \equiv M$ is super McDuff.
- (4) $M^{\mathcal{U}}$ is super McDuff for every nonprincipal ultrafilter $\mathcal{U}$ on $\mathbb{N}$.
- (5) $M^{\mathcal{U}}$ is super McDuff for some nonprincipal ultrafilter $\mathcal{U}$ on $\mathbb{N}$.
- (6) Every separable $N \equiv M$ is super McDuff.
- (7) M has the Brown property.

We then proceed to show that two particular families of super McDuff II_1 factors are in fact uniformly super McDuff (and hence also have the Brown property):

Theorem B. *Any II_1 factor that is elementarily equivalent to any of following separable II_1 factors is uniformly super McDuff:*

- (1) An *infinitely generic* II_1 factor.
- (2) $N \otimes \mathcal{R}$ where N is a separable full (that is, non-Gamma) II_1 factor.

The first family is a certain subclass of the family of existentially closed factors (see [15]) arising in the model theory of II_1 factors. Our result for this case improves upon the work of [6], where it was shown that infinitely generic II_1 factors are super McDuff. Note that our result also obtains uniform super McDuffness for factors elementarily equivalent to infinitely generic factors, some of which are not infinitely generic themselves (see Lemma 2.4).

The second family of factors in the above list are usually referred to as **strongly McDuff** factors and were first considered by Popa in [34]. Strongly McDuff II_1 factors were shown to be super McDuff in [1, Proposition 6.1.5].

Theorems A and B produce several new concrete examples of factors with the Brown property. Indeed, before this work, the only known examples of II_1 factors with the Brown property were those factors elementarily equivalent to $\mathcal{R}$ [1, 22], which do not fall into either of the two categories in the above theorem.

Having proven that the infinitely generic factors have the uniform super McDuff property, it is natural to inquire whether another important subclass of the existentially closed factors, namely the finitely generic factors, have this property. Here, a third property, which we call the (T)-factorial relative commutant property, or (T)-FRC property, becomes relevant. We say that a separable II_1 factor M has the **(T)-FRC property** if and only if every separable II_1 factor with property (T) embeds into M^U with factorial relative commutant. We show below that this property is an axiomatizable property of separable II_1 factors. In [18], it was shown that the infinitely generic factors have the (T)-FRC property. Here, we connect the question of whether or not finitely generic factors have the (T)-FRC property and whether or not they are uniformly super McDuff:

Theorem C. *If the finitely generic factors have the (T)-FRC property, then they are also uniformly super McDuff.*

We end this introduction by stating the main open question in connection to our work:

Question D. Is the class of II_1 factors with the super McDuff property closed under elementary equivalence? Equivalently, do the classes of super McDuff and uniformly super McDuff factors coincide?

Organization of the paper. In Section 2, we recall some background on the model theory of tracial von Neumann algebras, including a discussion of the classes of existentially closed, infinitely generic, and finitely generic factors. For a more detailed background on related matters, we direct the reader to [1, Section 2]. Section 3 is where the uniform super McDuff property is introduced and Theorem A is proved. Section 4 proves Theorem B via a unified approach involving a certain spectral gap criterion. In Section 5, Theorem C is proved and various remarks surrounding the problem of whether finitely generic II_1 factors are uniformly super McDuff are documented.

Acknowledgements. It is our pleasure to thank Adrian Ioana for suggesting to us the idea of using Hastings' result for proving Theorem B in Section 4.3. We thank the editor and referee for communicating to us the new argument added in Remark 5.10 (1).

2. PRELIMINARIES

2.1. Existentially closed, infinitely generic, and finitely generic factors.

Definition 2.1. A separable tracial von Neumann algebra M is said to be **existentially closed (e.c.)** if: for any tracial von Neumann algebra N containing M , there is an embedding $N \hookrightarrow M^{\mathcal{U}}$ that restricts to the diagonal embedding $M \hookrightarrow M^{\mathcal{U}}$.

The definition given above is the semantic definition and uses the separability assumption on M . The syntactic definition, which makes no separability assumption, reads: M is e.c. if and only if, for every tracial von Neumann algebra N extending M and every quantifier-free formula $\varphi(x)$ with parameters from M , we have

$$\inf\{\varphi(a)^M : a \in M\} = \inf\{\varphi(b)^N : b \in N\}.$$

E.c. tracial von Neumann algebras are McDuff II_1 factors (see, for example, [24, Section 2]); for this reason, we often refer to e.c. tracial von Neumann algebras as e.c. II_1 factors. Moreover, the class of e.c. tracial von Neumann algebras is **embedding universal**, meaning that every separable II_1 factor is contained in a separable e.c. II_1 factor. (This is a special case of a general model-theoretic fact; see [5, Lemma 3.5.7] for a proof in classical logic.)

Before moving on, we make a short digression to point out some facts about e.c. factors that follow from the results of [1] but were not explicitly pointed out there. In [1, Question 6.3.1], it was asked whether or not every e.c. factor is super McDuff. The following yields a criterion for when an e.c. factor is super McDuff:

Theorem 2.2. *For an e.c. factor M , the following are equivalent:*

- (1) M is super McDuff.
- (2) For any e.c. factor N with $M \subseteq N$, we have that $M' \cap N^{\mathcal{U}}$ is a factor.
- (3) There is an e.c. factor N with $M \subseteq N$ for which $M' \cap N^{\mathcal{U}}$ is a factor.

Proof. For the implication (1) implies (2), suppose that M is super McDuff and N is an e.c. factor with $M \subseteq N$. By considering the chain $M \subseteq M^{\mathcal{U}} \subseteq N^{\mathcal{U}}$, we see that $M' \cap N^{\mathcal{U}}$ is a factor using [1, Theorem 7.1.5(2), Lemma 7.2.1, and Lemma 7.2.8].

The implication (2) implies (3) is trivial. For the implication (3) implies (1), suppose that N is e.c., $M \subseteq N$, and $M' \cap N^{\mathcal{U}}$ is a factor. We then have that $M \subseteq M^{\mathcal{U}}$ is a factor using [1, Theorem 7.1.5(1) and Lemma 7.2.1]. $\square$

In [1, Question 6.3.10], it was asked if a union of a chain of super McDuff factors is once again super McDuff. Using Theorem 2.2, we can show that the answer to this question is positive if all of the super McDuff factors in the chain are also e.c. (which is a strengthening of [1, Proposition 6.3.17]):

Corollary 2.3. *Any union of a chain of e.c. super McDuff factors is (e.c.) super McDuff.*

Proof. Let $M_0 \subseteq M_1 \subseteq M_2 \dots$ be a chain of e.c. super McDuff factors, and let N denote their union. Then by Theorem 2.2, $M'_n \cap N^{\mathcal{U}}$ is a factor for every n . It follows that $N' \cap N^{\mathcal{U}}$ is a factor by [1, Proposition 6.3.12]. $\square$

We now return to the main thread of this section. In this paper, we will be concerned with specific subclasses of the class of e.c. factors. We first consider the class of **infinitely generic factors**, which is defined to be the largest class of II_1 factors that is embedding universal and which is **model-complete**, meaning any embedding $i : M \hookrightarrow N$ between two infinitely generic factors is elementary, that is, extends to an isomorphism $M^{\mathcal{U}} \rightarrow N^{\mathcal{U}}$. (The existence of such a class follows from [15, Corollary 5.15]. The terminology infinitely generic refers to the use of so-called infinite model-theoretic forcing in the construction of these factors; see [15, Section 5].) It follows from [15, Lemma 5.20] that any two infinitely generic II_1 factors are elementarily equivalent. That infinitely generic factors are e.c. follows from [15, Proposition 5.11].

In connection with Theorem 4.12 below, we need to record the following:

Lemma 2.4. *There is a II_1 factor M that is elementarily equivalent to the infinitely generic factors but is not itself infinitely generic.*

Proof. Let T^* denote the common theory of the infinitely generic factors. If the statement of the lemma is false, then the models of T^* are precisely the infinitely generic factors themselves. Since the collection of infinitely generic factors is embedding universal and model-complete, it follows that T^* is the model companion of the theory of tracial von Neumann algebras, contradicting the main result of [24]. $\square$

The other class of e.c. factors we will need to consider are the **finitely generic factors**. Any definition of these factors involves a discussion of finite model-theoretic forcing (see, for example, [15, Section 6] or [16, Section 3]). However, for our purposes, we will only need to know the following two properties of a finitely generic factor N (besides their existence):

- N is **locally universal**, meaning that every separable II_1 factor admits an embedding into $N^{\mathcal{U}}$;
- N has the **generalized Jung property**, meaning that every embedding $N \hookrightarrow N^{\mathcal{U}}$ is elementary.

The first item follows from [16, Propositions 2.10 and 3.9] and the fact that e.c. models of theories with the joint embedding property are locally universal (see [16, Subsection 1.1]). The second item follows from [16, Proposition 3.10].

Whether these two properties actually characterize being finitely generic is an open problem (see [1, Question 3.3.8.]).

It follows immediately from these two properties that any finitely generic factor is e.c. It is unknown whether or not the theories of finitely generic and infinitely generic II_1 factors coincide, or equivalently, whether or not any finitely generic factor must be infinitely generic. (In general, there need not be any connection between the finitely generic and infinitely generic models of a given theory.)

2.2. Definability. We freely use the terminology around definability adopted in [19]. Throughout this subsection, T denotes the theory of tracial von Neumann algebras in the appropriate continuous signature.

For a tracial von Neumann algebra M , we let $\mathcal{P}_2(M)$ denote the set of pairs of projections from M of the same trace. The relevance of the set $\mathcal{P}_2(M)$ is that M is a factor if and only if, for any $(p, q) \in \mathcal{P}_2(M)$, there is a unitary $u \in M$ with $upu^* = q$.

The assignment $M \mapsto \mathcal{P}_2(M)$ is a T -functor which is the zeroset of the T -formula

$$\xi(p, q) := \max(\|p^* - p\|_2, \|p^2 - p\|_2, \|q^* - q\|_2, \|q^2 - q\|_2, |\tau(p) - \tau(q)|).$$

Moreover, an easy functional calculus argument shows that

$$\mathcal{P}_2\left(\prod_{\mathcal{U}} M_i\right) = \prod_{\mathcal{U}} \mathcal{P}_2(M_i)$$

for all families $(M_i)_{i \in I}$ of tracial von Neumann algebras and all ultrafilters $\mathcal{U}$ on I , whence $\mathcal{P}_2$ is a T -definable set by [19, Theorem 2.13].

Consequently, for any T -formula $\varphi(p, q, \vec{z})$, the T -functor $\sup_{(p, q) \in \mathcal{P}_2} \varphi(p, q, \vec{z})$ is a T -formula. It will behoove us to understand the relationship between the quantifier-complexity of $\sup_{(p, q) \in \mathcal{P}_2} \varphi(p, q, \vec{z})$ and of $\varphi(p, q, \vec{z})$ itself. To understand this relationship, we first note that, as T -formulae, one has

$$\sup_{(p, q) \in \mathcal{P}_2} \varphi(p, q, \vec{z}) = \sup_{\vec{x}} (\varphi(\vec{x}, \vec{z}) \dot{-} \alpha(d(\vec{x}, \mathcal{P}_2))),$$

where $\alpha : [0, 1] \rightarrow [0, 1]$ is an increasing continuous function with $\alpha(0) = 0$ satisfying

$$|\varphi(\vec{x}, \vec{z}) - \varphi(\vec{y}, \vec{z})| \leq \alpha(d(\vec{x}, \vec{y}))$$

for all tuples $\vec{x}, \vec{y}, \vec{z}$ in models of T . Since ξ is quantifier-free, one can show that $d(\vec{x}, \mathcal{P}_2)$ is T -equivalent to an existential formula (see [23, Section 2], whence so is $\alpha(d(\vec{x}, \mathcal{P}_2))$ as α is increasing. As a result, if $\varphi(p, q, \vec{z})$ is $\forall_n$ (resp. $\exists_n$), then the $\sup_{(p, q) \in \mathcal{P}_2} \varphi(p, q, \vec{z})$ is $\forall_n$ (resp. $\forall_{n+1}$).

We will also find the need to take the infimum over the T -definable set U of unitaries, which is the zeroset of the T -formula

$$v(u) = \max(\|u^*u - 1\|_2, \|uu^* - 1\|_2).$$

In this case, for any T -formula $\varphi(u, \vec{z})$, there is a continuous function γ such that, as T -formulas, we have

$$\inf_{u \in U} \varphi(u, \vec{z}) = \inf_x (\varphi(x, \vec{z}) + \gamma(v(x)))$$

(see the proof of [19, Theorem 2.13]). As a result, if $\varphi(u, \vec{z})$ is $\forall_n$ (resp. $\exists_n$), then $\inf_{u \in U} \varphi(u, \vec{z})$ is $\exists_{n+1}$ (resp. $\exists_n$).

3. THE UNIFORM SUPER MCDUFF PROPERTY

The following is a finitary (ultraproduct-free) reformulation of having factorial relative commutant:

Proposition 3.1. *If M is a separable II_1 factor, then $M' \cap M^{\mathcal{U}}$ is a factor if and only if: for every $\epsilon > 0$ and every finite $F \subseteq (M)_1$, there are $\delta > 0$ and finite $G \subseteq (M)_1$ such that: for all $(p, q) \in \mathcal{P}_2(M)$, if $\|[x, p]\|_2, \|[x, q]\|_2 < \delta$ for all $x \in G$, then there is $u \in U(M)$ such that $\|upu^* - q\|_2 < \epsilon$ and $\|[x, u]\|_2 < \epsilon$ for all $x \in F$.*

Proof. First suppose that $M' \cap M^{\mathcal{U}}$ is a factor and yet, towards a contradiction, the condition fails for a particular ϵ and F . Let (G_n) denote a sequence of finite subsets of $(M)_1$ whose union is SOT-dense in $(M)_1$. Then by the contradiction assumption, for each $n \geq 1$, there are projections $(p_n, q_n) \in \mathcal{P}_2(M)$ witnessing that $\delta = \frac{1}{n}$ and $G = G_n$ do not satisfy the conclusion of the forward direction for this particular choice of ϵ and F . Set $p = (p_n)_{\mathcal{U}}$ and $q = (q_n)_{\mathcal{U}}$. Note then that $(p, q) \in \mathcal{P}_2(M' \cap M^{\mathcal{U}})$. Since M is super McDuff, there is $u \in U(M' \cap M^{\mathcal{U}})$ such that $upu^* = q$. Consequently, for $\mathcal{U}$ -almost all n , we have that $\|u_n p_n u_n^* - q_n\|_2 < \epsilon$ and $\max_{x \in F} \|[x, u_n]\|_2 < \epsilon$, a contradiction to the choice of (p_n, q_n) .

For the converse direction, suppose the condition holds. Fix $(p, q) \in \mathcal{P}_2(M' \cap M^{\mathcal{U}})$; we aim to show that p and q are unitarily conjugate in $M' \cap M^{\mathcal{U}}$. By saturation, it suffices to show that, for each $\epsilon > 0$ and finite $F \subseteq (M)_1$, there is $u \in U(M^{\mathcal{U}})$ such that $\|upu^* - q\|_2 \leq \epsilon$ and $\max_{x \in F} \|[x, u]\|_2 \leq \epsilon$. Let δ and G be as in the condition for ϵ and F . Write $p = (p_n)_{\mathcal{U}}$ and $q = (q_n)_{\mathcal{U}}$ with $(p_n, q_n) \in \mathcal{P}_2(M)$ for all $n \in \mathbb{N}$. Note that, for $\mathcal{U}$ -almost all n , we have $\max_{x \in G} \max(\|[x, p_n]\|_2, \|[x, q_n]\|_2) < \delta$. Consequently, for these n , there is $u_n \in U(M)$ such that $\|u_n p_n u_n^* - q_n\|_2 < \epsilon$ and $\max_{x \in F} \|[x, u_n]\|_2 < \epsilon$. Let $u = (u_n)_{\mathcal{U}} \in U(M^{\mathcal{U}})$. Then $\|upu^* - q\|_2 \leq \epsilon$ and $\max_{x \in F} \|[x, u]\|_2 \leq \epsilon$, as desired. $\square$

Remark 3.2. While the original definition of super McDuff was only for separable II_1 factors, we can use the finitary criterion appearing in Proposition 3.1 to give a definition of super McDuffness for arbitrary II_1 factors (using also that there is a finitary definition of McDuffness that makes sense for arbitrary tracial von Neumann algebras; see [14, Proposition 3.9]).

The condition in Proposition 3.1 lends itself naturally to a uniform version, namely that, in the criterion of Proposition 3.1, both δ and the size of G may be taken to depend only on ϵ and the size of F . Said somewhat differently:

Definition 3.3. A McDuff II_1 factor M is **uniformly super McDuff** if and only if, for all $n \geq 1$, there is $m \geq 1$ such that: for all $F \subseteq (M)_1$ with $|F| \leq n$, there is $G \subseteq (M)_1$ with $|G| \leq m$ for which, given any $(p, q) \in \mathcal{P}_2(M)$ with

$$\max_{x \in G} \max(\|[x, p]\|_2, \|[x, q]\|_2) < \frac{1}{m},$$

there is $u \in U(M)$ with $\|upu^* - q\|_2 < \frac{1}{n}$ and $\max_{x \in F} \|[x, u]\|_2 < \frac{1}{n}$.

Remark 3.4. We make the above definition only for M which is already McDuff. This is because the other condition in the definition does not force M to be McDuff. Indeed, if M is non-Gamma, then $M' \cap M^{\mathcal{U}} = \mathbb{C}$, which is a factor, and the “uniform factoriality” condition in our definition holds automatically. Indeed, given $n \geq 1$, there is a finite subset $K \subseteq (M)_1$ and $m \geq 1$ such that, for all contractions $x \in M$, if $\max_{y \in K} \|[x, y]\|_2 < \frac{1}{m}$, then $d(x, \mathbb{C}) < \frac{1}{2n}$. Consequently, if $(p, q) \in \mathcal{P}_2(M)$ are such that $\max_{y \in K} \max(\|[p, y]\|_2, \|[q, y]\|_2) < \frac{1}{m}$, then $d(p, \mathbb{C}), d(q, \mathbb{C}) < \frac{1}{2n}$, whence $\|p - q\|_2 < \frac{1}{n}$ and we may take $u = 1$ in the above definition. This is the reason why we specify the definition above only for a II_1 factor M which is already McDuff.

Now we are ready to prove that the Brown property is equivalent to the uniform super McDuff property, as well as several other conditions:

Theorem 3.5. *For a II_1 factor M , the following are equivalent:*

- (1) M is uniformly super McDuff.

- (2) Every $N \equiv M$ is uniformly super McDuff.
- (3) Every $N \equiv M$ is super McDuff.
- (4) $M^{\mathcal{U}}$ is super McDuff for every nonprincipal ultrafilter $\mathcal{U}$ on $\mathbb{N}$.
- (5) $M^{\mathcal{U}}$ is super McDuff for some nonprincipal ultrafilter $\mathcal{U}$ on $\mathbb{N}$.
- (6) Every separable $N \equiv M$ is super McDuff.
- (7) M has the Brown property.

Proof. (1) implies (2): Suppose that M is uniformly super McDuff. Fix $n \geq 1$ and take the corresponding $m \geq 1$ as in the definition of uniformly super McDuff. Consider the sentence $\sigma_{n,m} = \sup_{x_1, \dots, x_n} \inf_{y_1, \dots, y_m} \sup_{(p,q) \in \mathcal{P}_2} \varphi_{n,m}(\vec{x}, \vec{y}, p, q)$, where $\varphi_{n,m}(\vec{x}, \vec{y}, p, q)$ is the following formula:

$$\min \left(\frac{1}{m} \div \max_{j=1, \dots, m} \max(\|[p, y_j]\|_2, \|[q, y_j]\|_2), \inf_{u \in U(M)} \max(\|upu^* - q\|_2, \max_{i=1, \dots, n} \|[u, x_i]\|_2) \div \frac{1}{n} \right).$$

Then $\sigma_{n,m}^M = 0$ by assumption, whence the same is true for any elementarily equivalent structure N . We use this to show that N is uniformly super McDuff. Fix $\eta > 0$ with $\eta < \min(\frac{1}{n}, \frac{1}{m})$; we claim that $\delta := \frac{1}{m} - \eta$ and m witnesses the uniform super McDuffness of N for finite sets of size at most n and with $\epsilon = \frac{2}{n}$. To see this, fix $x_1, \dots, x_n \in N$ and take $y_1, \dots, y_m \in N$ so that $\sup_{(p,q) \in \mathcal{P}_2} \varphi_{n,m}(\vec{x}, \vec{y}, p, q) < \eta$. Now suppose that $(p, q) \in \mathcal{P}_2(N)$ is such that

$$\max_{j=1, \dots, m} \max(\|[p, y_j]\|_2, \|[q, y_j]\|_2) < \frac{1}{m} - \eta.$$

It follows that the first entry of $\varphi_{n,m}(\vec{x}, \vec{y}, p, q)$ is greater than η , whence the second entry is smaller than η . Consequently, there is a unitary which conjugates p to within $\frac{1}{n} + \eta < \frac{2}{n}$ of q and which $\frac{1}{n} + \eta < \frac{2}{n}$ commutes with each x_i , as desired.

The implications (2) implies (3), (3) implies (4), and (4) implies (5) are all trivial. To see that (5) implies (6), assume that $M^{\mathcal{U}}$ is super McDuff and fix a separable $N \equiv M$. By considering an elementary embedding $N \hookrightarrow M^{\mathcal{U}}$, we may assume that N is an elementary substructure of $M^{\mathcal{U}}$. Fix $n \geq 1$ and finitely many $x_1, \dots, x_n \in N$. Since $M^{\mathcal{U}}$ is super McDuff, we get a corresponding $\delta > 0$ and finite $G \subseteq (M^{\mathcal{U}})_1$. If $m \geq \frac{1}{\delta}$ and $G = \{y_1, \dots, y_m\}$ (possibly with repetitions), we have that $(\sup_{(p,q) \in \mathcal{P}_2} \varphi_{n,m}(\vec{x}, \vec{y}, p, q))^{M^{\mathcal{U}}} = 0$, where $\varphi_{n,m}$ is as in the implication (1) implies (2). Since N is an elementary substructure of $M^{\mathcal{U}}$, it follows that

$$\left(\inf_{\vec{y}} \sup_{(p,q) \in \mathcal{P}_2} \varphi_{n,m}(\vec{x}, \vec{y}, p, q) \right)^N = 0.$$

If $z_1, \dots, z_m \in N$ witness that the existential is less than $\eta < \min(\frac{1}{n}, \frac{1}{m})$, as in the implication (1) implies (2), we see that $\delta = \frac{1}{m}$ and $z_1, \dots, z_m$ witness the super McDuffness of N for $x_1, \dots, x_n$ and the error $\frac{2}{n}$.

(6) implies (4): Fix $n \geq 1$ and $x_1, \dots, x_n \in M^{\mathcal{U}}$. Take a separable elementary substructure N of $M^{\mathcal{U}}$ containing $x_1, \dots, x_n$. By (6), N is super McDuff. Let $m \geq 1$ be such that $\delta = \frac{1}{m}$ and $y_1, \dots, y_m \in N$ are as in the definition of super McDuffness for $\epsilon = \frac{1}{n}$ and $x_1, \dots, x_n$. It follows that $(\sup_{(p,q) \in \mathcal{P}_2} \varphi_{n,m}(\vec{x}, \vec{y}, p, q))^N = 0$. By elementarity, we have that $(\sup_{(p,q) \in \mathcal{P}_2} \varphi_{n,m}(\vec{x}, \vec{y}, p, q))^{M^{\mathcal{U}}} = 0$. Thus, $\frac{1}{m}$ and $y_1, \dots, y_m$ witness super McDuffness of $M^{\mathcal{U}}$ for $\epsilon = \frac{1}{n}$ and $x_1, \dots, x_n$.

(4) implies (1): Suppose, towards a contradiction, that M is not uniformly super McDuff. Then there is some $n \geq 1$ such that for each $m \geq 1$, there are $x_1^m, \dots, x_n^m \in M$ such that no $G \subseteq (M)_1$ of size at most m witnesses the truth of super McDuffness for M . We can then consider the elements $x_j = (x_j^m)_{\mathcal{U}} \in M^{\mathcal{U}}$. By assumption, $M^{\mathcal{U}}$ is super McDuff, so there are $\delta > 0$ and finite $G = \{y_1, \dots, y_t\} \in M^{\mathcal{U}}$ that witness super McDuffness for the finite subset $\{x_1, \dots, x_n\}$ of $M^{\mathcal{U}}$ and $\epsilon = \frac{1}{n}$. Write $y_j = (y_j^m)_{\mathcal{U}}$ for each $j = 1, \dots, t$. By assumption, for each $m \geq t$, there is $(p_m, q_m) \in \mathcal{P}_2(M)$ with $\max_{j=1, \dots, t} (\|p_m, y_j^m\|_2, \|q_m, y_j^m\|_2) < \frac{1}{m}$ and yet no unitary in M conjugates p_m to within $\frac{1}{n}$ of q_m while also $\frac{1}{n}$ -commuting with each x_j^m for $j = 1, \dots, n$. Let $p = (p_m)_{\mathcal{U}}$ and $q = (q_m)_{\mathcal{U}}$, whence $(p, q) \in \mathcal{P}_2(M^{\mathcal{U}})$ and $[p, y_j] = [q, y_j] = 0$ for all $j = 1, \dots, t$. It follows that there is $u \in U(M^{\mathcal{U}})$ such that $\|upu^* - q\|_2 < \frac{1}{n}$ and $\max_{j=1, \dots, n} \|[u, x_j]\|_2 < \frac{1}{n}$. Write $u = (u_m)_{\mathcal{U}}$ with each $u_m \in U(M)$. Then for $\mathcal{U}$ -almost all m , we have $\|u_m p_m u_m^* - q_m\|_2 < \frac{1}{n}$ and $\max_{j=1, \dots, n} \|[u_m, x_j^m]\|_2 < \frac{1}{n}$, contradicting the choice of p_m and q_m .

At this point, we have that (1)-(6) are equivalent. However, as mentioned in the introduction, it was shown in [1, Proposition 6.2.4] that (6) and (7) are equivalent, finishing the proof of the theorem. $\square$

Remark 3.6. By the comments made at the end of Subsection 2.2, we see that the sentences appearing in the above proof are $\forall_4$ -sentences. The quantifier-complexity of these concrete sentences agrees with the fact established in [22], proven using Ehrenfeucht-Fraïssé games, that any II_1 factor with the same four quantifier theory as a II_1 factor with the Brown property also has the Brown property.

Theorem 3.5 above, along with the fact that $\mathcal{R}$ has the Brown property by [4, Theorem 6.9], gives a proof that $\mathcal{R}$ is uniformly super McDuff. In Remark 5.10 below, we will give a direct argument that $\mathcal{R}$ is uniformly super McDuff.

4. SPECTRAL GAP AND THE UNIFORM SUPER MCDUFF PROPERTY

In this section, we describe a sufficient condition for the uniform super McDuff property in terms of spectral gap, and deduce that infinitely generic II_1 factors and strongly McDuff II_1 factors are both uniformly super McDuff.

4.1. Spectral gap criterion. We begin by recalling the definition of spectral gap for a subalgebra of a tracial factor.

Definition 4.1. Let M be a tracial factor and $A \subseteq M$ a von Neumann subalgebra. Then $A \subseteq M$ has **spectral gap** if there exists a finite $F \subseteq (A)_1$ and a constant $C > 0$ such that, for all $\xi \in L^2(M)$, we have

$$\|\xi - E_{A' \cap M}(\xi)\|_2^2 \leq C \sum_{x \in F} \|x\xi - \xi x\|_2^2.$$

If, in addition, $|F| \leq C$, then we call C a **spectral gap number** for $A \subseteq M$. We let $\text{SG}(A, M)$ denote the smallest natural number that is a spectral gap number for A inside M . We also write $\text{SG}(M) = \text{SG}(M, M)$.

Remark 4.2. In particular, $M \subseteq M$ has spectral gap if there exists a finite $F \subseteq (M)_1$ and a constant $C > 0$ such that, for all $\xi \in L^2(M)$, we have

$$\|\xi - \tau(\xi) \cdot 1\|_2^2 \leq C \sum_{x \in F} \|x\xi - \xi x\|_2^2.$$

Connes showed in [8, Theorem 2.1] that a tracial factor $M \subseteq M$ has spectral gap if and only if M is full (equivalently M does not have property Gamma in the case of a II_1 factor).

For a tracial von Neumann algebra M , subsets $X, Y \subseteq M$, and $\epsilon > 0$, we write $X \subseteq_\epsilon Y$ to mean that every element of X is within ϵ (with respect to the 2-norm) of some element of Y .

Definition 4.3. We say that a tracial factor M has the **factorial commutant limit spectral gap property** or **FC limit spectral gap property** if there exists some constant $s > 0$ such that, for any finite $F \subseteq M$ and $\epsilon > 0$, there is some von Neumann subalgebra $A \subseteq M$ such that $A' \cap M$ is a factor, $\text{SG}(A, M) \leq s$, and $F \subseteq_\epsilon A$.

The following proposition provides a sufficient criterion for a McDuff II_1 factor to be uniformly super McDuff:

Proposition 4.4. *If M is a McDuff II_1 factor with the FC limit spectral gap property, then M is uniformly super McDuff.*

Proof. Fix $n \geq 1$. We claim that any $m > 24sn$ witnesses that M is uniformly super McDuff. Towards that end, fix $F \subseteq (M)_1$ with $|F| \leq n$. Choose $A \subseteq M$ such that $A' \cap M$ is a factor, $\text{SG}(A, M) \leq s$, and F is contained in the $(1/2n)$ -neighborhood of A . Then there exists $G \subseteq (A)_1$ such that $|G| \leq s$ and

$$\|\xi - E_{A' \cap M}(\xi)\|_2^2 \leq s \sum_{x \in G} \|x\xi - \xi x\|_2^2.$$

In particular, if $(p, q) \in \mathcal{P}_2(M)$ and $\max_{x \in G} \max(\|[x, p]\|_2, \|[x, q]\|_2) \leq 1/m$, then

$$\|p - E_{A' \cap M}(p)\|_2^2 \leq \frac{s^2}{m^2}.$$

Set $a = E_{A' \cap M}(p)$. Since $\|a^2 - p^2\|_2 < 2s/m$, we get $\|a - a^2\|_2 < 3s/m$. Set $\tilde{p} = \mathbf{1}_{[1/2, 1]}(a)$, a projection in $A' \cap M$. Since $|t - \mathbf{1}_{[1/2, 1]}(t)| \leq 2t(1 - t)$ for $t \in [0, 1]$, we have $\|\tilde{p} - p\|_2 \leq 2\|a(1 - a)\|_2 < 6s/m$. Similarly, there is a projection $\tilde{q} \in A' \cap M$ such that $\|\tilde{q} - q\|_2 < 6s/m$.

Next notice that $|\tau(\tilde{p}) - \tau(\tilde{q})| \leq 12s/m$. Since $A' \cap M$ is a factor, there exists a unitary u in $A' \cap M$ such that $u\tilde{p}u^* - \tilde{q}$ is plus or minus a projection of trace $< 12s/m$. In particular, $\|u\tilde{p}u^* - \tilde{q}\|_2 < 12s/m$, which implies $\|upu^* - q\|_2 < 24s/m < 1/n$.

Moreover, since we assumed that F is contained in the $(1/2n)$ -neighborhood of A and u belongs to $A' \cap M$, we have that $\|[u, x]\|_2 < 1/n$ for $x \in F$. Finally, note that $|G| \leq s \leq 24sn < m$, finishing the proof. $\square$

Remark 4.5. It is interesting to note that in the previous proof, the size of G only depends on the constant s and does not depend on n .

4.2. Infinitely generic factors are uniformly super McDuff. In this section, we use Proposition 4.4 to show that infinitely generic factors are uniformly super McDuff. This result improves the result of Chifan, Drimbe, and Ioana from [6] showing that infinitely generic factors are super McDuff. A main ingredient in the proof of the latter result is the embedding universality of property (T) factors, also proved in [6]. Our proof in this subsection also utilizes this fact (and its proof).

Recall that a II_1 factor N has **property (T)** if there is a finite $F \subseteq N$ and $\delta > 0$ such that, whenever $\mathcal{H}$ is a normal N - N bimodule with a unit vector $v \in \mathcal{H}$ satisfying $\max_{x \in F} \|x\xi - \xi x\| < \delta$, then there is a nonzero vector $\eta \in \mathcal{H}$ such that $x\eta = \eta x$ for all $x \in N$. If Γ is an ICC group, then $L(\Gamma)$ has property (T) if and only if Γ has property (T). Connes and Jones [9] also showed that property (T) for factors is equivalent to the following, seemingly stronger, property.

Theorem 4.6. *A II_1 factor has property (T) if and only if there is a finite subset $F \subseteq N$ (called a **Kazhdan set**) and a constant $K > 0$ such that, for every normal N - N bimodule $\mathcal{H}$, we have*

$$\|\xi - P_{\text{central}}(\xi)\|_{\mathcal{H}}^2 \leq K \sum_{x \in F} \|x\xi - \xi x\|_{\mathcal{H}}^2,$$

where P_{central} is the projection onto the subspace of central vectors in $\mathcal{H}$.

In particular, if $N \subseteq M$ is an inclusion of II_1 factors and N has property (T), then $L^2(M)$ is a normal N - N -bimodule and therefore N has spectral gap inside M . Moreover, Tan [35] showed the converse implication, namely that if a II_1 factor N has spectral gap in M for every inclusion $N \subseteq M$, then N has property (T). Hence, it is natural to leverage property (T) subalgebras of M together with our spectral gap criterion for the uniform super McDuff property.

Parallel to the spectral gap number in the previous section, we introduce the following notion.

Definition 4.7. We say that K is a **Kazhdan number** for a II_1 factor N with property (T) if there exists a finite $F \subseteq (N)_1$ with $|F| \leq K$ such that for every normal N - N -bimodule $\mathcal{H}$,

$$\|\xi - P_{\text{central}}(\xi)\|_{\mathcal{H}}^2 \leq K \sum_{x \in F} \|x\xi - \xi x\|_{\mathcal{H}}^2.$$

We denote by $K(N)$ the smallest natural number that is a Kazhdan number for N .

Observe that if N is a II_1 factor with property (T) and $N \subseteq M$ is an inclusion of tracial von Neumann algebras, then $\text{SG}(N, M) \leq K(N)$.

Definition 4.8. We say that a II_1 factor M has the **uniform local property (T)** if there exists some constant s such that, for any finite $F \subseteq M$ and $\epsilon > 0$, there is some II_1 factor $N \subseteq M$ with property (T) such that $K(N) \leq s$ and $F \subseteq_{\epsilon} N$.

Proposition 4.9. *Let M be an e.c. II_1 factor with the uniform local property (T). Then M has the FC limit spectral gap property, and hence is uniformly super McDuff.*

Proof. Let s be as in the definition of uniform local property (T). If $F \subseteq M$ is a finite subset, then there exists some II_1 factor $N \subseteq M$ with property (T) such that $K(N) \leq s$ and $F \subseteq_{\epsilon} N$. Note $\text{SG}(N, M) \leq K(N) \leq s$. Moreover, since M is e.c. and N has property (T), it follows that $N' \cap M$ is a factor (see [18, Theorem 2.8]). Hence, M has the FC limit spectral gap property. Moreover, since e.c. II_1 factors are McDuff, Proposition 4.4 implies that M is uniformly super McDuff. $\square$

Next, we will show that the previous proposition applies to *some* infinitely generic II_1 factors.

Proposition 4.10. *There exists an infinitely generic II_1 factor with the uniform local property (T).*

To prove this, we will use the following Theorem, which is extracted from the discussion at the end of the introduction to [6].

Theorem 4.11 ([6]). *There is a fixed countable property (T) group Γ such that: given any II_1 factor M , there is a II_1 factor N containing M that is generated by a group homomorphism $\pi : \Gamma \rightarrow U(N)$. Consequently, N has property (T) and a Kazhdan set for N can be taken to be the image under π of a Kazhdan set for Γ with associated Kazhdan constant equal to that of Γ . As a result, the Kazhdan numbers of N are uniformly bounded (independently of M) by the Kazhdan number $K(\Gamma)$ of Γ .*

Proof of Proposition 4.10. We construct a chain of II_1 factors

$$M_0 \subseteq N_0 \subseteq M_1 \subseteq N_1 \subseteq \dots$$

inductively as follows. Let M_0 be an infinitely generic II_1 factor. Once M_i is chosen, let N_i be a property (T) factor constructed using Theorem 4.11, whence its Kazhdan number is bounded by $K(\Gamma)$. Once N_i is chosen, let M_{i+1} be an infinitely generic II_1 factor containing N_i , which exists because infinitely generic II_1 factors are embedding universal.

Let M be the inductive limit of the chain. Then M has uniform local property (T) by construction. It is also infinitely generic by [1, Lemma 6.3.16]. $\square$

Theorem 4.12. *Every infinitely generic II_1 factor is uniformly super McDuff. Consequently, any II_1 factor that is elementarily equivalent to an infinitely generic II_1 factor is uniformly super McDuff.*

Proof. As noted in the preliminaries, any two infinitely generic II_1 factors are elementarily equivalent. Since the uniform super McDuff property is preserved by elementary equivalence, it suffices to show that *some* infinitely generic II_1 factor is uniformly super McDuff. By Proposition 4.10, there exists an infinitely generic II_1 factor with uniform local property (T), which must be uniformly super McDuff by Proposition 4.9. $\square$

Theorem 4.12 generalizes Chifan, Drimbe, and Ioana's result, which showed that infinitely generic II_1 factors are super McDuff [6, Theorem 6.4]. In fact, it is a strict generalization because there are II_1 factors elementarily equivalent to the infinitely generic ones without themselves being infinitely generic (Lemma 2.4).

Theorem 4.12 also allows us to connect two open questions discussed in Subsection 2.1: whether every e.c. factor is super McDuff and whether the union of a chain of super McDuff factors is once again super McDuff.

Proposition 4.13. *If the union of every chain of uniformly super McDuff factors is (uniformly) super McDuff, then every e.c. factor is (uniformly) super McDuff.*

Proof. Let P be an e.c. factor and embed P into an infinitely generic factor M . By Theorem 4.12, M is uniformly super McDuff. Fix an embedding $M \hookrightarrow P^{\mathcal{U}}$ that restricts to the diagonal embedding $P \hookrightarrow P^{\mathcal{U}}$. We then have the chain

$$P \subseteq M \hookrightarrow P^{\mathcal{U}} \subseteq M^{\mathcal{U}} \hookrightarrow P^{2\mathcal{U}} \subseteq M^{2\mathcal{U}} \dots$$

Note that all the embeddings $P^{k\mathcal{U}} \hookrightarrow P^{(k+1)\mathcal{U}}$ are elementary (being iterated ultrapowers of the diagonal embedding). Consequently, the limit Q of the chain is an elementary extension of P . Since M is uniformly super McDuff and each $M^{k\mathcal{U}}$ is elementarily equivalent to M , they are also uniformly super McDuff. By hypothesis, Q is (uniformly) super McDuff. Since (uniform) super McDuffness passes to elementary substructures, the result follows. $\square$

4.3. Strongly McDuff implies uniformly super McDuff. The following result was suggested to us by Adrian Ioana. Recall that a factor M is *full* or *non-Gamma* if $M' \cap M^{\mathcal{U}} = \mathbb{C}$. Although this terminology is most often applied to infinite-dimensional factors, the matrix algebra $M_k(\mathbb{C})$ is also full since $M_k(\mathbb{C})' \cap M_k(\mathbb{C})^{\mathcal{U}} = M_k(\mathbb{C})' \cap M_k(\mathbb{C}) = \mathbb{C}$.

Theorem 4.14. *Let M be a full tracial factor. Then $M \otimes \mathcal{R}$ has the FC limit spectral gap property and hence is uniformly super McDuff.*

In particular, by taking $M = \mathbb{C}$, we obtain a direct argument that $\mathcal{R}$ is uniformly super McDuff (which was already known because $\mathcal{R}$ has the Brown property). When M is a full II_1 factor, $M \otimes \mathcal{R}$ is often called a **strongly McDuff** II_1 factor. Strongly McDuff II_1 factors were shown to be super McDuff in [1, Proposition 6.1.5] and Theorem 4.14 generalizes this latter result to obtain that strongly McDuff factors are uniformly super McDuff. As shown in [1, Proposition 6.3.2], e.c. factors are never strongly McDuff, whence Theorem 4.14 represents a completely different class of examples of uniformly super McDuff factors from the infinitely generic factors above.

The main ingredient in the proof of Theorem 4.14 is the following result of Hastings on quantum expanders [25]. Here we only state a special case of Hastings' result; in Hastings' terminology, this is the Hermitian case with $D = 4$.

Theorem 4.15 (Hastings [25]). *Fix an even $d \geq 4$. There exist unitaries $U_{k,1}, \dots, U_{k,d}$ in $M_k(\mathbb{C})$ such that the following properties hold: Let $\Phi_k : M_k(\mathbb{C}) \rightarrow M_k(\mathbb{C})$ be defined by*

$$\Phi_k(X) = \frac{1}{d} \sum_{j=1}^d U_{k,j} X U_{k,j}^*.$$

Let $\lambda_2(\Phi_k)$ be the second largest eigenvalue of Φ_k , which is the largest eigenvalue of Φ_k on the orthogonal complement of $\mathbb{C}1$. Then

$$\lim_{k \rightarrow \infty} \lambda_2(\Phi_k) = \frac{2\sqrt{d-1}}{d}.$$

For us, the importance of Hastings' result is the following:

Corollary 4.16. *The spectral gap numbers for $M_k(\mathbb{C})$ satisfy $\limsup_{k \rightarrow \infty} \text{SG}(M_k(\mathbb{C})) \leq 8$. In particular, $\sup_k \text{SG}(M_k(\mathbb{C})) < \infty$.*

Proof. Fix an even $d \geq 4$ (in the end, we will use $d = 8$), and let Φ_k be as in the previous theorem. By the definition of $\lambda_2(\Phi_k)$, we have

$$\|X - \text{tr}_k(X)1\|_2 \leq \frac{1}{1 - \lambda_2(\Phi_k)} \|X - \Phi_k(X)\|_2.$$

We also have

$$\|X - \Phi_k(X)\|_2 = \left\| X - \frac{1}{d} \sum_{j=1}^d U_{k,j} X U_{k,j}^* \right\|_2 \leq \frac{1}{d} \sum_{j=1}^d \|X - U_{k,j} X U_{k,j}^*\|_2 = \frac{1}{d} \sum_{j=1}^d \|U_{k,j} X - X U_{k,j}\|_2.$$

Thus, by the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \|X - \text{tr}_k(X)1\|_2^2 &\leq \frac{1}{(1 - \lambda_2(\Phi_k))^2} \left(\frac{1}{d} \sum_{j=1}^d \|U_{k,j} X - X U_{k,j}\|_2 \right)^2 \\ &\leq \frac{1}{(1 - \lambda_2(\Phi_k))^2} \frac{1}{d} \sum_{j=1}^d \|U_{k,j} X - X U_{k,j}\|_2^2. \end{aligned}$$

Therefore,

$$\limsup_{k \rightarrow \infty} \text{SG}(M_k(\mathbb{C})) \leq \limsup_{k \rightarrow \infty} \max \left(d, \frac{1}{4(1 - \lambda_2(\Phi_k))^2} \right) = \max \left(d, \frac{1}{d(1 - 2\sqrt{d-1}/d)^2} \right).$$

The second term inside the maximum equals $d(d + 2\sqrt{d-1})/(d-2)^2$, which is less than d for $d \geq 8$, and hence by taking $d = 8$ we get the result of the corollary. $\square$

We will also use the following elementary observations about spectral gap numbers and tensor products. These results are known to experts, but we include the proof here for completeness (and because we pay attention to the spectral gap numbers).

Lemma 4.17. *Let M be a full tracial factor. Then for any tracial von Neumann algebra N , $M \otimes \mathbb{C}$ has spectral gap inside $M \otimes N$. In fact, if F and C witness spectral gap for M inside M , then $F \otimes 1$ and C witness spectral gap for $M \otimes \mathbb{C}$ inside $M \otimes N$. In particular, $\text{SG}(M, M \otimes N) \leq \text{SG}(M, M)$.*

Proof. Let S be the operator $\text{id} - \tau$ on $L^2(M)$ and let $T : L^2(M) \rightarrow L^2(M)^F$ be the linear map given by $T(\xi) = (x\xi - \xi x)_{x \in F}$. Then the spectral gap property of $M \subseteq M$ can be expressed by

$$\|S\xi\|^2 \leq C\|T\xi\|^2 \text{ for } \xi \in L^2(M),$$

or equivalently, $S^*S \leq C \cdot T^*T$. Note that $L^2(M \otimes N) \cong L^2(M) \otimes L^2(N)$ and $E_{\mathbb{C} \otimes N} = \tau_M \otimes \text{id}_{L^2(N)}$ as an operator on $L^2(M) \otimes L^2(N)$. Since $S^*S \leq C \cdot T^*T$, we have

$$(S \otimes \text{id}_{L^2(N)})^*(S \otimes \text{id}_{L^2(N)}) \leq C \cdot (T \otimes \text{id}_{L^2(N)})^*(T \otimes \text{id}_{L^2(N)}),$$

which translates to

$$\|\xi - E_{\mathbb{C} \otimes N}(\xi)\|_2^2 \leq C \sum_{x \in F} \|(x \otimes 1)\xi - \xi(x \otimes 1)\|_2^2,$$

yielding the desired result. $\square$

The following lemma follows from [8, Corollary 2.3], but we provide a direct proof using spectral gap numbers here.

Lemma 4.18. *Let M_1 and M_2 be full tracial factors. Then $M_1 \otimes M_2$ is also full. More precisely, for $j = 1, 2$, let F_j and C_j witness the spectral gap of $M_j \subseteq M_j$. Then $(F_1 \otimes 1) \cup (1 \otimes F_2)$ and $\max(C_1, C_2)$ witness spectral gap for $M_1 \otimes M_2$. Hence, $\text{SG}(M_1 \otimes M_2) \leq \text{SG}(M_1) + \text{SG}(M_2)$.*

Proof. Fix $\xi \in L^2(M_1 \otimes M_2)$ and set $1 \otimes \eta = E_{\mathbb{C} \otimes M_2}(\xi)$. By the previous lemma,

$$\|\xi - 1 \otimes \eta\|_2^2 \leq C_1 \sum_{x \in F_1} \|(x \otimes 1)\xi - \xi(x \otimes 1)\|_2^2.$$

By the spectral gap of $M_2 \subseteq M_2$,

$$\begin{aligned} \|\eta - \tau_N(\eta)\|_2^2 &\leq C_2 \sum_{y \in F_2} \|y\eta - \eta y\|_2^2 = C_2 \sum_{y \in F_2} \|(1 \otimes y)(1 \otimes \eta) - (1 \otimes \eta)(1 \otimes y)\|_2^2 \\ &= C_2 \sum_{y \in F_2} \|(1 \otimes y)E_{1 \otimes M_2}(\xi) - E_{\mathbb{C} \otimes M_2}(\xi)(1 \otimes y)\|_2^2 \\ &= C_2 \sum_{y \in F_2} \|E_{\mathbb{C} \otimes M_2}((1 \otimes y)\xi) - E_{\mathbb{C} \otimes M_2}(\xi(1 \otimes y))\|_2^2 \\ &\leq C_2 \sum_{y \in F_2} \|(1 \otimes y)\xi - \xi(1 \otimes y)\|_2^2. \end{aligned}$$

Now $\xi - 1 \otimes \eta$ is orthogonal to $\mathbb{C} \otimes M_2$ and in particular to $1 \otimes \eta - \tau_{M_2}(\eta)$, hence

$$\begin{aligned} \|\xi - \tau_{M_1 \otimes M_2}(\xi)\|_2^2 &= \|\xi - E_{\mathbb{C} \otimes M_2}(\xi)\|_2^2 + \|\eta - \tau_{M_2}(\eta)\|_2^2 \\ &\leq C_1 \sum_{x \in F_1} \|(x \otimes 1)\xi - \xi(x \otimes 1)\|_2^2 + C_2 \sum_{y \in F_2} \|(1 \otimes y)\xi - \xi(1 \otimes y)\|_2^2 \\ &\leq \max(C_1, C_2) \left(\sum_{x \in F_1} \|(x \otimes 1)\xi - \xi(x \otimes 1)\|_2^2 + \sum_{y \in F_2} \|(1 \otimes y)\xi - \xi(1 \otimes y)\|_2^2 \right). \end{aligned}$$

This concludes the proof of the Lemma. $\square$

We are now ready to prove Theorem 4.14.

Proof of Theorem 4.14. Let $F \subseteq M \otimes \mathcal{R}$ be a finite subset and let $\epsilon > 0$. Set $s = \text{SG}(M) + \sup_k \text{SG}(M_k(\mathbb{C}))$. Then there exists some tensor decomposition $\mathcal{R} = M_k(\mathbb{C}) \otimes \tilde{\mathcal{R}}$ such that F is contained in the ϵ -neighborhood of $M \otimes M_k(\mathbb{C}) \otimes \mathbb{C}$. Note also that

$$(M \otimes M_k(\mathbb{C}) \otimes \mathbb{C})' \cap (M \otimes \mathcal{R}) = \mathbb{C} \otimes \mathbb{C} \otimes \tilde{\mathcal{R}},$$

which is a factor. By Lemmas 4.17 and 4.18,

$$\text{SG}(M \otimes M_k(\mathbb{C}) \otimes \mathbb{C}, M \otimes \mathcal{R}) \leq \text{SG}(M \otimes M_k(\mathbb{C})) \leq \text{SG}(M) + \text{SG}(M_k(\mathbb{C})) \leq s.$$

Therefore, $M \otimes \mathcal{R}$ has the FC limit spectral gap property, as desired. $\square$

In [1, Proposition 6.1.7], it was shown that an infinite tensor product of full II_1 factors is super McDuff. It is natural to wonder whether or not there is a uniform version of this result. The proof of Theorem 4.14 immediately yields the following:

Proposition 4.19. *Suppose that (M_k) is a sequence of full II_1 factors for which*

$$\sup_k \text{SG}(M_1 \otimes \cdots \otimes M_k) < \infty.$$

Then $\bigotimes_k M_k$ is uniformly super McDuff.

Remark 4.20. It is unknown whether or not there is a uniform bound on the spectral gap number of full II_1 factors or, equivalently, whether or not the class of full II_1 factors is closed under ultraproducts. (Note that if there was not a uniform bound on the spectral gap number of full II_1 factors, then by taking the ultraproduct of a sequence of II_1 factors with strictly increasing spectral gap numbers, one would exhibit that the class of full II_1 factors is not closed under taking ultraproducts.) Proposition 4.19 shows that a positive answer to this question would imply that an infinite tensor product of full factors is uniformly super McDuff. Said differently, if one can find an example of an infinite tensor product of full II_1 factors which is not uniformly super McDuff, then one can conclude that the spectral gap numbers for full II_1 factors is unbounded.

5. DO FINITELY GENERIC FACTORS HAVE THE UNIFORM SUPER MCDUFF PROPERTY?

The results concerning the infinitely generic factors above motivate the following questions for the finitely generic factors:

Question 5.1. Is there a finitely generic (uniformly) super McDuff factor? Are all finitely generic factors (uniformly) super McDuff?

Remark 5.2. The question of whether the finitely generic factors are uniformly super McDuff relates to the Connes Embedding Problem. Indeed, if we assume a positive solution to the Connes Embedding Problem, then $\mathcal{R}$ is the unique finitely generic factor (see [1, Corollary 3.2.6]). Let (σ_n) be a sequence of sentences witnessing that $\mathcal{R}$ is uniformly super McDuff. By [16, Theorem 2.14 and Proposition 3.9], for each $n \geq 1$, there is a unique real number r_n such that $\sigma_n^M = r_n$ in all finitely generic factors M . If $r_n > 0$ for some n , then we conclude that the finitely generic factors are not uniformly super McDuff. This result might lead to a potentially different proof that the Connes Embedding Problem has a negative solution (see [28]).

5.1. A sufficient criterion for finitely generic factors to be uniformly super McDuff. In this subsection, we present a sufficient criterion for a finitely generic factor to be uniformly super McDuff. We first need a preliminary lemma.

Lemma 5.3. *For any $K \geq 1$, there is a function $\alpha_K : (0, 1) \rightarrow (0, 1)$ with the following property: Suppose that N has property (T) with finite Kazhdan set G and Kazhdan constant at most K . Then for any II_1 factor M containing N and any $(p, q) \in \mathcal{P}_2(M)$ with $\max_{x \in G} \max(\|x, p\|_2, \|x, q\|_2) < \alpha_K(\epsilon)$, there is $(p', q') \in \mathcal{P}_2(N' \cap M)$ with $\|p - p'\|_2, \|q - q'\|_2 < \epsilon$.*

Proof. Suppose that $\delta < \frac{1}{K}$ and $(p, q) \in \mathcal{P}_2(M)$ is such that

$$\max_{x \in G} \max(\|x, p\|_2, \|x, q\|_2) < \delta.$$

Then by the definition of Kazhdan set, there is $(p_0, q_0) \in N' \cap M$ such that $\|p - p_0\|_2, \|q - q_0\|_2 < K\delta$. Recall from Subsection 2.2 the formula ξ whose zeroset is $\mathcal{P}_2$. Let $\beta : \mathbb{R}^{>0} \rightarrow \mathbb{R}^{>0}$ be the uniform continuity modulus for ξ and $\gamma : \mathbb{R}^{>0} \rightarrow \mathbb{R}^{>0}$ be the definability modulus for ξ . Then if $K\delta < \beta(\gamma(\frac{\epsilon}{2}))$, we have $\xi(p_0, q_0) < \gamma(\frac{\epsilon}{2})$, whence there is $(p', q') \in \mathcal{P}_2(N' \cap M)$ such that $\|p_0 - p'\|_2, \|q_0 - q'\|_2 < \frac{\epsilon}{2}$. Thus, $\alpha_K(\epsilon) := \min(\frac{1}{K}\beta(\gamma(\frac{\epsilon}{2})), \frac{\epsilon}{2K})$ is the desired function. $\square$

Proposition 5.4. *Suppose that P is finitely generic and there is a II_1 factor $N \supseteq P$ with property (T) for which there is an embedding $N \hookrightarrow P^{\mathcal{U}}$ with factorial relative commutant. Then P is uniformly super McDuff.*

Proof. We consider the sequence of embeddings

$$P \subseteq N \hookrightarrow P^{\mathcal{U}} \subseteq N^{\mathcal{U}} \hookrightarrow P^{2\mathcal{U}} \subseteq N^{2\mathcal{U}} \dots,$$

where for each $k \in \mathbb{N}$, we let $P^{k\mathcal{U}}$ denote the k^{th} -iterated ultrapower of P and similarly for other iterated ultrapowers and the embeddings $N^{k\mathcal{U}} \hookrightarrow P^{(k+1)\mathcal{U}}$ are induced by the original embedding $N \hookrightarrow P^{\mathcal{U}}$. We let Q denote the union of the chain. Since P is finitely generic, we have that the embedding $P \hookrightarrow P^{\mathcal{U}}$ is elementary, whence so are all of the iterated embeddings $P^{k\mathcal{U}} \hookrightarrow P^{(k+1)\mathcal{U}}$, and thus $Q \equiv P$. It thus suffices to show that Q is uniformly super McDuff.

Let G be a finite Kazhdan set for N with Kazhdan constant K .

Claim: Fix $\epsilon > 0$. For all $k \geq 1$, and all $(p, q) \in \mathcal{P}_2(P^{k\mathcal{U}})$, if $\|[x, p]\|_2, \|[x, q]\|_2 < \alpha_K(\frac{\epsilon}{2})$ for all $x \in G$ (viewed as elements of $P^{k\mathcal{U}}$ in the obvious way), then there is $u \in U((N^{(k-1)\mathcal{U}})' \cap P^{k\mathcal{U}})$ such that $\|upu^* - q\|_2 \leq \epsilon$.

We prove the claim by induction on k . First suppose that $k = 1$. If

$$\max_{x \in G} \max(\|[x, p]\|_2, \|[x, q]\|_2) < \alpha_K\left(\frac{\epsilon}{2}\right),$$

then there is $(p', q') \in \mathcal{P}_2(N' \cap P^{\mathcal{U}})$ with $\|p - p'\|_2, \|q - q'\|_2 < \frac{\epsilon}{2}$. Since $N' \cap P^{\mathcal{U}}$ is a factor, there is $u \in U(N' \cap P^{\mathcal{U}})$ with $up'u^* = q'$. It follows that $\|upu^* - q\|_2 \leq \epsilon$. Thus, the case $k = 1$ is proven.

We now prove the inductive step. Suppose the result is true for k and consider $(p, q) \in \mathcal{P}_2(P^{(k+1)\mathcal{U}})$ with $\|[x, p]\|_2, \|[x, q]\|_2 < \alpha_K(\frac{\epsilon}{2})$ for all $x \in G$. Write $p = (p_n)_{\mathcal{U}}$ and $q = (q_n)_{\mathcal{U}}$ with $(p_n, q_n) \in \mathcal{P}_2(P^{k\mathcal{U}})$. It follows that for $\mathcal{U}$ -almost all n we have $\|[x, p_n]\|_2, \|[x, q_n]\|_2 < \alpha_K(\frac{\epsilon}{2})$ for all $x \in G$, whence, by induction, for these n , there are $u_n \in U((N^{(k-1)\mathcal{U}})' \cap P^{k\mathcal{U}})$ with $\|u_n p_n u_n^* - q_n\|_2 \leq \epsilon$. Letting $u = (u_n)_{\mathcal{U}} \in U((N^{k\mathcal{U}})' \cap P^{(k+1)\mathcal{U}})$, we see that $\|upu^* - q\|_2 \leq \epsilon$, as desired. The claim is thus proven.

We now verify that Q is uniformly super McDuff. Fix $\epsilon > 0$. Fix $F \subseteq Q$ finite. Take $k \geq 1$ such that $F \subseteq_{\frac{\epsilon}{2}} N^{(k-1)\mathcal{U}}$. Now suppose that $(p, q) \in \mathcal{P}_2(Q)$ are such that $\max_{x \in G} \max(\|[x, p]\|_2, \|[x, q]\|_2) < \rho$ (for $\rho > 0$ to be determined). Take $l \geq k$ such that there are $(p', q') \in \mathcal{P}_2(P^{l\mathcal{U}})$ with $\|p - p'\|_2, \|q - q'\|_2 < \rho$. Then $\max_{x \in G} \max(\|[x, p']\|_2, \|[x, q']\|_2) < \alpha_K(\frac{\epsilon}{4})$ if ρ was small enough, whence by the claim there is $u \in U((N^{(l-1)\mathcal{U}})' \cap P^{l\mathcal{U}})$ with $\|up'u^* - q'\|_2 \leq \frac{\epsilon}{2}$. It follows that $\|upu^* - q\|_2 < \epsilon$ and $\|[u, x]\|_2 < \epsilon$ for all $x \in F$. $\square$

Remark 5.5. We note that, in the previous proposition, one can obtain the weaker conclusion that P is super McDuff using [1, Theorem 3.2.2].

5.2. The (T)-FRC property. To apply Proposition 5.4 to a finitely generic P , we would need a property (T) factor N containing P which admits an embedding into $P^{\mathcal{U}}$ with factorial relative commutant. In fact, one can ask whether every property (T) factor N admits such an embedding, motivating the following definition.

Definition 5.6. We say that a separable II_1 factor M has the **(T)-FRC property** if every II_1 factor with property (T) admits an embedding into $M^{\mathcal{U}}$ with factorial relative commutant.

It will be a consequence of Proposition 5.12 below that whether or not a given property (T) factor admits an embedding into $M^{\mathcal{U}}$ with factorial relative commutant is independent of the choice of

ultrafilter, making the (T)-FRC property well-defined. Another consequence of Proposition 5.12 is that the (T)-FRC property is preserved under elementary equivalence.

We may also relativize the (T)-FRC property to the setting of Connes embeddable factors:

Definition 5.7. A Connes embeddable separable II_1 factor M has the **embeddable (T)-FRC property** if every Connes embeddable property (T) factor admits an embedding into $M^{\mathcal{U}}$ with factorial relative commutant.

A well-known question of Popa asks whether or not $\mathcal{R}$ has the embeddable (T)-FRC property.

Proposition 5.4 together with embedding universality of Property (T) factors yields the following:

Corollary 5.8. *If the finitely generic factors have the (T)-FRC property, then they are also uniformly super McDuff.*

Corollary 5.9. *If the (T)-FRC property is enforceable, then being uniformly super McDuff is enforceable. In particular, if the enforceable factor exists and has the (T)-FRC property, then it is uniformly super McDuff.*

Remarks 5.10.

- (1) The conclusions of Proposition 5.4 and Corollary 5.8 also hold with the class of II_1 factors replaced by the class of Connes-embeddable II_1 factors. As the discussion following [17, Theorem 2] establishes, $\mathcal{R}$ is the unique separable II_1 factor that is finitely generic in the class of Connes-embeddable II_1 factors. Hence, to show that $\mathcal{R}$ is uniformly super McDuff, it suffices to find a II_1 factor N with property (T) with $\mathcal{R} \subseteq N \subseteq \mathcal{R}^{\mathcal{U}}$ such that $N' \cap \mathcal{R}^{\mathcal{U}}$ is a factor. For this purpose, let G be a property (T) group with irreducible finite-dimensional representations $\pi_k : G \rightarrow U(M_{n_k}(\mathbb{C}))$ with $n_k \rightarrow \infty$ (for instance $G = SL_3(\mathbb{Z})$). Write $\mathcal{R}^{\mathcal{U}} \cong \prod_{k \rightarrow \mathcal{U}} (M_{n_k}(\mathbb{C}) \otimes \mathcal{R})$, and let $\pi : G \rightarrow U(\prod_{k \rightarrow \mathcal{U}} M_{n_k}(\mathbb{C}) \otimes \mathcal{R})$ be given by $\pi(g) = [\pi_k(g)]_{k \rightarrow \mathcal{U}}$. Then $N = \pi(G)'' \subseteq \mathcal{R}^{\mathcal{U}}$ will be a II_1 factor with property (T). Moreover, N contains a copy of $\mathcal{R}$ and all copies of $\mathcal{R}$ in $\mathcal{R}^{\mathcal{U}}$ are unitarily conjugate, and hence we can assume without loss of generality that $\mathcal{R} \subseteq N \subseteq \mathcal{R}^{\mathcal{U}}$. This gives a more direct argument for $\mathcal{R}$ being uniformly super McDuff. It is still open whether $\mathcal{R}$ has the (T)-FRC property.
- (2) It is interesting to note that in the class of Connes embeddable II_1 factors, the finitely generic factor is uniformly super McDuff although we do not know if the infinitely generic factors are uniformly super McDuff. (It is an open question as to whether or not $\mathcal{R}$ is an infinitely generic embeddable factor.) On the other hand, in the class of II_1 factors, as shown in Theorem 4.12, the infinitely generic factors are uniformly super McDuff and we are unsure whether the finitely generic factors are uniformly super McDuff.

Given Corollary 5.8, it is now natural to ask:

Question 5.11. Do finitely generic factors have the (T)-FRC property?

In [18], it was shown that the infinitely generic factors have the (T)-FRC property. Thus, if the finitely and infinitely generic factors are elementarily equivalent, then the finitely generic factors also have the (T)-FRC property.

5.3. Model-theoretic aspects of the (T)-FRC property. In this section, we explain why proving the (T)-FRC property for infinitely generic factors is easier than proving the (T)-FRC property for an arbitrary e.c. factor, which in turn is easier than proving the (T)-FRC property for finitely generic factors. We first observe:

Proposition 5.12. *For any property (T) factor N , there is a family $(\sigma_{N,m})_{m \geq 1}$ of $\exists_3$ -sentences such that the following are equivalent for any II_1 factor M :*

- (1) $\sigma_{N,m}^M = 0$ for all $m \geq 1$.
- (2) For every ultrafilter $\mathcal{U}$, N embeds into $M^\mathcal{U}$ with factorial relative commutant.
- (3) For some ultrafilter $\mathcal{U}$, N embeds into $M^\mathcal{U}$ with factorial relative commutant.

Proof. Fix a property (T) factor N with generating set F of cardinality n and Kazhdan constant K . For each $m \geq 1$, consider the following formula $\psi_{N,m}(\vec{x})$ in the free variables $\vec{x} = (x_1, \dots, x_n)$:

$$\sup_{(p,q) \in \mathcal{P}_2} \min \left(\alpha_K \left(\frac{1}{2m} \right) \div \max_{i=1, \dots, n} \max(\|[x_i, p]\|_2, \|[x_i, q]\|_2), \right. \\ \left. \left(\inf_{u \in U} \max(\|upu^* - q\|_2, \max_{i=1, \dots, n} \|[u, x_i]\|_2) \div \frac{1}{m} \right) \right) = 0,$$

where α_K is as in Lemma 5.3. Enumerate a countable dense subset of $\text{qftp}^N(F)$ as $\{\theta_{N,i}(\vec{x}) : i \in \omega\}$ and let $\sigma_{N,m}$ be $\inf_x \max(\max_{i=1, \dots, m} \theta_i(x), \psi_{N,m}(x))$. Note that $\psi_{N,m}(\vec{x})$ is a $\forall_2$ -formula, whence $\sigma_{N,m}$ is a $\exists_3$ -sentence. We claim that these are the desired sentences.

First suppose that $\sigma_{N,m}^M = 0$ for all $m \geq 1$. Then by saturation, there is a tuple $\vec{a} = (a_1, \dots, a_n)$ in $M^\mathcal{U}$ satisfying $\text{qftp}^N(F)$ and for which $\psi_{N,m}^M(\vec{a}) = 0$ for all $m \geq 1$. There is thus an embedding $N \hookrightarrow M^\mathcal{U}$ obtained by sending the elements of F to the a_i 's. We claim that this embedding has factorial relative commutant. To see this, fix $(p, q) \in \mathcal{P}_2(N' \cap M^\mathcal{U})$ and $m \geq 1$. Then the first part of the minimum appearing in $\psi_{N,m}$ is $\alpha_K(\frac{1}{2m})$, whence the second part of the minimum must be 0. Thus, there is $u \in U(M^\mathcal{U})$ such that $\|upu^* - q\|_2 \leq \frac{1}{m}$ and $\|[u, a_i]\|_2 \leq \frac{1}{m}$. Since such a unitary exists for all $m \geq 1$, by saturation, there is $u \in U(N' \cap M^\mathcal{U})$ conjugating p to q , as desired.

Now suppose that N admits an embedding into $M^\mathcal{U}$ with factorial relative commutant for some ultrafilter $\mathcal{U}$. Let $a_1, \dots, a_n \in M^\mathcal{U}$ be the images of the elements of F under the embedding. Fix $m \geq 1$. We show that $\psi_{N,m}^M(\vec{a}) = 0$, whence $\sigma_{N,m}^M = \sigma_{N,m}^{M^\mathcal{U}} = 0$. Towards this end, fix $(p, q) \in \mathcal{P}_2(M^\mathcal{U})$ and suppose $\|[a_i, p]\|_2, \|[a_i, q]\|_2 < \alpha_K(\frac{1}{2m})$ for all $i = 1, \dots, n$. Arguing exactly as in the proof of Proposition 5.4, there is $u \in U(N' \cap M^\mathcal{U})$ satisfying $\|upu^* - q\|_2 \leq \frac{1}{m}$. This u realizes the inner infimum in $\psi_{N,m}$. $\square$

Corollary 5.13. *The (T)-FRC property and the embeddable (T)-FRC property are both $\exists_3$ -axiomatizable properties of separable II_1 factors.*

Remarks 5.14.

- (1) One could use the sentences in Proposition 5.12 to give a definition of the (T)-FRC property for arbitrary II_1 factors.

- (2) It is worth pointing out that the (T)-FRC property is axiomatizable while we currently only know that the Brown property is merely *local*, that is, preserved under elementary equivalence.

Let us recall the following various facts about truth of $\exists_3$ sentences in e.c. structures, which, for simplicity, we state only for II_1 factors (although the first item holds in general and the second and third item hold in any $\forall_2$ theory with the joint embedding property). The classical versions of these results appear as [26, Corollary 3.2.5, Exercise 5.3.12, Theorem 4.3.4].

Fact 5.15. *Suppose that σ is a $\exists_3$ -sentence in the language of tracial von Neumann algebras.*

- (1) *If $M \subseteq N$ are e.c. factors and $\sigma^M = 0$, then $\sigma^N = 0$. (Here, σ can even have parameters from M .)*
- (2) *If $\sigma^M = 0$ for some e.c. factor, then $\sigma^N = 0$ for all infinitely generic factors.*
- (3) *If $\sigma^M = 0$ for some finitely generic factor, then $\sigma^N = 0$ for all e.c. factors.*

Proof of (1) and (2). For (1), fix an embedding $i : N \hookrightarrow M^{\mathcal{U}}$ restricting to the diagonal embedding $M \hookrightarrow M^{\mathcal{U}}$. Write $\sigma = \inf_x \sup_y \inf_z \varphi(x, y, z)$, where φ is a quantifier-free formula with parameters from M . Since $\sigma^M = 0$, there is $a \in M$ such that $(\sup_y \inf_z \varphi(a, y, z))^M = 0$. Fix $b \in N$; we need $(\inf_z \varphi(a, b, z))^N = 0$. However, since N is e.c., this follows from $(\inf_z \varphi(a, i(b), z))^{M^{\mathcal{U}}} = 0$, which in turn follows from the fact that $(\sup_y \inf_z \varphi(a, y, z))^{M^{\mathcal{U}}} = 0$.

(2) follows from (1) together with the facts that the infinitely generic factors are e.c., embedding universal, and are all elementarily equivalent to one another. $\square$

The third item in the above proposition can be proven just as in the classical case, using the continuous versions of the necessary ingredients which were established in [16].

The second item in the previous fact explains why proving that the infinitely generic factors have the (T)-FRC property was the “easiest” possible while the third item explains why proving that $\mathcal{R}$ has the embeddable (T)-FRC property and the finitely generic factors have the (T)-FRC property are as “hard” as possible.

Item (1) of the above fact also yields the following fact:

Corollary 5.16. *If M is an e.c. factor that contains an infinitely generic factor, then M has the (T)-FRC property.*

We remark that it is still possible that all e.c. factors may be infinitely generic, whence the previous corollary would degenerate to the main result of [18].

Conflicts of interest: On behalf of all authors, the corresponding author states that there is no conflict of interest.

Data Availability: Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

REFERENCES

- [1] Scott Atkinson, Isaac Goldbring, and Srivatsav Kunnawalkam Elayavalli. “Factorial Relative Commutants and the Generalized Jung Property for II_1 Factors”. In: *Advances in Mathematics* 396 (2022), p. 108107.
- [2] Scott Atkinson and Srivatsav Kunnawalkam Elayavalli. “On ultraproduct embeddings and amenability for tracial von Neumann algebras”. In: *Int. Math. Res. Not. IMRN* 4 (2021), pp. 2882–2918. ISSN: 1073-7928. DOI: 10.1093/imrn/rnaa257.
- [3] Rémi Boutonnet, Ionuț Chifan, and Adrian Ioana. “ II_1 factors with nonisomorphic ultrapowers”. In: *Duke Math. J.* 166.11 (2017), pp. 2023–2051. ISSN: 0012-7094. DOI: 10.1215/00127094-0000017X.
- [4] Nathaniel P Brown. “Topological dynamical systems associated to II_1 -factors”. In: *Advances in Mathematics* 227.4 (2011), pp. 1665–1699.
- [5] C.C. Chang and H.J. Keisler. *Model Theory: Third Edition*. Dover Books on Mathematics. Dover Publications, 2013. ISBN: 9780486310954.
- [6] Ionuț Chifan, Daniel Drimbe, and Adrian Ioana. “Embedding universality for II_1 factors with property (T)”. In: *arXiv preprint arXiv:2205.07442* (2022).
- [7] Ionuț Chifan, Adrian Ioana, and Srivatsav Kunnawalkam Elayavalli. “An exotic II_1 factor without property Gamma”. In: *Geometric and Functional Analysis* (2023). DOI: 10.1007/s00039-023-00649-4.
- [8] Alain Connes. “Classification of injective factors. Cases II_1 , II_∞ , III_λ , $\lambda \neq 1$ ”. In: *Ann. of Math. (2)* 104.1 (1976), pp. 73–115. ISSN: 0003-486X. DOI: 10.2307/1971057.
- [9] Alain Connes and Vaughan Jones. “Property T for von Neumann algebras”. In: *Bulletin of the London Mathematical Society* 17.1 (1985), pp. 57–62.
- [10] Jacques Dixmier and E Christopher Lance. “Deux nouveaux facteurs de type II_1 ”. In: *Inventiones mathematicae* 7.3 (1969), pp. 226–234.
- [11] Junsheng Fang, Liming Ge, and Weihua Li. “Central sequence algebras of von Neumann algebras”. In: *Taiwanese J. Math.* 10.1 (2006), pp. 187–200. ISSN: 1027-5487. DOI: 10.11650/twjmath/1500403810.
- [12] Ilias Farah, Bradd Hart, and David Sherman. “Model theory of operator algebras I: stability”. In: *Bulletin of the London Mathematical Society* 45.4 (2013), pp. 825–838.
- [13] Ilias Farah, Bradd Hart, and David Sherman. “Model theory of operator algebras II: model theory”. In: *Israel Journal of Mathematics* 201.1 (2014), pp. 477–505.
- [14] Ilias Farah, Bradd Hart, and David Sherman. “Model theory of operator algebras III: elementary equivalence and II_1 factors”. In: *Bulletin of the London Mathematical Society* 46.3 (2014), pp. 609–628.
- [15] Ilias Farah et al. “Existentially closed II_1 factors”. In: *Fundamenta Mathematicae* 233 (2016), pp. 173–196.
- [16] Isaac Goldbring. “Enforceable operator algebras”. In: *Journal of the Institute of Mathematics of Jussieu* 20 (2021), pp. 31–63.
- [17] Isaac Goldbring. “Non-embeddable II_1 factors resembling the hyperfinite II_1 factor”. In: *Journal of Noncommutative Geometry* 17.1 (2023), pp. 233–239.
- [18] Isaac Goldbring. “On Popa’s factorial commutant embedding problem”. In: *Proceedings of the American Mathematical Society* 148.11 (2020), pp. 5007–5012.
- [19] Isaac Goldbring. “Spectral gap and definability”. In: *arXiv preprint arXiv:1805.02752* (2018).

- [20] Isaac Goldbring and Bradd Hart. *A survey on the model theory of tracial von Neumann algebras*. 2022. DOI: 10.48550/ARXIV.2210.15508.
- [21] Isaac Goldbring and Bradd Hart. “On the Theories of McDuff’s II_1 Factors”. In: *International Mathematics Research Notices* 2017.18 (2017), pp. 5609–5628.
- [22] Isaac Goldbring and Bradd Hart. “Properties expressible in small fragments of the theory of the hyperfinite II_1 factor”. In: *Confluentes Mathematici* 12.2 (2020), pp. 37–47.
- [23] Isaac Goldbring and Bradd Hart. “The Universal Theory Of The Hyperfinite II_1 Factor Is Not Computable”. In: *arXiv preprint arXiv:2006.05629* (2020).
- [24] Isaac Goldbring, Bradd Hart, and Thomas Sinclair. “The theory of tracial von Neumann algebras does not have a model companion”. In: *Journal of Symbolic Logic* 78.3 (2013), pp. 1000–1004.
- [25] M. B. Hastings. “Random unitaries give quantum expanders”. In: *Phys. Rev. A* 76 (3 Sept. 2007), p. 032315. DOI: 10.1103/PhysRevA.76.032315.
- [26] Wilfrid Hodges. *Building models by games*. Courier Corporation, 2006.
- [27] Adrian Ioana and Pieter Spaas. “ II_1 factors with exotic central sequence algebras”. In: *J. Inst. Math. Jussieu* 20.5 (2021), pp. 1671–1696. ISSN: 1474-7480. DOI: 10.1017/S1474748019000653.
- [28] Zhengfeng Ji et al. “MIP* = RE”. In: *Communications of the ACM* 64.11 (2021), pp. 131–138.
- [29] Kenley Jung. “Amenability, tubularity, and embeddings into $\mathcal{R}^\omega$ ”. In: *Math. Ann.* 338.1 (2007), pp. 241–248. ISSN: 0025-5831. DOI: 10.1007/s00208-006-0074-y.
- [30] Amine Marrakchi. “Stability of products of equivalence relations”. In: *Compos. Math.* 154.9 (2018), pp. 2005–2019. ISSN: 0010-437X. DOI: 10.1112/s0010437x18007388.
- [31] Dusa McDuff. “Central sequences and the hyperfinite factor”. In: *Proc. London Math. Soc. (3)* 21 (1970), pp. 443–461. ISSN: 0024-6115. DOI: 10.1112/plms/s3-21.3.443.
- [32] Dusa McDuff. “Uncountably many II_1 factors”. In: *Ann. of Math. (2)* 90 (1969), pp. 372–377. ISSN: 0003-486X. DOI: 10.2307/1970730.
- [33] F. J. Murray and J. von Neumann. “On rings of operators. IV”. In: *Ann. of Math. (2)* 44 (1943), pp. 716–808. ISSN: 0003-486X. DOI: 10.2307/1969107.
- [34] S. Popa. “On the classification of inductive limits of II_1 factors with spectral gap”. In: *Trans. Amer. Math. Soc.* 364.6 (2012), pp. 2987–3000.
- [35] Hui Tan. “Spectral gap characterizations of property (T) for II_1 factors”. preprint, arXiv:2202.06089. 2022.
- [36] G. Zeller-Meier. “Deux nouveaux facteurs de type II_1 ”. In: *Invent. Math.* 7 (1969), pp. 235–242. ISSN: 0020-9910. DOI: 10.1007/BF01404308.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, 340 ROWLAND HALL (BLDG.# 400),
IRVINE, CA 92697-3875

Email address: isaac@math.uci.edu

URL: <http://www.math.uci.edu/~isaac>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, 410 ROWLAND HALL (BLDG.# 400),
IRVINE, CA 92697-3875

Email address: jsapi@math.uci.edu

URL: <https://sites.uci.edu/jpi314/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SAN DIEGO, 9500 GILMAN DR, LA JOLLA, CA 92093

Email address: `djekel@ucsd.edu`

URL: `https://davidjekel.com/`

INSTITUTE OF PURE AND APPLIED MATHEMATICS, UCLA, 460 PORTOLA PLAZA, LOS ANGELES, CA 90095, USA

Email address: `srivatsav.kunnawalkam.elayavalli@vanderbilt.edu`

URL: `https://sites.google.com/view/srivatsavke/home`