



Cross-Coupling Hot Paper

How to cite:

International Edition: doi.org/10.1002/anie.202109146

German Edition: doi.org/10.1002/ange.202109146

Base-Activated Latent Heteroaromatic Sulfinates as Nucleophilic Coupling Partners in Palladium-Catalyzed Cross-Coupling Reactions

Xinlan A. F. Cook, Loïc R. E. Pantaine, David C. Blakemore, Ian B. Moses, Neal W. Sach, Andre Shavnya, and Michael C. Willis*

Abstract: Heteroaromatic sulfinates are effective nucleophilic reagents in Pd⁰-catalyzed cross-coupling reactions with aryl halides. However, metal sulfinate salts can be challenging to purify, solubilize in reaction media, and are not tolerant to multi-step transformations. Here we introduce base-activated, latent sulfinate reagents: β -nitrile and β -ester sulfones. We show that under the cross-coupling conditions, these species generate the sulfinate salt *in situ*, which then undergo efficient palladium-catalyzed desulfinate cross-coupling with (hetero)aryl bromides to deliver a broad range of biaryls. These latent sulfinate reagents have proven to be stable through multi-step substrate elaboration, and amenable to scale-up.

Introduction

Cross-coupling processes are ubiquitous in synthetic chemistry as a dependable method to construct biaryl scaffolds, and are established as a “go to” strategy for carbon–carbon bond formation.^[1] The Suzuki–Miyaura cross-coupling (SMC), which utilizes boron-derived species as nucleophilic coupling partners, is the most popular variant for C(sp²)–C(sp²) bond formation.^[2] This is due to the commercial accessibility of boronic acids and boronate reagents, the mild reaction conditions, and the plethora of research conducted into these transformations.^[2a,3] However, traditional cross-coupling tools, including the Suzuki–Miyaura

approach, fall short when it comes to tackling notoriously difficult heteroaryl–heteroaryl coupling reactions.^[4] Heteroaromatic boron reagents, in particular 2-pyridyl variants, are especially challenging to prepare and couple due to their propensity to undergo protodeboronation.^[5] 2-Arylpyridines, and bi(hetero)aryls in general, are prevalent as motifs in therapeutic and agrochemical molecules,^[6] as ligands in metal catalysis,^[7] and as key units in functional materials^[8] (Figure 1). Owing to the importance of these frameworks there has been significant research towards rectifying this problem,^[4,9] key examples of which include the development of slow release boronate reagents,^[10] and contractive coupling methodologies utilizing phosphorane^[9a,b] or sulfurane^[9c] intermediates. Correspondingly, the “2-pyridyl organometallic cross-coupling problem”^[9d] has become a benchmark for challenging biaryl cross-coupling reactions, and a springboard for innovation. However, a general approach is yet to be established, as many of the solutions reported are plagued with limited heterocycle scopes, difficult reagent preparation, or substrate-specific re-optimization.

Our laboratory has previously contributed to solving these challenges, establishing heteroaromatic sulfinate salts as efficient reagents in Pd-catalyzed desulfinate cross-coupling reactions with aryl halides (Figure 2a).^[11] Desulfinate cross-coupling processes avoid the use of stoichiometric organometallic reagents and only release SO₂ as a by-product. Although the use of aryl sulfinate salts in desulfinate cross-couplings was first reported in the early 1990s,^[12] only in recent years has substantial research into their use as nucleophilic coupling partners in Pd-mediated reactions been implemented.^[13,14] Importantly, heteroaromatic sulfinate salts are generally more air- and moisture-stable than the corre-

[*] X. A. F. Cook, Dr. L. R. E. Pantaine, Prof. M. C. Willis
Department of Chemistry
University of Oxford, Chemistry Research Laboratory
Mansfield Road, Oxford, OX1 3TA (UK)
E-mail: michael.willis@chem.ox.ac.uk

Dr. D. C. Blakemore, A. Shavnya
Medicine Design, Pfizer Inc.
Eastern Point Road, Groton, CT 06340 (USA)

I. B. Moses
Pharmaceutical sciences, Pfizer Inc.
Discovery Park, Ramsgate Road, CT13 9ND (UK)

N. W. Sach
Medicine Design, La Jolla Laboratories, Pfizer Inc.
10777 Science Center Drive, San Diego, CA 92121 (USA)

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202109146>.

© 2021 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.




Figure 1. Select examples of 2-pyridyl derivatives with varied applications.




Previous work in the Willis laboratory:

(a) Heteroaromatic metal sulfonates (2017)



(b) Heteroaromatic allylsulfones as latent sulfonates (2018)



Metal sulfonate salts

- Robust cross-coupling
- Insoluble salts
- Difficult to purify and isolate
- Not suitable for multi-step elaboration

Allylsulfones

- Efficient coupling via in situ sulfinate
- Stable & organic soluble
- Suitable for multi-step elaboration
- Sensitive to Pd-mediated chemistry
- Incompatible with alkene transformations

This work:

(c) Heteroaromatic β-nitrile and β-ester sulfones as latent sulfonates




Figure 2. Introduction to previous and current work: a) heteroaromatic sulfonate salts, b) heteroaromatic allylsulfones and c) this work: β-nitrile sulfones and β-ester sulfones in cross-coupling reactions.

sponding boronic acids, and are now becoming commercially available.^[15]

Despite being efficient coupling partners, the purification of metal sulfonate salts, as well as their poor solubility in organic media, pose challenges and hinder their ability to be carried through multi-step synthetic sequences. To address this, our laboratory pioneered the use of latent sulfonates in cross-coupling reactions. This concept exploits neutral, organosoluble, and easily accessible protected sulfonates that can be selectively unmasked under coupling conditions. The first generation of latent sulfonate reagents developed were allylsulfones (Figure 2b).^[16] Owing to the lability of the allyl group under Pd-catalysis, no additives were required for the release of the sulfinate coupling partner. However, this also limited the use of such substrates in other palladium-catalyzed processes, and in alkene functionalization reactions.


This work explores a new generation of latent *N*-heteroaromatic sulfonates: β-nitrile and β-ester sulfones (Figure 2c).^[9f,17,21] The active sulfinate species are released via E₁cB elimination under the basic cross-coupling conditions.^[18] These base-labile reagents, in contrast to the Pd-labile allylsulfones, are more amenable to orthogonal Pd-catalyzed reactions and typical alkene transformations.

Results and Discussion


Heteroaromatic sulfones can be readily accessed from commercially available starting materials, either directly from thiols, halides or via the sulfinate salt.^[19,21] The β-nitrile sulfones were primarily accessed from thiols through conjugate addition into acrylonitrile and subsequent oxidation (the order of these steps could also be reversed, Scheme 1a, b),^[11a,b] from aryl halides using the sodium 3-methoxy-3-oxopropene-1-sulfinate (SMOPS) reagent (Scheme 1c),^[21] or by utilizing SO₂ surrogates, such as DABSO, to form the parent sulfinate,^[19b,e,20] followed by conjugate addition (Scheme 1d).

The use of β-nitrile sulfones in Pd-mediated cross-coupling was initially investigated through the reaction of the pyridine sulfone **1a** with 4-bromoanisole **2a** (Table 1). Encouragingly, the desired biaryl product **3a** was obtained in a 29 % yield under the original conditions reported for the use of 2-pyridyl metal sulfinate salts (Entry 1).^[11b] A strong solvent dependence was quickly observed, as substituting 1,4-dioxane for toluene more than doubled the yield of biaryl **3a** (Entry 2). The presence of K₂CO₃ proved to be essential to reaction success, in agreement with the reported mechanism of the transformation.^[11c,d] A range of ligands were evaluated and electron-rich, bulky monodentate phosphine ligands performed best (see Supporting Information). Whilst the optimum ligand for the allylsulfone work, P(tBu)₂Me·HBF₄, showed good activity (Entry 3), cataCXium A (PAd₂Bu) was shown to be superior for this process giving biaryl **3a** in 84 % yield (Entry 4). Addition of acetic acid to the system was key for achieving high yields of the desired product at 130 °C (Entries 5–6). Other bases and acids were explored, however, none performed as well (see Supporting Information). Considering the poor solubility of sulfinate salts in toluene, the combination of base and acid could potentially create a buffered system, enabling a more gradual release of the sulfinate from the heteroaromatic sulfone. Lastly, by adjusting the stoichiometry of the pyridine sulfone to a slight excess


(a) Conjugate addition/oxidation




(b) Oxidation/conjugate addition




(c) Cu-mediated sulfone formation and sulfinate release/conjugate addition



(d) Metalation/sulfonation/conjugate addition



Scheme 1. Synthesis of β-nitrile sulfones. TIMSO = N-Me-pyrrolidine SO₂ adduct.


Table 1: Selected optimization studies of the desulfurative coupling.^[a]


Entry	Solvent	Ligand	Additive (1.0 equiv)	T [°C]	Yield [%] ^[b]
1	1,4-dioxane	PCy ₃	—	150	29 ^[c,d]
2	toluene	PCy ₃	—	150	67
3	toluene	P(^t Bu) ₂ Me-HBF ₄	—	150	76
4	toluene	cataCXium A	—	150	84
5	toluene	cataCXium A	—	130	59
6	toluene	cataCXium A	AcOH	130	84
7 ^[e]	toluene	cataCXium A	AcOH	130	91 ^[d]
8 ^[e]	toluene	cataCXium A	AcOH	120	88 ^[d]

[a] Reaction conditions: **1a** (0.20 mmol, 1.0 equiv), 4-bromoanisole (0.20 mmol, 1.0 equiv), K₂CO₃ (0.30 mmol, 1.5 equiv), Pd(OAc)₂ (5.0 mol%), ligand (10 mol%), solvent (0.10 M). [b] HPLC yields determined by using *p*-tolylether as an internal standard. [c] Run on a 0.40 mmol scale. [d] Isolated yield. [e] Using 1.1 equiv of **1a**. cataCXium A = Pd₂Bu.

(1.1 equiv), we were able to achieve an 88 % yield of biaryl **3a** at 120 °C (Entry 8). The acrylonitrile by-product was not observed and is assumed to be removed by evaporation.

We began the reaction scope exploration by varying the electrophilic partner in combination with sulfone **1a** (Table 2a). Aryl bromides were found to couple more efficiently than aryl chlorides, as shown by the 61 % yield of compound **3a** obtained from 4-chloroanisole, compared to 88 % using 4-bromoanisole. This difference in reactivity could be advantageously exploited and chlorobiaryl **3b** was obtained in a high 80 % yield. The reaction yield remained high when varying the position of the methyl substituent on the aryl bromide (**3c–3f**), although a slight increase in temperature to 130 °C was needed for some *ortho*-substituted aryl bromides (**3e**, **3f**). Under these reaction conditions, pyridine sulfone **1a** provides an efficient synthesis of the pyridine **3h**, used widely as a ligand in photocatalyst complexes and luminescent platinum complexes.^[7a,b] The reaction has a broad functional group tolerance, delivering high yields when employing aryl halides containing esters, ketones, nitriles, protected amines and

Table 2: Scope of the desulfurative cross-coupling of heteroaromatic β-nitrile sulfones with (hetero)aryl halides.^[a]


(a) Variation of (hetero)aryl halide coupling partners with β-nitrile sulfone 1a

3a, 88%, 61%^[b,c] **3b**, 80% **3c**, 80% **3d**, 95% **3e**, 91%^[b] **3f**, 65%^[b] **3g**, 91%^[b]

3h, 75%^[b] **3i**, 87% **3j**, 83% **3k**, 91% **3l**, 78% **3m**, 75%

3n, 74% **3o**, 76% **3p**, 87%^[b] **3q**, 75% **3r**, 97% **3s**, 80%

(b) Variation of heteroaromatic β-nitrile sulfones with (hetero)aryl halides

3t, 94%^[b] **3u**, 88% **3v**, 84% **3w**, 67%^[b] **3x**, 73%^[b] SEM **3y**, 94%^[b]

3z, 77% **3aa**, 95%^[b] **3ab**, 74%^[b] **3ac**, 83%^[b] **3ad**, 59%^[b] **3ae**, 90% **3af**, 93%^[b]

3ag, 49%^[b] **3ah**, 95% **3ai**, 98%^[b] **3aj**, 63%^[b] **3ak**, 82% **3al**, 64%

[a] Isolated yields. Reaction conditions: pyridine sulfone (0.22 mmol, 1.1 equiv), aryl halide (0.20 mmol, 1.0 equiv), K₂CO₃ (0.30 mmol, 1.5 equiv), AcOH (0.20 mmol, 1.0 equiv), Pd(OAc)₂ (5.0 mol%), cataCXium A (10 mol%), toluene (0.10 M), 120 °C, 18 h. [b] Reaction run at 130 °C. [c] Using 4-chloroanisole. Temperatures stated are those of the aluminium heating block.

alcohols, thiophenes, substituted pyridines, quinolines, quinoxalines and naphthalenes (**3i–3s**).


On variation of the nucleophilic coupling partner (Table 2b), we found that the 6- and 5-methyl substituted pyridine sulfones behave similarly to the electron-deficient 5-trifluoromethyl pyridine sulfone **1a** (**3t**, **3u**, **3z**). Pleasingly, the more challenging couplings to access biaryls, such as pyrimidines (**3v–3w**), SEM-protected pyrazole (**3x**) and imidazo[1,2-*a*]pyrazine (**3y**), were achieved in good to high yields. Increasing the steric hindrance at the reaction site via substitution at the 3-position of the pyridine, was also well tolerated, giving 95 % of 2-arylpyridine **3aa**. Increasing the reaction temperature to 130 °C was often necessary to achieve good yields when using unsubstituted pyridine sulfones (**3ab–3ag**). This is in good agreement with our previous mechanistic studies, which show that extrusion of SO₂ during the catalytic cycle is easier with electron-deficient or sterically bulky substrates.^[11c] Electron-rich pyridine sulfones also performed well under these reaction conditions (**3ah**, **3ai**). Notably, indole **3aj** and pyrazine **3ak** were synthesized in good to high yields. This methodology can also be extended to carbocyclic substrates, as shown with compound **3al**, obtained in 64 % from *p*-tolyl β-nitrile sulfone.

To demonstrate the generality and robustness of base-labile latent sulfonates, we extended our approach to explore the use of β-ester sulfones. In comparison to β-nitrile sulfones, the synthetic routes towards β-ester substrates are more straightforward, mainly due to β-methylester sulfones being established intermediates in the synthesis of metal sulfina-

tes.^[9f,19a,21] The aforementioned SMOPS reagent is commercially available and provides a one-step route to the β-methylester sulfone functionality.^[21] Additionally, commercially available methyl 3-mercaptopropionate can also be reacted directly with heteroaromatic halides, or facilitated by metal catalysis if required,^[22] which in turn can be followed by S-oxidation to give β-methylester sulfones. Thus, synthetic routes towards β-ester sulfones allow better use of the vast libraries of readily accessible heteroaryl halides.

In contrast to the nitrile derivatives, β-ester sulfones generally performed better when the AcOH additive was omitted from the system. Under such conditions, the reactivity and functional group tolerance of these alternative latent sulfonates were found to be comparable to the β-nitrile sulfones, giving the desired biaryls in high yields (Table 3, **3a**, **3b**, **3i**, **3k**, **3q**). The free alcohol product **3l-OH** was pleasingly obtained in a 65 % yield from the telescoped deprotection of protected alcohol **3l**. Similar to the β-nitrile sulfone system, raising the reaction temperature to 130 °C was required for efficient coupling of the unsubstituted pyridyl β-ester sulfone to give the biaryl **3ac**, was poor yielding under the optimized conditions, despite complete consumption of the sulfone. However, addition of acetic acid (1.5 equiv) into the reaction allowed for the dramatic increase in the yield of product **3ac** from 29 % to 91 %. The 4-pyridyl β-ester sulfone worked comparably to the 2-pyridyl reagent, giving 4-arylpyridines **3ag** and **3ao**.

Table 3: Scope of the desulfurative cross-coupling of heteroaromatic β-ester sulfones with (hetero)aryl halides.^[a]



[a] Isolated yields. Reaction conditions: sulfone (0.3 mmol, 1.5 equiv), aryl bromide (0.2 mmol, 1.0 equiv), $\text{Pd}(\text{OAc})_2$ (5.0 mol %), cataCXium A (10 mol %), K_2CO_3 (0.4 mmol, 2.0 equiv), toluene (0.10 M), 120 °C, 18 h. [b] Run at 130 °C. [c] Telescoped deprotection of protected phenol product directly after cross-coupling reaction: reaction temperature lowered to 85 °C, ethanol (2 mL), 2 M aq. solution of K_2CO_3 (0.2 mL, 0.4 mmol), 18 h. [d] Reaction run with AcOH (0.3 mmol, 1.5 equiv). Temperatures stated are those of the aluminium heating block.

Nucleophilic coupling partners with electron-withdrawing substituents on the pyridine core, such as a nitrile (**3ap**) or an ester (**3aq**), as well as quinoxaline sulfones (**3at**), performed well, as did electron-rich β -ester examples (**3ah**, **3ai**). Similar to the β -nitrile sulfone system, substituents next to the sulfinate coupling site were well tolerated (**3ar–3as**). More complex, nitrogen-rich frameworks **3au** and **3av**, were constructed from the corresponding 2-quinoline and 2-quinoxaline sulfones. The pyrimidine containing molecule **3aw**, an intermediate of a class of ROR γ modulators,^[23] was synthesized in a 99 % yield. Pleasingly, the challenging coupling of benzothiazole β -methyl ester sulfone to give benzothiazole **3ax**, a product related to PET scanning radiopharmaceutical Flutemetamol (¹⁸F), was achieved.^[24] Lastly, we were able to prepare complex biaryl **3ay**, containing a fragment of arthritis drug celecoxib, in 83 % yield.^[25]

The success of the latent sulfonates (Table 2 and Table 3) inspired the design and brief exploration of diethyl succinate sulfones **5**. Upon generation of the sulfinate salt, this alternative masking group releases non-toxic, high-boiling diethyl fumarate as the by-product, making this system more amenable to cross-coupling on larger scale. Additionally, the majority of these reagents can be readily isolated as solids. Encouragingly, the succinate pyridine sulfone **5** performed well under the same conditions employed for the β -methyl ester sulfone (Table 4). These reagents react efficiently and comparably to their β -nitrile and β -ester counterparts, as exemplified through a representative set of biaryls (Table 4).


The described reactions of our masked sulfonates were routinely performed on 0.2 mmol scale in sealed reaction vials. However, we were keen to show that the chemistry could be translated to larger, non-pressurized reaction vessels. To investigate this, the cross-coupling of sulfones **1a** or **4a** with 4-bromoanisole **2a** to give biaryl **3a** was re-visited. The key substitution of toluene for higher boiling anisole, and a reaction temperature of 130 °C allowed the reactions of both masked sulfonates species to be run effectively on a 1.0 mmol scale (Scheme 2). Pleasingly, an unoptimized 52 % yield of 2-

(*p*-tolyl)pyridine (**3ac**) was also obtained using the 2-pyridyl succinate sulfone **5a**.


To highlight the compatibility of β -nitrile and β -methyl ester sulfone with multi-step substrate functionalization, we investigated the stability and utility of these masking groups through a series of transformations around the pyridine core, primarily focusing on oxidative and palladium-mediated reactions. Firstly, using O-allyl pyridine sulfone **6**, we explored oxidative transformations at the alkene functionality (Scheme 3). Pyridine **6** was oxidized under Wacker-type conditions^[26] to give the desired ketone product **7** in a 44 % yield, and the resultant ketone was further reacted to access biaryl **3bb**. Furthermore, through a hydroboration-oxidation sequence we could achieve a 65 % yield of the combined alcohol products (**8a,b**); this is a transformation that would not be possible using the corresponding allylsulfone. The primary alcohol **8a** was then successfully coupled to give 60 % of the desired 2-arylpyridine **3bc**.

Lastly, we explored the bidirectional elaboration of masked pyridine reagents by installing two cross-coupling reaction sites that could be selectively transformed (Scheme 4). 5-Bromopyridine sulfones (**9a,b**) successfully underwent Suzuki–Miyaura cross-coupling (SMC) with phenyl boronic acid to give the biaryl sulfones (**10a,b**) in 72 % and 78 % yields, when using β -nitrile or β -ester masking groups, respectively.^[27] In both cases the latent sulfinate group was stable under these reaction conditions. Sequentially these products were submitted to desulfonative coupling conditions to give the desired diarylpyridine compound **11** in high yields. To further demonstrate the compatibility of this chemistry with transition metal-mediated processes, the boronate ester


Table 4: Scope of the desulfonative cross-coupling of succinate sulfones with aryl bromides.^[a]



[a] Isolated yields. Reaction conditions: pyridine succinate sulfone (0.3 mmol, 1.5 equiv), aryl bromide (0.2 mmol, 1.0 equiv), Pd(OAc)₂ (5.0 mol %), cataCXium A (10 mol %), K₂CO₃ (0.4 mmol, 2.0 equiv), toluene (0.1 M), 130 °C, 18 h. [b] Reaction run at 120 °C.



Scheme 2. Scale up of the cross-coupling reactions of base-labile latent sulfinate reagents. Reaction conditions: a) pyridine sulfone (1.5 mmol, 1.5 equiv), 4-bromoanisole (1.0 mmol, 1.0 equiv), Pd(OAc)₂ (5.0 mol %), cataCXium A (10 mol %), K₂CO₃ (2.0 mmol, 2.0 equiv), AcOH (1.5 mmol, 1.5 equiv), anisole (0.1 M), 130 °C, 18 h. b and c) pyridine sulfone (1.5 mmol, 1.5 equiv), 4-bromoanisole (1.0 mmol, 1.0 equiv), Pd(OAc)₂ (5.0 mol %), cataCXium A (10 mol %), K₂CO₃ (2.0 mmol, 2.0 equiv), anisole (0.1 M), 130 °C, 18 h. Isolated yields; internal temperatures measured.




Scheme 3. Oxidative transformations of O-allyl pyridine sulfone **6**. DMP = Dess–Martin periodinane.

containing substrate **12** was prepared using an Ir-catalyzed borylation reaction (Scheme 4b).^[28] The bifunctional pyridine **12** underwent SMC with 4-bromotoluene **2b** to give sulfone **13** in 62 % yield;^[29] notably, the masked sulfinate functionality remained intact. Finally, diarylpyridine **14** was obtained in excellent 92 % yield from biaryl sulfone **13** using our standard cross-coupling protocol.

Conclusion

We have designed and developed β -nitrile and β -ester sulfones as efficient base-labile latent sulfinate reagents in palladium-catalyzed cross-coupling reactions. This method has been used to construct challenging heteroaryl-(hetero)aryl linkages, allowing access to a diverse range of 2-arylpyridines and pharmaceutically relevant fragments. The scope of electrophilic partners is broad, displaying a good tolerance of multiple functional groups, and substitution patterns, delivering the desired cross-coupled products in good to high yields. Furthermore, both latent sulfinate groups are stable and can be carried through multistep elaboration, notably oxidative and palladium-mediated transformations. We have further shown the successful use of the succinate sulfone masking group, which releases a benign by-product on activation. Owing to the prevalence of the pyridine motif in medicinal chemistry and the robustness of this chemistry, we anticipate this method will find wide application.



Scheme 4. Transition-metal mediated, multi-step cross-coupling sequences of 2-pyridyl latent sulfinate reagents. tfp = tri(2-furyl)phosphine. dtbpy = 4,4'-di-*tert*-butyl-2,2'-dipyridyl. XPhos = 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl.

Acknowledgements

X.A.F.C. is grateful to the EPSRC Centre for Doctoral Training in Synthesis for Biology and Medicine (EP/L015838/1) for a studentship, generously supported by Pfizer, GlaxoSmithKline, Vertex, AstraZeneca, Diamond Light Source, Defence Science and Technology Laboratory, Evotec, Janssen, Novartis, Syngenta, Takeda and UCB. X.A.F.C. also thanks the Clarendon Fund for a scholarship and support. L.R.E.P. acknowledges financial support from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement DeCoChem No. 793155.

Conflict of Interest

The authors declare no conflict of interest.

Keywords: cross-coupling · desulfonation · palladium · pyridine · sulfone



- [1] a) L.-C. Campeau, N. Hazari, *Organometallics* **2019**, *38*, 3–35; b) M. J. Buskes, M.-J. Blanco, *Molecules* **2020**, *25*, 3493–3515.
- [2] a) D. Blakemore, *Synthetic Methods in Drug Discovery, Vol. 1*, The Royal Society of Chemistry, London, **2016**, pp. 1–69; b) N. Miyaura, A. Suzuki, *Chem. Rev.* **1995**, *95*, 2457–2483; c) S. D. Roughley, A. M. Jordan, *J. Med. Chem.* **2011**, *54*, 3451–3479; d) N. Miyaura, K. Yamada, A. Suzuki, *Tetrahedron Lett.* **1979**, *20*, 3437–3440.
- [3] a) A. J. J. Lennox, G. C. Lloyd-Jones, *Chem. Soc. Rev.* **2014**, *43*, 412–443; b) I. P. Beletskaya, F. Alonso, V. Tyurin, *Coord. Chem. Rev.* **2019**, *385*, 137–173.
- [4] X. A. F. Cook, A. de Gombert, J. McKnight, L. R. E. Pantaine, M. C. Willis, *Angew. Chem. Int. Ed.* **2021**, *60*, 11068–11091; *Angew. Chem.* **2021**, *133*, 11168–11191.
- [5] a) P. A. Cox, A. G. Leach, A. D. Campbell, G. C. Lloyd-Jones, *J. Am. Chem. Soc.* **2016**, *138*, 9145–9157; b) P. A. Cox, M. Reid, A. G. Leach, A. D. Campbell, E. J. King, G. C. Lloyd-Jones, *J. Am. Chem. Soc.* **2017**, *139*, 13156–13165; c) H. G. Kuivila, K. V. Nahabedian, *J. Am. Chem. Soc.* **1961**, *83*, 2159–2163.
- [6] Selected examples: a) S. L. Hargreaves, B. L. Pilkington, S. E. Russell, P. A. Worthington, *Tetrahedron Lett.* **2000**, *41*, 1653–1656; b) P. Fu, S. Wang, K. Hong, X. Li, P. Liu, Y. Wang, W. Zhu, *J. Nat. Prod.* **2011**, *74*, 1751–1756; c) S. P. Curtis, B. Bockow, C. Fisher, J. Olaleye, A. Compton, A. T. Ko, A. S. Reicin, *BMC Musculoskeletal Disord.* **2005**, *6*, 58; d) E. Vitaku, D. T. Smith, J. T. Njardarson, *J. Med. Chem.* **2014**, *57*, 10257–10274; e) J. J. Cui, M. Tran-Dubé, H. Shen, M. Nambu, P.-P. Kung, M. Pairish, L. Jia, J. Meng, L. Funk, I. Botrous, M. McTigue, N. Grodsky, K. Ryan, E. Padrique, G. Alton, S. Timofeevski, S. Yamazaki, Q. Li, H. Zou, J. Christensen, B. Mroczkowski, S. Bender, R. S. Kania, M. P. Edwards, *J. Med. Chem.* **2011**, *54*, 6342–6363; f) S. Y. Jung, S. Hwang, J. M. Clarke, T. M. Bauer, V. L. Keedy, H. Lee, N. Park, S.-J. Kim, J. I. Lee, *Invest. New Drugs* **2019**, *38*, 812–820.
- [7] Selected examples: a) K. Teegardin, J. I. Day, J. Chan, J. Weaver, *Org. Process Res. Dev.* **2016**, *20*, 1156–1163; b) L. Gao, J. Ni, M. Su, J. Kang, J. Zhang, *Dyes Pigm.* **2019**, *165*, 231–238; c) F. Havas, N. Leygue, M. Danel, B. Mestre, C. Galaup, C. Picard, *Tetrahedron* **2009**, *65*, 7673–7686.
- [8] Selected examples: a) C. L. Fraser, A. P. Smith, *J. Polym. Sci. Part A* **2000**, *38*, 4704–4716; b) V. Bekiari, P. Lianos, P. Judeinstein, *Chem. Phys. Lett.* **1999**, *307*, 310–316; c) V. Bekiari, G. Pistolis, P. Lianos, *Chem. Mater.* **1999**, *11*, 3189–3195; d) W. Sun, H. Zhu, P. M. Barron, *Chem. Mater.* **2006**, *18*, 2602–2610; e) D. Saccone, C. Magistris, N. Barbero, P. Quagliotto, C. Barolo, G. Viscardi, *Materials* **2016**, *9*, 137; f) E. Kreidt, C. Bischof, C. Platas-Iglesias, M. Seitz, *Inorg. Chem.* **2016**, *55*, 5549–5557; g) H. Zhang, J. Lee, A. D. Lammer, X. Chi, J. T. Brewster, V. M. Lynch, H. Li, Z. Zhang, J. L. Sessler, *J. Am. Chem. Soc.* **2016**, *138*, 4573–4579.
- [9] a) B. T. Boyle, M. C. Hilton, A. McNally, *J. Am. Chem. Soc.* **2019**, *141*, 15441–15449; b) M. C. Hilton, X. Zhang, B. T. Boyle, J. V. Alegre-Requena, R. S. Paton, A. McNally, *Science* **2018**, *362*, 799–804; c) M. Zhou, J. Tsien, T. Qin, *Angew. Chem. Int. Ed.* **2020**, *59*, 7372–7376; *Angew. Chem.* **2020**, *132*, 7442–7446; d) L.-C. Campeau, S. Rousseaux, K. Fagnou, *J. Am. Chem. Soc.* **2005**, *127*, 18020–18021; e) C. Fricke, G. J. Sherborne, I. Funes-Ardoiz, E. Senol, S. Guven, F. Schoenebeck, *Angew. Chem. Int. Ed.* **2019**, *58*, 17788–17795; *Angew. Chem.* **2019**, *131*, 17952–17959; f) J. Wei, H. M. Liang, C. F. Ni, R. Sheng, J. B. Hu, *Org. Lett.* **2019**, *21*, 937–940; g) T. Nokami, Y. Tomida, T. Kamei, K. Itami, J.-i. Yoshida, *Org. Lett.* **2006**, *8*, 729–731; h) L. Ackermann, H. K. Potukuchi, A. R. Kapdi, C. Schulzke, *Chem. Eur. J.* **2010**, *16*, 3300–3303; i) L. Chen, D. R. Sanchez, B. Zhang, B. P. Carrow, *J. Am. Chem. Soc.* **2017**, *139*, 12418–12421; j) M. R. Luzung, J. S. Patel, J. Yin, *J. Org. Chem.* **2010**, *75*, 8330–8332; k) X. Li, D. Zou, F. Leng, C. Sun, J. Li, Y. Wu, Y. Wu, *Chem. Commun.* **2013**, *49*, 312–314; l) D. C. Blakemore, L. A. Marples, *Tetrahedron Lett.* **2011**, *52*, 4192–4195.
- [10] a) K. L. Billingsley, S. L. Buchwald, *Angew. Chem. Int. Ed.* **2008**, *47*, 4695–4698; *Angew. Chem.* **2008**, *120*, 4773–4776; b) J. Z. Deng, D. V. Paone, A. T. Ginnetti, H. Kurihara, S. D. Dreher, S. A. Weissman, S. R. Stauffer, C. S. Burgey, *Org. Lett.* **2009**, *11*, 345–347; c) G. Q. Li, Y. Yamamoto, N. Miyaura, *Synlett* **2011**, 1769–1773; d) S. Sakashita, M. Takizawa, J. Sugai, H. Ito, Y. Yamamoto, *Org. Lett.* **2013**, *15*, 4308–4311; e) M. A. Oberli, S. L. Buchwald, *Org. Lett.* **2012**, *14*, 4606–4609; f) W. Ren, J. Li, D. Zou, Y. Wu, Y. Wu, *Tetrahedron* **2012**, *68*, 1351–1358; g) P. B. Hodgson, F. H. Salingue, *Tetrahedron Lett.* **2004**, *45*, 685–687; h) P. Gros, A. Doudouh, Y. Fort, *Tetrahedron Lett.* **2004**, *45*, 6239–6241; i) K. Chen, R. Peterson, S. K. Math, J. B. LaMunyon, C. A. Testa, D. R. Cefalo, *Tetrahedron Lett.* **2012**, *53*, 4873–4876; j) Y. Mutoh, K. Yamamoto, S. Saito, *ACS Catal.* **2020**, *10*, 352–357; k) H. Yoshida, M. Seki, S. Kamio, H. Tanaka, Y. Izumi, J. Li, I. Osaka, M. Abe, H. Andoh, T. Yajima, T. Tani, T. Tsuchimoto, *ACS Catal.* **2020**, *10*, 346–351; l) S. Kamio, I. Kageyuki, I. Osaka, H. Yoshida, *Chem. Commun.* **2019**, *55*, 2624–2627; m) G. R. Dick, E. M. Woerly, M. D. Burke, *Angew. Chem. Int. Ed.* **2012**, *51*, 2667–2672; *Angew. Chem.* **2012**, *124*, 2721–2726; n) Y. Yamamoto, M. Takizawa, X. Q. Yu, N. Miyaura, *Angew. Chem. Int. Ed.* **2008**, *47*, 928–931; *Angew. Chem.* **2008**, *120*, 942–945.
- [11] a) T. Markovic, B. N. Rocke, D. C. Blakemore, V. Mascitti, M. C. Willis, *Org. Lett.* **2017**, *19*, 6033–6035; b) T. Markovic, B. N. Rocke, D. C. Blakemore, V. Mascitti, M. C. Willis, *Chem. Sci.* **2017**, *8*, 4437–4442; c) A. de Gombert, A. I. McKay, C. J. Davis, K. M. Wheelhouse, M. C. Willis, *J. Am. Chem. Soc.* **2020**, *142*, 3564–3576; d) A. de Gombert, M. C. Willis, *Trends Chem.* **2020**, *2*, 865–866.
- [12] D. H. Ortgies, A. Hassanpour, F. Chen, S. Woo, P. Forgione, *Eur. J. Org. Chem.* **2016**, 408–425.
- [13] a) C. Zhou, Q. Liu, Y. Li, R. Zhang, X. Fu, C. Duan, *J. Org. Chem.* **2012**, *77*, 10468–10472; b) J. Liu, X. Zhou, H. Rao, F. Xiao, C.-J. Li, G.-J. Deng, *Chem. Eur. J.* **2011**, *17*, 7996–7999; c) D. H. Ortgies, A. Barthelme, S. Aly, B. Desharnais, S. Rioux, P. Forgione, *Synthesis* **2013**, *45*, 694–702; d) D. H. Ortgies, P. Forgione, *Synlett* **2013**, *24*, 1715–1721; e) C. Zhou, Y. Li, Y. Lu, R. Zhang, K. Jin, X. Fu, C. Duan, *Chin. J. Chem.* **2013**, *31*, 1269–1273.
- [14] a) W. Chen, P. Li, T. Miao, L.-G. Meng, L. Wang, *Org. Biomol. Chem.* **2013**, *11*, 420–424; b) S. Sévigny, P. Forgione, *New J. Chem.* **2013**, *37*, 589–592; c) S. Sévigny, P. Forgione, *Chem. Eur. J.* **2013**, *19*, 2256–2260; d) J. Shi, X.-D. Tang, Y.-C. Wu, H.-N. Li, L.-J. Song, Z.-Y. Wang, *Eur. J. Org. Chem.* **2015**, *1193*–1197; e) D. Mangel, C. Buonomano, S. Sevigny, G. Di Censo, G. Thevendran, P. Forgione, *Heterocycles* **2015**, *90*, 1228–1239; f) J. Colomb, T. Billard, *Tetrahedron Lett.* **2013**, *54*, 1471–1474.
- [15] D. Kaiser, I. Klose, R. Oost, J. Neuhaus, N. Maulide, *Chem. Rev.* **2019**, *119*, 8701–8780.
- [16] T. Markovic, P. R. D. Murray, B. N. Rocke, A. Shavnya, D. C. Blakemore, M. C. Willis, *J. Am. Chem. Soc.* **2018**, *140*, 15916–15923.
- [17] a) A. G. M. Barrett, S. M. Cramp, A. J. Hennessy, P. A. Procopiou, R. S. Roberts, *Org. Lett.* **2001**, *3*, 271–273; b) M. Gilligan, A. C. Humphries, T. Ladduwahetty, K. J. Merchant (Merch, Sharpe & Dohme), WO 095205, **2006**.
- [18] Related nitriles and esters have been used as protecting groups for thiols and alcohols, see for example: a) J.-M. Becht, A. Wagner, C. Mioskowski, *J. Org. Chem.* **2003**, *68*, 5758–5761; b) H. Saneyoshi, K. Ando, K. Seio, M. Sekine, *Tetrahedron* **2007**, *63*, 11195–11203; c) M. Janczyk, B. Appel, D. Springstube, H. J. Fritz, S. Muller, *Org. Biomol. Chem.* **2012**, *10*, 1510–1513.
- [19] a) S. Liang, K. Hofman, M. Friedrich, G. Manolikakes, *Eur. J. Org. Chem.* **2020**, 4664–4674; b) E. J. Emmett, B. R. Hayter,

- M. C. Willis, *Angew. Chem. Int. Ed.* **2013**, *52*, 12679–12683; *Angew. Chem.* **2013**, *125*, 12911–12915; c) A. Shavnya, K. D. Hesp, A. S. Tsai, *Adv. Synth. Catal.* **2018**, *360*, 1768–1774; d) K. M. Maloney, J. T. Kuethe, K. Linn, *Org. Lett.* **2011**, *13*, 102–105; e) A. S. Deeming, C. J. Russell, A. J. Hennessy, M. C. Willis, *Org. Lett.* **2014**, *16*, 150–153.
- [20] a) H. Woolven, C. González-Rodríguez, I. Marco, A. L. Thompson, M. C. Willis, *Org. Lett.* **2011**, *13*, 4876–4878; b) B. Skillinghaug, J. Rydfjord, L. R. Odell, *Tetrahedron Lett.* **2016**, *57*, 533–536.
- [21] J. M. Baskin, Z. Y. Wang, *Tetrahedron Lett.* **2002**, *43*, 8479–8483.
- [22] a) X. Zheng, P. Bauer, T. Baumeister, A. J. Buckmelter, M. Caligiuri, K. H. Clodfelter, B. Han, Y.-C. Ho, N. Kley, J. Lin, D. J. Reynolds, G. Sharma, C. C. Smith, Z. Wang, P. S. Dragovich, J. Gunzner-Toste, B. M. Liederer, J. Ly, T. O'Brien, A. Oh, L. Wang, W. Wang, Y. Xiao, M. Zak, G. Zhao, P.-W. Yuen, K. W. Bair, *J. Med. Chem.* **2013**, *56*, 6413–6433; b) W. Liu, W. Deng, S. Sun, C. Yu, X. Su, A. Wu, Y. Yuan, Z. Ma, K. Li, H. Yang, X. Peng, J. Dietrich, *Org. Lett.* **2019**, *21*, 9909–9913.
- [23] S. Das, L. A. Gharat, R. L. Harde, S. Y. Shelke, S. R. Pardeshi, A. Thomas, N. Khair-Atkar-Joshi, D. M. Shah, M. Bajpai (Glenmark Pharmaceuticals, SA), WO 021879 A1, **2017**.
- [24] a) K. Heurling, A. Leuzy, E. R. Zimmer, M. Lubberink, A. Nordberg, *Eur. J. Nucl. Med. Mol. Imaging* **2016**, *43*, 362–373; b) T. G. Beach, D. R. Thal, M. Zanette, A. Smith, C. Buckley, *J. Alzheimer's Dis.* **2016**, *52*, 863–873.
- [25] R. C. Hubbard, R. J. Koepp, S. Yu, S. Talwalker, G. S. Geis, C. W. Wiesenhutter, W. S. Makarowski, H. A. Paulus, *Arthritis Rheum.* **1996**, *39*, 1188–1188.
- [26] D. A. Chaudhari, R. A. Fernandes, *J. Org. Chem.* **2016**, *81*, 2113–2121.
- [27] D. Hédou, A. S. Voisin-Chiret, *Eur. J. Org. Chem.* **2020**, 3640–3649.
- [28] a) E. Demory, K. Devaraj, A. Orthaber, P. J. Gates, L. T. Pilarski, *Angew. Chem. Int. Ed.* **2015**, *54*, 11765–11769; *Angew. Chem.* **2015**, *127*, 11931–11935; b) S. A. Sadler, H. Tajuddin, I. A. I. Mkhaliid, A. S. Batsanov, D. Albesa-Jove, M. S. Cheung, A. C. Maxwell, L. Shukla, B. Roberts, D. C. Blakemore, Z. Lin, T. B. Marder, P. G. Steel, *Org. Biomol. Chem.* **2014**, *12*, 7318–7327; c) M. A. Larsen, J. F. Hartwig, *J. Am. Chem. Soc.* **2014**, *136*, 4287–4299.
- [29] T. S.-B. Lou, M. C. Willis, *Tetrahedron* **2020**, *76*, 130782.

Manuscript received: July 9, 2021

Accepted manuscript online: August 2, 2021

Version of record online: ■ ■ ■ ■ ■ ■ ■ ■ ■ ■


Research Articles



Cross-Coupling

X. A. F. Cook, L. R. E. Pantaine,
D. C. Blakemore, I. B. Moses, N. W. Sach,
A. Shavnya, M. C. Willis* — ■■■■–■■■■

Base-Activated Latent Heteroaromatic
Sulfonates as Nucleophilic Coupling
Partners in Palladium-Catalyzed Cross-
Coupling Reactions



Next-generation, base-activated latent sulfonate reagents have been successfully developed for use in the construction of heteroaromatic frameworks, such as 2-arylpyridines. Under Pd-catalyzed conditions, these species unmask to give the sulfinate *in situ*, which then undergoes efficient desulfonative cross-coupling with an array of (hetero)aryl halides.

