

Structural basis for specificity and promiscuity in a carrier protein/enzyme system from the sulfur cycle

Daniel B. Grabarczyk^a, Paul E. Chappell^b, Steven Johnson^b, Lukas S. Stelzl^a, Susan M. Lea^b, and Ben C. Berks^a.

^a Department of Biochemistry, University of Oxford, South Parks Road, Oxford OX1 3QU, United Kingdom.

^b Sir William Dunn School of Pathology, University of Oxford, South Parks Road, Oxford OX1 3RE, United Kingdom.

Corresponding author:

Ben C. Berks, Department of Biochemistry, University of Oxford, South Parks Road, Oxford OX1 3QU, United Kingdom.

Tel.: +44=1865-613294 E.mail: ben.berks@bioch.ox.ac.uk

Classification: Biological sciences – Biochemistry

Short title : Specificity in a carrier protein-enzyme system

Key words: promiscuous interactions; protein-protein interactions; structures of protein complexes; sulfur oxidation; Sox system; thiosulfate metabolism

Abstract

The bacterial Sox pathway is an important route for the oxidation of inorganic sulfur compounds. Intermediates in the Sox pathway are covalently attached to the heterodimeric carrier protein SoxYZ through conjugation to a cysteine on a protein swinging arm. We have investigated how the carrier protein shuttles intermediates between the enzymes of the Sox pathway using the interaction between SoxYZ and the enzyme SoxB as our model. The carrier protein and enzyme interact only weakly but we have trapped their complex using a 'suicide enzyme' strategy in which an engineered cysteine in the SoxB active site forms a disulfide bond with the incoming carrier arm cysteine. The structure of this trapped complex, together with calorimetric data, identifies sites of protein-protein interaction both at the entrance to the enzyme active site tunnel and at a second, distal, site. We find that the enzyme distinguishes between the substrate and product forms of the carrier protein through differences in their interaction kinetics and deduce that this behaviour arises from substrate-specific stabilization of a conformational change in the enzyme active site. Our analysis also suggests how the carrier arm-bound substrate group is able to out-compete the adjacent C-terminal carboxylate of the carrier arm for binding to the active site metal ions. We infer that similar principles underlie carrier protein interactions with other enzymes of the Sox pathway.

Significance Statement

Certain metabolic pathways use a carrier protein to shuttle covalently-attached intermediates between the active sites of enzymes. However, the details of the carrier protein-partner interactions have only been elucidated in a few cases. We have used biophysical methods and crystallography to obtain a molecular-level description of the interactions between a carrier protein and an enzyme involved in bacterial sulfur oxidation. Characterization of the contact sites between the two proteins suggests a basis for the promiscuous, but specific, binding interactions of the carrier protein. We also infer that the enzyme discriminates between the substrate- and product-bound forms of the carrier protein based on different interaction kinetics and link this behaviour to a structural change at the enzyme active site.

\body

Introduction

Intermediates in some metabolic pathways are covalently attached to a carrier protein to enhance their solubility or control their reactivity. Such intermediates are normally conjugated to a long flexible protein or cofactor arm on the carrier protein allowing the intermediate to be introduced into the buried active sites of the pathway enzymes (1). In some cases the carrier protein and partner enzymes form a permanent complex and may even be domains of a single polypeptide. Well-known examples include type I fatty acid synthases (2), non-ribosomal peptide synthetases (3), biotin carboxylases (4), and the mitochondrial α -keto acid dehydrogenases (1). In other cases the carrier protein and its partner enzymes are separate entities which form transient complexes to effect catalysis. Examples include the acyl carrier proteins interacting with enzymes involved in type II fatty acid synthesis and polyketide synthesis (5, 6). In reality these two types of carrier protein pathway are mechanistically similar because the carrier protein domain in the permanent complexes is normally itself flexibly tethered to the rest of the complex allowing significant freedom of movement between partner enzyme domains (1, 2).

An intriguing example of a carrier protein-dependent metabolic process is the Sox pathway located in the periplasmic compartment of many sulfur-oxidizing bacteria. This pathway oxidizes thiosulfate to produce electrons for use in respiratory energy generation or carbon fixation (7). Intermediates in the Sox pathway are covalently bound to the heterodimeric carrier protein SoxYZ (8, 9). The C-terminal peptide of SoxY forms a swinging arm bearing a Cys residue to which the pathway intermediates are conjugated (8, 9). Adjacent to this carrier arm is a conserved apolar pocket that is proposed to accommodate and protect labile intermediates (9). The current model for the Sox pathway postulates that thiosulfate is disulfide-linked to the

SoxYZ carrier arm Cys residue before being catabolized in a series of oxidative and hydrolytic steps catalysed by different enzymes (Fig. 1). Thus, SoxYZ carries a range of chemical species and must interact with multiple partner enzymes. Structures of the SoxYZ partner enzymes show that their active site access channels are wide enough to permit carrier arm-bound substrates to reach the catalytic groups (15-17). The proteins of the Sox pathway do not co-purify with each other from cell extracts. However, the independently-purified components can be mixed together to reconstitute Sox activity *in vitro* (18). These observations suggest that the interactions between the carrier protein and its enzymatic partners are weak and transient. Indeed, specific protein-protein interactions between SoxYZ and partner enzymes have yet to be experimentally demonstrated. A long, highly conserved, and flexible surface loop found immediately adjacent to the carrier arm on SoxYZ, termed the 'Z-loop', has been proposed to mediate SoxYZ interactions by packing on to partners of different structure (9).

A carrier protein, such as SoxYZ, forms part of the substrate of its enzyme partners. Thus, an understanding of catalysis in such systems requires an understanding of the protein-protein interactions made by the carrier protein. These interactions are distinct from normal enzyme-substrate interactions in two respects. Firstly, the substrate and product of an enzyme, as well as the other pathway intermediates, are all conjugated to the same carrier protein. Thus, any interactions the carrier protein makes with the enzyme which promote complex formation might correspondingly be expected to retard product release after catalysis. How this tension is resolved is not well understood in any carrier protein system. Secondly, the carrier protein must be able to make specific interactions with multiple, and usually structurally distinct, partners. This raises the question as to whether these promiscuous interactions are mediated by the same or different determinants on the carrier protein surface.



In this study we have used the interaction between the carrier protein SoxYZ and the Sox pathway enzyme SoxB as a model both for how the central component of the Sox pathway interacts with partners and as a test case for exploring carrier protein-partner interactions in

general. To stabilize the SoxB-SoxYZ complex for crystallization we developed a strategy in which the SoxY carrier arm was cross-linked to the SoxB active site. Combining structural and biophysical analysis we elucidate the molecular basis of the interaction between the carrier protein and its partner.

Results

Protein-protein interactions between SoxB and SoxYZ

In this study we have investigated carrier protein interactions in the Sox pathway using the SoxYZ and SoxB proteins from *Thermus thermophilus* as our model carrier protein-partner pair. SoxB is an enzyme that is inferred to catalyze the hydrolytic cleavage of thiosulfonate groups (-S-SO₃⁻) conjugated to the SoxY carrier arm (Fig. 1, Reaction 1)(16). This reaction has not been directly demonstrated because carrier protein-bound Sox pathway intermediates are not turned over unless all pathway components are present. However, we show that *T. thermophilus* SoxB will catalyse the hydrolysis of the small molecule substrate analogue trithionate (-O₃S-S-SO₃⁻) according to Reaction 2 (Fig. 2).



The observed trithionate hydrolase activity followed Michaelis-Menten kinetics with a K_M of 2.20 ± 0.15 mM (95% confidence limit, $n=3$), and a k_{cat} of 2 ± 0.1 s⁻¹ (SEM). SoxB contains a pair of Mn(II) ions at the active site to which the sulfone group (-SO₃⁻) of the non-hydrolyzable substrate analogue thiosulfate (-S-SO₃⁻) is seen to bind in a co-crystal (16, 19). The trithionate hydrolase activity of SoxB decreased in the presence of the metal chelator EDTA, but was increased when the assay mix was supplemented with Mn²⁺ ions, consistent with catalysis by the active site metal ions. The previously-reported SoxB-thiosulfate co-crystal structure suggests that active site residue Arg416 is involved in substrate binding and transition-state stabilisation (16). In agreement with this hypothesis we found that an Arg416Gly variant had undetectable trithionate hydrolase activity (Table 1).

We investigated whether the *T. thermophilus* SoxYZ has specific binding interactions with SoxB. In these experiments we used a SoxYZ variant in which the carrier arm Cys residue had been substituted with a Ser (SoxY_{C151S}Z). This variant allowed us to assess the protein-protein interactions between SoxB and SoxYZ independent of the contribution from the substrate-binding residue. Complex formation between SoxB and SoxY_{C151S}Z was observed at room temperature by both native PAGE (Fig. 3A) and size-exclusion chromatography in-line with multi-angle laser light scattering (SEC-MALLS) (Fig. 3B). The absolute molecular mass of the complex determined by MALLS was concentration dependent such that even at the highest protein concentration tested (40 μ M of each protein loaded on the column) the measured mass of 68 kDa was lower than the 90 kDa mass calculated for a 1:1 SoxB:SoxYZ complex. This behaviour indicates that the complex is unstable and partially dissociates during chromatography.

Isothermal titration calorimetry (ITC) was used to quantify the interaction between SoxB and SoxY_{C151S}Z. The binding isotherm was well fit by a simple 1:1 association model with a K_D of 3 μ M (Fig. 4A). This figure is consistent with the low micromolar K_M for the SoxYZ protein reported for a reconstituted Sox system (21). We took advantage of the inability of the isolated SoxB component to turn over its substrate to determine the contribution the SoxYZ-bound substrate group makes to the interaction between SoxB and SoxYZ. To this end the ITC experiment was repeated using the S-thiosulfonate derivative of SoxYZ ('SoxY(SSO₃)Z') (Fig. 4B). Addition of the thiosulfonate group made no significant difference to the affinity of SoxB for SoxYZ (compare Figs 4A and 4B). However, there was a large difference in the ΔH of the interaction, changing from exothermic (-3 kcal/mol) for SoxY_{C151S}Z to endothermic for SoxY(SSO₃)Z (+10 kcal/mol). Calculation shows that this enthalpy change was balanced by a change in T ΔS from -5.2 kcal/mol to -17.4 kcal/mol. These observations suggest that conjugation of substrate to SoxYZ alters some aspect of the mechanism of interaction with SoxB.

To further probe the influence that the group bound to the carrier arm of SoxYZ has on interactions with SoxB we repeated the ITC analysis with SoxYZ conjugated either to an S-carboxymethyl group ('SoxY(Ac)Z') or to its amide derivative ('SoxY(Am)Z'). The carboxymethyl group ($-\text{CH}_2\text{-CO}_2^-$) has physicochemical similarity to S-thiosulfonate. However, amidating this species produces a functional group ($-\text{CH}_2\text{-CONH}_2$) that, like SoxY_{C151S}Z, would be unable to provide the bidentate ligation of the active site manganese ions exhibited by the substrate analogue thiosulfate (16). SoxY(Ac)Z exhibits an endothermic enthalpy change on interaction with SoxB that is identical to that measured for SoxY(SSO₃⁻)Z (Fig. 4B,C) while SoxY(Am)Z shows an exothermic enthalpy change that is very close to that observed with SoxY_{C151S}Z (Fig. 4A,D). Thus, there is an apparent correlation between the thermodynamic characteristics of the SoxB-SoxYZ interaction and the potential of the SoxY-conjugated group to act as a bidentate ligand to the di-manganese centre.


We used surface plasmon resonance (SPR) to compare the interaction kinetics of SoxB with the substrate and product forms of SoxYZ. Equilibrium SPR experiments measured dissociation constants of 2.4 μM for the SoxB-SoxY(SSO₃⁻)Z substrate complex and 2.7 μM for the SoxB-SoxY_{C151S}Z product complex (Fig. S1). These are close to the values of 5 μM and 3 μM determined for the same complexes by ITC (Table 1) showing that the SPR technique reproduces the binding interactions seen in solution. Binding of the product analogue SoxY_{C151S}Z to SoxB was extremely rapid with association and dissociation complete within 0.5 seconds at all concentrations (Fig. 4E). By contrast, the kinetics of both association and dissociation between SoxB and the putative substrate, SoxY(SSO₃⁻)Z were markedly slower with association and dissociation now occurring over 10 seconds (Fig. 4F). The dissociation kinetics for this complex showed a small burst phase (Fig. S2). While this burst phase may indicate a two-step dissociation mechanism it more likely corresponds to the rapid dissociation of a small subpopulation of underivatized SoxYZ molecules. For both types of SoxB-SoxYZ complexes the association and dissociation phases were too fast to allow reliable extraction of kinetic parameters using standard multiple parameter fitting methods. However, the dissociation phase

kinetics could be fitted using simple exponential decay models giving an apparent rate constant of $5.7 \pm 0.8 \text{ s}^{-1}$ for dissociation of the SoxB-SoxY_{C151S}Z complex and apparent rate constants of $0.34 \pm 0.01 \text{ s}^{-1}$ and $4.5 \pm 0.7 \text{ s}^{-1}$, respectively, for the major dissociation phase and minor burst phase of the SoxB-SoxY(SSO₃)Z complex (Fig. S2). These figures reinforce the qualitative conclusion that k_{off} is significantly slower for the interaction involving the putative substrate. The observed dissociation rates would be compatible with the rate of turnover of 0.16 s^{-1} (thiosulfate) measured for the reconstituted Sox system of *P. pantotrophus* (22).

Preparation of a disulfide-linked SoxB-SoxYZ complex

To identify the specific molecular contacts involved in the interaction between SoxYZ and SoxB a structure of the SoxB-SoxYZ complex was required. Crystallization trials with the purified complex were unsuccessful, most likely due to the relatively low affinity of the interaction between the components. To overcome this problem we developed a strategy to covalently link the two proteins through formation of a disulfide bond between the Cys of the SoxY carrier arm and a Cys residue engineered into the SoxB active site. The rationale behind this approach was that a covalent linkage at the active site would prevent dissociation of the complex but would not interfere with protein-protein interactions outside the active site because the two proteins would be tethered together through the highly flexible SoxY carrier arm.

We targeted SoxB active site residue Trp175 for Cys substitution. Trp175 stabilizes the sulfane group of the substrate analogue thiosulfate in the previously determined SoxB-thiosulfate complex structure (16) and so was inferred to be close to the carrier arm Cys sulfur atom in the native SoxB-SoxYZ complex. To provide a leaving group in the disulfide bond-forming reaction we used the S-thiosulfonate derivative of the partner SoxYZ protein (Reaction 3).



Incubation of the SoxB Trp175Cys variant at 70°C with the S-thiosulfonate derivative of SoxYZ resulted in a high yield of disulfide-linked SoxB-SoxYZ complexes (Fig. S3). Mass spectrometry of the purified crosslinked product before and after incubation with the reductant DTT confirmed the expected disulfide linkage between SoxYZ and SoxB.

Structure of the disulfide-linked SoxB-SoxYZ complex

The crosslinked SoxB-SoxYZ complex was crystallized at pH 8.3 and the structure solved to a resolution of 3.3 Å (Fig. 5A, Table S1) using the high-resolution structures of the isolated components to define the complex and constrain the geometry during refinement. All four SoxB subunits in the asymmetric unit were well ordered and displayed interpretable electron density. By contrast, the SoxY and SoxZ subunits exhibited high B-factors and more ambiguous electron density in all but one copy of the complex (Fig. S4, Table S1). Nevertheless, all copies of the complex showed the same overall arrangement of the components. This was also true of the four copies of the complex in a 4.1 Å resolution structure obtained in a different space group (*P*12₁1) from crystals grown at pH 6.5. The structure of the most highly ordered copy of the SoxB-SoxYZ is used in the structural interpretation below and the quality of the electron density for this structure is shown in Figs. S5-S8.

The interface between SoxYZ and SoxB buries a surface area of 1220 Å² including the carrier arm (Fig. 5B). The relatively small size of this interface is consistent with the low stability of the complex (24). The binding interface shows two main sites of interaction (Fig. 5B). One site involves the area around the active site channel of SoxB. This we term the 'Y-patch'. It covers 81% of the total interface area including the active site channel (990 Å²). The other area, which we term the 'Z-patch', involves contacts between a SoxB surface loop and the face of a β-sheet in SoxZ. The Z-patch covers the remaining 19% of the interface area (234 Å²). These interaction

surfaces contain most of the conserved surface residues of both proteins (Fig. 5C). The presence of such multiple independent interaction patches is a common theme in protein-protein interactions (25).

SoxZ contains a large polypeptide loop ('the Z-loop') that was previously predicted to be involved in interactions with SoxYZ partner proteins (9). However, in the SoxB-SoxYZ complex structure the Z-loop is positioned well away from SoxB and does not participate in inter-protein contacts (Fig. 5A). Consistent with the interaction surfaces seen *in crystallo*, deletion of the Z-loop did not reduce the affinity of SoxYZ for SoxB in solution (Table 1, Fig. S9A).

Interactions between SoxZ and SoxB

The Z-patch on SoxZ is formed from the outward facing side chains of strands $\beta 1$, $\beta 2$ and $\beta 5$. A hydrophobic pocket is formed by Ile6, Val25, and Leu74 (Fig. 6A). Two arginine residues, SoxZ Arg8 and SoxY Arg117, are positioned at the edge of the pocket and the non-polar portion of their side chains contribute to the hydrophobicity of the pocket. The interacting surface on SoxB is formed from a surface loop with the sequence ²⁰²DDLFGD²⁰⁷. This loop creates a hydrophobic bulge surrounded by negatively charged residues that is complementary to the structure of the Z-patch on SoxZ. The non-polar SoxB residues Leu204 and Phe205 insert into the hydrophobic pocket of the SoxZ Z-patch whilst the side chains of SoxB Asp207 and SoxZ Arg8 are positioned close enough to form a salt-bridge at the centre of the Z-patch (Fig. 6A). It is conceivable that the other charged residues at the interface could form alternative salt bridges if there is some plasticity in the SoxB-SoxYZ interaction or could provide long range electrostatic interactions to aid docking as the interacting sites have opposite charges (Fig. 6B). The importance of long range electrostatic interactions in protein complex formation is well-established (27). The salt bridge and the presence of an aromatic residue at position 205 in SoxB are conserved in the SoxB and SoxZ proteins from the distantly related model organism *P. pantotrophus*, as expected

if these interactions are of biological relevance (Fig. 6C). Additionally, the basic electrostatic potential of the Z-patch is conserved in the structure of SoxYZ from *P. pantotrophus* (Fig. 6B).

It was important to establish whether the structure of the artificially crosslinked SoxB-SoxYZ complex seen in the crystals resembles the structure of the native SoxB-SoxYZ complex found in solution. To do this we designed SoxB variants that would disrupt the Z-patch interface seen in the crystallographic complex and then assessed their interaction with SoxYZ by ITC. Correct folding of the SoxB variants was verified by measuring their trithionate hydrolase activity (Table 1). Either a Phe205 to Ser substitution, which removes the hydrophobic 'knob' from the Z-patch interaction, or an Asp207 to arginine substitution, which introduces electrostatic repulsion across the Z-patch interface, abolished binding to SoxY_{C151S}Z (Table 1, Fig. S9C,E). Thus, single amino acid substitutions targeting the Z-patch interaction seen in the crystal structure also affect SoxYZ-SoxB interactions in solution, implying that the Z-patch is involved in native complex formation. The SoxB Phe205Ser substitution also prevented interaction with SoxY(Ac)Z (Table 1, Fig. S9D) indicating that SoxYZ binds via the Z-patch irrespective of the species conjugated to the carrier arm.

Interaction between SoxY and the SoxB substrate channel

The structure of *T. thermophilus* SoxYZ in the SoxB-SoxYZ complex can be compared with that of the isolated SoxYZ complex previously crystallized from *P. pantotrophus* (Fig. 7A). The only significant structural change in the SoxYZ protein on complex formation is that the carrier arm moves from a pocket formed between SoxY and SoxZ to the active site channel of SoxB (Fig. 7B). The C-terminal carboxylate of the SoxY polypeptide lies at the end of the carrier arm and bidently co-ordinates the active site manganese ions in a similar way to the sulfonate group of the substrate analogue thiosulfate in the structure of the thiosulfate-SoxB complex determined earlier (16). The SoxB active site channel is relatively wide compared to the thickness of the SoxY carrier arm, which runs along one side of the channel (Fig. 7B). The conformational change

undergone by the SoxYZ carrier arm on complex formation with SoxB is reminiscent of the 'switchblade' mechanism employed by acyl carrier proteins in which the non-polar substrate molecule bound to the phosphopantetheine arm is protected within a hydrophobic pocket on the side of the carrier protein, but swung fully out of the pocket to insert into the partner enzyme on complex formation [6].

The SoxB channel accommodates the C-terminal ¹⁴⁷TVGGCG-COOH portion of the SoxY carrier arm. SoxY residues Arg145, Ser143, and the loop comprising residues ⁶⁶AIAES⁷⁰ contact the surface of SoxB around the mouth of the active site channel (Fig. 7C). Residues Ala66 and Ile67 rest in a pocket formed by SoxB residues Tyr232, Val235, Asn463, Tyr471 and Gln473, together with SoxY Thr147 (Fig. 7C). Apart from this pocket, and the manganese coordination, the interaction surface between SoxY and SoxB is formed by a sparse hydrogen bonding network involving six direct interactions (Fig. 7C). Additional hydrogen bonding interactions are probably mediated by waters not seen at this resolution. Removing the C-terminal carrier arm from SoxY increased the K_D of the SoxB-SoxYZ interaction 50-fold (Table 1, Fig. S9B) confirming that carrier arm contacts are important for the association of SoxYZ with SoxB.

Conformational changes in the SoxB mobile loop

Our earlier structures (16) showed that binding of the substrate analogue thiosulfate to SoxB is associated with a widening of the active site channel through movement of a loop (residues 463 to 478) containing the highly conserved motif ⁴⁷²QQGGD⁴⁷⁶ (Fig. 8A). This conformational change is mediated through residue Asp476 in the mobile loop. In the unliganded SoxB structure the mobile loop is anchored close to the active site by a salt bridge between Asp476 and the catalytically important residue Arg416 (Fig. 8B, magenta). However, in the thiosulfate complex Arg416 co-ordinates the sulfonate part of thiosulfate in preference to Asp476 which, in turn, now forms a salt bridge with Arg385 (Fig. 8B, cyan). This switch in Asp476 bonding interactions causes the mobile loop to move away from the active site.

The SoxB-SoxYZ complex also shows a change in the conformation of the mobile loop relative to unliganded SoxB (Fig. 8A). However, this movement is small and in a different direction to the displacement induced by thiosulfate binding. The structural change induced by SoxYZ appears to be driven by the need to alleviate a number of steric clashes which would otherwise exist between SoxY and the SoxB mobile loop at the Y patch interface, namely those between SoxY Ile67 and SoxB Gln473, between SoxY Ala66 and SoxB Tyr471, and between SoxY Arg145 and SoxB Asn463 (Fig. S10). Further into the SoxB tunnel this displacement of the mobile loop leads to Asp476 pairing with Arg385, as in the SoxB-thiosulfate complex, rather than sequestering the catalytic residue Arg416, as occurs in the unliganded enzyme (Fig. 8B). The consequence of this structural re-arrangement is that Arg416 is made available to ligate the sulfonate moiety of the substrate.

Elucidation of the structural changes in SoxB induced by SoxYZ docking allows us to propose a molecular explanation for the observation that the substrate-conjugated form of SoxYZ is released from SoxB more slowly than the substrate-free form (Fig. 4E). We suggest that the presence of the substrate group stabilizes the displaced conformer of the SoxB mobile loop by providing SoxB Arg416 with a binding partner to replace the Asp476 interaction present in the resting state. Because mobile loop displacement minimizes the steric clashes between SoxB and SoxYZ, the substrate stabilization of the displaced conformer would be expected to lead to a reduction in the rate of SoxB-SoxYZ complex dissociation.

Basis for the selective binding of sulfonate to the active site manganese ions

The previously-determined structure of SoxB in complex with the substrate analogue thiosulfate suggests the sulfonate group of the substrate molecule must co-ordinate the SoxB active site manganese ions for catalysis to occur (16). However, the C-terminal carboxylate of the SoxY carrier arm is also present in the active site channel and will compete with the sulfonate group for binding to the metal ions. Indeed, active site co-ordination by the C-terminal

carboxylate is seen in the SoxB-SoxYZ complex structure (Fig. 7B,C). To gain insight into how SoxB is able to resolve this competition in favour of sulfonate ligation we used the disulfide-linked SoxB-SoxYZ structure to produce models of the physiological thiosulfonated complex with the manganese ions ligated either by the thiosulfonate group or the C-terminal carboxylate (Fig. 9).

Comparison of the two models shows that the different binding configurations place chemical groups of different character in contact with a ring of aromatic and non-polar residues which provide access to the manganese ions. When the SoxY C-terminal carboxylate co-ordinates the metal ions, the C-terminal SoxY Cys-Gly peptide is within the aromatic ring and participates in hydrogen-bonding interactions with the ring residues (Fig. 9A). However, in the substrate complex model, it is the hydrophobic side chain of cysteine-S-thiosulfonate that is within the aromatic ring and the interactions are now non-polar in nature (Fig. 9B). These different modes of interaction of the carrier arm with the SoxB aromatic ring provide a plausible mechanism for the change from net enthalpic to net entropic association between SoxB with SoxYZ observed following substrate conjugation to the SoxY carrier arm (Fig. 4A,B, Table 1). This is because the hydrogen bonding interactions seen with the C-terminal carboxylate interaction are enthalpically driven, while the non-polar interactions seen with the cysteine-S-thiosulfonate side chain are entropically driven.

In our structural models the S-thiosulfonate group and C-terminal carboxylate both use two terminal oxygen atoms to provide bi-dentate co-ordination to the manganese ions (Fig. 9). However, the S-thiosulfonate has an additional terminal oxygen atom which forms bonding interactions with SoxB Arg416 (Fig. 9B). To investigate whether this modelled interaction plays a role in selective binding of the S-thiosulfonate group by the active site we investigated the effect of substituting SoxB Arg416 with Gly on the thermodynamics of the interaction between SoxB and S-thiosulfonated SoxYZ. The mode of association changed from entropic to enthalpic (Table 1, Fig. S11). Given the correlation between binding mode and thermodynamics outlined

above this indicates that the carrier arm no longer ligates the active site through the thiosulfonate group but instead binds the metal ions by the C-terminal carboxylate. Thus, Arg416 is critical for correct positioning of the substrate group at the active site.

Discussion

Intermediates in the Sox protein transport system are conjugated to a Cys residue on the flexible C-terminal arm of the SoxYZ carrier protein and undergo reactions in the buried active sites of multiple partner proteins (Fig. 1). Using the SoxB-SoxYZ pair as our model we have, for the first time, been able to demonstrate binding interactions between the SoxYZ carrier protein and a partner enzyme. The measured low micromolar binding affinity between these proteins is appropriate for partners that have to engage in specific, but reversible, interactions and where the concentrations of the interacting proteins are likely to be of this order in the cell. It is highly likely that SoxYZ engages in equivalent binding interactions with the other enzymes of the Sox system.

Our work addresses three key questions about the way the SoxYZ protein interacts with its partners. Firstly, how does a single carrier protein specifically interact with multiple different enzymes? Secondly, how do the partner enzymes distinguish substrate-bearing carriers from those conjugated to other species? Thirdly, how does SoxB discriminate between the substrate sulfonate group and the carrier arm carboxy terminus?

Our SoxB-SoxYZ complex structure shows that SoxYZ and SoxB interact through multiple specific contact points on the surface of each molecule (Fig. 5). One set of interactions occur where the SoxY carrier arm enters the SoxB active site tunnel. At this contact point a SoxY surface loop adjacent to the carrier arm inserts into a pocket on SoxB. SoxB engages in sparse hydrogen bonding interactions with this loop, with the base of the carrier arm, and with the carrier arm itself within the active site tunnel (Fig. 7C). These carrier arm-associated interactions are supplemented by a second point of contact located away from the SoxB active site which involves insertion of a SoxB surface loop into a depression on the surface of SoxZ (Fig. 6). This 'Z-patch' contact is stabilized by matched hydrophobic and electrostatic interactions between the two interacting protein surfaces. Protein engineering experiments validate the interactions seen in the crystal structure by confirming that the association of

SoxYZ with SoxB in solution requires both the Z-patch and SoxY carrier arm contacts (Table 1), but does not involve the 'Z-loop' previously suggested to be the contact site (9).

Proteins that engage in promiscuous interactions commonly recognizing a partner protein using a surface binding site that is well-separated from the site of functional interaction (sometimes termed 'dual recognition')(28, 29). Distributing the binding interaction either partly or wholly to a distal site reduces the number of specific interactions that the functional domain needs to make with the partner protein. In this way the interaction of the functional domain with multiple, structurally distinct, partners is facilitated. In the case of SoxYZ the Z-patch interaction with SoxB permits the SoxY carrier arm to have only limited bonding interactions with the SoxB active site tunnel (Fig. 7C). In combination with the inherent flexibility of the carrier arm this would allow the carrier arm to be accommodated within active sites of different structure. A paucity of carrier arm interactions with the partner enzyme active site has also been observed for phosphopantetheine-containing carrier proteins suggesting that this is a general feature of carrier protein interactions (6,30-32).

In summary, our structural data show that SoxB has been adapted for interaction with SoxYZ by the straightforward alteration of a surface loop to provide the Z-patch interaction. Such a limited structural change could evolve without affecting the protein fold or catalytic site of the progenitor enzyme.

Are other SoxYZ-partner complexes stabilized in the same way as the SoxB-SoxYZ complex? This question cannot be definitively answered without determining the structure of each complex. It is, nevertheless, striking that the SoxB contact regions on SoxYZ correspond almost exactly to the regions of highest surface sequence conservation (Fig. 5B,C). This suggests that other partners interact with the same parts of SoxYZ that SoxB does. Inspection of the surfaces of the partner enzymes SoxAX and SoxCD shows that there is high sequence conservation around the mouths of the active site channels as expected if these enzymes, like SoxB, interact with the base of the SoxY carrier arm (Fig. S12). In each case there is also a highly conserved

surface patch located at a similar distance from the active site channel as the Z-patch is in SoxB. These patches resemble the SoxB Z-patch in being formed by protruding loops bearing a surface-exposed aromatic side chain surrounded by negatively charged amino acids. Thus, surface features resembling those used by SoxB to contact SoxYZ are present in other SoxYZ partner enzymes and it is plausible that SoxYZ interacts with all its partners by broadly the same mechanisms characterized here for the SoxB-SoxYZ interaction.

It has not previously been possible to exclude the possibility that the components of the Sox pathway form a permanent supercomplex *in vivo* that is disrupted by purification. The structure of the SoxYZ-SoxB complex precludes this possibility since the structure shows that SoxYZ would need to dissociate from SoxB to allow the carrier arm to access another enzyme active site.

For the Sox pathway to operate efficiently it is important that the enzyme partners of SoxYZ are able to selectively interact with SoxYZ carrying the substrate form of the relevant pathway intermediate. The Y-patch and Z-patch interactions just discussed would not be affected by the species conjugated to the SoxYZ carrier arm and so an additional mechanism is required to enable SoxB to preferentially interact with the substrate form of the carrier protein. Our data strongly suggest that this selectivity is kinetically determined because we find that the apparent dissociation rate constant for the substrate form of SoxYZ is an order of magnitude lower than that of substrate-free SoxYZ (Fig. 4E,F). Since dissociation is a unimolecular reaction this difference in rates will be maintained independent of variation of protein concentration or macromolecular crowding effects. Our structural data suggests that the observed differences in dissociation kinetics can be explained by the operation of an induced fit mechanism at the SoxB active site. In the resting SoxB enzyme the catalytically essential residue Arg416 is sequestered through ion pairing to a mobile loop that lies above the active site. Upon SoxYZ binding to SoxB the mobile loop is displaced as a consequence of steric clashes with the incoming carrier protein (Figs. 8A and S10). This change in mobile loop conformation results in the release of the Arg416

side chain which is then able to bind the sulfonate group of the substrate. Interaction with the substrate group discourages Arg416 from re-forming the ion pair with the mobile loop. This has the effect of shifting the loop conformer equilibrium further towards the displaced state (Fig. 8B). Thus, the mobile loop co-operatively couples SoxYZ binding at the surface of SoxB to substrate binding at the active site. Following catalysis the substrate group is lost and so no longer contributes to stabilizing (through interaction with Arg416) the displaced conformer of the mobile loop. Because it is the displaced loop conformer that minimizes steric clashes with SoxY, disfavoring this conformer has the effect of accelerating the rate at which the SoxB-SoxYZ complex dissociates. This model provides a molecular explanation for why the substrate form of the SoxYZ carrier protein leaves SoxB more slowly than the substrate-free form.

From the point of view of enzyme specificity our model implies that only the substrate form of SoxYZ stabilizes the catalytically relevant conformer of the SoxYZ-SoxB complex and so only this form of the carrier protein is kinetically partitioned into the catalytically productive forward reaction path rather than reversing the initial association event (33). By similar logic the destabilization of the catalytic conformer that occurs following product formation can be viewed as promoting product release relative to reversal of the catalytic reaction.

The conformational changes associated with substrate binding may also assist in preventing competitive inhibition of SoxB by the substrate analogue thiosulfate, a molecule which will always be present during the operation of the Sox pathway. The crystal structure of SoxB in complex with thiosulfate shows that thiosulfate binding, like binding of the substrate form of SoxYZ, involves displacement of the mobile loop (Fig. 8)(16). However, while for the physiological substrate this conformational change is aided by a steric clash with the SoxY carrier arm (above), similar assistance is not available during thiosulfate binding. Thus the necessity for conformational change at the active site on ligand binding presents a significantly greater energetic barrier to the binding of thiosulfate than it does to the binding of carrier arm-conjugated S-thiosulfonate.

The final selectivity issue that must be overcome in order for SoxB to efficiently interact with SoxYZ during catalysis is that the enzyme must ensure that the catalytic metal ions bind the substrate sulfonate group in preference to the adjacent carrier arm C-terminal carboxylate group. Our structural, modelling, and biophysical data suggest two mechanisms by which this competition is resolved in favour of the substrate group. Firstly, the extra oxygen atom found in the sulfonate group forms an ion pair with Arg416 and so increases the binding strength of the sulfonate group over the carboxylate moiety. Secondly, it is more energetically favourable to place the relatively non-polar cysteine-S-thiosulfonate side chain of the substrate group in the hydrophobic environment above the active site metal ions than it is to move the more polar carboxy terminus of the carrier arm to this position.

Materials and Methods

Genetic constructs and protein production

Plasmid construction, mutagenesis, protein expression and purification, as well as modelling the native SoxB-SoxYZ complex are described in SI Materials and Methods.

Protein chemistry

The S-thiosulfonate derivative of SoxYZ [$\text{SoxY}(\text{SSO}_3^-)\text{Z}$] was generated by reacting SoxYZ with potassium tetrathionate (Fluka) as described (16). The S-carboxymethyl and S-carboxyamidomethyl derivatives of SoxYZ [$\text{SoxY}(\text{Ac})\text{Z}$ and $\text{SoxY}(\text{Am})\text{Z}$] were generated by reaction with, respectively, iodoacetate or iodoacetamide (both Sigma-Aldrich) using a published protocol (34). Following derivatization small molecules were removed from SoxYZ by size-exclusion chromatography on a Superdex 75 10/300 column (GE Healthcare).

The disulfide-linked SoxB-SoxYZ complex was produced by incubating 50 μM SoxB(W175C) with 50 μM $\text{SoxY}(\text{SSO}_3^-)\text{Z}$ in ITC Buffer for one hour at 70°C. Precipitate was removed by centrifugation at 16000 x g, and the complex was purified by loading on a 1 mL histrap HP column (GE Healthcare) and eluting with a 10 column volume gradient from 25 to 210 mM imidazole in ITC buffer. The complex was further purified by size-exclusion chromatography on a Superdex 200 10/300 column (GE Healthcare) equilibrated in 30 mM Tris-HCl pH 8.0. For crystallization trials the purified sample was diluted to achieve a final buffer concentration of 10 mM Tris-HCl pH 8.0 and the protein complex then concentrated with a Millipore Amicon Ultra 2 ml 10K spin concentrator.

Electrospray-ionisation mass spectrometry (ESI-MS) was used to confirm that SoxYZ samples had been correctly and quantitatively derivatized, and to confirm successful formation of the disulfide linked SoxB-SoxYZ complex.

Analysis of trithionate hydrolysis

Sodium trithionate was synthesised by reacting sodium thiosulfate with hydrogen peroxide (35). Trithionate hydrolysis assays were carried out at 70°C and contained 1.3 mM trithionate, 10 mM HEPES-NaOH pH 6.8, 1 mM MnCl₂. Activities were corrected for the rate of non-enzymatic hydrolysis. Trithionate and thiosulfate concentrations were quantified by cyanolysis using sodium thiosulfate standards of known concentration (35). Sulfate concentrations were determined using the barium sulfate assay calibrated with sodium sulfate standards of known concentration (36). Michaelis-Menten parameters were determined using 0.5 µM SoxB. Initial velocities were estimated from the thiosulfate concentration after 10 min. K_M and k_{cat} were calculated by non-linear curve-fitting to the standard Michaelis-Menten equation using OriginPro 8.5.1 (OriginLab). The activities of SoxB variants were assessed as sulfate production from 5 mM sodium trithionate in 10 minutes.

Biophysical techniques

ITC, surface plasmon resonance (SPR), ESI-MS and size-exclusion chromatography multi-angle laser light scattering (SEC-MALLS) were all performed using standard procedures which are described in detail in SI Materials and Methods. The methods used to fit the SPR dissociation curves are described in the caption of Fig. S2.

Protein crystallisation, X-ray data collection, structure solution, and refinement

Crystals of the disulfide-linked SoxB-SoxYZ complex were obtained by the vapour-diffusion method using 0.55 μ l sitting drops containing 70% protein solution (93 μ M) and 30% mother liquor (0.1 M Tris-HCl, 0.2 M (NH₄)₂SO₄, 8.55% PEG 8000, pH 8.3). Drops were incubated at 20°C and equilibrated against 120 μ l of mother liquor.

Crystals were cryoprotected in 30% ethylene glycol and 70% mother liquor and flash cooled in liquid nitrogen. Multiple crystals diffracted to \sim 4.5 Å but a single crystal diffracted to better than 4 Å. Diffraction data were collected at 100K to 3.28 Å on the i04 beamline at the Diamond Light Source, Oxfordshire, England. Automatic data processing was carried out with the Xia2 package (37). The limited resolution and high Wilson B Factor (83 Å²) of the diffraction data suggest that inherent mobility of the molecules within the crystal lattice limit order. Molecular replacement utilised Phaser (38) to sequentially find search models derived from *T. thermophilus* SoxB (PDB:2WDF), *T. thermophilus* SoxZ (PDB:1V8H), and *P. pantotrophus* SoxY (PDB:20X5). The SoxY carrier arm was omitted from the molecular replacement model but difference density for the arm could be clearly seen within the SoxB active site following placement of the remaining protein components.

Cycles of refinement and rebuilding were carried out using the computer graphics program Coot (39) and the autoBUSTER (Global Phasing) refinement package, with non-crystallographic symmetry restraints and targeting of the structure to the previously determined, high resolution, structures of the individual components of the complex (40) to help condition the refinement. Although characterised by different thermal mobilities (see the by-chain average B factors in Table S1 and residue-by-residue B factors in Fig. S4) there are no significant structural rearrangements between the crystallographically independent copies of each protein chain (average RMSD C α between equivalent chains 0.5 \pm 0.2 Å; residue-by-residue RMSD analysis in Fig. S13) even though the high mobility means that the models for Sox Z are not complete in the more poorly ordered copies. Whilst at this resolution side chains in the mobile regions of the

proteins are poorly defined, side chains at the protein-protein interfaces important for this study are resolved. Close inspection of the 7 Ramachandran outliers revealed that 4 outliers were the independent copies of SoxB residue 174 that coordinates a Mn^{2+} . This residue is also a Ramachandran outlier in the high resolution structure of SoxB deposited as PDB:2WDF. The other three outliers likely reflect errors in the model and are outliers in a single copy of each chain. RMSDs to the models used for molecular replacement are 0.3 ± 0.0 Å to 2WDF (SoxB), 1.5 ± 0.3 Å to 1V8H (SoxZ) and 1.9 ± 0.5 Å to 2OX5 (SoxY). The structure and diffraction data have been deposited in the Protein Data Bank (PDB:4UWQ). Structural figures were produced using the PyMol Molecular Graphics System, Version 1.3 (Schrödinger, LLC).

Acknowledgements

We thank Colin Kleanthous for comments on the manuscript and Martin Krehenbrink, Pietro Roversi, David Staunton, and Mark Sansom for technical advice. DG, PC, and LS were supported by studentships from the Biotechnology and Biological Sciences Research Council. SJ was supported by Medical Research Council grant G0900888. This work used the University of Oxford Department of Biochemistry Biophysics Facility. We acknowledge Diamond Light Source for time on beam line I04 under proposal MX9306.

References

1. Perham RN (2000) Swinging arms and swinging domains in multifunctional enzymes: Catalytic machines for multistep reactions. *Annual review of biochemistry* 69:961-1004.
2. Leibundgut M, Maier T, Jenni S, & Ban N (2008) The multienzyme architecture of eukaryotic fatty acid synthases. *Curr Opin Struc Biol* 18(6):714-725.
3. Tanovic A, Samel SA, Essen LO, & Marahiel MA (2008) Crystal structure of the termination module of a nonribosomal peptide synthetase. *Science* 321(5889):659-663.
4. Tong L (2013) Structure and function of biotin-dependent carboxylases. *Cellular and Molecular Life Sciences* 70(5):863-891.
5. Crosby J & Crump MP (2012) The structural role of the carrier protein - active controller or passive carrier. *Nat Prod Rep* 29(10):1111-1137.
6. Nguyen C, *et al.* (2014) Trapping the dynamic acyl carrier protein in fatty acid biosynthesis. *Nature* 505(7483):427-431.
7. Frigaard NU & Dahl C (2009) Sulfur metabolism in phototrophic sulfur bacteria. *Advances in microbial physiology* 54:103-200.
8. Quentmeier A & Friedrich CG (2001) The cysteine residue of the SoxY protein as the active site of protein-bound sulfur oxidation of *Paracoccus pantotrophus* GB17. *FEBS letters* 503(2-3):168-172.
9. Sauve V, Bruno S, Berks BC, & Hemmings AM (2007) The SoxYZ complex carries sulfur cycle intermediates on a peptide swinging arm. *The Journal of biological chemistry* 282(32):23194-23204.
10. Hensen D, Sperling D, Truper HG, Brune DC, & Dahl C (2006) Thiosulphate oxidation in the phototrophic sulphur bacterium *Allochromatium vinosum*. *Molecular microbiology* 62(3):794-810.
11. Welte C, *et al.* (2009) Interaction between Sox proteins of two physiologically distinct bacteria and a new protein involved in thiosulfate oxidation. *FEBS letters* 583(8):1281-1286.
12. Chen ZW, *et al.* (1994) The structure of flavocytochrome *c* sulfide dehydrogenase from a purple phototrophic bacterium. *Science* 266(5184):430-432.
13. Bardischewsky F, Quentmeier A, & Friedrich CG (2006) The flavoprotein SoxF functions in chemotrophic thiosulfate oxidation of *Paracoccus pantotrophus* *in vivo* and *in vitro*. *FEMS microbiology letters* 258(1):121-126.
14. Dahl C, Franz B, Hensen D, Kesselheim A, & Zigann R (2013) Sulfite oxidation in the purple sulfur bacterium *Allochromatium vinosum*: identification of SoeABC as a major player and relevance of SoxYZ in the process. *Microbiology* 159(Pt 12):2626-2638.
15. Bamford VA, *et al.* (2002) Structural basis for the oxidation of thiosulfate by a sulfur cycle enzyme. *The EMBO journal* 21(21):5599-5610.
16. Sauve V, *et al.* (2009) Mechanism for the hydrolysis of a sulfur-sulfur bond based on the crystal structure of the thiosulfohydrolase SoxB. *The Journal of biological chemistry* 284(32):21707-21718.
17. Zander U, *et al.* (2011) Structural basis for the oxidation of protein-bound sulfur by the sulfur cycle molybdohemo-enzyme sulfane dehydrogenase SoxCD. *The Journal of biological chemistry* 286(10):8349-8360.

18. Rother D, Henrich HJ, Quentmeier A, Bardischewsky F, & Friedrich CG (2001) Novel genes of the *sox* gene cluster, mutagenesis of the flavoprotein SoxF, and evidence for a general sulfur-oxidizing system in *Paracoccus pantotrophus* GB17. *Journal of bacteriology* 183(15):4499-4508.
19. Epel B, Schafer KO, Quentmeier A, Friedrich C, & Lubitz W (2005) Multifrequency EPR analysis of the dimanganese cluster of the putative sulfate thiohydrolase SoxB of *Paracoccus pantotrophus*. *J Biol Inorg Chem* 10(6):636-642.
20. Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227(5259):680-685.
21. Welte C, *et al.* (2009) Interaction between Sox proteins of two physiologically distinct bacteria and a new protein involved in thiosulfate oxidation. *FEBS Letters* 583(8):1281-1286.
22. Bardischewsky F, *et al.* (2005) Sulfur dehydrogenase of *Paracoccus pantotrophus*: the heme-2 domain of the molybdoprotein cytochrome *c* complex is dispensable for catalytic activity. *Biochemistry* 44(18):7024-7034.
23. Celniker G, *et al.* (2013) ConSurf: Using Evolutionary Data to Raise Testable Hypotheses about Protein Function. *Israel Journal of Chemistry* 53(3-4):199-206.
24. Kastiris PL, *et al.* (2011) A structure-based benchmark for protein-protein binding affinity. *Protein Sci* 20(3):482-491.
25. Reichmann D, *et al.* (2005) The modular architecture of protein-protein binding interfaces. *Proceedings of the National Academy of Sciences of the United States of America* 102(1):57-62.
26. Baker NA, Sept D, Joseph S, Holst MJ, & McCammon JA (2001) Electrostatics of nanosystems: Application to microtubules and the ribosome. *Proceedings of the National Academy of Sciences* 98(18):10037-10041.
27. Schreiber G, Haran G, & Zhou HX (2009) Fundamental aspects of protein-protein association kinetics. *Chemical reviews* 109(3):839-860.
28. Schreiber G & Keating AE (2011) Protein binding specificity versus promiscuity. *Curr Opin Struct Biol* 21(1):50-61.
29. Meenan NA, *et al.* (2010) The structural and energetic basis for high selectivity in a high-affinity protein-protein interaction. *Proceedings of the National Academy of Sciences of the United States of America* 107(22):10080-10085.
30. Liu Y, Zheng TF, & Bruner SD (2011) Structural basis for phosphopantetheinyl carrier domain interactions in the terminal module of nonribosomal peptide synthetases. *Chem Biol* 18(11):1482-1488.
31. Sundlov JA, Shi C, Wilson DJ, Aldrich CC, & Gulick AM (2012) Structural and functional investigation of the intermolecular interaction between NRPS adenylation and carrier protein domains. *Chem Biol* 19(2):188-198.
32. Agarwal V, Lin S, Lukk T, Nair SK, Cronan JE (2012) Structure of the enzyme-acyl carrier protein (ACP) substrate gatekeeper complex required for biotin synthesis. *Proceedings of the National Academy of Sciences of the United States of America* 109(43):17406-11.

33. Johnson KA (2008) Role of induced fit in enzyme specificity: a molecular forward/reverse switch. *The Journal of biological chemistry* 283(39):26297-26301.
34. Crankshaw MW & Grant GA (2001) Modification of cysteine. *Current Protocols in Protein Science*, (John Wiley & Sons, Inc.).
35. Kelly DP, Wood AP, & Harry D. Peck JLL (1994) [35] Synthesis and determination of thiosulfate and polythionates. *Methods in Enzymology*, (Academic Press), Vol Volume 243, pp 475-501.
36. Kolmert Å, Wikström P, & Hallberg KB (2000) A fast and simple turbidimetric method for the determination of sulfate in sulfate-reducing bacterial cultures. *Journal of Microbiological Methods* 41(3):179-184.
37. Winter G (2010) xia2: an expert system for macromolecular crystallography data reduction. *J Appl Crystallogr* 43(Pt 1): 186-190.
38. McCoy AJ, *et al.* (2007) Phaser crystallographic software. *J Appl Crystallogr* 40(Pt 4):658-674.
39. Emsley P, Lohkamp B, Scott WG, & Cowtan K (2010) Features and development of Coot. *Acta crystallographica. Section D, Biological crystallography* 66(Pt 4):486-501.
40. Blanc E, *et al.* (2004) Refinement of severely incomplete structures with maximum likelihood in BUSTER-TNT. *Acta crystallographica. Section D, Biological crystallography* 60(Pt 12 Pt 1):2210-2221.

Figure Legends

Figure 1. Model for the oxidation of thiosulfate by the Sox system. Pathway intermediates are carried by the SoxYZ complex through conjugation to a cysteine residue on the SoxY swinging arm (shown as -S). A small *c*-type cytochrome acts as the direct electron acceptor from the Sox enzymes. In some sulfur-oxidizing bacteria the Sox pathway is cyclical and thiosulfate is completely oxidized to sulfate (solid arrows). Other sulfur oxidizing bacteria lack the SoxCD enzyme and the sulfane intermediate is used to produce polymeric sulfur species in the form of sulfur globules which are subsequently oxidized to sulfate by a cytoplasmic pathway (dotted arrow)(10). Sulfane transfer between SoxYZ and the sulfur globules is likely mediated by the rhodanese-like protein SoxL (11). As well as partnering the canonical Sox enzymes, SoxYZ probably also interacts with SoxEF/FccAB (12), a flavocytochrome *c* with a poorly-defined function in the Sox mechanism (13). Additionally, SoxYZ has been implicated in sulfite oxidation in *Allochromatium vinosum* where it is suggested sulfite is carried as a SoxY-cysteine-S-sulfinate derivative which then interacts with unidentified partners (14).

Figure 2. SoxB possesses trithionate hydrolase activity. (A) Reaction containing 0.5 μ M SoxB. (B) Non-enzymatic control. Error bars show confidence limits ($p = 0.05$) from 3 experiments.

Figure 3. SoxYZ forms a weak complex with SoxB. (A) Comparative native PAGE analysis of SoxY_{C151S}Z, SoxB, or a mixture of the two proteins (left). Each sample analysed contained 10 μ M (per component) protein. Native PAGE employed the Laemmli buffer system (20) and a 7% polyacrylamide gel. A major band present only in the mixed sample is indicated with *. The polypeptide composition of this band was determined by excising the band, dehydrating the gel slice in acetonitrile followed by boiling in SDS-containing buffer and analysis on an SDS-PAGE

gel (right). Note that *T. thermophilus* SoxY and SoxZ have identical electrophoretic mobilities in SDS-PAGE. **(B)** SEC-MALLS analysis of SoxY_{C151S}Z, SoxB, or mixtures of the two proteins. Measured average molar masses are shown for loading concentrations (per protein component) of 10 μ M, 20 μ M and 40 μ M (dashed lines, with the loading concentrations indicated for the SoxB/SoxY_{C151S}Z mixture). The normalized excess Rayleigh ratio is shown for the 40 μ M concentration samples only (solid lines).

Figure 4. Biophysical analysis of the interaction between SoxYZ and SoxB. (A-D)

Isothermal calorimetry experiments. In each case integrated heats from a representative experiment are shown. The fit to the data and corresponding K_D and enthalpy change values are from duplicate experiments. Experiments differ in the derivatization of the SoxY carrier arm cysteine residue as shown schematically by the cartoon in each panel. **(A)** SoxY_{C151S}Z titrated into 47 μ M SoxB. **(B)** SoxY(SSO₃⁻)Z titrated into 50 μ M SoxB. **(C)** SoxY(Ac)Z titrated into 50 μ M SoxB. **(D)** SoxY(Am)Z titrated into 58 μ M SoxB. **(E-F)** Representative surface plasmon resonance sensograms showing the kinetics of the association and dissociation of **(E)** SoxY_{C151S}Z or **(F)** SoxY(SSO₃⁻)Z to a SoxB-coated sensor chip. The concentrations of injected SoxYZ samples are indicated.

Figure 5. Structure of a disulfide-linked SoxB-SoxYZ complex. (A)

Overall structure of the complex with proteins in cartoon representation and the manganese ions shown as purple spheres. Stick representation is used to show the disulfide bond between the SoxY carrier arm Cys and SoxB Cys175, and for SoxB residue Phe205 that contributes to the Z-patch. **(B)** Surface representation of the interacting faces of the SoxB and SoxYZ proteins. The surfaces which are buried upon interaction are shown in blue. **(C)** The same views of the SoxB and SoxYZ proteins as in *B* but with the surface coloured according to sequence conservation using the program Consurf (25). Magenta indicates areas of highest sequence conservation and cyan the most

variable sequences. Note that the Z-patch conservation in SoxB and SoxYZ is probably under reported due to alignment difficulties caused by insertions and deletions in adjacent sequences in the proteins from Purple Sulfur Bacteria and Green Sulfur Bacteria.

Figure 6. The SoxB-SoxZ interface. (A) Molecular details of the SoxB-SoxZ interface. The SoxB surface is colored gray, the SoxY surface orange, and the SoxZ surface yellow. (B) The surface potentials around the Z-patch in the SoxYZ proteins of *T. thermophilus* (from the SoxB-SoxYZ complex) and *P. pantotrophus* (2OX5), and around the Z-patch of *T. thermophilus* SoxB. The surface potential was calculated using Adaptive Poisson-Boltzmann Solver (26). (C) Sequence of the Z-patch loop in SoxB and the Z-patch region in SoxZ for the proteins of *T. thermophilus* (Tt) and *P. pantotrophus* (Pp) with structurally important residues marked (•).

Figure 7. The SoxB-SoxY interface. (A) Backbone alignment of *T. thermophilus* SoxYZ from the SoxB-SoxYZ complex (SoxY orange; SoxZ yellow; carrier arm red) with *P. pantotrophus* SoxYZ (PDB:2OX5; SoxY cyan; SoxZ light blue; carrier arm blue). The RMSD between the structures is 1.6 Å over 78 equivalent Cα atoms. (B) Position of the SoxY carrier arm within the SoxB active site channel. SoxY (orange) and SoxZ (yellow) are shown in ribbon representation with the carrier arm in sticks representation. SoxB is shown as a section through a space-filling model. The manganese atoms are shown as purple spheres. (C) Molecular details of the interactions between SoxY and SoxB with relevant residues shown in stick representation and the manganese atoms as purple spheres. The SoxY ⁶⁵PAIAES⁷⁰ loop is highlighted in green.

Figure 8. Conformational changes related to the SoxB mobile loop. (A) Structures of SoxB in complex with SoxYZ (this work), thiosulfate (PDB:2WDE), or unliganded (PDB:2WDC), were backbone aligned and are displayed with the SoxB mobile loop in gray (SoxYZ complex), cyan

(thiosulfate complex), or magenta (unliganded structure). The remainder of the SoxB molecule is shown in gray and the SoxYZ molecule in orange and yellow. **(B)** Overlay of the mobile loop-interacting salt-bridge network in the three structures shown in **(A)** with the same coloring scheme as in **(A)**. Thiosulfate and relevant amino acid residues are shown in stick representation where oxygen is red, nitrogen blue, and sulfur yellow. The manganese ions are represented by purple spheres.

Figure 9. Models of the active site of the SoxB-SoxY(SSO₃⁻)Z complex. Models were constructed based on the native SoxB-SoxYZ structure as described in SI Materials and Methods. The carbon atoms of SoxB are shown in grey and of SoxY in orange. Oxygen atoms are red, nitrogen atoms blue, and sulfur atoms yellow. The manganese ions are represented by purple spheres. **(A)** Complex with the SoxY C-terminal carboxylate co-ordinating the SoxB manganese ions. The interaction of SoxB with the SoxY C-terminal Cys-Gly peptide is stabilised by hydrogen bonding. **(B)** Complex with the S-thiosulfonate group co-ordinating the manganese ions. The interaction of SoxB with the cysteine-S-thiosulfonate is stabilised by an arc of hydrophobic residues.