

## Hippocampal mGluR1-dependent long-term potentiation requires NAADP-mediated acidic store $\text{Ca}^{2+}$ signaling

William J. Foster<sup>1\*</sup>, Henry B. C. Taylor<sup>1</sup>, Zahid Padamsey<sup>1</sup>, Alexander F. Jeans<sup>1</sup>, Antony Galione<sup>1\*</sup>, and Nigel J. Emptage<sup>1\*</sup>

<sup>1</sup>Department of Pharmacology, University of Oxford, Mansfield Road, Oxford, OX1 3QT, United Kingdom. \*Corresponding author. Email: [william.foster@pharm.ox.ac.uk](mailto:william.foster@pharm.ox.ac.uk) (WJF); [nigel.emptage@pharm.ox.ac.uk](mailto:nigel.emptage@pharm.ox.ac.uk) (NJE); [antony.galione@pharm.ox.ac.uk](mailto:antony.galione@pharm.ox.ac.uk) (AG).

### Abstract:

Acidic organelles, such as endosomes and lysosomes, store  $\text{Ca}^{2+}$  that is released in response to intracellular increases in the second messenger nicotinic acid adenine dinucleotide phosphate (NAADP). In neurons, NAADP and  $\text{Ca}^{2+}$  signaling contribute to synaptic plasticity, a process of activity-dependent long-term potentiation (LTP) [or, alternatively, long-term depression (LTD)] of synaptic strength and neuronal transmission that is critical for neuronal function as well as memory formation. We explored the function of and mechanisms regulating acidic  $\text{Ca}^{2+}$  store signaling in murine hippocampal neurons. We found that metabotropic glutamate receptor 1 (mGluR1) was coupled to NAADP signaling that elicited  $\text{Ca}^{2+}$  release from acidic stores. In turn, this released  $\text{Ca}^{2+}$  mediated mGluR1-dependent LTP by transiently inhibiting SK-type  $\text{K}^+$  channels, possibly through the activation of protein phosphatase 2A. Genetically removing two-pore channels (TPCs), which are endo-lysosomal-specific ion channels, switched the polarity of plasticity from LTP to LTD, indicating the importance of specific receptor-store coupling and providing mechanistic insight into how mGluR1 can produce both synaptic potentiation and synaptic depression.

## Introduction:

Acidic storage vesicles, such as lysosomes and endosomes, have traditionally been viewed as compartments for degradation and recycling of cellular metabolites. However, they have also been recognised as having additional signalling roles, including in intracellular  $\text{Ca}^{2+}$  signalling, as through their release of  $\text{Ca}^{2+}$  in response to the second messenger nicotinic acid adenine dinucleotide phosphate (NAADP) [1-6]. At present, the functions of NAADP signalling in the central nervous system (CNS) are incompletely understood. NAADP evokes  $\text{Ca}^{2+}$  release from brain microsomes [7], and NAADP-binding sites are present throughout the brain [8], suggesting that NAADP is an important signalling molecule in the CNS. Thus far, several key areas in which NAADP signalling plays a role have been identified: (i) in differentiation and growth of [9-11] and augmentation of neurite outgrowth in [12] cortical neurons; (ii) in autophagy in astrocytes, in which NAADP promotes autophagosome formation, which can be reduced in the presence of a dominant negative TPC2, a lysosomal two-pore channel isoform targeted by NAADP [13] or with the NAADP antagonist Ned-19 [14]; and (iii), in regulating neuronal excitability, neurotransmitter release, and synaptic plasticity. The application of membrane-permeable NAADP (NAADP-AM) reportedly causes membrane depolarization in medullary neurons [15] and application of NAADP enhances neurotransmitter release in *Aplysia californica* [16] and at neuromuscular junctions in frogs [17]. Work from our laboratory has shown that acidic store  $\text{Ca}^{2+}$  signalling enhances neurotransmitter release [18] and is essential for dendritic spine growth associated with late-phase long term potentiation (LTP) [19] in hippocampal CA1 pyramidal neurons from rodents.

The activation of various metabotropic receptors reportedly induces NAADP synthesis in non-neuronal cell types [20-24], and the application of glutamate, which can activate metabotropic glutamate receptors (mGluRs), also reportedly stimulates NAADP synthesis in both neurons [25] and astrocytes [14]. However, the specific glutamate receptor subtype(s) responsible for NAADP synthesis has thus far remained elusive. Here, we examined NAADP-mediated changes in neuronal excitability and found that these NAADP-mediated effects are driven by mGluR1 activation. Critically, this pathway led us to identify a unique role for NAADP in the genesis of mGluR1-mediated synaptic potentiation.

## **Results:**

### **NAADP-AM causes membrane depolarization in pyramidal neurons of the hippocampus**

NAADP is a charged molecule, making it necessary to bond it to an acetoxymethyl ester group (AM) for it to be membrane permeable [26]. To explore the effect of NAADP on membrane excitability, we introduced this engineered molecule (NAADP-AM) extracellularly, via a glass pipette, onto pyramidal neurons in CA3 and CA1 regions of hippocampal slice cultures. We chose cells that were localized to the surface of the tissue to improve the delivery of NAADP-AM.

Extracellular application of NAADP-AM (pipette concentration of 300  $\mu$ M) produced transient membrane depolarizations (denoted by  $\Delta V_M$ ; **Figure 1, B to D**) in both CA3 and CA1 pyramidal neurons of the hippocampus. The effect was highly consistent as the application of NAADP-AM at 30-second intervals produced reproducible membrane depolarization. Application of vehicle alone or NAADP alone, in other words the “no AM” group, failed to produce depolarization (**Figure 1, B and C**).

To then examine the mechanism by which NAADP-AM produced the membrane depolarization, we performed a series of pharmacological manipulations in combination with the extracellular application of NAADP-AM. A characteristic feature of NAADP signalling in mammals is that at high concentrations NAADP itself can desensitize the NAADP receptor [27]. This feature allows for a specific pharmacological inhibition of NAADP signalling. Desensitization of the NAADP receptor with 1 mM NAADP in the patch pipette abolished responses to NAADP-AM (**Figure 1D**) as did pharmacological antagonism of the NAADP receptor with Ned-19 [28] (**Figure 1D**). These data show that NAADP-mediated depolarization depends on canonical mechanisms of NAADP signalling.

We next examined whether an NAADP-mediated increase in intracellular  $\text{Ca}^{2+}$  concentration was required for the depolarization we had observed. By introducing the fast  $\text{Ca}^{2+}$  chelator BAPTA [1,2-bis (o-aminophenoxy) ethane-N,N,N',N'-tetraacetic acid] via the patch pipette (15 mM), we clamped intracellular  $\text{Ca}^{2+}$ , and depolarization by NAADP-AM was inhibited (**Figure 1D**). Whereas it was now clear that depolarization was dependent both upon  $\text{Ca}^{2+}$  and NAADP, the intracellular source of the  $\text{Ca}^{2+}$  required confirmation. NAADP reportedly causes  $\text{Ca}^{2+}$  release from acidic stores [29, 30]. Bafilomycin A1 is reported to abrogate endolysosomal  $\text{Ca}^{2+}$  release by inhibiting vacuolar  $\text{H}^{+}$ -ATPases and thereby preventing  $\text{Ca}^{2+}$  loading into the store [31, 32]. Pre-incubation with bafilomycin A1 abolished depolarization by NAADP-AM (**Figure 1D**), suggesting that NAADP-mediated membrane depolarization requires  $\text{Ca}^{2+}$  release from an acidic store.

Amplification of acidic store  $\text{Ca}^{2+}$  signals by  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release (CICR) from ryanodine receptors (RyR) has been reported in neurons of the medulla [15]; therefore, we examined whether this signalling motif was conserved in pyramidal neurons. Ryanodine inhibits RyRs in

the hippocampus at micromolar concentrations [33]. Bath application of ryanodine at 30  $\mu$ M inhibited NAADP-mediated membrane depolarization (**Figure 1D**), suggesting that the CICR amplification motif is conserved.

$\text{Ca}^{2+}$ -mobilizing second messengers are not generally associated with driving membrane depolarization; thus, we explored whether NAADP is unique in its ability to do so. We dialyzed the known  $\text{Ca}^{2+}$ -mobilizing second messengers [NAADP, inositol 1,4,5-trisphosphate ( $\text{IP}_3$ ), and cyclic adenosine diphosphate ribose (cADPR)] into CA1 pyramidal neurons via a patch pipette and measured membrane potential. We found that only when NAADP was included in the pipette was membrane depolarization observed (**Figure 2, A and B**). Critically, the magnitude of depolarization generated in response to dialysis with NAADP produced a “bell-shaped” distribution, a hallmark of NAADP signalling where supra-maximal concentrations of NAADP cause inhibition of the associated response [27, 34, 35]. The half-maximal effective concentration ( $\text{EC}_{50}$ ) was determined to be 12.5 nM, and the half maximal inhibitory concentration ( $\text{IC}_{50}$ ) was 7.07  $\mu$ M. The Hill coefficients were +0.527 for stimulation and -0.927 for inhibition. Furthermore, the maximal  $\Delta V_M$  achieved was at the concentration of 100 nM NAADP, which is within the range previously reported for maximum  $\text{Ca}^{2+}$  mobilisation from an acidic store [4, 36]. Neither  $\text{IP}_3$  nor cADPR, at any concentration, produced significant depolarization (**Figure 2B**). These results suggest that NAADP is unique in its capacity to produce a membrane depolarization amongst  $\text{Ca}^{2+}$ -mobilizing second messenger family.

### **Activation of mGluR1 causes NAADP-dependent membrane depolarization**

In order that we might begin to understand the significance of the NAADP-mediated depolarization, we sought to identify the mechanism by which NAADP elevations are

triggered physiologically. The excitatory neurotransmitter glutamate has previously been shown to generate the synthesis of NAADP in neurons and astrocytes (Pandey, et al. [25], Pereira, et al. [14]), although the specific glutamate receptor subtype(s) were not identified. In a variety of other tissue types, NAADP synthesis is reported to occur following activation of metabotropic receptors [20-23, 37], we therefore looked to metabotropic glutamate receptors as likely candidates. To examine this we pharmacologically isolated the mGluRs with antagonists of ionotropic glutamate and GABA receptors (50  $\mu$ M AP5, 10  $\mu$ M NBQX, 100  $\mu$ M picrotoxin and 2  $\mu$ M CGP 55845) and delivered patterned electrical stimulation (4x, 20 Hz), known to activate mGluRs in the hippocampus [38], whilst measuring the membrane potential from CA1 pyramidal neurons (**Figure 3A**). Under these conditions, we observe a depolarization  $\Delta V_m$  of  $+ 2.09 \pm 0.15$  mV (**Figure 3, B and C**). To confirm that this was a consequence of mGluR activation the mGluR antagonist LY341495 (100  $\mu$ M) was added to the bath, at a concentration reported to block all members of the mGluR family [39]. This blocked the electrically-induced membrane depolarization (**Figure 3, B and C**). Now confident that mGluRs are able to produce depolarization we sought to determine the specific mGluR subtype(s) that mediated the response.

The mGluR family contains 8 members, divided into three groups based on their pharmacological and functional profiles. Group I (mGluR1, mGluR5), Group II (mGluR2, mGluR3) and Group III (mGluR4, mGluR6, mGluR7, mGluR8) [40, 41]. We systematically isolated members of the mGluR family using a series of mGluR subgroup and/or subtype specific pharmacological antagonists. LY341495 acts as an mGluR Group II and III antagonist at 100 nM, whereas at 100  $\mu$ M it acts as a pan-mGluR antagonist [39]. The addition of 100 nM LY341495 produced no significant reduction in the depolarization observed upon electrical stimulation), whereas 100  $\mu$ M LY341495 blocked the membrane depolarization (**Figure 3C**).

Thus, the depolarization is a Group I mGluR-dependent effect. Group 1 contains two mGluR subtypes mGluR1 and mGluR5. MPEP, a selective mGluR5 antagonist [42], did not block the depolarization, whereas JNJ16259685, a selective mGluR1 antagonist [43], abolished the depolarization (**Figure 3C**). Based on the pharmacological dissection of the response we conclude that of the mGluR family of receptors, only mGluR1 activation produces membrane depolarization in CA1 pyramidal neurons.

We next sought to examine a link between the mGluR1-mediated membrane depolarization, NAADP signalling and acidic store  $\text{Ca}^{2+}$  release. We examined their relationship by isolating the mGluR response as described above prior to the introduction of further manipulations. Desensitization of the NAADP receptor with NAADP (1 mM) inside patch pipette abolished mGluR-mediated depolarization, whereas, 1 mM nicotinic acid adenosine dinucleotide (NAAD), an inactive metabolite, had no effect (**Figure 3D**), suggesting that the action of NAADP is specific. Pre-incubation with the NAADP receptor antagonist Ned-19 also significantly reduced responses to mGluR depolarization, and disruption of the lysosomes with acute application of glycyl-L-phenylalanine 2-naphthylamide (GPN) abolished mGluR-mediated depolarization (**Figure 3D**). GPN prevents acidic store  $\text{Ca}^{2+}$  signalling without compromising cell health [19]; thus, together, these data suggest mGluR membrane depolarization is dependent on NAADP and  $\text{Ca}^{2+}$  release from acidic stores.

### **mGluR1-dependent membrane depolarization does not require dendritic $\text{IP}_3\text{Rs}$**

Our data suggest that NAADP and  $\text{Ca}^{2+}$  release from acidic stores drives the membrane depolarization. It therefore seems reasonable that  $\text{Ca}^{2+}$  signalling via the acidic store is intimately linked to this process. Several studies report an amplification of acidic store  $\text{Ca}^{2+}$

by the ER and we show the RyRs are essential for NAADP-mediated depolarization (**Figure 1D**). However, we are also mindful that Group 1 mGluRs are thought to be involved in IP<sub>3</sub>R-mediated Ca<sup>2+</sup> release [41], therefore, we needed to explore this relationship in greater detail to better understand these Ca<sup>2+</sup> signalling events.

First, we confirmed the requirement of the acidic Ca<sup>2+</sup> stores for mGluR1-mediated depolarization by pharmacologically isolating mGluR1s (as previously described) and recording membrane potential whilst glutamate was bath applied (**Figure 4, A and B**). As expected, acute application of GPN abolished mGluR1-mediated depolarization (**Figure 4C**). We next wanted to determine whether the mGluR1-mediated depolarization was also dependent on Ca<sup>2+</sup> release from the ER. Therefore, we examined whether RyRs or IP<sub>3</sub>Rs are essential for mGluR1-mediated membrane depolarization. Acute application of desensitizing concentrations of ryanodine prevented mGluR1-mediated depolarization (**Figure 4C**). In contrast, Xestospongine C, a potent and selective inhibitor of IP<sub>3</sub>Rs [44], had no effect on the mGluR1-mediated depolarization (**Figure 4C**).

Membrane contact sites between the acidic Ca<sup>2+</sup> stores and the ER are reported and are suggested to mediate microdomain signalling between these two organelles [45]. To assess whether microdomain Ca<sup>2+</sup> signalling is likely to occur with mGluR1-mediated membrane depolarization we compared the action of two Ca<sup>2+</sup> chelators: ethylene glycol-bis (β-aminoethyl ether)-N, N, N', N'-tetraacetic acid (EGTA) and BAPTA. EGTA has an “on” rate for binding Ca<sup>2+</sup> of 3-10 μM<sup>-1</sup> s<sup>-1</sup>, whereas BAPTA is several orders of magnitude faster at 100–1000 μM<sup>-1</sup> s<sup>-1</sup> [46, 47]. Consequently, EGTA is thought not to chelate Ca<sup>2+</sup> at a rate fast enough to inhibit microdomain Ca<sup>2+</sup> signalling [48], whereas BAPTA can prevent all but elementary Ca<sup>2+</sup> signalling events [49]. Bath application of membrane-permeable EGTA (EGTA-AM) did



not affect mGluR1-mediated cellular depolarization (**Figure 4C**), whereas membrane permeable BAPTA (BAPTA-AM) prevented mGluR1-mediated depolarization (**Figure 4C**). These data suggest that mGluR1-mediated depolarization is dependent on microdomain  $\text{Ca}^{2+}$  signalling.

### **Dendritic mGluR1-dependent $\text{Ca}^{2+}$ transients require NAADP signalling and RyRs but not $\text{IP}_3\text{Rs}$**

To assess more directly the intracellular  $\text{Ca}^{2+}$  signalling events initiated by mGluR1 activation we used confocal microscopy to visualise  $\text{Ca}^{2+}$  signals in the dendrites of CA1 pyramidal neurons. Neurons were loaded with the  $\text{Ca}^{2+}$  sensitive dye Oregon Green 488 BAPTA-1 (OGB-1) and mGluR1s were pharmacologically isolated as described above.

Electrical stimulation was applied to generate dendritic  $\text{Ca}^{2+}$  signals. These  $\text{Ca}^{2+}$  signals could be blocked by pharmacologically inhibiting either the NAADP receptor with Ned-19 (**Figure 4, D to F**) or all members of the mGluR family with LY341495(**Figure 4, E and F**), again indicating that mGluR1-mediated  $\text{Ca}^{2+}$  signals depend on NAADP signalling. We found that ryanodine abolished mGluR1-dependent  $\text{Ca}^{2+}$  signals (**Figure 4, E and F**), whereas addition of the  $\text{IP}_3$  receptor antagonists Xestospongin C or 2-APB did not reduce mGluR1 mediated  $\text{Ca}^{2+}$  signalling (**Figure 4, E and F**). Together, these data suggest that mGluR1-mediated  $\text{Ca}^{2+}$  signals are dependent on  $\text{Ca}^{2+}$  release from both acidic  $\text{Ca}^{2+}$  stores and from the ER via RyR, but not  $\text{IP}_3\text{Rs}$ . As a positive control, we confirmed that Xestospongin C inhibited  $\text{IP}_3\text{R}$ -mediated  $\text{Ca}^{2+}$  release by visualising  $\text{Ca}^{2+}$  signals in CA1 pyramidal neurons patch clamped with an internal solution containing  $\text{IP}_3$  under control conditions and after pre-exposure to Xestospongin C (**fig. S1, A to C**).

## **mGluR1-dependent depolarization is mediated by the inactivation of SK channels following dephosphorylation**

We wished to understand the biophysical basis of the mGluR1-mediated membrane depolarization. A number of studies suggest that Group 1 mGluRs inhibit K<sup>+</sup> channels in CA1 pyramidal neurons of the hippocampus [42, 50-57] with Tigaret, et al. [58] providing evidence that mGluR1 activation inhibits SK channels, an important step for the induction of mGluR1-mediated long-term potentiation (LTP) in the hippocampus.

SK channels are present in the dendrites of CA1 pyramidal neurons of the hippocampus [59] where their activation produces action potential after-hyperpolarisation currents ( $I_{AHP}$ ). SK channels have also been implicated in regulating dendritic excitability [58, 60, 61].

We sought to determine whether the allosteric inhibitor of SK channels, apamin [62] inhibited mGluR1-mediated depolarization. mGluR1s were pharmacologically isolated (as described above) and membrane potential was recorded from CA1 pyramidal neurons while a patterned of electrical stimulation, was delivered. The experiment reveals that synaptic activation of mGluR1 produced membrane depolarization (**Figure 5, A and B**) and apamin significantly reduced the amplitude of depolarization (**Figure 5B**). GPN was then introduced to determine whether a common signalling pathway was being utilised. GPN was found to have no further effect on reducing the mGluR1-mediated depolarization (**Figure 5B**). These data suggest that inhibition of the SK channels by acidic store Ca<sup>2+</sup> is the key intermediate step in producing mGluR1-mediated depolarization.

To ensure that the addition of apamin and/or GPN had not interfered with presynaptic glutamate release we repeated this experiment with transient bath application of L-glutamate

rather than electrical stimulation. The mGluR1s were again pharmacologically isolated and neurotransmission prevented with TTX. Extracellular glutamate caused membrane depolarization (**Figure 5C**) and was significantly reduced by the addition of apamin (**Figure 5D**). Again, we found that application of GPN had no further effect on reducing the amount of mGluR1 depolarization after application of apamin (**Figure 5D**). Apamin did not affect the resting membrane potential of CA1 neurons of the hippocampus (**figure S2, A and B**).

SK channels are subject to modulation. Multiple sites for phosphorylation have been reported [63], with SK channels shown to form macromolecular complexes with protein kinase 2 (CK2) and protein phosphatase 2A (PP2A) [64, 65]. Some members of the PP2A family are  $\text{Ca}^{2+}$  sensitive [66] and the presence of  $\text{Ca}^{2+}$  binding EF-hand domains is noted on regulatory B-type subunits [67]. We therefore used the inhibitor of PP2A (okadaic acid) to determine whether SK channel modulation could affect mGluR1-mediated depolarization. We found that acute application of okadaic acid significantly reduced mGluR1-mediated depolarization (**Figure 5, E and F**). Therefore, we suggest that activation of mGluR1 and the subsequent  $\text{Ca}^{2+}$  signals evoked are key in activating one or more members of the PP2A family.

Finally, it has also been suggested that mGluR1-mediated depolarization may occur via activation of a non-selective cation or TRP channel [68, 69]. To assess this possibility, we used the broad-spectrum antagonists of non-selective cation channels, Flufenamic acid and  $\text{La}^{3+}$ , to determine whether mGluR1-mediated depolarization could be achieved via these channels. We found that neither had any effect on mGluR1-mediated depolarization (**figure S3, A-C**).

**mGluR1-dependent plasticity requires inhibition of SK channels by acidic store  $\text{Ca}^{2+}$**

Several studies suggest that hippocampal long-term potentiation can occur following the activation of mGluR1 [58, 70-73]. We wished to determine whether the signalling pathway we describe is required for mGluR1-dependent plasticity. We therefore implemented a spike timing dependent plasticity (STDP) protocol known to produce mGluR1-dependent plasticity [58].

The STDP induction protocol produced LTP of around 150% (**Figure 6, A**) in CA1 pyramidal neurons. To ensure the protocol induced mGluR1-dependent LTP we repeated the experiment in the presence of an mGluR1 specific antagonist (JNJ16259685), which blocked the LTP (**Figure 6, B**).

Next, we selectively abolished NAADP signalling with a desensitizing concentration of NAADP [74, 75] in the patch pipette. We found this manipulation prevented LTP (**Figure 6, C**), indicating that mGluR1-dependent LTP requires NAADP-acidic store  $\text{Ca}^{2+}$  signalling.

As the inhibition of SK channels via mGluR1 is thought to be required for mGluR1-mediated LTP, pharmacological inactivation of SK channels with apamin should have no impact on the LTP; SK channels should be inhibited by mGluR1 activation, thereby occluding the apamin's action. Indeed, we found that apamin alone had no effect on the magnitude of LTP compared to control experiments (**Figure 6, D**). Critically, we also found that antagonism of mGluR1s with JNJ16259685 while simultaneously inhibiting SK channels with apamin rescues the ability of the STDP protocol to produce mGluR1-dependent plasticity (**Figure 6, E**). These findings (summarized **Figure 6F**) indicates that activation of mGluR1 can produce LTP in the hippocampus and that LTP is achieved via the modulation of SK channel activity.

**Two-pore channels are essential for mGluR1-dependent membrane depolarization and LTP**

We have shown that NAADP signalling is required for mGluR1-mediated depolarization and LTP. Here we add to the pharmacological evidence by genetically manipulating the signalling pathway. Two-pore channels (TPC1 and TPC2) are localized to acidic  $\text{Ca}^{2+}$  stores in mammalian cells and have been shown to be essential for NAADP-mediated  $\text{Ca}^{2+}$  release [1-3, 35]. First, we wanted to determine if TPCs were required for NAADP-mediated events in our experimental preparation. Therefore, we dialyzed NAADP into CA1 pyramidal neurons, via a patch pipette at the maximally effective concentration (determined in data shown in **Figure 2B**), whilst recording membrane potential. NAADP-mediated depolarization was reduced in neurons from *Tpc1*<sup>-/-</sup> and abolished in neurons from *Tpc2*<sup>-/-</sup> (**figure S4, A and B**).

Next, we explored if either *Tpc1* and *Tpc2* was required for mGluR1-mediated depolarization and mGluR1-dependent LTP. To achieve this, we first assessed whether mGluR1-dependent depolarization could be produced in CA1 neurons of wild-type (WT) mice. We pharmacologically isolated mGluR1s (as described above) and delivered electrical stimulation confirming that mGluR1 activation produced depolarization (**Figure 7, A and B**). In contrast, depolarization was not observed in CA1 pyramidal neurons of either *Tpc1*<sup>-/-</sup> or *Tpc2*<sup>-/-</sup> animals.

Next, we sought to determine whether the TPCs were important for mGluR1-dependent LTP. We confirmed that was is observed in CA1 pyramidal neurons (**Figure 7, C and D**). Upon assessing LTP in knock-out mice, we were surprised to find that the STDP protocol that had generated LTP instead induced LTD in both *Tpc1*<sup>-/-</sup> and *Tpc2*<sup>-/-</sup> animals (**Figure 7, C and D**). It would therefore appear that the change in the  $\text{Ca}^{2+}$  signalling profile in these neurons resulted in a switch in the polarity of the plasticity for our STDP stimulus regime.

## Discussion:

In this work, we show that the metabotropic receptor, mGluR1, is specifically linked to the NAADP signalling cascade, a pathway that uniquely mobilises the release of  $\text{Ca}^{2+}$  from acidic organelles such as lysosomes and late endosomes. We found that activation of mGluR1 produced NAADP-dependent  $\text{Ca}^{2+}$  release and also membrane depolarization, both of which requiring  $\text{Ca}^{2+}$  amplification via the RyRs. As we found cADPR alone was not able to produce depolarization we suggest that mGluR1/NAADP-mediated  $\text{Ca}^{2+}$  release from the acidic stores is the critical first step in this signalling pathway. Importantly, the pathway is quite specific, as we find that mGluR5, the second member of Group 1 mGluR group, fails to trigger NAADP-mediated cellular depolarization. We reveal that this pathway is intimately linked to the induction of mGluR1-mediated LTP. The mechanism by which this occurs appears to be a transient modulation of SK channel function following dephosphorylation by PP2A.

In a previous study, we found that acidic stores were essential for the maintenance of spine growth following LTP induction [19]. We now find that in addition to LTP maintenance, acidic stores play a role in the induction of mGluR1-dependent forms of LTP. Whether LTP requires mGluR1; however, depends on the parameters of stimulation, with stronger stimulation protocols capable of bypassing the requirement of mGluRs and, more generally, store  $\text{Ca}^{2+}$  release [76-78].

Perhaps the least intuitive of our results is the observation that mGluR1 activation modulates SK channel function to produce *depolarization* of the membrane. Canonically, SK channels are thought to underlie after-hyperpolarising potentials (AHPs). In this context, the channels are closed and become active as  $\text{Ca}^{2+}$  enters neurons following activity. However, it would appear that the regulation of SK channels by  $\text{Ca}^{2+}$  is more complex than once thought, with SK

channels differentially regulated by  $\text{Ca}^{2+}$  from different sources.  $\text{Ca}^{2+}$  that enters via R-Type voltage-dependent  $\text{Ca}^{2+}$  channels (VDCCs) or N-methyl-D-aspartate receptors (NMDARs) acts to potentiate SK channel opening causing hyperpolarization. In this context, hyperpolarisation helps reinstate the  $\text{Mg}^{2+}$  block of NMDARs, reducing the amount of  $\text{Ca}^{2+}$  entry and consequently reducing the magnitude of LTP [79, 80]. Conversely, we find that  $\text{Ca}^{2+}$  release from acidic stores reduces SK channel activity. We do not fully describe the mechanism as to how this occurs; however, we show that PP2A is required to produce mGluR1-mediated depolarization. PP2A form complexes with SK channels and are only active whilst the channels are in an open state; the result of their activity is to increase the sensitivity of SK channels to  $\text{Ca}^{2+}$  [64, 65]. Perhaps this increased sensitivity allows detection of acidic store  $\text{Ca}^{2+}$  and in turn reduce SK channel activity. Further work needs to be undertaken to elucidate this signalling mechanism. Additionally, we found apamin had no effect on resting membrane potential of hippocampal neurons. This is consistent with previous findings [81-84] and might be explained by apamin's allosteric mode of action [62], which perhaps locks SK channels in their current state and thus prevents them from being activated by any source of  $\text{Ca}^{2+}$ .

Nevertheless, the reduction in SK channel activity is a key step in the induction of metabotropic receptor-dependent forms of LTP [58, 85]. In good agreement with *Tigaret et al.* [58] we confirm that inhibition of SK channels is essential for mGluR1-mediated LTP, but critically we identify the intermediate step and link mGluR1-mediated depolarization to the NAADP signalling pathway. Underpinning our link to this pathway is the data showing that genetic knock out of either *Tpc1* or *Tpc2* removes both mGluR1-mediated depolarization and mGluR1-mediated LTP. The importance of both *Tpc1* and *Tpc2* for mGluR1-mediated LTP may indicate that mGluR1 is interacting with heteromeric complexes of *Tpc1* and *Tpc2*, an arrangement reported in other cell types [86].

Roles for mGluR1 in both long-term depression (LTD) [38, 72, 87-89] and long-term potentiation (LTP) [70, 90-92] have been reported at CA3-CA1 synapses. Whether LTD or LTP is observed seems to be dependent on the stimulation pattern delivered to the neurons, with stronger stimulation regimes generating LTP and weaker ones LTD [72, 76, 93]. Interestingly, mGluR1 can produce both hyperpolarization and depolarization in dopaminergic neurons, with strong stimulation causing depolarization and weak stimulation hyperpolarisation [94]. One clear illustration of the significance of recruiting specific signalling pathways to generate a specific plasticity outcome was seen in our data that showed that when TPCs are genetically removed our LTP induction protocol induced LTD not LTP. Collectively, these data suggest that at least two intracellular signalling pathways can be stimulated by mGluR1 activation and that the pathway engaged, and the consequent polarity of plasticity, is dependent on the stimulation pattern.

The importance of understanding the different mGluR1 signalling pathways is clear when interpreting work in vivo. mGluR1 antagonists impair spatial learning [95] and the deletion of the gene encoding mGluR1 reduces LTP in CA1 region of the hippocampus and impairs context-specific associative learning [96] and spatial memory [97]. Impaired Group 1 mGluR signalling is also implicated in the pathogenesis of Fragile X Syndrome [98, 99] as yet an untreatable disorder, thus the signalling pathway we reveal offers new opportunities towards understanding diverse behavioural phenotypes and routes for novel therapeutic intervention.

## **Materials and methods:**

### *Hippocampal slice preparation*

All animal work was carried out in accordance with the Animals (Scientific Procedures) Act, 1986 (UK) and under the project and personal licenses approved by the Home Office (UK).



Slice cultures of the hippocampus were prepared from male Wistar rats (P6-8). The hippocampi were isolated in ice-cold Earle's balanced salt solution (EBSS) with added: HEPES (21 mM), D-(+)-Glucose (27.8 mM) pH adjusted to 7.2 - 7.4 with NaOH and cut into slices of 350  $\mu$ m thickness with a McIlwain tissue chopper. Slices were placed into Millicell CM culture plate inserts (PTFE filter, pore size 0.4  $\mu$ m, diam. 12 mm) in a six-well Millicell culture plate (both supplied by Merck Millipore) with 1 mL of culture medium and stored at 34.5°C at 5% CO<sub>2</sub>. Culture medium composed of 78.8% minimum essential medium (MEM) with GlutaMAX (Gibco), 20% heat-inactivated horse serum, 1% B27 with added CaCl<sub>2</sub> (1 mM), HEPES (30 mM,) D-(+)-Glucose (26 mM,) NaHCO<sub>3</sub> (5.8 mM) and MgSO<sub>4</sub> (2 mM). Culture media was renewed every 3-4 days.

During experiments, slices (10–14 days in vitro [DIV]) were perfused (1–2 mL/min) with heated (32°C–34°C) artificial cerebrospinal fluid (ACSF) which comprised of NaCl (145 mM), KCl (2.5 mM), KH<sub>2</sub>PO<sub>4</sub> (1.2 mM), NaHCO<sub>3</sub> (16.0 mM), glucose (11.0 mM), CaCl<sub>2</sub> (3.0 mM) and MgCl<sub>2</sub> (2.0 mM) aerated with 95% O<sub>2</sub>, 5% CO<sub>2</sub>.

### *Electrophysiology*

Whole-cell patch clamp recordings were performed on pyramidal neurons from either CA3 or CA1 neurons of hippocampal slice cultures. To minimize intracellular dialysis high-resistance patch electrodes (16–20 M $\Omega$ ) were used where mGluR1 was isolated pharmacologically and all other experiments were undertaken with lower resistance electrodes (5–8 M $\Omega$ ). Voltage signals were detected using either an Axoclamp 2B (Axon Instruments/Molecular devices) amplifier, signals were digitised using a Digidata 1440A then recorded digitally with WinWCP V4.7.9 (Strathclyde Electrophysiology Software), 50 Hz noise was eliminated with a Hum Bug (Quest Scientific). The internal solution contained (135 mM KGluconate, 10 mM KCl, 10 mM

HEPES, 2 mM MgCl<sub>2</sub>, 2 mM Na<sub>2</sub>ATP, and 0.4 mM Na<sub>3</sub>GTP; pH = 7.2–7.4). Pharmacological agents/second messengers added to the internal solution were conjugated to K<sup>+</sup> salts, and when present, the KGluconate concentration was reduced by equal molarity in order to maintain osmolarity. Electrical stimulation was applied using a tungsten stimulating electrode with an isolated constant current stimulator (Digitimer Ltd). To produce mGluR1-mediated depolarization the stimulation pattern used consisted of four pulses at 20 Hz. The mGluR1 LTP protocol consisted of causally pairing one pre-synaptic stimulus (to produce a subthreshold excitatory postsynaptic potential (EPSP)) with two backpropagating action potentials (bAPs) (100 Hz), elicited in the postsynaptic neuron via current injection, at a 10 ms interval. This paired induction protocol was repeated 300 times at 5 Hz and delivered within 5 minutes of whole cell breakthrough to prevent dialysis of factors required for mGluR1-dependent LTP.

#### *Ca<sup>2+</sup> imaging*

Pyramidal neurons in the CA1 region of hippocampal slice cultures, 10-14 D.I.V. were filled with Oregon BAPTA Green 488 (OGB-1) via a patch clamp in whole cell configuration for 1 minute. The patch electrode's internal solution consisted of: 135 mM K-gluconate, 10 mM KCL, 10 mM HEPES, 1 mM MgCl<sub>2</sub>, 1 mM OGB-1, 10 mM QX314. Apical dendrites were imaged using confocal laser-scanning microscopy and a 488-nm argon laser whilst an isolated constant current stimulator was used to deliver electrical stimulation (4 pulses, 20 Hz) presynaptic neurons.

#### *Reagents*

NAADP-AM & NAADP were synthesised in house [26, 100]). Other drugs were purchased from the following suppliers: Abcam: LY341495, CGP55845, D-AP5, Ned-19, JNJ16259685, MPEP, QX314, TTX. Sigma Aldrich: Ryanodine, bafilomycin A1, picrotoxin, NBQX, nicotinic acid

adenine dinucleotide (NAAD) & BAPTA, BAPTA-AM, EGTA-AM. Santa Cruz Biotechnology: GPN. Fisher Scientific: OBG-1. Tocris: U73122, U73343.

### Supplementary materials

Figure S1: Xestospongine C inhibits somatic IP<sub>3</sub> mediated Ca<sup>2+</sup> release in CA1 pyramidal neurons in the hippocampus.

Figure S2: Apamin does not affect resting membrane potential of CA1 pyramidal neurons.

Figure S3: mGluR1-mediated depolarization unlikely to occur via transient receptor potential (TRP) channel activation.

Figure S4: NAADP-mediated membrane depolarization in CA1 pyramidal neurons requires *Tpc1* and *Tpc2*.

### References and notes:

1. Brailoiu, E., Churamani, D., Cai, X., Schrlau, M.G., Brailoiu, G.C., Gao, X., Hooper, R., Boulware, M.J., Dun, N.J., Marchant, J.S., et al. (2009). Essential requirement for two-pore channel 1 in NAADP-mediated calcium signaling. *J Cell Biol* 186, 201-209.
2. Calcraft, P.J., Ruas, M., Pan, Z., Cheng, X., Arredouani, A., Hao, X., Tang, J., Rietdorf, K., Teboul, L., Chuang, K.T., et al. (2009). NAADP mobilizes calcium from acidic organelles through two-pore channels. *Nature* 459, 596-600.
3. Zong, X., Schieder, M., Cuny, H., Fenske, S., Gruner, C., Rotzer, K., Griesbeck, O., Harz, H., Biel, M., and Wahl-Schott, C. (2009). The two-pore channel TPCN2 mediates NAADP-dependent Ca<sup>2+</sup>-release from lysosomal stores. *Pflugers Arch* 458, 891-899.
4. Galione, A. (2011). NAADP receptors. *Cold Spring Harb Perspect Biol* 3, a004036.
5. Morgan, A.J., Platt, F.M., Lloyd-Evans, E., and Galione, A. (2011). Molecular mechanisms of endolysosomal Ca<sup>2+</sup> signalling in health and disease. *Biochem J* 439, 349-374.
6. Menteyne, A., Burdakov, A., Charpentier, G., Petersen, O.H., and Cancela, J.-M. (2006). Generation of Specific Ca<sup>2+</sup> Signals from Ca<sup>2+</sup> Stores and Endocytosis by Differential Coupling to Messengers. *Current Biology* 16, 1931-1937.
7. Bak, J., White, P., Timár, G., Missiaen, L., Genazzani, A.A., and Galione, A. (1999). Nicotinic acid adenine dinucleotide phosphate triggers Ca<sup>2+</sup> release from brain microsomes. *Current Biology* 9, 751-754.
8. Patel, S., Churchill, G.C., Sharp, T., and Galione, A. (2000). Widespread distribution of binding sites for the novel Ca<sup>2+</sup>-mobilizing messenger, nicotinic acid adenine dinucleotide phosphate, in the brain. *J Biol Chem* 275, 36495-36497.
9. Brailoiu, E., Churamani, D., Pandey, V., Brailoiu, G.C., Tuluc, F., Patel, S., and Dun, N.J. (2006). Messenger-specific role for nicotinic acid adenine dinucleotide phosphate in neuronal differentiation. *J Biol Chem* 281, 15923-15928.
10. Zhang, Z.H., Lu, Y.Y., and Yue, J. (2013). Two pore channel 2 differentially modulates neural differentiation of mouse embryonic stem cells. *PLoS One* 8, e66077.

11. Hao, B., Webb, S.E., Miller, A.L., and Yue, J. (2016). The role of Ca<sup>2+</sup> signaling on the self-renewal and neural differentiation of embryonic stem cells (ESCs). *Cell Calcium* 59, 67-74.
12. Brailoiu, E., Hoard, J.L., Filipeanu, C.M., Brailoiu, G.C., Dun, S.L., Patel, S., and Dun, N.J. (2005). Nicotinic acid adenine dinucleotide phosphate potentiates neurite outgrowth. *J Biol Chem* 280, 5646-5650.
13. Pereira, G.J., Hirata, H., Fimia, G.M., do Carmo, L.G., Bincoletto, C., Han, S.W., Stilhano, R.S., Ureshino, R.P., Bloor-Young, D., Churchill, G., et al. (2011). Nicotinic acid adenine dinucleotide phosphate (NAADP) regulates autophagy in cultured astrocytes. *J Biol Chem* 286, 27875-27881.
14. Pereira, G.J., Antonioli, M., Hirata, H., Ureshino, R.P., Nascimento, A.R., Bincoletto, C., Vescovo, T., Piacentini, M., Fimia, G.M., and Smaili, S.S. (2016). Glutamate induces autophagy via the two-pore channels in neural cells. *Oncotarget* 5.
15. Brailoiu, G.C., Brailoiu, E., Parkesh, R., Galione, A., Churchill, G.C., Patel, S., and Dun, N.J. (2009). NAADP-mediated channel 'chatter' in neurons of the rat medulla oblongata. *Biochem J* 419, 91-97, 92 p following 97.
16. Chameau, P., Van de Vrede, Y., Fossier, P., and Baux, G. (2001). Ryanodine-, IP3- and NAADP-dependent calcium stores control acetylcholine release. *Pflugers Arch* 443, 289-296.
17. Brailoiu, E., Miyamoto, M.D., and Dun, N.J. (2001). Nicotinic acid adenine dinucleotide phosphate enhances quantal neurosecretion at the frog neuromuscular junction: Possible action on synaptic vesicles in the releasable pool. *Molecular Pharmacology* 60, 718-724.
18. McGuinness, L., Bardo, S.J., and Emptage, N.J. (2007). The lysosome or lysosome-related organelle may serve as a Ca<sup>2+</sup> store in the boutons of hippocampal pyramidal cells. *Neuropharmacology* 52, 126-135.
19. Padamsey, Z., McGuinness, L., Bardo, S.J., Reinhart, M., Tong, R., Hedegaard, A., Hart, M.L., and Emptage, N.J. (2017). Activity-Dependent Exocytosis of Lysosomes Regulates the Structural Plasticity of Dendritic Spines. *Neuron* 93, 132-146.
20. Soares, S., Thompson, M., White, T., Isbell, A., Yamasaki, M., Prakash, Y., Lund, F.E., Galione, A., and Chini, E.N. (2007). NAADP as a second messenger: neither CD38 nor base-exchange reaction are necessary for in vivo generation of NAADP in myometrial cells. *Am J Physiol Cell Physiol* 292, C227-239.
21. Arredouani, A., Evans, A.M., Ma, J., Parrington, J., Zhu, M.X., and Galione, A. (2010). An emerging role for NAADP-mediated Ca<sup>2+</sup> signaling in the pancreatic beta-cell. *Islets* 2, 323-330.
22. Rah, S.Y., Mushtaq, M., Nam, T.S., Kim, S.H., and Kim, U.H. (2010). Generation of cyclic ADP-ribose and nicotinic acid adenine dinucleotide phosphate by CD38 for Ca<sup>2+</sup> signaling in interleukin-8-treated lymphokine-activated killer cells. *J Biol Chem* 285, 21877-21887.
23. Mushtaq, M., Nam, T.S., and Kim, U.H. (2011). Critical role for CD38-mediated Ca<sup>2+</sup> signaling in thrombin-induced procoagulant activity of mouse platelets and hemostasis. *J Biol Chem* 286, 12952-12958.
24. Cosker, F., Cheviron, N., Yamasaki, M., Menteyne, A., Lund, F.E., Moutin, M.J., Galione, A., and Cancela, J.M. (2010). The ecto-enzyme CD38 is a nicotinic acid adenine dinucleotide phosphate (NAADP) synthase that couples receptor activation to Ca<sup>2+</sup> mobilization from lysosomes in pancreatic acinar cells. *J Biol Chem* 285, 38251-38259.
25. Pandey, V., Chuang, C.C., Lewis, A.M., Aley, P.K., Brailoiu, E., Dun, N.J., Churchill, G.C., and Patel, S. (2009). Recruitment of NAADP-sensitive acidic Ca<sup>2+</sup> stores by glutamate. *Biochem J* 422, 503-512.
26. Parkesh, R., Lewis, A.M., Aley, P.K., Arredouani, A., Rossi, S., Tavares, R., Vasudevan, S.R., Rosen, D., Galione, A., Dowden, J., et al. (2008). Cell-permeant NAADP: a novel chemical tool enabling the study of Ca<sup>2+</sup> signalling in intact cells. *Cell Calcium* 43, 531-538.
27. Cancela, J.M., Churchill, G.C., and Galione, A. (1999). Coordination of agonist-induced Ca<sup>2+</sup>-signalling patterns by NAADP in pancreatic acinar cells. *Nature* 398, 74-76.

28. Naylor, E., Arredouani, A., Vasudevan, S.R., Lewis, A.M., Parkesh, R., Mizote, A., Rosen, D., Thomas, J.M., Izumi, M., Ganesan, A., et al. (2009). Identification of a chemical probe for NAADP by virtual screening. *Nat Chem Biol* 5, 220-226.
29. Churchill, G.C., Okada, Y., Thomas, J.M., Genazzani, A.A., Patel, S., and Galione, A. (2002). NAADP Mobilizes Ca<sup>2+</sup> from Reserve Granules, Lysosome-Related Organelles, in Sea Urchin Eggs. *Cell* 111, 703-708.
30. Yamasaki, M., Masgrau, R., Morgan, A.J., Churchill, G.C., Patel, S., Ashcroft, S.J., and Galione, A. (2004). Organelle selection determines agonist-specific Ca<sup>2+</sup> signals in pancreatic acinar and beta cells. *J Biol Chem* 279, 7234-7240.
31. Yoshimori, T., Yamamoto, A., Moriyama, Y., Futai, M., and Tashiro, Y. (1991). Bafilomycin A1, a specific inhibitor of vacuolar-type H(+)-ATPase, inhibits acidification and protein degradation in lysosomes of cultured cells. *J Biol Chem* 266, 17707-17712.
32. Christensen, K.A., Myers, J.T., and Swanson, J.A. (2002). pH-dependent regulation of lysosomal calcium in macrophages. *J Cell Sci* 115, 599-607.
33. Emptage, N., Bliss, T.V.P., and Fine, A. (1999). Single Synaptic Events Evoke NMDA Receptor-Mediated Release of Calcium from Internal Stores in Hippocampal Dendritic Spines. *Neuron* 22, 115-124.
34. Davis, L.C., Morgan, A.J., Chen, J.L., Snead, C.M., Bloor-Young, D., Shenderov, E., Stanton-Humphreys, M.N., Conway, S.J., Churchill, G.C., Parrington, J., et al. (2012). NAADP activates two-pore channels on T cell cytolytic granules to stimulate exocytosis and killing. *Curr Biol* 22, 2331-2337.
35. Ruas, M., Davis, L.C., Chen, C.C., Morgan, A.J., Chuang, K.T., Walseth, T.F., Grimm, C., Garnham, C., Powell, T., Platt, N., et al. (2015). Expression of Ca(2+)-permeable two-pore channels rescues NAADP signalling in TPC-deficient cells. *EMBO J* 34, 1743-1758.
36. Guse, A.H., Ernst, I.M., and Fliegert, R. (2013). NAADP signaling revisited. *Curr Top Med Chem* 13, 2978-2990.
37. Menteyne, A., Burdakov, A., Charpentier, G., Petersen, O.H., and Cancela, J.-M. (2006). Generation of Specific Ca<sup>2+</sup> Signals from Ca<sup>2+</sup> Stores and Endocytosis by Differential Coupling to Messengers. Volume 16. pp. 1931-1937.
38. Fan, W., Ster, J., and Gerber, U. (2010). Activation conditions for the induction of metabotropic glutamate receptor-dependent long-term depression in hippocampal CA1 pyramidal cells. *J Neurosci* 30, 1471-1475.
39. Kingston, A.E., Ornstein, P.L., Wright, R.A., Johnson, B.G., Mayne, N.G., Burnett, J.P., Belagaje, R., Wu, S., and Schoepp, D.D. (1998). LY341495 is a nanomolar potent and selective antagonist of group II metabotropic glutamate receptors. *Neuropharmacology* 37, 1-12.
40. Ferraguti, F., and Shigemoto, R. (2006). Metabotropic glutamate receptors. *Cell Tissue Res* 326, 483-504.
41. Niswender, C.M., and Conn, P.J. (2010). Metabotropic Glutamate Receptors: Physiology, Pharmacology, and Disease. *Annual review of pharmacology and toxicology* 50, 295-322.
42. Mannaioni, G., Marino, M.J., Valenti, O., Traynelis, S.F., and Conn, P.J. (2001). Metabotropic glutamate receptors 1 and 5 differentially regulate CA1 pyramidal cell function. *J Neurosci* 21, 5925-5934.
43. Lavreysen, H., Wouters, R., Bischoff, F., Nobrega Pereira, S., Langlois, X., Blokland, S., Somers, M., Dillen, L., and Lesage, A.S. (2004). JNJ16259685, a highly potent, selective and systemically active mGlu1 receptor antagonist. *Neuropharmacology* 47, 961-972.
44. Gafni, J., Munsch, J.A., Lam, T.H., Catlin, M.C., Costa, L.G., Molinski, T.F., and Pessah, I.N. (1997). Xestospongins: potent membrane permeable blockers of the inositol 1,4,5-trisphosphate receptor. *Neuron* 19, 723-733.
45. Levine, T.P., and Patel, S. (2016). Signalling at membrane contact sites: two membranes come together to handle second messengers. *Current opinion in cell biology* 39, 77-83.

46. Tsien, R.Y. (1980). New calcium indicators and buffers with high selectivity against magnesium and protons: design, synthesis, and properties of prototype structures. *Biochemistry* 19, 2396-2404.
47. Naraghi, M., and Neher, E. (1997). Linearized Buffered  $\text{Ca}^{2+}$  Diffusion in Microdomains and Its Implications for Calculation of  $[\text{Ca}^{2+}]$  at the Mouth of a Calcium Channel. *The Journal of Neuroscience* 17, 6961.
48. Dargan, S.L., and Parker, I. (2003). Buffer kinetics shape the spatiotemporal patterns of IP(3)-evoked  $\text{Ca}^{2+}$  signals. *The Journal of Physiology* 553, 775-788.
49. Horne, J.H., and Meyer, T. (1997). Elementary calcium-release units induced by inositol trisphosphate. *Science* 276, 1690-1693.
50. Charpak, S., Gahwiler, B.H., Do, K.Q., and Knopfel, T. (1990). Potassium conductances in hippocampal neurons blocked by excitatory amino-acid transmitters. *Nature* 347, 765-767.
51. Desai, M.A., and Conn, P.J. (1991). Excitatory effects of ACPD receptor activation in the hippocampus are mediated by direct effects on pyramidal cells and blockade of synaptic inhibition. *J Neurophysiol* 66, 40-52.
52. Guerinéau, N.C., Gahwiler, B.H., and Gerber, U. (1994). Reduction of Resting  $\text{K}^{+}$  Current by Metabotropic Glutamate and Muscarinic Receptors in Rat CA3 Cells - Mediation by G-Proteins. *J Physiol-London* 474, 27-33.
53. Luthi, A., Gahwiler, B.H., and Gerber, U. (1996). A slowly inactivating potassium current in CA3 pyramidal cells of rat hippocampus in vitro. *J Neurosci* 16, 586-594.
54. Takeshita, Y., Harata, N., and Akaike, N. (1996). Suppression of  $\text{K}^{+}$  conductance by metabotropic glutamate receptor in acutely dissociated large cholinergic neurons of rat caudate putamen. *J Neurophysiol* 76, 1545-1558.
55. Wu, R.L., and Barish, M.E. (1999). Modulation of a slowly inactivating potassium current,  $\text{I}(\text{D})$ , by metabotropic glutamate receptor activation in cultured hippocampal pyramidal neurons. *J Neurosci* 19, 6825-6837.
56. Chuang, S.C., Bianchi, R., and Wong, R.K. (2000). Group I mGluR activation turns on a voltage-gated inward current in hippocampal pyramidal cells. *J Neurophysiol* 83, 2844-2853.
57. Ireland, D.R., and Abraham, W.C. (2002). Group I mGluRs increase excitability of hippocampal CA1 pyramidal neurons by a PLC-independent mechanism. *J Neurophysiol* 88, 107-116.
58. Tigaret, C.M., Olivo, V., Sadowski, J.H., Ashby, M.C., and Mellor, J.R. (2016). Coordinated activation of distinct  $\text{Ca}^{2+}$  sources and metabotropic glutamate receptors encodes Hebbian synaptic plasticity. *Nat Commun* 7, 10289.
59. Allen, D., Bond, C.T., Lujan, R., Ballesteros-Merino, C., Lin, M.T., Wang, K., Klett, N., Watanabe, M., Shigemoto, R., Stackman, R.W., Jr., et al. (2011). The SK2-long isoform directs synaptic localization and function of SK2-containing channels. *Nat Neurosci* 14, 744-749.
60. Faber, E.S. (2009). Functions and modulation of neuronal SK channels. *Cell Biochem Biophys* 55, 127-139.
61. Faber, E.S., and Sah, P. (2007). Functions of SK channels in central neurons. *Clin Exp Pharmacol Physiol* 34, 1077-1083.
62. Lamy, C., Goodchild, S.J., Weatherall, K.L., Jane, D.E., Liégeois, J.-F., Seutin, V., and Marrion, N.V. (2010). Allosteric Block of  $\text{KCa}_2$  Channels by Apamin. *Journal of Biological Chemistry* 285, 27067-27077.
63. Köhler, M., Hirschberg, B., Bond, C.T., Kinzie, J.M., Marrion, N.V., Maylie, J., and Adelman, J.P. (1996). Small-Conductance, Calcium-Activated Potassium Channels from Mammalian Brain. *Science* 273, 1709-1714.
64. Bildl, W., Strassmaier, T., Thurm, H., Andersen, J., Eble, S., Oliver, D., Knipper, M., Mann, M., Schulte, U., Adelman, J.P., et al. (2004). Protein kinase CK2 is coassembled with small conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^{+}$  channels and regulates channel gating. *Neuron* 43, 847-858.

65. Allen, D., Fakler, B., Maylie, J., and Adelman, J.P. (2007). Organization and regulation of small conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^{+}$  channel multiprotein complexes. *J Neurosci* 27, 2369-2376.
66. Wlodarchak, N., Guo, F., Satyshur, K.A., Jiang, L., Jeffrey, P.D., Sun, T., Stanevich, V., Mumby, M.C., and Xing, Y. (2013). Structure of the  $\text{Ca}^{2+}$ -dependent PP2A heterotrimer and insights into Cdc6 dephosphorylation. *Cell Res* 23, 931-946.
67. Janssens, V., Jordens, J., Stevens, I., Van Hoof, C., Martens, E., De Smedt, H., Engelborghs, Y., Waelkens, E., and Goris, J. (2003). Identification and functional analysis of two  $\text{Ca}^{2+}$ -binding EF-hand motifs in the B"/PR72 subunit of protein phosphatase 2A. *J Biol Chem* 278, 10697-10706.
68. Congar, P., Leinekugel, X., BenAri, Y., and Crepel, V. (1997). A long-lasting calcium-activated nonselective cationic current is generated by synaptic stimulation or exogenous activation of group I metabotropic glutamate receptors in CA1 pyramidal neurons. *Journal of Neuroscience* 17, 5366-5379.
69. Gee, C.E., Benquet, P., and Gerber, U. (2003). Group I metabotropic glutamate receptors activate a calcium-sensitive transient receptor potential-like conductance in rat hippocampus. *J Physiol* 546, 655-664.
70. Lapointe, V., Morin, F., Ratte, S., Croce, A., Conquet, F., and Lacaille, J.C. (2004). Synapse-specific mGluR1-dependent long-term potentiation in interneurons regulates mouse hippocampal inhibition. *J Physiol* 555, 125-135.
71. Volk, L.J., Daly, C.A., and Huber, K.M. (2006). Differential roles for group 1 mGluR subtypes in induction and expression of chemically induced hippocampal long-term depression. *J Neurophysiol* 95, 2427-2438.
72. Neyman, S., and Manahan-Vaughan, D. (2008). Metabotropic glutamate receptor 1 (mGluR1) and 5 (mGluR5) regulate late phases of LTP and LTD in the hippocampal CA1 region in vitro. *Eur J Neurosci* 27, 1345-1352.
73. Fan, W. (2013). Group I metabotropic glutamate receptors modulate late phase long-term potentiation in hippocampal CA1 pyramidal neurons: comparison of apical and basal dendrites. *Neurosci Lett* 553, 132-137.
74. Aarhus, R., Dickey, D.M., Graeff, R.M., Gee, K.R., Walseth, T.F., and Lee, H.C. (1996). Activation and inactivation of  $\text{Ca}^{2+}$  release by NAADP<sup>+</sup>. *J Biol Chem* 271, 8513-8516.
75. Genazzani, A.A., Empson, R.M., and Galione, A. (1996). Unique inactivation properties of NAADP-sensitive  $\text{Ca}^{2+}$  release. *J Biol Chem* 271, 11599-11602.
76. Wilsch, V.W., Behnisch, T., Jager, T., Reymann, K.G., and Balschun, D. (1998). When are class I metabotropic glutamate receptors necessary for long-term potentiation? *J Neurosci* 18, 6071-6080.
77. Padamsey, Z., Tong, R., and Emptage, N. (2017). Glutamate is required for depression but not potentiation of long-term presynaptic function. *eLife* 6, e29688.
78. Padamsey, Z., Foster, W.J., and Emptage, N.J. (2018). Intracellular  $\text{Ca}^{2+}$  Release and Synaptic Plasticity: A Tale of Many Stores. *The Neuroscientist : a review journal bringing neurobiology, neurology and psychiatry*, 1073858418785334.
79. Ngo-Anh, T.J., Bloodgood, B.L., Lin, M., Sabatini, B.L., Maylie, J., and Adelman, J.P. (2005). SK channels and NMDA receptors form a  $\text{Ca}^{2+}$ -mediated feedback loop in dendritic spines. *Nat Neurosci* 8, 642-649.
80. Bloodgood, B.L., and Sabatini, B.L. (2007). Nonlinear regulation of unitary synaptic signals by  $\text{CaV}(2.3)$  voltage-sensitive calcium channels located in dendritic spines. *Neuron* 53, 249-260.
81. Criado-Marrero, M., Santini, E., and Porter, J.T. (2014). Modulating fear extinction memory by manipulating SK potassium channels in the infralimbic cortex. *Frontiers in Behavioral Neuroscience* 8, 96.
82. Lappin, S.C., Dale, T.J., Brown, J.T., Trezise, D.J., and Davies, C.H. (2005). Activation of SK channels inhibits epileptiform bursting in hippocampal CA3 neurons. *Brain Research* 1065, 37-46.

83. Lotshaw, D.P. (1997). Effects of K<sup>+</sup> channel blockers on K<sup>+</sup> channels, membrane potential, and aldosterone secretion in rat adrenal zona glomerulosa cells. *Endocrinology* 138, 4167-4175.
84. Suzuki, K., Ito, K.M., Minayoshi, Y., Suzuki, H., Asano, M., and Ito, K. (1993). Modification by charybdotoxin and apamin of spontaneous electrical and mechanical activity of the circular smooth muscle of the guinea-pig stomach. *Br J Pharmacol* 109, 661-666.
85. Buchanan, K.A., Petrovic, M.M., Chamberlain, S.E., Marrion, N.V., and Mellor, J.R. (2010). Facilitation of long-term potentiation by muscarinic M(1) receptors is mediated by inhibition of SK channels. *Neuron* 68, 948-963.
86. Rietdorf, K., Funnell, T.M., Ruas, M., Heinemann, J., Parrington, J., and Galione, A. (2011). Two-pore channels form homo- and heterodimers. *J Biol Chem* 286, 37058-37062.
87. Kemp, N., McQueen, J., Faulkes, S., and Bashir, Z.I. (2000). Different forms of LTD in the CA1 region of the hippocampus: role of age and stimulus protocol. *Eur J Neurosci* 12, 360-366.
88. Wang, W., Zhang, Z., Shang, J., Jiang, Z.Z., Wang, S., Liu, Y., and Zhang, L.Y. (2008). Activation of group I metabotropic glutamate receptors induces long-term depression in the hippocampal CA1 region of adult rats in vitro. *Neurosci Res* 62, 43-50.
89. Luscher, C., and Huber, K.M. (2010). Group 1 mGluR-dependent synaptic long-term depression: mechanisms and implications for circuitry and disease. *Neuron* 65, 445-459.
90. Bortolotto, Z.A., Bashir, Z.I., Davies, C.H., and Collingridge, G.L. (1994). A molecular switch activated by metabotropic glutamate receptors regulates induction of long-term potentiation. *Nature* 368, 740-743.
91. Raymond, C.R., Thompson, V.L., Tate, W.P., and Abraham, W.C. (2000). Metabotropic glutamate receptors trigger homosynaptic protein synthesis to prolong long-term potentiation. *J Neurosci* 20, 969-976.
92. Raymond, C.R., and Redman, S.J. (2002). Different calcium sources are narrowly tuned to the induction of different forms of LTP. *J Neurophysiol* 88, 249-255.
93. Nagaraja, R.Y., Becker, A., Reymann, K.G., and Balschun, D. (2005). Repeated administration of group I mGluR antagonists prevents seizure-induced long-term aberrations in hippocampal synaptic plasticity. *Neuropharmacology* 49 Suppl 1, 179-187.
94. Fiorillo, C.D., and Williams, J.T. (1998). Glutamate mediates an inhibitory postsynaptic potential in dopamine neurons. *Nature* 394, 78-82.
95. Balschun, D., Manahan-Vaughan, D., Wagner, T., Behnisch, T., Reymann, K.G., and Wetzel, W. (1999). A specific role for group I mGluRs in hippocampal LTP and hippocampus-dependent spatial learning. *Learning & memory (Cold Spring Harbor, N.Y.)* 6, 138-152.
96. Aiba, A., Chen, C., Herrup, K., Rosenmund, C., Stevens, C.F., and Tonegawa, S. (1994). Reduced hippocampal long-term potentiation and context-specific deficit in associative learning in mGluR1 mutant mice. *Cell* 79, 365-375.
97. Conquet, F., Bashir, Z.I., Davies, C.H., Daniel, H., Ferraguti, F., Bordi, F., Franz-Bacon, K., Reggiani, A., Matarese, V., Conde, F., et al. (1994). Motor deficit and impairment of synaptic plasticity in mice lacking mGluR1. *Nature* 372, 237-243.
98. Bear, M.F., Huber, K.M., and Warren, S.T. (2004). The mGluR theory of fragile X mental retardation. *Trends Neurosci* 27, 370-377.
99. Dolen, G., Osterweil, E., Rao, B.S., Smith, G.B., Auerbach, B.D., Chattarji, S., and Bear, M.F. (2007). Correction of fragile X syndrome in mice. *Neuron* 56, 955-962.
100. Aarhus, R., Graeff, R.M., Dickey, D.M., Walseth, T.F., and Lee, H.C. (1995). ADP-ribosyl cyclase and CD38 catalyze the synthesis of a calcium-mobilizing metabolite from NADP. *J Biol Chem* 270, 30327-30333.



**Acknowledgements:** The authors thank Dr Grant Churchill and Clive Garnham for synthesizing and providing us the NAADP-AM. **Funding:** This work was funded by the a Wellcome Trust Senior Investigator Enhancement Grant, “A messenger role for NAADP in the central nervous system” to AG, by an Alison Brading Scholarship from Lady Margaret Hall, Oxford to WJF, and by a grant from the BBSRC to NJE. **Author Contributions:** W.J.F. and H.T. performed and analysed the experiments. W.J.F, Z.P, A.F.J. designed the experiments. W.J.F. and N.J.E wrote the manuscript. A.G. and N.J.E supervised the project. **Competing interests:** The authors declare the have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper or the Supplementary Materials.

## Figure legends

**Figure 1. NAADP causes membrane depolarization in pyramidal neurons of the hippocampus in a manner dependent on acidic store signaling, intracellular  $\text{Ca}^{2+}$ , and RyR.**

**(A)** Diagram showing the experimental configuration to record membrane potential of CA1 or CA3 pyramidal neurons in hippocampal slices whilst NAADP-AM was applied locally. **(B)** Example voltage traces recorded whilst applying NAADP-AM, NAADP, or vehicle. Arrows indicate the start of delivery and grey bar indicates the total time of application. **(C)** Transient membrane depolarization ( $\Delta V_M$ ) upon application of NAADP-AM (300  $\mu\text{M}$ , n=12 cells), NAADP (300  $\mu\text{M}$ , n=5), or vehicle (n=6). Data are mean  $\pm$ SEM. **(D)** Mean  $\Delta V_M$  upon application of NAADP-AM (300  $\mu\text{M}$ , n=11 cells) alone or (left to right) in combination with a desensitizing concentration of NAADP (1 mM) inside the internal solution of the patch pipette (n=6);

preincubation with the NAADP antagonist Ned-19 (100  $\mu$ M, 40 min, n=6); preincubation with the vacuolar H<sup>+</sup> ATPase inhibitor bafilomycin (4  $\mu$ M, 40 min, n=5); with BAPTA (15 mM) inside the internal solution of the patch pipette (n=5); or pre-incubation with ryanodine (30  $\mu$ M, 40 min, n=4). Significance was assessed with; Data are mean  $\pm$ SEM; n = single cells. \*\*\* P < 0.005 by Kruskal-Wallis and post hoc Dunn's tests.

**Figure 2. NAADP is unique among second messengers in its ability to depolarize hippocampal pyramidal neurons. (A)** Example voltage traces for dialysis of CA1 pyramidal neurons patched with internal solutions containing various concentrations of the Ca<sup>2+</sup>-mobilizing second messengers NAADP, IP<sub>3</sub> and cADPR. Changes to membrane potential were recorded over time as the second messengers dialysed into the patched cell. **(B)** Transient membrane depolarization ( $\Delta V_M$ ) of the cells described in (A) in response to increasing concentrations of the Ca<sup>2+</sup>-mobilizing second messengers. Data are mean  $\pm$  SEM; n = single cells, indicated above each column; n > 4 for all concentrations of second messengers. \*\*\* P < 0.005 and \* P < 0.05 by Kruskal-Wallis and post hoc Dunn's tests.

**Figure 3. Activation of mGluR1 in CA1 pyramidal neurons causes a membrane depolarization that depends on NAADP signalling and acidic store Ca<sup>2+</sup> signalling. (A)** Diagram showing the experimental configuration to record membrane potential of CA1 pyramidal neurons in hippocampal slices whilst mGluRs were pharmacologically isolated (50  $\mu$ M AP5, 10  $\mu$ M NBQX, 100  $\mu$ M picrotoxin, and 2  $\mu$ M CGP 55845) and electrical stimulation was applied to Schaffer collaterals (4 pulses, 20 Hz), (n=11 cells). **(B)** Typical voltage recordings from single cells upon electrical stimulation with pharmacological isolation of mGluRs or plus

antagonism of group 2 and 3 mGluRs (LY341485 at 100 nM; n=6), pan-mGluRs (LY341485 at 100  $\mu$ M; n=6), mGluR5 (MPEP, 10  $\mu$ M; n=5) or mGluR1 (JNJ16259685, 300 nM; n=5) (top to bottom). The red lines indicate where membrane potentials were compared before and after stimulation. **(C and D)** Transient membrane depolarization ( $\Delta V_M$ ) of CA1 pyramidal neurons after electrical stimulation alone (control; n=7 cells) or with the presence of (C) the mGluR antagonism described in (B), or (D) pan-mGluR antagonist [LY341485, 100  $\mu$ M; n=6; cells and data set are independent from those in (C)], desensitizing concentration of NAADP inside the internal solution of the patch pipette (1 mM; n=6), NAAD inside the internal solution of the patch pipette (1 mM; n=6), preincubation with the NAADP antagonist Ned-19 (100  $\mu$ M, 40 min; n=6), or acute administration of the lysosomal disrupting agent GPN (200  $\mu$ M; n=5). Data are mean  $\pm$  SEM; n = single cells. \*\* P < 0.01 and \* P < 0.05 by Kruskal-Wallis and post hoc Dunn's tests.

**Figure 4. In CA1 pyramidal neurons, mGluR1-dependent membrane depolarization and Ca<sup>2+</sup> release require acidic store signalling and Ca<sup>2+</sup> release from the ER via ryanodine receptors but not IP<sub>3</sub> receptors. (A)** Diagram showing the experimental configuration. The membrane potential of CA1 pyramidal neurons in hippocampal slices was recorded whilst mGluR1 was pharmacologically isolated and extracellular glutamate was applied. **(B)** Typical voltage recordings recorded upon bath application of glutamate (300  $\mu$ M, 120 s) or the vehicle. **(C)** Columns show mean  $\Delta V_M$  of CA1 pyramidal neurons before and after extracellular glutamate application under control conditions (n=8) and in the presence of the lysosomal disrupting agent GPN (200  $\mu$ M; n=6), ryanodine receptor antagonist ryanodine (40  $\mu$ M, 15 min; n=6), IP<sub>3</sub> receptor antagonist Xestospongine C (2  $\mu$ M, 15 min; n=6), 'fast' Ca<sup>2+</sup> chelator BAPTA (20  $\mu$ M,

15 min; n=6), or the 'slow'  $\text{Ca}^{2+}$  chelator EGTA (20  $\mu\text{M}$ , 15 min; n=6). **(D)** Time series images of CA1 neurons filled with  $\text{Ca}^{2+}$  indicator OGB-1 (1 mM) were recorded whilst mGluR1 was pharmacologically isolated (50  $\mu\text{M}$  AP5, 10  $\mu\text{M}$  NBQX, 100  $\mu\text{M}$  picrotoxin, and 2  $\mu\text{M}$  CGP 55845) and electrical stimulation was applied (4 pulses, 20 Hz). Images top to bottom: z-stack of the dendritic branch being imaged (green);  $\text{Ca}^{2+}$  signal at baseline, before stimulation;  $\text{Ca}^{2+}$  signal 300 ms after stimulation; subtraction of  $\text{Ca}^{2+}$  at 300 ms from baseline (purple). Scale bar, 0.5  $\mu\text{m}$ . **(E)**  $\Delta F/F$  over the imaging time course where mGluR1 was pharmacologically isolated (n=20 cells) in combination with acute application of LY341495 (100  $\mu\text{M}$ , 10 min; n=5), pre-incubation with Ned-19 (100  $\mu\text{M}$ , 1 hour; n=6), or acute application of ryanodine (20  $\mu\text{M}$ , 10 min; n=5), xestospongine C (2  $\mu\text{M}$ , 15 min; n=5), or 2-APB (50  $\mu\text{M}$ , 15 min; n=5). **(F)** Columns show mean  $\Delta F/F$  before and after electrical stimulation for each pharmacological manipulation undertaken. Significance was assessed with Kruskal-Wallis and post hoc Dunn's tests; Error bars denote SEM, n = single cell. Significant differences indicated by asterisks where \*\*\* =  $P < 0.005$  \*\* =  $P < 0.01$  and \* =  $P < 0.05$ .

**Figure 5. In CA1 pyramidal neurons, mGluR1-dependent depolarization occurs through the inactivation of SK channels by possibly protein phosphatase 2A (PP2A).** **(A and B)** Representative voltage recordings (A) and mean  $\Delta V_M$  (B) upon electrical stimulation (4x, 20 Hz) of CA1 neurons whilst mGluR1 was pharmacologically isolated, then subsequent addition of apamin (200 nM, 15 min) and finally GPN (200  $\mu\text{M}$ , 10 min); n=6 cells. **(C and D)** Representative voltage recordings (C) and mean  $\Delta V_M$  (D) upon bath application of glutamate (red arrowhead; 300  $\mu\text{M}$ , 120 s) of CA1 neurons whilst mGluR1 was pharmacologically isolated, then subsequent addition of apamin (200 nM, 15 min) and finally GPN (200  $\mu\text{M}$ , 10

min); n=6 cells. **(E and F)** Representative voltage recordings (E) and mean  $\Delta V_M$  (F) upon electrical stimulation (4x, 20 Hz) of CA1 neurons whilst mGluR1 was pharmacologically isolated in the absence or presence of okadaic acid (100 nM, 15 min); n=6 cells. Data are means + SEM, each from n = 6 single cells. \*\* P < 0.01 and \* P < 0.05 (relative to mGluR1 isolation-alone condition), and n.s. = no significant difference; by Friedman's test with post hoc Dunn's tests (B and D) or a Wilcoxon pair matched signed ranks test (F).

**Figure 6. In CA1 pyramidal neurons, mGluR1-dependent synaptic plasticity requires inhibition of SK channels via NAADP signalling. (A)** A causal spike-timing-dependent plasticity protocol was used to induce mGluR1-dependent LTP, in which one causal presynaptic stimulation is paired with two backpropagating action potentials (100 Hz), at a 10 ms interval. The induction protocol is delivered where t = 0, indicated by the black triangles. Example EPSP traces before (black) and after (red) STDP induction are shown at the top right of each graph. Scale bar, 5 mV x 50 ms. This STDP protocol produces LTP lasting at least 30 min (n=7 cells). **(B)** LTP in the STDP protocol described in (A) with mGluR1-specific antagonism with JNJ16259685 (300 nM; n=5 cells). **(C)** LTP as described in (A) upon prevention of NAADP/acidic store Ca<sup>2+</sup> signalling with a desensitizing concentration of NAADP (5 mM; n=5 cells). **(D)** Magnitude of LTP upon induction of STDP in the presence of SK channel antagonist apamin (200 nM; n=7 cells). **(E)** LTP as described in (A) in the presence of apamin and JNJ16259685 (300 nM; n=6 cells). **(F)** Mean change in synaptic strength at 25-30 min, expressed as a % of the baseline. Data are means ± SEM, n = single cell. \* P < 0.05 by Kruskal-Wallis and post hoc Dunn's tests.

**Figure 7. In CA1 pyramidal neurons, Two-pore channels are required for mGluR1-mediated membrane depolarization and mGluR1-dependent LTP. (A)** Representative average (5 traces) voltage recordings from CA1 neurons in hippocampal slice preparations from wild-type (WT), *Tpc1*<sup>-/-</sup>, and *Tpc2*<sup>-/-</sup> mice (n=6 mice each). Recordings were obtained upon pharmacological isolation of mGluR1 (50  $\mu$ M AP5, 10  $\mu$ M NBQX, 100 nM LY341495, 10  $\mu$ M MPEP, 100  $\mu$ M picrotoxin, and 2  $\mu$ M CGP 55845) and electrical stimulation (4x, 20 Hz) of afferent fibres in stratum radiatum. Dashed red line indicates where membrane potentials were compared before and after stimulation. **(B)** Columns show mean  $\Delta V_M$  of CA1 pyramidal neurons before and after electrical stimulation described in (A). **(C)** A causal spike-time-dependent plasticity protocol was used to induce mGluR1-dependent LTP after a baseline of EPSPs were recorded for 5 min (indicated by marker at 0 min) in WT (n=4), *Tpc1*<sup>-/-</sup> (n=5) and *Tpc2*<sup>-/-</sup> (n=7) animals. One casual presynaptic stimulation is paired with two backpropagating action potentials (100 Hz) at a 10 ms interval. **(D)** Mean change in synaptic strength at 25-30 min shown/described in (C), expressed as a % of the baseline (red dashed line). Data are means  $\pm$  SEM, n = single cell. \*\* P < 0.01 and \* P < 0.05 by Kruskal-Wallis and post hoc Dunn's tests.

**Figure 8. Proposed model for mGluR1-dependent plasticity. (A)** Model of SK channel activation, wherein (i) synaptic glutamate activates GluA receptors to produce (ii) membrane depolarization and (iii) Ca<sup>2+</sup> entry via VDCCs. This causes (iv) activation of SK channels and local hyperpolarisation, resulting in inhibition of GluNs by reinstating Mg<sup>2+</sup> block, thereby reducing Ca<sup>2+</sup> entry through the GluNs and reducing the probability of LTP induction. Where synaptic activity is sufficiently strong, the mGluR1 receptors are recruited. **(B)** The proposed

model for SK channel inhibition mediated by mGluR1 signaling; GluA/VDCC regulation of SK channels is also present but not shown. (i) Glutamate activates mGluR1 receptors and causes (ii) NAADP synthesis, which results in (iii) acidic store  $\text{Ca}^{2+}$  release, which is amplified through activation of ryanodine receptors (RyR) in the ER. This somehow inactivates SK channels (iv), which in turn prevents local hyperpolarization and (v) allows greater  $\text{Ca}^{2+}$  entry through the GluN receptors, which facilitates the induction of LTP.