
Corrigendum: The plasma virome of febrile adult Kenyans shows frequent parvovirus B19 infections and a novel arbovirus (Kadipiro virus)

Carolyn N. Ngoi,^{1,2} Juliana Siqueira,^{2,3} Linlin Li,² Xutao Deng,² Peter Mugo,¹ Susan M. Graham,^{1,4} Matt A. Price,^{5,6} Eduard J. Sanders^{1,7} and Eric Delwart^{2,8,*}

J. Gen. Virol. 2016;97:3359–3367, doi: 10.1099/jgv.0.000644

There was an error in the Results section of the published article.

The samples from the plasma pool that yielded the Kadipiro virus (KDV) sequences using metagenomics were individually tested for KDV RNA using RT-nested PCR. All PCRs were negative using several primer sets. We are therefore unable to confirm the presence of KDV RNA in human plasma. The possible source of this contamination remains unidentified; KDV RNA may be a sporadic contaminant of one of the reagents or of the QIAamp Viral RNA Mini Kit (Qiagen) kit used.

The authors apologize for any inconvenience.

Received 1 March 2017; Accepted 1 March 2017

Author affiliations: ¹Centre for Geographic Medicine Research – Coast, Kenya Medical Research Institute, Kilifi, Kenya; ²Blood Systems Research Institute, San Francisco, CA, USA; ³Programa de Oncovirologia, Instituto Nacional de Câncer, Rio de Janeiro, Brazil; ⁴University of Washington, Seattle, WA, USA; ⁵International AIDS Vaccine Initiative, New York, NY, USA; ⁶Department of Epidemiology and Biostatistics, University of California at San Francisco, CA, USA; ⁷Nuffield Department of Medicine, University of Oxford, Headington, UK; ⁸Department of Laboratory Medicine, University of California at San Francisco, CA, USA.

***Correspondence:** Eric Delwart, delwarte@medicine.ucsf.edu