

ROUGH PATH METRICS ON A BESOV–NIKOLSKII-TYPE SCALE

PETER K. FRIZ AND DAVID J. PRÖMEL

ABSTRACT. It is known, since the seminal work [T. Lyons, Differential equations driven by rough signals, *Rev. Mat. Iberoamericana*, 14 (1998)], that the solution map associated to a controlled differential equation is locally Lipschitz continuous in q -variation, resp., $1/q$ -Hölder-type metrics on the space of rough paths, for any regularity $1/q \in (0, 1]$.

We extend this to a new class of Besov–Nikolskii-type metrics, with arbitrary regularity $1/q \in (0, 1]$ and integrability $p \in [q, \infty]$, where the case $p \in \{q, \infty\}$ corresponds to the known cases. Interestingly, the result is obtained as a consequence of known q -variation rough path estimates.

1. INTRODUCTION

We are interested in controlled differential equations of the type

$$(1.1) \quad dY_t = V(Y_t) dX_t, \quad t \in [0, T],$$

where $X = (X_t)$ is a suitable (n -dimensional) driving signal, $Y = (Y_t)$ is the (m -dimensional) output signal, and $V = (V_1, \dots, V_n)$ are vector fields of suitable regularity. A fundamental question concerns the continuity of the solution map $X \mapsto Y$, strongly dependent on the used metric.

A decisive answer is given by rough path theory, which identifies a cascade of good metrics, determined by some *regularity parameter* $\delta \equiv 1/q \in (0, 1]$, and essentially given by q -variation, resp., δ -Hölder-type metrics. As long as the driving signal X possesses sufficient regularity, say X is a continuous path of finite q -variation for $q \in [1, 2)$ (in symbols $X \in C^{q\text{-var}}([0, T]; \mathbb{R}^n)$), Lyons [Lyo94] showed that the solution map $X \mapsto Y$ associated to equation (1.1) is a locally Lipschitz continuous map with respect to the q -variation topology. However, this strong regularity assumption on X excludes many prominent examples from probability theory as sample paths of stochastic processes like (fractional) Brownian motion, martingales, or many Gaussian processes.

In order to restore the continuity of the solution map associated to a controlled differential equation for continuous paths X of finite q -variation for arbitrary large

Received by the editors October 21, 2016, and, in revised form, March 30, 2017.

2010 *Mathematics Subject Classification.* Primary 34A34, 60H10; Secondary 26A45, 30H25, 46N20.

Key words and phrases. Controlled differential equation, Besov embedding, Besov space, Itô–Lyons map, p -variation, Riesz-type variation, rough path.

The first author was partially supported by the European Research Council through CoG-683164 and DFG research unit FOR2402.

The second author gratefully acknowledges financial support of the Swiss National Foundation under Grant No. 200021_163014.

$q < \infty$, it is not sufficient anymore to consider a path “only” taking in the Euclidean space \mathbb{R}^n ; cf. [Lyo91, LCL07]. Instead, X must be viewed as a $\lfloor q \rfloor$ -level rough path, which in particular means X takes values in a step- $\lfloor q \rfloor$ free nilpotent group $G^{\lfloor q \rfloor}(\mathbb{R}^n)$: Let us recall that for $Z \in C^{1\text{-var}}(\mathbb{R}^n)$ its $\lfloor q \rfloor$ -step signature is given by

$$S_{\lfloor q \rfloor}(Z)_{s,t} := \left(1, \int_{s < u < t} dZ_u, \dots, \int_{s < u_1 < \dots < u_{\lfloor q \rfloor} < t} dZ_{u_1} \otimes \dots \otimes dZ_{u_{\lfloor q \rfloor}} \right).$$

The corresponding space of all these lifted paths is

$$G^{\lfloor q \rfloor}(\mathbb{R}^n) := \{S_{\lfloor q \rfloor}(Z)_{0,T} : Z \in C^{1\text{-var}}([0,T]; \mathbb{R}^n)\} \subset \bigoplus_{k=0}^{\lfloor q \rfloor} (\mathbb{R}^n)^{\otimes k},$$

which we equip with the Carnot–Carathéodory metric d_{cc} ; see Subsection 3.2 for more details. While in the case of $q \in [1, 2)$ this reduces to a classical path $X: [0, T] \rightarrow \mathbb{R}^n$, in the case of $q > 2$ this means, intuitively, X is a path enhanced with the information corresponding to the “iterated integrals” up to order $\lfloor q \rfloor$. In the context of rough path theory, the solution map $X \mapsto Y$, taking now a $\lfloor q \rfloor$ -level rough path X (in symbols $X \in C^{q\text{-var}}([0, T], G^{\lfloor q \rfloor}(\mathbb{R}^n))$) as input, is often called the *Itô–Lyons map*.

In most applications, the output is regarded as path, $Y \in C^{q\text{-var}}([0, T]; \mathbb{R}^m)$, although—depending on the route one takes—it can be seen as rough path [Lyo98, LQ02, FV10] or controlled rough path [Gub04, FH14]. It is a fundamental property of rough path theory that solving differential equations, that is, applying the Itô–Lyons map entails no loss of regularity: if the driving signal enjoys δ -Hölder (resp., q -variation) regularity, then so does the output signal.

Let us explain the basic idea which underlies this work. To this end (only estimates matter) take X smooth and rewrite (1.1) in the classical form $\dot{Y} = V(Y)\dot{X}$. Take L^p -norms on both sides to arrive at

$$(1.2) \quad \|Y\|_{W^{1,p};[0,T]} \leq \|V\|_{\infty} \|X\|_{W^{1,p};[0,T]}$$

in terms of the semi-norm $\|X\|_{W^{1,p};[0,T]} := (\int_0^T |\dot{X}_t|^p dt)^{1/p}$. Here, of course, we have regularity $\delta = 1$ ($\Leftrightarrow q = 1$), and the extreme cases $p \in \{1, \infty\}$ ($= \{q, \infty\}$) amount exactly to the variation, resp., Hölder estimates

$$(1.3) \quad \begin{aligned} \|Y\|_{1\text{-var};[0,T]} &\leq \|V\|_{\infty} \|X\|_{1\text{-var};[0,T]}, \\ \|Y\|_{1\text{-Höl};[0,T]} &\leq \|V\|_{\infty} \|X\|_{1\text{-Höl};[0,T]}, \end{aligned}$$

since indeed $\|X\|_{1\text{-var};[0,T]} \approx \|X\|_{W^{1,1}[0,T]}$, resp., $\|X\|_{1\text{-Höl};[0,T]} = \|X\|_{W^{1,\infty};[0,T]}$. Conversely, one may view (1.2) as *interpolation* of the estimates (1.3), by regarding $W^{1,p}$, for any $p \in [1, \infty]$, as interpolation space of $W^{1,1}$ and $W^{1,\infty}$. This discussion suggests moreover that the solution map $X \mapsto Y$ is also continuous in $W^{1,p}$, even locally Lipschitz in the sense

$$(1.4) \quad \|Y^1 - Y^2\|_{W^{1,p};[0,T]} \lesssim \|X^1 - X^2\|_{W^{1,p};[0,T]},$$

as indeed may be seen by some fairly elementary analysis. (Mind, however, that the solution map $X \mapsto Y$ is highly non-linear so that there is little hope to appeal to some “general theory of interpolation”.)

The estimates (1.3) and (1.4), in case $p = 1$ and $p = \infty$, are well known (e.g., [Lyo98, LQ02, FV10]) to extend to arbitrarily low regularity $\delta \equiv 1/q \in (0, 1]$, provided that, essentially, $\|\cdot\|_{1\text{-var};[0,T]}$ is replaced by $\|\cdot\|_{q\text{-var};[0,T]}$ (with the correct rough path interpretation on the right-hand sides above).

The question arises if the well-studied q -variation and δ -Hölder formulation of rough path theory are not the extreme cases of a more flexible formulation of the theory, that comes—in the spirit of Besov (Nikolskii) spaces—with an additional *integrability parameter* $p \in [q, \infty]$. (Here, having q as lower bound on p is quite natural in view of known Besov embeddings: in the Besov-scale $(B_r^{\delta,p})$, with additional fine-tuning parameter r , one has, always with $\delta = 1/q$, $C^{q\text{-var}} \approx N^{\delta,q} \equiv B_{\infty}^{\delta,q}$, in the form of tight (but strict) inclusions $N^{\delta+\varepsilon,q} \subset C^{q\text{-var}} \subset N^{\delta,q}$; see Remark 2.1.)

The first contribution of this paper is to give an affirmative answer to the above question, in the generality of arbitrarily low regularity $\delta > 0$. With focus on the interesting case of regularity $\delta < 1$, we have, loosely stated, the following theorem.

Theorem 1.1. *Let $\delta \equiv 1/q \in (0, 1]$ and $p \in [q, \infty]$. Then, for Lip^γ vector fields V with $\gamma > q$, the Itô–Lyons map (as defined below in (3.5)) is locally Lipschitz continuous from a Besov–Nikolskii-type (rough) path space with regularity/integrability (δ, p) into a Besov–Nikolskii-type path space of identical regularity/integrability (δ, p) .*

Somewhat surprisingly, it is possible to prove this via delicate application of classical q -variation estimates¹ in rough path theory; that is, morally, from the case $p = q$. On the other hand, a precise definition of the involved spaces—to make this reasoning possible—is a subtle matter. First, care is necessary for rough paths take values in a non-linear space, the step- $[q]$ free nilpotent group $G^{[q]}(\mathbb{R}^n)$ equipped with the Carnot–Carathéodory metric d_{cc} , which is a no standard setting for classical Besov, resp., Nikolskii spaces $(B_r^{\delta,p})$, resp., $N^{\delta,p} := B_{\infty}^{\delta,p}$. Another and quite serious difficulty is the lack of super-additivity of Nikolskii norms. Recall that the “control”

$$\omega(s, t) := \|X\|_{p\text{-var};[s,t]}^p$$

has the most desirable property of super-additivity, i.e., $\omega(s, t) + \omega(t, u) \leq \omega(s, u)$, a simple fact that is used throughout Lyons’ theory. For instance, as a typical consequence

$$\|X\|_{p\text{-var};[0,T]}^p = \sup_{\mathcal{P} \subset [0,T]} \sum_{[u,v] \in \mathcal{P}} \|X\|_{p\text{-var};[u,v]}^p,$$

where the supremum is taking over all partition of the interval $[0, T]$. Several other (rough) path space norms also have this property, as exploited, e.g., in [FV06]. However, this convenient property fails for the Besov spaces of consideration (unless $\delta = 1$) and indeed, in general with strict inequality,

$$\|X\|_{N^{\delta,p};[0,T]}^p \leq \sup_{\mathcal{P} \subset [0,T]} \sum_{[u,v] \in \mathcal{P}} \|X\|_{N^{\delta,p};[u,v]}^p =: \|X\|_{\hat{N}^{\delta,p};[0,T]}^p.$$

This leads us to use the *Besov–Nikolskii-type space* $\hat{N}^{\delta,p}$, defined as those X for which the right-hand side above is finite, as the correct space (in rough path or path space incarnation) to which we refer in Theorem 1.1, at least in the new regimes $\delta < 1$, $p \in (q, \infty)$.

¹The use of control functions is pure notational convenience.

A better understanding of these spaces is compulsory, and this is the second contribution of this paper. For instance, it is reassuring that one has tight inclusions of the form $N^{\delta+\varepsilon,p} \subset \hat{N}^{\delta,p} \subset N^{\delta,p}$ (Corollary 2.12). In fact, an exact characterization is possible in terms of *Riesz-type variation* spaces, in reference to Riesz [Rie10], who considered such spaces (although with regularity parameter $\delta = 1$). We have the following theorem.

Theorem 1.2. *Consider $\delta = 1/q < 1$ and $p \in (q, \infty)$. Then the Besov–Nikolskii-type space $\hat{N}^{\delta,p}$ coincides with the Riesz-type variation spaces $V^{\delta,p}$ and $\tilde{V}^{\delta,p}$ defined, respectively, via finiteness of*

$$\begin{aligned} \|X\|_{V^{\delta,p}}^p &:= \sup_{\mathcal{P} \subset [0,T]} \sum_{[u,v] \in \mathcal{P}} \frac{d_{cc}(X_v, X_u)^p}{|v-u|^{\delta p-1}}, \\ \|X\|_{\tilde{V}^{\delta,p}}^p &:= \sup_{\mathcal{P} \subset [0,T]} \sum_{[u,v] \in \mathcal{P}} \frac{\|X\|_{\frac{1}{\delta}\text{-var};[u,v]}^p}{|v-u|^{\delta p-1}}, \end{aligned}$$

for a rough path X and the Carnot–Carathéodory distance d_{cc} . More generally, this is also true for arbitrary metric spaces instead of $G^{[q]}(\mathbb{R}^n)$.

Moreover, all associated inhomogenous rough path distances are locally Lipschitz equivalent.

Let us also note that the above introduced Riesz-type variation spaces agree (trivially) with the q -variation space in the extreme case of $p = q \equiv 1/\delta$. (In the Besov scale, this usually fails. For instance, we have the strict inclusion $W^{1,1} \subset C^{1\text{-var}}$; not every rectifiable path is absolutely continuous.)

We conclude this introduction with some pointers to previous works. The case of regularity $\delta > 1/2$, essentially a Young regime, was considered in [Zäh98, Zäh01]. Our result can also be regarded as an extension of [PT16], which effectively dealt with regularity $\delta = 1/q > 1/3$ and accordingly integrability $p \geq q = 3$. We note that path spaces with “mixed” Hölder-variation regularity, similar in spirit to the Riesz-type spaces (with tilde) also appear as tangent spaces to Hölder rough path spaces [FV10, p. 209]; see also [Aid16]. Moreover, regularity of Cameron–Martin spaces associated to Gaussian processes with “Hölder dominated ρ -variation of the covariance” (a key condition in Gaussian rough path theory; cf. [FH14, Ch. 10], [FGGR16]) can be expressed with the help of “mixed” Hölder-variation regularity; see, e.g., [FH14, p. 151].

Organization of the paper. In Section 2 we define and give various characterizations of our spaces, starting for the reader’s convenience with the (much) simpler situation $\delta = 1$. In particular, Theorem 1.2 is an effective summary of Theorem 2.11 and Lemmas 3.4, 3.6, and 3.7. Section 3 is devoted to establishing the local Lipschitz continuity of the Itô–Lyons map in suitable rough path metrics and Theorem 1.1 can be found in Theorem 3.3 and Corollaries 3.5 and 3.8.

2. RIESZ-TYPE VARIATION

In this section we introduce a class of function spaces which unifies the notions of Hölder and q -variation regularity. For this purpose we generalize an old version of variation due to F. Riesz and provide two alternative but equivalent characterizations of the so-called Riesz-type variation and additionally various embedding

results. As explained in the introduction, the latter application in the rough path framework requires us to set up all the function spaces for paths taking values in a metric space.

Let us briefly fix some basic notation: \mathcal{P} is called a partition of an interval $[s, t] \subset [0, T]$ if $\mathcal{P} = \{[t_i, t_{i+1}] : s = t_0 < t_1 < \cdots < t_n = t, n \in \mathbb{N}\}$. In this case we write $\mathcal{P} \subset [s, t]$ indicating that \mathcal{P} is a partition of the interval $[s, t]$. Furthermore, for such a partition \mathcal{P} and a function $\chi : \{(u, v) : s \leq u < v \leq t\} \rightarrow \mathbb{R}$ we use the abbreviation

$$\sum_{[u,v] \in \mathcal{P}} \chi(u, v) := \sum_{i=0}^{n-1} \chi(t_i, t_{i+1}).$$

If not otherwise specified, (E, d) denotes a metric space, $T \in (0, \infty)$ is a finite real number, and $C([0, T]; E)$ stands for the set of all continuous functions $f : [0, T] \rightarrow E$.

Two frequently used topologies to measure the regularity of functions are the Hölder continuity and the q -variation:

The *Hölder continuity* of a function $f \in C([0, T]; E)$ is measured by

$$\|f\|_{\delta\text{-Hö};[s,t]} := \sup_{u,v \in [s,t], u < v} \frac{d(f_u, f_v)}{|v - u|^\delta}, \quad \delta \in (0, 1],$$

and $C^{\delta\text{-Hö}}([0, T]; E)$ stands for the set of all functions $f \in C([0, T]; E)$ such that $\|f\|_{\delta\text{-Hö}} := \|f\|_{\delta\text{-Hö};[0,T]} < \infty$. The case $\delta = 1$, that is, the Hölder continuity of order 1, is usually referred to as Lipschitz continuity.

The *q -variation* of a function $f \in C([0, T]; E)$ is defined by

$$(2.1) \quad \|f\|_{q\text{-var};[s,t]} := \left(\sup_{\mathcal{P} \subset [s,t]} \sum_{[u,v] \in \mathcal{P}} d(f_u, f_v)^q \right)^{\frac{1}{q}}, \quad q \in [1, \infty),$$

where the supremum is taken over all partitions \mathcal{P} of the interval $[s, t]$. The set of all functions $f \in C([0, T]; E)$ with $\|f\|_{q\text{-var}} := \|f\|_{q\text{-var};[0,T]} < \infty$ is denoted by $C^{q\text{-var}}([0, T]; E)$. The notion of q -variation can be traced back to N. Wiener [Wie24]. The special case of 1-variation is also called bounded variation. A comprehensive list of generalizations of q -variation and further references can be found in [DN99].

Remark 2.1. Classical function spaces as fractional Sobolev or more general Besov spaces do not provide a unifying framework simultaneously covering the space of Hölder continuous functions and the space of continuous functions with finite q -variation. For example, let us replace for a moment (E, d) by the Euclidean space $(\mathbb{R}, |\cdot|)$ and denote the homogeneous Besov spaces by $B_r^{\delta,p}([0, T]; \mathbb{R})$. While the Hölder space $C^{\delta\text{-Hö}}([0, T]; \mathbb{R})$ is a special case of Besov spaces, namely the homogeneous Besov space $B_\infty^{\delta,\infty}([0, T]; \mathbb{R})$, for $\delta \in (0, 1)$, the q -variation space $C^{q\text{-var}}([0, T]; \mathbb{R})$ is not covered by the wide class of Besov spaces. Indeed, classical embedding theorems, [You36] and [LY38], yield the following continuous embeddings:

$$B_\infty^{\alpha,p}([0, T]; \mathbb{R}) \subset C^{p\text{-var}}([0, T]; \mathbb{R}) \subset B_\infty^{1/p,p}([0, T]; \mathbb{R})$$

for $p \in (1, \infty)$ and $\alpha \in (1/p, 1)$; see also [Sim90] and [FV06]. In particular, it is known that the second embedding is not an equality. An example can be found in [Ter67]. The relation between the space of functions with finite q -variation and Besov spaces was investigated in the literature for a long time; see for example [MS61], [Pee76], [BLS06], and [Ros09]. For a comprehensive introduction to function spaces we refer to [Tri10].

To set up a class of function spaces covering precisely and simultaneously the Hölder spaces and the q -variation spaces, we introduce a generalized version of a variation due to F. Riesz [Rie10]. For $\delta \in (0, 1]$ and $p \in [1/\delta, \infty)$ the *Riesz-type variation* of a function $f \in C([0, T]; E)$ is given by

$$(2.2) \quad \|f\|_{V^{\delta,p};[s,t]} := \left(\sup_{\mathcal{P} \subset [s,t]} \sum_{[u,v] \in \mathcal{P}} \frac{d(f_u, f_v)^p}{|v-u|^{\delta p-1}} \right)^{\frac{1}{p}}$$

for a subinterval $[s, t] \subset [0, T]$ and for $p = \infty$ we set

$$(2.3) \quad \|f\|_{V^{\delta,\infty};[s,t]} := \sup_{u,v \in [s,t], u < v} \frac{d(f_u, f_v)}{|v-u|^\delta}.$$

The set $V^{\delta,p}([0, T]; E)$ denotes all continuous functions $f \in C([0, T]; E)$ such that $\|f\|_{V^{\delta,p}} := \|f\|_{V^{\delta,p};[0,T]} < \infty$. The case of $\delta = 1$ was originally defined by F. Riesz [Rie10] and a similar generalization as given in (2.2) was already mentioned in [Pee76, p. 114, (14')].

Proposition 2.2. *Let (E, d) be a metric space and $T \in (0, \infty)$. For $\delta \in (0, 1]$ and $p \in [1/\delta, \infty]$ one has the following relations:*

$$\begin{aligned} C^{\delta\text{-Hö}}([0, T]; E) &= V^{\delta,\infty}([0, T]; E) \subset V^{\delta,p}([0, T]; E) \subset V^{\delta,1/\delta}([0, T]; E) \\ &= C^{1/\delta\text{-var}}([0, T]; E). \end{aligned}$$

More precisely, the $1/\delta$ -variation of a function $f \in V^{\delta,p}([0, T]; E)$ satisfies the bound

$$\|f\|_{1/\delta\text{-var};[s,t]} \leq \|f\|_{V^{\delta,p};[s,t]} |t-s|^{\delta-\frac{1}{p}}$$

for every subinterval $[s, t] \subset [0, T]$.

Before we come to the proof, we need the following remark about super-additive functions.

Remark 2.3. Setting $\Delta_T := \{(s, t) : 0 \leq s \leq t \leq T\}$ a function $\omega : \Delta_T \rightarrow [0, \infty)$ is called super-additive if

$$\omega(s, t) + \omega(t, u) \leq \omega(s, u) \quad \text{for } 0 \leq s \leq t \leq u \leq T.$$

Furthermore, if ω and $\tilde{\omega}$ are super-additive and $\alpha, \beta > 0$ with $\alpha + \beta \geq 1$, then $\omega^\alpha \tilde{\omega}^\beta$ is super-additive. The proof works as [FV10, Exercise 1.8 and 1.9].

Proof of Proposition 2.2. The identities

$$C^{\delta\text{-Hö}}([0, T]; E) = V^{\delta,\infty}([0, T]; E) \quad \text{and} \quad V^{\delta,1/\delta}([0, T]; E) = C^{1/\delta\text{-var}}([0, T]; E)$$

are ensured by the definitions of the involved function spaces.

The first embedding can be seen by

$$\|f\|_{V^{\delta,p}}^p = \sup_{\mathcal{P} \subset [0,T]} \sum_{[u,v] \in \mathcal{P}} \left(\frac{d(f_u, f_v)}{|v-u|^\delta} \right)^p |v-u| \leq T \|f\|_{C^\delta;[0,T]}^p, \quad f \in C^\delta([0, T]; E).$$

The second embedding is trivial for $\delta = 1/p$. For $\delta > 1/p$ we first observe that

$$(2.4) \quad d(f_s, f_t) \leq \left(\frac{d(f_s, f_t)^p}{|t-s|^{\delta p-1}} \right)^{\frac{1}{p}} |t-s|^{\delta-\frac{1}{p}} \leq \|f\|_{V^{\delta,p};[s,t]} |t-s|^{\delta-\frac{1}{p}}, \quad [s, t] \subset [0, T],$$

for $f \in V^{\delta,p}([0, T]; E)$, and thus

$$d(f_s, f_t)^{\frac{1}{\delta}} \leq \|f\|_{V^{\delta,p};[s,t]}^{\frac{1}{\delta}} |t-s|^{1-\frac{1}{\delta p}} =: \omega(s, t).$$

Since $\|f\|_{V^{\delta,p};[s,t]}^p$ and $|t-s|$ are super-additive as functions in $(s,t) \in \Delta_T$ and $(\delta p)^{-1} + 1 - (\delta p)^{-1} \geq 1$, ω is a super-additive by Remark 2.3. Hence, using the super-additivity of ω , we arrive at the claimed estimate

$$\|f\|_{1/\delta\text{-var};[s,t]} \leq \|f\|_{V^{\delta,p};[s,t]} |t-s|^{\delta-\frac{1}{p}}. \quad \square$$

The next lemma justifies the definition of the Riesz-type variation in the case of $p = \infty$, cf. (2.3), and collects some embedding results of these sets of functions.

Lemma 2.4. *Let (E, d) be a metric space, $T \in (0, \infty)$ and $[s, t] \subset [0, T]$. Suppose $\delta \in (0, 1)$ and $p \in [1/\delta, \infty]$.*

(1) *If $\delta > 1/p$, then $V^{\delta,p}([0, T]; E) \subset C^{(\delta-1/p)\text{-H\"{o}l}}([0, T]; E)$ with the estimate*

$$d(f_s, f_t) \leq \|f\|_{V^{\delta,p};[s,t]} |t-s|^{\delta-\frac{1}{p}}, \quad f \in V^{\delta,p}([0, T]; E).$$

(2) *If $\delta, \delta' \in (0, 1)$ and $p, p' \in [1/\delta, \infty]$ with $\delta' < \delta$ and $p' < p$, then one has*

$$V^{\delta,p}([0, T]; E) \subset V^{\delta',p}([0, T]; E) \quad \text{and} \quad V^{\delta,p}([0, T]; E) \subset V^{\delta,p'}([0, T]; E)$$

with the estimates for $f \in V^{\delta,p}([0, T]; E)$

$$\|f\|_{V^{\delta',p};[s,t]} \leq (t-s)^{\delta-\delta'} \|f\|_{V^{\delta,p};[s,t]} \quad \text{and} \quad \|f\|_{V^{\delta,p'};[s,t]} \leq (t-s)^{\frac{1}{p'}-\frac{1}{p}} \|f\|_{V^{\delta,p};[s,t]}.$$

(3) *For every $f \in V^{\delta,\infty}([0, T]; E)$ one has*

$$\lim_{p \rightarrow \infty} \|f\|_{V^{\delta,p};[s,t]} = \|f\|_{V^{\delta,\infty};[s,t]}.$$

Proof. (1) The first assertion follows directly by the estimate (2.4).

(2) Let \mathcal{P} be a partition of the interval $[s, t] \subset [0, T]$. For $f \in V^{\delta,p}([0, T]; E)$ the estimates

$$\sum_{[u,v] \in \mathcal{P}} \frac{d(f_u, f_v)^p}{|v-u|^{\delta'p-1}} \leq |t-s|^{(\delta-\delta')p} \sum_{[u,v] \in \mathcal{P}} \frac{d(f_u, f_v)^p}{|v-u|^{\delta p-1}}$$

and (using Hölder's inequality)

$$\begin{aligned} \sum_{[u,v] \in \mathcal{P}} \frac{d(f_u, f_v)^{p'}}{|v-u|^{\delta'p'-1}} &= \sum_{[u,v] \in \mathcal{P}} \left(\frac{d(f_u, f_v)}{|v-u|^{\delta-\frac{1}{p}}} \right)^{p'} |v-u|^{1-\frac{p'}{p}} \\ &\leq |t-s|^{1-\frac{p'}{p}} \left(\sum_{[u,v] \in \mathcal{P}} \frac{d(f_u, f_v)^p}{|v-u|^{\delta p-1}} \right)^{\frac{p'}{p}} \end{aligned}$$

lead to (2) by taking the supremum over all partitions of $[s, t]$.

(3) Due to Lemma 2.4 (1), we have

$$\|f\|_{V^{\delta,\infty};[s,t]} \leq \liminf_{p \rightarrow \infty} \|f\|_{V^{\delta,p};[s,t]}, \quad f \in V^{\delta,\infty}([0, T]; E).$$

Furthermore, for $p > q \geq 1$ we get

$$\|f\|_{V^{\delta,p};[s,t]} = \left(\sup_{\mathcal{P} \subset [s,t]} \sum_{[u,v] \in \mathcal{P}} \frac{d(f_u, f_v)^q}{|v-u|^{\delta q-1}} \frac{d(f_u, f_v)^{p-q}}{|v-u|^{\delta(p-q)}} \right)^{\frac{1}{p}} \leq \|f\|_{V^{\delta,q};[s,t]}^{\frac{q}{p}} \|f\|_{V^{\delta,\infty};[s,t]}^{1-\frac{q}{p}}$$

and thus

$$\limsup_{p \rightarrow \infty} \|f\|_{V^{\delta,p};[s,t]} \leq \|f\|_{V^{\delta,\infty};[s,t]}. \quad \square$$

In the following we introduce two different but equivalent characterizations of the Riesz-type variation (2.2). The first one is based on the classical notion of q -variation due to Wiener and thus is particularly convenient for applications in rough path theory. The second one relies on certain Besov spaces, namely Nikolskii spaces, which allows us to relate the Riesz-type variation spaces to classical function spaces as fractional Sobolev spaces. See Lemma 2.6 and Theorem 2.11 for the equivalence.

In order to give a characterization of Riesz-type variation of a function $f \in C([0, T]; E)$ in terms of q -variation, we introduce a *mixed Hölder-variation regularity* by

$$(2.5) \quad \|f\|_{\tilde{V}^{\delta,p};[s,t]} := \left(\sup_{\mathcal{P} \subset [s,t]} \sum_{[u,v] \in \mathcal{P}} \frac{\|f\|_{\frac{1}{\delta}\text{-var};[u,v]}^p}{|v-u|^{\delta p-1}} \right)^{\frac{1}{p}}, \quad \delta \in (0, 1], p \in [1/\delta, \infty),$$

for a subinterval $[s, t] \subset [0, T]$ and in the case of $p = \infty$ we define

$$\|f\|_{\tilde{V}^{\delta,\infty};[s,t]} := \sup_{\mathcal{P} \subset [s,t]} \sup_{[u,v] \in \mathcal{P}} \frac{\|f\|_{\frac{1}{\delta}\text{-var};[u,v]}}{|v-u|^\delta}.$$

Moreover, we denote by $\tilde{V}^{\delta,p}([0, T]; E)$ the set of all functions $f \in C([0, T]; E)$ such that $\|f\|_{\tilde{V}^{\delta,q}} := \|f\|_{\tilde{V}^{\delta,q};[0,T]} < \infty$.

An alternative way to measure Riesz-type variation of a function $f \in C([0, T]; E)$ is related to homogeneous Nikolskii spaces. Hence, we briefly recall the notation of homogeneous *Nikolskii spaces*, which correspond to the homogeneous Besov spaces $B_\infty^{\delta,p}([0, T]; E)$. For $\delta \in (0, 1]$ and $p \in [1, \infty)$ we define

$$\|f\|_{N^{\delta,p};[s,t]} := \sup_{|t-s| \geq h > 0} h^{-\delta} \left(\int_s^{t-h} d(f_u, f_{u+h})^p du \right)^{\frac{1}{p}}$$

for a subinterval $[s, t] \subset [0, T]$ and for $p = \infty$ we further set

$$\|f\|_{N^{\delta,\infty};[s,t]} := \sup_{|t-s| \geq h > 0} h^{-\delta} \sup_{u \in [s, t-h]} d(f_u, f_{u+h}).$$

The set of all functions $f \in C([0, T]; E)$ such that $\|f\|_{N^{\delta,p}} := \|f\|_{N^{\delta,p};[0,T]} < \infty$ is denoted by $N^{\delta,p}([0, T]; E)$.

Using the definition of Nikolskii regularity, we introduce a *refined Nikolskii-type regularity* by

$$(2.6) \quad \|f\|_{\hat{N}^{\delta,p};[s,t]} := \left(\sup_{\mathcal{P} \subset [s,t]} \sum_{[u,v] \in \mathcal{P}} \|f\|_{N^{\delta,p};[u,v]}^p \right)^{\frac{1}{p}}, \quad \delta \in (0, 1], p \in [1, \infty),$$

for $f \in C([0, T]; E)$ and a subinterval $[s, t] \subset [0, T]$. For $p = \infty$ we set

$$\|f\|_{\hat{N}^{\delta,\infty};[s,t]} := \sup_{\mathcal{P} \subset [s,t]} \sup_{[u,v] \in \mathcal{P}} \|f\|_{N^{\delta,\infty};[u,v]}.$$

Furthermore, $\hat{N}^{\delta,p}([0, T]; E)$ stands for the set of all functions $f \in C([0, T]; E)$ such that $\|f\|_{\hat{N}^{\delta,q}} := \|f\|_{\hat{N}^{\delta,q};[0,T]} < \infty$.

Remark 2.5. While $\|\cdot\|_{\hat{N}^{\delta,p};[s,t]}^p$ is a super-additive function in $(s, t) \in \Delta_T$ by its definition, this is not true for the Nikolskii regularity $\|\cdot\|_{N^{\delta,p};[s,t]}$ itself if $\delta \in (0, 1)$. The latter can be seen particularly by Remark 2.13.

In the next two subsections we show that the just introduced two ways of measuring path regularity are indeed equivalent to the Riesz-type variation. We start by considering the special case of regularity $\delta = 1$, that is, the space $V^{1,p}$, in Subsection 2.1. The equivalence for general Riesz-type variation spaces is the content of Subsection 2.2

2.1. Characterization of the space $V^{1,p}$. The special case $\delta = 1$ or in other words the set $V^{1,p}([0, T]; \mathbb{R}^n)$ coincides with the original definition due to F. Riesz [Rie10] and is already fairly well understood. For the sake of completeness we present here the full picture assuming $E = \mathbb{R}^n$ since it will be general enough for the latter applications concerning the solution map associated to a controlled differential equation; see Subsection 3.1.

It is well known that the Riesz-type variation space $V^{1,p}([0, T], \mathbb{R}^n)$ corresponds to the classical Sobolev space $W^{1,p}([0, T]; \mathbb{R}^n)$; see, e.g., [FV10, Proposition 1.45]. Let us recall the definition of the Sobolev space $W^{1,p}([0, T]; \mathbb{R}^n)$ (cf. [FV10, Definition 1.41]). For $p \in [1, \infty]$ and $T \in (0, \infty)$ a function $f \in C([0, T]; \mathbb{R}^n)$ is in $W^{1,p}([0, T]; \mathbb{R}^n)$ if and only if f is of the form

$$f_t = f_0 + \int_0^t f'_s \, ds, \quad t \in [0, T],$$

for some $f' \in L^p([0, T]; \mathbb{R}^n)$. Moreover, we define $\|f\|_{W^{1,p}} := \|f'\|_{L^p}$ for $f \in W^{1,p}([0, T]; \mathbb{R}^n)$.

Including the three known characterizations of $V^{1,p}([0, T], \mathbb{R}^n)$, we end up with the following five different ways to measure the Riesz-type variation.

Lemma 2.6. *Let $T \in (0, \infty)$, $p \in (1, \infty)$ and \mathbb{R}^n be equipped with the Euclidean norm $|\cdot|$. The space $V^{1,p}([0, T]; \mathbb{R}^n)$ has the following different characterizations:*

$$\begin{aligned} V^{1,p}([0, T]; \mathbb{R}^n) &= \tilde{V}^{1,p}([0, T]; \mathbb{R}^n) = \hat{N}^{1,p}([0, T]; \mathbb{R}^n) \\ &= N^{1,p}([0, T]; \mathbb{R}^n) = W^{1,p}([0, T]; \mathbb{R}^n) \end{aligned}$$

with

$$\|f\|_{V^{1,p}} = \|f\|_{\tilde{V}^{1,p}} = \|f\|_{\hat{N}^{1,p}} = \|f\|_{W^{1,p}} = \|f\|_{N^{1,p}} \quad \text{for } f \in C([0, T]; \mathbb{R}^n).$$

Proof. For $f \in C([0, T]; \mathbb{R}^n)$ and $p \in (1, \infty)$ the identities

$$\|f\|_{V^{1,p}} = \|f\|_{W^{1,p}} = \|f\|_{N^{1,p}}$$

can be found in [FV10, Proposition 1.45] and [Leo09, Theorem 10.55].

Next we observe that

$$\|f\|_{W^{1,p}} = \|f\|_{\tilde{V}^{1,p}}^p \leq \|f\|_{\hat{V}^{1,p}}^p \leq \sup_{\mathcal{P} \subset [0, T]} \sum_{[u, v] \in \mathcal{P}} \frac{\|f\|_{W^{1,p}; [u, v]}^p |v - u|^{p-1}}{|v - u|^{p-1}} \leq \|f\|_{W^{1,p}}^p,$$

where we used [FV10, Theorem 1.44] (see also [FV06, Theorem 1]) for the second estimate and the super-additivity of $\|f\|_{W^{1,p}; [u, v]}^p$ as a function in $(u, v) \in \Delta_T$ in the last one.

As a last step note that

$$\|f\|_{\hat{N}^{1,p}}^p = \sup_{\mathcal{P} \subset [0,T]} \sum_{[u,v] \in \mathcal{P}} \|f\|_{W^{1,p};[u,v]}^p$$

due to [Leo09, Theorem 10.55], which implies

$$\|f\|_{W^{1,p}}^p = \|f\|_{\hat{N}^{1,p}}^p \leq \|f\|_{W^{1,p}}^p$$

using once more the super-additivity of $\|f\|_{W^{1,p};[u,v]}^p$ as a function in $(u, v) \in \Delta_T$. \square

2.2. Characterizations of Riesz-type variation. While Sobolev spaces and Nikolskii spaces coincide with the Riesz-type variation spaces for regularity $\delta = 1$, this is not true anymore for the fractional regularity $\delta \in (0, 1)$. However, the characterizations of Riesz-type variation via q -variation due to Wiener (2.5) and via classical Nikolskii spaces (2.6) still work as we will see in this subsection.

We start by recalling the definition of fractional Sobolev spaces. For $\delta \in (0, 1)$ and $p \in [1, \infty)$ the *fractional Sobolev* (also called *Sobolev-Slobodeckij*) regularity of a function $f \in C([0, T]; E)$ is given by

$$\|f\|_{W^{\delta,p};[s,t]} := \left(\iint_{[s,t]^2} \frac{d(f_u, f_v)^p}{|v - u|^{1+\delta p}} du dv \right)^{\frac{1}{p}}$$

for a subinterval $[s, t] \subset [0, T]$ and we abbreviate $\|\cdot\|_{W^{\delta,p}} := \|\cdot\|_{W^{\delta,p};[0,T]}$. The set of all functions $f \in C([0, T]; E)$ such that $\|f\|_{W^{\delta,p}} < \infty$ is denoted by $W^{\delta,p}([0, T]; E)$.

As an auxiliary result we first need an explicit embedding of Nikolskii regular functions $N^{\delta',p}([0, T]; E)$ into the set of functions with fractional Sobolev regularity $W^{\delta,p}([0, T]; E)$.

Lemma 2.7. *Suppose that (E, d) is a metric space and $T \in (0, \infty)$. Let $p \in [1, \infty)$ and $\delta, \delta' \in (0, 1)$ be such that $\delta' > \delta$. For $f \in N^{\delta',p}([0, T]; E)$ it holds*

$$\|f\|_{W^{\delta,p};[s,t]} \leq \left(\frac{2}{(\delta' - \delta)p} \right)^{\frac{1}{p}} \|f\|_{N^{\delta',p};[s,t]} (t - s)^{\delta' - \delta}$$

for any $s, t \in [0, T]$ with $s < t$. In particular, $N^{\delta',p}([0, T]; E) \subset W^{\delta,p}([0, T]; E)$.

Proof. The fractional Sobolev regularity can be rewritten as

$$\|f\|_{W^{\delta,p};[s,t]}^p = \iint_{[s,t]^2} \frac{d(f_u, f_v)^p}{|v - u|^{1+\delta p}} du dv = 2 \int_0^{t-s} \int_s^{t-h} \frac{d(f_u, f_{u+h})^p}{|h|^{1+\delta p}} du dh$$

for $s, t \in [0, T]$ with $s < t$ and for every $f \in W^{\delta,p}([0, T]; E)$. Since $f \in N^{\delta',p}([0, T]; E)$, one has

$$\int_s^{t-h} d(f_u, f_{u+h})^p du \leq \|f\|_{N^{\delta',p};[s,t]}^p h^{\delta' p}.$$

Therefore, we conclude for $\delta' > \delta > 0$ that

$$\|f\|_{W^{\delta,p};[s,t]}^p \leq 2 \int_0^{t-s} \frac{\|f\|_{N^{\delta',p};[s,t]}^p h^{\delta' p}}{|h|^{1+\delta p}} dh \leq \frac{2}{(\delta' - \delta)p} \|f\|_{N^{\delta',p};[s,t]}^p (t - s)^{(\delta' - \delta)p}$$

for every interval $[s, t] \subset [0, T]$, and thus $N^{\delta',p}([0, T]; E) \subset W^{\delta,p}([0, T]; E)$. \square

The next proposition presents that functions of refined Nikolskii-type regularity are also of finite q -variation and Hölder continuous. It can be seen as a refinement of [FV06, Theorem 2].

For the sake of notational brevity, we use in the following $A_\vartheta \lesssim B_\vartheta$, for a generic parameter ϑ , meaning that $A_\vartheta \leq CB_\vartheta$ for some constant $C > 0$ independent of ϑ .

Proposition 2.8. *Suppose that (E, d) is a metric space and $T \in (0, \infty)$. Let $\delta \in (0, 1)$ and $p \in (1, \infty)$ be such that $\alpha := \delta - 1/p > 0$, and set $q := \frac{1}{\delta}$.*

(1) *If $f \in N^{\delta,p}([0, T]; E)$, then $f \in C^{\alpha\text{-Hö}}([0, T]; E)$ and*

$$d(f_s, f_t) \lesssim \|f\|_{N^{\delta,p};[s,t]}(t-s)^{\delta-\frac{1}{p}}, \quad [s, t] \subset [0, T].$$

(2) *The q -variation of any $f \in \hat{N}^{\delta,p}([0, T]; E)$ can be estimated by*

$$\|f\|_{q\text{-var};[s,t]} \lesssim \|f\|_{\hat{N}^{\delta,p};[s,t]}(t-s)^\alpha, \quad [s, t] \subset [0, T],$$

and one has $\hat{N}^{\delta,p}([0, T]; E) \subset C^{\alpha\text{-Hö}}([0, T]; E)$ and

$$\hat{N}^{\delta,p}([0, T]; E) \subset C^{q\text{-var}}([0, T]; E).$$

Proof. (1) Choose $\gamma < \delta$ such that $\gamma - 1/p > 0$. Because $f \in N^{\delta,p}([0, T]; E)$, Lemma 2.7 yields $f \in W^{\gamma,p}([0, T]; E)$ and we have

$$\|f\|_{W^{\gamma,p};[s,t]}^p = F_{s,t} := \iint_{[s,t]^2} \left(\frac{d(f_u, f_v)}{|v-u|^{1/p+\gamma}} \right)^p du dv, \quad [s, t] \subset [0, T].$$

Applying the Garsia–Rodemich–Rumsey inequality with $\Psi(\cdot) = (\cdot)^p$ and $p(\cdot) = (\cdot)^{1/p+\gamma}$ gives

$$d(f_s, f_t) \leq 8 \int_0^{t-s} \left(\frac{F_{s,t}}{u^2} \right)^{\frac{1}{p}} dp(u) = \frac{8}{(\gamma - 1/p)} \|f\|_{W^{\gamma,p};[s,t]}(t-s)^{\gamma-\frac{1}{p}},$$

using $\gamma - \frac{1}{p} > 0$; see for instance [FV10, Theorem A.1] for a version of the Garsia–Rodemich–Rumsey lemma suitable for functions with values in a metric space. Furthermore, Lemma 2.7 yields

$$\begin{aligned} (2.7) \quad d(f_s, f_t) &\leq \frac{8}{(\gamma - 1/p)} \left(\frac{2}{(\delta - \gamma)p} \right)^{\frac{1}{p}} \|f\|_{N^{\delta,p};[s,t]}(t-s)^{\delta-\gamma}(t-s)^{\gamma-\frac{1}{p}} \\ &\lesssim \|f\|_{N^{\delta,p};[s,t]}(t-s)^{\delta-\frac{1}{p}}, \end{aligned}$$

which gives $f \in C^{(\delta-1/p)\text{-Hö}}([0, T]; E)$.

(2) Assuming $f \in \hat{N}^{\delta,p}([0, T]; E)$ the estimate (2.7) leads to

$$d(f_s, f_t) \lesssim \|f\|_{\hat{N}^{\delta,p};[s,t]}(t-s)^{\delta-\frac{1}{p}}, \quad [s, t] \subset [0, T].$$

Recalling $\alpha = \delta - 1/p > 0$ and $q = \frac{1}{\delta}$, we note that

$$\omega(s, t) := \|f\|_{\hat{N}^{\delta,p};[s,t]}^q (t-s)^{\alpha q}, \quad 0 \leq s \leq t \leq T,$$

is super-additive. Indeed, since $\|f\|_{\hat{N}^{\delta,p};[s,t]}^p$ and $|t-s|$ are super-additive as functions in $(s, t) \in \Delta_T$ and $q/p + 1 - 1/\delta p \geq 1$, Remark 2.3 ensures the super-additivity of ω .

Hence, by the super-additivity of ω we deduce that

$$\|f\|_{q\text{-var};[s,t]}^q \leq C\omega(s, t) = C\|f\|_{\hat{N}^{\delta,p};[s,t]}^q |t-s|^{\alpha q},$$

for some constant $C > 0$ depending only on δ and p .

In particular, we have proven that $\hat{N}^{\delta,p}([0, T]; E) \subset C^{\alpha\text{-Hö}l}([0, T]; E)$ and

$$\hat{N}^{\delta,p}([0, T]; E) \subset C^{q\text{-var}}([0, T]; E). \quad \square$$

Remark 2.9. Proposition 2.8 (2) does not hold for $\|\cdot\|_{\hat{N}^{\delta,p};[s,t]}$ replaced by $\|\cdot\|_{N^{\delta,p};[s,t]}$; see Remark 2.13 below.

Remark 2.10. Alternatively to the given proofs of Lemma 2.7 and Proposition 2.8, one could use the abstract Kuratowski embedding to extend the known Besov embeddings from Banach spaces to general metric spaces and then proceed further as presented above. For example note, if $(E, \|\cdot\|)$ is a Banach space, then classical Besov embeddings lead to

$$\|f_t - f_s\| \leq \sup_{|t-s| \geq h > 0} \left(\frac{\|f_{s+h} - f_s\|}{|h|^{\delta-1/p}} \right) |t-s|^{\delta-1/p} \leq C \|f\|_{N^{\delta,p};[s,t]} |t-s|^{\delta-1/p}$$

for every $f \in N^{\delta,p}([0, T]; E)$, $\delta \in (0, 1)$, $p \in (1, \infty)$ such that $\delta > 1/p$, and some constant $C > 0$; cf. [Sim90, Theorem 10]. However, we prefer here to give direct proofs.

On the other hand, the embedding $N^{\delta,p}([0, T]; E) \subset W^{\delta,p}([0, T]; E)$ does not hold true, which prevents us from deducing Proposition 2.8 as a corollary of [FV06, Theorem 2]. Hence, the elaborated embedding of Lemma 2.7 is essential to obtain Proposition 2.8.

The next theorem is the main result of the first part: the characterization of Riesz-type variation via $\|\cdot\|_{\tilde{V}^{\delta,p}}$ and $\|\cdot\|_{\hat{N}^{\delta,p}}$.

Theorem 2.11. *Let $T \in (0, \infty)$ and (E, d) be a metric space. Suppose that $\delta \in (0, 1)$ and $p \in (1, \infty)$ such that $\delta > 1/p$. Then, $\|\cdot\|_{V^{\delta,p}}$, $\|\cdot\|_{\tilde{V}^{\delta,p}}$ and $\|\cdot\|_{\hat{N}^{\delta,p}}$ are equivalent, that is,*

$$\|f\|_{V^{\delta,p}} \lesssim \|f\|_{\tilde{V}^{\delta,p}} \lesssim \|f\|_{\hat{N}^{\delta,p}} \lesssim \|f\|_{V^{\delta,p}}$$

for every function $f \in C([0, T]; E)$, and thus

$$V^{\delta,p}([0, T]; E) = \tilde{V}^{\delta,p}([0, T]; E) = \hat{N}^{\delta,p}([0, T]; E).$$

Proof. For a function $f \in C([0, T]; E)$ and an interval $[s, t] \subset [0, T]$ recall that

$$\|f\|_{N^{\delta,p};[s,t]} = \left(\sup_{|t-s| \geq h > 0} h^{-\delta p} \int_s^{t-h} d(f_u, f_{u+h})^p du \right)^{\frac{1}{p}}.$$

Let us fix $h \in (0, t-s]$ and take a partition $\mathcal{P}(h) := \{[t_i, t_{i+1}] : s = t_0 < \dots < t_M = t-h\}$ such that

$$|t_M - t_{M-1}| \leq h \quad \text{and} \quad |t_{i+1} - t_i| = h \quad \text{for } i = 0, \dots, M-2, \quad M \in \mathbb{N}.$$

Since $\sup_{u \in [t_i, t_{i+1}]} d(f_u, f_{u+h})^p \leq \|f\|_{1/\delta\text{-var}; [t_i, t_{i+2}]}^p$ for $i = 0, \dots, M-1$ with $t_{M+1} := t-h$, we observe that

$$\begin{aligned} \int_s^{t-h} |d(f_u, f_{u+h})|^p du &= \sum_{i=0}^{M-1} \int_{t_i}^{t_{i+1}} d(f_u, f_{u+h})^p du \\ &\leq \sum_{i=0}^{M-1} \sup_{u \in [t_i, t_{i+1}]} d(f_u, f_{u+h})^p (t_{i+1} - t_i) \\ &\leq \frac{1}{2} (2h)^{\delta p} \sum_{i=0}^{M-1} \frac{\|f\|_{1/\delta\text{-var}; [t_i, t_{i+2}]}^p}{(2h)^{\delta p-1}} \lesssim h^{\delta p} \|f\|_{\tilde{V}^{\delta,p}; [s,t]}^p, \end{aligned}$$

which implies $\|f\|_{N^{\delta,p}; [s,t]}^p \lesssim \|f\|_{\tilde{V}^{\delta,p}; [s,t]}^p$. Therefore, the super-additivity of $\|f\|_{\tilde{V}^{\delta,p}; [s,t]}^p$ as a function in $(s, t) \in \Delta_T$ reveals

$$\|f\|_{\hat{N}^{\delta,p}} \lesssim \|f\|_{\tilde{V}^{\delta,p}}.$$

For the converse inequality Proposition 2.8 gives

$$\|f\|_{\frac{1}{\delta}\text{-var}; [u,v]} \lesssim \|f\|_{\hat{N}^{\delta,p}; [u,v]} |v-u|^{\delta-\frac{1}{p}}, \quad 0 \leq u < v \leq T,$$

for $\delta \in (0, 1)$ and $p \in (1, \infty)$ such that $\delta > 1/p$, which leads to

$$\begin{aligned} \|h\|_{\tilde{V}^{\delta,p}}^p &= \sup_{\mathcal{P} \subset [0,T]} \sum_{[u,v] \in \mathcal{P}} \frac{\|h\|_{\frac{1}{\delta}\text{-var}; [u,v]}^p}{|v-u|^{\delta p-1}} \\ &\leq \sup_{\mathcal{P} \subset [0,T]} \sum_{[u,v] \in \mathcal{P}} \frac{\|h\|_{\hat{N}^{\delta,p}; [u,v]}^p |v-u|^{\delta p-1}}{|v-u|^{\delta p-1}} \leq \|h\|_{\hat{N}^{\delta,p}}^p, \end{aligned}$$

where we applied the super-additivity of $\|f\|_{\hat{N}^{\delta,p}; [s,t]}^p$ as a function in $(s, t) \in \Delta_T$.

It remains to show

$$\|f\|_{V^{\delta,p}} \lesssim \|f\|_{\tilde{V}^{\delta,p}} \lesssim \|f\|_{V^{\delta,p}}, \quad f \in C([0, T], E).$$

The first inequality follows immediately from the definitions and the observation

$$d(f_u, f_v)^p \leq \|f\|_{1/\delta\text{-var}; [u,v]}^p, \quad [u, v] \subset [0, T].$$

The second inequality can be deduced from Proposition 2.2, which gives the estimate

$$\|f\|_{1/\delta\text{-var}; [u,v]}^p \leq \|f\|_{V^{\delta,p}; [u,v]}^p |v-u|^{\delta p-1},$$

and the super-additivity of $\|f\|_{V^{\delta,p}; [s,t]}^p$ as a function in $(s, t) \in \Delta_T$. \square

As a next step we briefly want to understand how the set $V^{\delta,p}([0, T]; E)$ of functions with finite Riesz-type variation are related to other types of measuring the regularity of functions. The characterization of Riesz-type variation in terms of Nikolskii regularity allows us to deduce the following result connecting the set $V^{\delta,p}([0, T]; E)$ with the notion of classical fractional Sobolev and Nikolskii regularity.

Corollary 2.12. *Let $T \in (0, \infty)$ and (E, d) be a metric space. If $\delta \in (0, 1)$ and $p \in (1, \infty)$ such that $\delta > 1/p$, then one has the inclusions*

$$(2.8) \quad W^{\delta,p}([0, T]; E) \subset V^{\delta,p}([0, T]; E) \subset N^{\delta,p}([0, T]; E)$$

and

$$N^{\delta+\epsilon,p}([0, T]; E) \subset \hat{N}^{\delta,p}([0, T]; E) \subset N^{\delta,p}([0, T]; E)$$

for $\epsilon \in (0, 1 - \delta)$.

Proof. For the first embedding let $f \in W^{\delta,p}([0, T]; E)$. Applying Theorem 2.11 and [Sim90, Theorem 11], which can be extended to general metric spaces by Kuratowski's embedding theorem, we get

$$\begin{aligned} \|f\|_{V^{\delta,p}}^p &\lesssim \|f\|_{\hat{N}^{\delta,p}}^p = \sup_{\mathcal{P} \subset [0, T]} \sum_{[s, t] \in \mathcal{P}} \|f\|_{N^{\delta,p}; [s, t]}^p \\ &\lesssim \sup_{\mathcal{P} \subset [0, T]} \sum_{[s, t] \in \mathcal{P}} \|f\|_{W^{\delta,p}; [s, t]}^p \leq \|f\|_{W^{\delta,p}}^p. \end{aligned}$$

For the second embedding let $f \in N^{\delta,p}([0, T]; E)$ and we apply again Theorem 2.11 to obtain

$$\|f\|_{N^{\delta,p}}^p \leq \|f\|_{\hat{N}^{\delta,p}}^p \lesssim \|f\|_{V^{\delta,p}}^p.$$

The first embedding for the refined Nikolskii-type space $\hat{N}^{\delta+\epsilon,p}$ is a consequence of Theorem 2.11 and the embedding

$$N^{\delta+\epsilon,p}([0, T]; E) \subset W^{\delta,p}([0, T]; E) \subset V^{\delta,p}([0, T]; E),$$

where we used Lemma 2.7 and (2.8).

The second embedding for the refined Nikolskii-type space $\hat{N}^{\delta,p}$ follows directly from its definition. \square

Remark 2.13. Both embeddings are proper embeddings, which means in both cases the equality does not hold.

Indeed, an example of a set of functions which are included in $V^{1/2+H,2}([0, T]; \mathbb{R})$ but not in $W^{1/2+H,2}([0, T]; \mathbb{R})$ consists of the Cameron–Martin space of a fractional Brownian motion with Hurst index $H \in (0, 1/2)$; see [FV06] and [FH14, Section 11] and the references therein.

To see that the second embedding is not an equality, we recall that the sample paths of a Brownian motion belong in the Nikolskii space $N^{1/2,p}([0, T]; \mathbb{R})$ for $p \in (2, \infty)$, which was proven by [Roy93] (cf. [Ros09, Proposition 1]). However, it is also well known that sample paths of a Brownian motion are not contained in $C^{2\text{-var}}([0, T]; \mathbb{R})$. In other words, they cannot be contained in $V^{1/2,p}([0, T]; \mathbb{R})$ for $p \in (2, \infty)$ since this is a subset of $C^{2\text{-var}}([0, T]; \mathbb{R})$ by Proposition 2.2.

2.3. Separability considerations. In order to embed the Riesz-type variation spaces into separable Banach spaces, we need to restrict the general metric space E and focus here on the case $E = \mathbb{R}^n$ equipped with the Euclidean norm $|\cdot|$. As usual $|\cdot|$ induces the metric $d(x, y) := |y - x|$ for $x, y \in \mathbb{R}^n$ and thus $\|\cdot\|_{V^{\delta,p}}$, $\|\cdot\|_{\hat{V}^{\delta,p}}$ and $\|\cdot\|_{\hat{N}^{\delta,p}}$ become semi-norms, which can be easily modified to proper norms by adding for instance the Euclidean norm of the functions evaluated at zero; cf. (2.9). An immediate consequence of Theorem 2.11 is the following equivalence.

Corollary 2.14. *Let $T \in (0, \infty)$ and \mathbb{R}^n be equipped with the Euclidean norm $|\cdot|$. If $\delta \in (0, 1)$ and $p \in (1, \infty)$ are such that $\delta > 1/p$, then the semi-norms $\|\cdot\|_{V^{\delta,p}}$, $\|\cdot\|_{\hat{V}^{\delta,p}}$, and $\|\cdot\|_{\hat{N}^{\delta,p}}$ are equivalent.*

In order to turn $C^{\delta\text{-H\"{o}l}}([0, T]; \mathbb{R}^n)$ and $C^{p\text{-var}}([0, T]; \mathbb{R}^n)$ into Banach spaces, one usually introduces the norms

$$(2.9) \quad |f(0)| + \|f\|_{\delta\text{-H\"{o}l}} \quad \text{and} \quad |g(0)| + \|g\|_{p\text{-var}}$$

for $f \in C^{\delta\text{-H\"{o}l}}([0, T]; \mathbb{R}^n)$ and $g \in C^{p\text{-var}}([0, T]; \mathbb{R}^n)$, respectively. These Banach spaces are not separable; see [FV10, Theorem 5.25].

To restore the separability, one can consider the closure of smooth paths. Let $C^\infty([0, T]; \mathbb{R}^n)$ be the space of smooth functions $f \in C([0, T]; \mathbb{R}^n)$. For $\delta \in (0, 1)$ and $p \in (1, \infty)$ we define

$$C^{0, \delta\text{-H\"{o}l}}([0, T]; \mathbb{R}^n) := \overline{C^\infty([0, T]; \mathbb{R}^n)}^{\|\cdot\|_{\delta\text{-H\"{o}l}}}$$

and

$$C^{0, p\text{-var}}([0, T]; \mathbb{R}^n) := \overline{C^\infty([0, T]; \mathbb{R}^n)}^{\|\cdot\|_{p\text{-var}}}.$$

These two Banach spaces are separable and one has the obvious embeddings

$$C^{0, \delta\text{-H\"{o}l}}([0, T]; \mathbb{R}^n) \subset C^{\delta\text{-H\"{o}l}}([0, T]; \mathbb{R}^n)$$

and

$$C^{0, p\text{-var}}([0, T]; \mathbb{R}^n) \subset C^{p\text{-var}}([0, T]; \mathbb{R}^n).$$

The Riesz-type variation space $V^{\delta, p}([0, T]; \mathbb{R}^n)$ can be embedded into

$$C^{0, \alpha\text{-H\"{o}l}}([0, T]; \mathbb{R}^n)$$

and

$$C^{0, p\text{-var}}([0, T]; \mathbb{R}^n).$$

Lemma 2.15. *Let $T \in (0, \infty)$ and \mathbb{R}^n be equipped with the Euclidean norm $|\cdot|$. If $\delta \in (0, 1)$ and $p \in (1, \infty)$ are such that $\delta > 1/p$, then one has the embeddings*

$$V^{\delta, p}([0, T]; \mathbb{R}^n) \subset C^{0, p\text{-var}}([0, T]; \mathbb{R}^n) \quad \text{and} \quad V^{\delta, p}([0, T]; \mathbb{R}^n) \subset C^{0, \alpha\text{-H\"{o}l}}([0, T]; \mathbb{R}^n)$$

for $\alpha \in (0, \delta - 1/p)$.

Proof. For $f \in V^{\delta, p}([0, T]; \mathbb{R}^n)$ and $\delta > 1/p$ we apply Lemma 2.4 to obtain

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \sup_{\mathcal{P} \subset [0, T], |\mathcal{P}| < \varepsilon} \sum_{[s, t] \in \mathcal{P}} |f_t - f_s|^p &\lesssim \lim_{\varepsilon \rightarrow 0} \sup_{\mathcal{P} \subset [0, T], |\mathcal{P}| < \varepsilon} \sum_{[s, t] \in \mathcal{P}} \|f\|_{V^{\delta, p}; [s, t]}^p |t - s|^{\delta p - 1} \\ &\leq \|f\|_{V^{\delta, p}}^p \lim_{\varepsilon \rightarrow 0} \varepsilon^{\delta p - 1} = 0, \end{aligned}$$

where $|\mathcal{P}|$ denotes the mesh size of the partition \mathcal{P} , and thus $f \in C^{0, p\text{-var}}([0, T]; \mathbb{R}^n)$ due to Wiener's characterization of $C^{0, p\text{-var}}([0, T]; \mathbb{R}^n)$; see [FV10, Theorem 5.31].

Using Wiener's characterization of $C^{0, \alpha\text{-H\"{o}l}}([0, T]; \mathbb{R}^n)$ for $\alpha \in (0, \delta - 1/p)$, we get the second embedding because of

$$\lim_{\varepsilon \rightarrow 0} \sup_{[s, t] \subset [0, T], |t - s| < \varepsilon} \frac{|f_t - f_s|}{|t - s|^\alpha} \leq \lim_{\varepsilon \rightarrow 0} \sup_{[s, t] \subset [0, T], |t - s| < \varepsilon} \left(\frac{|f_t - f_s|^p}{|t - s|^{\delta p - 1}} \right)^{1/p} \varepsilon^{\delta - 1/p - \alpha} = 0$$

for $f \in V^{\delta, p}([0, T]; \mathbb{R}^n)$. □

3. CONTINUITY OF THE ITÔ–LYONS MAP

The dynamics of a controlled differential equation driven by a path $X: [0, T] \rightarrow \mathbb{R}^n$ of finite q -variation is formally given by

$$(3.1) \quad dY_t = V(Y_t) dX_t, \quad Y_0 = y_0, \quad t \in [0, T],$$

where $y_0 \in \mathbb{R}^m$ is the initial condition, $V: \mathbb{R}^m \rightarrow \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ is a smooth enough vector field and $T \in (0, \infty)$. Here $\mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ denotes the space of linear operators from \mathbb{R}^n to \mathbb{R}^m . If the driving signal $X \in C^{p\text{-var}}([0, T]; \mathbb{R}^n)$ for $p \in [1, 2)$, Lyons [Lyo94] first established the existence and uniqueness of a solution Y to the equation (3.1). Moreover, he proved that the Itô–Lyons map is a locally Lipschitz continuous map with respect to the q -variation topology. In order to restore the continuity for more irregular paths X , say $X \in C^{q\text{-var}}([0, T]; \mathbb{R}^n)$ for an arbitrarily large $q < \infty$, Lyons introduced the notion of rough paths in his seminal paper [Lyo98]; see Subsection 3.2. Based on Lyons’ estimate, one can deduce the local Lipschitz continuity of the Itô–Lyons map with respect to a Hölder topology; see for example [Fri05].

The aim of this section is to particularly unify these two results by establishing the local Lipschitz continuity of the Itô–Lyons map on Riesz-type variation spaces. For this purpose we combine Lyons’ estimates with our characterization of Riesz-type variation in terms to q -variation to deduce the locally Lipschitz continuity of the Itô–Lyons map with respect to an inhomogeneous Riesz-type distance. See Proposition 3.1 for the continuity result in the regime of bounded variation paths. For the result in the general rough path setting we refer to Theorem 3.3 and Corollaries 3.5 and 3.8.

To quantify the regularity of the vector field V in the controlled differential equation (3.1), we introduce for $\alpha > 0$ the space $\text{Lip}^\alpha := \text{Lip}^\alpha(\mathbb{R}^m; \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m))$ in the sense of E. Stein; cf. [FV10, Definition 10.2]. For $\alpha > 0$ and

$$[\alpha] := \max\{n \in \mathbb{N} : n \leq \alpha\}$$

the space Lip^α consists of all maps $V: \mathbb{R}^m \rightarrow \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ such that V is $[\alpha]$ -times continuously differentiable with $(\alpha - [\alpha])$ -Hölder continuous partial derivatives of order $[\alpha]$ (or with continuous partial derivatives of order α in the case $\alpha = [\alpha]$). On the space Lip^α we introduce the usual norm $\|\cdot\|_{\text{Lip}^\alpha}$ and further denote the supremum norm by $\|\cdot\|_\infty$. For the supremum norm on $C([0, T]; \mathbb{R}^n)$ we write $\|\cdot\|_{\infty; [0, T]} := \sup_{0 \leq t \leq T} \|\cdot\|$.

3.1. Continuity w.r.t. $\tilde{V}^{1,p}$. In this subsection we derive the local Lipschitz continuity of the solution map on the Riesz-type variation spaces $V^{1,p}([0, T]; \mathbb{R}^n)$. To that end the equivalent characterization of $V^{1,p}([0, T]; \mathbb{R}^n)$ given by $\tilde{V}^{1,p}([0, T]; \mathbb{R}^n)$ turns out to be particularly convenient. The solution map Φ is defined by

$$(3.2) \quad \Phi: \mathbb{R}^m \times \text{Lip}^1 \times \tilde{V}^{1,p}([0, T]; \mathbb{R}^n) \rightarrow \tilde{V}^{1,p}([0, T]; \mathbb{R}^m) \quad \text{via} \quad \Phi(y_0, V, X) := Y,$$

where Y denotes the solution to the integral equation

$$(3.3) \quad Y_t = y_0 + \int_0^t V(Y_s) dX_s, \quad t \in [0, T].$$

First notice that the integral appearing in equation (3.3) can be defined as a classical Riemann–Stieltjes integral with respect to bounded variation functions because

of the embedding $\tilde{V}^{1,p}([0, T]; \mathbb{R}^n) \subset C^{1\text{-var}}([0, T]; \mathbb{R}^n)$ for all $p \in (1, \infty)$ due to Proposition 2.2 and Lemma 2.6.

Proposition 3.1. *For $X \in \tilde{V}^{1,p}([0, T]; \mathbb{R}^n)$ with $p \in (1, \infty)$, $V \in \text{Lip}^1$ and every initial condition $y_0 \in \mathbb{R}^m$, the controlled differential equation (3.3) has a unique solution $Y \in \tilde{V}^{1,p}([0, T]; \mathbb{R}^n)$ and the solution map Φ as defined in (3.2) is locally Lipschitz continuous.*

More precisely, for $y_0^i \in \mathbb{R}^m$, $X^i \in \tilde{V}^{1,p}([0, T]; \mathbb{R}^n)$, $V^i \in \text{Lip}^1$ such that

$$\|X^i\|_{\tilde{V}^{1,p}} \leq b \quad \text{and} \quad \|V^i\|_{\text{Lip}^1} \leq l, \quad i = 1, 2,$$

for some $b, l > 0$ and corresponding solution Y^i , there exist a constant $C = C(b, l, p) \geq 1$ such that

$$\|Y^1 - Y^2\|_{\tilde{V}^{1,p}} \leq C(\|V^1 - V^2\|_\infty + |y_0^1 - y_0^2| + \|X^1 - X^2\|_{\tilde{V}^{1,p}}).$$

Proof. Since $X^i \in \tilde{V}^{1,p}([0, T]; \mathbb{R}^n)$, X^i is in particular of bounded variation and thus the integral equation (3.3) is well defined and admits a unique solution $Y^i \in C^{1\text{-var}}([0, T]; \mathbb{R}^n)$ for each $i = 1, 2$. Moreover, for every subinterval $[s, t] \subset [0, T]$ the local Lipschitz continuity of the solution map Φ in 1-variation, cf. [FV10, Theorem 3.18] and [FV10, Remark 3.19], yields

$$\begin{aligned} & \|Y^1 - Y^2\|_{1\text{-var};[s,t]} \\ & \leq 2 \exp(3bl) (|y_s^1 - y_s^2| c(s, t) + \|V^1 - V^2\|_\infty c(s, t) + l \|X^1 - X^2\|_{1\text{-var};[s,t]}), \end{aligned}$$

where $c(s, t)$ can be chosen such that

$$\|X^1\|_{1\text{-var};[s,t]} + \|X^2\|_{1\text{-var};[s,t]} \leq c(s, t) \lesssim k(s, t)(t - s)^{1-1/p}$$

with $k(s, t) := \|X^1\|_{\tilde{V}^{1,p};[s,t]} + \|X^2\|_{\tilde{V}^{1,p};[s,t]}$. Dividing both sides by $|t - s|^{1-1/p}$ and taking them to the power p leads to

$$\begin{aligned} & \frac{\|Y^1 - Y^2\|_{1\text{-var};[s,t]}^p}{|t - s|^{p-1}} \\ & \lesssim \exp(3blp) \left(|y_s^1 - y_s^2|^{lp} k(s, t)^p + \|V^1 - V^2\|_\infty^p k(s, t)^p + l^p \frac{\|X^1 - X^2\|_{1\text{-var};[s,t]}^p}{|t - s|^{p-1}} \right). \end{aligned}$$

From this inequality we deduce, by summing over a partition of $[0, T]$ and taking then the supremum over all partitions, that

$$\|Y^1 - Y^2\|_{\tilde{V}^{1,p}} \lesssim \exp(3vl) (\|V^1 - V^2\|_\infty b + \|Y^1 - Y^2\|_{\infty;[0,T]} bl + l \|X^1 - X^2\|_{\tilde{V}^{1,p}}),$$

where we used the super-additivity of $k(s, t)^p$ and $k(0, T) \leq 2b$. Finally, $\|Y^1 - Y^2\|_{\infty;[0,T]}$ can be estimated by [FV10, Theorem 3.15] to complete the proof. \square

Remark 3.2. An immediate consequence of Lemma 2.6 is that the local Lipschitz continuity as stated in Proposition 3.1 of the solution map Φ given by (3.2) also holds with respect to the (equivalent) Sobolev or Nikolskii metric.

3.2. Continuity w.r.t. general Riesz-type variation. In order to give a meaning to the controlled differential equation (3.1) for driving signals X which are not of bounded variation, we introduce here the basic framework of rough path theory. For more comprehensive monographs about rough path theory we refer to [LQ02, FV10, FH14], and for the convenience of the reader the following definitions are mainly borrowed from [FV10].

As already explained in the introduction, a rough path takes values in the metric space $(G^N(\mathbb{R}^n), d_{cc})$ and not “only” in the Euclidean space $(\mathbb{R}^n, \|\cdot\|)$. Let us recall the basic ingredients:

For $N \in \mathbb{N}$ and a path $Z \in C^{1\text{-var}}(\mathbb{R}^n)$ its N -step signature is given by

$$S_N(Z)_{s,t} := \left(1, \int_{s < u < t} dZ_u, \dots, \int_{s < u_1 < \dots < u_N < t} dZ_{u_1} \otimes \dots \otimes dZ_{u_N} \right) \\ \in T^N(\mathbb{R}^n) := \bigoplus_{k=0}^N (\mathbb{R}^n)^{\otimes k},$$

where $(\mathbb{R}^n)^{\otimes k}$ denotes the k -tensor space of \mathbb{R}^n and $\mathbb{R}^{\otimes 0} := \mathbb{R}$. We note that $T^N(\mathbb{R}^n)$ is an algebra (“level- N truncated tensor algebra”) under the tensor product \otimes . The corresponding space of all these lifted paths is the step- N free nilpotent group (w.r.t. \otimes)

$$G^N(\mathbb{R}^n) := \{S_N(Z)_{0,T} : Z \in C^{1\text{-var}}([0, T]; \mathbb{R}^n)\} \subset T^N(\mathbb{R}^n).$$

For every $g \in G^N(\mathbb{R}^n)$ the so-called “Carnot–Carathéodory norm”

$$\|g\|_{cc} := \inf \left\{ \int_0^T \|d\gamma_s\| : \gamma \in C^{1\text{-var}}([0, T]; \mathbb{R}^n) \text{ and } S_N(\gamma)_{0,T} = g \right\},$$

where $\int_0^T \|d\gamma_s\|$ is the length of γ based on the Euclidean distance, is finite and the infimum is attained; see [FV10, Theorem 7.32]. This leads to the *Carnot–Carathéodory metric* d_{cc} via

$$d_{cc}(g, h) := \|g^{-1} \otimes h\|_{cc}, \quad g, h \in G^N(\mathbb{R}^n),$$

where g^{-1} is the inverse of g in the sense $g^{-1} \otimes g = 1$; see [FV10, Proposition 7.36 and Definition 7.41]. Hence, $(G^N(\mathbb{R}^n), d_{cc})$ is a metric space.

The space of all *weakly geometric rough paths* of finite q -variation is then given by

$$\Omega^q := C^{q\text{-var}}([0, T]; G^{\lfloor q \rfloor}(\mathbb{R}^n)) := \left\{ \mathbf{X} \in C([0, T]; G^{\lfloor q \rfloor}(\mathbb{R}^n)) : \|\mathbf{X}\|_{q\text{-var}} < \infty \right\},$$

where $\|\cdot\|_{q\text{-var}}$ is the q -variation with respect to the metric space $(G^{\lfloor q \rfloor}(\mathbb{R}^n), d_{cc})$ as defined in (2.1) and $\lfloor q \rfloor := \max\{n \in \mathbb{N} : n \leq q\}$. Note that $\|\cdot\|_{q\text{-var}}$ on Ω^q is commonly called the homogeneous (rough path) norm since it is homogeneous with respect to the dilation map on $T^{\lfloor q \rfloor}(\mathbb{R}^n)$; cf. [FV10, Definition 7.13].

Coming back to the controlled differential equation (3.1), we first need to introduce a solution concept suitable for this equation given the driving signal is now a weakly geometric rough path. Let $V: \mathbb{R}^m \rightarrow \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ be a sufficiently smooth vector field and $y_0 \in \mathbb{R}^m$ be some initial condition. For a weakly geometric rough path $\mathbf{X} \in C^{q\text{-var}}([0, T]; G^{\lfloor q \rfloor}(\mathbb{R}^n))$ we call $Y \in C([0, T]; \mathbb{R}^m)$ a solution to the controlled differential equation (also called rough differential equation)

$$(3.4) \quad dY_t = V(Y_t) d\mathbf{X}_t, \quad Y_0 = y_0, \quad t \in [0, T],$$

if there exist a sequence $(X^n) \subset C^{1\text{-var}}([0, T]; \mathbb{R}^n)$ such that

$$\lim_{n \rightarrow \infty} \sup_{0 \leq s < t \leq T} d_{cc}(S_{\lfloor q \rfloor}(X^n)_{s,t}, \mathbf{X}_{s,t}) = 0, \quad \sup_n \|S_{\lfloor q \rfloor}(X^n)\|_{q\text{-var}; [0, T]} < \infty,$$

and the corresponding solutions Y^n to equation (3.3) converge uniformly on $[0, T]$ to Y as n tends to ∞ ; cf. [FV10, Definition 10.17].

On the space $\Omega^q = C^{q\text{-var}}([0, T]; G^N(\mathbb{R}^n))$ the classical way to restore to the continuity of the solution map associated to a controlled differential equation (3.4) (also called the Itô–Lyons map) is to introduce the *inhomogeneous variation distance*

$$\rho_{q\text{-var}}(\mathbf{X}^1, \mathbf{X}^2) := \max_{k=1, \dots, N} \rho_{q\text{-var}; [0, T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)$$

for $\mathbf{X}^1, \mathbf{X}^2 \in C^{q\text{-var}}([0, T]; G^N(\mathbb{R}^n))$ and $q \in [1, \infty)$, with

$$\rho_{q\text{-var}; [s, t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2) := \sup_{\mathcal{P} \subset [s, t]} \left(\sum_{[u, v] \in \mathcal{P}} |\pi_k(\mathbf{X}_{u, v}^1 - \mathbf{X}_{u, v}^2)|^{\frac{q}{k}} \right)^{\frac{k}{q}}, \quad [s, t] \subset [0, T],$$

for $k = 1, \dots, N$, where $\pi_k: T^N(\mathbb{R}^n) \rightarrow (\mathbb{R}^n)^{\otimes k}$ is the projection to the k th tensor level and each tensor level $(\mathbb{R}^n)^{\otimes k}$ is equipped with the Euclidean structure. Here we recall that $\mathbf{X}_{u, v} := \mathbf{X}_u^{-1} \otimes \mathbf{X}_v$. Note that the distance $\rho_{q\text{-var}}$ is not homogeneous anymore with respect to the dilation map on $T^N(\mathbb{R}^n)$ as indicated by its name.

In the seminal paper [Lyo98], Lyons showed that the solution map Φ given by

$$\Phi: \mathbb{R}^m \times \text{Lip}^\gamma \times C^{1/\delta\text{-var}}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n)) \rightarrow C^{1/\delta\text{-var}}([0, T]; \mathbb{R}^m)$$

via

$$\Phi(y_0, V, \mathbf{X}) := Y,$$

where Y denotes the solution to equation (3.4) given the input (y_0, V, \mathbf{X}) , is local Lipschitz continuity with respect to the inhomogeneous variation distance $\rho_{1/\delta\text{-var}}$ for any finite regularity $1/\delta > 1$.

In the spirit of our characterization (2.5) of the Riesz-type variation we introduce for $\delta \in (0, 1)$ and $p \in [1/\delta, \infty)$ the inhomogeneous *mixed Hölder-variation distance* by

$$\rho_{\tilde{V}^{\delta, p}}(\mathbf{X}^1, \mathbf{X}^2) := \max_{k=1, \dots, N} \rho_{\tilde{V}^{\delta, p}; [0, T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2),$$

for $\mathbf{X}^1, \mathbf{X}^2 \in \tilde{V}^{\delta, p}([0, T]; G^N(\mathbb{R}^n))$ and for $k = 1, \dots, N$, we set

$$\rho_{\tilde{V}^{\delta, p}; [s, t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2) := \sup_{\mathcal{P} \subset [s, t]} \left(\sum_{[u, v] \in \mathcal{P}} \frac{\rho_{1/\delta\text{-var}; [u, v]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}}}{|u - v|^{\delta p - 1}} \right)^{\frac{k}{p}}, \quad [s, t] \subset [0, T].$$

Furthermore, we define the *Riesz-type geometric rough path space* by

$$\begin{aligned} \Omega^{\delta, p} &:= V^{\delta, p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n)) = \tilde{V}^{\delta, p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n)) \\ &= \hat{N}^{\delta, p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n)), \end{aligned}$$

for $\delta \in (0, 1)$, $p \in (1/\delta, \infty)$. The identities hold due to Theorem 2.11 since $G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n)$ is a metric space equipped with the Carnot–Carathéodory metric d_{cc} .

Relying on the equivalent characterization of Riesz-type variation in terms of q -variation (cf. Theorem 2.11) and on the inhomogeneous mixed Hölder-variation distances for Riesz-type geometric rough paths, the local Lipschitz continuity of the Itô–Lyons map Φ given by

$$(3.5) \quad \Phi: \mathbb{R}^m \times \text{Lip}^\gamma \times \tilde{V}^{\delta, p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n)) \rightarrow \tilde{V}^{\delta, p}([0, T]; \mathbb{R}^m) \quad \text{via} \quad \Phi(y_0, V, \mathbf{X}) := Y,$$

where Y denotes the solution to equation (3.4) given the input (y_0, V, \mathbf{X}) , can be obtained with respect to the inhomogeneous mixed Hölder-variation distance.

Theorem 3.3. *Let $\delta \in (0, 1)$ and $\gamma, p \in (1, \infty)$ be such that $\delta > 1/p$ and $\gamma > 1/\delta$. For a Riesz-type geometric rough path $\mathbf{X} \in \tilde{V}^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$, for $V \in \text{Lip}^\gamma$ and for every initial condition $y_0 \in \mathbb{R}^m$, the controlled differential equation (3.4) has a unique solution $Y \in \tilde{V}^{\delta,p}([0, T]; \mathbb{R}^m)$.*

Furthermore, the Itô–Lyons map Φ as defined in (3.5) is locally Lipschitz continuous, that is, for $y_0^i \in \mathbb{R}^m$, $X^i \in \tilde{V}^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$ and $V^i \in \text{Lip}^\gamma$ satisfying

$$\|\mathbf{X}^i\|_{\tilde{V}^{\delta,p}} \leq b \quad \text{and} \quad \|V^i\|_{\text{Lip}^\gamma} \leq l, \quad i = 1, 2,$$

for some $b, l > 0$, with corresponding solution Y^i , there exist a constant $C = C(b, l, \gamma, \delta, p) \geq 1$ such that

$$\|Y^1 - Y^2\|_{\tilde{V}^{\delta,p}} \leq C(\|V^1 - V^2\|_{\text{Lip}^{\gamma-1}} + |y_0^1 - y_0^2| + \rho_{\tilde{V}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2)).$$

Before we come to the proof, we recall that a continuous function $\omega: \Delta_T \rightarrow [0, \infty)$ is called a *control function* if $\omega(s, s) = 0$ for $s \in [0, T]$ and ω is super-additive; cf. Remark 2.3.

Proof. Due to Proposition 2.8 and Theorem 2.11, the assumption

$$\mathbf{X} \in \tilde{V}^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$$

implies $\mathbf{X} \in C^{1/\delta\text{-var}}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$ as $\delta > 1/p$. Therefore, the controlled differential equation (3.4) has a unique solution $Y \in C^{1/\delta\text{-var}}([0, T]; \mathbb{R}^m)$ given the regularity of the vector field $V \in \text{Lip}^\gamma$ with $\gamma > 1/\delta$; see [FV10, Theorem 10.26].

It remains to show the Riesz-type variation of Y and the local Lipschitz continuity of the Itô–Lyons map Φ as defined in (3.5). For this purpose we choose a suitable control function ω (see (3.9) for the specific definition) and introduce the distance

$$\rho_{1/\delta-\omega}(\mathbf{X}^1, \mathbf{X}^2) := \max_{k=1, \dots, \lfloor 1/\delta \rfloor} \rho_{1/\delta-\omega; [0, T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)$$

for $\mathbf{X}^1, \mathbf{X}^2 \in \tilde{V}^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$ and

$$\rho_{1/\delta-\omega; [0, T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2) := \sup_{0 \leq s < t \leq T} \frac{|\pi_k(\mathbf{X}_{s,t}^1 - \mathbf{X}_{s,t}^2)|}{\omega(s, t)^{k\delta}}, \quad k = 1, \dots, \lfloor 1/\delta \rfloor.$$

As one can see in the proof of [FV10, Theorem 10.26], one has the following two estimates for the control function ω and a constant $C = C(\gamma, \delta) > 0$:

$$\begin{aligned} & \|Y^1 - Y^2\|_{\infty; [0, T]} \\ (3.6) \quad & \leq C \left(|y_0^1 - y_0^2| + \frac{1}{l} \|V^1 - V^2\|_{\text{Lip}^{\gamma-1}} + \rho_{1/\delta-\omega}(\mathbf{X}^1, \mathbf{X}^2) \right) \exp(Cl^{1/\delta}\omega(0, T)) \end{aligned}$$

and, for all $u < v$ in $[0, T]$,

$$\begin{aligned} (3.7) \quad & |Y_{u,v}^1 - Y_{u,v}^2| \leq C \left(l|Y_u^1 - Y_u^2| + \|V^1 - V^2\|_{\text{Lip}^{\gamma-1}} + l\rho_{1/\delta-\omega}(\mathbf{X}^1, \mathbf{X}^2) \right) \omega(u, v)^\delta \\ & \times \exp(Cl^{1/\delta}\omega(0, T)). \end{aligned}$$

From inequality (3.7) we deduce that

$$\begin{aligned} \|Y^1 - Y^2\|_{1/\delta\text{-var};[s,t]}^{\frac{1}{\delta}} &= \sup_{\mathcal{P} \subset [s,t]} \sum_{[u,v] \in \mathcal{P}} |Y_{u,v}^1 - Y_{u,v}^2|^{1/\delta} \\ &\lesssim \left(l \|Y^1 - Y^2\|_{\infty;[s,t]} + \|V^1 - V^2\|_{\text{Lip}^{\gamma-1}} + l \rho_{1/\delta-\omega}(\mathbf{X}^1, \mathbf{X}^2) \right)^{1/\delta} \omega(s, t) \\ &\quad \times \exp(C l^{1/\delta} \omega(0, T))^{1/\delta}, \end{aligned}$$

which further leads to

$$\begin{aligned} \|Y^1 - Y^2\|_{\tilde{V}^{\delta,p}}^p &= \sup_{\mathcal{P} \subset [0,T]} \sum_{[s,t] \in \mathcal{P}} \frac{\|Y^1 - Y^2\|_{1/\delta\text{-var};[s,t]}^p}{|t - s|^{\delta p - 1}} \\ &\lesssim \left(l \|y^1 - y^2\|_{\infty;[0,T]} + \|V^1 - V^2\|_{\text{Lip}^{\gamma-1}} + l \rho_{1/\delta-\omega}(\mathbf{X}^1, \mathbf{X}^2) \right)^p \\ &\quad \times \exp(C l^{1/\delta} \omega(0, T))^p \left(\sup_{\mathcal{P} \subset [0,T]} \sum_{[s,t] \in \mathcal{P}} \frac{\omega(s, t)^{\delta p}}{|t - s|^{\delta p - 1}} \right)^p. \end{aligned}$$

Plugging in estimate (3.6) in the last inequality gives

$$\begin{aligned} \|Y^1 - Y^2\|_{\tilde{V}^{\delta,p}} &\lesssim \left(l |y_0^1 - y_0^2| + \|V^1 - V^2\|_{\text{Lip}^{\gamma-1}} + l \rho_{1/\delta-\omega}(\mathbf{X}^1, \mathbf{X}^2) \right) \\ (3.8) \quad &\quad \times \exp(C l^{1/\delta} \omega(0, T)) \left(\sup_{\mathcal{P} \subset [0,T]} \sum_{[s,t] \in \mathcal{P}} \frac{\omega(s, t)^{\delta p}}{|t - s|^{\delta p - 1}} \right). \end{aligned}$$

In order to complete the proof, we consider the control function

$$(3.9) \quad \omega(s, t) := \|\mathbf{X}^1\|_{1/\delta\text{-var};[s,t]}^{\frac{1}{\delta}} + \|\mathbf{X}^2\|_{1/\delta\text{-var};[s,t]}^{\frac{1}{\delta}} + \sum_{k=1}^{\lfloor 1/\delta \rfloor} \omega_{\mathbf{X}^1, \mathbf{X}^2}^{(k)}(s, t), \quad (s, t) \in \Delta_T,$$

where

$$\omega_{\mathbf{X}^1, \mathbf{X}^2}^{(k)}(s, t) := \left(\frac{\rho_{1/\delta\text{-var};[s,t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)}{\rho_{\tilde{V}^{\delta,p};[0,T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)} \right)^{\frac{1}{\delta k}}$$

with the convention $0/0 := 0$, and investigate some properties of ω . First notice that ω fulfills all the requirements to apply [FV10, Theorem 10.26]. Moreover, it is easy to see that

$$(3.10) \quad \rho_{1/\delta-\omega}(\mathbf{X}^1, \mathbf{X}^2) \lesssim \rho_{\tilde{V}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2)$$

as one has for $k = 1, \dots, \lfloor 1/\delta \rfloor$ and $0 \leq s < t \leq T$ the following estimate:

$$|\pi_k(\mathbf{X}_{s,t}^1 - \mathbf{X}_{s,t}^2)| \leq \frac{\rho_{1/\delta-\omega;[s,t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)}{\rho_{\tilde{V}^{\delta,p};[0,T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)} \rho_{\tilde{V}^{\delta,p};[0,T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2) \leq \omega(s, t)^{\delta k} \rho_{\tilde{V}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2).$$

The last observation on ω we need is

$$(3.11) \quad \sup_{\mathcal{P} \subset [0,T]} \sum_{[s,t] \in \mathcal{P}} \frac{\omega(s, t)^{\delta p}}{|t - s|^{\delta p - 1}} \lesssim \|\mathbf{X}^1\|_{\tilde{V}^{\delta,p}}^p + \|\mathbf{X}^2\|_{\tilde{V}^{\delta,p}}^p + 1.$$

Indeed, by Proposition 2.8 we have for every partition \mathcal{P} of $[0, T]$

$$\sum_{[s,t] \in \mathcal{P}} \frac{\|\mathbf{X}^i\|_{1/\delta\text{-var};[s,t]}^p}{|t-s|^{\delta p-1}} \lesssim \sum_{[s,t] \in \mathcal{P}} \frac{\|\mathbf{X}^i\|_{\tilde{V}^{\delta,p};[s,t]}^p |t-s|^{\delta p-1}}{|t-s|^{\delta p-1}} \lesssim \|\mathbf{X}^i\|_{\tilde{V}^{\delta,p}}^p, \quad i = 1, 2,$$

and further using

$$\begin{aligned} \rho_{1/\delta\text{-var};[s,t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2) &\leq \left(\frac{\rho_{1/\delta\text{-var};[s,t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}}}{|t-s|^{\delta p-1}} \right)^{\frac{k}{p}} |t-s|^{(\delta-1/p)k} \\ &\leq \rho_{\tilde{V}^{\delta,p};[s,t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2) |t-s|^{(\delta-1/p)k}, \quad \text{for } k = 1, \dots, \lfloor 1/\delta \rfloor, \end{aligned}$$

we arrive at

$$\begin{aligned} \sum_{[s,t] \in \mathcal{P}} \frac{\omega_{\mathbf{X}^1, \mathbf{X}^2}^{(k)}(s, t)^{\delta p}}{|t-s|^{\delta p-1}} &= \rho_{\tilde{V}^{\delta,p};[0,T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{-\frac{p}{k}} \sum_{[s,t] \in \mathcal{P}} \frac{\rho_{1/\delta\text{-var};[s,t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}}}{|t-s|^{\delta p-1}} \\ &\leq \rho_{\tilde{V}^{\delta,p};[0,T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{-\frac{p}{k}} \sum_{[s,t] \in \mathcal{P}} \rho_{\tilde{V}^{\delta,p};[s,t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}} \leq 1. \end{aligned}$$

By combining these estimates we deduce (3.11).

Therefore, estimate (3.8) together with (3.10) and (3.11) reveals

$$\begin{aligned} \|Y^1 - Y^2\|_{\tilde{V}^{\delta,p}} &\lesssim \left(l|y_0^1 - y_0^2| + \|V^1 - V^2\|_{\text{Lip}^{\gamma-1}} + l\rho_{\tilde{V}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2) \right) \\ &\quad \times \exp(Cl^{1/\delta}(2b+1)(2b+1)), \end{aligned}$$

which completes the proof. \square

3.3. Inhomogeneous Riesz-type distance. In the context of rough path theory it is very convenient to work with the characterization of Riesz-type variation in terms of classical q -variation and to introduce the corresponding inhomogeneous mixed Hölder-variation distance $\rho_{\tilde{V}^{\delta,p}}$, as we have seen in the previous subsection. However, also the other characterizations of Riesz-type variation allow for obtaining the local Lipschitz continuity of the Itô–Lyons map Φ as defined in (3.5).

Keeping in mind the Riesz-type variation (2.2), we define for $\delta \in (0, 1)$ and $p \in [1/\delta, \infty)$ inhomogeneous *Riesz-type distance* by

$$\rho_{V^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2) := \max_{k=1, \dots, \lfloor 1/\delta \rfloor} \rho_{V^{\delta,p};[0,T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2),$$

for $\mathbf{X}^1, \mathbf{X}^2 \in V^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$, where we set

$$\rho_{V^{\delta,p};[s,t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2) := \sup_{\mathcal{P} \subset [s,t]} \left(\sum_{[u,v] \in \mathcal{P}} \frac{|\pi_k(\mathbf{X}_{u,v}^1 - \mathbf{X}_{u,v}^2)|^{\frac{p}{k}}}{|u-v|^{\delta p-1}} \right)^{\frac{k}{p}}, \quad [s, t] \subset [0, T]$$

for $k = 1, \dots, \lfloor 1/\delta \rfloor$. Indeed, the inhomogeneous Riesz-type distance and inhomogeneous mixed Hölder-variation distance are equivalent.

Lemma 3.4. *If $\delta \in (0, 1)$ and $p \in (1, \infty)$ with $\delta > 1/p$, then*

$$\rho_{V^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2) \lesssim \rho_{\tilde{V}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2) \lesssim \rho_{V^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2)$$

for $\mathbf{X}^1, \mathbf{X}^2 \in V^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$.

Proof. The first inequality directly follows from

$$|\pi_k(\mathbf{X}_{u,v}^1 - \mathbf{X}_{u,v}^2)|^{\frac{p}{k}} \leq \rho_{1/\delta\text{-var};[u,v]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}}, \quad [u, v] \subset [0, T]$$

for $k = 1, \dots, \lfloor 1/\delta \rfloor$.

For the second inequality we notice

$$\begin{aligned} |\pi_k(\mathbf{X}_{u,v}^1 - \mathbf{X}_{u,v}^2)|^{\frac{1}{\delta k}} &\leq \left(\frac{|\pi_k(\mathbf{X}_{u,v}^1 - \mathbf{X}_{u,v}^2)|^{\frac{p}{k}}}{|u-v|^{\delta p-1}} \right)^{\frac{1}{\delta p}} |u-v|^{1-\frac{1}{\delta p}} \\ &\leq \rho_{V^{\delta,p};[u,v]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{1}{\delta k}} |u-v|^{1-\frac{1}{\delta p}}, \quad [u, v] \subset [0, T] \end{aligned}$$

for $k = 1, \dots, \lfloor 1/\delta \rfloor$. Due to Remark 2.3 the function

$$\Delta_T \ni (u, v) \mapsto \rho_{V^{\delta,p};[u,v]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{1}{\delta k}} |u-v|^{1-\frac{1}{\delta p}}$$

is super-additive and thus we get

$$\rho_{1/\delta\text{-var};[u,v]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}} \leq \rho_{V^{\delta,p};[u,v]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2) |u-v|^{\delta p-1}.$$

Therefore, using the super-additive of $\rho_{V^{\delta,p};[u,v]}^{(k)}(\mathbf{X}^1, \mathbf{X}^1)^{\frac{p}{k}}$ as functions in $(u, v) \in \Delta_T$, we obtain

$$\rho_{\tilde{V}^{\delta,p};[u,v]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}} \leq \rho_{V^{\delta,p};[u,v]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}}$$

for $k = 1, \dots, \lfloor 1/\delta \rfloor$, which implies the second inequality. \square

Applying the equivalence of the inhomogeneous distances $\rho_{V^{\delta,p}}$ and $\rho_{\tilde{V}^{\delta,p}}$ (Lemma 3.4) and the characterization of Riesz-type spaces (Theorem 2.11), the local Lipschitz continuity of the Itô–Lyons map (3.5) with respect to $\rho_{V^{\delta,p}}$ is an immediate consequence of Theorem 3.3:

Corollary 3.5. *Let $\delta \in (0, 1)$ and $\gamma, p \in (1, \infty)$ be such that $\delta > 1/p$ and $\gamma > 1/\delta$.*

The Itô–Lyons map

$$\Phi: \mathbb{R}^m \times \text{Lip}^\gamma \times V^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n)) \rightarrow V^{\delta,p}([0, T]; \mathbb{R}^m) \quad \text{via} \quad \Phi(y_0, V, \mathbf{X}) := Y,$$

where Y denotes the solution to controlled differential equation (3.4) given the input (y_0, V, \mathbf{X}) , is locally Lipschitz continuous with respect to inhomogeneous Riesz-type distance $\rho_{V^{\delta,p}}$.

3.4. Inhomogeneous Nikolskii-type distance. To complete the picture, we provide in this subsection an inhomogeneous distance in terms of Nikolskii regularity (cf. (2.6)), which is locally Lipschitz equivalent to the inhomogeneous Riesz-type distance and which ensures the local Lipschitz continuity of the Itô–Lyons map Φ as defined in (3.5).

To that end we introduce the inhomogeneous Nikolskii-type distance as follows: For $\mathbf{X}^1, \mathbf{X}^2 \in \tilde{N}^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$ we set

$$\rho_{N^{\delta,p};[u,v]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2) := \sup_{|v-u| \geq h > 0} h^{-\delta k} \left(\int_u^{v-h} |\pi_k(\mathbf{X}_{r,r+h}^1 - \mathbf{X}_{r,r+h}^2)|^{\frac{p}{k}} dr \right)^{\frac{k}{p}}, \quad [u, v] \subset [0, T],$$

and

$$\rho_{\tilde{N}^{\delta,p};[s,t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2) := \sup_{\mathcal{P} \subset [s,t]} \left(\sum_{[u,v] \in \mathcal{P}} \rho_{N^{\delta,p};[u,v]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}} \right)^{\frac{k}{p}}, \quad [s, t] \subset [0, T],$$

for $k = 1, \dots, \lfloor 1/\delta \rfloor$. The *inhomogeneous Nikolskii-type distance* $\rho_{\hat{N}^{\delta,p}}$ is defined by

$$\rho_{\hat{N}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2) := \max_{k=1, \dots, N} \rho_{\hat{N}^{\delta,p}; [0, T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2).$$

In the next two lemmas (Lemmas 3.6 and 3.7) we establish that the two ways of introducing an inhomogeneous distance on $C([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$, given by the inhomogeneous mixed Hölder-variation distance and the inhomogeneous Nikolskii-type distance, are locally equivalent.

Lemma 3.6. *If $\delta \in (0, 1)$ and $p \in (1, \infty)$ with $\delta > 1/p$, then*

$$\rho_{\hat{N}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2) \lesssim \rho_{\tilde{V}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2)$$

for $\mathbf{X}^1, \mathbf{X}^2 \in \tilde{V}^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$.

Proof. Let $\mathbf{X}^1, \mathbf{X}^2 \in \tilde{V}^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$ and $k = 1, \dots, N$. For $[s, t] \subset [0, T]$ and $h \in (0, t - s]$ fixed we consider the dissection $\mathcal{P}(h) := \{[t_i, t_{i+1}] : s = t_0 < \dots < t_M = t - h\}$ such that

$$|t_M - t_{M-1}| \leq h \quad \text{and} \quad |t_{i+1} - t_i| = h \quad \text{for } i = 0, \dots, M-2, \quad M \in \mathbb{N}.$$

Since $\sup_{u \in [t_i, t_{i+1}]} |\pi_k(\mathbf{X}_{u, u+h}^1 - \mathbf{X}_{u, u+h}^2)| \leq \rho_{1/\delta\text{-var}; [t_i, t_{i+2}]}(\mathbf{X}^1, \mathbf{X}^2)$ for $i = 1, \dots, M-1$ with $t_{M+1} := t - h$, we deduce that

$$\begin{aligned} \int_s^{t-h} |\pi_k(\mathbf{X}_{u, u+h}^1 - \mathbf{X}_{u, u+h}^2)|^{\frac{p}{k}} du &\leq \sum_{i=1}^{M-1} \sup_{u \in [t_i, t_{i+1}]} |\pi_k(\mathbf{X}_{u, u+h}^1 - \mathbf{X}_{u, u+h}^2)|^{\frac{p}{k}} h \\ &\leq \frac{1}{2} (2h)^{\delta p} \sum_{i=1}^{M-1} \frac{\rho_{1/\delta\text{-var}; [t_i, t_{i+2}]}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}}}{|t_i - t_{i+2}|^{\delta p - 1}} \\ &\lesssim h^{\delta p} \rho_{\tilde{V}^{\delta,p}; [s, t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}}. \end{aligned}$$

Therefore, by the super-additivity of $\rho_{\hat{N}^{\delta,p}; [s, t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}}$ and $\rho_{\tilde{V}^{\delta,p}; [s, t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}}$ as functions in $(s, t) \in \Delta_T$ we obtain

$$\rho_{\hat{N}^{\delta,p}; [0, T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2) \lesssim \rho_{\tilde{V}^{\delta,p}; [0, T]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)$$

for every $k = 1, \dots, \lfloor 1/\delta \rfloor$, which implies that $\rho_{\hat{N}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2) \lesssim \rho_{\tilde{V}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2)$. \square

Lemma 3.7. *Let $\delta \in (0, 1)$ and $p \in (1, \infty)$ with $\delta > 1/p$. For Riesz-type geometric rough paths $\mathbf{X}^1, \mathbf{X}^2 \in \hat{N}^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$ there exists a constant*

$$C := C(\delta, p, \|\mathbf{X}^1\|_{\hat{N}^{\delta,p}}, \|\mathbf{X}^2\|_{\hat{N}^{\delta,p}}) \geq 1,$$

depending only on δ, p and the upper bound of $\|\mathbf{X}^1\|_{\hat{N}^{\delta,p}}$ and $\|\mathbf{X}^2\|_{\hat{N}^{\delta,p}}$, such that

$$\rho_{\tilde{V}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2) \leq C \rho_{\hat{N}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2).$$

Proof. Let $\mathbf{X}^1, \mathbf{X}^2 \in \hat{N}^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n))$ be Riesz-type geometric rough paths. In order to prove the inequality in Lemma 3.7, it is sufficient to show that there exists a constant $C = C(\delta, p, \|\mathbf{X}^1\|_{\hat{N}^{\delta,p}}, \|\mathbf{X}^2\|_{\hat{N}^{\delta,p}}) \geq 1$ such that

$$(3.12) \quad |\pi_j(\mathbf{X}_{s,t}^1 - \mathbf{X}_{s,t}^2)|^{\frac{1}{\delta j}} \leq C \left(\sum_{i=1}^j \rho_{\hat{N}^{\delta,p}; [s, t]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{1}{\delta p}} |t - s|^{1 - \frac{1}{\delta p}} =: \omega^{(j)}(s, t)$$

for all $s, t \in [0, T]$ with $s < t$ and for every $j = 1, \dots, \lfloor 1/\delta \rfloor$.

Indeed, for each $j = 1, \dots, \lfloor 1/\delta \rfloor$ the function $\omega^{(j)}: \Delta_T \rightarrow [0, \infty)$ is the super-additive:

$$\begin{aligned} \omega^{(j)}(s, t) + \omega^{(j)}(t, u) &\leq C \left(\sum_{i=1}^j \rho_{\hat{N}^{\delta,p};[s,t]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} + \rho_{\hat{N}^{\delta,p};[u,t]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{1}{\delta p}} (|t-s| + |u-t|)^{1-\frac{1}{\delta p}} \\ &\leq C \left(\sum_{i=1}^j \rho_{\hat{N}^{\delta,p};[s,u]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{1}{\delta p}} |u-s|^{1-\frac{1}{\delta p}} = \omega^{(j)}(s, u), \end{aligned}$$

for $0 \leq s \leq t \leq u \leq T$, where we used Hölder's inequality and $p/j \geq 1$. This implies

$$\begin{aligned} \rho_{1/\delta\text{-var};[s,t]}^{(j)}(\mathbf{X}^1, \mathbf{X}^2) &= \sup_{\mathcal{P} \subset [s,t]} \left(\sum_{[u,v] \in \mathcal{P}} |\pi_j(\mathbf{X}_{u,v}^1 - \mathbf{X}_{u,v}^2)|^{\frac{1}{\delta j}} \right)^{j\delta} \\ &\leq C \left(\sum_{i=1}^j \rho_{\hat{N}^{\delta,p};[s,t]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{j}{p}} |t-s|^{\frac{j}{p}(\delta p-1)}, \end{aligned}$$

where the super-additivity of $\omega^{(j)}$ is applied in the last line. Hence, we get further

$$\begin{aligned} \rho_{\hat{V}^{\delta,p};[0,T]}^{(j)}(\mathbf{X}^1, \mathbf{X}^2) &= \sup_{\mathcal{P} \subset [0,T]} \left(\sum_{[u,v] \in \mathcal{P}} \frac{\rho_{1/\delta\text{-var};[u,v]}^{(j)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}}}{|u-v|^{\delta p-1}} \right)^{\frac{j}{p}} \\ &\leq C \sup_{\mathcal{P} \subset [0,T]} \left(\sum_{[u,v] \in \mathcal{P}} \sum_{i=1}^j \rho_{\hat{N}^{\delta,p};[u,v]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{j}{p}} \\ &\leq C \sum_{i=1}^j \rho_{\hat{N}^{\delta,p};[0,T]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2), \end{aligned}$$

which immediately gives by taking the maximum over $j = 1, \dots, \lfloor 1/\delta \rfloor$ that

$$\rho_{\hat{V}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2) \leq C \rho_{\hat{N}^{\delta,p}}(\mathbf{X}^1, \mathbf{X}^2).$$

In order to prove inequality (3.12) for each $j = 1, \dots, \lfloor 1/\delta \rfloor$, we argue via induction. For $j = 1$ inequalities (3.12) is an easy consequence of Theorem 2.11. We now assume that (3.12) is true for the levels $j = 1, \dots, k-1$ and establish the inequality for level k . Let us fix $s, t \in [0, T]$ with $s < t$ and define

$$Z_u^s := \pi_k(\mathbf{X}_{s,s+u}^1 - \mathbf{X}_{s,s+u}^2), \quad u \in [0, t-s].$$

For $u, h \in [0, t-s]$ with $u+h \in [0, t-s]$ and using

$$\mathbf{X}_{s,s+u+h}^1 - \mathbf{X}_{s,s+u}^1 = \mathbf{X}_{s,s+u}^1 \otimes (\mathbf{X}_{s+u,s+u+h}^1 - 1), \quad \pi_0(\mathbf{X}_{s+u,s+u+h}^1 - 1) = 0$$

and

$$\pi_j(\mathbf{X}_{s+u,s+u+h}^1 - 1) = \pi_j(\mathbf{X}_{s+u,s+u+h}^1) \quad \text{for } j > 0,$$

we obtain

$$\begin{aligned}
 Z_{u+h}^s - Z_u^s &= \pi_k(\mathbf{X}_{s,s+u+h}^1 - \mathbf{X}_{s,s+u}^1) - \pi_k(\mathbf{X}_{s,s+u+h}^2 - \mathbf{X}_{s,s+u}^2) \\
 &= \sum_{j=1}^k \pi_{k-j}(\mathbf{X}_{s,s+u}^1 \otimes \pi_j(\mathbf{X}_{s+u,s+u+h}^1) - \sum_{j=1}^k \pi_{k-j}(\mathbf{X}_{s,s+u}^2 \otimes \pi_j(\mathbf{X}_{s+u,s+u+h}^2)) \\
 &= \sum_{j=1}^k \pi_{k-j}(\mathbf{X}_{s,s+u}^1 \otimes \pi_j(\mathbf{X}_{s+u,s+u+h}^1 - \mathbf{X}_{s+u,s+u+h}^2) \\
 &\quad + \sum_{j=1}^k \pi_{k-j}(\mathbf{X}_{s,s+u}^1 - \mathbf{X}_{s,s+u}^2) \otimes \pi_j(\mathbf{X}_{s+u,s+u+h}^2).
 \end{aligned}$$

Keeping in mind $\pi_0(\mathbf{X}_{s,s+u}^1 - \mathbf{X}_{s,s+u}^2) = 0$, we arrive at

$$\begin{aligned}
 Z_{u+h}^s - Z_u^s &= \sum_{j=1}^{k-1} \pi_{k-j}(\mathbf{X}_{s,s+u}^1 \otimes \pi_j(\mathbf{X}_{s+u,s+u+h}^1 - \mathbf{X}_{s+u,s+u+h}^2) \\
 &\quad + \sum_{j=1}^{k-1} \pi_{k-j}(\mathbf{X}_{s,s+u}^1 - \mathbf{X}_{s,s+u}^2) \otimes \pi_j(\mathbf{X}_{s+u,s+u+h}^2) \\
 &\quad + \pi_k(\mathbf{X}_{s+u,s+u+h}^1 - \mathbf{X}_{s+u,s+u+h}^2).
 \end{aligned}$$

Hence, we get the following estimate:

$$\sup_{|t-s| \geq h > 0} h^{-\delta} \left(\int_s^{t-h} |Z_{u+h}^s - Z_u^s|^p du \right)^{\frac{1}{p}} \lesssim \Delta_1 + \Delta_2 + \Delta_3$$

where we set

$$\begin{aligned}
 \Delta_1 &:= \sum_{j=1}^{k-1} \sup_{|t-s| \geq h > 0} h^{-\delta} \left(\int_s^{t-h} \|\mathbf{X}_{s,s+u}^1\|^{p(k-j)} |\pi_j(\mathbf{X}_{s+u,s+u+h}^1 - \mathbf{X}_{s+u,s+u+h}^2)|^p du \right)^{\frac{1}{p}}, \\
 \Delta_2 &:= \sum_{j=1}^{k-1} \sup_{|t-s| \geq h > 0} h^{-\delta} \left(\int_s^{t-h} |\pi_{k-j}(\mathbf{X}_{s,s+u}^1 - \mathbf{X}_{s,s+u}^2)|^p \|\mathbf{X}_{s+u,s+u+h}^2\|^{pj} du \right)^{\frac{1}{p}}, \\
 \Delta_3 &:= \sup_{|t-s| \geq h > 0} h^{-\delta} \left(\int_s^{t-h} |\pi_k(\mathbf{X}_{s+u,s+u+h}^1 - \mathbf{X}_{s+u,s+u+h}^2)|^p du \right)^{\frac{1}{p}}.
 \end{aligned}$$

Due to Proposition 2.8 and $\delta > 1/p$, we have

$$\|\mathbf{X}_{s,s+u}^1\|^{p(k-j)} \leq \|\mathbf{X}^1\|_{1/\delta\text{-var};[s,t]}^{p(k-j)} \lesssim \|\mathbf{X}^1\|_{\hat{N}^{\delta,p};[s,t]}^{p(k-j)} |t-s|^{(\delta-\frac{1}{p})p(k-j)}.$$

Moreover, the induction hypothesis gives

$$\begin{aligned}
 &|\pi_j(\mathbf{X}_{s+u,s+u+h}^1 - \mathbf{X}_{s+u,s+u+h}^2)|^{(1-\frac{1}{j})p} \\
 &\lesssim \left(\sum_{i=1}^j \rho_{\hat{N}^{\delta,p};[s,t]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{j-1} |t-s|^{(j-1)(\delta-\frac{1}{p})p}.
 \end{aligned}$$

Therefore, Δ_1 can be estimated by

$$\begin{aligned}
\Delta_1 &\lesssim \sum_{j=1}^{k-1} \|\mathbf{X}^1\|_{\hat{N}^{\delta,p};[s,t]}^{k-j} |t-s|^{(k-j)(\delta-\frac{1}{p})} \\
&\quad \times \left(\sum_{i=1}^j \rho_{\hat{N}^{\delta,p};[s,t]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{j-1}{p}} |t-s|^{(j-1)(\delta-\frac{1}{p})} \\
&\quad \times \sup_{|t-s|\geq h>0} h^{-\delta} \left(\int_s^{t-h} |\pi_j(\mathbf{X}_{s+u,s+u+h}^1 - \mathbf{X}_{s+u,s+u+h}^2)|^{\frac{p}{j}} du \right)^{\frac{1}{p}} \\
&\leq \sum_{j=1}^{k-1} \|\mathbf{X}^1\|_{\hat{N}^{\delta,p};[s,t]}^{k-j} \rho_{\hat{N}^{\delta,p};[s,t]}^{(j)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{1}{j}} \\
&\quad \times \left(\sum_{i=1}^j \rho_{\hat{N}^{\delta,p};[s,t]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{j-1}{p}} |t-s|^{(k-1)(\delta-\frac{1}{p})} \\
&\leq \sum_{j=1}^{k-1} \|\mathbf{X}^1\|_{\hat{N}^{\delta,p};[s,t]}^{k-j} \left(\sum_{i=1}^j \rho_{\hat{N}^{\delta,p};[s,t]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{j}{p}} |t-s|^{(\delta-\frac{1}{p})(k-1)}.
\end{aligned}$$

For Δ_2 we first observe again due to Proposition 2.8 that

$$\|\mathbf{X}_{s+u,s+u+h}^2\|^{pj} \lesssim \|\mathbf{X}_{s+u,s+u+h}^2\|^p \|\mathbf{X}^2\|_{\hat{N}^{\delta,p};[s,t]}^{p(j-1)} |t-s|^{(\delta-\frac{1}{p})(j-1)p}$$

and by the induction hypothesis that

$$|\pi_{k-j}(\mathbf{X}_{s,s+u}^1 - \mathbf{X}_{s,s+u}^2)|^p \lesssim \left(\sum_{i=1}^{k-j} \rho_{\hat{N}^{\delta,p};[s,t]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k-j}} \right)^{k-j} |t-s|^{(\delta-\frac{1}{p})(k-j)p}.$$

Combining the last two estimates, we get

$$\begin{aligned}
\Delta_2 &\lesssim \sum_{j=1}^{k-1} \|\mathbf{X}^2\|_{\hat{N}^{\delta,p};[s,t]}^{j-1} |t-s|^{(\delta-\frac{1}{p})(j-1)} \\
&\quad \times \sup_{|t-s|\geq h>0} h^{-\delta} \left(\int_s^{t-h} |\pi_{k-j}(\mathbf{X}_{s,s+u}^1 - \mathbf{X}_{s,s+u}^2)|^p \|\mathbf{X}_{s+u,s+u+h}^2\|^p du \right)^{\frac{1}{p}} \\
&\lesssim \sum_{j=1}^{k-1} \|\mathbf{X}^2\|_{\hat{N}^{\delta,p};[s,t]}^j |t-s|^{(\delta-\frac{1}{p})(k-1)} \left(\sum_{i=1}^{k-j} \rho_{\hat{N}^{\delta,p};[s,t]}^{(i)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k-j}} \right)^{\frac{k-j}{p}}.
\end{aligned}$$

For Δ_3 we briefly need to introduce the inhomogeneous Hölder distance for level k by

$$\rho_{(\delta-1/p)\text{-Hö};[s,t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2) := \sup_{u,v \in [s,t]; u \neq v} \frac{|\pi_k(\mathbf{X}_{s+u,s+v}^1 - \mathbf{X}_{s+u,s+v}^2)|}{|u-v|^{(\delta-\frac{1}{p})k}}.$$

This time we simply estimate

$$\Delta_3 \leq \rho_{\hat{N}^{\delta,p};[s,t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{1}{k}} \rho_{(\delta-1/p)\text{-Hö};[s,t]}^{(k)}(\mathbf{X}^1, \mathbf{X}^2)^{1-\frac{1}{k}} |t-s|^{(\delta-\frac{1}{p})(k-1)}.$$

Applying Proposition 2.8 to Z^s we get

$$|\pi_k(\mathbf{X}_{s,t}^1 - \mathbf{X}_{s,t}^2)| = |Z_0^s - Z_{t-s}^s| \lesssim \sup_{|t-s| \geq h > 0} h^{-\delta} \left(\int_s^{t-h} |Z_{u+h}^s - Z_u^s|^p \, du \right)^{\frac{1}{p}} |t-s|^{\delta - \frac{1}{p}}.$$

Putting the estimates for Δ_1 , Δ_2 and Δ_3 , we deduce further

$$\begin{aligned} & |\pi_k(\mathbf{X}_{s,t}^1 - \mathbf{X}_{s,t}^2)| \\ & \leq \tilde{C} |t-s|^{\delta - \frac{1}{p}} \left[\sum_{j=1}^{k-1} \left(\sum_{i=1}^j \rho_{\hat{N}^{\delta,p};[s,t]}^{(i)} (\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{j}{p}} |t-s|^{(\delta - \frac{1}{p})(k-1)} \right. \\ & \quad \left. + \rho_{\hat{N}^{\delta,p};[s,t]}^{(k)} (\mathbf{X}^1, \mathbf{X}^2)^{\frac{1}{k}} \rho_{(\delta-1/p)\text{-Hö};[s,t]}^{(k)} (\mathbf{X}^1, \mathbf{X}^2)^{1-\frac{1}{k}} |t-s|^{(\delta - \frac{1}{p})(k-1)} \right] \\ & \lesssim \tilde{C} |t-s|^{(\delta - \frac{1}{p})k} \left[\sum_{j=1}^k \rho_{\hat{N}^{\delta,p};[s,t]}^{(j)} (\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right. \\ & \quad \left. + \left(\sum_{j=1}^k \rho_{\hat{N}^{\delta,p};[s,t]}^{(j)} (\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{1}{p}} \rho_{(\delta-1/p)\text{-Hö};[s,t]}^{(k)} (\mathbf{X}^1, \mathbf{X}^2)^{1-\frac{1}{k}} \right] \\ & \lesssim \tilde{C} |t-s|^{(\delta - \frac{1}{p})k} \left[\left(\sum_{j=1}^k \rho_{\hat{N}^{\delta,p};[s,t]}^{(j)} (\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{k}{p}} \right. \\ & \quad \left. + \left(\sum_{j=1}^k \rho_{\hat{N}^{\delta,p};[s,t]}^{(j)} (\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{1}{p}} \rho_{(\delta-1/p)\text{-Hö};[s,t]}^{(k)} (\mathbf{X}^1, \mathbf{X}^2)^{1-\frac{1}{k}} \right], \end{aligned}$$

for some constant $\tilde{C} = \tilde{C}(\delta, p, \|\mathbf{X}^1\|_{\hat{N}^{\delta,p}}, \|\mathbf{X}^2\|_{\hat{N}^{\delta,p}}) \geq 1$, which can be rewritten as

$$\begin{aligned} \frac{|\pi_k(\mathbf{X}_{s,t}^1 - \mathbf{X}_{s,t}^2)|}{|t-s|^{(\delta - \frac{1}{p})k}} & \lesssim \tilde{C} \left[\left(\sum_{j=1}^k \rho_{\hat{N}^{\delta,p};[s,t]}^{(j)} (\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{k}{p}} \right. \\ & \quad \left. + \left(\sum_{j=1}^k \rho_{\hat{N}^{\delta,p};[s,t]}^{(j)} (\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{1}{p}} \rho_{(\delta-1/p)\text{-Hö};[s,t]}^{(k)} (\mathbf{X}^1, \mathbf{X}^2)^{1-\frac{1}{k}} \right]. \end{aligned}$$

In other words, we showed that

$$\frac{\rho_{(\delta-1/p)\text{-Hö};[s,t]}^{(k)} (\mathbf{X}^1, \mathbf{X}^2)}{\tilde{\omega}^{(k)}(s, t)} \lesssim \tilde{C} \left[1 + \left(\frac{\rho_{(\delta-1/p)\text{-Hö};[s,t]}^{(k)} (\mathbf{X}^1, \mathbf{X}^2)}{\tilde{\omega}^{(k)}(s, t)} \right)^{1-\frac{1}{k}} \right],$$

with

$$\tilde{\omega}^{(k)}(s, t) := \left(\sum_{j=1}^k \rho_{\hat{N}^{\delta,p};[s,t]}^{(j)} (\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{j}} \right)^{\frac{k}{p}}.$$

Hence, there exists a constant $C = C(\delta, p, \|\mathbf{X}^1\|_{\hat{N}^{\delta,p}}, \|\mathbf{X}^2\|_{\hat{N}^{\delta,p}}) \geq 1$ such that

$$\frac{\rho_{(\delta-1/p)\text{-Hö};[s,t]}^{(k)} (\mathbf{X}^1, \mathbf{X}^2)}{\tilde{\omega}^{(k)}(s, t)} \leq C.$$

In particular, we have

$$|\pi_k(\mathbf{X}_{s,t}^1 - \mathbf{X}_{s,t}^2)| \leq C \left(\sum_{j=1}^k \rho_{\hat{N}^{\delta,p};[s,t]}^{(j)}(\mathbf{X}^1, \mathbf{X}^2)^{\frac{p}{k}} \right)^{\frac{k}{p}} |t - s|^{(\delta - \frac{1}{p})k},$$

which implies (3.12) for level k and the proof is complete. \square

Combining the local equivalence of the inhomogeneous distances $\rho_{\tilde{V}^{\delta,p}}$ and $\rho_{\hat{N}^{\delta,p}}$ (Lemmas 3.6 and 3.7) with the local Lipschitz continuity of the Itô–Lyons map (3.5) with respect to $\rho_{\tilde{V}^{\delta,p}}$ (Theorem 3.3), we deduce same continuity result with respect to $\rho_{\hat{N}^{\delta,p}}$:

Corollary 3.8. *Let $\delta \in (0, 1)$ and $\gamma, p \in (1, \infty)$ be such that $\delta > 1/p$ and $\gamma > 1/\delta$. The Itô–Lyons map*

$\Phi: \mathbb{R}^m \times \text{Lip}^\gamma \times \hat{N}^{\delta,p}([0, T]; G^{\lfloor 1/\delta \rfloor}(\mathbb{R}^n)) \rightarrow \hat{N}^{\delta,p}([0, T]; \mathbb{R}^m)$ *via* $\Phi(y_0, V, \mathbf{X}) := Y$, *where Y denotes the solution to controlled differential equation (3.4) given the input (y_0, V, \mathbf{X}) , is locally Lipschitz continuous with respect to inhomogeneous Nikolskii-type distance $\rho_{\hat{N}^{\delta,p}}$.*

ACKNOWLEDGMENT

Both authors are grateful for the excellent hospitality of the Hausdorff Research Institute for Mathematics, where the work was initiated.

REFERENCES

- [Aid16] Shigeki Aida, *Reflected rough differential equations*, Stochastic Process. Appl. **125** (2015), no. 9, 3570–3595. MR3357620
- [BLS06] Gérard Bourdaud, Massimo Lanza de Cristoforis, and Winfried Sickel, *Superposition operators and functions of bounded p -variation*, Rev. Mat. Iberoam. **22** (2006), no. 2, 455–487. MR2294787
- [DN99] Richard M. Dudley and Rimas Norvaiša, *Differentiability of six operators on non-smooth functions and p -variation*, Lecture Notes in Mathematics, vol. 1703, Springer-Verlag, Berlin, 1999. With the collaboration of Jinghua Qian. MR1705318
- [FGGR16] Peter K. Friz, Benjamin Gess, Archil Gulisashvili, and Sebastian Riedel, *The Jain–Monrad criterion for rough paths and applications to random Fourier series and non-Markovian Hörmander theory*, Ann. Probab. **44** (2016), no. 1, 684–738. MR3456349
- [FH14] Peter K. Friz and Martin Hairer, *A course on rough paths*, Universitext, Springer, Cham, 2014. With an introduction to regularity structures. MR3289027
- [Fri05] Peter K. Friz, *Continuity of the Itô-map for Hölder rough paths with applications to the support theorem in Hölder norm*, Probability and partial differential equations in modern applied mathematics, IMA Vol. Math. Appl., vol. 140, Springer, New York, 2005, pp. 117–135. MR2202036
- [FV06] Peter Friz and Nicolas Victoir, *A variation embedding theorem and applications*, J. Funct. Anal. **239** (2006), no. 2, 631–637. MR2261341
- [FV10] Peter K. Friz and Nicolas B. Victoir, *Multidimensional stochastic processes as rough paths*, Cambridge Studies in Advanced Mathematics, vol. 120, Cambridge University Press, Cambridge, 2010. Theory and applications. MR2604669
- [Gub04] M. Gubinelli, *Controlling rough paths*, J. Funct. Anal. **216** (2004), no. 1, 86–140. MR2091358
- [LCL07] Terry J. Lyons, Michael Caruana, and Thierry Lévy, *Differential equations driven by rough paths*, Lecture Notes in Mathematics, vol. 1908, Springer, Berlin, 2007. Lectures from the 34th Summer School on Probability Theory held in Saint-Flour, July 6–24, 2004; With an introduction concerning the Summer School by Jean Picard. MR2314753

- [Leo09] Giovanni Leoni, *A first course in Sobolev spaces*, Graduate Studies in Mathematics, vol. 105, American Mathematical Society, Providence, RI, 2009. MR2527916
- [LQ02] Terry Lyons and Zhongmin Qian, *System control and rough paths*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2002. Oxford Science Publications. MR2036784
- [LY38] E. R. Love and L. C. Young, *On Fractional Integration by Parts*, Proc. London Math. Soc. (2) **44** (1938), no. 1, 1–35. MR1575481
- [Lyo91] Terry Lyons, *On the nonexistence of path integrals*, Proc. Roy. Soc. London Ser. A **432** (1991), no. 1885, 281–290. MR1116958
- [Lyo94] Terry Lyons, *Differential equations driven by rough signals. I. An extension of an inequality of L. C. Young*, Math. Res. Lett. **1** (1994), no. 4, 451–464. MR1302388
- [Lyo98] Terry J. Lyons, *Differential equations driven by rough signals*, Rev. Mat. Iberoamericana **14** (1998), no. 2, 215–310. MR1654527
- [MS61] J. Musielak and Z. Semadeni, *Some classes of Banach spaces depending on a parameter*, Studia Math. **20** (1961), 271–284. MR0137985
- [Pee76] Jaak Peetre, *New thoughts on Besov spaces*, Mathematics Department, Duke University, Durham, N.C., 1976. Duke University Mathematics Series, No. 1. MR0461123
- [PT16] David J. Prömel and Mathias Trabs, *Rough differential equations driven by signals in Besov spaces*, J. Differential Equations **260** (2016), no. 6, 5202–5249. MR3448778
- [Rie10] Friedrich Riesz, *Untersuchungen über Systeme integrierbarer Funktionen* (German), Math. Ann. **69** (1910), no. 4, 449–497. MR1511596
- [Ros09] Mathieu Rosenbaum, *First order p -variations and Besov spaces*, Statist. Probab. Lett. **79** (2009), no. 1, 55–62. MR2483397
- [Roy93] Bernard Roynette, *Mouvement brownien et espaces de Besov* (French, with English summary), Stochastics Stochastics Rep. **43** (1993), no. 3-4, 221–260. MR1277166
- [Sim90] Jacques Simon, *Sobolev, Besov and Nikol'skii fractional spaces: imbeddings and comparisons for vector valued spaces on an interval*, Ann. Mat. Pura Appl. (4) **157** (1990), 117–148. MR1108473
- [Ter67] A. P. Terehin, *Integral smoothness properties of periodic functions of bounded p -variation* (Russian), Mat. Zametki **2** (1967), 289–300. MR0223512
- [Tri10] Hans Triebel, *Theory of function spaces*, Modern Birkhäuser Classics, Birkhäuser/Springer Basel AG, Basel, 2010. Reprint of 1983 edition [MR0730762]; Also published in 1983 by Birkhäuser Verlag [MR0781540]. MR3024598
- [Wie24] Norbert Wiener, *The Quadratic Variation of a Function and its Fourier Coefficients*, J. of Math. and Physics **3** (1924), no. 2, 72–94.
- [You36] L. C. Young, *An inequality of the Hölder type, connected with Stieltjes integration*, Acta Math. **67** (1936), no. 1, 251–282. MR1555421
- [Zäh98] M. Zähle, *Integration with respect to fractal functions and stochastic calculus. I*, Probab. Theory Related Fields **111** (1998), no. 3, 333–374. MR1640795
- [Zäh01] M. Zähle, *Integration with respect to fractal functions and stochastic calculus. II*, Math. Nachr. **225** (2001), 145–183. MR1827093

TECHNISCHE UNIVERSITÄT BERLIN AND WEIERSTRASS INSTITUTE BERLIN, GERMANY

EIDGENÖSSISCHE TECHNISCHE HOCHSCHULE ZÜRICH, SWITZERLAND

Current address: Mathematical Institute, University of Oxford, Oxford OX2 6GG, United Kingdom