

Feature

A clinician's guide to probabilistic suicide risk prediction tools: cautions and pitfalls

Alastair Cockburn and Matthew Large

Summary

There are a growing number of new tools designed to predict suicide risk. One, OxSATS, developed in Oxford (UK) using Swedish data, produces a probabilistic risk of suicide in people who have self-harmed. It is accompanied by a web-based calculator, and states that it can 'accurately predict 12-month risk of suicide'. It represents a departure from longstanding research arguing that risk prediction provides insufficient information to be clinically useful.

We analyse the use of OxSATS from a clinician's perspective using eight illustrative vignettes. For each, we use the OxSATS online tool to calculate the 12-month risk of suicide and consider how clinicians might interpret or act on the results. We highlight several potential harms to patients arising from the tool's use.

In our discussion, we explore broader limitations of OxSATS and similar tools, some of which are insidious. These tools can shift resources towards perceived higher-risk patients, often older men, diverting attention away from prevention, younger women and even the treatment of mental illness. Their reductionist approach

misunderstands the complexity and stochastic nature of suicide. Tools tend to be disliked by patients and can subvert a clinician's role away from helping patients, towards mitigating perceived risk.

We conclude that tools such as OxSATS should be treated with significant caution and require careful scrutiny before being considered for clinical use. At present, psychosocial assessments and understanding patients' narratives remain at the heart of good care for suicidal patients.

Keywords

Suicide; risk assessment; prognostic/prediction modelling; self-harm; risk prediction.

Copyright and usage

© The Author(s), 2025. Published by Cambridge University Press on behalf of Royal College of Psychiatrists. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Suicide prediction tools, traditionally pen-and-paper tests that derive a score by the accumulation of suicide risk factors, have been used for decades in an attempt to identify patients at particularly significant risk of suicide.^{1–4} Many researchers and some healthcare systems have concluded that suicide risk prediction tools do not provide sufficient information to be clinically useful,^{5–9} chiefly because they produce far too many false positives and miss too many future suicides.^{10,11} In recent years a new wave of tools, some assisted by machine learning, have been published^{12–14} and advocated for.¹⁵

The appeal of such tools – the promise to identify which patients are likely to go on to die by suicide so that they can be targeted for treatment – appears clear. However, these tools also have a number of limitations and hidden dangers. Here, we explore what clinicians might make of the next generation of tools by considering one of the recently published and accessible suicide risk assessment scales, OxSATS.¹³ We do so using eight plausible but fictional clinical vignettes.

OxSATS

OxSATS is a tool developed to assess risk of suicide in people who have presented to an accident and emergency department with self-harm. It uses 11 items (Table 1), determined by a model from 35 possible items, to produce a probabilistic (percentage) estimate of suicide for each person within 6 and 12 months. It was developed and tested on the record data of 53 172 patients in Sweden between 2008 and 2012.

OxSATS can be completed quickly, does not rely on a detailed assessment and is objective. What is new about OxSATS is that, rather than producing risk categories, it provides a probability estimate expressed as percentage suicide mortality within 6 and 12 months, for each patient. This score can be derived using a freely available online calculator.¹³ OxSATS selects variables through stepwise selection rather than machine learning. Unlike some machine learning tools, it is transparent in letting us see how risk is

scored. It is designed to be used alongside clinicians as a clinical adjunct, not as a replacement. However, the authors suggest that it could be used to allocate treatment to those most at risk.

While OxSATS' statistical performance is not the focus of this paper, and no single statistic captures the value of a tool, its predictive accuracy is comparable to older methods when thresholds are applied; such thresholds would probably be required for use in clinical practice. Using a 2% risk threshold, OxSATS' sensitivity (43%) and specificity (81%) closely match those in Pokorny's 1983 study (45 and 84%, respectively), although OxSATS has a lower positive predictive value (PPV) (2.5 v. 3.7%).¹ OxSATS' sensitivity here (43%) means that more than half of those who died by suicide fell below the threshold; at a 1% threshold, OxSATS' odds ratio (5.36) aligns with that of a previous meta-analysis (4.84).¹⁶ OxSATS' area under the curve (AUC) score (0.75) is similar to that of traditional tools (0.71).⁵

Vignettes

In Box 1 we present eight fictionalised clinical vignettes of typical patients who present to accident and emergency departments with self-harm. None require overnight admission to a medical bed. The information is brief, as might be recorded in accident and emergency triage, but has sufficient data to calculate OxSATS scores. For each of the presentations we have recorded the 12-month estimated probability of suicide according to OxSATS. A discussion follows about the probabilistic estimates produced by OxSATS and how they might affect patients, clinicians and the systems in which they are used.

Results

Once seen by clinicians, risk scores will seem salient, and most clinicians would accept that probability estimates generated by

Table 1 Items used by OxSATS

Domains assessed	Factors assessed in each domain
Demographics	Age Gender (male/female)
Severity of method	Hanging, strangulation or suffocation Use of psychotropics Overnight medical admission
Substance use	Current or lifetime alcohol use Current or lifetime drug use
Psychiatric history	Psychiatric diagnosis (yes/no) Psychotropic medication (past 3 months) Prior self-harm (ever) Prior self-harm (past 12 months)

Box 1 Case vignettes of eight fictionalised patients and their OxSATS scores**Vignette 1**

A 13-year-old girl who self-harms and has emotional dysregulation but no psychiatric diagnosis, and is absent from school. Overdoses on 35 paracetamol with suicidal intent.

Estimated suicide probability in 12 months, 0.0%

Vignette 2

A 34-year-old woman with EUPD and lifelong self-harm worsening over the past 12 months presents with vertical cuts to her wrists with suicidal intent.

Estimated suicide probability in 12 months, 0.6%

Vignette 3

A 22-year-old man who cut his wrists in response to transient auditory hallucinations. He has no current diagnosis and no history of self-harm, but has dropped out of university and is struggling with relationships.

Estimated suicide probability in 12 months, 0.3%

Vignette 4

A 54-year-old man, experiencing marital difficulties and currently living away from partner and two children. No psychiatric history prior to his GP diagnosing new-onset depression and alcohol use disorder; started on sertraline. Impulsive and regretted overdose of his sertraline.

Estimated suicide probability in 12 months, 3.0%

Vignette 5

A 65-year-old man with bipolar, no previous self-harm, overdosing on his prescribed lithium with suicidal intent.

Estimated suicide probability in 12 months, 3.2%

Vignette 6

A 25-year-old woman with bipolar, no previous self-harm, overdosing on her prescribed lithium with suicidal intent.

Estimated suicide probability in 12 months, 0.9%

Vignette 7

A 48-year-old woman with depression, previous self-harm and alcohol use disorder presents with an overdose of her antidepressants.

Estimated suicide probability in 12 months, 1.7%

Vignette 8

A 45-year-old man with schizophrenia, on risperidone, presents with chaotic suffocation attempt after stopping risperidone. Historical substance use disorder and self-harm and unpleasant previous psychiatric hospital admission, but stable for many years.

Estimated suicide probability in 12 months, 7.7%

EUPD, emotionally unstable personality disorder (ICD-11: 6D11.5); GP, general practitioner.

Vignette 1: false reassurance, risk of storing up problems for the future

This 13-year-old girl clearly needs a careful assessment to understand, and help her understand, the causes of her distress. Would knowing that her 12-month risk score is 0.0% help her, help her clinician or help her access the support she needs? OxSATS authors suggest that, in future, tools could 'act as a screen for more detailed assessment' and 'allow for more efficient allocation of clinical resources'.¹³ Should clinicians take this to mean that the patient in vignette 1 might not need a full psychosocial assessment or follow-up support? Even if her immediate risk of suicide is low, her distress is real and the need for intervention urgent. The focus of mental healthcare should be on her many and immediate needs.

This case demonstrates the risk that such a tool would de-prioritise 'lower-risk' patients. Unseen and untreated, will her risk stay at 0%? Today's lower-risk patients will become tomorrow's 'higher-risk' patients,¹⁷ let alone tomorrow's unhappy person failing to thrive. As such, in cases like these the use of OxSATS may provide unwarranted reassurance and increase the likelihood of leaving illness untreated and distress unexplored, only to present more severely in the future.

Vignette 2: underestimation of groups and vulnerability to exclusion culture

This vignette represents a patient diagnosed with emotionally unstable personality disorder (EUPD) experiencing a crisis. At present, around 10% of patients with EUPD die by suicide in their lifetime.¹⁸ Clinicians may feel that 0.6% represents a significant underestimation of risk for this patient, who needs careful assessment, validation, psychoeducation and referral to appropriate treatment pathways. Furthermore, although 0.6% is below the median score in the OxSATS sample, it translates as 65 times the 2021 global suicide rate (9.1/100 000).¹⁹

This vignette illustrates a danger regarding the way in which OxSATS scores could be used in overstretched services. When attempting onward referral, would her relatively low suicide risk score be used as an excuse for exclusion?²⁰ We fear that it may be.

Furthermore, would the illusions of certainty, precision and objectivity from the score particularly affect less experienced clinicians who may rely on the number and overlook or ignore worrying aspects of the presentation?

Vignette 3: missed opportunities, prioritising perceived suicide risk over other debilitating mental illness experiences

The young man here probably has emergent schizophrenia. He too needs careful assessment to identify, explain and treat the underlying psychiatric cause. Were OxSATS to be used as a screen for more detailed assessment, he may not get that assessment at all. The role of psychiatric assessment in accident and emergency departments is about much more than predicting risk. In this case, assessment serves as a hugely important screen for a treatable mental illness. As in vignette 1, untreated, this patient's risk of suicide is only likely to rise.

However, the most pressing problems faced by the patient in vignette 3 are not suicide – he has significant functional impairments and is falling off his developmental trajectory. Identifying the underlying cause, and understanding and addressing his social withdrawal and academic difficulties, can emerge only in a more complete, relational, assessment.

OxSATS provide some information about future suicide risk. The tool is both intended to, and in all likelihood would, influence clinician decision-making: sometimes nudging clinicians' decision-making in the direction of the tool's score, sometimes overtly influencing or even obligating decisions and actions by clinicians. It is intended to help decision-making, but the actual potential effect on both clinicians and patients requires careful consideration.

Vignettes 4 and 8: priming overtreatment and iatrogenic harm of higher-risk patients

The patient in vignette 4 may well be experiencing an adjustment reaction to a significant social stressor (marital breakdown). He requires an assessment to screen for mental illness, identify strengths and form a mutually agreed plan. The patient in vignette 8 is probably experiencing a psychotic relapse after stopping risperidone, and needs assessment of why he stopped and how and where to restart.

High scores according to OxSATS may alert staff to high-risk individuals who might otherwise be missed. However, this benefit comes with a cost. On seeing either of these OxSATS scores, clinicians, who are already prone to being risk averse,²⁰ may feel obliged to act – or be seen to act – on the high-risk scores. At the most restrictive end this may involve admitting someone to hospital, perhaps coercively or directly against their will. Alternatively, it may involve inappropriate prescribing or interventions and referrals that the patient does not want. Treatment such as involuntary admission may serve to help a patient but may also represent defensive psychiatry,²¹ serving not to benefit that patient but limit a clinician's liability.^{22,23}

Hospitalisation does not prevent suicide, and it may even increase the risk.²⁴ Suicide rates are notably high following discharge from psychiatric hospital and subsequently remain elevated for decades. The suicide rate among discharged patients is around 25 times that in the general population, even over 10 years following discharge.²⁵ Suicide risk following discharge for patients admitted with suicidal thoughts/behaviours is particularly high – around 200 times the baseline rate (2078/100 000).²⁵ The extent to which this is the effect of selecting a very high-risk group, versus hospitalisation at times being harmful, is unclear.²⁶ However, it is clear that the benefits of hospitalisation need to be carefully weighed against its risks, and it is possible that hospitalisation may delay – or even magnify – the high-risk period following self-harm. In this case, admitting the man in vignette 4, perhaps influenced by his OxSATS score, would probably be inappropriate, unhelpful and may even serve to increase, not decrease, his risk of suicide.

As has always been the case, the great majority of so-called higher-risk patients do not go on to die by suicide. We do not have effective, low-side-effect interventions for suicide²⁷ and, if we did, they should not be denied to lower-risk patients, among whom many suicides occur. With our present interventions, it is these higher-risk patients, as in vignette 4, who pay the cost of interventions, including coercive treatment and occasionally harmful hospitalisation, even when their individual likelihood of suicide is low.

Vignettes 5 and 6: systematic de-prioritisation of demographic groups (especially young and adolescent women)

Aside from demographic factors, these are identical presentations. It would be clear to clinicians that vignette 5 carries a higher risk of death by suicide than vignette 6. To what extent should that determine assessment or treatment? If used clinically, the influence of OxSATS on clinicians would probably introduce a systematic gender and age bias, prioritising services towards attending to the risk of older men over the suffering of younger women. Because adolescent women are the fastest-growing demographic with mental illness,²⁸ de-prioritising them would be contentious to say the least. If clinicians and services wanted to introduce such tools, their introduction of such biases would necessitate candid, thoughtful discussions among professionals, patients and health-care funders.

Vignette 7: impact of OxSATS score on patients

With this vignette, what are we to make of her score of 1.7%? What degree of confidence can we have in this estimate? Should we give her this information? And, if so, how should she understand it and what would she make of it? She may feel that the score is high. Perhaps it serves as an important message that she needs change in her life; but perhaps the score implies a mechanical, destined process that disempowers her and takes away her agency. Alternatively, she may feel that the score is low. Perhaps it devalues her feelings and experiences, making her feel unheard. Perhaps it could make her think that the only way she can get the help she needs is to increase her risk score. There are many ways in which communicating this score to patients could be harmful. In truth, we cannot truly know how an algorithmically assigned score would impact a patient in significant distress in the aftermath of a suicide attempt in a hospital emergency department.

Discussion

There are a variety of risk prediction tools published and under development in this area.²⁹ They vary in terms of their setting, the population being screened and the outcomes they predict. In taking people who have presented following self-harm, OxSATS is trying to parse a high-risk, very distressed group in which we argue that all patients, even those scoring 0% risk, require careful assessment and consideration. Importantly, the vignettes demonstrate where tools like OxSATS have the potential to do harm. These also risk overshadowing the role of the clinical assessment of the needs of people who self-harm. That assessment is about much more than attempting to predict the risk of suicide.

Psychiatry's perceived lack of risk prediction tools is sometimes compared unfavourably with the many risk prediction tools in general medicine.¹⁵ A particular comparison made is with cardiovascular disease when using tools such as Framingham or QRISK,¹⁵ which give patients a percentage risk of having a cardiovascular event within 10 years.³⁰ The closest analogy would be that a suicide attempt (rather than death by suicide) is akin to a cardiovascular event. OxSATS is designed for use following a suicide attempt. This is like using a cardiac risk score after a patient experiences a heart attack.

More importantly, we think the comparison with cardiovascular disease fundamentally misunderstands what suicide is. Cardiovascular disease, in addition to well-described risk factors, has a known, well-understood final pathway to an event, which follows biological and largely predictable processes (insufficient perfusion of tissue, usually via a clot). No such comparison exists for suicidal thoughts and behaviours. These have innumerable pathways leading to them and are part of a complex syndrome with biological, psychological, social and existential dimensions, many origins and huge stochasticity.^{31,32} As such, likening cardiovascular disease to suicide is inappropriate and a category error. Furthermore, it carries implications of disempowering people as being at the mercy of their neurobiology or destiny, rather than promoting their agency.

Comparison of these suicide risk tools to QRISK is also currently of limited use. QRISK allows patients to explore how their risk of cardiovascular events would alter if they were to make changes – lose weight, lower blood pressure, stop smoking, etc. OxSATS is purely predictive and cannot do this. If vignette 5 or 6 tried to use the tool in this way, they would perversely find that stopping their psychotropic (lithium) had reduced their risk score, despite randomised trials showing that lithium reduces suicide risk.³³ OxSATS does not help clinicians in the search with our patients for modifiable factors in their presentation.

Importantly, there is no equivalent of a statin for suicide, no low-side-effect intervention that can be safely and effectively given to all higher-risk patients. In contrast, specific interventions to treat suicidal thoughts and behaviours are limited, with small effect sizes across outcomes, a situation largely unchanged for over 50 years.²⁷ Classifying people by perceived risk has little value if we lack interventions for those identified as being at greater risk.

Suicide is as much a cultural phenomenon as a medical one. To what extent can a Swedish sample from 2008 to 2012 be applied to a different country today? World Health Organization data from Sweden and the UK in 2012 show significant differences in suicide rates.³⁴ Overall age-adjusted rates (per 100 000) were nearly twice as high in Sweden (11.1 v. 6.2), with a particular increase in suicide in Swedish versus UK women ($\times 2.35$) and different rates in each country by age distribution.³⁴ OxSATS was validated on a sample of the same Swedish population on which it was developed.¹³ Given the difference in populations and interventions, its performance on patients in other countries is likely to be limited. In practice, if applied to the UK, the scores are likely to overestimate risk in most people, particularly women.

OxSATS is based on observational data; the patients on which it was trained would have received assessments and interventions routinely. As such, lower-risk patients may be lower risk because they responded well to the intervention. It is not possible to justify removing the intervention based on a model developed from such observational data.

Tools such as OxSATS are usually deliberately designed to achieve high reliability by including only easily collected, unambiguous, quantitative data. However, a person's true risk is likely formed of a combination of (a) nomothetic data – generalisable, group-level data, (b) idiographic data – based on the unique, variable circumstances of each individual and (c) data that cannot be known or accessed. The acknowledgement that tools can only be adjuncts to clinicians is partially because tools cannot easily access idiographic data. However, even the best assessment will leave unknowns that contribute to a patient's true risk. Five minutes after leaving accident and emergency, a person may go through a relationship breakdown, or have an old friend unexpectedly reach out, or find their home burgled, or receive good news on a much-longed-for job. None of us can predict these unknowns, but clinicians can get a sense of the idiographic data and the patient in front of them. Experienced clinicians, who also implicitly use idiographic data, outperform some predictive tools in physical healthcare.³⁵ Whether clinicians outperform suicide and self-harm tools is very hard to test, because clinicians will inevitably do all they can to reduce risk in patients they think are high-risk, in effect trying to prove themselves wrong. Given this, we should keep an open mind that good clinicians can and will outperform these tools.

As discussed in vignette 7, tools like this risk disempowering patients and tend to be disliked by them.^{7,11} We must be conscious too that the focus on suicide risk over and above other aspects of mental illness can disempower clinicians. We become risk predictors and managers, rather than professionals using our experience and skills to develop a therapeutic relationship, diagnose and treat mental illness and help people in distress. We must guard against a culture that renders patients bearers of risk to be feared, rather than people whom we have the opportunity to help.^{36,37}

Alternative

At present, the key intervention for people who have self-harmed, attempted suicide or are acutely suicidal is an individualised therapeutic assessment. There is convergence on this point from

Box 2 Suggestions for areas to focus on during a psychosocial assessment, drawing on Large et al and Hawton et al^{7,10}

Key areas of a psychosocial assessment


Form a **therapeutic relationship** with the patient in front of you.
Is there a **diagnosis** and can it guide treatment?
Are there **modifiable factors** in the presentation?
Draw out the patient's **strengths**, resources and values.
Look for opportunities to instil **reasonable hope**.

colleagues from different perspectives of suicide research.^{7,10,38} Gibbons describes suicide as an 'action take(ing) the place of feeling'. She posits that one role of professionals is to help patients 'put the feelings into words to encourage expression of pain', ultimately aiming to 'reduce the risk of action'.³⁶ Each presentation of a suicidal person is unique, with different complexities, circumstances, risk factors and best approaches to management. Key things for clinicians to bear in mind during assessments are shown in Box 2.

There are other important areas for clinicians and researchers to focus on, such as: how to build better therapeutic relationships; training clinical skills in helping patients manage strong emotions and crises; carefully assessing the effectiveness of newer interventions such as safety plans; incorporating patient experiences into practice; reducing burnout in clinicians; working to reduce bureaucracy so that more clinician time can be spent with patients and less completing documentation (just 17% of nursing time on in-patient wards was spent on direct care in one study³⁹); and, as a society, how to help the 74% of people who are not known to psychiatric services in the year prior to suicide.⁴⁰

It may be that, in the future, new tools – perhaps aided by machine learning or artificial intelligence – will allow us to identify certain patients, in certain settings, at particularly high or low risk of suicide. However, the impact and value of such tools would need to be very carefully considered, with a number of pitfalls to avoid.⁴¹ As we have argued here, implementation of tools may even be harmful and the role of clinicians is much broader than risk prediction.

Presently it remains the case that tools are not able to give clinicians sufficient meaningful information regarding who will go on to die by suicide to guide assessment or management. Our conclusions here align with and support the current National Institute for Health and Care Excellence guidance to that effect.⁶ Suicidal actions are expressions of distress that demand our respect and attention. Every patient who presents with self-harm and suicidal behaviour should be met with a comprehensive, sensitive, patient-centred, psychosocial assessment, with onward treatment decisions made collaboratively on the basis of person-specific needs. The efforts of many healthcare services around the world to deliver this standard of care should be celebrated.

Alastair Cockburn , Department of Psychiatry, University of Oxford, UK; and Oxford Health NHS Foundation Trust, Warneford Hospital, Oxford, UK; **Matthew Large**, Discipline of Psychiatry and Mental Health, UNSW Medicine and Health, Kensington, NSW, Australia

Correspondence: Alastair Cockburn. Email: alastair.cockburn1@oxfordhealth.nhs.uk

First received 25 Apr 2025, final revision 24 Jul 2025, accepted 22 Sep 2025

Data availability

The data produced by the OxSATS tool can be recreated at its website (at the time of publishing), <https://oxrisk.com/oxsats/>, or are available on request from the corresponding author (A.C).

Acknowledgements

A.C. thanks the following colleagues for comments on a draft: Dr Karen Lascelles, psychiatry nurse consultant; Julie Cooper, expert by experience; Fiona Brand, psychiatric liaison and research nurse; and Dr Kezia Lange, MRCPsych.

Author contributions

Both authors contributed to the drafting and final version of the manuscript, and approved it for submission.

Funding

This research received no specific grant from any funding agency, commercial or not-for-profit sectors. A.C. is supported by an Academic Clinical Fellowship funded by the University of Oxford Department of Psychiatry.

Declaration of interest

A.C., none. M.L. has provided evidence in legal matters following suicides.

References

- Pokorny AD. Prediction of suicide in psychiatric patients. Report of a prospective study. *Arch Gen Psychiatry* 1983; **40**: 249–57.
- Motto JA. An integrated approach to estimating suicide risk. *Suicide Life Threat Behav* 1991; **21**: 74–89.
- Beck AT, Brown GK, Steer RA, Dahlsgaard KK, Grisham JR. Suicide ideation at its worst point: a predictor of eventual suicide in psychiatric outpatients. *Suicide Life Threat Behav* 1999; **29**: 1–9.
- Steed S, Kapur N, Webb R, Applegate E, Stewart SLK, Hawton K, et al. The development of a population-level clinical screening tool for self-harm repetition and suicide: the ReACT Self-Harm Rule. *Psychol Med* 2012; **42**: 2383–94.
- Steed S, Quinlivan L, Nowland R, Carroll R, Casey D, Clements C, et al. Accuracy of risk scales for predicting repeat self-harm and suicide: a multicentre, population-level cohort study using routine clinical data. *BMC Psychiatry* 2018; **18**: 113.
- National Institute for Health and Care Excellent. *Self-Harm: Assessment, Management and Preventing Recurrence. NICE Guideline NG225*. NICE, 2022.
- Hawton K, Lascelles K, Pitman A, Gilbert S, Silverman M, Keith P. Assessment of suicide risk in mental health practice: shifting from prediction to therapeutic assessment, formulation, and risk management. *Lancet Psychiatry* 2022; **9**: 922–8.
- Belsher BE, Smolenski DJ, Pruitt LD, Bush NE, Beech EH, Workman DE, et al. Prediction models for suicide attempts and deaths: a systematic review and simulation. *JAMA Psychiatry* 2019; **76**: 642–51.
- Ribeiro JD, Franklin JC, Fox KR, Bentley KH, Kleiman EM, Chang BP, et al. Self-injurious thoughts and behaviors as risk factors for future suicide ideation, attempts, and death: a meta-analysis of longitudinal studies. *Psychol Med* 2016; **46**: 225–36.
- Large MM, Ryan CJ, Carter G, Kapur N. Can we usefully stratify patients according to suicide risk? *BMJ* 2017; **359**: j4627.
- Graney J, Hunt IM, Quinlivan L, Rodway C, Turnbull P, Gianatsi M, et al. Suicide risk assessment in UK mental health services: a national mixed-methods study. *Lancet Psychiatry* 2020; **7**: 1046–53.
- Kessler RC, Hwang I, Hoffmire CA, McCarthy JF, Petukhova MV, Rosellini AJ. Developing a practical suicide risk prediction model for targeting high-risk patients in the Veterans Health Administration. *Int J Methods Psychiatr Res* 2017; **26**: e1575.
- Fazel S, Vazquez-Montes MDLA, Molero Y, Runeson B, D'Onofrio BM, Larsson H, et al. Risk of death by suicide following self-harm presentations to healthcare: development and validation of a multivariable clinical prediction rule (OxSATS). *BMJ Mental Health* 2023; **26**: e300673.
- Kennedy CJ, Kearns JC, Geraci JC, Gildea SM, Hwang IH, King AJ, et al. Predicting suicides among us army soldiers after leaving active service. *JAMA Psychiatry* 2024; **81**: 1215–24.
- Seyedsalehi A, Fazel S. Suicide risk assessment tools and prediction models: new evidence, methodological innovations, outdated criticisms. *BMJ Ment Health* 2024; **27**: e300990.
- Large M, Kanesson M, Myles N, Myles H, Gunaratne P, Ryan C. Meta-analysis of longitudinal cohort studies of suicide risk assessment among psychiatric patients: heterogeneity in results and lack of improvement over time. *PLOS One* 2016; **11**: e0156322.
- Gianatsi M, Burns H, Hunt IM, Ibrahim S, Windfuhr K, While D, et al. Treatment of mental illness prior to suicide: a national investigation of 12,909 patients, 2001–2016. *Psychiatr Serv* 2020; **71**: 772–8.
- Paris J. Suicidality in borderline personality disorder. *Medicina* 2019; **55**: 223.
- World Health Organization. *World Health Statistics 2024: Monitoring Health for the SDGs, Sustainable Development Goals*. WHO, 2024 (<https://iris.who.int/bitstream/handle/10665/376869/9789240094703-eng.pdf?sequence=1> [cited 10 Apr 2025]).
- Beale C. Magical thinking and moral injury: exclusion culture in psychiatry. *BJPsych Bull* 2022; **46**: 16–9.
- Reuveni I, Pelov I, Reuveni H, Bonne O, Canetti L. Cross-sectional survey on defensive practices and defensive behaviours among Israeli psychiatrists. *BMJ Open* 2017; **7**: e014153.
- Riolo A, Babici F. An ethical perspective. *Eur Psychiatry* 2017; **41**: S582.
- Brown J, Rayne JT. Some ethical considerations in defensive psychiatry: a case study. *Am J Orthopsychiatry* 1989; **59**: 534–41.
- Large M, Ryan C, Walsh G, Stein-Parbury J, Patfield M. Nosocomial suicide. *Austr Psychiatry* 2014; **22**: 118–21.
- Chung DT, Ryan CJ, Hadzi-Pavlovic D, Singh SP, Stanton C, Large MM. Suicide rates after discharge from psychiatric facilities: a systematic review and meta-analysis. *JAMA Psychiatry* 2017; **74**: 694–702.
- Large MM, Kapur N. Psychiatric hospitalisation and the risk of suicide. *Br J Psychiatry* 2018; **212**: 269–73.
- Fox KR, Huang X, Guzmán EM, Funsch KM, Cha CB, Ribeiro JD, et al. Interventions for suicide and self-injury: a meta-analysis of randomized controlled trials across nearly 50 years of research. *Psychol Bull* 2020; **146**: 1117–45.
- McManus S, Bebbington P, Jenkins R, Brugha T (eds). *Adult Psychiatric Morbidity Survey: Survey of Mental Health and Wellbeing, England, 2014*. NHS Digital, 2016 (<https://digital.nhs.uk/data-and-information/publications/statistical/adult-psychiatric-morbidity-survey/adult-psychiatric-morbidity-survey-of-mental-health-and-wellbeing-england-2014>).
- Pigoni A, Delvecchio G, Turtulici N, Madonna D, Pietrini P, Cecchetti L, et al. Machine learning and the prediction of suicide in psychiatric populations: a systematic review. *Transl Psychiatry* 2024; **14**: 140.
- Hippisley-Cox J, Coupland C, Brindle P. Development and validation of QRISK3 risk prediction algorithms to estimate future risk of cardiovascular disease: prospective cohort study. *BMJ* 2017; **357**: j2099.
- Mental and Behavioural Disorders, Department of Mental Health, World Health Organization. *Preventing Suicide: A Resource for General Physicians*. WHO, 2000 (https://apps.who.int/iris/bitstream/handle/10665/67165/WHO_MNH_MBD_00.1.pdf [cited 28 Mar 2025]).
- Soper CA, Ocejo PM, Large MM. On the randomness of suicide: an evolutionary, clinical call to transcend suicide risk assessment. In *Evolutionary Psychiatry: Current Perspectives on Evolution and Mental Health* (eds R Abed, P St John-Smith): 134–52. Cambridge University Press, 2022.
- Cipriani A, Hawton K, Stockton S, Geddes JR. Lithium in the prevention of suicide in mood disorders: updated systematic review and meta-analysis. *BMJ* 2013; **346**: f3646.
- World Health Organization. *Preventing Suicide: A Global Imperative*. WHO, 2014 (<https://www.who.int/publications/i/item/9789241564779> [cited 28 Mar 2025]).
- Penaloza A, Verschuren F, Meyer G, Quentin-Georget S, Soulie C, Thys F, et al. Comparison of the unstructured clinician gestalt, the wells score, and the revised Geneva score to estimate pretest probability for suspected pulmonary embolism. *Ann Emerg Med* 2013; **62**: 117–24.
- Gibbons R. Eight 'truths' about suicide. *BJPsych Bull* 2024; **48**: 350–4.
- Appel JM. The archives of psychiatry commentary helping vs. preventing harm: reversing mission creep in psychiatry. *Arch Psychiatry* 2024; **2**: 59–62.
- Kapur N, Tham SG, Turnbull P, Richards N, Rodway C, Clements C, et al. Quality improvement for suicide prevention and self-harm intervention: addressing the implementation gap and saving lives. *Lancet Psychiatry* 2025; **12**: 303–9.
- Glantz A, Sunnqvist C. The time, places, and activities of nurses in a psychiatric inpatient context: a time and motion study with a time-geographic perspective. *Issues Ment Health Nurs* 2023; **44**: 387–95.
- Health Quality Improvement Partnership. *National Confidential Inquiry into Suicide and Safety in Mental Health. Annual Report 2024: UK Patient and General Population Data 2011–2021*. University of Manchester, 2024 (<https://sites.manchester.ac.uk/ncish/reports/annual-report-2024/> [cited 28 Mar 2025]).
- Kirtley OJ, van Mens K, Hoogendoorn M, Kapur N, de Beurs D. Translating promise into practice: a review of machine learning in suicide research and prevention. *Lancet Psychiatry* 2022; **9**: 243–52.