Biochemical studies on *Pseudomonas* spp. ethylene-forming enzymes



A thesis presented to the Board of the Faculty of Mathematical, Physical, and Life Sciences in partial fulfilment of the requirements for the degree of Doctor of Philosophy at the University of Oxford

Siddhant Dhingra

St. Edmund Hall, University of Oxford

Hilary Term 2025

To my parents,

Karan & Richa Dhingra,

my guardian angels.

Your love and light live on in me in every step.

bravery builds empires, so be brave and take the first step.

Acknowledgments

Firstly, I would like to thank Prof. Christopher J. Schofield for giving me the opportunity to join such a vibrant research group and introducing me to the world of 2OG oxygenases. I am deeply grateful for his supervision, mentorship, and guidance across multiple projects.

I would like to thank Dr. Lennart Brewitz for his mentorship, friendship, and all his help from troubleshooting experiments to manuscript revisions. Thanks to him for pushing me when I was almost giving up or losing direction. From my first-ever NMR reaction in 2021 to proofreading this thesis, I am beyond grateful to him for his supervision even over the weekends and late-night hours. I am also grateful to Chris & Lennart for their support and understanding during some of the toughest personal moments I faced during the DPhil. I would also like to thank the Ineos Oxford Institute for Antimicrobial Research for funding my DPhil studies and providing a dynamic platform for scientific exchange.

I would like to thank Dr. Zhihong Zhang for training me in microbiology and molecular biology techniques, for instilling in me the passion for exploring ethylene biosynthesis and biochemical nuances of the enzymes involved.

I would like to extend my gratitude to Prof. Christina Redfield, Prof. Tim Claridge, Dr. Nader Amin, Dr. Coral Mycroft, Dr. James Montgomery, and Dr. Harry Mackenzie for the keeping the world of NMR running. I am particularly grateful to Tim and Nader for patiently training me on the NMR machines and answering countless questions. Thanks to Christina, Coral, James, and Harry for helping me with experimental setups and troubleshooting. I am beyond grateful for all the patience and understanding Christina and Coral had when I needed more NMR time (which was often!).

I owe massive thanks to Dr. Mark Allen for his invaluable help with solving crystals structures and helping us optimise (and reoptimise) crystallisation conditions. To Dr. Anthony Tumber and Dr. Eidarus Salah for always supporting me and guiding me through different stages of my DPhil.

I would like to thank the wonderful people I had the opportunity to collaborate with and train in the lab during my DPhil, Mr. Yihong Sun, Mr. Luke Harrington, Ms. Ilinca Farcas, Mr. Ethan Sip, Ms. Sakshi Sharma - who will forever stay my favourite *et al*. The EFE projects would not have progressed as far and concluded without the dedication of Yihong, Luke, and Ethan. The HIS1 project, too, matured within just two years thanks to the hard work of Ilinca

and Sakshi. Thanks to Mr. Faisal Alshref for our collaboration on HPPD and HIS1 projects; this would not have been possible without his friendship and curiosity. To all of them, thank you for sharing the passion I had for my work and taking the projects to new heights.

I would like to extend my gratitude to Prof. James McCullagh and Ms. Isabel Legge for all their advice on the MS aspects of my projects. Thanks to Izzy for collaborating on the isotope characterisation of KMBA reaction products and for her help with data analysis.

I am grateful to Prof. Hiroaki Suga, Dr. Wei-En Huang, Dr. Junshi Chang for welcoming me to their lab at the University of Tokyo for a research placement during my DPhil. Their training and patience introduced me to a whole new world of mRNA display and macrocyclic peptides, enabling the PBP and Ldt screens to come to fruition. I am also grateful to Mr. Junsei Yamashita for a wonderful time in and out of the lab, both in Oxford and Tokyo. Thanks also to Dr. Tika R. Malla and Dr. Mariska de Munnik for introducing me to transpeptidases, to Ms. Christine Arndt for her ongoing work on the Ldt screen; To Dr. Helen Smith, Dr. Monisha Singha, and Dr. Ignacio Tobal Salcedo for collaborating on the PBPs with such patience and generosity. I would also like to thank Ms. Gwendoline Denning, Mrs. Zahn Gowar, Dr. Liz Bentley, and Dr. Adam Hardy for all that they do for us to keep such a massive group up and running, while making it all seem so effortless.

This DPhil would not have been the same without the support of my friends in the CJS group: Dr. Gayatri, Ms. Sakshi Sharma, Mr. Yihong Sun, Dr. Liam Wilson, Ms. Christine Arndt, Ms. Dora Laczi, Ms. Sophia Schoenbauer, Ms. Yicheng Wang, Dr. Patrick Rabe, Dr. Karina Calvopina Tapia, Dr. Monisha Singha, Dr. Lewis Ibbotson, Dr. Tom Corner, and Dr. Tika R. Malla. Thanks to them for making every day full of reasons to smile and motivating me when things seemed too uphill. Special thanks to Liam, Patrick, Karina, and Tika for their guidance and scientific advice.

I am grateful to St. Edmund Hall and the MCR for a fantastic time and Oxford experience. The MCR became one of my happiest places and I am super glad that it happened. Serving on the MCR committees for four years has been among the most enriching parts of my time here. Special thanks to Mr. Duncan Lyster for being an excellent and organised Vice President during my presidency (2024-25) and for supporting me while I had to be away.

I don’t think this journey could have been completed without my friends, who had been instrumental at every step. To ‘Familia’, Kai, Dyuti, Atharva, Nancy, Linda, and Giovanni for helping me settle in Oxford and build a home away from home – 19 Norham Gardens will

forever be one of my fondest memories. To ‘Support System’, Sakshi, Prasita, and Shreya, for all the long-nights, dinners, and all that they came with – I cannot imagine Oxford any other way. To ‘SNS’, Shubham and Nilanjana, for all the choicest memories, tea, and sharing the DPhil journey – Crick Road shall go down in history! To Disha, Seerat, and Aditi, coming all the way from Stephen’s to the UK, it’s been a privilege sharing these experiences and growth with you. To Niharika, for making the weekends happier and organising (?) our trips! Oxford, for me, will always mean love and friendship, and it would have been impossible without this extraordinary bunch — the reason I made it through despite all the challenges of the latter half of my DPhil. I am grateful to Kavneet, Jigyasa, Prachi, Sazia, Shreyansh, Shubh, Roshni, and Anurima for being there for me in some of the toughest times and for forgiving me when I fell short as a friend amidst all the responsibilities and time commitments. It has been a privilege growing up with them and to continue a friendship that has lasted through years and still counting.

Finally, I am forever indebted to my guardian angles. To my dad, the late Mr. Karan Dhingra, whom I lost during my DPhil, for the sacrifices, the vision, and the character he imbued in me. It would not have been possible without his dream of sending his son to Oxford that I would have travelled miles and made him proud. I could not have imagined even a fraction of this if it wasn’t for him as my pillar of support, my rock, and my best friend. To my mom, the late Mrs. Richa Dhingra, whom I lost at the age of seven, for always watching over me and giving me strength when I needed it the most.

I am grateful to my grandparents, Adv. J. K. Dhingra and Mrs. Neelam Dhingra, for being my wings. It has been their upbringing and emphasis on education in my formative years that I had the capability to ask questions and explore my potential. Their love and sacrifice, staying miles away from me, carries me through each day. Thanks to my grandparents, the late Mr. J. C. Sharma and the late Mrs. Rajni Sharma for raising me (and spoiling me!) and watching over me still.

I am grateful to my uncle, the late Mr. Aviral Sharma, and my aunt, Mrs. Jiny Soni, for being friends before relatives and supporting me through these years. I would also like to thank my cousins - my little sisters, Avishi, Yashita, and Saisha - for never letting me feel alone and all their warmth (and chaos!).

Lastly, I thank myself. For taking the road not taken. For the resilience. For the sensitivity. For making it through both the good and the bad.

Declaration

I, Siddhant Dhingra, declare that the thesis I am submitting is entirely my own work except where indicated in the text, caption, footnote, or bibliography. This thesis has not been submitted in whole or in part for any other academic degree or professional qualification.

Date: 01/10/2025

Signature: 

List of publications

This integrated thesis has been based upon the following five manuscripts, my contributions to which are described in the Statement of Authorship at the end of each chapter:

1. **Dhingra, S**.; Zhang, Z.; Lohans, C. T.; Brewitz, L.; Schofield, C. J. Substitution of 2-oxoglutarate alters reaction outcomes of the *Pseudomonas savastanoi* ethylene-forming enzyme. *J. Biol. Chem.* **2024**, *300*, 107546
2. Sun, Y.; **Dhingra, S**.; Allen, M. D.; Cheng, E. S. Z.; Zhang, Z.; Schofield, C. J.; Brewitz, L. Structural and functional validation of *Pseudomonas savastanoi* ethylene forming enzymes reveals flexibility in 2-oxoglutarate binding mode and conformation. **Manuscript under review in** *ChemBioChem*.
3. **Dhingra, S**.; Sun, Y.; Zhang, Z.; Schofield, C. J.; Brewitz, L. Expanding the alkene forming capability of the *Pseudomonas savastanoi* ethylene-forming enzyme. **Manuscript under review in** *ChemCatChem*.
4. **Dhingra, S**.; Harrington, L; Sun, Y.; Legge, I.; Zhang, Z.; McCullagh, J. S.; Schofield, C. J.; Brewitz, L. Oxygenase-catalyzed production of ethylene from the methionine transamination product α - keto - γ - methylthiobutyric acid. **Manuscript in preparation.**
5. **Dhingra, S**.; Sharma, S.; Farcas, I.M.; Alshref, F. M.; Allen, M. D.; Redfield, C.; Schofield, C. J.; Brewitz, L. Crystallographic and biochemical characterisation of HIS1 and OsHSL1 inform on strategies to evade resistance-associated β-triketone herbicide metabolism. **Manuscript in preparation.**

Abstract

2-oxoglutarate (2OG)-dependent oxygenases are non-haem Fe(II) enzymes that catalyse a diverse repertoire of oxidative transformations. The ethylene-forming enzyme from *Pseudomonas savastanoi* (PsEFE) catalyses a bifurcating reaction: the C5 hydroxylation of *L*-arginine coupled to the oxidative decarboxylation of 2OG to succinate and the conversion of 2OG to ethylene and CO_2. This thesis presents biochemical studies on PsEFE and two other plant oxygenases, HIS1 and OsHSL1, involved in conferring herbicide resistance in plants; the research described in this thesis explores the potential applications of 2OG oxygenases in biocatalysis and agriculture.

Chapter 1 introduces 2OG oxygenases and ethylene biosynthesis in a comprehensive literature review. **Chapter 2** discuses substrate scope studies on PsEFE and explores how substitution patterns of 2OG derivatives affect reaction outcomes. **Chapter 3** validates EFEs from two additional *P. savastanoi* strains, 1449B and glycinea PsEFE; this chapter also reveals a crystallographically-observed atypical binding mode of 2OG in 1449B PsEFE.

Chapter 4 expands the work on PsEFE and 2OG derivatives, describing PsEFE active site variants that enable the production of alkenes other than ethylene from substituted 2OG derivatives. **Chapter 5** investigates the interaction of PsEFE with hydrophobic 2-oxoacids, derived from proteinogenic and non-proteinogenic amino acids. This chapter also reports on PsEFE active site variants that enable ethylene production from 2-keto-4-methylthiobutyrate (KMBA) and 2-keto-4-ethylthiobutyrate (KEBA).

Chapter 6 describes biochemical and structural studies of HIS1 and OsHSL1, herbicide resistance enzymes from *Oryza sativa*, which confer resistance to widely-used β-triketone herbicides via hydroxylation. This work highlights how catalytic promiscuity in the form of resistance poses challenges to global food security. **Chapter 7** summarises the results and discusses the biocatalytic potential and promiscuity of 2OG oxygenases.

Collectively, the work described in this thesis advances the biochemical understanding of PsEFE and discusses how the (co)substrate structure and active-site architecture govern reaction partitioning. Guided by substrate scope analyses, targeted mutagenesis, and structural characterisation, this work demonstrates how PsEFE reactivity has the potential to be tuned, improved, and harnessed for biocatalytic and biotechnological applications.

List of abbreviations

Abbreviation	Full form
2OG	2-Oxoglutarate
2YT	2× Yeast Tryptone medium
ACC	1-Aminocyclopropane-1-carboxylate
ACCO	ACC oxidase
AspH	Aspartyl/asparaginyl β-hydroxylase
BSA	Bovine serum albumin
CAS	Clavaminic acid synthase
DAOCS	Deacetoxycephalosporin C synthase
DMSO	Dimethylsulphoxide
DSBH	Double-stranded β-helix
E. coli	*Escherichia coli*
EDTA	Ethylenediaminetetraacetic acid
EFE	Ethylene-forming enzyme
FIH	Factor inhibiting HIF
HIF	Hypoxia-inducible factor
HIS1	HPPD-inhibitor sensitive 1 (*Oryza sativa* 2OG oxygenase)
HPLC	High performance liquid chromatography
HPPD	4-Hydroxyphenylpyruvate dioxygenase
IC_{50}	Half maximum inhibitory concentration
IPTG	Isopropyl β-*D*-1-thiogalactopyranoside
JmjC	Jumonji C domain protein
Kan	Kanamycin
KDM	Lysine demethylase
KMBA	2-Keto-4-(methylthio)butanoic acid
L-asc	*L*-Ascorbic acid
LC-MS	Liquid chromatography–mass spectrometry
MSA	Methanesulphonic acid
MS	Mass spectrometry
MW	Molecular weight
NMR	Nuclear magnetic resonance
NOG	*N*-Oxalylglycine
OD_{600}	Optical density at 600 nm
OsHSL	*Oryza sativa* HIS1-like 1
PDB	Protein Data Bank
PEG	Polyethylene glycol
ppm	Parts per million
PsEFE	*Pseudomonas savastanoi* ethylene-forming enzyme
PCR	Polymerase chain reaction

Abbreviation	Full form
RMSD	Root-mean-square deviation
RNA	Ribonucleic acid
SDS	Sodium dodecyl sulphate
SDS-PAGE	Sodium dodecyl sulphate–polyacrylamide gel electrophoresis
SEC	Size exclusion chromatography
SUMO	Small ubiquitin-like modifier
Tris	Tris(hydroxymethyl)aminomethane
UV–vis	Ultraviolet–visible
v/v	Volume per volume
wt	Wild type

Standard 3 letter or single letter abbreviations for amino acids, standard SI unit abbreviations, and chemical element symbols are used throughout this thesis.

Table of Contents

Acknowledgements ..... **iv-vi**
Declaration ..... **vii**
List of Publications ..... **viii**
Abstract ..... **ix**
List of Abbreviations ..... **x-xi**
Table of Contents ..... **xii-xiv**

CHAPTER 1

1.1 Introduction **3**
1.2 Industrial importance of ethylene **3**
1.3 Biological significance of ethylene **4**
1.4 Ethylene biosynthetic pathways: overview **11**
1.5 The nitrogenase-like ethylene-forming enzymes **17**
1.6 ACC-Oxidase (ACCO) **20**
1.7 PsEFE **29**
1.8 Biocatalytic applications of ethylene-forming enzymes **46**
1.9 Thesis overview **51**
1.10 References **53**

CHAPTER 2

2.1 Overview **64**
2.2 Abstract **65**
2.3 Introduction **66**
2.4 Results **69**
2.5 Discussion **84**
2.6 Experimental Procedures **89**
2.7 Acknowledgements **92**
2.8 Author Contributions **93**
2.9 References **93**
2.10 Supporting Information **99**
2.11 Statement of Authorship **161**

CHAPTER 3

3.1 Overview **164**
3.2 Table Contents Graphic **165**
3.3 Abstract **165**

3.4 Introduction **166**
3.5 Results **169**
3.6 Discussion **182**
3.7 Conclusion **185**
3.8 Experimental Procedures **185**
3.9 Acknowledgements **187**
3.10 Author Contributions **187**
3.11 References **188**
3.12 Supporting Information **191**
3.12.1 Supporting Figures **191**
3.12.2 Supporting Tables **201**
3.12.3 Supporting Information References **204**
3.13 Statement of Authorship **205**

CHAPTER 4

4.1 Overview **208**
4.2 Table of Contents Graphic **209**
4.3 Abstract **209**
4.4 Introduction **210**
4.5 Results **214**
4.6 Discussion **222**
4.7 Conclusions **224**
4.8 Experimental Section **225**
4.9 Acknowledgements **226**
4.10 Author Contributions **226**
4.11 References **227**
4.12 Supporting Information **230**
4.13 Statement of Authorship **249**

CHAPTER 5

5.1 Overview **252**
5.2 Abstract **253**
5.3 Introduction **253**
5.4 Results **257**
5.5 Discussion **272**
5.6 Methods **280**

5.7 Acknowledgements **281**
5.8 Author Contributions **282**
5.9 References **282**
5.10 Supporting Figures **288**
5.11 Supporting Tables **372**
5.12 Supporting Information References **374**
5.13 Statement of Authorship **375**

CHAPTER 6

6.1 Overview **378**
6.2 Abstract **379**
6.3 Introduction **379**
6.4 Results **382**
6.5 Discussion **396**
6.6 Experimental Procedures **400**
6.7 Acknowledgements **403**
6.8 Author Contributions **403**
6.9 References **404**
6.10 Supporting Figures **408**
6.11 Supporting Tables **430**
6.12 Supporting Information References **433**
6.13 Statement of Authorship **435**

CHAPTER 7

7.1 Catalytic versatility of 2OG oxygenases **438**
7.2 PsEFE: bifurcating catalysis, substrate scope, and engineering new reactivity **440**
7.3 KMBA and KEBA as ethylene precursors **442**
7.4 HIS1 and OsHSL1: catalytic promiscuity - a herbicide resistance challenge **447**
7.5 Concluding Remarks **449**
7.6 References **449**



CHAPTER 1

INTRODUCTION



Contents

1.1 Introduction ... 3
1.2 Industrial importance of ethylene ... 3
1.2.1 Efforts towards sustainable ethylene production ... 4
1.3 Biological significance of ethylene ... 4
1.3.1 Ethylene signalling in plants ... 5
1.3.1.1 Ethylene in plant stress responses ... 9
1.3.2 Ethylene signalling in microbes ... 9
1.3.3 Applications of ethylene in agroindustry ... 10
1.4 Ethylene biosynthetic pathways: overview ... 11
1.4.1 Discovery of the ACC pathway ... 13
1.4.2 Yang cycle ... 14
1.4.3 Discovery of the 2OG/ethylene-forming enzyme pathway ... 16
1.5 The nitrogenase-like ethylene-forming enzymes ... 17
1.5.1 Mo- and V-dependent nitrogenases: CO reduction to ethylene ... 18
1.5.2 Nitrogenase-like reductases: sulphur salvage-linked ethylene formation ... 19
1.6 ACC-Oxidase (ACCO) ... 20
1.6.1 The 2-oxoglutarate (2OG) oxygenase superfamily ... 21
1.6.2 ACCO catalysis ... 23
1.6.3 Regulation of the ACC pathway ... 27
1.7 PsEFE ... 29
1.7.1 Pathogenic context of PsEFE ... 31
1.7.2 Active site of PsEFE ... 33
1.7.3 Substrate scope of PsEFE ... 35
1.7.4 Mechanistic insights and oxygen reactivity of PsEFE ... 37
1.7.5 Branch points in PsEFE catalysis ... 38
1.7.6 Mutagenesis studies on PsEFE ... 41
1.7.7 Sequence and structure relationships of EFEs and ACCOs ... 44
1.8 Biocatalytic applications of ethylene-forming enzymes ... 46
1.8.1 Production of ethylene and other gaseous alkenes ... 46
1.8.2 PsEFE as a tool for abiological transformations ... 49
1.9 Thesis overview ... 51
1.10 References ... 53

1.1 Introduction

Small gaseous molecules, such as dioxygen (O_2) and ethylene (C_2H_4), are of enormous importance in biology and industry.[1-3]A biological role for ethylene (C_2H_4, 28 Da) was first identified following observations that 'illuminating gas', a term given to gas containing ethylene in the 19th century owing to its usage in lighting as a component of coal gas, promoted plant growth.[4] Ethylene was subsequently shown to be the active component responsible for this effect, triggering investigations into its signalling function.[4,5] Ethylene is also one of the most important commodities in the chemical industry, being the highest-volume organic chemical produced globally,[4] with current industrial consumption at approximately 200 million tonnes per annum and an estimated annual production projected to exceed 230 million tonnes by 2030.[6,7]

1.2 Industrial importance of ethylene

Industrially, ethylene is produced by steam-cracking, a process in which hydrocarbons, including naphtha, are subjected to high temperatures (~800–900°C) to break chemical bonds and generate ethylene.[8] The process of cracking requires substantial amounts of energy which is currently, predominantly, obtained by combustion of fossil fuels, leading to high greenhouse gas emissions. This accounts for approximately 1.5–2.0 tonnes of CO_2 for every ton of ethylene produced; given the scale of ethylene production, cracking is responsible for a substantial amount of global energy consumption and greenhouse gas emissions.[7,8] Cracking is also environmentally detrimental, because it generates volatile nitrogen oxides (NO_x) in addition to CO_2, compounding its contribution to air pollution and climate change.[8]

Ethylene is a building block for a diverse range of products, most notably polyethylene, which is the most widely produced plastic worldwide.[9] Polyethylene is extensively used in packaging,

construction materials, household goods, and medical applications. Ethylene also serves as an indirect precursor for other synthetic polymers such as polyvinyl chloride (PVC), polystyrene, and polyester fibres.[9] Ethylene serves as a key feedstock for the synthesis of multiple industrially important chemicals, including ethylene oxide, which is used in the synthesis of ethylene glycol for antifreeze and polyester production,[10] and ethylbenzene, which is a precursor to styrene,[11] the monomer used in the production of polystyrene plastics. Ethylene is also employed in the industrial production of ethanol, acetaldehyde, and acetic acid, which serve as crucial intermediates in solvent and fuel production.[12]

1.2.1 Efforts towards sustainable ethylene production

To alleviate the significant burden that industrial ethylene production places on the environment, alternative ethylene sources are being investigated. Catalytic cracking of biomass has been employed to produce ethylene from biomass-derived feedstocks, such as ethanol and waste-oil, by dehydration and catalytic pyrolysis.[7,12] Recent advances in electrocatalysis have demonstrated the potential to convert CO_2 into ethylene using renewable electricity.[13,14] This process involves the electrochemical production of syngas, an energy-rich mixture of hydrogen and carbon monoxide, from CO_2, followed by conversion of syngas to methanol and subsequently to ethylene (and other high value chemicals).[13,14] Leveraging biological pathways, such as those catalysed by ethylene-forming enzymes found in nature, also presents a promising avenue for low-energy and relatively green production of ethylene.[15]

1.3 Biological significance of ethylene

Historically, ethylene was used as an anaesthetic and ‘illuminating gas’ in the form of gas mixtures. Biologically, ethylene is a widely studied phytohormone, regulating processes in higher plants, including senescence and ageing. It promotes the ripening of climacteric fruits

such as bananas, tomatoes, and apples. Ethylene also modulates seed dormancy and lateral root formation.[16,17] Ethylene influences the intricate hormonal balance in plants and thus regulates maturity traits (such as seed dormancy);[18,19] at least in part, ethylene functions by promoting gibberellin (GA) biosynthesis, a process which promotes seed germination, and which counteracts the effects of abscisic acid (ABA), a phytohormone which inhibits seed germination.[18,19] Ethylene is also reported to trigger ageing of leaves and flowers, eventually leading to programmed cell death.[4]

1.3.1 Ethylene signalling in plants

Ethylene signalling in seeding plants is driven by ethylene receptors localised to the endoplasmic reticulum (ER) membrane.[17,20-22] Ethylene receptor structures in plants are conserved, comprising an *N*-terminal transmembrane domain facing the ER lumen that binds ethylene and a *C*-terminal signal output domain. A widely occurring protein domain present in cyclic **G**MP (cGMP)-regulated cyclic nucleotide phosphodiesterases, **a**denylyl cyclases, and the bacterial transcription factor **F**hlA, GAF domain,[23] is centrally positioned between the *N*- and *C*-termini; however, the function of the GAF domain is poorly understood. The transmembrane *N*-terminal domain plays an important role in membrane localisation and coordinates Cu(I) ions which are proposed to coordinate ethylene and enable ethylene sensing (**Figure 1.1**).[17,24] Cu(I) ions are essential cofactors for ethylene receptors and are indispensable for ethylene signalling.[25] Although the ethylene receptors are functionally active as dimers, one Cu(I) ion binds to each dimer, indicating that one receptor dimer binds only one ethylene molecule.[17,24,25] The *C*-terminal portion of ethylene receptors has domains that are reminiscent of histidine kinases.[17,20-22]

Studies aimed at elucidating the molecular basis of ethylene signalling have used *Arabidopsis thaliana* as a model, which has five different ethylene receptor isoforms: ETRs (ethylene

receptors ETR1 and ETR2), ERSs (ethylene response sensors ERS1 and ERS2), and EIN2 (ethylene-insensitive 2).[17] ETRs and EIN2 act as primary receptors, bearing a receiver domain, while ERSs play a modulatory role in the signalling pathway.[26] Structurally, ETRs are similar to the bacterial two-component systems having two activated sites – a response regulator and a sensory histidine kinase.[17,27] Similarities between the model ethylene receptors and the bacterial two-component systems, indicate that the ethylene receptor machinery may have been acquired from a *cyanobacterium* species that gave rise to plastids, acquired alongside other signalling modules like phytochromes and cytokinin receptors.[17,28]

The two central protein components of ethylene signalling are Constitutive Triple Response I (CTR1) and Ethylene Insensitive 2 (EIN2).[16,17] The ethylene signalling pathway involves a sequence of molecular events initiated by ethylene binding to Cu(I) ions localised in ETRs and ERSs. Typically, this process involves: (i) ethylene perception by its membrane-bound receptors, (ii) the inactivation of its negative regulator, CTR1, (iii) self-cleavage of EIN2, and (iv) subsequent activation of transcription factors, EIN3 and EILs. It is to be noted that ethylene acts as a suppressor of its receptor(s), thereby making the receptor a negative regulator of ethylene responses.[16,17]

In the absence of ethylene, the ETRs and ERSs keep CTR1 active while it can interact with the kinase (sensory) domain of ETRI (**Figure 1.1**).[17,22] CTR1 is involved in the direct phosphorylation of the ER-localised protein EIN2, leading to its ubiquitination and subsequent proteolysis. As a downstream consequence, transcription factors (EIN3, EIL1 (EIN3-LIKE1), and EIL2 (EIN3-LIKE2)) are targeted for ubiquitination by F-box proteins (EBF1 and EBF2); this breakdown prevents ethylene responses. Thus, by reducing the levels of EIN2, ethylene blocks the signal transduction pathway.[17,20,21]

In the presence of ethylene, CTR1 is inhibited leading to an increase in EIN2 levels.[17,22] It is unclear whether ethylene binding to its receptors reduces the catalytic activity of CTR1 or whether it triggers conformational changes in ethylene receptors that reduce the interaction between CTR1 and EIN2 (**Figure 1.1**).[17,20,21] Whatever the mechanism, the binding of ethylene to its receptors reduces the phosphorylation (and subsequent degradation by the proteasome) of EIN2, thereby increasing its cellular levels. Consequently, EIN2 undergoes self-cleavage to release its *C*-terminal portion, EIN2-C, while EIN2-N remains membrane bound. EIN2-C provides both transcriptional and translational control. Firstly, it binds the mRNAs that encode for EBF1 and EBF2 and directs their degradation by exoribonucleases. Secondly, EIN2-C contains a nuclear localisation sequence which helps in modulating EIN3-dependent transcription.[17,20,21]




Figure 1.1: Ethylene signal transduction and the activity of Constitutive Triple Response I (CTR1) and Ethylene Insensitive 2 (EIN2) in the absence (A) and presence (B) of ethylene.[17,22] **A)** CTR1 is active in the absence of ethylene and enables the phosphorylation, and subsequent degradation, of EIN2 and other downstream transcription factors. Thus, in the absence of ethylene, signal transduction is blocked. **B)** Ethylene is a repressor of its receptor and ethylene's interaction with ETRs prevents EIN2 degradation by inhibiting EIN2 phosphorylation; this enables ethylene-stimulated regulation of transcription in the nucleus. The figure was created using BioRender.com.

There are several studies supporting the existence of a non-canonical ethylene signalling pathway wherein ethylene receptors and downstream transcription factors are involved in other signalling processes.[17,22] It is proposed that ETR1 catalyses a phosphorelay to components of cytokinin signalling pathway.[17,29] However, the role of different transcription factors, EIN3, EIL1, and EIL2, has not yet been assigned to ethylene-related traits.[17]

1.3.1.1 Ethylene in plant stress responses

Ethylene production in plants is dynamically regulated in response to biotic and abiotic environmental stresses.[30-33] Abiotic stresses that promote ethylene production in plants include flooding and light exposure.[30-33] Ethylene signalling promotes the opening of stomatal pores, aerenchyma (air spaces) formation, and adventitious root formation to improve oxygen transport during waterlogging to cope with hypoxic conditions.[30-33] During drought, ethylene slows down plant transpiration to conserve water. Ethylene production is also upregulated in response to the recognition of microbe-associated molecular patterns (MAMPs), activating immune responses.[34] In response to physical trauma, ethylene production is also upregulated.[30-33]

1.3.2 Ethylene signalling in microbes

Ethylene biosynthesis is not exclusive to higher plants—microbes, including plant-infecting bacteria and fungi, also produce ethylene.[35-37] Plant-like ethylene receptors with a histidine-kinase-like domain similar to ETRs have been identified in proteobacteria and cyanobacteria.[17,27,35-39] ETR homologues, similar to those in higher plants, have also been identified in eukaryotes, such as fungal plant symbionts. The presence of Cu(I) in these receptor homologues has not been biochemically validated; however, the conservation of the ethylene-binding domain and the presence of functional ethylene receptors in bacteria suggests that microbial ethylene receptors may similarly utilise Cu(I) as a cofactor for ethylene sensing.[38,39]

It has been proposed that plant pathovars might have evolved ethylene-producing capabilities to facilitate invasion of plant tissues by hijacking the plant's innate ethylene-response mechanisms.[22,40,41] This hypothesis is supported by the observation that, in certain ecosystems, microbially produced ethylene has deleterious effects on plants.[34,36] Ethylene is also known to

influence biofilm formation in bacteria.[39] The earliest studies on microbial ethylene production date back to the 1980s in Japan, where cell-free systems were first shown to produce ethylene.[42] These microbial processes are of increasing interest due to their potential applications in biocontrol, plant-pathogen interactions, and biocatalytic ethylene synthesis**.**

1.3.3 Applications of ethylene in agroindustry

Regulating ethylene responses is common practice in horticulture and agriculture, because it affects traits influencing plant vigour, post-harvest physiology, and storage.[4,5,16,17,28] While amplifying the effects of ethylene can force ripening of fruits, induce flowering, and initiate reproductive development in plants, reducing its effects can effectively prolong vase life of flowers, shelf-life of fruits, and delay ripening.[4] The agricultural industry exploits ethylene regulation for crop management in multiple ways, including by:

- **Amplifying ethylene responses for regulating plant growth and ripeness**: Plant growth regulators, like Ethephon, are widely employed to accelerate fruit ripening.[43] Ethephon is metabolised to ethylene in the plant and helps induce flowering, hasten ripening, and reduce seed formation.[43,44]
- **Amplifying ethylene responses for weed management**: Ethylene responses, such as increased senescence, have been harnessed to control parasitic weeds such as *Striga* which cause annual crop-damage in sub-Saharan Africa.[45,46] *Striga* utilises ethylene as a cue to germinate in the presence of other crops in a field. 'Suicidal germination', a strategy to stimulate seed germination in the absence of a host, has been harnessed to control weed outbreaks.[45] For example, ethylene can directly be applied to fields prone to *Striga* to induce growth prior to sowing crops.
- **Designing ethylene insensitive plants**: Heterologous expression of the *etr1-1* gene, a variant of the *etr1* gene with a C65Y mutation that is unable to bind copper and ethylene,

induced ethylene insensitivity in model organisms like *A. thaliana.*[17] However, *etr1-1* expression was linked to adverse outcomes such as increased susceptibility to stress and pathogens, and reduction in pollen viability.[17] This strategy may also enhance the biosynthesis of ethylene, potentially affecting neighbouring non-mutant plants.

- **Blocking ethylene responses**: Controlling ethylene levels is indispensable for the agriculture industry.[4] Ethylene receptor inhibitors are widely used to control fruit ripening and extend their shelf-life.[47] Metal ions, including Ag(I), competitively inhibit the ethylene transduction pathway by disrupting interactions of ethylene with the receptor-bound Cu(I) ions and are proposed to sequester ethylene, delaying its physiological responses.[17] Strained alkenes are ethylene antagonists which competitively bind to Cu(I). 1-Methylcyclopropene (1-MCP), 2,5-norbornadiene, 1-(2,2-dimethylpropyl)cyclopropene, *cis*- and *trans*-cyclooctene have shown excellent potency as ethylene antagonists;[48,49] however, only 1-MCP is commercially used for prolonging the shelf life of flowers and fruits as a cyclodextrin inclusion complex, EthylBoc.[48-50] EthylBoc is commercially marketed and widely used in floriculture and horticulture.[50]

1.4 Ethylene biosynthetic pathways: overview

The ethylene biosynthesis pathway in higher plants is relatively well-characterised: ethylene is produced from 1-aminocyclopropane-1-carboxylic acid (ACC) in a reaction catalysed by ACC oxidase (ACCO). ACCO employs Fe(II) as a cofactor and requires ACC, O_2, and ascorbate to produce ethylene (**Figure 1.2**).[15,22,51]

Certain marine algae utilise dimethylsulphoniopropionate (DMSP) to produce ethylene in a two-step process: DMSP lyase catalyses fragmentation of DMSP to give acrylate, which is subsequently converted to ethylene by acrylate decarboxylase.[52-54]

Pseudomonas spp. catalyse ethylene production from 2-oxoglutarate (2OG) rather than ACC (**Figure 1.2),** employing an Fe(II)-dependent oxygenase structurally related to ACCOs: PsEFE.[15,55,56] PsEFE-like enzymes are apparently not present in plants. Recent evidence implies that the photosynthetic alphaproteobacteria *Rhodospirillum rubrum* employs a reductive pathway to produce ethylene from 2-(methylthio)ethanol, a reaction catalysed by the nitrogenase-like enzyme MarBHDK under anaerobic conditions (**Figure 1.2)**.[57-60]




Figure 1.2: Validated/proposed pathways for biological ethylene production: (**i**) ACCO catalysis in higher plants;[15,22,51] (**ii**) EFE catalysis in *Pseudomonas spp*.,;[15,55,56] (**iii**) non-enzymatic ethylene production from KMBA, also observed in *Penicillium digitatum* cell extracts;[61-63] (**iv**) a nitrogenase-like reductase catalysing ethylene production in *Rhodospirillum rubrum*.[57-60]

Interestingly, lower plants, including mosses[64] (*Physcomitrella patens*) and ferns[65] (*Ceratopteris richardii*) contain ethylene receptors but, apparently, do not encode for

ACCOs.[64,65] It is unclear whether they contain ethylene-forming enzymes other than ACCOs.[64,65] In pioneering work, the transamination product of methionine (α-keto-γ-methylthiobutyric acid, KMBA) was shown to be a precursor for ethylene production in several bacteria and fungi (**Figure 1.2)**.[61-63] Subsequent work implied that the KMBA reaction may occur non-enzymatically; oxidation of KMBA in the presence of reactive oxygen species (ROS), such as hydrogen peroxide or metal catalysts, fragmented KMBA to ethylene, methanethiol, and CO_2.[42] Thus, KMBA formation has been proposed as a method of salvaging the amino nitrogen from methionine, generating KMBA-derived ethylene.[66]

Enzyme-catalysed oxidation of KMBA (**Figure 1.2** and **Figure 1.3**) has also been proposed.[62,63,67] An uncharacterised NADH:Fe(III):EDTA-dependent oxidoreductase from *Cryptococcus albidus* is proposed to generate ROS under nitrogen-limiting conditions, thereby, indirectly enhancing ethylene formation from KMBA.[62] The fungus *Botrytis cinerea* is also known to produce ethylene from KMBA by photo-oxidation during invasion.[61] Cellular extracts of green mould, *Penicillium digitatum*, supplemented with KMBA and methionine were also found to manifest high ethylene levels.[63,67] However, no enzyme capable of directly converting KMBA to ethylene has yet been characterised.[15] The KMBA route for ethylene production has been largely overlooked in biochemical literature, but not in the biological literature.

Notably, ethylene formation, as a consequence of lipid peroxidation, has also been observed in human cells during systemic inflammation.[68]

1.4.1 Discovery of the ACC pathway

Pioneering studies implied that levels of ethylene production in fruit and flower tissues varied with *L*-methionine availability.[69] A role for methionine in plant ethylene biosynthesis was shown by studies in which [^{14}C]-labelled *L*-methionine produced [^{14}C]-labelled ethylene in a

process requiring O_2. In the absence of oxygen, no ethylene formation was observed but the formation of an 'amino acid-like intermediate', with an isoelectric point of 6.8, was detected. This 'intermediate' reacted with Sanger's reagent (2,4-dinitrofluorobenzene) to generate α-aminobutyrate upon hydrogenation.[69] Systematic analysis of all potential forms of desaturated α-aminobutyrate demonstrated the identity of this intermediate to be 1-aminocyclopropane-1-carboxylic acid (ACC).[69] The enzymes involved in plant ethylene biosynthesis, starting from *L*-methionine, are now well defined.[15,22,51]

1.4.2 Yang cycle

Ethylene biosynthesis in higher plants is achieved through the Yang cycle (**Figure 1.3**),[70-73] in which *L*-methionine is converted into *S*-adenosyl-*L*-methionine (SAM), as catalysed by SAM synthase. SAM is then converted into ACC by a pyridoxal phosphate (PLP)-dependent enzyme, ACC synthase (ACS), releasing methylthioadenosine as a by-product to be recycled back to methionine in the Yang cycle.[74] ACC synthesis is the rate-limiting step in the biosynthesis of ethylene in plants, as the cellular levels of ACS are tightly regulated.[70] ACCO, then, finally catalyses conversion of ACC into ethylene.[15,22,51,70-73] It has also been shown that ACC can act as a signalling molecule independently of ethylene, with roles in pathogen interactions as well as pollen tube attraction.[71] It is striking that the ACC route to ethylene, the reductive route to ethylene from *R. rubrum*, and the potential KMBA route to ethylene, all involve methionine as an indirect ethylene precursor (**Figure 1.3**), suggesting an evolutionary relationship between methionine and ethylene.




Figure 1.3: Methionine (red) is a block for the biosynthesis of ethylene precursors (blue), including via the Yang cycle.[70-73] The Yang cycle involves the production of *S*-adenosyl-methionine (SAM) by SAM-synthase which is then converted to 5'-*S*-methyl-5'-thioadenosine (MTA) and 1-aminocyclopropane-1-carboxylic acid (ACC) by ACC synthase (ACS). MTA is then deadenanted to 5-methylthioribose (MTR) by a nucleosidase (MTN). MTR kinase (MTK) then phosphorylates MTR to form 5-methylthioribose-1-phosphate (MTR-P) which is subsequently isomerised by MTR-P isomerase (MTI) to 5-methylthioribulose-1-phosphate (MTRu-P). MTRu-P is converted to 1,2-dihidroxy-3-keto-5-methylthiopentene (DHKMP) by dehydratase-enolase-phosphatase (DEP). DHKMP is converted to α-keto-γ-methylthiobutyrate (KMBA) by acireductone dioxygenase (ARD). Ultimately, KMBA is transaminated back to methionine by an aminotransferase. It is to be noted that the converse reaction (methionine to KMBA) is also feasible in living systems enabled by oxidases.

1.4.3 Discovery of the 2OG/ethylene-forming enzyme pathway

Penicillium digitatum, referred to as green mould[75], has been reported to accelerate plant senescence, indicating that it produces a plant growth hormone. Cultures of *P. digitatum* were also found to induce epinasty in orange and grapefruit plants.[76,77] Epinasty has been characterised as a key physiological effect of ethylene, auxin, and abscisic acid.[78-81] Epinasty involves the increased growth of the upper surface of an organ relative to its lower side and causes it to bend downwards.[78-81] Chemical characterisation of the green mould vapours confirmed the identity of the growth hormone being produced by *P. digitatum* as ethylene (**Figure 1.4**). It was thus hypothesised that *P. digitatum* produced ethylene, however its mechanism was not defined.[78]




Figure 1.4. Chemical identification of ethylene gas produced by *Penicillium digitatum*.[80] **a**) Ethylene (red) forms a complex with aqueous Hg(II) as the 'green mould' vapours are passed through a solution of mercuric perchlorate. **b**) Ethylene ligands are displaced by chloride ions and released quantitatively into a reaction flask. **c**) The released ethylene decolourises bromine (brown) forming 1,2-dibromoethane (colourless). **d**) 1,2,-dibromoethane reacts with aniline

forming *N,N'*-diphenyl ethylenediamine (DPEDA). **e**) DPEDA further reacts with phenyl isothiocyanate producing *N,N'*-diphenyl-*N,N'*-dianilinothioformyl-ethylenediamine. The identity of products formed in reactions **d**) and **e**) was confirmed by melting point analysis of ethylene-derived authentic samples of the corresponding derivatives.

It was then proposed by Jacobsen *et al.* that the origin of ethylene production by *P. digitatum* is linked to the Krebs cycle.[80,82] To investigate this hypothesis, *P. digitatum* cultures were grown in the presence of [^{14}C]-labelled Krebs cycle intermediates (fumarate, succinate, malate, acetate); [^{14}C] was found to be incorporated into the ethylene skeleton from the 'middle carbon atoms' of dicarboxylic acids.[82] Ethylene labelling was also observed in the same study when [^{14}C]-labelled methionine, acrylic acid, and β-alanine were supplemented to *P. digitatum* growth media, possibly indirectly via the Krebs cycle acids.[80,82]

The characterisation of an ethylene-forming enzyme (EFE) from *Pseudomonas syringae pv. phaseolicola* PK2 (PsEFE) which produces ethylene from 2-oxoglutarate (2OG), a Krebs cycle intermediate, in the presence of *L*-arginine, confirmed the hypothesis that ethylene production may be linked to the Krebs cycle.[83] Notably, a 2OG-dependent EFE from *Penicillium digitatum* (PdEFE), sharing ~50% sequence identity with PsEFE has been recently isolated and characterised.[84] Biochemical and structural analyses to date reveal no striking differences between PsEFE and PdEFE.[84]

1.5 The nitrogenase-like ethylene-forming enzymes

Beyond 2OG oxygenases, metalloenzymes of the nitrogenase superfamily have been implicated in ethylene biosynthesis.[57,60,85-90] While nitrogenases typically catalyse reduction of N_2 to ammonia, they have been shown to catalyse the formation of ethylene and other small molecules like propylene, ethane and propane, under anaerobic conditions by the reduction of CO (**Figure 1.5**).[57,58,60,85-90] Nitrogenases have also been shown to catalyse reduction of

acetylene (C_2H_2) and H^+ to ethylene and molecular hydrogen, respectively (**Figure 1.5**). Nitrogenase-like reductases, evolved from nitrogenases, have been implicated in photosynthetic alphaproteobacteria *Rhodospirillum rubrum* and *Rhodopseudomonas palustris*, apparently repurposed for reductive chemistries beyond nitrogen fixation, particularly in sulphur metabolism.[57,58,60,85]




Figure 1.5. Reactions catalysed by nitrogenases and nitrogenase-like reductases.[58,86] Stoichiometric reactions for the reduction of: **a**) molecular nitrogen (N_2), to ammonia, and acetylene to ethylene, by canonical nitrogenases[86] (FeMoco, FeVco, FeFeco), **b**) CO, to hydrocarbons, including ethylene, by canonical nitrogenases (FeMoco, FeVco, FeFeco),[90] and **c**) volatile organic sulphur compounds, 2-(methylthio)ethanol, 2-(methylthio)ethane, and dimethyl sulphide, by nitrogenase-like reductases (MarBHDK) under anaerobic conditions.[57,58,60]

1.5.1 Mo- and V-dependent nitrogenases: CO reduction to ethylene

The nitrogenase-family of ethylene-forming enzymes comprises two homologous members, *i.e.*, the Mo- and V-dependent nitrogenases, distinguished primarily by the metal ion composition of their cofactors (FeMoco or FeVco, respectively).[86-90] Iron-only nitrogenases

(FeFeco) are also present in some species.[86,87] Nitrogenases are composed of a specific reductase (*nif*H- or *vnf*H-encoded Fe protein) and a catalytic component (*nif*DK-encoded MoFe protein or *vnf*DGK-encoded VFe protein); the two components form a functional complex during catalysis.[86-90] ATP-dependent electron transfer from the reductase to the cofactor site of the catalytic component facilitates substrate reduction.

A V-dependent nitrogenase from *Azotobacter vinelandii* reduces CO to various hydrocarbons, including ethylene (**Figure 1.5b**).[90] Following the same dual-component machinery used for N_2 reduction, the nitrogenase diverts electrons towards C-C bond formation in the presence of CO. GC-MS and isotope-labelling experiments have validated CO as the direct carbon source for ethylene. Notably, among all hydrocarbons produced, ethylene is the dominant product for both V- and Mo-nitrogenases.[89,90] However, the hydrocarbon-forming ability and product partition of Mo-nitrogenases was found to be significantly lower than for their V counterparts.[89,90] The analogous mechanism of CO reduction suggests that early nitrogenases may have had broader substrate preferences and likely a dual role in carbon and nitrogen fixation.[86-90]

1.5.2 Nitrogenase-like reductases: sulphur salvage-linked ethylene formation

The anaerobic salvage of sulphur from volatile organic sulphur compounds has recently been characterised as another ethylene-producing pathway, catalysed by nitrogenase-like enzymes (**Figure 1.5c**).[57,58,60,85] Under sulphur-limiting conditions, such as in soils with low sulphate availability, *R. rubrum* and *R. palustris* express the MarBHDK gene cluster which encodes for a methyl-thio alkane reductase.[58,60] This reductase catalyses the conversion of (2-methylthio)ethanol to methanethiol and ethylene.[58,60] The products, methanethiol and gaseous hydrocarbons, reflect a coupling of sulphur assimilation with hydrocarbon formation. Unlike canonical Mo- and V-dependent nitrogenases, this reductive pathway is independently

regulated by sulphur availability, via the transcription factor SalR.[58,60] Despite being structurally-related to canonical nitrogenases, with an ATP-binding site and [4Fe-4S] cluster motifs, nitrogenase-like reductases do not exhibit nitrogen-reduction capabilities.[85] Interestingly, methylthio-alkane reductases retain the ability to reduce the alternative nitrogenase substrate, acetylene, to ethylene (**Figure 1.5**).[57,58,60,85] A P-cluster and an [8Fe-9S-C]-cluster has also been identified in the reductases belonging to the MarBHDK gene cluster.[85]

The nitrogenase-like reductases form a distinct phylogenetic clade within the nitrogenase superfamily.[58,60] MarBHDK homologues have been identified in diverse soil and aquatic bacteria, suggesting that the reductive anaerobic route to ethylene production may be widespread.[57,58,60] The characterisation of this pathway also explains the accumulation of ethylene in oxygen-depleted soils, an observation that had previously lacked a clear biochemical basis.[58,60]

1.6 ACC-Oxidase (ACCO)

ACCO, a member of the 2OG oxygenase structural superfamily, catalyses the final step in the plant ethylene biosynthetic pathway.[15,22,51] Unlike most members of the 2OG oxygenase superfamily, ACCO does not employ 2OG as a co-substrate; instead, ACCO catalytically converts ACC into ethylene in the presence of oxygen and ascorbate, releasing carbon dioxide and hydrogen cyanide (HCN) as by-products (**Figure 1.2**).[15,22,51] β-Cyanoalanine synthase detoxifies cyanide by converting it into β-cyanoalanine,[91-93] which is subsequently hydrolysed to asparagine or aspartic acid.[93]

1.6.1 The 2-oxoglutarate (2OG) oxygenase superfamily

2-Oxoglutarate (2OG)-dependent oxygenases form one of the most expansive and functionally diverse oxygenase families. They catalyse reactions including hydroxylation, dealkylation, halogenation, desaturation, ring formation, and ring expansion processes.[94] 2OG oxygenases often utilise Fe(II) as a cofactor and 2OG as a co-substrate, coupling oxidative decarboxylation of 2OG, to form succinate and carbon dioxide, with substrate oxidation.[94] 2OG functions as a source of electrons that reduce and activate dioxygen in order to facilitate the oxidation of a primary substrate.

In plants, enzymes belonging to the 2OG-oxygenase structural superfamily catalyse processes including hormone biosynthesis (*e.g.*, GA20-oxidases produce gibberellins,[94,95] ACCO produces ethylene[15,22,51]). While in humans, 2OG oxygenases are involved in modulating responses to hypoxia (*e.g.*, prolyl hydroxylases[96]), epigenetic regulation (*e.g.*, Jumonji-C domain-containing histone demethylases[97,98]), collagen biosynthesis (*e.g.*, prolyl hydroxylases[96]), and transcriptional regulation (*e.g.*, Factor Inhibiting HIF, FIH[99]). Members of the 2OG-oxygenase superfamily are also widely distributed in microbes,[94] with diverse functions including in penicillin and cephalosporin production (*e.g.*, isopenicillin *N* synthase, IPNS).[100,101]

Despite their functional diversity, 2OG-dependent oxygenases share a conserved distorted double-stranded beta helix (DSBH) core (**Figure 1.6**).[94,102,103] The DSBH core typically comprises eight β-strands, arranged in a β-barrel which help to form the active-site. The Fe(II) in most 2OG-oxygenases is bound by a conserved 'facial triad' motif of two histidine residues and an aspartate or glutamate residue (**Figure 1.6**). In this superfamily of oxygenases, 2OG binds to a different pocket from the primary substrate, chelating Fe(II) via its α-ketoacid

moiety.[94,102,103] The C5 carboxylate of 2OG is typically bound to a polar lysine- or arginine-residue at the catalytic site and commonly an alcohol-containing residue (Ser or Tyr).[94,102,103]




Figure 1.6: Structural views from representative members of the 2-oxoacid-dependent oxygenase (2OADO) superfamily. A-C) Views from structures of 2OADOs from plants, humans, and microbes, in complex with their native substrates or (co)substrates, displaying the conserved DSBH fold (highlighted in teal): **A**) *Arabidopsis thaliana* Jasmonate-induced oxygenase (JOX2, PDB 6LSV[104]), **B**) *Homo sapiens* Jumonji-C domain-containing 7[105] **(**JMJD7, PDB 5NFO**),** and **C)** *Aspergillus nidulans* Isopenicillin N-synthase[106] (IPNS, PDB 6Y0O[106]). **D-F**) Close-up views of the conserved '2-His-1-carboxylate' facial triad in **D**) JOX2, **E**) JMJD7, and **F**) IPNS. Fe(II) and Fe(II)-substituting metal ions are shown as magenta spheres (Mn(II) for JMJD7, and Fe(III) for IPNS).

Notably, some enzymes of the 2OG-oxygenase superfamily, including ACCO, and isopenicillin *N* synthase (IPNS), which possess the characteristic DSBH fold, do not employ 2-oxoglutarate (2OG) as a cosubstrate.[15,22,51,106,107] IPNS catalyses the formation of isopenicillin N from *N*-[(5*S*)-5-amino-5-carboxypentanoyl]-*L*-cysteinyl-*D*-valine (ACV).[106]

Another notable example of a 2-oxoacid-dependent oxygenase that does not require 2OG as a cosubstrate is the 4-hydroxyphenylpyruvate dioxygenase (HPPD). HPPD catalyses the oxidative decarboxylation of 4-hydroxyphenylpyruvic acid (HPPA) to homogentisic acid (HGA), which is a key precursor for tocopherol (vitamin E) and plastoquinone in plants.[107,108] Note that HPPD is of agricultural importance as it is the primary target of β-triketone herbicides, widely used for weed management by disrupting carotenoid synthesis, leading to bleaching of the plant tissue and eventually plant death.[108] 2OG-dependent oxygenases conferring herbicide resistance in plants have also been identified, *e.g.*, HPPD Inhibitor Sensitive 1 (HIS1), highlighting an adaptive enzymatic role in xenobiotic metabolism.[109-112]

1.6.2 ACCO catalysis

Pioneering studies on ACCO purified from apple fruit demonstrated the need for Fe(II), *L*-ascorbate, and carbon dioxide/bicarbonate for ethylene formation.[113-116] The precise mechanism of ACCO catalysis has not been fully experimentally elucidated, but biochemical and computational studies considering alternative substrates, solvent isotope effects, and competitive oxygen kinetic isotope effects have led to a proposed mechanism (**Figure 1.7**).[15,22,51,117] Crystal structure of *Petunia hybrida* ACCO in complex with ACC (PDB: 5TCV[118]) and Q-band electron nuclear double resonance (ENDOR) studies have confirmed that the α-amino and α-carboxylate groups of ACC directly coordinate to Fe(II).[119]




Figure 1.7: Proposed mechanism for ACCO catalysis.[15,22,51,117] Binding (sequential) of ACC and molecular oxygen in the ACCO active site generates an Fe(IV)-oxo/N-radical intermediate upon reduction and protonation. Radical rearrangement yields an Fe(III)-hydroxide (dual-radical) species which eventually reacts to form ethylene. Product release restores ACCO for the next catalytic cycle.[120-122] The proposed roles of bicarbonate and *L*-ascorbate are shown in blue; bicarbonate facilitates proton transfer and ascorbate helps to meet the redox requirements of ACCO catalysis.[120-122] Note, *L*-ascorbate is required for the completion of catalytic cycles; it also promotes the reaction of O_2.[120-122]

ACCO-bound Fe(II) chelates ACC upon the displacement of two water molecules followed by the monodentate binding of an O_2 molecule, subsequently forming an Fe(III)-superoxide intermediate. An Fe(III)-hydroxyperoxide species is formed upon protonation/reduction, the formation of which is proposed to be accelerated by bicarbonate.[121] This step is followed by the formation of a transient Fe(IV)-oxo intermediate with a nitrogen radical, where the transfer of a hydrogen radical that reduces Fe(IV) back to Fe(III) initiates a decarboxylation-like

reaction, releasing ethylene and forming the labile cyanoformate species. ACCO is restored for the next catalytic cycle upon product release and metal reduction.[15,22,51,117,120-122] Intriguingly, bicarbonate or carbon dioxide have been shown to enhance ACCO's catalytic activity.[113,122] Circular dichroism studies indicated that carbon dioxide stabilises a six-coordinate state of the metal in the ACCO active site until ascorbate is supplemented.[123] In the absence of CO_2, ACCO rapidly reacts with oxygen and is deactivated, at least *in vitro*.[113-116] The presence of bicarbonate/CO_2 was also found to impede Fe(II) dissociation from ACCO.[121]

Unlike 2OG-dependent oxygenases where oxidative decarboxylation of 2OG meets the electron requirements for catalysis, ACCO uses *L*-ascorbate as a co-substrate/effector to meet its redox requirements (**Figure 1.7**), not shown in plants.[120-122] Most proposals for the ACCO mechanism have postulated that *L*-ascorbate acts as an electron donor (reductant) during catalysis. *L*-ascorbate can not only restore the oxidised iron species to the reactive Fe(II) state at the end of each catalytic cycle, while itself being oxidised to dehydroascorbate, but may also provide electrons for the initial reduction of Fe(II)-bound O_2.[120-122] Using steady-state kinetic studies, Thrower *et al.*, proposed a mechanism including *L*-ascorbate binding as a crucial step in ACCO catalysis; their findings suggest that ascorbate binding occurs after ACC and oxygen binding.[122] This sequential pattern also limits the oxidative damage to the enzyme, since O_2 binding occurs only in the presence of ACC.[122] Note that no crystal structure of ACCO in complex with ascorbate is known to date.




Figure 1.8. ACC binding to *Petunia hybrida* ACCO (PhACCO, PDB: 5TCV[118]). The overall structure of PhACCO in complex with Co(II) and ACC (DSBH core is highlighted in teal). ACCO residues in proximity to the substrate are shown in pink; ACC is in brown. The facial triad is in purple; Co(II) is in magenta.

An ACCO crystal structure in complex with Co(II) (as a catalytically inactive Fe(II) surrogate) and ACC has revealed a hydrophobic binding pocket with residues similar to those of the active site of IPNS (**Figure 1.8**).[106,124,125] The ACC cyclopropane ring binds in a pocket formed by Ser249, Asn216, Val236, and Ala248 (**Figure 1.8**). Reported mutagenesis studies on *Malus domestica* ACCO1 have shown that Thr157, Lys158, Arg175, Arg244, and Ser246 are critical for activity;[71,126] equivalent residues in *Petunia hybrida* ACCO[118] are highlighted in **Figure 1.8**. Arg244 and Ser246 are part of an RXS motif which is conserved within the ACCO structural family;[94] note that Arg244 and Ser246 are proposed to be involved in ascorbate binding.[122]

1.6.3 Regulation of the ACC pathway

Ethylene formation via the ACCO pathway is tightly regulated to ensure that it is produced in a controlled manner in response to internal developmental cues of the plant and/or in response to biotic/abiotic stimuli, which occur via different processes, as listed below.

- **Transcriptional control**: The *ACS* and *ACCO* genes are transcriptionally regulated in response to environmental signals such as flooding, drought, absence of sunlight, and mechanical damage.[31,71,127,128]. The responses to biotic and abiotic stress are governed by plant kinases, MKK9 and MPK3/6, that are involved in the phosphorylation of downstream transcription factors in the ethylene signalling pathway.[129,130]
- **Post-translational modifications**: ACS activity is regulated by phosphorylation, dephosphorylation, and ubiquitination, thereby governing ACC levels in plants.[18,19,71] These events are determined by the hormonal crosstalk of indole-3-acetic acid (IAA), ABA, GA, and ethylene to control ACC levels.[18,19,71] Cellular levels of ACC are also regulated by the amino acid transporter LHT1 (Lysine Histidine Transporter 1). While primarily involved in the uptake and transport of lysine and histidine in plants, LHT1 also mediates ACC transport and regulates its cellular availability.[131,132]
- **Formation of ACC derivatives**: ACC can undergo conjugation to form malonyl-ACC (M-ACC), γ-glutamyl-ACC (G-ACC) and jasmonyl-ACC (JA-ACC) by ACC-*N*-malonyl transferase (AMT), γ-glutamyl-transpeptidase (GGT), and jasmonic acid resistance 1 (JAR1) respectively, modulating the cellular concentration of free ACC available for ethylene biosynthesis by ACCO.[71,131,133]




Figure 1.9. The cellular concentration of free ACC for ethylene biosynthesis is enzymatically regulated.[71,131,133-138] ACC can form three conjugates: 1-malonyl-ACC (M-ACC) is formed by ACC-*N*-malonyl transferase (AMT), γ-glutamyl-ACC (G-ACC) is formed by γ-glutamyl-transpeptidase (GGT) using glutathione, and jasmonyl-ACC (JA-ACC) is formed by jasmonic acid resistance 1 (JAR1).[71,131,133] ACC can also be degraded to α-ketobutyric acid and ammonia by ACC deaminase.[134-137]

- **Feedback inhibition**: Ethylene can downregulate its biosynthesis through negative feedback loops. As mentioned previously, ethylene, contrary to most signal transduction ligands that activate their receptors to relay signals, suppresses its receptors, self-regulating its extracellular levels.[17,20-22]
- **Pathogen interactions**: Plant growth-promoting pathogenic bacteria also influence ethylene levels in plants. These bacteria encode for an ACC deaminase, which cleaves ACC, producing ammonia and α-ketobutyrate, that serve as nitrogen and carbon sources for bacterial growth, respectively.[134-137] ACC deaminase is proposed to aid the pathogen to overcome ACC-mediated immune responses of the plant.[138] Studies have shown that plant-infecting pathogens produce IAA which triggers the expression of ACS in plants and thereby elevates ACC levels; this is in accord with the observation that ethylene levels are upregulated upon IAA administration.[135,136]

1.7 PsEFE

The *Pseudomonas syringae* pv. *phaseolicola* PK2 ethylene-forming enzyme (PsEFE) is one of the most studied and well-characterised ethylene-forming enzymes.[3,15,55,56,139-142] Like ACCO, PsEFE is a member of the Fe(II)/2-oxoglutarate (2OG) oxygenase structural superfamily. Unlike other members of this enzyme family, which primarily catalyse the hydroxylation of a primary substrate coupled to the oxidative decarboxylation of 2OG to succinate, PsEFE simultaneously oxidises two different substrates—*L*-arginine and 2OG, the latter to give ethylene via an unprecedented bifurcating reaction (**Figure 1.10**).[3,15,55,56,139-142]




Figure 1.10. Schematic representation of PsEFE catalysis of 2OG.[56,139-142] PsEFE catalyses the conversion of 2OG to give (i) succinic acid and CO_2, an oxidative decarboxylation reaction coupled to the C5 hydroxylation of *L*-arginine that fragments to guanidine and *L*-Δ^1-pyrroline-5-carboxylate (P5C), and (ii) ethylene and three molecules of CO_2, a Grob-type oxidative fragmentation reaction, apparently uncoupled from the C5 hydroxylation of *L*-arginine.

PsEFE couples the oxidative decarboxylation of 2OG to succinate with the C5 hydroxylation of *L*-arginine; the resultant 5-hydroxyarginine non-enzymatically decomposes to guanidine and pyrroline-5-carboxylic acid (P5C). Concomitantly, the enzyme catalyses the four-electron oxidation of 2OG, dependent on the presence of *L*-arginine, to give ethylene.[3,15,55,56,139-142] Note that ethylene formation pathway is not coupled to *L*-arginine hydroxylation but only proceeds in the presence of *L*-arginine.[3,15]

The overall PsEFE structure comprises of 10 α-helices and 14 β-strands.[55] PsEFE has a classical DSBH core or the 'cupin' fold, characteristic of the 2OG oxygenase structural superfamily, formed by eight β-strands (**Figure 1.11**). It bears a canonical '2-His-1-carboxylate' Fe(II)-binding motif (His189, His268, and Asp191).[56,140,142] The 2OG binding pocket is unusually hydrophobic (lined by side chains of Leu173, Phe175, Ile186, Ala199, Leu206, Val270, Ala279, Ala281, and Phe283), with only a single polar residue (Arg277) located at the Fe(II)-distal end of the pocket (**Figure 1.11**).[55] The DSBH core forms the scaffold of the active site, including a loop between the two strands where the primary substrate of PsEFE, *L*-arginine binds.[55]




Figure 1.11. The hydrophobic binding pocket of PsEFE.[55] View of the active site of PsEFE in complex with Mn(II), 2OG, and *L*-arginine (PDB ID: 5LUN[55]). 2OG (green), binds in a pocket lined by hydrophobic residues (raspberry). The canonical '2-His 1-carboxylate' motif (salmon) and the 2OG C1 carboxylate and C2 carbonyl coordinate the Fe(II) ion in an octahedral arrangement. *L*-Arginine (blue) is bound through electrostatic interactions at the opposite end of the active site (orange).

1.7.1 Pathogenic context of PsEFE

Pseudomonas syringae pv. *phaseolicola* causes bean halo blight, a disease marked by interveinal chlorosis caused by the accumulation of ornithine and deficiency of arginine in the host plant tissues.[143,144] The phytotoxin responsible for driving chlorosis is phaseolotoxin, a tripeptide produced by *Pseudomonas* spp., consisting of ornithine, alanine, and homoarginine, [N^{δ}-(*N′*-sulphodiaminophosphinyl)-ornithyl-alanylhomoarginine] (**Figure 1.12)**.[143-145] Phaseolotoxin is metabolised in plants to N^{δ}-(*N′*-sulphodiaminophosphinyl)-*L*-ornithine or octicidine; octicidine inhibits ornithine transcarbamoylase (OCTase), an enzyme that converts ornithine to arginine via citrulline (**Figure 1.12**).[143-145] Inhibition of OCTase disrupts arginine biosynthesis in the host and causes a systemic collapse of the plant tissue. The pathogen itself encodes for an octicidine-resistant OCTase isoform, ArgK, to protect itself.[143-145] The presence of an *efe* gene in the same pathogen, *P. syringae*, raises the possibility of synergistic action where PsEFE may further deplete the host-arginine pools to exacerbate the effects of phaseolotoxin on the plant, amplifying chlorosis. Although this interplay has not been validated, it suggests that PsEFE may have a role in virulence beyond ethylene production.




Figure 1.12. *Pseudomonas* spp. disrupt host arginine biosynthesis.[143-145] Phaseolotoxin is metabolised to octicidine in the plant, which inhibits ornithine transcarbamoylase, OCTase, an enzyme crucial for citrulline biosynthesis from *L*-ornithine; *L*-citrulline is the precursor to arginine in biological systems.[143-145]

The *efe* gene of *P. syringae* is proposed to be carried on a conjugative plasmid rather than being chromosomally encoded.[146-148] Weingart *et al.*, showed that the *efe* gene is horizontally transferred across strains and resides on a mobile plasmid.[146] The sporadic presence of *efe* gene across phylogenetically-related strains, even within the same pathovar, supports the hypothesis of the horizontal transfer of *efe* gene. The fact that virulent *Pseudomonas* strains, such as *tomato* and *maculicola* (tomato- and Arabidopsis-infecting pathovars) do not encode an *efe* suggests that the ethylene-forming ability is not detrimental for virulence across *P. syringae*.[149] Functional studies using gene knockout have demonstrated that *efe* contributes to virulence in *pv. glycinea*, but has little effect in *pv. phaseolicola*.[146] Ethylene production may represent an auxiliary virulence factor and may be beneficial in specific host-pathogen interactions providing an edge during infection of some plant hosts, but not crucial for all pathogenic strains.[146-148] Ethylene signalling across bacterial colonies and its physiological effects remain poorly understood. Note that chromosomal integration of the *efe* gene in engineered systems,

including *Pseudomonas putida* and *Synechococcus elongatus*, has also been shown to manifest ethylene production.[150,151]

1.7.2 Active site of PsEFE

As revealed by the high-resolution crystal structures[15,55,142,152] of the PsEFE active site, *L*-arginine binds in an extended conformation stabilised by a network of hydrogen-bonding and electrostatic interactions involving Arg171, Asp191, Tyr192, and His189 (**Figure 1.11**). *L*-Arginine binding to PsEFE induces conformational rearrangements in the β4–β5 loop and α8–α10 helices, supporting an 'induced' fit for 2OG.[15,55,142,152] Arg171 reorients to stack with the guanidinium group of *L*-arginine and interact electrostatically with both Glu84 (β4–β5 loop) and the C1 carboxylate of 2OG.[152] Glu84 itself undergoes a ~90° rotation to engage in a stabilising network of hydrogen bonds; mutations at either Arg171 or Glu84 abolish ethylene production (**Table 1.1**), underscoring their essential roles. Tyr192 reorients into the active site to form a hydrogen bond with the *L*-arginine carboxylate and an offset π-π interaction with its guanidino group (**Figure 1.13**).[152] This event leads to a twisted peptide bond between Tyr192 and Asp191, an event deemed crucial for PsEFE catalysis.[152] Mutation of Tyr192 significantly reduces activity (~5% of wild type, **Table 1.1**), confirming its critical role in *L*-arginine stabilisation (**Figure 1.13**).[152] Further, the α8–α10 helical region undergoes up to 3 Å of movement towards the active site, particularly repositioning Arg316 to engage in salt-bridge formation with *L*-arginine.[152] These shifts support a substrate-induced redesign of the active site cavity.




Figure 1.13. The Tyr192 side chain reorients to stabilise *L*-arginine in a conformation crucial for PsEFE catalysis.[152] Active-site comparisons of PsEFE in complex with Mn(II) and 2OG in **A**) the absence (PDB: 5V2X[152]) and **B**) the presence (PDB: 5V2Y[152]) of *L*-arginine. *L*-arginine binding leads to bidentate metal coordination by 2OG and introduces conformational changes in the active site. Metal-chelating residues are shown in salmon; 2OG is shown in green; *L*-arginine is shown in cyan; twisted residues are shown in orange and yellow before and after arginine-binding, respectively; Mn(II) is shown as a magenta sphere.

In the absence of *L*-arginine, 2OG coordinates Mn(II) in a monodentate fashion in two different conformations, one involving its C1 carboxylate and the other its C5 carboxylate (**Figure 1.13A**).[55,84,140,142,152] In the presence of *L*-arginine, 2OG chelates the metal in a bidentate manner (through its α-ketoacid moiety) and the final ligand, water or molecular oxygen, occupies the remaining position in the octahedral coordination sphere of the metal (**Figure 1.13**). A single polar residue, Arg277, anchors the C5 carboxylate of 2OG. Despite the hydrophobicity of the 2OG binding pocket, several structured water molecules populate the active site, forming hydrogen bonds with the C5 carboxylate and connecting to a channel that may serve as a O_2 entry or CO_2 exit route.[55] 2OG, apparently, adopts a strained, non-coplanar conformation in PsEFE. Unlike typical 2OG oxygenases where the C5 carboxylate lies

coplanar with the adjacent carbon atoms (C4 and C3), PsEFE stabilises torsion angles of ~40°–70°, as observed by Zhang *et al.*[55] This geometric distortion likely orients 2OG for its Grob-type fragmentation-like reaction, enabling ethylene release. Such a strained conformation has not been reported in other 2OG oxygenases and may be a defining feature responsible for PsEFE's unique catalytic bifurcation towards ethylene formation rather than the *L*-arginine hydroxylation pathway.

1.7.3 Substrate scope of PsEFE

The substrate scope of PsEFE has been explored both with respect to its primary substrate, *L*-arginine, and its co-substrate, 2OG.[55,56,142,152,153] Ethylene formation is strictly dependent on *L*-arginine with no activity observed for *D*-arginine.[55] Biochemical studies by Martinez *et al.* and Zhang *et al.* have independently shown that PsEFE only accepts close *L*-arginine derivatives as alternative substrates, with substantially lower ethylene production rates than in the presence of *L*-arginine.[55,142] The active *L*-arginine analogues include *L*-ornithine, *L*-canavanine, agmatine, *L*-homoarginine, *L*-argininamide, and *N*-hydroxylated/methylated guanidinium group derivatives of *L*-arginine (**Figure 1.14**).[55,142]




Figure 1.14. *L*-Arginine analogues accepted as alternative PsEFE substrates, enabling ethylene formation from 2OG. The structural differences from *L*-arginine are highlighted in blue.[55,142]

Various studies have demonstrated that 2OG oxygenases can employ 2OG derivatives as alternative co-substrates.[56,154-156] For example, kinetic studies with human aspartate/asparagine-β-hydroxylase (AspH) have shown that some C3- and/or C4-substituted derivatives can sustain catalysis as co-substrates, while others function as inhibitors or enhancers.[155] If the co-substrate scope of PsEFE extends to these derivatives, gaseous alkenes heavier than ethylene could be enzymatically produced. Efforts were, thus, probed in this direction to explore the co-substrate flexibility of PsEFE, as discussed extensively in **Chapter 2.** Studies reported by Copeland *et al.*, during the course of my work, identified five alternative C4-substituted 2OG derivatives and a novel reaction pathway, that was not observed with 2OG.[153] This involved the formation of an abortive elimination (alcohol) product, a corresponding 3-hydroxypropionate (3HP) derivative, alongside the corresponding succinate derivative (diacid) (**Figure 1.15**).[56,153] However, alkene formation was detected for none of the derivatives except with 4-fluoro-2OG, which produced fluoroethylene.[153] Notably, substrate-scope studies by Martinez *et al*., have shown that hydrophobic α-ketoacids lacking the distal carboxylate moiety (pyruvic acid, 2-oxobutyric acid, 2-oxovaleric acid) are not PsEFE (co)substrates.[142]




Figure 1.15. PsEFE catalysed oxidation of C3- and C4-substituted 2OG derivatives.[56,153] PsEFE catalyses the conversion of C3/C4-substituted 2OG derivatives to give (ii) the corresponding diacid and CO_2, the formation of which is coupled to the C5 hydroxylation of arginine (as observed for 2OG), and (iii) the corresponding alcohol product alongside two molecules of CO_2.

1.7.4 Mechanistic insights and oxygen reactivity of PsEFE

The partitioning of PsEFE catalytic flux between *L*-arginine hydroxylation (RO), ethylene formation (EF), and 3HP formation complicates mechanistic studies.[55,56,142,152,153]. Despite multiple points of uncertainty, a consensus on a few key mechanistic points in EFE catalysis has now been reached, partly by using 2OG derivatives as alternative substrates.[56,142,152,153,157,158]

Calorimetric and UV-Visible (UV-Vis) spectroscopic studies have shown that the binding of *L*-arginine enhances chelation of PsEFE:Fe(II) with 2OG under anaerobic conditions. In the absence of *L*-arginine, 2OG is only bound weakly (and often monodentately, as supported by structural evidence), with a smaller enthalpy change as compared to cooperative binding of 2OG to PsEFE in the presence of *L*-arginine.[55,142,152] The bidentate coordination of 2OG to Fe(II) in the presence of *L*-arginine displaces Fe(II)-bound water molecules and primes the active site for dioxygen activation.

Molecular dynamics (MD) simulations have identified two primary dioxygen access channels leading to the PsEFE active site.[15,159] The apparently energetically favoured pathway involves an 'off-line' geometry where O_2 binds trans to His189, away from *L*-arginine,[153] atypically positioned away from the substrate.[160] The thermodynamically less feasible route, in contrast, involves an 'in-line' Fe(III)-superoxo intermediate formation directed towards *L*-arginine.[153] The proposed PsEFE reaction mechanisms thus utilise the off-line geometry as a basis and aim to identify crucial branch points for competing reaction pathways.

1.7.5 Branch points in PsEFE catalysis

Biochemical studies on the PsEFE (co)substrate scope have helped to improve mechanistic understanding of PsEFE.[56,142,152,153,157,158,160] The use of isotopically labelled substrates, alternative co-substrates, and targeted mutagenesis has shown how perturbations in substrate/cofactor binding can disproportionately affect one pathway (*i.e.*, *L*-arginine hydroxylation versus ethylene formation) over the other. For example, 4-methyl-2OG undergoes oxidative decarboxylation coupled to *L*-arginine oxidation to form 2-methylsuccinate, but it does not produce propylene.[56,153]

L-Arginine hydroxylation (**RO**), 3HP formation (**HP**), and ethylene formation (**EF**), are proposed to be initiated from the same Fe(III)-superoxo intermediate (pink, **Figure 1.16**) to which 2OG is bound in a bidentate manner, but these pathways diverge at the level of oxygen activation and substrate orientation.[158] For the RO pathway (top green, **Figure 1.16**), a succinate-bound ferryl intermediate (**Figure 1.16,** RO1) was identified using stopped-flow UV-Vis spectroscopy; the Fe(IV)-oxo intermediate (**Figure 1.16,** RO2) flips to face *L*-arginine

(**Figure 1.16,** RO3) and mediates a hydrogen-atom abstraction and hydroxyl radical rebound to hydroxylate *L*-arginine, forming succinic acid alongside (**Figure 1.16,** RO4).[153]




Figure 1.16. Mechanistic studies have identified two key branch points (green and blue) in the PsEFE-catalysed turnover of 2OG (and 2OG derivatives) producing three products: succinate (coupled to *L*-arginine hydroxylation, RO1-RO4), ethylene (EF1-EF4), and 3HP (HP1-HP2).[56,152,153,157,158]

The ethylene-producing pathway (bottom green, **Figure 1.16**) is proposed to involve oxygen insertion into the C1-C2 bond of 2OG generating a carbonate-peroxysuccinate anhydride intermediate (**Figure 1.16,** EF1), the O-O bond of which homolytically cleaves to yield a succinyl radical and a Fe(III)-carbonate species (**Figure 1.16,** EF2).[153] Decarboxylation of the succinyl radical generates a propion-3-yl radical (**Figure 1.16,** EF3) which can combine with the Fe(III)-carbonate species to give an Fe(II)-3-(carboxyoxy)propionic acid species (**Figure 1.16,** EF4(i)).[153] The 3-(carboxyoxy)propionyl species then undergoes fragmentation to release ethylene and CO_2 (**Figure 1.16,** EF4(ii)). Fragmentation of the 3-(carboxyoxy)propionyl species is fostered by the non-coplanar conformation of the 2OG backbone and the unusually hydrophobic binding pocket of PsEFE, which stabilises the hydrophobic free radical intermediates.[153,157,158] The formation of 3HP (or its derivatives) in recent studies from 2OG

(and derivatives) supports this mechanistic proposal, where the ferrous intermediate (**Figure 1.16,** HP1) utilises Fe(II)-bound water molecules for carbonate hydrolysis, enabling 3HP/3HP-derivative formation (**Figure 1.16,** HP2).[153]

Recent studies have suggested a second possible branch point in the PsEFE mechanism governing the partition between EF and 3HP (blue, **Figure 1.16):**[158] QM/MM simulations by Rifayee *et al.*, suggested that ethylene is not produced via the ferrous intermediate (**Figure 1.16,** EF4(i), discussed above), but from the Fe(III)-propion-3-yl radical (**Figure 1.16,** EF3) that precedes it, undergoing sequential cleavage of the C2-C3 and C4-C5 bonds.[158] Whether this radical fragments to ethylene or couples with the Fe(III)-carbonate to form the ferrous intermediate depends upon the stabilisation of the radical, which is dependent on the hydrophobic environment of the active site. The ferrous intermediate forms the precursor to 3HP (or its derivatives). These results also highlight the critical role of second-sphere residues in determining product partition, through the specific example of A198L mutant.[158] Mutations altering the second coordination sphere (Leu173, Ala198, and Ile186) were proposed to influence the fate of the propionyl radical.[158] Increased hydrophobic bulk in the A198L mutant stabilised the radical permitting the formation of the ferrous intermediate, consequently forming increased levels of 3HP from 2OG.[158]

Hybrid quantum mechanics/molecular mechanics (QM/MM) calculations from two independent studies support the bifurcating reaction pathway.[15] These studies suggest that the conformational differences in *L*-arginine, and not 2OG alone, dictate product outcomes. Computational studies have also explored the potential of using electric-field switches to modulate PsEFE activity to favour one pathway over the other.[15] Application of an external electric field that lowers the intrinsic field of EFE directs the reaction flux to favour ethylene formation over *L*-arginine hydroxylation.[15]

1.7.6 Mutagenesis studies on PsEFE

Mutagenesis studies on PsEFE, aimed at improving its mechanistic understanding and influencing the partition ratios of its bifurcating reactions, have been conducted.[55,139,152,161,162] Site-directed mutagenesis has been focused on residues in proximity to the 2OG-binding site and on surface residues proposed to influence PsEFE stability.[162] Variants that disrupted the hydrophobic environment of the PsEFE active site[152] (A281R, F283R, V196R) or disrupted metal coordination[139] (H189A, D191A, H268A, D191E) showed reduced P5C and ethylene production (**Table 1.1**). Hydrophobic variants of arginine residues that stabilise 2OG (Arg277 and Arg171) highlighted the critical nature of these residues for catalysis. Interestingly, the charge-conservative R277K PsEFE variant was found to produce both ethylene and 3HP from 2OG, while sustaining *L*-arginine hydroxylation to P5C.[161] Mutagenesis studies have also demonstrated the potential of L206V PsEFE variant to produce propene from 4-methyl-2OG.[157]

Note that modest loss in PsEFE activity was also observed for variants with substitutions in residues distant from the active site (Leu73, Arg184, Glu213).[152] **Table 1.1** summarises key PsEFE variants, their proposed role, and reported catalytic outcomes.

Table 1.1. Summary of PsEFE variants (from *Pseudomonas syringae pv. phaseolicola*) the proposed role of the involved residues, and observed catalytic outcomes.

Mutation	Proposed role of residue	Outcome
L22M[139]	Surface residue	Conserved activity
L73K/R[152]	Surface residue	Reduced activity
S81R/Y[152]	Surface residue	Reduced activity
E84A/Q[55,152]	*L*-arginine binding	Ablated activity
E84D[55]	*L*-arginine binding	No ethylene production
T86S/V[55]	*L*-arginine binding	Reduced EF:RO partition ratio
H116Q[162]	Structural (interior/surface)	Reduced activity and thermostability
H168Q[162]	Structural (interior/surface)	Reduced activity and thermostability
H169Q[162]	Structural (interior)	Reduced activity and thermostability
R171A[152]	*L*-arginine binding	No ethylene production
R171K[55]	*L*-arginine binding	Ablated activity
V172T[139]	Surface residue	Reduced EF:RO partition ratio
F175Y[55]	2OG binding	Reduced EF:RO partition ratio
R184A[152]	Surface residue	Reduced activity
H189A[139]	Metal-binding ligand	Ablated activity
H189Q[162]	Metal-binding ligand	Ablated activity
D191A[139]	Metal-binding ligand	Ablated activity
D191E[161]	Metal-binding ligand	Reduced EF:RO partition ratio
Y192F[55]	*L*-arginine binding	Reduced ethylene production (4% WT)
Y192R/W[152]	*L*-arginine binding	Reduced activity
V196F[152]	2OG binding (hydrophobic pocket)	Co-purified with inclusion bodies, activity not assessed
V196R[152]	2OG binding (hydrophobic pocket)	Reduced activity; reduction in EF:RO partition ratio
A198V[152]	2OG binding (hydrophobic pocket)	Reduced activity; reduction in EF:RO partition ratio
A199G[139]	Surface residue	Reduced activity
L206V[157]	2OG binding (hydrophobic pocket)	Reduced ethylene formation from 2OG; propylene formation observed (4-Me-2OG)
E213A[152]	Surface residue (‘lid’)	Reduced activity
E215A[152]	Surface residue (‘lid’)	Reduced activity
H233A[139]	Surface residue	Enhanced activity (125% wt)
H233Q[162]	Surface residue	Ablated activity
E235D[139]	Surface residue	Reduced EF:RO partition ratio
I254M[139]	Surface residue	Conserved activity
H268A[139]	Metal-binding ligand	Ablated activity

Mutation	Proposed role of residue	Outcome
H268Q[162]	Metal-binding ligand	Reduced activity and thermostability
V270T[55]	2OG binding (hydrophobic pocket)	Reduced ethylene production (4% WT)
R277A[152]	2OG binding	Co-purified with inclusion bodies, activity not assessed
R277K[161]	2OG binding	3HP formation from 2OG
F278Y[139]	Surface residue	Conserved activity
C280F[55]	Surface/possible interaction with M54	Reduced activity
A281R[152]	2OG binding (hydrophobic pocket)	Reduced activity
A281V[152]	2OG binding (hydrophobic pocket)	Reduced activity; reduction in EF:RO partition ratio
F283A/R/V/Y[152]	2OG binding (hydrophobic pocket)	Reduced EF:RO partition ratio
H284Q[162]	Structural (interior)	Reduced activity and thermostability
E285A[152]	Surface residue ('lid')	Reduced EF:RO partition ratio
E285Q[152]	Structural role (H-bond to Y306 in aromatic network)	Reduced EF:RO partition ratio
I304N[139]	Surface residue	Ablated activity
H305Q[162]	Surface residue	Reduced activity
H309Q[162]	Structural (interior)	Reduced activity and thermostability
Y306A/F[152]	Aromatic network with Y192 and F283	Reduced activity
F310R[152]	Aromatic network with Y192 and F283	Reduced activity
F310W[152]	Aromatic network with Y192 and F284	Conserved activity
R316A/K[55]	*L*-arginine binding	Reduced ethylene production (4% and 13% WT)
C317A[55]	*L*-arginine binding	Reduced activity; reduction in EF:RO partition ratio
C317N[152]	*L*-arginine binding	Reduced activity
C317S[55]	*L*-arginine binding	Reduced activity; reduction in EF:RO partition ratio
Y318F[55]	*L*-arginine binding	Reduced ethylene production (35% WT)
R326S[139]	Surface residue	Reduced activity
I322V[139]	Surface residue	Enhanced activity (110% wt)
H335Q[162]	Surface residue	Reduced activity

1.7.7 Sequence and structure relationships of EFEs and ACCOs

Despite mechanistic and structural insights into EFEs and ACCOs, the evolutionary basis of their divergent reactivities remains poorly understood. Comparative sequence analysis of four representative enzymes (**Figure 1.17**): the 2OG-dependent ethylene-forming enzymes from *Pseudomonas savastanoi* pv. *phaseolicola* PK2 (PsEFE[56,142,152,153,157]) and *Penicillium digitatum* (PdEFE[84]) and ACCOs from *Petunia x hybrida* (PhACCO[15]) and *Dictyostelium discoideum* (DdACCO[15]), reveals several important features underscoring functional conservation.

Despite catalysing the same reactions – *L*-arginine hydroxylation and ethylene formation – PdEFE and PsEFE share only ~50% sequence identity. PsEFE has ~24% identity with DdACCO (slime mould ACCO) and only ~20% identity with PhACCO, showing evolutionary divergence from bacteria to higher organisms. Interestingly, PhACCO and DdACCO share only ~25% sequence identity. Notably, the Fe(II)-binding residues are conserved across all four sequences (shown in green circles). As discussed above, EFEs and ACCOs are structurally similar, with both enzymes exhibiting a DSBH core;[15] PsEFE comprises fourteen β-strands and ten α-helices while the PhACCO main chain contains eleven α-helices and twelve β-strands, eight of which constitute the conserved distorted DSBH core.




Figure 1.17. Sequence comparison of EFE sequences from *Pseudomonas syringae* pv. *phaseolicola* PK2 and *Penicillium digitatum* (P32021 and A0A7T7BQH3 respectively) and ACCO sequences from *Petunia x hybrida* and *Dictyostelium discoideum* (Q08506 and Q76NT9 respectively), highlighting the secondary structure of PK2 EFE (PsEFE). α-helices and β-strands are sequentially numbered. Identical residues are highlighted in red and the metal-binding ligands are indicated by green circles. The figure was generated using ESPript 3.0.[163]




Figure 1.18. Structural comparison of PhACCO and PsEFE. Cartoon depiction and active site of **A**) PhACCO (PDB: 5TCV[15]) and **B**) PsEFE (PDB: 5V2Y[152]). Metal-binding ligands are shown as sticks and the Fe-substituting catalytically inactive metals, Ni (PhACCO) and Mn (PsEFE), are shown as magenta spheres. Substrates and (co)substrates, ACC (brown), 2OG (green), and *L*-arginine (cyan), are shown as sticks.

1.8 Biocatalytic applications of ethylene-forming enzymes

The potential of natural ethylene-forming enzymes, including ACCOs, nitrogenases, and EFEs, has been shown to extend beyond their role in ethylene biosynthesis, including the production of higher olefins.[15]

1.8.1 Production of ethylene and other gaseous alkenes

The V- and Mo-dependent nitrogenases produce hydrocarbons, such as, propylene, propane, and ethane, alongside ethylene, during CO reduction, highlighting their broad catalytic potential. Nitrogenases are of particular industrial interest as they link two industrially significant processes, Haber–Bosch (nitrogen fixation) and Fischer–Tropsch (CO reduction), within a single biological framework.[86-89]

ACC oxidases purified from *Solanum lycopersicum* have also been shown to catalyse 1-butene production using 1-amino-2-ethylcyclopropane (AEC) as a substrate in a manner similar to

ethylene formation.[71,126,164-168] Different stereoisomers of the same analogue were found to produce 1-butene with varying efficiency; (1*R*,2*S*)- and (1*S*, 2*R*)-AEC were ~21 times more efficiently turned over by ACCO than (1*R*, 2*R*)- and (1*S*, 2*S*)-AEC (**Figure 1.19**).[167,168]




Figure 1.19. *Solanum lycopersicum* ACCO preferentially converts mixtures of (1*R*,2*S*)- and (1*S*, 2*R*)-AEC to 1-butene over mixtures of (1*R*, 2*R*)- and (1*S*, 2*S*)-AEC.[167,168]

ACCO catalysis is associated with stoichiometric production of toxic cyanide,[15,22,51] which limits its use for biocatalytic ethylene/olefin synthesis. PsEFE neither produces cyanide as a by-product (unlike ACCO)[12] nor does it require reducing and sulphur-limiting conditions (unlike MarBHDK),[57,58,60] potentially overcoming major limitations for biotechnological applications.




Figure 1.20. L206V PsEFE can catalyse the formation of propylene from 4-methyl-2OG.[157] L206V PsEFE can accommodate the additional steric bulk of the 4-methyl substituent and produce propylene alongside the other products of PsEFE catalysis observed with wild-type PsEFE (2-methylsuccinic acid and 3-hydroxyisobutyric acid).

PsEFE has been shown to produce propylene, from non-native methyl-substituted 2OG derivative, 4-methyl-2OG (**Figure 1.20**).[157] The feasibility of this strategy is highly dependent upon the ability of the enzyme to accommodate additional substituents and stabilise bulkier intermediates (**Chapter 2** and **4**). Bulk-diminishing substitutions in the 2OG-binding pocket could reduce the steric hindrance and better accommodate the antiperiplanar conformation necessary for the fragmentation reaction. One study explored the L206V PsEFE variant which displayed improved propylene formation from 4-methyl-2OG compared to wild-type PsEFE (**Figure 1.20**)[157]. In contrast, wild-type PsEFE predominantly formed the corresponding alcohol product. This is in accord with the proposed mechanistic model in which subtle modifications around the 2OG-binding pocket influence the second branch point (**Figure 1.16**), *i.e.*, the stabilisation of the 2-(carboxyethyl)carbonato-Fe(II) intermediate and its subsequent fragmentation to produce olefins.[158] L206V PsEFE achieves this reactivity enhancement without diverting flux towards the hydroxylation pathway.[157] These findings suggest that PsEFE may be engineered for biocatalytic production of alkenes beyond ethylene.

1.8.2 PsEFE as a tool for abiological transformations

Reported substrate-scope and mutagenesis studies on PsEFE have highlighted the potential for tuning its bifurcating reaction.[55,56,153] This flexibility can be employed for the production of substituted succinic acid and β-hydroxypropionate derivatives, including some widely used in pharmaceuticals (γ-hydroxybutyric acid, β-hydroxybutyric acid) and cosmetics (malic acid).[56,153]

PsEFE has also shown promise as a biocatalyst for C-N bond-forming reactions, including olefin aziridination and C-H bond insertion (**Figure 1.21**).[169-171] Goldberg *et al*. demonstrated that PsEFE can be used as a non-haem catalyst capable of catalysing nitrene-transfer reactions – a reaction historically restricted to engineered haem proteins.[170] PsEFE was shown to catalyse olefin aziridination and C-H bond insertion using *para*-toluenesulphonyl azide as a nitrene precursor.[170] The study also reported enhanced activity for engineered PsEFE variants obtained by directed evolution approaches like site-saturation mutagenesis.[170]




Figure 1.21. PsEFE is capable of catalysing nitrene transfer upon ligand-mediated activation of its non-haem iron centre.[169-171]

The aziridination reaction, unlike the canonical Fe(IV)-oxo intermediate,[15] proceeds via a putative Fe-nitrene intermediate.[170] The aziridination ability of PsEFE was significantly enhanced when the likely native co-substrate 2OG was replaced by small-molecule ligands such as acetate or *N*-oxalylglycine (NOG).[170] These ligands might influence catalysis by modulating the metal-coordination environment for enabling non-native reactivity modes. Directed evolution of PsEFE also yielded a highly optimised variant with enhanced aziridination activity.[170] The five-residue mutant (MLHMM) also showed excellent enantioselectivity and ligand selectivity towards acetate, with no activity observed with 2OG,[170] highlighting the tunability of the metal-ligand interface of PsEFE. Other PsEFE variants demonstrated high chemo- and enantio-selectivity in C-H insertions. Note that the aziridination activity of PsEFE was retained under aerobic conditions, a notable advantage over haem-based nitrene transferases.[170]

These examples combined with the ease of expression of PsEFE in simple *E. coli* hosts and its observable catalytic activity in whole cells and lysates, underscore its biotechnological potential as an engineerable biocatalyst. However, despite its versatility, the use of PsEFE as a biocatalyst remains challenging because of its associated wasteful flux towards competing reaction pathways which poses challenges for maximising the yield of the desired product.

1.9 Thesis overview

This chapter **(Chapter 1)** gives a background to the research described in this thesis and summarises the reported structural and biochemical analyses on characterised ethylene-forming enzymes (PsEFE, PdEFE, and ACCO), involving X-ray crystallography, mutagenesis, isotope-labelling experiments, and substrate scope studies.

Chapter 2 (Manuscript I[56]) describes ^{1}H NMR studies on the *Pseudomonas savastanoi* pv. *phaseolicola* PK2 ethylene-forming enzyme, PsEFE, substrate scope with >30 C3- and C4-substituted 2OG derivatives. The results show how the 2OG substitution pattern can affect catalytic efficiency and the outcome of the PsEFE reaction. These studies demonstrate that the outcome of PsEFE's bifurcating reaction can be modulated through substrate structure for biocatalytic applications.

Chapter 3 (Manuscript II, under review in *ChemBioChem***)** reports biochemical and crystallographic studies on *Pseudomonas savastanoi* EFEs from two additional strains, glycinea and 1449B, validating their ethylene-forming capacity. This chapter discusses an unusual inverted binding-mode of 2OG in complex with 1449B PsEFE and Mn(II).

Chapter 4 (Manuscript III, under review in *ChemCatChem*) extends the mutagenesis studies on PsEFE to variants capable of producing alkenes other than ethylene, including propene and

1-butene. This chapter describes bump-and-hole studies on Leu206 PsEFE variants designed to accommodate the steric bulk of C4-substituted 2OG derivatives for alkene formation.

Chapter 5 (Manuscript IV, in preparation) describes the reactivity of PsEFE with respect to hydrophobic 2-oxoacids. It describes structure-guided site-directed mutagenesis studies aimed at tuning the PsEFE co-substrate scope to enable alkene formation from 2-oxoacids, in particular KMBA. This chapter provides the first description of enzyme-catalysed conversion of KMBA to ethylene. It also describes the identification of a novel KMBA-derived, naturally occurring small-molecule ethylene precursor, i.e., α-keto-γ-ethylthiobutyric acid (KEBA).

Chapter 6 (Manuscript V, in preparation) describes structural and biochemical studies on 2OG- and Fe(II)-dependent oxygenases conferring herbicide resistance in rice, HIS1 and OsHSL1. HIS1 and OsHSL1 hydroxylate and subsequently degrade widely used herbicides such as tembotrione and mesotrione. The identification of enzymes conferring herbicide-resistance has raised concerns with respect to food security and, therefore, this work aims to guide the development of potent HIS1 inhibitors and/or herbicides resistant to HIS1-mediated degradation.

Chapter 7 summarises the results and outlines future research directions.

1.10 References

1 Aresta, M. & Dibenedetto, A. in *Developments and Innovation in Carbon Dioxide (CO2) Capture and Storage Technology* Vol. 2 (ed M. Mercedes Maroto-Valer) 377–410 (Woodhead Publishing, 2010).

2 Hancock, J. T. Oxygen Is Instrumental for Biological Signaling: An Overview. *Oxygen* **1**, 3–15 (2021).

3 Walewska, A., Szewczyk, A. & Koprowski, P. Gas Signaling Molecules and Mitochondrial Potassium Channels. *Int. J. Mol. Sci.* **19**, 3227–3247 (2018). https://doi.org/10.3390/ijms19103227

4 Schaller, G. E. & Kieber, J. J. Ethylene. *Arabidopsis Book* **1**, e0071 (2002). https://doi.org/10.1199/tab.0071

5 Dillard, M. M. Ethylene - The New General Anesthetic. *J. Natl. Med. Assoc.* **22**, 10 (1930).

6 Tilsted, J. P., Bauer, F., Deere Birkbeck, C., Skovgaard, J. & Rootzén, J. Ending fossil-based growth: Confronting the political economy of petrochemical plastics. *One Earth* **6**, 607–619 (2023). https://doi.org/https://doi.org/10.1016/j.oneear.2023.05.018

7 Shin, W., Lin, B., Lai, H., Ibrahim, G. & Zang, G. Decarbonization approaches for ethylene production: comparative techno-economic and life-cycle analysis. *Green Chem.* **27**, 3655–3675 (2025). https://doi.org/10.1039/D4GC04538F

8 Bajpai, P. in *Biomass to Energy Conversion Technologies* (ed Pratima Bajpai) 153–167 (Elsevier, 2020).

9 Egbuhuzor, M. O., Ogbobe, P. & Umunnakwe, C. in *Waste Material Recycling in the Circular Economy - Challenges and Developments* (ed Dimitris S. Achilias) (IntechOpen, 2022).

10 Liang, J. *et al.* Security analysis of ethylene glycol production pathways. *Chem. Eng. Res. Des.* **214**, 65–77 (2025). https://doi.org/10.1016/j.cherd.2024.12.024

11 Gupta, R., Uslu, H. & Majumder, S. Production of Styrene from Dehydrogenation of Ethylbenzene. *Chem. Eng. Technol.* **45**, 817–823 (2022). https://doi.org/https://doi.org/10.1002/ceat.202100577

12 Wu, L., Moteki, T., Gokhale, Amit A., Flaherty, David W. & Toste, F. D. Production of Fuels and Chemicals from Biomass: Condensation Reactions and Beyond. *Chem* **1**, 32–58 (2016). https://doi.org/https://doi.org/10.1016/j.chempr.2016.05.002

13 Schreiber, M. W. Industrial CO2 Electroreduction to Ethylene: Main Technical Challenges. *Curr. Opin. Electrochem.* **44**, 101438–101445 (2024). https://doi.org/https://doi.org/10.1016/j.coelec.2023.101438

14 Berkelaar, L. *et al.* Electrochemical Conversion of Carbon Dioxide to Ethylene: Plant Design, Evaluation and Prospects for the Future. *Chem. Eng. Res. Des.* **182**, 194–206 (2022). https://doi.org/https://doi.org/10.1016/j.cherd.2022.03.034

15 Hausinger, R. P. *et al.* Biological Formation of Ethylene. *RSC Chem. Biol.* **4**, 635–646 (2023). https://doi.org/10.1039/D3CB00066D

16 Ecker, J. R. The Ethylene Signal Transduction Pathway in Plants. *Science* **268**, 667–675 (1995). https://doi.org/doi:10.1126/science.7732375

17 Binder, B. M. Ethylene signaling in plants. *J. Biol. Chem.* **295**, 7710–7725 (2020). https://doi.org/10.1074/jbc.REV120.010854

18 Beaudoin, N., Serizet, C., Gosti, F. & Giraudat, J. Interactions between abscisic acid and ethylene signaling cascades. *Plant Cell* **12**, 1103–1115 (2000). https://doi.org/10.1105/tpc.12.7.1103

19 Van de Poel, B., Smet, D. & Van Der Straeten, D. Ethylene and Hormonal Cross Talk in Vegetative Growth and Development. *Plant Physiol.* **169**, 61–72 (2015). https://doi.org/10.1104/pp.15.00724

20 Schaller, G. E. Localization of the Ethylene-Receptor Signaling Complex to the Endoplasmic Reticulum: Analysis by Two-Phase Partitioning and Density-Gradient Centrifugation. *Methods Mol. Biol.* **1573**, 113–131 (2017). https://doi.org/10.1007/978-1-4939-6854-1_10

21 Park, H. L. *et al.* Ethylene-triggered Subcellular Trafficking of CTR1 enhances the Response to Ethylene Gas. *Nat. Commun.* **14**, 365 (2023). https://doi.org/10.1038/s41467-023-35975-6

22 Wang, K. L., Li, H. & Ecker, J. R. Ethylene Biosynthesis and Signaling Networks. *Plant Cell* **14 Suppl**, S131–151 (2002). https://doi.org/10.1105/tpc.001768
23 Ho, Y. S., Burden, L. M. & Hurley, J. H. Structure of the GAF domain, a Ubiquitous Signaling Motif and a New Class of Cyclic GMP Receptor. *EMBO J.* **19**, 5288–5299 (2000). https://doi.org/10.1093/emboj/19.20.5288
24 Rodríguez, F. I. *et al.* A Copper Cofactor for the Ethylene Receptor ETR1 from Arabidopsis. *Science* **283**, 996–998 (1999). https://doi.org/10.1126/science.283.5404.996
25 Rodríguez, F. I. *et al.* A Copper Cofactor for the Ethylene Receptor ETR1 from *Arabidopsis*. *Science* **283**, 996–998 (1999). https://doi.org/10.1126/science.283.5404.996
26 Müller-Dieckmann, H.-J., Grantz, A. A. & Kim, S.-H. The Structure of the Signal Receiver Domain of the *Arabidopsis thaliana* Ethylene Receptor ETR1. *Structure* **7**, 1547–1556 (1999). https://doi.org/10.1016/S0969-2126(00)88345-8
27 Gao, R. & Stock, A. M. Biological Insights from Structures of Two-component Proteins. *Annu. Rev. Microbiol.* **63**, 133–154 (2009). https://doi.org/10.1146/annurev.micro.091208.073214
28 Lacey, R. F. & Binder, B. M. Ethylene Regulates the Physiology of the *Cyanobacterium Synechocystis* sp. PCC 6803 via an Ethylene Receptor. *Plant Physiol.* **171**, 2798–2809 (2016). https://doi.org/10.1104/pp.16.00602
29 Zdarska, M. *et al.* ETR1 Integrates Response to Ethylene and Cytokinins into a Single Multistep Phosphorelay Pathway to Control Root Growth. *Mol. Plant* **12**, 1338–1352 (2019). https://doi.org/10.1016/j.molp.2019.05.012
30 Fatma, M. *et al.* Ethylene Signaling under Stressful Environments: Analyzing Collaborative Knowledge. *Plants* **11**, 2211–2240 (2022). https://doi.org/10.3390/plants11172211
31 Wang, X., Wen, H., Suprun, A. & Zhu, H. Ethylene Signaling in Regulating Plant Growth, Development, and Stress Responses. *Plants* **14**, 309 (2025).
32 Dubois, M., Van den Broeck, L. & Inzé, D. The Pivotal Role of Ethylene in Plant Growth. *Trends Plant Sci.* **23**, 311–323 (2018). https://doi.org/https://doi.org/10.1016/j.tplants.2018.01.003
33 Chen, H., Bullock, D. A., Jr., Alonso, J. M. & Stepanova, A. N. To Fight or to Grow: The Balancing Role of Ethylene in Plant Abiotic Stress Responses. *Plants* **11**, 33–57 (2021). https://doi.org/10.3390/plants11010033
34 Shekhawat, K., Fröhlich, K., García-Ramírez, G. X., Trapp, M. A. & Hirt, H. Ethylene: A Master Regulator of Plant-Microbe Interactions under Abiotic Stresses. *Cells* **12**, 31–45 (2022). https://doi.org/10.3390/cells12010031
35 Primrose, S. B. & Dilworth, M. J. Ethylene Production by Bacteria. *Microbiology* **93**, 177–181 (1976). https://doi.org/10.1099/00221287-93-1-177
36 Ravanbakhsh, M., Sasidharan, R., Voesenek, L. A. C. J., Kowalchuk, G. A. & Jousset, A. Microbial Modulation of Plant Ethylene Signaling: Ecological and Evolutionary Consequences. *Microbiome* **6**, 52–61 (2018). https://doi.org/10.1186/s40168-018-0436-1
37 Fukuda, H., Ogawa, T. & Tanase, S. in *Advances in Microbial Physiology* Vol. 35 (ed A. H. Rose) 275–306 (Academic Press, 1993).
38 Papon, N. & Binder, B. M. An Evolutionary Perspective on Ethylene Sensing in Microorganisms. *Trends Microbiol.* **27**, 193–196 (2019). https://doi.org/10.1016/j.tim.2018.12.002
39 Carlew, T. S. *et al.* Ethylene Signals through an Ethylene Receptor to Modulate Biofilm Formation and Root Colonization in a Beneficial Plant-associated Bacterium. *PLoS Genet.* **21**, e1011587 (2025). https://doi.org/10.1371/journal.pgen.1011587
40 Ali, J. *et al.* Multifaceted Roles of Ethylene in Plants: From Traditional Functions to Modern Insights into Tritrophic Interactions for Sustainable Agriculture. *Plant Stress* **17**, 100925–100940 (2025). https://doi.org/https://doi.org/10.1016/j.stress.2025.100925
41 Fortt, J. *et al.* Bacterial Modulation of the Plant Ethylene Signaling Pathway Improves Tolerance to Salt Stress in Lettuce (Lactuca sativa L.). *Front. Sustain. Food Syst.* **6**, 768250 (2022). https://doi.org/10.3389/fsufs.2022.768250
42 Fukuda, H., Fujii, T. & Ogawa, T. Preparation of a Cell-free Ethylene-forming System from *Penicillium digitatum*. *Agric. Biol. Chem.* **50**, 977–981 (1986). https://doi.org/10.1271/bbb1961.50.977

43 Kowsalya, K. *et al.* Unraveling the Multifaceted role of Ethephon in Plant Physiology: from Seed Germination to Crop Maturation and Harvesting. *J. Plant Biochem. Biotechnol.* **34**, 639–664 (2025). https://doi.org/10.1007/s13562-024-00953-6

44 Iqbal, N. *et al.* Ethylene Role in Plant Growth, Development and Senescence: Interaction with Other Phytohormones. *Front. Plant. Sci.* **8**, 475 (2017). https://doi.org/10.3389/fpls.2017.00475

45 Uraguchi, D. *et al.* A Femtomolar-range Suicide Germination Stimulant for the Parasitic Plant *Striga hermonthica*. *Science* **362**, 1301–1305 (2018). https://doi.org/10.1126/science.aau5445

46 Runo, S. & Kuria, E. K. Habits of a Highly Successful Cereal Killer, Striga. *PLoS Pathog.* **14**, e1006731 (2018). https://doi.org/10.1371/journal.ppat.1006731

47 Asrey, R. *et al.* Biological and Postharvest Interventions to Manage the Ethylene in Fruit: a Review. *Sustain. Food Technol.* **1**, 803–826 (2023). https://doi.org/10.1039/D3FB00037K

48 Choi, S.-J., Ahn, G.-H., Lee, K. H. & Jahng, W. J. Synthesis of 1-(2,2-Dimethylpropyl)-Cyclopropene (1-DCP) as an Ethylene Antagonist. *ACS Omega.* **8**, 29770–29778 (2023). https://doi.org/10.1021/acsomega.3c04220

49 Pirrung, M. C. *et al.* Ethylene Receptor Antagonists: Strained Alkenes Are Necessary but Not Sufficient. *Chem. Biol.* **15**, 313–321 (2008). https://doi.org/10.1016/j.chembiol.2008.02.018

50 Ariyanto, H. D. & Yoshii, H. in *Functionality of Cyclodextrins in Encapsulation for Food Applications* (eds Thao M. Ho, Hidefumi Yoshii, Keiji Terao, & Bhesh R. Bhandari) 299–313 (Springer International Publishing, 2021).

51 Houben, M. & Van de Poel, B. 1-Aminocyclopropane-1-Carboxylic Acid Oxidase (ACO): The Enzyme That Makes the Plant Hormone Ethylene. *Front. Plant. Sci.* **10**, 695 (2019). https://doi.org/10.3389/fpls.2019.00695

52 Li, C. Y. *et al.* A novel ATP Dependent Dimethylsulfoniopropionate Lyase in Bacteria that releases Dimethyl Sulfide and Acryloyl-CoA. *Elife* **10**, e64045 (2021). https://doi.org/10.7554/eLife.64045

53 Shaw, D. K., Sekar, J. & Ramalingam, P. V. Recent Insights into Oceanic Dimethylsulfoniopropionate Biosynthesis and Catabolism. *Environ. Microbiol.* **24**, 2669–2700 (2022). https://doi.org/https://doi.org/10.1111/1462-2920.16045

54 Li, C. Y. *et al.* Molecular Insight into Bacterial Cleavage of Oceanic Dimethylsulfoniopropionate into Dimethyl Sulfide. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 1026–1031 (2014). https://doi.org/10.1073/pnas.1312354111

55 Zhang, Z. *et al.* Structural and Stereoelectronic Insights into Oxygenase-catalyzed Formation of Ethylene from 2-Oxoglutarate. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 4667–4672 (2017). https://doi.org/10.1073/pnas.1617760114

56 Dhingra, S., Zhang, Z., Lohans, C. T., Brewitz, L. & Schofield, C. J. Substitution of 2-Oxoglutarate Alters Reaction Outcomes of the *Pseudomonas savastanoi* Ethylene-forming Enzyme. *J. Biol. Chem.* **300**, 107546 (2024). https://doi.org/https://doi.org/10.1016/j.jbc.2024.107546

57 Lago-Maciel, A. *et al.* Methylthio-alkane Reductases use Nitrogenase Metalloclusters for Carbon-sulfur Bond Cleavage. *bioRxiv* (2024). https://doi.org/10.1101/2024.10.19.619033

58 North, J. A. *et al.* A Nitrogenase-like Enzyme System Catalyzes Methionine, Ethylene, and Methane Biogenesis. *Science* **369**, 1094–1098 (2020). https://doi.org/10.1126/science.abb6310

59 Miller, A. R., North, J. A., Wildenthal, J. A. & Tabita, F. R. Two Distinct Aerobic Methionine Salvage Pathways Generate Volatile Methanethiol in *Rhodopseudomonas palustris*. *mBio* **9**, e00407–00418 (2018). https://doi.org/10.1128/mBio.00407-18

60 North, J. A., Miller, A. R., Wildenthal, J. A., Young, S. J. & Tabita, F. R. Microbial Pathway for Anaerobic 5'-Methylthioadenosine Metabolism Coupled to Ethylene Formation. *Proc. Natl. Acad. Sci. U. S. A.* **114**, e10455–e10464 (2017). https://doi.org/10.1073/pnas.1711625114

61 Chagué, V., Elad, Y., Barakat, R., Tudzynski, P. & Sharon, A. Ethylene Biosynthesis in *Botrytis cinerea*. *FEMS Microbiol. Ecol.* **40**, 143–149 (2002). https://doi.org/10.1111/j.1574-6941.2002.tb00946.x

62 Fukuda, H., Takahashi, M., Fujii, T., Tazaki, M. & Ogawa, T. An NADH : Fe(III):EDTA Oxidoreductase from *Cryptococcus albidus:* an Enzyme involved in Ethylene Production *in vivo*? *FEMS Microbiol. Lett.* **60**, 107–111 (1989).

63 Yang, J. *et al.* An Insight on the Ethylene Biosynthetic Pathway of Two Major Fruit Postharvest Pathogens with different Host Specificity: P*enicillium digitatum* and *Penicillium expansum*. *Eur. J. Plant Pathol.* **149**, 575–585 (2017). https://doi.org/10.1007/s10658-017-1205-x

64 Yasumura, Y., Pierik, R., Fricker, M. D., Voesenek, L. A. & Harberd, N. P. Studies of *Physcomitrella patens* reveal that Ethylene-mediated Submergence Responses arose relatively early in Land-plant Evolution. *Plant J.* **72**, 947–959 (2012). https://doi.org/10.1111/tpj.12005

65 Linh T. B., Amelia H., Erin E. I. & C., C.-L. The Effects of Sugars and Ethylene on Apospory and Regeneration in *Ceratopteris richardii*. *Am. J. Plant Sci.* **3**, 953–960 (2012).

66 Lennicke, C. & Cochemé, H. M. Redox Metabolism: ROS as Specific Molecular Regulators of Cell Signaling and Function. *Mol. Cell* **81**, 3691–3707 (2021). https://doi.org/https://doi.org/10.1016/j.molcel.2021.08.018

67 Ballester, A. R. & González-Candelas, L. EFE-Mediated Ethylene Synthesis Is the Major Pathway in the Citrus Postharvest Pathogen *Penicillium digitatum* during Fruit Infection. *J. Fungi* **6**, 175–192 (2020). https://doi.org/10.3390/jof6030175

68 Paardekooper, L. M. *et al.* Ethylene, an Early Marker of Systemic Inflammation in Humans. *Sci. Rep.* **7**, 6889–6898 (2017). https://doi.org/10.1038/s41598-017-05930-9

69 Adams, D. O. & Yang, S. F. Ethylene biosynthesis: Identification of 1-aminocyclopropane-1-carboxylic acid as an intermediate in the conversion of methionine to ethylene. *Proc Natl Acad Sci U S A* **76**, 170–174 (1979). https://doi.org/10.1073/pnas.76.1.170

70 Pattyn, J., Vaughan-Hirsch, J. & Van de Poel, B. The Regulation of Ethylene Biosynthesis: a Complex Multilevel Control Circuitry. *New Phytol.* **229**, 770–782 (2021). https://doi.org/https://doi.org/10.1111/nph.16873

71 Khan, S., Alvi, A. F., Saify, S., Iqbal, N. & Khan, N. A. The Ethylene Biosynthetic Enzymes, 1-Aminocyclopropane-1-Carboxylate (ACC) Synthase (ACS) and ACC Oxidase (ACO): The Less Explored Players in Abiotic Stress Tolerance. *Biomolecules* **14**, 90–123 (2024). https://doi.org/10.3390/biom14010090

72 Bradford, K. J. Shang Fa Yang: Pioneer in Plant Ethylene Biochemistry. *Plant Sci.* **175**, 2–7 (2008). https://doi.org/https://doi.org/10.1016/j.plantsci.2008.01.005

73 Chen, H. *et al.* The Yang cycle in plants: a journey of methionine recycling with fascinating metabolites and enzymes. *Plant Hormones* **1** (2025). https://doi.org/10.48130/ph-0025-0007

74 Chen, H. *et al.* The Yang Cycle in Plants: a journey of Methionine Recycling with Fascinating Metabolites and Enzymes. *Plant Hormones* **1**, e007 (2025). https://doi.org/10.48130/ph-0025-0007

75 Bhatta, U. K. Alternative Management Approaches of Citrus Diseases Caused by *Penicillium digitatum* (Green Mold) and *Penicillium italicum* (Blue Mold). *Front. Plant. Sci.* **12**, 833328 (2021). https://doi.org/10.3389/fpls.2021.833328

76 Eckert, C. *et al.* Ethylene-forming Enzyme and Bioethylene Production. *Biotechnol. Biofuels* **7**, 33–42 (2014). https://doi.org/10.1186/1754-6834-7-33

77 Miller, E. V., Winston, J. & Fisher, D. Production of Epinasty by Emanations from Normal and Decaying Citrus Fruits and from *Penicillium digitatum*. *J. Agric. Res* **60**, 269–277 (1940).

78 Biale, J. B. Effect of Emanations from Several Species of Fungi on Respiration and Color Development of Citrus Fruits. *Science* **91**, 458–459 (1940). https://doi.org/doi:10.1126/science.91.2367.458.b

79 Primrose, S. B. Ethylene and Agriculture: the Role of the Microbe. *J. Appl. Bacteriol.* **46**, 1–25 (1979). https://doi.org/10.1111/j.1365-2672.1979.tb02579.x

80 Thomas, T. H. Hormonal Regulation of Development. III. Role of Environment Factors. *J. Exp. Bot.* **39**, 1327–1329 (1988).

81 Ajdanian, L. *et al.* in *The Role of Growth Regulators and Phytohormones in Overcoming Environmental Stress* (eds Anket Sharma *et al.*) 113–135 (Academic Press, 2023).

82 Jacobsen, D. W. & Wang, C. H. The Biogenesis of Ethylene in *Penicillium digitatum*. *Plant Physiol.* **43**, 1959–1966 (1968). https://doi.org/10.1104/pp.43.12.1959

83 Nagahama, K. *et al.* Purification and Properties of an Ethylene-forming Enzyme from *Pseudomonas syringae* pv. *phaseolicola* PK2. *J. Gen. Microbiol.* **137**, 2281–2286 (1991). https://doi.org/10.1099/00221287-137-10-2281

84 Chatterjee, S. *et al.* Biochemical, Structural, and Conformational Characterization of a Fungal Ethylene-Forming Enzyme. *Biochemistry* **64**, 2054–2067 (2025). https://doi.org/10.1021/acs.biochem.5c00038
85 Barney, B. M., Igarashi, R. Y., Dos Santos, P. C., Dean, D. R. & Seefeldt, L. C. Substrate Interaction at an Iron-Sulfur Face of the FeMo-cofactor during Nitrogenase Catalysis. *J. Biol. Chem.* **279**, 53621–53624 (2004). https://doi.org/https://doi.org/10.1074/jbc.M410247200
86 Seefeldt, L. C. *et al.* Reduction of Substrates by Nitrogenases. *Chem. Rev.* **120**, 5082–5106 (2020). https://doi.org/10.1021/acs.chemrev.9b00556
87 Hu, Y. *et al.* Enzymatic Fischer-Tropsch-Type Reactions. *Chem. Rev.* **123**, 5755–5797 (2023). https://doi.org/10.1021/acs.chemrev.2c00612
88 Lee, C. C., Hu, Y. & Ribbe, M. W. Tracing the Hydrogen Source of Hydrocarbons formed by Vanadium Nitrogenase. *Angew. Chem., Int. Ed. Engl.* **50**, 5545–5547 (2011). https://doi.org/10.1002/anie.201100869
89 Hu, Y., Lee, C. C. & Ribbe, M. W. Extending the carbon chain: hydrocarbon formation catalyzed by vanadium/molybdenum nitrogenases. *Science* **333**, 753–755 (2011). https://doi.org/10.1126/science.1206883
90 Lee, C. C., Hu, Y. & Ribbe, M. W. Vanadium Nitrogenase Reduces CO. *Science* **329**, 642–642 (2010). https://doi.org/doi:10.1126/science.1191455
91 Xu, F., Peng, Y., He, Z.-Q. & Yu, L.-L. The role of Cyanoalanine Synthase and Alternative Oxidase in Promoting Salt Stress Tolerance in *Arabidopsis thaliana*. *BMC Plant Biol.* **23**, 163 (2023). https://doi.org/10.1186/s12870-023-04167-1
92 Elias, M., Sudhakaran, P. R. & Nambisan, B. Purification and Characterisation of β-Cyanoalanine Synthase from Cassava Tissues. *Phytochemistry* **46**, 469–472 (1997). https://doi.org/https://doi.org/10.1016/S0031-9422(97)00305-1
93 Machingura, M., Salomon, E., Jez, J. M. & Ebbs, S. D. The β-Cyanoalanine Synthase Pathway: Beyond Cyanide Detoxification. *Plant Cell Environ.* **39**, 2329–2341 (2016). https://doi.org/https://doi.org/10.1111/pce.12755
94 *2-Oxoglutarate-Dependent Oxygenases*. (The Royal Society of Chemistry, 2015).
95 Cheng, J. *et al.* Functional Analysis of the Gibberellin 2-Oxidase Gene Family in Peach. *Front. Plant Sci.* **12**, 619158 (2021). https://doi.org/10.3389/fpls.2021.619158
96 Markolovic, S., Wilkins, S. E. & Schofield, C. J. Protein Hydroxylation Catalyzed by 2-Oxoglutarate-dependent Oxygenases. *J. Biol. Chem.* **290**, 20712–20722 (2015). https://doi.org/10.1074/jbc.R115.662627
97 Fletcher, S. C. & Coleman, M. L. Human 2-Oxoglutarate-dependent Oxygenases: Nutrient Sensors, Stress Responders, and Disease Mediators. *Biochem. Soc. Trans.* **48**, 1843–1858 (2020). https://doi.org/10.1042/bst20190333
98 Hancock, R. L., Dunne, K., Walport, L. J., Flashman, E. & Kawamura, A. Epigenetic Regulation by Histone Demethylases in Hypoxia. *Epigenomics* **7**, 791–811 (2015). https://doi.org/10.2217/epi.15.24
99 Choi, H. *et al.* A Human Protein Hydroxylase that Accepts D-Residues. *Commun. Chem.* **3**, 52 (2020). https://doi.org/10.1038/s42004-020-0290-5
100 Crüsemann, M. Highlights in (Bio)Synthesis and Applications of Bioactive Metabolites: The 37th Irsee Natural Product Symposium. *Eur. J. Org. Chem.* **28**, 2500350 (2025). https://doi.org/https://doi.org/10.1002/ejoc.202500350
101 Zhao, S., Wu, L., Xu, Y. & Nie, Y. Fe(II) and 2-Oxoglutarate-dependent Dioxygenases for Natural Product Synthesis: Molecular Insights into Reaction Diversity. *Nat. Prod. Rep.* **42**, 67–92 (2024). https://doi.org/https://doi.org/10.1039/d4np00030g
102 Clifton, I. J. *et al.* Structural Studies on 2-Oxoglutarate Oxygenases and Related Double-stranded β-helix Fold Proteins. *J. Inorg. Biochem.* **100**, 644–669 (2006). https://doi.org/https://doi.org/10.1016/j.jinorgbio.2006.01.024
103 Martinez, S. & Hausinger, R. P. Catalytic Mechanisms of Fe(II)- and 2-Oxoglutarate-dependent Oxygenases. *J. Biol. Chem.* **290**, 20702–20711 (2015). https://doi.org/10.1074/jbc.R115.648691
104 Zhang, X. *et al.* Structure-guided Analysis of *Arabidopsis* JASMONATE-INDUCED OXYGENASE (JOX) 2 Reveals Key Residues for Recognition of Jasmonic Acid Substrate by

Plant JOXs. *Mol. Plant* **14**, 820–828 (2021). https://doi.org/https://doi.org/10.1016/j.molp.2021.01.017
105 Bilgin, N. *et al.* Substrate Selectivity and Inhibition of the Human Lysyl Hydroxylase JMJD7. *Protein Sci.* **33**, e5162 (2024). https://doi.org/https://doi.org/10.1002/pro.5162
106 Rabe, P. *et al.* Anaerobic Fixed-target Serial Crystallography. *IUCrJ* **7**, 901–912 (2020). https://doi.org/doi:10.1107/S2052252520010374
107 Dong, J. *et al.* Structural Insights of 4-Hydrophenylpyruvate dioxygenase Inhibition by Structurally Diverse Small Molecules. *Adv Agrochem.* **1**, 174–181 (2022). https://doi.org/https://doi.org/10.1016/j.aac.2022.10.002
108 Matringe, M., Sailland, A., Pelissier, B., Rolland, A. & Zink, O. p-Hydroxyphenylpyruvate Dioxygenase Inhibitor-resistant Plants. *Pest Manag. Sci.* **61**, 269–276 (2005). https://doi.org/10.1002/ps.997
109 Maeda, H. *et al.* A Rice Gene that Confers Broad-spectrum Resistance to β-Triketone Herbicides. *Science* **365**, 393–396 (2019). https://doi.org/doi:10.1126/science.aax0379
110 Dong, J. *et al.* An Artificially Evolved Gene for Herbicide-resistant Rice Breeding. *Proc. Natl. Acad. Sci. U. S. A.* **121**, e2407285121 (2024). https://doi.org/doi:10.1073/pnas.2407285121
111 Wu, J. *et al.* Directed Evolution of HIS1-Like Proteins to Enhance Herbicide Metabolism in Crops. *Adv. Agrochem.* **in press** (2025). https://doi.org/https://doi.org/10.1016/j.aac.2025.05.005
112 Takamura, N. *et al.* Catalytic Promiscuity of Rice 2-Oxoglutarate/Fe(II)-dependent Dioxygenases Supports Xenobiotic Metabolism. *Plant Physiol.* **187**, 816–828 (2021). https://doi.org/10.1093/plphys/kiab293
113 Smith, J. J. & John, P. Activation of 1-Aminocyclopropane-1-carboxylate Oxidase by Bicarbonate/carbon dioxide. *Phytochemistry* **32**, 1381–1386 (1993). https://doi.org/https://doi.org/10.1016/0031-9422(93)85142-E
114 Poneleit, L. S. & Dilley, D. R. Carbon Dioxide Activation of 1-Aminocyclopropane-1-carboxylate (ACC) Oxidase in Ethylene Biosynthesis. *Postharvest Biol. Technol.* **3**, 191–199 (1993). https://doi.org/https://doi.org/10.1016/0925-5214(93)90055-8
115 Fernandezmaculet, J. C., Dong, J. G. & Yang, S. F. Activation of 1-Aminocyclopropane-1-carboxylate Oxidase by Carbon Dioxide. *Biochem. Biophys. Res. Commun.* **193**, 1168–1173 (1993). https://doi.org/https://doi.org/10.1006/bbrc.1993.1748
116 Dong, J. G., Fernández-Maculet, J. C. & Yang, S. F. Purification and Characterization of 1-Aminocyclopropane-1-carboxylate Oxidase from Apple Fruit. *Proc. Natl. Acad. Sci. U. S. A.* **89**, 9789–9793 (1992). https://doi.org/doi:10.1073/pnas.89.20.9789
117 Baldwin, J. E. *et al.* On the Biosynthesis of Ethylene: Further Evidence for Stepwise Enzymatic Cyclopropane Ring Cleavage. *J. Chem. Soc., Chem. Commun.*, 775–777 (1988). https://doi.org/10.1039/C39880000775
118 Gunawardana, D. M., Yosaatmadja, Y., Leung, I. K. & Squire, C. J. 5TCV: ACC Oxidase Complex with Substrate 1-Aminocyclopropane-1-carboxylic acid. *Protein Data Bank* (2017). https://doi.org/10.2210/pdb5TCV/pdb
119 Rocklin, A. M. *et al.* Role of the Nonheme Fe(II) Center in the Biosynthesis of the Plant Hormone Ethylene. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 7905–7909 (1999). https://doi.org/10.1073/pnas.96.14.7905
120 Zhang, Z., Ren, J.-S., Clifton, I. J. & Schofield, C. J. Crystal Structure and Mechanistic Implications of 1-Aminocyclopropane-1-Carboxylic Acid Oxidase—The Ethylene-Forming Enzyme. *Chem. Biol.* **11**, 1383–1394 (2004). https://doi.org/https://doi.org/10.1016/j.chembiol.2004.08.012
121 Rocklin, A. M., Kato, K., Liu, H.-w., Que, L. & Lipscomb, J. D. Mechanistic Studies of 1-Aminocyclopropane-1-carboxylic Acid Oxidase: Single Turnover Reaction. *J. Biol. Inorg. Chem.* **9**, 171–182 (2004). https://doi.org/10.1007/s00775-003-0510-3
122 Thrower, J. S., Blalock, R. & Klinman, J. P. Steady-State Kinetics of Substrate Binding and Iron Release in Tomato ACC Oxidase. *Biochemistry* **40**, 9717–9724 (2001). https://doi.org/10.1021/bi010329c
123 Zhou, J., Rocklin, A. M., Lipscomb, J. D., Que, L. & Solomon, E. I. Spectroscopic Studies of 1-Aminocyclopropane-1-carboxylic Acid Oxidase: Molecular Mechanism and CO_2 Activation

in the Biosynthesis of Ethylene. *J. Am. Chem. Soc.* **124**, 4602–4609 (2002). https://doi.org/10.1021/ja017250f
124 Chapman, N. C. & Rutledge, P. J. Isopenicillin *N* Synthase: Crystallographic Studies. *ChemBioChem* **22**, 1687–1705 (2021). https://doi.org/https://doi.org/10.1002/cbic.202000743
125 Elkins, J. M. *et al.* Crystallographic Studies on the Reaction of Isopenicillin *N* Synthase with an Unsaturated Substrate Analogue. *Org. Biomol. Chem.* **1**, 1455–1460 (2003). https://doi.org/10.1039/B212270G
126 Dilley, D. R. *et al.* 1-Aminocyclopropane-1-carboxylic Acid Oxidase Reaction Mechanism and Putative Post-translational Activities of the ACCO Protein. *AoB Plants* **5**, plt031 (2013). https://doi.org/10.1093/aobpla/plt031
127 Thilakarathne, A. S., Liu, F. & Zou, Z. Plant Signaling Hormones and Transcription Factors: Key Regulators of Plant Responses to Growth, Development, and Stress. *Plants* **14**, 1070–1124 (2025). https://doi.org/10.3390/plants14071070
128 Ahmadizadeh, M., Chen, J. T., Hasanzadeh, S., Ahmar, S. & Heidari, P. Insights into the Genes Involved in the Ethylene Biosynthesis pathway in *Arabidopsis thaliana* and *Oryza sativa*. *J. Genet. Eng. Biotechnol.* **18**, 62–81 (2020). https://doi.org/10.1186/s43141-020-00083-1
129 Xu, J. *et al.* Activation of MAPK Kinase 9 Induces Ethylene and Camalexin Biosynthesis and Enhances Sensitivity to Salt Stress in Arabidopsis*. *J. Biol. Chem.* **283**, 26996–27006 (2008). https://doi.org/https://doi.org/10.1074/jbc.M801392200
130 Hahn, A. & Harter, K. Mitogen-activated Protein Kinase Cascades and Ethylene: Signaling, Biosynthesis, or Both? *Plant Physiol.* **149**, 1207–1210 (2009). https://doi.org/10.1104/pp.108.132241
131 Vanderstraeten, L. & Van Der Straeten, D. Accumulation and Transport of 1-Aminocyclopropane-1-Carboxylic Acid (ACC) in Plants: Current Status, Considerations for Future Research and Agronomic Applications. *Front. Plant Sci.* **8**, 38 (2017). https://doi.org/10.3389/fpls.2017.00038
132 Westfall, C. S., Muehler, A. M. & Jez, J. M. Enzyme Action in the Regulation of Plant Hormone Responses. *J. Biol. Chem.* **288**, 19304–19311 (2013). https://doi.org/10.1074/jbc.R113.475160
133 Nascimento, F. X., Rossi, M. J. & Glick, B. R. Ethylene and 1-Aminocyclopropane-1-carboxylate (ACC) in Plant-Bacterial Interactions. *Front. Plant. Sci.* **9**, 114–130 (2018). https://doi.org/10.3389/fpls.2018.00114
134 Shahid, M. *et al.* Bacterial ACC Deaminase: Insights into Enzymology, Biochemistry, Genetics, and Potential Role in Amelioration of Environmental Stress in Crop Plants. *Front. Microbiol.* **14**, e1132770 (2023). https://doi.org/10.3389/fmicb.2023.1132770
135 Glick, B. R. *et al.* Promotion of Plant Growth by Bacterial ACC Deaminase. *Crit. Rev. Plant Sci.* **26**, 227–242 (2007). https://doi.org/10.1080/07352680701572966
136 Gamalero, E. & Glick, B. R. Bacterial Modulation of Plant Ethylene Levels. *Plant Physiol.* **169**, 13–22 (2015). https://doi.org/10.1104/pp.15.00284
137 Gamalero, E., Lingua, G. & Glick, B. R. Ethylene, ACC, and the Plant Growth-Promoting Enzyme ACC Deaminase. *Biology* **12**, 1043–1060 (2023). https://doi.org/10.3390/biology12081043
138 Singh, R. P., Shelke, G. M., Kumar, A. & Jha, P. N. Biochemistry and genetics of ACC deaminase: a weapon to "stress ethylene" produced in plants. *Front. Microbiol.* **6**, 937 (2015). https://doi.org/10.3389/fmicb.2015.00937
139 Johansson, N., Persson, K. O., Larsson, C. & Norbeck, J. Comparative Sequence Analysis and Mutagenesis of Ethylene Forming Enzyme (EFE) 2-Oxoglutarate/Fe(II)-dependent Dioxygenase Homologs. *BMC Biochem.* **15**, 22–29 (2014). https://doi.org/10.1186/1471-2091-15-22
140 Xue, J., Lu, J. & Lai, W. Mechanistic Insights into a Non-heme 2-Oxoglutarate-dependent Ethylene-forming Enzyme: Selectivity of Ethylene-formation versus L-Arg Hydroxylation. *Phys. Chem. Chem. Phys.* **21**, 9957–9968 (2019). https://doi.org/10.1039/C9CP00794F
141 Cui, Y. *et al.* Discovery of Five New Ethylene-Forming Enzymes for Clean Production of Ethylene in *E. coli*. *Int. J. Mol. Sci.* **23**, 4500–4515 (2022).
142 Martinez, S. & Hausinger, R. P. Biochemical and Spectroscopic Characterization of the Non-Heme Fe(II)- and 2-Oxoglutarate-Dependent Ethylene-Forming Enzyme from *Pseudomonas*

syringae pv. *phaseolicola* PK2. *Biochemistry* **55**, 5989–5999 (2016). https://doi.org/10.1021/acs.biochem.6b00890

143 Mitchell, R. E. & Bieleski, R. L. Involvement of Phaseolotoxin in Halo Blight of Beans: Transport and Conversion to Functional Toxin. *Plant Physiol.* **60**, 723–729 (1977). https://doi.org/10.1104/pp.60.5.723

144 Arvizu-Gómez, J. L., Hernández-Morales, A., Campos-Guillén, J., González-Reyes, C. & Pacheco-Aguilar, J. R. Phaseolotoxin: Environmental Conditions and Regulatory Mechanisms Involved in Its Synthesis. *Microorganisms* **12**, 1300–1321 (2024). https://doi.org/10.3390/microorganisms12071300

145 Langley, D. B., Templeton, M. D., Fields, B. A., Mitchell, R. E. & Collyer, C. A. Mechanism of Inactivation of Ornithine Transcarbamoylase by N δ-(N′-Sulfodiaminophosphinyl)-l-ornithine, a True Transition State Analogue?: CRYSTAL STRUCTURE AND IMPLICATIONS FOR CATALYTIC MECHANISM*. *J. Biol. Chem.* **275**, 20012–20019 (2000). https://doi.org/https://doi.org/10.1074/jbc.M000585200

146 Weingart, H., Ullrich, H., Geider, K. & Völksch, B. The Role of Ethylene Production in Virulence of *Pseudomonas syringae* pvs. *glycinea* and *phaseolicola*. *Phytopathology* **91**, 511–518 (2001). https://doi.org/10.1094/phyto.2001.91.5.511

147 Guan, R. *et al.* Multilayered Regulation of Ethylene Induction Plays a Positive Role in *Arabidopsis* Resistance against *Pseudomonas syringae*. *Plant Physiol.* **169**, 299–312 (2015). https://doi.org/10.1104/pp.15.00659

148 Watanabe, K., Nagahama, K. & Sato, M. A Conjugative Plasmid Carrying the *efe* Gene for the Ethylene-Forming Enzyme Isolated from *Pseudomonas syringae* pv. *glycinea*. *Phytopathology* **88**, 1205–1209 (1998). https://doi.org/10.1094/phyto.1998.88.11.1205

149 Weingart, H. & Volksch, E. Ethylene Production by *Pseudomonas syringae* Pathovars *In Vitro* and *In Planta*. *Appl. Environ. Microbiol.* **63**, 156–161 (1997).

150 Veetil, V. P., Angermayr, S. A. & Hellingwerf, K. J. Ethylene production with engineered Synechocystis sp PCC 6803 strains. *Microb Cell Fact* **16**, 34 (2017). https://doi.org/10.1186/s12934-017-0645-5

151 Wang, J. P. *et al.* Metabolic Engineering for Ethylene Production by Inserting the Ethylene-forming Enzyme Gene (efe) at the 16S rDNA Sites of *Pseudomonas putida* KT2440. *Bioresour. Technol.* **101**, 6404–6409 (2010). https://doi.org/10.1016/j.biortech.2010.03.030

152 Martinez, S. *et al.* Structures and Mechanisms of the Non-Heme Fe(II)- and 2-Oxoglutarate-Dependent Ethylene-Forming Enzyme: Substrate Binding Creates a Twist. *J. Am. Chem. Soc.* **139**, 11980–11988 (2017). https://doi.org/10.1021/jacs.7b06186

153 Copeland, R. A. *et al.* Hybrid Radical-polar Pathway for Excision of Ethylene from 2-Oxoglutarate by an Iron Oxygenase. *Science* **373**, 1489–1493 (2021). https://doi.org/doi:10.1126/science.abj4290

154 Liu, X. *et al.* Natural and Synthetic 2-Oxoglutarate Derivatives are Substrates for Oncogenic Variants of Human Isocitrate Dehydrogenase 1 and 2. *J. Biol. Chem.* **299**, 102873–102889 (2023). https://doi.org/10.1016/j.jbc.2023.102873

155 Brewitz, L., Nakashima, Y. & Schofield, C. J. Synthesis of 2-Oxoglutarate Derivatives and their Evaluation as Cosubstrates and Inhibitors of Human Aspartate/asparagine-β-hydroxylase. *Chem. Sci.* **12**, 1327–1342 (2021). https://doi.org/10.1039/D0SC04301J

156 Nakashima, Y., Brewitz, L., Tumber, A., Salah, E. & Schofield, C. J. 2-Oxoglutarate Derivatives can Selectively Enhance or Inhibit the Activity of Human Oxygenases. *Nat. Commun.* **12**, 6478–6494 (2021). https://doi.org/10.1038/s41467-021-26673-2

157 Burke, E. J., Copeland, R. A., Dixit, Y., Krebs, C. & Bollinger, J. M., Jr. Steric Perturbation of the Grob-like Final Step of Ethylene-Forming Enzyme Enables 3-Hydroxypropionate and Propylene Production. *J. Am. Chem. Soc.* **146**, 1977–1983 (2024). https://doi.org/10.1021/jacs.3c09733

158 Jaber Sathik Rifayee, S. B., Thomas, M. G. & Christov, C. Z. Revealing the Nature of the Second Branch Point in the Catalytic Mechanism of the Fe(II)/2OG-dependent Ethylene Forming Enzyme. *Chem. Sci.* **16**, 7667–7684 (2025). https://doi.org/10.1039/D4SC08378D

159 Chaturvedi, S. S. *et al.* Dioxygen Binding Is Controlled by the Protein Environment in Non-heme Fe(II) and 2-Oxoglutarate Oxygenases: A Study on Histone Demethylase PHF8 and an

Ethylene-Forming Enzyme. *Chemistry* **29**, e202300854 (2023). https://doi.org/10.1002/chem.202300854
160 Hausinger, R. P. Fe II/Alpha-ketoglutarate-dependent Hydroxylases and Related Enzymes. *Crit. Rev. Biochem. Mol. Biol.* **39**, 21–68 (2004). https://doi.org/10.1080/10409230490440541
161 Copeland, R. A. *et al.* An Iron(IV)–Oxo Intermediate Initiating l-Arginine Oxidation but Not Ethylene Production by the 2-Oxoglutarate-Dependent Oxygenase, Ethylene-Forming Enzyme. *J. Am. Chem. Soc.* **143**, 2293–2303 (2021). https://doi.org/10.1021/jacs.0c10923
162 Nagahama, K. *et al.* Site-directed Mutagenesis of Histidine Residues in the Ethylene-forming Enzyme from *Pseudomonas syringae*. *J. Ferment. Bioeng.* **85**, 255–258 (1998). https://doi.org/https://doi.org/10.1016/S0922-338X(97)85671-1
163 Robert, X. & Gouet, P. Deciphering Key Features in Protein Structures with the New ENDscript Server. *Nucleic Acids Res.* **42**, W320–W324 (2014). https://doi.org/10.1093/nar/gku316
164 McGarvey, D. J. & Christoffersen, R. E. Characterization and Kinetic Parameters of Ethylene-forming Enzyme from Avocado Fruit. *J. Biol. Chem.* **267**, 5964–5967 (1992). https://doi.org/https://doi.org/10.1016/S0021-9258(18)42649-X
165 Fernández-Maculet, J. C. & Yang, S. F. Extraction and Partial Characterization of the Ethylene-forming Enzyme from Apple Fruit. *Plant Physiol.* **99**, 751–754 (1992). https://doi.org/10.1104/pp.99.2.751
166 Zhang, Z., Schofield, C. J., Baldwin, J. E., Thomas, P. & John, P. Expression, Purification and Characterization of 1-Aminocyclopropane-1-carboxylate Oxidase from Tomato in *Escherichia coli*. *Biochem. J.* **307 (Pt 1)**, 77–85 (1995). https://doi.org/10.1042/bj3070077
167 McKeon, T. A. & Shang Fa, Y. A Comparison of the Conversion of 1-Amino-2-ethylcyclopropane-1-carboxylic Acid Stereoisomers to 1-Butene by Pea Epicotyls and by a Cell-free System. *Planta* **160**, 84–87 (1984). https://doi.org/10.1007/bf00392470
168 Zhang, Z., Schofield, C. J., Baldwin, J. E., Thomas, P. & John, P. Expression, Purification and Characterization of 1-Aminocyclopropane-1-carboxylate Oxidase from Tomato in *Escherichia coli*. *Biochem. J.* **307**, 77–85 (1995). https://doi.org/10.1042/bj3070077
169 Dequina, H. J., Jones, C. L. & Schomaker, J. M. Recent Updates and Future Perspectives in Aziridine Synthesis and Reactivity. *Chem.* **9**, 1658–1701 (2023). https://doi.org/https://doi.org/10.1016/j.chempr.2023.04.010
170 Goldberg, N. W., Knight, A. M., Zhang, R. K. & Arnold, F. H. Nitrene Transfer Catalyzed by a Non-Heme Iron Enzyme and Enhanced by Non-Native Small-Molecule Ligands. *J. Am. Chem. Soc.* **141**, 19585–19588 (2019). https://doi.org/10.1021/jacs.9b11608
171 Ren, X. & Fasan, R. Engineered and Artificial Metalloenzymes for Selective C–H Functionalization. *Curr. Opin. Green Sustain. Chem.* **31**, 100494–100507 (2021). https://doi.org/https://doi.org/10.1016/j.cogsc.2021.100494

CHAPTER 2

MANUSCRIPT I

Substitution of 2-oxoglutarate alters reaction outcomes of the *Pseudomonas savastanoi* ethylene-forming enzyme

Contents

2.1 Overview **64**
2.2 Abstract **65**
2.3 Introduction **66**
2.4 Results **69**
2.4.1 NMR assays for monitoring PsEFE catalysis **69**
2.4.2 PsEFE catalyzes conversion of 2-oxoadipate to γ-hydroxybutyrate **72**
2.4.3 The 2OG substitution pattern affects the reaction outcomes of PsEFE catalysis **76**
2.5 Discussion **84**
2.6 Experimental Procedures **89**
2.7 Acknowledgements **92**
2.8 Author Contributions **93**
2.9 References **93**
2.10 Supporting Information **99**
2.10.1 General Information **99**
2.10.2 Supporting Figures **100**
2.10.3 Supporting Tables **159**
2.10.4 Supporting Information References **160**
2.11 Statement of Authorship **161**

2.1 Overview

2-Oxoglutarate (2OG) oxygenases catalyse diverse oxidative transformations, typically by coupling substrate hydroxylation with oxidative decarboxylation of 2OG to succinic acid[1]. The *Pseudomonas savastanoi* pv. *phaseolicola* PK2 ethylene-forming enzyme, PsEFE, is a unique member of the 2OG oxygenase structural superfamily (Figure 2.1). The reaction of PsEFE with substituted 2OG derivatives led to the discovery of a novel reaction pathway, involving the formation of an alcohol,[44] not observed for 2OG, alongside the expected diacid product (Section 1.7).

Structural and mechanistic studies on PsEFE have expanded our understanding of its reactivity; this chapter contains a manuscript describing studies on the impact of substrate structure on the partitioning of bifurcating reaction pathways of PsEFE. ^{1}H NMR assays were used to study the effect of C3- and C4-substitution of 2OG on PsEFE catalysis. The results demonstrate that cosubstrate structure can bias the outcome of PsEFE catalysis, influencing product partition ratios between competing reaction pathways. The results suggest that rational substrate design can modulate bifurcating reactions of PsEFE, and potentially other 2OG oxygenases, for biocatalytic applications.

This work is reported in the *Journal of Biological Chemistry* as:

Dhingra, S.; Zhang, Z.; Lohans, C. T.; Brewitz, L.; Schofield, C. J. Substitution of 2-oxoglutarate alters reaction outcomes of the *Pseudomonas savastanoi* ethylene-forming enzyme. *J. Biol. Chem.* **2024**, *300*, 107546.

Each author contribution is described in the Author Contributions (Section 2.8). A Statement of Authorship (Section 2.11) is given at the end of the chapter.

2.2 Abstract

In seeding plants, biosynthesis of the phytohormone ethylene, which regulates processes including fruit ripening and senescence, is catalyzed by 1-aminocyclopropyl-1-carboxylic acid (ACC) oxidase. The plant pathogen *Pseudomonas savastanoi* (previously classified as: *P. syringae*) employs a different type of ethylene-forming enzyme (PsEFE), though from the same structural superfamily as ACC oxidase, to catalyze ethylene formation from 2-oxoglutarate (2OG) in an arginine dependent manner. PsEFE also catalyzes the more typical oxidation of arginine to give L-Δ^1-pyrroline-5-carboxylate (P5C), a reaction coupled to oxidative decarboxylation of 2OG giving succinate and CO_2. We report on the effects of C3 and/or C4 substituted 2OG derivatives on the reaction modes of PsEFE. ^{1}H NMR assays, including using the pure shift method, reveal that, within our limits of detection, none of the tested 2OG derivatives is converted to an alkene; some are converted to the corresponding β-hydroxypropionate or succinate derivatives, with only the latter being coupled to arginine oxidation. The NMR results reveal that the nature of 2OG derivatization can affect the outcome of the bifurcating reaction, with some 2OG derivatives exclusively favoring the arginine oxidation pathway. Given that some of the tested 2OG derivatives are natural products, the results are of potential biological relevance. There are also opportunities for therapeutic or biocatalytic regulation of the outcomes of reactions catalyzed by 2OG-dependent oxygenases by the use of 2OG derivatives.

2.3 Introduction

2-Oxoglutarate (2OG, **1**)- and non-heme Fe(II)-dependent oxygenases are widely distributed in aerobic biology; catalysis by them typically involves coupling the oxidative decarboxylation of 2OG to give succinate and CO_2 with the 2 electron oxidation of their prime substrates (1). There is a wealth of biochemical data showing the efficiency of 2OG as a cosubstrate for the 2OG oxygenases (1). It is, however, feasible that 2OG oxygenases can employ other (co)substrates instead of, or as well as, 2OG in cells; indeed, 2-oxoacids other than 2OG are natural products and many are abundant in cells (2-5). This proposal is supported by the flexibility of 2OG oxygenases with respect to the type of their prime substrate oxidations, including hydroxylation-, dealkylation-, halogenation-, desaturation-, ring formation-, and ring expansion-reactions (1). The flexibility of 2OG oxygenase-type active sites is further highlighted by the existence of enzymes belonging to the same structural cupin fold subfamily as the 2OG oxygenases, but which catalyze oxidations without a need for 2OG as a cosubstrate. Two prominent examples are isopenicillin N synthase (IPNS) (6, 7) and 1-aminocyclopropyl-1-carboxylic acid (ACC) oxidase (ACCO), the latter of which catalyzes the biosynthesis of the plant hormone ethylene (Figure 2.1a) (8, 9), which regulates processes including fruit ripening and senescence (9-11).




Figure 2.1. The bifurcating reaction of 2OG during PsEFE catalysis. (**a**) Plant ACCOs catalyze ethylene formation from 1-aminocyclopropyl-1-carboxylic acid (ACC) (8, 9). The role(s) of L-ascorbate and bicarbonate in ACCO catalysis in cells is unclear; (**b**) PsEFE catalyzes: (i) the oxidative decarboxylation of 2OG (**1**) coupled to C5 oxidation of L-arginine to give 5-hydroxyarginine which fragments to guanidine and L-Δ^1-pyrroline-5-carboxylate (P5C), and (ii) fragmentation of 2OG (**1**) to give ethylene and CO_2, a reaction outcome in which Arg is not modified (11, 37, 39-41).

The human lysine and tryptophan catabolite 2-oxoadipate (2OA) (12, 13), the carbon scaffold of which is elongated by one methylene compared to 2OG, can sustain catalysis *in vitro* by some human and bacterial 2OG oxygenases (14-21). We and others have shown that the cosubstrate scope of some, but not all, human 2OG oxygenases extends to C3 and/or C4 substituted 2OG derivatives, some of which are natural products (22-26). Recently reported mass spectrometry (MS)-based assays with isolated recombinant human aspartate/asparagine-β-hydroxylase (AspH) and factor inhibiting hypoxia-inducible factor HIF-α (FIH), which both catalyze the 2OG-dependent post-translational hydroxylation of aspartate and asparagine residues in proteins (27-34), have revealed that AspH and FIH prefer structurally distinct sets of C3 and/or C4 substituted 2OG derivatives as cosubstrates(23). Kinetic studies implied that some of the tested 2OG derivatives react as efficiently as 2OG with AspH, FIH, and human Jumonji-C (JmjC) domain-containing protein 5 (JMJD5) (22-24). By contrast, the human JmjC histone lysine-specific N^{ε}-demethylases 4A and 4E (KDM4A and KDM4E) have not been observed to react efficiently with any of the tested 2OG derivatives other than 2OG (24, 25).

The combined evidence indicates the potential of specific 2OG oxygenases to selectively react with and, in some cases, be inhibited by, 2OG derivatives (23).

We have proposed that 2OG derivatives may react differently with the same enzyme to give different products/product ratios to those observed with 2OG (23). To investigate this possibility, we envisioned that the 2OG-dependent ethylene/succinate-forming enzyme from *Pseudomonas savastanoi* (previously classified as: *Pseudomonas syringae*) pv. phaseolicola (PsEFE; EC:1.13.12.19) (35, 36) would be a particularly useful model, because with 2OG it catalyzes a bifurcating reaction (37), *i.e.*, (i) the oxidative decarboxylation of 2OG coupled to the oxidation of L-arginine (Arg) (38) to give succinate, CO_2, and 5-hydroxyarginine that fragments giving guanidine and L-Δ^1-pyrroline-5-carboxylate (P5C), and (ii) a Grob-type fragmentation of 2OG to give ethylene and three molecules of CO_2 (Figure 2.1b) (11, 37, 39-41). PsEFE thus has potential for the biocatalytic production of ethylene and other industrially important alkenes (35, 36, 42, 43).

Limited studies with PsEFE and 2OG derivatives have been reported (39, 44, 45). PsEFE can catalyze hydroxylation of Arg using 2OA as a cosubstrate, but it apparently does not catalyze the conversion of 2OA into propene (39). The reactivity of PsEFE with five C4 substituted 2OG derivatives was recently reported using infrared (IR) and LCMS assays, the results of which indicated that wild-type (wt) PsEFE does not catalyze the fragmentation of C4-alkyl- and/or C4-hydroxyl-substituted 2OG derivatives to give alkenes, though it did catalyze fragmentation of 4-fluoro-2OG into fluoroethylene (44). All of the five tested C4 substituted 2OG derivatives sustained the PsEFE-catalyzed hydroxylation of Arg to give the corresponding succinate derivatives; additionally, PsEFE catalyzed the decarboxylative β-oxidation of all of the five tested C4 substituted 2OG derivatives to give the corresponding β-hydroxypropionate products (44).

Here we report ^{1}H NMR studies investigating the reactivity of PsEFE with a set of 18 C3 and/or C4 substituted 2OG derivatives; the reaction of C3 substituted 2OG derivatives with PsEFE has not been previously tested. Oxidative decarboxylation of 2OG derivatives to give diacid (succinate) derivatives coupled to Arg oxidation was observed in some cases. Although no evidence for alkene formation was accrued, some of the 2OG derivatives reacted to give alcohol (β-hydroxypropionate) derivatives. Notably, the results provide clear evidence that the ratio of bifurcating reaction outcomes can be affected by the nature of the 2OG derivatization, supporting the proposal that the outcome of 2OG oxygenase catalysis can be altered in a 2-oxoacid (co)substrate dependent manner.

2.4 Results

2.4.1 NMR assays for monitoring PsEFE catalysis

^{1}H NMR assays were employed to investigate the reactivity of C3 and/or C4 substituted 2OG derivatives with PsEFE, because they can simultaneously: (i) monitor substrate depletion and product formation, (ii) enable quantification of substrate turnover by comparison with an internal standard, and (iii) enable the structural characterization of reaction products. Prior to investigating the reactivity of C3 and/or C4 substituted 2OG derivatives with PsEFE, we optimized the reported ^{1}H NMR assay conditions using 2OG as acosubstrate (40). The optimized standard conditions were: *Pseudomonas savastanoi* (previously: *Pseudomonas syringae*) pv. phaseolicola ethylene/succinate-forming enzyme (PsEFE; UNIPROT ID: P32021; EC:1.13.12.19) (40) (10 or 30 μM), L-arginine (Arg; 500 μM), L-ascorbate (LAA; 400 μM), 2OG/2OG derivative (400 μM), and Fe(II) (50 μM) in buffer (50 mM sodium phosphate, 10%$_{v/v}$$D_2O$, pH 7.4) at 25 °C (Supporting Figure 2.S1). Exchanging the previously reported Tris-d_{11} buffer (40) for phosphate buffer improved detection of P5C by ^{1}H NMR

(singlet, δ ~ 7.70 ppm). 3-(Trimethylsilyl)-2,2,3,3-tetradeuteropropionic acid (46) (TMSP-*d*$_4$; 800 μM) was used as an inert internal standard to enable quantification of turnover.

No reaction products other than succinate (singlet, δ ~ 2.32 ppm), ethylene (singlet, δ ~ 5.35 ppm), and P5C (5-H: singlet, δ ~7.70 ppm) were observed by ^{1}H NMR (600 MHz) analysis of the crude reaction mixture under the optimized conditions when using 2OG as a cosubstrate (Figure 2.1); note that all peaks are assigned in the PsEFE-catalyzed conversion of 2OG (Supporting Figure 2.S2). This result was confirmed by pure shift ^{1}H NMR (47) (700 MHz) analysis of the reaction mixture (Supporting Figure 2.S3). The pure shift method increases the sensitivity for detection of reaction products by broadband homonuclear decoupling that collapses proton signals with multiplet patterns into singlets at near the respective chemical shift (48-50). Arg depletion could not be quantified by ^{1}H NMR analysis, because the Arg signals overlapped with those of the P5C product. Signals corresponding to dehydroascorbate, arising from oxidation of LAA, were observed at low levels (Supporting Figure 2.S4).

The results of integrating the relevant peaks of ^{1}H NMR assays of up to >1 h duration after PsEFE addition imply that the formation of succinate and P5C are coupled, *i.e.*, they are produced in a ~1:1 ratio, at least in early time points (Figure 2.2). The P5C signal was, however, observed to decrease at later time points, likely due to relatively slow non-enzymatic reaction(s). The ^{1}H NMR-based data for ethylene production obtained by integration of the ethylene peak should be regarded as approximate, because reactions were performed in open 3mm NMR tubes to enable sufficient O_2 supply. Ethylene formation was thus indirectly estimated by ^{1}H NMR analysis of the crude reaction mixtures by subtracting the normalized integral for succinate (obtained via quantitative comparison with that for TMSP-*d*$_4$; T_1: 2 s) from that for 2OG. Analysis using this method showed that succinate production correlates stoichiometrically with that of Arg hydroxylation to give P5C (Supporting Figure 2.S5). This method was found to be more reliable than direct integration of the ethylene signal; note that

quantitative comparison of the 2OG integrals with those of TMSP-d_4 confirms the 1:2 stoichiometry of 2OG and TMSP-d_4 in the initial reaction mixture. Using this method of analysis, the ratio of ethylene:succinate formation was estimated to be 1.0:1.0 ± 0.1 (n = 3; mean ± standard deviation, SD), which differs from reported ethylene:succinate ratios obtained by scintillation or gas chromatography (GC) methods, *i.e.* ~2:1 (37, 39, 41). The observed difference in the ethylene:succinate ratios may be the result of the different assay employed and/or the different reaction conditions, as recently proposed (11).




Figure 2.2. Reaction conditions affect the product ratio of PsEFE catalysis when using 2OG as a (co)substrate. The ethylene:succinate ratios estimated by ^{1}H NMR (600 MHz) analysis of the crude reaction mixtures vary with pH: (**a**) 1.5 ± 0.1:1.0 at pH 6.5, (**b**) 1.0 ± 0.1:1.0 at pH 7.4, and (**c**) 0.5 ± 0.1:1.0 at pH 8.5. Ethylene formation was indirectly estimated by subtracting the normalized integral of succinate (obtained via comparison with TMSP-d_4) from that of 2OG. The time scales were calibrated to the end of the acquisition of the first ^{1}H NMR experiment following PsEFE addition to the reaction mixture (t = 0 min), by which time low levels of conversion were manifest. Conditions: 400 µM 2OG, 500 µM L-arginine, 500

μM L-ascorbate, 50 μM Fe(II), 800 μM TMSP-*d*$_4$, and 2 μM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C; note that full conversion was observed after ~12 h or when increasing PsEFE concentrations. Reaction profiles are a mean of three independent runs (n = 3; mean ± SD).

The possibility that the ratio of ethylene:succinate formation varies with the reaction conditions was tested by monitoring PsEFE catalysis at different pH values, using 2OG as a cosubstrate. The results reveal that the estimated ethylene:succinate ratios vary with the pH, *i.e.*, the ethylene:succinate ratio apparently decreases from ~1.5:1.0 at pH 6.5 to ~1:1 at pH 7.4, then to ~0.5:1.0 at pH 8.5 (Figure 2.2). Note that the product ratios appeared to be constant over the time course of analyses at a given pH. Thus, it appears that more acidic conditions (pH 6.5) favor ethylene over succinate formation, although other factors than the pH may affect the ethylene:succinate ratios considering that they are reported to be constant ~2:1 between pH 6.0 and 8.5 when using different conditions and assay methods (39). Increased levels of turnover were also observed under more acidic conditions (pH 6.5). Although the underlying reason(s) for the observed changes in the ethylene:succinate ratios at different pHs are presently unclear, these observations may be of interest from a biocatalytic perspective for optimizing conversion of 2OG to ethylene rather than succinate.

2.4.2 PsEFE catalyzes conversion of 2-oxoadipate to γ-hydroxybutyrate

We then reinvestigated the reaction of PsEFE with 2-oxoadipate (2OA, **2**), which binds in a similar manner as 2OG (PDB ID: 5V2Z (51)) and which is a reported PsEFE cosubstrate (39). Comparison of reaction profiles over >1 h reveals ~60% consumption of 2OA under the standard conditions (Figures 2.3 and 2.4b; Supporting Figures 2.S6-2.S8). Neither increasing the PsEFE concentration (from 10 to 30 μM), nor increasing the reaction time to ~20 h improved overall levels of 2OA consumption, suggesting possible product inhibition, though assays in the presence of added glutarate (**2a**), γ-hydroxybutyrate (**2b**) or guanidine (400 μM)

suggested that these do not affect PsEFE catalysis. As reported, our 2OA results reveal production of glutarate(39), the formation of which, as for 2OG (38), is tightly coupled to that of P5C (Figure 2.4b). No evidence for propene formation was observed, including by pure shift ^{1}H NMR analysis (47-50) (Figure 2.3b), even after prolonged reaction (>20 h), in accord with the literature (39). Note that for the pure shift ^{1}H NMR analysis, the observed peak may not necessarily be at the center of the peak obtained when using standard NMR analysis (47-50). Time-dependent formation of γ-hydroxybutyrate (**2b**) was observed (Figures 2.3 and 2.4b), in support of a report appearing during the course of our investigations (44). Formation of β-hydroxypropionate, the corresponding putative product arising from decarboxylative β-oxidation of 2OG, was not observed by ^{1}H NMR with 2OG as a cosubstrate (Supporting Figures 2.S2 and 2.S3), in accord with previous results (39, 44).

The results with 2OA show that the γ-hydroxybutyrate (**2b**):glutarate (**2a**, produced from 2OA) and ethylene (**1b**):succinate (**1a**, produced from 2OG) ratios differ. The ratio of γ-hydroxybutyrate:glutarate formation from 2OA (*i.e.*, 1.0:2.1 ± 0.1; n = 3 ± SD) differs from that estimated for ethylene:succinate formation from 2OG (*i.e.,* 1:0:1.0 ± 0.1). Note that this relationship is not affected by loss of ethylene from solutions and that the γ-hydroxybutyrate:glutarate ratio with 2OA measured by ^{1}H NMR differs from that reported using an LCMS based assay (*i.e.*, ~1:1) (44), possibly reflecting different assay conditions, as previously observed when using 2OG as a (co)substrate (Figure 2.2). The observations with 2OA encouraged us to carry out studies on the reactivity of PsEFE with synthetic 2OG derivatives, with the objective of exploring the possibility of alterations in product ratios.




Figure 2.3. PsEFE catalyzes the bifurcating reaction of 2-oxoadipate to give glutarate and γ-hydroxybutyrate. (**a**) PsEFE catalyzes the bifurcating decarboxylative oxidation of 2-oxoadipate (2OA, **2**) to give either (i) glutarate (**2a**) or (ii) γ-hydroxybutyrate (**2b**); propene was not observed; (**b**) analysis (1.4 to 3.6 ppm) of the crude reaction mixture by pure shift ^{1}H NMR (700 MHz) (47-50) 16 h post incubation with PsEFE indicates that PsEFE does not catalyze propene formation from 2OA. Note that for the pure shift ^{1}H NMR analysis, the observed peak may not necessarily be at the center of the peak obtained when using standard NMR analysis (47-50); (**c**-**h**) ^{1}H NMR (600 MHz) analysis (1.4 to 3.7 ppm) of (**c**) a reaction mixture of PsEFE and 2OA spiked with an authentic sample of **2b** (standard) 16 h post addition of PsEFE, (**d**) **2b** (standard) in the absence of 2OA under standard conditions; (**e**) a reaction mixture of PsEFE and 2OA spiked with an authentic sample of **2a** (standard) 16 h post addition of PsEFE, (**f**) **2a** (standard) in the absence of 2OA under standard conditions; (**g**) a reaction mixture of PsEFE and 2OA 16 h post addition of PsEFE; (**h**) 2OA under standard conditions in the absence of PsEFE. Conditions: 400 μM 2OA, 500 μM L-arginine, 500 μM L-ascorbate, 50 μM Fe(II), 800 μM TMSP-d_4, 10 μM PsEFE, and, if appropriate, 2 mM of an appropriate authentic standard in buffer (50 mM phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Representative ^{1}H NMR spectra from three independent repeats are shown (n = 3).




Figure 2.4. ^{1}H NMR assays enable measurement of product ratios of PsEFE-catalyzed reactions with 2OG derivatives. Reaction profiles of the PsEFE-catalyzed oxidation of (**a**) 2OG (**1**), (**b**) 2OA (**2**), (**c**) 4-methyl-2OG (**3**), (**d**) 4-hydroxy-2OG (**4**), (**e**) 4-hydroxy-4-methyl-2OG (**5**), (**f**) 4,4-dimethyl-2OG (**6**), (**g**) 4-ethyl-2OG (**7**), (**h**) 4-propyl-2OG (**8**), (**i**) 4-butyl-2OG (**9**), and (**j**) 3-methyl-2OG (**10**). Conditions: 400 μM 2-oxoacid, 500 μM L-arginine, 500 μM L-ascorbate, 50 μM Fe(II), 800 μM TMSP-d_4, 2 μM PsEFE (for 2OG) or 10-30 μM PsEFE (for 2OA and 2OG derivatives) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O) at 25 °C. 2OG derivatives were employed as racemic mixtures. The time scales were calibrated to the end of the first ^{1}H NMR experiment following PsEFE addition to the reaction mixture (t

= 0 min), by which time low levels of conversion were manifest. Results are the mean of independent triplicates (n = 3; mean ± SD); graphs were plotted using GraphPad Prism 9.3.0 using non-linear regression analysis and a single-phase decay model. Representative ^{1}H NMR spectra are shown in Supporting Figures 2.S9-2.S42.

2.4.3 The 2OG substitution pattern affects the reaction outcomes of PsEFE catalysis

The effects of 18 C3 and/or C4 substituted 2OG derivatives (22) on PsEFE catalysis were investigated in the absence of 2OG; note that the chiral 2OG derivatives were used as racemates.Analysis of the reaction mixtures by ^{1}H NMR reveals that PsEFE does not catalyze the conversion of any of the tested 2OG derivatives to alkenes, at least within our limits of detection using ^{1}H NMR (600 MHz) (Figure 2.5; Supporting Figures 2.S9-2.S42). This observation is consistent with results reported during the course of our work, showing that, although PsEFE employed 4-fluoro-2OG as a substrate to give fluoroethylene, it did not catalyze alkene formation from 2OG derivatives bearing alkyl or hydroxy substituents at the C4 position; note that alkene/diacid and alcohol/diacid product ratios were not reported, except for 4-hydroxy-2OG (**4**) (44). However, using IR spectroscopy-based assays, it was recently reported that PsEFE catalyzes the formation of low levels of propylene from 4-methyl-2OG (**3**) (45). We did not accrue evidence for propylene formation from **3** or **10** by NMR under the tested conditions, even when using pure shift ^{1}H NMR (48-50) (Figure 2.6 and Supporting Figure 2.S42), which increases sensitivity for propene detection (47). The difference is likely a consequence of the different reaction conditions employed and/or the intrinsic alkene detection limitations of our ^{1}H NMR-based PsEFE assays.




Figure 2.5. The substitution pattern of C3 and/or C4 substituted 2OG derivatives affects reaction outcomes of PsEFE catalysis. (**a**) Reaction outcomes of PsEFE (10 μM; for 2OG: 2 μM; for **9**: 30 μM) catalysis with C4 substituted 2OG derivatives (400 μM). (**b**) Reaction outcomes of PsEFE (10 μM; for 2OG: 2 μM) catalysis with C3 substituted 2OG derivatives (400 μM). (**c**) 2OG derivatives which sustain PsEFE-catalyzed Arg hydroxylation as cosubstrates, but which do not give alcohol products. In each case, the identity of the succinate derivative product was confirmed by spiking the reaction mixture with an authentic synthetic sample. However, levels of conversion could not be accurately determined using ^{1}H NMR analyses (600 MHz) due to overlapping signals of starting material and product. Spiking with the potential corresponding alcohol products revealed that they are not formed, within limits of detection (Supporting Figures 2.S43-2.S48). Evidence for the corresponding alcohol product was also not observed by pure shift ^{1}H NMR (Supporting Figure 2.S48). (**d**) 2OG derivatives which do not react with PsEFE (30 μM) under the tested conditions (Supporting Figures 2.S49-2.S56). 2OG derivatives were typically prepared from cyanosulfur ylids as racemic mixtures as reported (22); **14** was used as a mixture of diastereomers, dr (*trans*:*cis*) ~ 5:1. Note it is possible that stereoselective conversion(s) may occur in some cases. Results are the mean of independent triplicates (n = 3; mean ± SD).




Figure 2.6. Analysis of the reaction outcome of PsEFE-catalyzed conversion of 4-methyl-2OG using pure shift ^{1}H NMR. (**a**) PsEFE catalyzes the oxidation of 4-methyl-2OG (**3**) to give either (i) 2-methylsuccinate (**3a**) or (ii) β-hydroxyisobutyrate (**3b**); propene was not observed. Analysis (0.9 to 3.5 ppm) of the reaction of PsEFE with **3** 30 min post addition of PsEFE using (**b**) pure shift ^{1}H NMR (700 MHz) (47-50) and (**c**) ^{1}H NMR (600 MHz) indicates that reaction products other than **3a**,**3b**, and P5C are not being formed in substantial amounts under the tested conditions (Supporting Figures 2.S9-2.S12). Analysis of the reaction mixture with **3** in the absence of PsEFE using (**d**) pure shift ^{1}H NMR (700 MHz) (47-50) and (**e**) ^{1}H NMR (600 MHz). **3** was used as a racemic mixture; it is possible that stereoselective conversion may occur. Note that the signals observed between ~1.1 to ~1.3 ppm in (**c**) originate from the PsEFE sample. Note that for the pure shift ^{1}H NMR analysis, the observed peak may not necessarily be at the center of the peak obtained when using standard NMR analysis (47-50). Conditions: 400 µM **3**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).

Although PsEFE did not catalyze the formation of alkenes from the tested C3 and/or C4 substituted 2OG derivatives, ^{1}H NMR analyses of the reaction profiles >1 h post addition of PsEFE reveal that 10 of the 18 tested 2OG derivativeswere (co)substrates of PsEFE, *i.e.*, 4-methyl-2OG (**3**), 4-hydroxy-2OG (**4**), 4-hydroxy-4-methyl-2OG (**5**), 4,4-dimethyl-2OG (**6**), 4-ethyl-2OG (**7**), 4-propyl-2OG (**8**), 4-butyl-2OG (**9**), 4-benzyl-2OG (**15**), 3-methyl-2OG (**10**), and the carbocycle **14** (Figure 2.5; Supporting Figures 2.S9-2.S42). Interestingly, whilst PsEFE catalyzed diacid formation from the 2OG derivatives **3**-**10**, **14**, and **15** in a manner apparently coupled to Arg (substrate) oxidation, as evidenced by P5C formation and by comparison with authentic samples of the corresponding diacid products, it only catalyzed alcohol formation from a subset of the identified substrates, *i.e.*, **3**-**10**, which thus mimic the bifurcating reactivity of 2OA (**2**) with PsEFE (Figures 2.4 and 2.5; Supporting Figures 2.S9-2.S42).

Although evidence for the PsEFE-catalyzed formation of alcohol products from 4-butyl-2OG (**9**), 4-benzyl-2OG (**15**), and from the carbocycle **14** was not accrued under the tested conditions (Figures 2.5 and 2.7), the levels of their PsEFE-catalyzed conversion to the corresponding diacid products were relatively low and could not be quantified for **14** and **15**, because of signals overlapping in ^{1}H NMR which hampered accurate integration; diacid formation was only supported by analysis of P5C formation and by the addition of the corresponding authentic synthetic diacid sample to the reaction mixture (Figure 2.7; Supporting Figures 2.S42-2.S49). This combination of factors may perturb data analysis. Formation of an apparent single diacid product for **9**, **14**¸and **15** was supported using pure shift ^{1}H NMR (700 MHz) analyses, which also revealed no evidence for formation of alcohol products (Figure 2.7; Supporting Figures 2.S37 and 2.S48).

All the chiral 2OG derivatives tested were racemic; it is possible that the absolute/relative configurations of the chiral derivatives, including of the carbocycle **14**, may affect the efficiency and outcome of their reaction with PsEFE. In this regard, it is of interest that PsEFE

catalyzed conversion of **14**, but not of those (bi)cyclic 2OG derivatives which have a more restricted conformational flexibility, *i.e.*, the 3- and 4-membered carbocycles **17** and **18**, as well as **19** and **20** which bear an aromatic or bicyclic carbon scaffold, respectively (Figure 2.5). Likely, **17**-**20** cannot occupy a conformation which results in productive PsEFE catalysis when bound to the active site or they do not bind to the active site at all. The combined observations indicate that the substitution pattern of 2OG derivatives can alter the reaction outcome in their bifurcating reaction with PsEFE.




Figure 2.7. PsEFE employs carbocycle 14 as a cosubstrate to sustain Arg oxidation. (**a**) PsEFE catalyzes Arg oxidation using carbocycle **14** as a cosubstrate, but does not catalyze the competing decarboxylative β-oxidation of **14** to give **14b**; (**b**) analysis (0.8 to 3.0 ppm) of the crude reaction mixture by pure shift ^{1}H NMR (700 MHz) (47-50) 12 h post incubation with PsEFE indicates that PsEFE does not catalyze alcohol (**14b**) formation from **14**; (**c**-**g**) ^{1}H NMR (600 MHz) analysis (0.8 to 3.0 ppm) of (**c**) 4-hyxroxycyclohexanecarboxylate (**14b**, standard) in buffer, (**d**) a reaction mixture of PsEFE and **14** spiked with cyclohexane-1,4-dicarboxylate (**14a**, standard) 12 h post addition of PsEFE, (**e**) **14a** (standard) under standard conditions in the presence of PsEFE; (**f**) a reaction mixture of PsEFE and **14** 12 h post addition of PsEFE; (**g**) **14** under standard conditions in the absence of PsEFE. Conditions: 400 μM **14**, 500 μM L-arginine, 500 μM L-ascorbate, 50 μM Fe(II), 800 μM TMSP-d_4, 30 μM PsEFE, and, if appropriate, 2mM of an appropriate authentic standard (50 mM phosphate, pH 7.4, 10%$_{v/v}$ D_2O). **14** was used as a mixture of diastereomers, dr (*trans*:*cis*) ~ 5:1; note it is possible that stereoselective conversion may occur. Note that for the pure shift ^{1}H NMR analysis, the observed peak may not necessarily be at the center of the peak obtained when using standard NMR analysis (47-50). A representative set of ^{1}H NMR spectra from three independent repeats is shown (n = 3).

To investigate the effect of the steric bulk of the 2OG C4 substituent on the outcome of the bifurcating PsEFE reaction with 2OG derivatives, we analyzed the product ratios of diacid versus alcohol formation for the C4 alkyl-substituted 2OG derivatives(Figure 2.5a; Supporting Table S1). The results reveal that substituting the methyl group of 4-methyl-2OG (**3**) for an ethyl group results in a ~2-fold increase in the PsEFE-catalyzed formation of the corresponding succinate derivative over the β-hydroxypropionate derivative from ~1.8:1.0 to ~3.2:1.0. Substituting the methyl group of **3** for a propyl group results in a ~3-fold increase of the corresponding succinate derivative (~4.1:1.0), while its substitution for bulkier butyl or benzyl groups completely suppresses formation of the potential alcohol product (Figure 2.5a).

The comparison of the product ratio obtained for 4-methyl-2OG (**3**) with those obtained for 4,4-dimethyl-2OG (**6**) and 3-methyl-2OG (**10**) indicates that both the addition of a second methyl substituent at the C4 position and moving the methyl substituent to the C3 position favor formation of the alcohol product (Figure 2.5; Supporting Table 2.S1). 3-Ethyl-2OG (**11**), 3-propyl-2OG (**12**), 3-butyl-2OG (**13**), and 3-benzyl-2OG (**16**) do not react with PsEFE under the tested conditions (even after prolonged incubation times, ~20 h), by contrast with their corresponding C4 regioisomers **7**-**9** and **15** (Figure 2.5). These combined results show that both the steric bulk and the position of the 2OG substituents affects the product ratio in PsEFE catalysis. Note that, by contrast with the observed pH dependence of the product ratio when using 2OG as a (co)substrate (Figure 2.2), altering the pH from 6.5 to 8.5 did not affect the product ratio of PsEFE catalysis using 2OG derivatives **3** and **10** (Supporting Figures 2.S57-2.S58).

In addition to analyzing the product ratios, we analyzed the efficiencies of product formation and the overall levels of conversion at ~1.5 h post addition of PsEFE for the PsEFE-catalyzed reaction of 4-methyl-2OG (**3**), 4-ethyl-2OG (**7**), 4-propyl-2OG (**8**), and 4-butyl-2OG (**9**) (Figure 2.4, Supporting Table 2.S1). The results reveal that the specific activities of the 2OG

derivatives with PsEFE (*i.e.*, the efficiency of catalysis) decreases with increasing steric bulk of the C4 substituent: substrate conversion was >95% for 2OG, **3**, and **7**, but decreased to ~70% for **8**, and to ~50% for **9**. The reduced efficiency of conversion correlates with the observed trend in the product ratios, which favors diacid formation coupled to Arg oxidation for sterically bulky C4 substituents (Figure 2.5, Supporting Table 2.S1). The levels of conversion of **8** and **9** did not increase after >20 h incubation, an observation which may indicate (at least for **9**) that the absolute configuration of the C4 stereocenter affects the efficiency and/or the outcome of the reaction of 2OG derivatives with PsEFE, because the substrates used were racemic.

The combined evidence implies that the steric bulk of the 2OG substituents not only affects the product ratio, but also the overall PsEFE efficiency, potentially due to less efficient binding at the active site and/or their reduced conformational flexibility. This observation implies that active site mutagenesis may alter the reactivity of PsEFE active site variants with 2OG derivatives, as precedented by studies on the reactivity of C4 substituted 2OG derivatives with active site variants of the JmjC lysine-specific N^{ε}-demethylases KDM4A (25); a proposal validated by a recent proof-of-concept study on the reactivity of the L206V PsEFE active site variant which catalyzed propene formation from (4*R*)-4-methyl-2OG more efficiently than wt PsEFE (45).

We next compared the diacid:alcohol ratios obtained for 4-methyl-2OG (**3**), 4-hydroxy-2OG (**4**), 4-hydroxy-4-methyl-2OG (**5**), and 4,4-dimethyl-2OG (**6**) to investigate whether, in addition to the steric bulk of the 2OG C4 substituent, the nature of the substituent affects product ratios. The results imply that substituting the C4 methyl group of **3** with a hydroxy group does not substantially affect either the diacid:alcohol ratio or the extent of conversion after 1 h incubations (Figures 2.4 and 2.5). Note that the observed diacid:alcohol ratio of **4** was ~2:1 in our ^{1}H NMR assay (Supporting Table S1), whereas it was ~3:1 in reported LC-MS

assays (44). Similarly, substituting one of the two methyl groups of **6** for a hydroxy group did not affect the product ratio during PsEFE catalysis, within the limits of detection (Figure 2.4), but reduced the overall levels of conversion from ~80% to ~30% over the analyzed time period (Figure 2.4, Supporting Table 2.S1).

2.5 Discussion

The *P. savastanoi* (previously: *P. syringae*) pv. phaseolicola ethylene-forming enzyme (PsEFE) (35, 36) is of interest from both biocatalytic and mechanistic perspectives. When using 2OG (**1**) as a (co)substrate, its reaction bifurcates resulting in two types of oxidative decarboxylation reactions, *i.e.* (i) a Grob-type fragmentation of 2OG to give ethylene and three molecules of CO_2, and (ii) the oxidation of 2OG to give succinate (and CO_2) coupled to Arg oxidation giving guanidine and P5C (Figure 2.1) (37, 38, 40, 41). It is also reported that PsEFE can catalyze the oxidation of some C4 substituted 2OG derivatives to give the corresponding β-hydroxypropionate derivatives (Figure 2.5) (44). It is desirable to understand the factors which govern the product ratio for the (co)substrates of PsEFE, *inter alia*: (i) to improve PsEFE catalysis for biocatalytic applications, including ethylene production from renewable resources, (ii) to predict the reactivity of other 2OG oxygenases with 2OG derivatives, and (iii) to inform on the, likely context-dependent, biological roles of PsEFE.

The biological functions of ethylene in plants are, at least in part, understood, whereas the reasons why plant pathogens, including *P. savastanoi*, synthesize ethylene are less well characterized (52). Bacteria may have evolved to produce ethylene to interact with plant ethylene receptors in an ACC-independent manner (53, 54), *e.g.*, to perturb the plant metabolism and/or immune response. *P. savastanoi* employs 2OG rather than ACC as an ethylene precursor; although PsEFE and ACCO are structurally related, their mechanisms are different (Figure 2.1) (11). To our knowledge, no 2OG oxygenase other than PsEFE (and

analogous EFEs in related bacteria) is reported to produce ethylene from 2OG, including the bacterial 2OG oxygenases VioC and OrfP which also catalyze hydroxylation of isolated arginine, however, unlike PsEFE, at the arginine C3 position (55, 56).

PsEFE catalysis has potential to directly and indirectly regulate cellular levels of 2-oxoacids and associated metabolites, including α-amino acids. The PsEFE-catalyzed C5 hydroxylation of Arg to give P5C provides a further link to 2-oxoacid/amino acid metabolism, because P5C, which is in equilibrium with glutamate-5-semialdehyde, is metabolically linked to glutamate, proline, and ornithine (57). P5C is a substrate of various bacterial and plant enzymes, though above threshold levels is reported to be cytotoxic (58-62). The physiological roles of guanidine formation following arginine C5 hydroxylation are unclear, but recent evidence suggests that guanidine is a plant metabolite indicating that PsEFE may perturb plant guanidine levels (63). It has also been shown that specific 2OG-dependent oxygenases in plants and algae catalyze C6 hydroxylation of homoarginine, resulting in spontaneous guanidine formation; however, these enzymes do not catalyze ethylene formation. Interestingly, despite the homoarginine hydroxylase and PsEFE catalyzing closely related reactions, their sequences show only ~25% similarity (63).

Our NMR studies on the (co)substrate scope of PsEFE validate prior studies using different methods showing that the lysine metabolite 2OA is an efficient substrate of PsEFE in the absence of 2OG (39, 44). *Pseudomonal* 2OG oxygenases other than PsEFE are also reported to employ 2OA as a cosubstrate, however, with variable efficiencies compared to 2OG. For example, catalysis of the *P. putida* S-313 oxygenativealkylsulfatase with 2OA as a cosubstrate was ~20% less efficient than with 2OG (17), catalysis of the *P. stutzeri* WM88 hypophosphite:2-oxoglutarate dioxygenase (HtxA) with 2OA was ~70% less efficient than with 2OG (18), whereas catalysis of *P. aeruginosa* γ-butyrobetaine hydroxylase (BBOX) with 2OA did not proceed at all (64). The combined observations thus suggest that increased 2OA

levels may affect catalysis of *Pseudomonal* 2OG oxygenases in an, at least, partly selective manner. It is possible that 2OA or, by implication, other 2OG derivatives may be the preferred (co)substrate of some 2OG oxygenases or that competition between 2OG and 2OA regulates catalysis in cells (22, 23).

Our observations that PsEFE accepts C3 and/or C4 substituted 2OG derivatives as (co)substrates in addition to 2OA, complements recently reported studies on PsEFE substrate scope using five C4 substituted 2OG derivatives (Figure 2.5) (44). Notably, both increasing the steric bulk of the 2OG C4 substituent and using conformationally rigid carbocyclic 2OG derivatives apparently shifts the product ratio of PsEFE catalysis in favor of or completely to Arg hydroxylation/diacid formation over β-hydroxypropionate derivative/alcohol formation (Figures 2.5 and 2.8).

The bifurcating reaction of PsEFE with 2OG derivatives as prime substrates to give variable product ratios is precedented by studies on the mechanistically distinct iron-dependent cytochrome P450 peroxidase $OleT_{JE}$ (CYP152L1) that catalyzes formation of both alcohol and terminal alkene products via oxidation of fatty acids (65, 66). Notably, as proposed for PsEFE (44), the bifurcating reactivity of $OleT_{JE}$ is reported to be a result of substrate-centered radical intermediates (67-70), the reaction fate of which can differ between reaction with an iron-bound/carbonate oxygen atom to give an alcohol and β-decarboxylation to give an alkene. The observed variations in product ratios may thus reflect different stabilities of radical intermediates and/or restricted conformations of intermediates favoring particular reaction outcomes.




Figure 2.8. 2OG derivatization influences the outcome of PsEFE catalysis. (**a**) C3 and/or C4 substituted 2OG derivatives can be (co)substrates of PsEFE in a bifurcating reaction: (i) to catalyze Arg hydroxylation resulting in formation of P5C and the corresponding succinate derivative and/or (ii) to give a β-hydroxypropionate derivative/alcohol. None of the tested 2OG derivatives manifested alkene formation, but differences in the ratios of P5C/succinate derivative formation versus β-hydroxypropionate derivative formation were observed. Increasing the steric bulk of the C3 substituent (R_1) beyond a methyl group apparently ablates PsEFE catalysis. Increasing the steric bulk of the C4 substituent (R_2) shifts the product ratio to favor P5C/succinate derivative formation over β-hydroxypropionate derivative formation, and reduces overall levels of catalysis. Cyclic 2OG derivatives (*i.e.*, linking R_1 and R_2) either were inactive (**17**-**20**) or only catalyzed P5C/diacid formation (**14**); (**b**) proposed outline mechanism of PsEFE (11, 44).

It is of interest that several of the alternative PsEFE substrates identified here are natural products, *i.e.*, 4-methyl-2OG (**3**) (71-74), 4-hydroxy-2OG (**4**) (75, 76), 4-hydroxy-4-methyl-2OG (**5**) (77-79), and 3-methyl-2OG (**10**) (80), some of which have been detected in *Pseudomonas* species (81, 82), implying that the primary or sole biological role of PsEFE catalysis may not be ethylene production. PsEFE catalyzes the conversion of the natural product 3-methyl-2OG (**10**) (80) to give β-hydroxybutyrate, which is a reported metabolite of *Pseudomonas* species (83) and which is a building block in the biosynthesis of the biodegradable polyester poly(3-hydroxybutyrate) (84). It is possible that the reaction of PsEFE

with 2-oxoacids other than 2OG could help *P. savastanoi* sense and/or adapt to internal and/or external stimuli, possibly in a context-dependent manner based on (co)substrate availability. There is precedent for roles of 2OG oxygenases and ACCO to act in a sensing capacity, in particular in terms of regulating the response to hypoxia in animals and plants, respectively (1, 10, 85).

The apparent (co)substrate promiscuity of PsEFE is precedented *e.g.*, by work on (i) the cosubstrate scope of human 2OG oxygenases which revealed that C3 and/or C4 substituted 2OG derivatives can selectively sustain and/or inhibit catalysis by the protein hydroxylases AspH, FIH, and JMJD5 (22-24), (ii) plant ACCOs, which employ a range of ACC derivatives to give products other than ethylene (86-88), and (iii) transaminases and oxidoreductases, which mechanistically and structurally distinct from 2OG oxygenases but some of which employ 2OG as a substrate (89-91). The apparent ability of PsEFE to accept 2OG derivatives with a range of C3 and/or C4 substituents as (co)substrates (Figure 2.5), contrasts with its reported relatively narrow scope of accepting 'prime' substrates other than Arg (39, 40, 92). These observations indicate that the fragmentation of Arg itself may be of biological importance (see above). In apparent contrast with PsEFE, both human AspH and FIH not only accept a diverse set of 2OG derivatives as cosubstrate (22, 23), but also catalyze oxidation of multiple protein substrates and catalyze different types of oxidation (28, 30, 34, 93, 94).

The biocatalytic production of ethylene from renewable resources is desirable from a sustainability perspective (42, 43). However, the potential of PsEFE for biocatalytic ethylene production on an industrial scale is in part currently compromised by the bifurcating reaction of 2OG, as ethylene formation completes with succinate formation, a process coupled to arginine hydroxylation (Figure 2.1) (95). Our results showing that the product ratio of ethylene versus succinate formation is affected by the reaction conditions are thus of interest, in particular the observation that more acidic conditions (*i.e.*, pH 6.5) favor ethylene over

succinate production (Figure 2.2). The reasons for the observed dependance of the ethylene:succinate ratio on the pH are currently unclear and may relate to different protonation states of residues proximate the 2OG binding site. Refining the residues involved in 2OG and/or Arg binding may help in research aimed at engineering PsEFE in a manner to enhance ethylene production (51).

The observation that PsEFE catalyzes formation of alcohol products from some of the tested C3 and, in particular, C4 substituted 2OG derivatives highlights the potential of PsEFE biocatalytic applications beyond ethylene formation. For example, PsEFE catalyzes the formation of β-hydroxybutyrate from the natural product 3-methyl-2OG (**10**) (80), affording a useful building block for the industrial synthesis of biodegradable polymers (84). The PsEFE-catalyzed decarboxylative β-oxidation of 2OG derivatives to give alcohol products complements prior work on the utility of 2OG derivatives for the biocatalytic synthesis of alcohols, which revealed that oxidoreductases can catalyze the reduction of 2OG derivatives to give the corresponding hydroxyglutarate derivatives (91).

2.6 Experimental Procedures

Chemicals. 2-Oxoglutarate (2OG), DL-4-hydroxy-2-oxoglutarate (4-OH-2OG, **4**), and 2-oxoadipate (2OA, **2**) were commercially-sourced (Sigma Aldrich Inc.), all the other tested C3 and/or C4-substituted 2OG derivatives were synthesized as racemic mixtures as reported (22). All other chemicals used for this work were commercially-sourced (Sigma Aldrich Inc., Fluorochem Ltd.) and used without further purification.

PsEFE production and purification. Isolated recombinant PsEFE was produced and purified as reported (40). A pETite N-His SUMO Kan plasmid encoding for *Pseudomonas savastanoi* (previously: *Pseudomonas syringae*) pv. phaseolicola ethylene/succinate-forming enzyme (PsEFE; EC:1.13.12.19; UNIPROT ID: P32021) (40), codon optimized for expression in *E.*

coli, was transformed into competent T7 Express LysY competent *E. coli* cells (new England Biolabs). Cells were grown in 2TY media supplemented with kanamycin (30 μg/mL) at 37 °C with agitation (180 rpm), until PsEFE expression was induced at an OD_{600} ~0.8 by addition of isopropyl-β-D-thiogalactopyranoside (final concentration: 0.5 mM). Cells were incubated at 18 °C for 16 h, before being harvested by centrifugation (11,395 x g, 15 min, 4 °C); the resultant cell pellet was stored at –80 °C.

The frozen cells were resuspended in ice-cold buffer (50 mM Tris, pH 7.8, 500 mM NaCl, 20 mM imidazole) supplemented with DNase I (bovine pancreas, Sigma-Aldrich) and lysed at 4 °C using a high-pressure cell disrupter (Continuous Flow Cell Disruptor CF1; Constant Systems Ltd) followed by centrifugation (58545 x g, 30 min, 4 °C). The supernatant was purified at 4 °C using Ni(II)-affinity chromatography (HisTrap HP column, GE Healthcare) using an ÄKTA Pure chromatography system(GE Healthcare) employing an elution gradient with increasing buffer (50 mM Tris, pH 7.8, 500 mM NaCl) imidazole concentration (from 20 mM to 500 mM imidazole). Fractions containing PsEFE were combined, concentrated using Amicon Ultra centrifugal filters (3082 x g, 4 °C), and buffer exchanged using a PD-10 desalting column (Cytiva) to remove imidazole (25 mM Tris, pH 7.8, 500 mM NaCl, 10%$_{v/v}$ glycerol). The N-terminal His_6-SUMO tag was cleaved overnight using sentrin-specific protease 2 (SenP2; Supporting Figure 2.S59) at 4 °C; and the resultant mixture was purified using Ni(II)-affinity chromatography (50 mM Tris, pH 7.8, 500 mM NaCl, 20 mM imidazole). The flow-through was collected, concentrated, and purified using size-exclusion chromatography (300 mL HiLoad 26/60 Superdex 75 pg column; ÄKTA Pure chromatography system; 25 mM Tris, pH 7.8, 100 mM NaCl; 1 mL/min). Eluted fractions containing >95% pure PsEFE (determined by SDS-PAGE analysis) were pooled, then concentrated using Amicon Ultra centrifugal filters (3082 x g, 4 °C) and aliquoted; PsEFE was and had the anticipated mass, *i.e.*, 39369.5 Da (calculated mass: 39370.5 Da). The aliquots were flash frozen in liquid N_2 and stored at –80 °C.

The PsEFE concentration was determined using a NanoDrop One microvolume UV-Vis spectrophotometer (Thermo Scientific), assuming that the concentration of active PsEFE equals the total PsEFE concentration (note that potent covalent or tight-binding PsEFE inhibitors suitable for active site titrations are not yet reported); fresh aliquots of PsEFE were used for all biochemical experiments.

^{1}H NMR assays. Aqueous stock solutions of L-arginine (Arg; 50 mM), 2OG or a 2OG derivatives (40 mM), L-ascorbate (LAA; 50 mM), and ferrous ammonium sulphate (FAS; 100 mM) were freshly prepared the day the experiment was performed. These stock solutions, together with a solution (100 mM) of (trimethylsilyl)propionic-2,2,3,3-*d*$_4$ acid sodium salt (TMSP-*d*$_4$) (46), which was used as an internal standard (–0.11 ppm), were diluted in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O, pH 7.4) to give Arg (5 mM), 2OG/2OG derivative (4 mM), LAA (4 mM), FAS (0.2 mM), and TMSP-*d*$_4$ (8 mM) solutions for use in the NMR assays. ^{1}H NMR (600 MHz) assays were performed using a Bruker AVIII 600 instrument with a Prodigy nitrogen broadband cryoprobe. Pure shift ^{1}H NMR (700 MHz) analyses were performed as reported (47), using a Bruker AVIII 700 MHz NMR spectrometer with a 5-mm inverse triple-resonance-inverse cryoprobe.

Assays were performed in a reaction volume of 180 µL with the following final concentrations: 500 µM L-arginine, 400 µM L-ascorbate, 400 µM co-substrate, 50 µM Fe(II), and 2, 10 or 30 µM PsEFE. The indicated stock solutions were strictly added in the following order into a 1.5 mL Eppendorf tube: aqueous buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O, pH 7.4; variable volume), then Arg (5 mM; 18 µL), then LAA (4 mM; 18 µL), then 2OG/2OG derivative (4 mM; 18 µL), then FAS (0.5 mM; 18 µL), then TMSP-*d*$_4$ (8 mM; 18 µL), then PsEFE (variable volume). The assay mixture was carefully mixed with a glass Pasteur pipette, transferred to a 3mm Bruker NMR tube, and hand centrifuged.

The NMR tubes were transferred as rapidly as possible to the spectrometer sample exchanger and inserted into the probe of a Bruker AVIII 600 instrument. The probe was locked to D_2O ($10\%_{v/v}$ in water), followed by auto-tuning ('atma' command in Bruker TopSpin 4.3.1). The sample was then shimmed (tshim), receiver gain was adjusted (rga), and consecutive acquisitions were started. The variable delay was set to zero and the number of scans per acquisition was set to 64 scans per acquisition to ensure a sufficient signal-to-noise ratio; the relaxation delay was minimized to reduce the time-gap between acquisitions. For ^1H experiments, water suppression was achieved using perfect echo-modified WATERGATE experiments (96). Digital resolution of the free induction delay (FID) was adjusted to 0.3 Hz for accuracy in integration. Peak integrals were normalized via comparison with that of TMSP-*d*$_4$. Assigned products were validated by spiking the reactions 12-24 h post PsEFE addition with 5 µL of a freshly prepared 40 mM stock solution (in: 50 mM sodium phosphate, $10\%_{v/v}$ D_2O, pH 7.4) of the relevant commercially-sourced authentic standards.

Spectra were phased and corrected automatically for peak symmetry and baseline correction. Integral precision was attenuated by zero filling which adjusted the size of the spectra to the number of data points used. Spectra were integrated using MestReNova x86_64 (Mestrelab research) and integrals calibrated and scaled to the internal standard TMSP-*d*$_4$. Data was plotted using GraphPad Prism 9. All time course data was obtained in independent triplicates.

2.7 Acknowledgements

This research was funded in part by the Wellcome Trust (106244/Z/14/Z). For the purpose of open access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission. We thank the Biotechnology and Biological

Sciences Research Council (BB/J003018/1 and BB/R000344/1) for funding. We thank Prof. T. D. W. Claridge, Dr. N. Amin, Dr. C. Mycroft, and Ms. N. K. Goodyear for helpful discussions.

2.8 Author Contributions

C. J. S., L. B., Z. Z., S. D. writing–review & editing; C. J. S., L. B., S. D. writing–original draft; C. J. S., L. B., Z. Z. supervision; C. J. S. funding acquisition; C. J. S., L. B. conceptualization. L. B., S. D. visualization, L. B., C. T. L., Z. Z., S. D. investigation; C. T. L., S. D. methodology; Z. Z. resources; S. D. validation.

2.9 References

1. *2-Oxoglutarate-dependent oxygenases*. (2015). Hausinger, R. P., and Schofield, C. J. (Eds.). Cambridge, United Kingdom: The Royal Society of Chemistry.
2. Mooney, B. P., Miernyk, J. A., and Randall, D. D. (2002) The complex fate of α-ketoacids. *Annu. Rev. Plant Biol.***53**, 357-375
3. Luo, Z., Yu, S., Zeng, W., and Zhou, J. (2021) Comparative analysis of the chemical and biochemical synthesis of keto acids. *Biotechnol. Adv.***47**, 107706
4. Cooper, A. J. L., Ginos, J. Z., and Meister, A. (1983) Synthesis and properties of the α-keto acids. *Chem. Rev.***83**, 321-358
5. Song, Y., Li, J., Shin, H.-d., Liu, L., Du, G., and Chen, J. (2016) Biotechnological production of alpha-keto acids: current status and perspectives. *Bioresour. Technol.***219**, 716-724
6. Pang, C. P., Chakravarti, B., Adlington, R. M., Ting, H. H., White, R. L., Jayatilake, G. S.*, et al.* (1984) Purification of isopenicillin N synthetase. *Biochem. J.***222**, 789-795
7. Rabe, P., Kamps, J. J. A. G., Schofield, C. J., and Lohans, C. T. (2018) Roles of 2-oxoglutarate oxygenases and isopenicillin N synthase in β-lactam biosynthesis. *Nat. Prod. Rep.***35**, 735-756
8. Houben, M., and Van de Poel, B. (2019) 1-Aminocyclopropane-1-carboxylic acid oxidase (ACO): the enzyme that makes the plant hormone ethylene. *Front. Plant Sci.***10**, 695
9. Bleecker, A. B., and Kende, H. (2000) Ethylene: a gaseous signal molecule in plants. *Annu. Rev. Cell Dev. Biol.***16**, 1-18
10. Binder, B. M. (2020) Ethylene signaling in plants. *J. Biol. Chem.***295**, 7710-7725
11. Hausinger, R. P., Rifayee, S. B. J. S., Thomas, M. G., Chatterjee, S., Hu, J., and Christov, C. Z. (2023) Biological formation of ethylene. *RSC Chem. Biol.***4**, 635-646
12. Nishizuka, Y., Ichiyama, A., and Hayaishi, O. (1970) [58] Metabolism of the benzene ring of tryptophan (mammals). *Methods Enzymol.***17**, 463-491
13. Danhauser, K., Sauer, S. W., Haack, T. B., Wieland, T., Staufner, C., Graf, E.*, et al.* (2012) DHTKD1 mutations cause 2-aminoadipic and 2-oxoadipic aciduria. *Am. J. Hum. Genet.***91**, 1082-1087
14. Mukherji, M., Kershaw, N. J., MacKinnon, C. H., Clifton, I. J., Wierzbicki, A. S., Schofield, C. J., and Lloyd, M. D. (2001) 'Chemical co-substrate rescue' of phytanoyl-CoA 2-hydroxylase mutants causing Refsum's Disease. *Chem. Commun.* 972-973

15. Cunliffe, C. J., Franklin, T. J., Hales, N. J., and Hill, G. B. (1992) Novel inhibitors of prolyl 4-hydroxylase. 3. Inhibition by the substrate analog N-oxaloglycine and its derivatives. *J. Med. Chem.***35**, 2652-2658
16. Majamaa, K., Hanauske-Abel, H. M., Günzler, V., and Kivirikko, K. I. (1984) The 2-oxoglutarate binding site of prolyl 4-hydroxylase. *Eur. J. Biochem.***138**, 239-245
17. Kahnert, A., and Kertesz, M. A. (2000) Characterization of a sulfur-regulated oxygenative alkylsulfatase from Pseudomonas putida S-313. *J. Biol. Chem.***275**, 31661-31667
18. White, A. K., and Metcalf, W. W. (2002) Isolation and biochemical characterization of hypophosphite/ 2-oxoglutarate dioxygenase: a novel phosphorus-oxidizing enzyme from Pseudomonas Stutzeri WM88. *J. Biol. Chem.***277**, 38262-38271
19. Eichhorn, E., van der Ploeg, J. R., Kertesz, M. A., and Leisinger, T. (1997) Characterization of α-ketoglutarate-dependent taurine dioxygenase from Escherichia coli. *J. Biol. Chem.***272**, 23031-23036
20. Fukumori, F., and Hausinger, R. P. (1993) Purification and characterization of 2,4-dichlorophenoxyacetate/alpha-ketoglutarate dioxygenase. *J. Biol. Chem.***268**, 24311-24317
21. Müller, T. A., Fleischmann, T., Meer, J. R. v. d., and Kohler, H.-P. E. (2006) Purification and characterization of two enantioselective α-ketoglutarate-dependent dioxygenases, RdpA and SdpA, from *Sphingomonas herbicidovorans* MH. *Appl. Environ. Microbiol.***72**, 4853-4861
22. Brewitz, L., Nakashima, Y., and Schofield, C. J. (2021) Synthesis of 2-oxoglutarate derivatives and their evaluation as cosubstrates and inhibitors of human aspartate/asparagine-β-hydroxylase. *Chem. Sci.***12**, 1327-1342
23. Nakashima, Y., Brewitz, L., Tumber, A., Salah, E., and Schofield, C. J. (2021) 2-Oxoglutarate derivatives can selectively enhance or inhibit the activity of human oxygenases. *Nat. Commun.***12**, 6478
24. Tumber, A., Salah, E., Brewitz, L., Corner, T., and Schofield, C. J. (2023) Kinetic and inhibition studies on human Jumonji-C (JmjC) domain-containing protein 5. *RSC Chem. Biol.***4**, 399-413
25. Breski, M., Dey, D., Obringer, S., Sudhamalla, B., and Islam, K. (2016) Engineering biological C–H functionalization leads to allele-specific regulation of histone demethylases. *J. Am. Chem. Soc.***138**, 13505-13508
26. Tiwari, A. D., Guan, Y., Grabowski, D. R., Maciejewski, J. P., Jha, B. K., and Phillips, J. G. (2021) SAR insights into TET2 catalytic domain inhibition: Synthesis of 2-hydroxy-4-methylene-pentanedicarboxylates. *Bioorg. Med. Chem.***39**, 116141
27. Stenflo, J., Holme, E., Lindstedt, S., Chandramouli, N., Tsai Huang, L. H., Tam, J. P., and Merrifield, R. B. (1989) Hydroxylation of aspartic acid in domains homologous to the epidermal growth factor precursor is catalyzed by a 2-oxoglutarate-dependent dioxygenase. *Proc. Natl. Acad. Sci. USA***86**, 444-447
28. Hansson, K., and Stenflo, J. (2005) Post-translational modifications in proteins involved in blood coagulation. *J. Thromb. Haemost.***3**, 2633-2648
29. Pfeffer, I., Brewitz, L., Krojer, T., Jensen, S. A., Kochan, G. T., Kershaw, N. J.*, et al.* (2019) Aspartate/asparagine-β-hydroxylase crystal structures reveal an unexpected epidermal growth factor-like domain substrate disulfide pattern. *Nat. Commun.***10**, 4910
30. Brewitz, L., Onisko, B. C., and Schofield, C. J. (2022) Combined proteomic and biochemical analyses redefine the consensus sequence requirement for epidermal growth factor-like domain hydroxylation. *J. Biol. Chem.***298**, 102129
31. Brasnett, A., Pfeffer, I., Brewitz, L., Chowdhury, R., Nakashima, Y., Tumber, A.*, et al.* (2021) Human oxygenase variants employing a single protein Fe(II) ligand are catalytically active. *Angew. Chem. Int. Ed.***60**, 14657-14663
32. Mahon, P. C., Hirota, K., and Semenza, G. L. (2001) FIH-1: a novel protein that interacts with HIF-1α and VHL to mediate repression of HIF-1 transcriptional activity. *Genes Dev.***15**, 2675-2686

33. Koivunen, P., Hirsilä, M., Günzler, V., Kivirikko, K. I., and Myllyharju, J. (2004) Catalytic properties of the asparaginyl hydroxylase (FIH) in the oxygen sensing pathway are distinct from those of its prolyl 4-hydroxylases. *J. Biol. Chem.***279**, 9899-9904
34. Cockman, M. E., Webb, J. D., Kramer, H. B., Kessler, B. M., and Ratcliffe, P. J. (2009) Proteomics-based identification of novel factor inhibiting hypoxia-inducible factor (FIH) substrates indicates widespread asparaginyl hydroxylation of ankyrin repeat domain-containing proteins. *Mol. Cell. Proteomics***8**, 535-546
35. Nagahama, K., Ogawa, T., Fujii, T., Tazaki, M., Tanase, S., Morino, Y., and Fukuda, H. (1991) Purification and properties of an ethylene-forming enzyme from Pseudomonas syringae pv. phaseolicola PK2. *J. Gen. Microbiol.***137**, 2281-2286
36. Fukuda, H., Ogawa, T., Ishihara, K., Fujii, T., Nagahama, K., Omata, T.*, et al.* (1992) Molecular cloning in Escherichia coli, expression, and nucleotide sequence of the gene for the ethylene-forming enzyme of Pseudomonas syringae pv. phaseolicola PK2. *Biochem. Biophys. Res. Commun.***188**, 826-832
37. Fukuda, H., Ogawa, T., Tazaki, M., Nagahama, K., Fujii, T., Tanase, S., and Morino, Y. (1992) Two reactions are simultaneously catalyzed by a single enzyme: The arginine-dependent simultaneous formation of two products, ethylene and succinate, from 2-oxoglutarate by an enzyme from Pseudomonas syringae. *Biochem. Biophys. Res. Commun.***188**, 483-489
38. Nagahama, K., Ogawa, T., Fujii, T., Tazaki, M., Goto, M., and Fukuda, H. (1991) L-Arginine is essential for the formation in vitro of ethylene by an extract of Pseudomonas syringae. *J. Gen. Microbiol.***137**, 1641-1646
39. Martinez, S., and Hausinger, R. P. (2016) Biochemical and spectroscopic characterization of the non-heme Fe(II)- and 2-oxoglutarate-dependent ethylene-forming enzyme from pseudomonas syringae pv. phaseolicola PK2. *Biochemistry***55**, 5989-5999
40. Zhang, Z., Smart, T. J., Choi, H., Hardy, F., Lohans, C. T., Abboud, M. I.*, et al.* (2017) Structural and stereoelectronic insights into oxygenase-catalyzed formation of ethylene from 2-oxoglutarate. *Proc. Natl. Acad. Sci. USA***114**, 4667-4672
41. Copeland, R. A., Davis, K. M., Shoda, T. K. C., Blaesi, E. J., Boal, A. K., Krebs, C., and Bollinger, J. M., Jr. (2021) An iron(IV)–oxo intermediate initiating L-arginine oxidation but not ethylene production by the 2-oxoglutarate-dependent oxygenase, ethylene-forming enzyme. *J. Am. Chem. Soc.***143**, 2293-2303
42. Wilson, J., Gering, S., Pinard, J., Lucas, R., and Briggs, B. R. (2018) Bio-production of gaseous alkenes: ethylene, isoprene, isobutene. *Biotechnol. Biofuels***11**, 234
43. Eckert, C., Xu, W., Xiong, W., Lynch, S., Ungerer, J., Tao, L.*, et al.* (2014) Ethylene-forming enzyme and bioethylene production. *Biotechnol. Biofuels***7**, 33
44. Copeland, R. A., Zhou, S., Schaperdoth, I., Shoda, T. K. C., Bollinger, J. M., and Krebs, C. (2021) Hybrid radical-polar pathway for excision of ethylene from 2-oxoglutarate by an iron oxygenase. *Science***373**, 1489-1493
45. Burke, E. J., Copeland, R. A., Dixit, Y., Krebs, C., and Bollinger, J. M., Jr. (2024) Steric perturbation of the Grob-like final step of ethylene-forming enzyme enables 3-hydroxypropionate and propylene production. *J. Am. Chem. Soc.***146**, 1977-1983
46. Pohl, L., and Eckle, M. (1969) Sodium 3-trimethylsilyltetradeuteriopropionate, a new water-soluble standard for ^{1}H-NMR. *Angew. Chem. Int. Ed.***8**, 381
47. Gates, E. L., Bradley, J. P., Berry, D. B. G., Nilsson, M., Morris, G. A., Adams, R. W., and Castañar, L. (2024) Solvent suppression in pure shift NMR. *Anal. Chem.***96**, 3879-3885
48. Castañar, L. (2017) Pure shift ^{1}H NMR: what is next? *Magn. Reson. Chem.***55**, 47-53
49. Foroozandeh, M., Morris, G. A., and Nilsson, M. (2018) PSYCHE pure shift NMR spectroscopy. *Chem. Eur. J.***24**, 13988-14000

50. Zangger, K. (2015) Pure shift NMR. *Prog. Nucl. Magn. Reson. Spectrosc.***86-87**, 1-20
51. Martinez, S., Fellner, M., Herr, C. Q., Ritchie, A., Hu, J., and Hausinger, R. P. (2017) Structures and mechanisms of the non-heme Fe(II)- and 2-oxoglutarate-dependent ethylene-forming enzyme: substrate binding creates a twist. *J. Am. Chem. Soc.***139**, 11980-11988
52. Weingart, H., Ullrich, H., Geider, K., and Völksch, B. (2001) The role of ethylene production in virulence of Pseudomonas syringae pvs. glycinea and phaseolicola. *Phytopathol.***91**, 511-518
53. North, J. A., Narrowe, A. B., Xiong, W., Byerly, K. M., Zhao, G., Young, S. J.*, et al.* (2020) A nitrogenase-like enzyme system catalyzes methionine, ethylene, and methane biogenesis. *Science***369**, 1094-1098
54. Ogawa, T., Takahashi, M., Fujii, T., Tazaki, M., and Fukuda, H. (1990) The role of NADH:Fe(III)EDTA oxidoreductase in ethylene formation from 2-keto-4-methylthiobutyrate. *J. Ferment. Bioeng.***69**, 287-291
55. Yin, X., and Zabriskie, T. M. (2004) VioC is a non-heme iron, α-ketoglutarate-dependent oxygenase that catalyzes the formation of 3S-hydroxy-L-arginine during viomycin biosynthesis. *ChemBioChem***5**, 1274-1277
56. Chang, C.-Y., Lyu, S.-Y., Liu, Y.-C., Hsu, N.-S., Wu, C.-C., Tang, C.-F.*, et al.* (2014) Biosynthesis of streptolidine involved two unexpected intermediates produced by a dihydroxylase and a cyclase through unusual mechanisms. *Angew. Chem. Int. Ed.***53**, 1943-1948
57. Pérez-Arellano, I., Carmona-Álvarez, F., Martínez, A. I., Rodríguez-Díaz, J., and Cervera, J. (2010) Pyrroline-5-carboxylate synthase and proline biosynthesis: from osmotolerance to rare metabolic disease. *Protein Sci.***19**, 372-382
58. Hellmann, H., Funck, D., Rentsch, D., and Frommer, W. B. (2000) Hypersensitivity of an Arabidopsis sugar signaling mutant toward exogenous proline application. *Plant Physiol.***123**, 779-789
59. Deuschle, K., Funck, D., Hellmann, H., Däschner, K., Binder, S., and Frommer, W. B. (2001) A nuclear gene encoding mitochondrial Δ1-pyrroline-5-carboxylate dehydrogenase and its potential role in protection from proline toxicity. *Plant J.***27**, 345-356
60. Qamar, A., Mysore, K., and Senthil-Kumar, M. (2015) Role of proline and pyrroline-5-carboxylate metabolism in plant defense against invading pathogens. *Front. Plant Sci.***6**, 503
61. Chalecka, M., Kazberuk, A., Palka, J., and Surazynski, A. (2021) P5C as an interface of proline interconvertible amino acids and its role in regulation of cell survival and apoptosis. *Int. J. Mol. Sci.***22**, 11763
62. Sleeman, M. C., and Schofield, C. J. (2004) Carboxymethylproline synthase (CarB), an unusual carbon-carbon bond-forming enzyme of the crotonase superfamily involved in carbapenem biosynthesis. *J. Biol. Chem.***279**, 6730-6736
63. Funck, D., Sinn, M., Forlani, G., and Hartig, J. S. (2024) Guanidine production by plant homoarginine-6-hydroxylases. *Elife***12**, RP91458
64. Ng, S. F., Hanauske-Abel, H. M., and Englard, S. (1991) Cosubstrate binding site of Pseudomonas sp. AK1 gamma-butyrobetaine hydroxylase. Interactions with structural analogs of alpha-ketoglutarate. *J. Biol. Chem.***266**, 1526-1533
65. Rude, M. A., Baron, T. S., Brubaker, S., Alibhai, M., Del Cardayre, S. B., and Schirmer, A. (2011) Terminal olefin (1-alkene) biosynthesis by a novel P450 fatty acid decarboxylase from Jeotgalicoccus species. *Appl. Environ. Microbiol.***77**, 1718-1727
66. Liu, Y., Wang, C., Yan, J., Zhang, W., Guan, W., Lu, X., and Li, S. (2014) Hydrogen peroxide-independent production of α-alkenes by OleTJE P450 fatty acid decarboxylase. *Biotechnol. Biofuels***7**, 28
67. Grant, J. L., Hsieh, C. H., and Makris, T. M. (2015) Decarboxylation of fatty acids to terminal alkenes by cytochrome P450 compound I. *J. Am. Chem. Soc.***137**, 4940-4943

68. Belcher, J., McLean, K. J., Matthews, S., Woodward, L. S., Fisher, K., Rigby, S. E. J., *et al.* (2014) Structure and biochemical properties of the alkene producing cytochrome P450 $OleT_{JE}$ (CYP152L1) from the Jeotgalicoccus sp. 8456 bacterium. *J. Biol. Chem.***289**, 6535-6550
69. Matthews, S., Belcher, J. D., Tee, K. L., Girvan, H. M., McLean, K. J., Rigby, S. E. J., *et al.* (2017) Catalytic determinants of alkene production by the cytochrome P450 peroxygenase OleTJE. *J. Biol. Chem.***292**, 5128-5143
70. Grant, J. L., Mitchell, M. E., and Makris, T. M. (2016) Catalytic strategy for carbon−carbon bond scission by the cytochrome P450 OleT. *Proc. Natl. Acad. Sci. USA***113**, 10049-10054
71. Mukherjee, D., and Laloraya, M. M. (1974) Metabolism of γ-methyl-α-ketoglutaric acid, γ-methylene-α-ketoglutaric acid and other keto acids during the seedling growth in Tamarindus indica. *Biochem. Physiol. Pflanz.***166**, 429-436
72. Virtanen, A. I., and Berg, A.-M. (1955) New aminodicarboxylic acids and corresponding α-keto acids in Phyllitis scolopendrium. *Acta Chem. Scand.***9**, 553-554
73. Díaz, R., Gallart-Ayala, H., Sancho, J. V., Nuñez, O., Zamora, T., Martins, C. P. B., *et al.* (2016) Told through the wine: a liquid chromatography–mass spectrometry interplatform comparison reveals the influence of the global approach on the final annotated metabolites in non-targeted metabolomics. *J. Chromatogr. A***1433**, 90-97
74. C. Winter, H., and E. Dekker, E. (1987) An unusual distribution of 4-substituted glutamic acids in Sophora japonica. *Phytochemistry***26**, 2477-2480
75. Rabara, R. C., Tripathi, P., Reese, R. N., Rushton, D. L., Alexander, D., Timko, M. P., *et al.* (2015) Tobacco drought stress responses reveal new targets for Solanaceae crop improvement. *BMC Genomics***16**, 484
76. Riedel, T. J., Johnson, L. C., Knight, J., Hantgan, R. R., Holmes, R. P., and Lowther, W. T. (2011) Structural and biochemical studies of human 4-hydroxy-2-oxoglutarate aldolase: implications for hydroxyproline metabolism in primary hyperoxaluria. *PLOS ONE***6**, e26021
77. Mazurkewich, S., Brott, A. S., Kimber, M. S., and Seah, S. Y. K. (2016) Structural and kinetic characterization of the 4-carboxy-2-hydroxymuconate hydratase from the gallate and protocatechuate 4,5-cleavage pathways of Pseudomonas putida KT2440. *J. Biol. Chem.***291**, 7669-7686
78. Maruyama, K. (1990) Purification and properties of 4-hydroxy-4-methyl-2-oxoglutarate aldolase from *Pseudomonas ochraceae* grown on phthalate. *J. Biochem.***108**, 327-333
79. Mazurkewich, S., Wang, W., and Seah, S. Y. K. (2014) Biochemical and structural analysis of RraA proteins To decipher their relationships with 4-hydroxy-4-methyl-2-oxoglutarate/4-carboxy-4-hydroxy-2-oxoadipate aldolases. *Biochemistry***53**, 542-553
80. Wilkins, A. L., Lu, Y., and Tan, S.-T. (1995) Extractives from New Zealand honeys. 5. Aliphatic dicarboxylic acids in New Zealand rewarewa (Knightea excelsa) honey. *J. Agric. Food Chem.***43**, 3021-3025
81. Dennis, D. A., Chapman, P. J., and Dagley, S. (1973) Degradation of protocatechuate in *Pseudomonas testosteroni* by a pathway involving oxidation of the product of *meta*-fission. *J. Bacteriol.***113**, 521-523
82. Tack, B. F., Chapman, P. J., and Dagley, S. (1972) Purification and properties of 4-hydroxy-4-methyl-2-oxoglutarate aldolase. *J. Biol. Chem.***247**, 6444-6449
83. Frimmersdorf, E., Horatzek, S., Pelnikevich, A., Wiehlmann, L., and Schomburg, D. (2010) How Pseudomonas aeruginosa adapts to various environments: a metabolomic approach. *Environ. Microbiol.***12**, 1734-1747
84. Verlinden, R. A. J., Hill, D. J., Kenward, M. A., Williams, C. D., and Radecka, I. (2007) Bacterial synthesis of biodegradable polyhydroxyalkanoates. *J. Appl. Microbiol.***102**, 1437-1449
85. Fiorini, G., and Schofield, C. J. (2024) Biochemistry of the hypoxia-inducible factor hydroxylases. *Curr. Opin. Chem. Biol.***79**, 102428

86. Hoffman, N. E., Yang, S. F., Ichihara, A., and Sakamura, S. (1982) Stereospecific conversion of 1-aminocyclopropanecarboxylic acid to ethylene by plant tissues 1: conversion of stereoisomers of 1-amino-2-ethylcyclopropanecarboxylic acid to 1-butene. *Plant Physiol.***70**, 195-199
87. Pirrung, M. C. (1999) Ethylene biosynthesis from 1-aminocyclopropanecarboxylic acid. *Acc. Chem. Res.***32**, 711-718
88. Baldwin, J. E., Adlington, R. M., Lajoie, G. A., and Rawlings, B. J. (1985) On the biosynthesis of ethylene. Determination of the stereochemical course using modified substrates. *J. Chem. Soc., Chem. Commun.* 1496-1498
89. Hélaine, V., Rossi, J., Gefflaut, T., Alaux, S., and Bolte, J. (2001) Synthesis of 4,4-disubstituted L-glutamic acids by enzymatic transamination. *Adv. Synth. Catal.***343**, 692-697
90. Xian, M., Alaux, S., Sagot, E., and Gefflaut, T. (2007) Chemoenzymatic synthesis of glutamic acid analogues: substrate specificity and synthetic applications of branched chain aminotransferase from Escherichia coli. *J. Org. Chem.***72**, 7560-7566
91. Liu, X., Reinbold, R., Liu, S., Herold, R. A., Rabe, P., Duclos, S.*, et al.* (2023) Natural and synthetic 2-oxoglutarate derivatives are substrates for oncogenic variants of human isocitrate dehydrogenase 1 and 2. *J. Biol. Chem.***299**, 102873
92. Chatterjee, S., Fellner, M., Rankin, J., Thomas, M. G., J S Rifayee, S. B., Christov, C. Z.*, et al.* (2024) Structural, spectroscopic, and computational insights from canavanine-bound and two catalytically compromised variants of the ethylene-forming enzyme. *Biochemistry***63**, 1038-1050
93. Brewitz, L., Tumber, A., and Schofield, C. J. (2020) Kinetic parameters of human aspartate/asparagine-β-hydroxylase suggest that it has a possible function in oxygen sensing. *J. Biol. Chem.***295**, 7826-7838
94. Yang, M., Hardy, A. P., Chowdhury, R., Loik, N. D., Scotti, J. S., McCullagh, J. S. O.*, et al.* (2013) Substrate selectivity analyses of factor inhibiting hypoxia-inducible factor. *Angew. Chem. Int. Ed.***52**, 1700-1704
95. Wang, B., Xu, Y., Wang, X., Yuan, J. S., Johnson, C. H., Young, J. D., and Yu, J. (2021) A guanidine-degrading enzyme controls genomic stability of ethylene-producing cyanobacteria. *Nat. Commun.***12**, 5150
96. Adams, R. W., Holroyd, C. M., Aguilar, J. A., Nilsson, M., and Morris, G. A. (2013) "Perfecting" WATERGATE: clean proton NMR spectra from aqueous solution. *Chem. Commun.***49**, 358-360
97. Wishart, D. S., Knox, C., Guo, A. C., Eisner, R., Young, N., Gautam, B.*, et al.* (2009) HMDB: a knowledgebase for the human metabolome. *Nucleic Acids Res.***37**, D603-D610

2.10 Supporting Information

2.10.1 General Information


^{1}H NMR assays (600 MHz) were performed using a Bruker AVIII 600 instrument with a Prodigy nitrogen broadband cryoprobe, Pure shift ^{1}H NMR (700 MHz) analysis of reaction mixtures was performed as described (1) using a Bruker AVIII 700 MHz NMR spectrometer equipped with a 5-mm inverse triple-resonance-inverse cryoprobe. 4-Hydroxy-2OG (**4**) was obtained as a racemate from Merck UK Ltd; all other chiral 2OG derivatives were prepared as racemates as reported (2).

For ^{1}H NMR experiments, water suppression at ~4.7 ppm was achieved using the perfect echo-modified WATERGATE method (3). 3-(Trimethylsilyl)-2,2,3,3-tetradeuteropropionic acid (4) (TMSP-d_4) was used as an internal standard (–0.11 ppm) to enable product quantification via comparison of peak integrals with that of TMSP-d_4. The stereochemistry of P5C was not assigned, note that it may epimerize under the assay conditions.


Symbols are used to annotate signals in the ^{1}H NMR spectra as shown below:

Annotation	Molecule(s)
squares	L-ascorbate (LAA)
pentagons	L-arginine (Arg)
inverted triangle	L-Δ^1-pyrroline-5-carboxylate (P5C)
circles	2OG/2OG derivatives
diamonds/triangles	PsEFE reaction products


Colors are used to distinguish between chemically distinct protons, as shown below:




2.10.2 Supporting Figures




Supporting Figure 2.S1. ^{1}H NMR analysis of PsEFE catalysis using 2-oxoglutarate (2OG) as a cosubstrate under optimized assay conditions. a-e) ^{1}H NMR analysis of the reaction of PsEFE with 2OG (**1**) after: **a**) 60 min, **b**) 40 min, **c**) 20 min, **d**) 10 min, and **e**) 5 min. **f**) ^{1}H NMR analysis of the reaction mixture in the absence of PsEFE. Conditions: 400 µM 2OG (**1**), 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4 (4), and 2 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S2. ^{1}H NMR analysis implies that 3-hydroxypropionate (1c) is not a product of PsEFE catalysis using 2OG (1) and L-Arg as substrates. a) ^{1}H NMR analysis of the reaction of PsEFE with 2OG (**1**) and L-Arg spiked with a synthetic standard of 3-hydroxypropionate (**1c**) indicates that ethylene (5.35 ppm), succinate (2.32 ppm), and P5C (7.70 ppm) are products, no evidence for 3-hydroxypropionate (**1c**) was observed; **b**) ^{1}H NMR spectrum of synthetic 3-hydroxypropionate (**1c**) under standard conditions in the absence of PsEFE; **c**) ^{1}H NMR analysis of the reaction of PsEFE with 2OG (**1**) 12 h post addition of PsEFE; **d**) ^{1}H NMR spectrum of 2OG (**1**) under standard conditions in the absence of PsEFE. Conditions: 400 µM 2OG (**1**), 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, 2 µM PsEFE, and, if appropriate, 500 µM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The singlet observed at ~4.4 ppm originates from the enzyme sample; it was observed in the enzyme-only control. The two low level triplets at ~ 2.4 and ~ 2.5 ppm originate from the commercially obtained sample of **1c**.




Supporting Figure 2.S3. Analysis of the outcome of the reaction of PsEFE with 2OG (1) using pure shift ^{1}H NMR. Analysis of the reaction of PsEFE with 2OG (**1**) 1 h post addition of PsEFE using: **a**) pure shift ^{1}H NMR (1, 5-7) and **b**) ^{1}H NMR. The results indicate that substantial amounts of reaction products other than ethylene, succinate, and P5C (Supporting Figures 2.S1 and 2.S2) are not formed under the tested conditions. Analysis of the reaction mixture with 2OG (**1**) in the absence of PsEFE using: **c**) pure shift ^{1}H NMR (1, 5-7) and **d**) ^{1}H NMR. Conditions: 400 µM 2OG (**1**), 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, and 2 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Note that for the pure shift ^{1}H NMR analysis, the observed peak may not necessarily be at the center of the peak obtained when using standard NMR analysis (1, 5-7). The singlet observed at ~4.4 ppm originates from the enzyme sample; it was observed in the enzyme-only control.




Supporting Figure 2.S4. Low levels of dehydroascorbate (1d) are formed by the oxidation of L-ascorbate during PsEFE catalysis. a) ^{1}H NMR analysis of the reaction of PsEFE with 2OG (**1**) spiked with synthetic dehydroascorbate indicates that low levels of dehydroascorbate (**1d**) are present in the reaction mixture; **b**) ^{1}H NMR spectrum of **1d** in buffer; **c**) ^{1}H NMR analysis of the reaction of PsEFE with 2OG 12 h post addition of PsEFE. Conditions: 400 µM 2OG, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, and 2 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The unassigned singlets observed in the authentic synthetic sample of dehydroascorbate at ~2.6, ~2.8, and ~2.95 ppm are also present in the reported ^{1}H NMR spectrum of **1d** (8).




Supporting Figure 2.S5. Quantification of ethylene formation using ^{1}H NMR. a) The indirect quantification of ethylene formation by subtracting the normalized integral of succinate (obtained via comparison with TMSP-d_4) from that of 2OG (**1**) was more reliable than the direct integration of the ethylene signal (~5.35 ppm), possibly because ethylene evaporates from the reaction mixture; **b**) ^{1}H NMR analysis of the crude reaction mixture implies that PsEFE-catalyzed succinate formation correlates stoichiometrically with Arg hydroxylation to give P5C, suggesting that subtracting the normalized integral of succinate (obtained via comparison with TMSP-d_4) from that of 2OG is a valid method to estimate ethylene formation; note that substantial levels of products other than succinate and ethylene were not detected from 2OG (**1**) during PsEFE catalysis (Supporting Figures 2.S1 and 2.S2). Conditions: 400 μM 2OG (**1**), 500 μM *L*-arginine, 500 μM *L*-ascorbate, 50 μM Fe(II), 800 μM TMSP-d_4, and 2 μM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S6. PsEFE catalyses the conversion of *L*-arginine to P5C using 2OA (2) as a cosubstrate. **a**) ^{1}H NMR analysis of the reaction of PsEFE with 2OA (**2**) as a cosubstrate reveals formation of P5C; **b**) ^{1}H NMR spectrum of **2** under standard conditions in the absence of PsEFE. Conditions: 400 µM **2**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S7. Time-dependent PsEFE-catalyzed conversion of 2OA (2). a-f) Representative ^{1}H NMR spectra (~1.4 to ~3.6 ppm) monitoring the reaction of PsEFE with 2OA (**2**) after: **a**) 60 min, **b**) 40 min, **c**) 20 min, **d**) 15 min, **e**) 10 min, and **f**) 5 min. **g**) ^{1}H NMR analysis of the reaction mixture in the absence of PsEFE. Conditions: 400 µM **2**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S8. ^{1}H NMR analysis of 2OA (2) and authentic samples of its products formed during PsEFE catalysis. a-c) ^{1}H NMR spectra (~1.4 to ~3.6 ppm) of: **a**) γ-hydroxybutyrate (GHB, **2b**), **b**) glutarate (**2a**), and **c**) 2OA (**2**) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S9. PsEFE catalyzes the conversion of *L*-arginine to P5C using racemic 4-methyl-2OG (3) as a cosubstrate. a) ^{1}H NMR analysis of the reaction of PsEFE with 4-methyl-2OG (**3**) reveals formation of P5C; **b**) ^{1}H NMR spectrum of **3** under standard conditions in the absence of PsEFE. Conditions: 400 µM **3**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control. The stereochemistry of P5C was not assigned, as it may epimerize under the assay conditions.




Supporting Figure 2.S10. Time-dependent PsEFE-catalyzed conversion of racemic 4-methyl-2OG (3). a-f) Representative ^{1}H NMR spectra (~0.7 to ~3.7 ppm) monitoring the reaction of PsEFE with 4-methyl-2OG (**3**) after: **a**) 60 min, **b**) 40 min, **c**) 20 min, **d**) 15 min, **e**) 10 min, and **f**) 5 min. **g**) ^{1}H NMR analysis of the reaction mixture in the absence of PsEFE. Conditions: 400 µM **3**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The doublet observed at ~1.2 ppm originates from the compound-only control of 4-methyl 2OG (**3**).




Supporting Figure 2.S11. ^{1}H NMR analysis of racemic 4-methyl-2OG (3) and authentic samples of its products formed during PsEFE catalysis. a-c) ^{1}H NMR spectra (~0.7 to ~3.7 ppm) of: **a**) β-hydroxyisobutyrate (**3b**), **b**) 2-methylsuccinate (**3a**), and **c**) 4-methyl 2OG (**3**) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S12. Analysis of PsEFE catalysis using racemic 4-methyl-2OG (3) as a substrate. ^{1}H NMR analysis (~0.7 to ~3.7 ppm) of: **a**) a reaction mixture of PsEFE and racemic 4-methyl-2OG (**3**) spiked with β-hydroxyisobutyrate (**3b**), **b**) **3b** in the absence of **3** under standard conditions; **c**) a reaction mixture of PsEFE and **3** spiked with 2-methylsuccinate (**3a**), **d**) **3a** in the absence of **3** under standard conditions; **e**) a reaction mixture of PsEFE and **3** 80 min post addition of PsEFE; **f**) **3** under standard conditions in the absence of PsEFE. Conditions: 400 μM **3**, 500 μM L-arginine, 500 μM L-ascorbate, 50 μM Fe(II), 800 μM TMSP-d_4, 10 μM PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The singlet observed at ~4.4 ppm originates from the enzyme sample; it was observed in the enzyme-only control.




Supporting Figure 2.S13. PsEFE catalyzes the conversion of *L*-arginine to P5C using racemic 4-OH-2OG (4) as a cosubstrate. a) ^{1}H NMR analysis of the reaction of PsEFE with racemic 4-OH-2OG (**4**) reveals formation of P5C; **b**) ^{1}H NMR spectrum of **4** under standard conditions in the absence of PsEFE. Conditions: 400 µM **4**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S14. Time-dependent PsEFE-catalyzed conversion of racemic 4-OH-2OG (4). a-f) Representative ^{1}H NMR spectra (~2.1 to ~4.5 ppm) monitoring the reaction of PsEFE with racemic 4-OH-2OG (**4**) after: **a**) 60 min, **b**) 40 min, **c**) 20 min, **d**) 15 min, **e**) 10 min, and **f**) 5 min. **g**) ^{1}H NMR analysis of the reaction mixture in the absence of PsEFE. Conditions: 400 µM **4**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The singlet observed at ~4.4 ppm originates from the enzyme sample; it was observed in the enzyme-only control.




Supporting Figure 2.S15. ^{1}H NMR analysis of racemic 4-OH-2OG (4) and authentic samples of its products formed during PsEFE catalysis. a-c) ^{1}H NMR spectra (~2.1 to ~4.5 ppm) of: **a**) malate (**4a**), **b**) glycerate (**4b**), and **c**) racemic 4-OH-2OG (**4**) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S16. Analysis of PsEFE catalysis using racemic 4-OH-2OG (4) as a substrate. ^{1}H NMR analysis (~2.1 to ~4.5 ppm) of: **a**) a reaction mixture of PsEFE and racemic 4-OH-2OG (**4**) spiked with glycerate (**4b**), **b**) **4b** in the absence of **4** under standard conditions; **c**) a reaction mixture of PsEFE and **4** spiked with malate (**4a**), **d**) **4a** in the absence of **4** under standard conditions; **e**) a reaction mixture of PsEFE and **4** 80 min post addition of PsEFE; **f**) **4** under standard conditions in the absence of PsEFE. Conditions: 400 µM **4**, 500 µM L-arginine, 500 µM L-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM PsEFE, and, if appropriate, 2 mM of an appropriate authentic standard in buffer (50 mM phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The singlet observed at ~4.4 ppm originates from the enzyme sample; it was observed in the enzyme-only control.




Supporting Figure 2.S17. PsEFE catalyzes the conversion of *L*-arginine to P5C using racemic 4-hydroxy-4-methyl-2OG (5) as a cosubstrate. a) ^{1}H NMR analysis of the reaction of PsEFE with racemic 4-hydroxy-4-methyl-2OG (**5**) as a cosubstrate reveals formation of P5C; **b**) ^{1}H NMR spectrum of **5** under standard conditions in the absence of PsEFE. Conditions: 400 µM **5**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S18. Time-dependent PsEFE-catalyzed conversion of racemic 4-hydroxy-4-methyl-2OG (5). a-f) Representative ^{1}H NMR spectra (~0.8 to ~3.7 ppm) monitoring the reaction of PsEFE with racemic 4-hydroxy-4-methyl-2OG (**5**) after: **a**) 60 min, **b**) 40 min, **c**) 20 min, **d**) 15 min, **e**) 10 min, and **f**) 5 min. **g**) ^{1}H NMR analysis of the reaction mixture in the absence of PsEFE. Conditions: 400 µM **5**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The doublet observed at ~1.2 ppm originates from the enzyme sample; it was observed in the enzyme-only control.




Supporting Figure 2.S19. ^{1}H NMR analysis of racemic 4-hydroxy-4-methyl-2OG (5) and authentic samples of its products formed during PsEFE catalysis. a-c) ^{1}H NMR spectra (~0.8 to ~3.7 ppm) of: **a**) racemic α,β-dihydroxyisobutyrate (**5b**), **b**) racemic citramalate (**5a**), and **c**) racemic 4-hydroxy-4-methyl-2OG (**5**) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S20. Analysis of PsEFE catalysis using racemic 4-hydroxy-4-methyl-2OG (5) as a substrate. ^{1}H NMR analysis (~0.8 to ~3.7 ppm) of: **a**) a reaction mixture of PsEFE and racemic 4-hydroxy-4-methyl-2OG (**5**) spiked with racemic α,β-dihydroxyisobutyrate (**5b**), **b**) **5b** in the absence of **5** under standard conditions; **c**) a reaction mixture of PsEFE and **5** spiked with racemic citramalate (**5a**), **d**) **5a** in the absence of **5** under standard conditions; **e**) a reaction mixture of PsEFE and **5** 80 min post addition of PsEFE; **f**) **5** under standard conditions in the absence of PsEFE. Conditions: 400 µM **5**, 500 µM L-arginine, 500 µM L-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, 10 µM PsEFE, and, if appropriate, 2 mM of an appropriate authentic standard in buffer (50 mM phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The doublet observed at ~1.2 ppm originates from the enzyme sample; it was observed in the enzyme-only control.




Supporting Figure 2.S21. PsEFE catalyzes the conversion of *L*-arginine to P5C using 4,4-dimethyl-2OG (6) as a cosubstrate. a) ^{1}H NMR analysis of the reaction of PsEFE with 4,4-dimethyl-2OG (**6**) as a cosubstrate reveals formation of P5C; **b**) ^{1}H NMR spectrum of **5** under standard conditions in the absence of PsEFE. Conditions: 400 µM **6**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S22. Time-dependent PsEFE-catalyzed conversion of 4,4-dimethyl-2OG (6). a-f) Representative ^{1}H NMR spectra (~0.7 to ~3.5 ppm) monitoring the reaction of PsEFE with 4,4-dimethyl-2OG (**6**) after: **a**) 60 min, **b**) 40 min, **c**) 20 min, **d**) 15 min, **e**) 10 min, and **f**) 5 min. **g**) ^{1}H NMR analysis of the reaction mixture in the absence of PsEFE. Conditions: 400 µM **6**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S23. ^{1}H NMR analysis of 4,4-dimethyl-2OG (6) and authentic samples of its products formed during PsEFE catalysis. a-c) ^{1}H NMR spectra (~0.7 to ~3.7 ppm) of: **a**) hydroxypivalate (**6b**), **b**) 2,2-dimethylsuccinate (**6a**), and **c**) 4,4-dimethyl-2OG (**6**) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S24. Analysis of PsEFE catalysis using 4,4-dimethyl-2OG (6) as a substrate. ^{1}H NMR analysis (~0.7 to ~3.7 ppm) of: **a**) a reaction mixture of PsEFE and 4,4-dimethyl-2OG (**6**) spiked with 2,2-dimethylsuccinate (**6a**), **b**) **6a** in the absence of **6** under standard conditions; **c**) a reaction mixture of PsEFE and **6** spiked with hydroxypivalate (**6b**), **d**) **6b** in the absence of **6** under standard conditions; **e**) a reaction mixture of PsEFE and **6** 80 min post addition of PsEFE; **f**) **6** under standard conditions in the absence of PsEFE. Conditions: 400 µM **6**, 500 µM L-arginine, 500 µM L-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM PsEFE, and, if appropriate, 2 mM of an appropriate authentic standard in buffer (50 mM phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S25. PsEFE catalyzes the conversion of *L*-arginine to P5C using racemic 4-ethyl-2OG (7) as a cosubstrate. a) ^{1}H NMR analysis of the reaction of PsEFE with racemic 4-ethyl-2OG (**7**) as a cosubstrate reveals formation of P5C; **b**) ^{1}H NMR spectrum of **7** under standard conditions in the absence of PsEFE. Conditions: 400 µM **7**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S26. Time-dependent PsEFE-catalyzed conversion of racemic 4-ethyl-2OG (7). a-f) Representative ^{1}H NMR spectra (~0.7 to ~3.4 ppm) monitoring the reaction of PsEFE with racemic 4-ethyl-2OG (**6**) after: **a**) 60 min, **b**) 40 min, **c**) 20 min, **d**) 15 min, **e**) 10 min, and **f**) 5 min. **g**) ^{1}H NMR analysis of the reaction mixture in the absence of PsEFE. Conditions: 400 µM **7**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The doublet observed at ~1.2 ppm originates from the enzyme sample; it was observed in the enzyme-only control.




Supporting Figure 2.S27. ^{1}H NMR analysis of racemic 4-ethyl-2OG (7) and authentic samples of its products formed during PsEFE catalysis. a-c) ^{1}H NMR spectra (~0.4 to ~4.0 ppm) of: **a**) racemic 2-(hydroxymethyl)butyrate (**7b**), **b**) racemic 2-ethylsuccinate (**7a**), and **c**) racemic 4-ethyl-2OG (**7**) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S28. Analysis of PsEFE catalysis using racemic 4-ethyl-2OG (7) as a substrate. ^{1}H NMR analysis (~0.7 to ~3.7 ppm) of: **a**) a reaction mixture of PsEFE and racemic 4-ethyl-2OG (**7**) spiked with 2-(hydroxymethyl)butyrate (**7b**), **b**) **7b** in the absence of **7** under standard conditions; **c**) a reaction mixture of PsEFE and **7** spiked with 2-ethylsuccinate (**7a**), **d**) **7a** in the absence of **7** under standard conditions; **e**) a reaction mixture of PsEFE and **7** 160 min post addition of PsEFE; **f**) **7** under standard conditions in the absence of PsEFE. Conditions: 400 µM **7**, 500 µM L-arginine, 500 µM L-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM PsEFE, and, if appropriate, 2 mM of an appropriate authentic standard in buffer (50 mM phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The doublet observed at ~1.2 ppm originates from the enzyme sample; it was observed in the enzyme-only control.




Supporting Figure 2.S29. PsEFE catalyses the conversion of *L*-arginine to P5C using racemic 4-propyl-2OG (8) as a cosubstrate. a) ^{1}H NMR analysis of the reaction of PsEFE with racemic 4-propyl-2OG (**8**) as a cosubstrate reveals formation of P5C; **b**) ^{1}H NMR spectrum of **8** under standard conditions in the absence of PsEFE. Conditions: 400 µM **8**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S30. Time-dependent PsEFE-catalyzed conversion of racemic 4-propyl-2OG (8). a-f) Representative ^{1}H NMR spectra (~0.7 to ~3.7 ppm) monitoring the reaction of PsEFE with racemic 4-propyl-2OG (**8**) after: **a**) 120 min, **b**) 60 min, **c**) 40 min, **d**) 20 min, **e**) 10 min, and **f**) 5 min. **g**) ^{1}H NMR analysis of the reaction mixture in the absence of PsEFE. Conditions: 400 µM **8**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S31. ^{1}H NMR analysis of racemic 4-propyl-2OG (8) and authentic samples of its products formed during PsEFE catalysis. a-c) ^{1}H NMR spectra (~0.4 to ~4.0 ppm) of: **a**) 2-(hydroxymethyl)pentanoate (**8b**), **b**) 2-propylsuccinate (**8a**), and **c**) racemic 4-propyl-2OG (**8**) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S32. Analysis of PsEFE catalysis using racemic 4-propyl-2OG (8) as a substrate. ^{1}H NMR analysis (~0.7 to ~3.4 ppm) of: **a**) a reaction mixture of PsEFE and racemic 4-propyl-2OG (**8**) spiked with 2-(hydroxymethyl)pentanoate (**8a**) 120 min post-incubation with PsEFE, **b**) **8a** in the absence of **8** under standard conditions; **c**) a reaction mixture of PsEFE and **8** spiked with 2-propylsuccinate (**8b**) 120 min post-incubation with PsEFE, **d**) **8b** in the absence of **8** under standard conditions; **e**) a reaction mixture of PsEFE and **8** 120 min post addition of PsEFE; **f**) **8** under standard conditions in the absence of PsEFE. Conditions: 400 µM **8**, 500 µM L-arginine, 500 µM L-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM PsEFE, and, if appropriate, 2 mM of an appropriate authentic standard in buffer (50 mM phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The doublet observed at ~1.2 ppm originates from the enzyme sample; it was observed in the enzyme-only control.




Supporting Figure 2.S33. PsEFE catalyses the conversion of *L*-arginine to P5C using racemic 4-butyl-2OG (9) as a cosubstrate. a) ^{1}H NMR analysis of the reaction of PsEFE with racemic 4-butyl-2OG (**9**) as a cosubstrate reveals formation of P5C; **b**) ^{1}H NMR spectrum of **9** under standard conditions in the absence of PsEFE. Conditions: 400 µM **9**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S34. Time-dependent PsEFE-catalyzed conversion of racemic 4-butyl-2OG (9). a-f) Representative ^{1}H NMR spectra (~0.7 to ~3.4 ppm) monitoring the reaction of PsEFE with racemic 4-butyl-2OG (**9**) after: **a**) 160 min, **b**) 80 min, **c**) 40 min, **d**) 20 min, **e**) 10 min, and **f**) 5 min. **g**) ^{1}H NMR analysis of the reaction mixture in the absence of PsEFE. Conditions: 400 µM **9**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Note the two doublets at ~1.07 and ~0.97 ppm increase over time, but do not appear to be a direct product of **9**, because the sum of integrals for **9** and **9a** implies the presence of 400 µM **9** and **9a** (*i.e.*, the starting concentration of **9**) at any given time point during the assay (Figure 2.5i). The species giving rise to the signals at ~1.07 and ~0.97 ppm, however, has not been assigned.




Supporting Figure 2.S35. ^{1}H NMR analysis of racemic 4-butyl-2OG (9) and authentic sample of its products formed during PsEFE catalysis. a-c) ^{1}H NMR spectra (~0.4 to ~4.0 ppm) of: **a**) 2-(hydroxymethyl)hexanoate (**9b**), **b)** 2-butylsuccinate (**9a**), and **c**) racemic 4-butyl-2OG (**9**) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S36. Analysis of PsEFE catalysis using racemic 4-butyl-2OG (9) as a substrate. ^{1}H NMR analysis (~0.7 to ~3.4 ppm) of: **a**) 2-(hydroxymethyl)hexanoate (**9a**) in the absence of racemic 4-butyl-2OG (**9**) under standard conditions, **b)** a reaction mixture of PsEFE and **9** spiked with 2-butylsuccinate (**9b**), **c**) **9b** in the absence of **9** under standard conditions; **d**) a reaction mixture of PsEFE and **9** 180 min post addition of PsEFE; **e**) **9** under standard conditions in the absence of PsEFE. Conditions: 400 µM **9**, 500 µM L-arginine, 500 µM L-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM PsEFE, and, if appropriate, 2 mM of an appropriate authentic standard in buffer (50 mM phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Note the two doublets at ~1.07 and ~0.97 ppm increase over time, but do not appear to be a direct product of **9**, because the sum of integrals for **9** and **9a** implies the presence of 400 µM **9** and **9a** (*i.e.*, the starting concentration of **9**) at any given time point during the assay (Figure 2.5i). The species giving rise to the signals at ~1.07 and ~0.97 ppm, however, has not been assigned. Note that the signals at ~3.5 ppm observed originate from the enzyme sample; it was observed in the enzyme-only control and the integrals do not increase over time (Supporting Figure 2.S34).




Supporting Figure 2.S37. Analysis of the reaction outcome of PsEFE catalyzed conversion of racemic 4-butyl 2OG (9) using pure shift ^{1}H NMR. Analysis of the reaction of PsEFE with racemic 4-butyl 2OG (**9**) 1 h post addition of PsEFE using: **a**) pure shift ^{1}H NMR (1, 5-7) and **b**) ^{1}H NMR results indicate that substantial amounts of reaction products other than 2-butylsuccinate (**9a**) (Supporting Figures 2.S33-2.36) are not being formed under the tested conditions. Analysis of the reaction mixture with **9** in the absence of PsEFE using: **c**) pure shift ^{1}H NMR (1, 5-7) and **d**) ^{1}H NMR. Conditions: 400 µM **9**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Note that for the pure shift ^{1}H NMR analysis, the observed peak may not necessarily be at the center of the peak obtained when using standard NMR analysis (1, 5-7). The singlet observed at ~4.4 ppm originates from the enzyme sample; it was observed in the enzyme-only control. Note the two doublets at ~1.07 and ~0.97 ppm increase over time, but do not appear to be a direct product of **9**, because the sum of integrals for **9** and **9a** implies the presence of 400 µM **9** and **9a** (*i.e.*, the starting concentration of **9**) at any given time point during the assay (Figure 2.5i). The species giving rise to the signals at ~1.07 and ~0.97 ppm, however, has not been assigned.




Supporting Figure 2.S38. PsEFE catalyzes the conversion of *L*-arginine to P5C using racemic 3-methyl-2OG (10) as a cosubstrate. a) ^{1}H NMR analysis of the reaction of PsEFE with racemic 3-methyl-2OG (**10**) as a cosubstrate reveals formation of P5C; **b**) ^{1}H NMR spectrum of **10** under standard conditions in the absence of PsEFE. Conditions: 400 µM **10**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The singlet observed at ~4.4 ppm originates from the enzyme sample; it was observed in the enzyme-only control.




Supporting Figure 2.S39. Time-dependent PsEFE-catalyzed conversion of racemic 3-methyl-2OG (10). a-f) Representative ^{1}H NMR spectra (~0.8 to ~4.2 ppm) monitoring the reaction of PsEFE with racemic 3-methyl-2OG (**10**) after: **a**) 60 min, **b**) 40 min, **c**) 20 min, **d**) 15 min, **e**) 10 min, and **f**) 5 min. **g**) ^{1}H NMR analysis of the reaction mixture in the absence of PsEFE. Conditions: 400 µM **10**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S40. ^{1}H NMR analysis of racemic 3-methyl-2OG (10) and authentic samples of its products formed during PsEFE catalysis. a-c) ^{1}H NMR spectra (~0.8 to ~4.2 ppm) of: **a**) β-hydroxybutyrate (**10b**), **b**) 2-methylsuccinate (**3a**), and **c**) racemic 3-methyl-2OG (**10**) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S41. Analysis of PsEFE catalysis using racemic 3-methyl-2OG (10) as a substrate. ^{1}H NMR analysis (~0.8 to ~4.2 ppm) of: **a**) a reaction mixture of PsEFE and racemic 3-methyl-2OG (**10**) spiked with β-hydroxybutyrate (**10b**), **b**) **10b** in the absence of **10** under standard conditions; **c**) a reaction mixture of PsEFE and **10** spiked with 2-methylsuccinate (**3a**), **d**) **3a** in the absence of **10** under standard conditions; **e**) a reaction mixture of PsEFE and **10** 100 min post addition of PsEFE; **f**) **10** under standard conditions in the absence of PsEFE. Conditions: 400 µM **10**, 500 µM L-arginine, 500 µM L-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM PsEFE, and, if appropriate, 2 mM of an appropriate authentic standard in buffer (50 mM phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S42. Analysis of the reaction outcome of PsEFE catalysed conversion of racemic 3-methyl-2OG (10) using pure shift ^{1}H NMR. Analysis of the reaction of PsEFE with racemic 3-methyl-2OG (**10**) 30 min post addition of PsEFE using: **a**) pure shift ^{1}H NMR (1, 5-7) and **b**) ^{1}H NMR results indicate that substantial amounts of reaction products other than β-hydroxyisobutyrate (**10b**), 2-methylsuccinate (**3a**), and P5C (Supporting Figures 2.S38-2.S41) are not being formed under the tested conditions. Analysis of the reaction mixture with **10** in the absence of PsEFE using: **c**) pure shift ^{1}H NMR (1, 5-7) and **d**) ^{1}H NMR. Conditions: 400 µM **10**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Note that for the pure shift ^{1}H NMR analysis, the observed peak may not necessarily be at the center of the peak obtained when using standard NMR analysis (1, 5-7). The singlet observed at ~4.4 ppm originates from the enzyme sample; it was observed in the enzyme-only control.




Supporting Figure 2.S43. PsEFE catalyzes the conversion of *L*-arginine to P5C using 4-(carboxycarbonyl)cyclohexane-1-carboxylate (14) as a cosubstrate. a) ^{1}H NMR analysis of the reaction of PsEFE with 4-(carboxycarbonyl)cyclohexane-1-carboxylate (**14**) as a cosubstrate reveals formation of P5C; **b**) ^{1}H NMR spectrum of **14** under standard conditions in the absence of PsEFE. Conditions: 400 µM **14**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 30 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). **14** was used as a mixture of diastereomers, dr (*trans*:*cis*) = 5:1. Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S44. ^{1}H NMR analysis of 4-(carboxycarbonyl)cyclohexane-1-carboxylic acid (14) and authentic samples of its products formed during PsEFE catalysis. a-c) ^{1}H NMR spectra (~0.8 to ~3.0 ppm) of: **a**) a *cis*/*trans* isomeric mixture of 4-hydroxycyclohexanecarboxylate (**14b**), **b**) a *cis*/*trans* isomeric mixture of cyclohexane-1,4-dicarboxylate (**14a**), and **c**) 4-(carboxycarbonyl)cyclohexane-1-carboxylate (**14**) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). **14** was used as a mixture of diastereomers, dr (*trans*:*cis*) = 5:1 (2).




Supporting Figure 2.S45. PsEFE catalyzes the conversion of *L*-arginine to P5C using racemic 4-benzyl-2OG (15) as a cosubstrate. a) ^{1}H NMR analysis of the reaction of PsEFE with racemic 4-benzyl-2OG (**15**) as a cosubstrate reveals formation of P5C; **b**) ^{1}H NMR spectrum of **15** under standard conditions in the absence of PsEFE. Conditions: 400 µM **15**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 30 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S46. ^{1}H NMR analysis of racemic 4-benzyl-2OG (15) and authentic samples of its products formed during PsEFE catalysis. a-c) ^{1}H NMR spectra (~2.0 to ~7.8 ppm) of: **a**) racemic 2-benzyl-3-hydroxypropanoate (**15b**), **b**) racemic 2-benzylsuccinate (**15a**), and **c**) racemic 4-benzyl-2OG (**15**) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O).




Supporting Figure 2.S47. Analysis of PsEFE catalysis using racemic 4-benzyl-2OG (15) as a substrate. ^{1}H NMR analysis (~2.0 to ~7.8 ppm) of: **a**) racemic 2-benzyl-3-hydroxypropanoate (**15b**) in the absence of racemic 4-benzyl-2OG (**15**) under standard conditions, **b**) a reaction mixture of PsEFE and **15** spiked with racemic 2-benzylsuccinate (**15a**), **c**) **15a** in the absence of **15** under standard conditions; **d**) a reaction mixture of PsEFE and **15** 12 h post addition of PsEFE; **e**) **15** under standard conditions in the absence of PsEFE. Conditions: 400 µM **15**, 500 µM L-arginine, 500 µM L-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, 30 µM PsEFE, and, if appropriate, 2 mM of an appropriate authentic standard in buffer (50 mM phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The singlet observed at ~4.4 ppm originates from the enzyme sample; it was observed in the enzyme-only control.




Supporting Figure 2.S48. Analysis of the reaction outcome of PsEFE catalyzed conversion of racemic 4-benzyl 2OG (15) using pure shift ^{1}H NMR. Analysis of the reaction of PsEFE with racemic 4-benzyl 2OG (**15**) 30 min post addition of PsEFE using: **a**) pure shift ^{1}H NMR (1, 5-7) and **b**) ^{1}H NMR results indicate that substantial amounts of reaction products other than 2-benzylsuccinate and P5C (Supporting Figures 2.S44-2.S46) are not being formed under the tested conditions. Analysis of the reaction mixture with **15** in the absence of PsEFE using: **c**) pure shift ^{1}H NMR (1, 5-7) and **d**) ^{1}H NMR. Conditions: 400 µM **15**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 30 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Note that for the pure shift ^{1}H NMR analysis, the observed peak may not necessarily be at the center of the peak obtained when using standard NMR analysis (1, 5-7).




Supporting Figure 2.S49. Analysis of PsEFE catalysis using racemic 3-ethyl-2OG (11) as a potential substrate. Representative ^{1}H NMR spectra monitoring the reaction of PsEFE with racemic 3-ethyl-2OG (**11**) after: **a**) overnight incubation and **b**) 5 min. **c**) ^{1}H NMR spectrum of **11** under standard conditions in the absence of PsEFE; **d**) ^{1}H NMR spectrum of **11** in buffer. Conditions: 400 µM **11**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 30 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 11%$_{v/v}$ D_2O). Formation of new peaks was not observed, even after incubating the reaction mixture overnight. Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S50. Analysis of PsEFE catalysis using racemic 3-propyl-2OG (12) as a potential substrate. Representative ^{1}H NMR spectra monitoring the reaction of PsEFE with racemic 3-propyl-2OG (**12**) after: **a**) overnight incubation and **b**) 5 min. **c**) ^{1}H NMR spectrum of **12** under standard conditions in the absence of PsEFE; **d**) ^{1}H NMR spectrum of **12** in buffer. Conditions: 400 µM **12**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 30 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Formation of new peaks was not observed, even after incubating the reaction mixture overnight. Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S51. Analysis of PsEFE catalysis using racemic 3-butyl-2OG (13) as a potential substrate. Representative ^{1}H NMR spectra monitoring the reaction of PsEFE with racemic 3-butyl-2OG (**13**) after: **a**) overnight incubation and **b**) 5 min. **c**) ^{1}H NMR spectrum of **13** under standard conditions in the absence of PsEFE; **d**) ^{1}H NMR spectrum of **13** in buffer. Conditions: 400 µM **13**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 30 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Formation of new peaks was not observed, even after incubating the reaction mixture overnight. Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S52. Analysis of PsEFE catalysis using racemic 3-benzyl-2OG (16) as a potential substrate. Representative ^{1}H NMR spectra monitoring the reaction of PsEFE with racemic 3-benzyl-2OG (**16**) after: **a**) overnight incubation and **b**) 5 min. **c**) ^{1}H NMR spectrum of **16** under standard conditions in the absence of PsEFE; **d**) ^{1}H NMR spectrum of **16** in buffer. Conditions: 400 µM **16**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 30 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Formation of new peaks was not observed, even after incubating the reaction mixture overnight. Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S53. Analysis of PsEFE catalysis using 2-(carboxycarbonyl)cyclopropane-1-carboxylate (17) as a potential substrate. Representative ^{1}H NMR spectra monitoring the reaction of PsEFE with 2-(carboxycarbonyl)cyclopropane-1-carboxylate (**17**) after: **a**) overnight incubation and **b**) 5 min. **c**) ^{1}H NMR spectrum of **17** under standard conditions in the absence of PsEFE; **d**) ^{1}H NMR spectrum of **17** in buffer. Conditions: 400 µM **17**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 30 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Formation of new peaks was not observed, even after incubating the reaction mixture overnight. Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S54. Analysis of PsEFE catalysis using 3-(carboxycarbonyl)cyclobutane-1-carboxylate (18) as a potential substrate. Representative ^{1}H NMR spectra monitoring the reaction of PsEFE with 3-(carboxycarbonyl)cyclobutane-1-carboxylate (**18**) after: **a**) overnight incubation and **b**) 5 min. **c**) ^{1}H NMR spectrum of **18** under standard conditions in the absence of PsEFE; **d**) ^{1}H NMR spectrum of **18** in buffer. Conditions: 400 µM **18**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 30 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Formation of new peaks was not observed, even after incubating the reaction mixture overnight. Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S55. Analysis of PsEFE catalysis using 3-(carboxycarbonyl)benzoate (19) as a potential substrate. Representative ^{1}H NMR spectra monitoring the reaction of PsEFE with 3-(carboxycarbonyl)benzoate (**19**) after: **a**) overnight incubation and **b**) 5 min. **c**) ^{1}H NMR spectrum of **19** under standard conditions in the absence of PsEFE; **d**) ^{1}H NMR spectrum of **19** in buffer. Conditions: 400 µM **19**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 30 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Formation of new peaks was not observed, even after incubating the reaction mixture overnight. Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.




Supporting Figure 2.S56. Analysis of PsEFE catalysis using 4-(carboxycarbonyl)bicyclo[2,2,2]octane-1-carboxylate (20) as a potential substrate. Representative ^{1}H NMR spectra monitoring the reaction of PsEFE with 4-(carboxycarbonyl)bicyclo[2,2,2]octane-1-carboxylate (**20)** after: **a**) overnight incubation and **b**) 5 min. **c**) ^{1}H NMR spectrum of **20** under standard conditions in the absence of PsEFE; **d**) ^{1}H NMR spectrum of **20** in buffer. Conditions: 400 µM **20**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe (II), 800 µM TMSP-d_4, and 30 µM PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Formation of new peaks was not observed, even after incubating the reaction mixture overnight. Both the doublet observed at ~1.2 ppm and the singlet observed at ~4.4 ppm originate from the enzyme sample; both signals were observed in the enzyme-only control.



Supporting Figure 2.S57. The product ratio of PsEFE catalysis with racemic 4-methyl 2OG (3) as a (co)substrate is not affected by the tested conditions. The 2-methylsuccinate (**3a**):β-hydroxyisobutyrate (**3b**) ratios (~1.8:1) estimated by ^{1}H NMR (600 MHz) analysis of the crude reaction mixtures do not appear to vary within the tested pH range: (**a**) pH 6.5, (**b**) pH 7.4, and (**c**) pH 8.5. The time scales were calibrated to the end of acquisition of the first ^{1}H NMR experiment post addition of PsEFE to the reaction mixture (t = 0 min), by which time low levels of conversion were manifest. (**d**) 2-Methylsuccinate (**3a**):β-hydroxyisobutyrate (**3b**) ratios. Conditions: 400 µM **3**, 500 µM L-arginine, 500 µM L-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C.













Supporting Figure 2.S58. The product ratio of PsEFE catalysis with racemic 3-methyl 2OG (10) as a (co)substrate is not affected by the tested conditions. The 2-methylsuccinate (**3a**):β-hydroxybutyrate (**10b**) ratios (~1.2:1) estimated by ^{1}H NMR (600 MHz) analysis of the crude reaction mixtures do not appear to vary within the tested pH range: (**a**) pH 6.5, (**b**) pH 7.4, and (**c**) pH 8.5. The time scales were calibrated to the end of acquisition of the first ^{1}H NMR experiment post addition of PsEFE to the reaction mixture (t = 0 min), by which time low levels of conversion were manifest. (**d**) 2-Methylsuccinate (**3a**):β-hydroxyisobutyrate (**3b**) ratios. Conditions: 400 µM **10**, 500 µM L-arginine, 500 µM L-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C.









Supporting Figure 2.S59. DNA and amino acid sequence of the SenP2 protease construct used during the production of PsEFE.

2.10.3 Supporting Tables

Supporting Table 2.S1. Product ratios of PsEFE-catalyzed reactions with 2OG derivatives.[a]

	2OG derivative	diacid:alkene ratio	diacid:alcohol ratio	conversion
i	 **2OG (1)**	succinate / ethylene: 1.0 ± 0.1:1	no alcohol product formed under the tested conditions	>95%
ii	 **2OA (2)**	no alkene product formed under the tested conditions	**2a** / **2b**: 2.1 ± 0.1 : 1	~60%
iii	 **3**	no alkene product formed under the tested conditions	**3a** / **3b**: 1.8 ± 0.1 : 1	>95%
iv	 **4**	no alkene product formed under the tested conditions	**4a** / **4b**: 2.3 ± 0.1 : 1	~90%
v	 **5**	no alkene product formed under the tested conditions	**5a** / **5b**: 1.2 ± 0.1 : 1	~30%
vi	 **6**	no alkene product formed under the tested conditions	**6a** / **6b**: 1.3 ± 0.1 : 1	~80%
vii	 **7**	no alkene product formed under the tested conditions	**7a** / **7b**: 3.2 ± 0.1 : 1	>95%
viii	 **8**	no alkene product formed under the tested conditions	**8a** / **8b**: 4.1 ± 0.1 : 1	~70%
ix	 **9**	no alkene product formed under the tested conditions	**9a** / **9b**: ~1 : 0	~50%
x	 **10**	no alkene product formed under the tested conditions	**3a** / **10b**: 1.2 ± 0.1 : 1	>75%

	2OG derivative	diacid:alkene ratio	diacid:alcohol ratio		conversion
[b]xi	 **14** ***trans*:*cis*, 5:1**	no alkene product formed under the tested conditions	 **14a** ***isomeric mixture***	 **14b** ***isomeric mixture***	[c]not determined
			~1 : 0		
xii	 **15**	no alkene product formed under the tested conditions	 **15a**	 **15b**	[d]not determined
			~1 : 0		

a) 2OG derivatives were typically prepared from cyanosulfur ylids as racemic mixtures as reported (2). Note, there is a possibility for stereoselective conversion; b) **14** was used as a mixture of diastereomers, dr (*trans*:*cis*) = 5:1 (2); c) the identity of the diacid reaction product was confirmed by spiking the reaction mixture with a synthetic sample of **14a**. Levels of conversion could not be accurately determined by ^{1}H NMR due to overlapping signals of starting material (**14**) and product (**14a**). Spiking with the potential corresponding authentic alcohol product (**14b**) revealed that it is not formed, within limits of detection. **14a** and **14b** were commercially obtained as a mixture of *cis* and *trans* isomers; d) the identity of the diacid reaction product was confirmed by spiking the reaction mixture with a synthetic sample of **15a**. Levels of conversion could not be accurately determined by ^{1}H NMR due to overlapping signals of starting material (**15**) and product (**15a**). Spiking with the potential corresponding authentic alcohol product (**15b**) revealed that it is not formed, within limits of detection. Conditions: 400 µM 2-oxoacid, 500 µM L-arginine, 500 µM L-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, 2 µM psEFE (for 2OG) or 10-30 µM PsEFE (for 2OA and 2OG derivatives) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O) at 25 °C. The results are the mean of independent triplicates (n = 3; mean ± SD).

2.10.4 Supporting Information References

1. Gates, E. L., Bradley, J. P., Berry, D. B. G., Nilsson, M., Morris, G. A., Adams, R. W., and Castañar, L. (2024) Solvent suppression in pure shift NMR. *Anal. Chem.* **96**, 3879-3885

2. Brewitz, L., Nakashima, Y., and Schofield, C. J. (2021) Synthesis of 2-oxoglutarate derivatives and their evaluation as cosubstrates and inhibitors of human aspartate/asparagine-β-hydroxylase. *Chem. Sci.* **12**, 1327-1342

3. Adams, R. W., Holroyd, C. M., Aguilar, J. A., Nilsson, M., and Morris, G. A. (2013) "Perfecting" WATERGATE: clean proton NMR spectra from aqueous solution. *Chem. Commun.* **49**, 358-360

4. Pohl, L., and Eckle, M. (1969) Sodium 3-trimethylsilyltetradeuteriopropionate, a new water-soluble standard for ^{1}H-NMR. *Angew. Chem. Int. Ed.* **8**, 381

5. Castañar, L. (2017) Pure shift ^{1}H NMR: what is next? *Magn. Reson. Chem.* **55**, 47-53

6. Foroozandeh, M., Morris, G. A., and Nilsson, M. (2018) PSYCHE pure shift NMR spectroscopy. *Chem. Eur. J.* **24**, 13988-14000


7. Zangger, K. (2015) Pure shift NMR. *Prog. Nucl. Magn. Reson. Spectrosc.* **86-87**, 1-20

8. Wishart, D. S., Knox, C., Guo, A. C., Eisner, R., Young, N., Gautam, B., et al. (2009) HMDB: a knowledgebase for the human metabolome. *Nucleic Acids Res.* **37**, D603-D610

2.11 Statement of Authorship

Title of Paper	**Substitution of 2-oxoglutarate alters reaction outcomes of the *Pseudomonas savastanoi* ethylene-forming enzyme**
Publication Status	X Published □ Accepted for Publication □Submitted for Publication work □ Unpublished and unsubmitted work written in a manuscript style
Publication Details	Dhingra, S.; Zhang, Z.; Lohans, C. T.; Brewitz, L.; Schofield, C. J. Substitution of 2-oxoglutarate alters reaction outcomes of the *Pseudomonas savastanoi* ethylene-forming enzyme. *J. Biol. Chem.* **2024**, 300, 107546

Student Confirmation

Student Name:	Siddhant Dhingra		
Contribution to the Paper	I conducted all the experimental work, data analysis, and data visualisation described here. The manuscript was prepared by myself and Dr. Lennart Brewitz under the guidance of Prof. Christopher J. Schofield. Dr. Zhihong Zhang provided the PsEFE plasmid and the initial training for protein expression and purification. Dr. Christopher Lohans, Dr. Lennart Brewitz, and Prof. Christopher J. Schofield conceptualised the project.		
Signature 		Date	05-10-2025

Supervisor Confirmation

By signing the Statement of Authorship, you are certifying that the candidate made a substantial contribution to the publication, and that the description described above is accurate.

Supervisor name and title: Prof. Christopher J. Schofield		
Supervisor comments I'm happy to state that the candidate made the major and a very substantial contribution to the publication, and that the description described above is accurate.		
Signature C. J. Schofield	Date	25th September 2025

CHAPTER 3

MANUSCRIPT II

Structural and functional validation of *Pseudomonas savastanoi* ethylene forming enzymes reveals flexibility in 2-oxoglutarate binding mode and conformation

Contents

3.1 Overview 164
3.2 Table Contents Graphic 165
3.3 Abstract 165
3.4 Introduction 166
3.5 Results 169
3.5.1 Comparison of PsEFE sequences 169
3.5.2 The *glycinea* and 1449B PsEFEs catalyse ethylene production from 2OG 170
3.5.3 The *glycinea* and 1449B PsEFEs can accept 2OG derivatives as (co)substrates 173
3.5.4 Crystallography 176
3.6 Discussion 182
3.7 Conclusion 185
3.8 Experimental Procedures 185
3.9 Acknowledgements 187
3.10 Author Contributions 187
3.11 References 188
3.12 Supporting Information 191
3.12.1 Supporting Figures 191
3.12.2 Supporting Tables 201
3.12.3 Supporting Information References 204
3.13 Statement of Authorship 205

3.1 Overview

Pseudomonas savastanoi pv *phaseolicola* PK2 EFE has been extensively studied (Section 1.7, Chapter 2); EFEs from other *Pseudomonas* pathovars and strains, however, have been less explored. This chapter contains a manuscript describing biochemical and structural studies on two putative EFE orthologues from *Pseudomonas savastanoi* pv *phaseolicola* 1449B (1449B PsEFE) and *Pseudomonas savastanoi* pv *glycinea* (*glycinea* PsEFE). Structural features of PK2 and 1449B PsEFEs are compared.

^{1}H NMR assays validated the assignment of both of the studied PsEFE orthologues as ethylene-forming enzymes, however, with variable catalytic efficiencies and partition ratios. The *glycinea* PsEFE, like PK2 PsEFE, produces ethylene and succinic acid in approximately equal proportions, while 1449B PsEFE was found to favour ethylene formation relative to PK2 PsEFE. A high-resolution crystal structure of 1449B in complex with 2OG (PDB ID: 9SCF), obtained during these studies led to the observation that 2OG adapts an atypical binding mode, including with respect to Mn(II) coordination. This observation is in accord with previously reported crystal structures of 2OG oxygenases in complex with Mn(II), including PsEFE, showing atypical binding modes and conformations with respect to metal coordination by 2-oxoacids. Collectively, studies described in this chapter further our understanding on how sequence variations, despite being distant from the active site, can influence catalytic efficiency and product outcomes of ethylene-forming enzymes.


At the time of writing this thesis, Manuscript II has been submitted to *ChemBioChem* and is under review.

Sun, Y.; Dhingra, S.; Allen, M. D.; Cheng, E. S. Z.; Zhang, Z.; Schofield, C. J.; Brewitz, L. Structural and functional validation of *Pseudomonas savastanoi* ethylene forming enzymes

reveals flexibility in 2-oxoglutarate binding mode and conformation **Manuscript under review in** *ChemBioChem*.

Each author's contribution is described in the Author Contributions (Section 3.10). A Statement of Authorship is appended at the end of the chapter (Section 3.13).

3.2 Table Contents Graphic



3.3 Abstract

Pseudomonas savastanoi pv. *phaseolicola* PK2 employs an Fe(II)-dependent ethylene/succinate-forming enzyme (PK2 PsEFE) to produce ethylene from 2-oxoglutarate (2OG). Here we report NMR-based assays showing that the putative *P. savastanoi* pv. *glycinea* PsEFE, which differs from PK2 PsEFE by a single residue, and the *P. savastanoi* pv.1449B PsEFE, which differs from PK2 PsEFE by 28 residues and a C-terminal 13-residue truncation, catalyse ethylene production from 2OG. Like the PK2 PsEFE, they catalyse oxidation of naturally occurring 2OG derivatives to give alcohol and diacid products. Crystallographic analysis demonstrates that the overall fold and active site of 1449B PsEFE is similar to that of PK2 PsEFE. Interestingly, 2OG was observed to adopt an atypical inverse metal ion binding mode in complex with 1449B PsEFE and Mn in which its 2-oxoacid group is positioned to interact with the guanidinium group of R277, but not the Mn ion (substituting for catalytically active Fe(II)). Together with reported crystallographic results, this observation indicates that 2OG metal ion binding modes and conformations at the active sites of 2OG oxygenases can vary, possibly in a functionally relevant manner.

3.4 Introduction

Ethylene is produced by plants, fungi, and bacteria.[1] In seeding plants, 1-aminocyclopropane-1-carboxylate (ACC) oxidase (ACCO) catalyses ethylene production from ACC.[2] By contrast, organisms other than seeding plants do not typically employ ACCOs for ethylene production, but use mechanistically distinct enzymes which do not employ ACC as a substrate,[3] including: (i) nitrogenases,[3a-f] (ii) nitrogenase-like enzymes,[3g-i] and (iii) ethylene/succinate-forming enzymes (EFEs).[3j-m] Nitrogenases catalyse ethylene production from *inter alia* CO and acetylene,[3a-f] while the nitrogenase-like enzyme methylthio-alkane reductase (Mar) catalyses ethylene production from methionol.[3g-i] EFEs catalyse oxidation of 2-oxoglutarate (2OG) to give ethylene and CO_2 (Figure 3.1);[3l,3m] the EFE of *Pseudomonas savastanoi* pv. *phaseolicola* PK2 (PK2 PsEFE) requires *L*-arginine for this reaction to proceed efficiently.[4]

PK2 PsEFE also catalyses the competing oxidation of 2OG to give succinic acid and CO_2, a reaction coupled to the C5 hydroxylation of *L*-arginine; 5-hydroxyarginine fragments to give guanidine and *L*-Δ^1-pyrroline-5-carboxylic acid (P5C).[3j,5] The PK2 PsEFE (co)substrate scope extends to C3- and/or C4-substituted 2OG derivatives which can sustain catalysis in the absence of 2OG to give a diacid product and an alcohol product, but, at least in most cases, not an alkene product.[6]

PK2 PsEFE belongs to the Fe(II)-dependent 2OG oxygenase structural superfamily which normally employ an HXD/E…H triad of ligands embedded within a distorted double-stranded β-helix (DSBH) fold to bind the Fe(II) cofactor (Figure 3.1).[7] At 2OG oxygenase active sites, 2OG typically adopts an Fe(II) binding mode in which it chelates to the Fe(II) by its C1 carboxylate and C2 ketone groups, with its C5 carboxylate engaging in interactions with residues distal to the Fe(II) binding HXD/E…H triad.[5b,8] 2OG oxygenases normally catalyse 2 electron substrate oxidations, most commonly hydroxylation or *N*-methyl demethylation reactions (via initial hydroxylation), coupled to the oxidative decarboxylation of 2OG to give

succinic acid.[8b] Note that PK2 PsEFE bears a relatively apolar active site, a feature possibly contributing to its unusual reaction with 2OG to give ethylene (Figure 3.1).[9]




Figure 3.1. PK2 PsEFE catalysis and structure. (**a**) Reactions of PK2 PsEFE with 2OG. (**b**) The DSBH fold (light green) of PK2 PsEFE observed in a reported PK2 PsEFE:Mn:2OG structure (PDB ID: 5V2X[9b]); R326 is in purple (see below). (**c**) Active site view of a reported PK2 PsEFE:Mn:2OG structure (PDB ID: 5V2X[9b]) showing that, in addition to the typical binding mode (green), 2OG can adopt an atypical inverted metal ion binding mode (yellow), in the latter case coordinating to Mn via its C5 carboxylate rather than its 2-oxoacid group. (**d**) Active site view of a PK2 PsEFE:Mn:2OG:*L*-arginine structure (PDB ID: 5V2Y[9b]) showing that, in the presence of *L*-arginine, 2OG adopts a metal ion binding mode in which it coordinates to Mn via its 2-oxoacid group, as typically observed with 2OG oxygenases.[5b,8]

Reported crystal structures of PK2 PsEFE in complex with 2OG and catalytically inert Mn(II), substituting for catalytically active Fe(II), indicate that 2OG can adopt different metal ion binding modes at the active site. In one structure, the 2OG is positioned to coordinate Mn via

its 2-oxoacid group and to interact via its C5 carboxylate group with the guanidinium group of R277 via a salt-bridge (Figure 3.1),[9b] in a manner typically observed with 2OG oxygenases.[5b,8] In a structure obtained in the absence of *L*-arginine, 2OG was reported to additionally adopt an alternative metal ion binding mode (~50% occupancy) in which it is positioned to interact with Mn via its C5 carboxylate group and to interact with the guanidinium group of R277 with its 2-oxoacid group (Figure 3.1c).[9b]

P. syringae and *P. savastanoi* strains apparently carry the *EFE* gene in form of plasmid rather than chromosomal DNA;[10] even closely related *P. syringae* strains can differ in ethylene-producing capacity.[11] Pioneering work on PK2 PsEFE variants has revealed that minor sequence variations can ablate their ethylene producing capacity (Supporting Table 3.S1), including single residue variations both within and outside the active site.[6a,9,12] Thus, the ethylene-producing capacity of enzymes which have been assigned as EFEs based on their structural and/or sequence similarity with PK2 PsEFE should be experimentally validated by investigating the reactivity of the corresponding isolated enzymes.[3k,13]

Here we describe spectroscopic studies on the predicted EFEs of the *P. savastanoi* strains pv. *glycinea* and 1449B (hereafter, *glycinea* PsEFE and 1449B PsEFE) revealing that they catalyse ethylene production from 2OG, as reported for PK2 PsEFE.[3l,3m] The reactivities of the *glycinea* and 1449B PsEFEs with naturally occurring 2OG derivatives are similar to that of PK2 PsEFE,[6] as is the crystallographically observed fold of 1449B PsEFE.[9] The combined results validate the assignment of *glycinea* and 1449B PsEFE as ethylene-producing enzymes.

3.5 Results

3.5.1 Comparison of PsEFE sequences

We investigated the ethylene-producing capacity of predicted EFEs from two *P. savastanoi* strains, *i.e.*, *P. savastanoi* pv. *glycinea* (*glycinea* PsEFE; UNIPROT ID: Q7BS32) and pv. 1449B (1449B PsEFE; UNIPROT ID: Q9Z3T0), which differ by 1 residue (R326S; *glycinea* PsEFE) and which have ~88% sequence similarity (1449B PsEFE), respectively, compared with the validated ethylene-producing PK2 PsEFE[3l,3m] (UNIPROT ID: P32021; EC:1.13.12.19) (Figure 3.2), because the ethylene-producing capacity of *Pseudomonas* strains differs[11] and because sequence variations can affect the ethylene producing capacity of PK2 PsEFE (Supporting Table 3.S1).[6a,9,12]

The R326S variation is located on the surface of the *glycinea* PsEFE (Figure 3.1b), far from the active site, and appears to be conserved in different *P. savastanoi* and *P. syringae* strains.[9b,11b] To our knowledge, the effect of the R326S variation on catalysis of isolated recombinant PK2 PsEFE has not been defined (Supporting Table 3.S1). 1449B PsEFE lacks the 13 C-terminal residues of the PK2 and *glycinea* PsEFEs and differs by 28 additional residues from PK2 PsEFE: the 28 substitutions involve residues both close to and distant the active site (Figure 3.2). Note that the 1449B PsEFE sequence is identical to that of the *P. syringae* pv. *pisi* GSPB1206 EFE.[9b,11b] The sequences of the *glycinea*, 1449B, and PK2 PsEFEs differ substantially from those of functionally and structurally validated Fe(II)-dependent ethylene-producing enzymes of other organisms, including the EFE of *Penicillium digitatum*[13b,14] as well as *Arabidopsis thaliana* and *Petunia hybrida* ACCO1 (Supporting Figure 3.S1).[15]




Figure 3.2. Comparison of EFE sequences from *P. savastanoi* pv. *phaseolicola* PK2, pv. *glycinea*, and pv. 1449B. α-Helices and β-strands of PK2 PsEFE are numbered based on a PK2 PsEFE structure (PDB ID: 5MOF[9a]); β-turns: TT. Cyan triangles: β-strands forming the DSBH fold of PK2 PsEFE; green circles: the PK2 PsEFE HXD…H Fe(II)-binding motif; blue circles: polar PK2 PsEFE residues crystallographically observed to directly interact with 2OG; purple pentagon: the R326S variation distinguishes the *glycinea* PsEFE from the PK2 PsEFE; brown squares: 28 residues distinguishing the 1449B PsEFE from the PK2 PsEFE; orange rectangle: 13 C-terminal PK2 PsEFE residues (338-350) missing in 1449B PsEFE. Identical residues are in red; residues of similar polarity are in yellow. The Figure 3.2 was generated using ESPript 3.0.[16]

3.5.2 The *glycinea* and 1449B PsEFEs catalyse ethylene production from 2OG

The isolated *glycinea* and 1449B PsEFEs were recombinantly produced in *E. coli* cells and purified to high purity (>90% by SDS-PAGE and mass spectrometric analyses; Supporting Figure 3.S2). To investigate their ethylene-producing capabilities, we used ^{1}H NMR assays, which have been employed to monitor ethylene formation/2OG depletion by PK2 PsEFE.[6b,9a]

^{1}H NMR analysis (700 MHz) of the reaction of the isolated *glycinea* and 1449B PsEFEs with

2OG reveals that both catalyse ethylene production from 2OG in the presence of *L*-arginine, under conditions reported to give ethylene with PK2 PsEFE (Figure 3.3),[6b] confirming their assignment as EFEs. The *glycinea* and 1449B PsEFEs also catalysed oxidation of 2OG to succinic acid in a reaction likely coupled to C5 oxidation of *L*-arginine to give P5C, as implied by formation of the characteristic P5C singlet at ~7.70 ppm (Figure 3.3). The *glycinea* and 1449B PsEFEs did not catalyse efficient 2OG oxidation in the absence of *L*-arginine (Supporting Figure 3.S3).




Figure 3.3. The *glycinea* and 1449B PsEFEs catalyse ethylene production. (**a**) Reactions of PK2, *glycinea*, and 1449B PsEFE with 2OG. (**b**) Outcomes of the reaction of 1449B and *glycinea* PsEFE with 2OG compared to that reported for PK2 PsEFE.[6b] Ethylene formation was estimated by subtracting the integral (normalized to TMSP-d_4) of succinic acid from that of 2OG. Note that we did not observe 3-hydroxypropanoic acid formation by ^{1}H NMR.[12a,13a] (**c-e**) ^{1}H NMR spectra (~0.8 to ~4.4 ppm and ~5.4 to ~8.0 ppm) of a 2OG reaction mixture, ~90 min post addition of: (**c**) 1449B PsEFE or (**d**) *glycinea* PsEFE, compared to that reported for: (**e**) PK2 PsEFE[6b]; (**f**) ^{1}H NMR spectrum 2OG in the absence of PsEFE. Conditions: 400 µM 2OG, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and PsEFE (2 µM for PK2;[6b] 5 µM for 1449B; 10 µM for *glycinea*) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Note that low levels of dehydroascorbate (~4.3 and ~4.2 ppm),[6b] succinic acid (~2.4 ppm), and the hydrated ketone form of 2OG (~2.0 and ~2.2 ppm, dashed circles)[17] were observed. The peak at ~3.75 ppm (labelled: *) corresponds to Tris. Time scales were calibrated to the end of the acquisition of the first ^{1}H NMR experiment following PsEFE addition to the reaction mixture (t = 0 min), by which time low levels of turnover were manifest. Results are means of independent triplicates (n = 3; mean ± standard deviation, SD).

The *glycinea* and 1449B PsEFEs were not observed to catalyse formation of 2OG-derived products other than succinic acid and ethylene (Figure 3.3), in accord with our reported ^{1}H NMR observations on the reactivity of PK2 PsEFE with 2OG.[6b,9a] This observation contrasts with the reported ability of PK2 PsEFE and related EFEs to catalyse formation of low levels of 3-hydroxypropionic acid, possibly because of different assay conditions and/or because ^{1}H NMR-based assays are less sensitive than LCMS/infrared-based assays previously used to detect 3-hydroxypropionic acid.[12a,13]

Specific activities were calculated for the PK2, 1449B, and *glycinea* PsEFEs by analysing 2OG depletion ~10 min post incubation with PsEFE. The results reveal that the specific activities decrease in the order: PK2 PsEFE > 1449B PsEFE > *glycinea* PsEFE, assuming that the purified PsEFEs were fully active (Table 3.1). Although ethylene formation was not directly quantified because ethylene may evaporate from the reaction mixture, time-course studies measuring 2OG depletion and succinic acid formation imply that ethylene was continuously formed from 2OG during the reaction with *glycinea* and 1449B PsEFE (Supporting Figures 3.S4-3.S5). The ethylene:succinic acid ratios were estimated to be ~3:2 for 1449B PsEFE, but ~1:1 for both glycinea and PK2 PsEFE (Figure 3.3b).

3.5.3 The *glycinea* and 1449B PsEFEs can accept 2OG derivatives as (co)substrates

The reactivities of the *glycinea* and 1449B PsEFEs with four naturally occurring 2OG derivatives, *i.e.*, 2-oxoadipate (2OA, **2**),[18] 4-methyl-2OG (**3**),[19] 3-methyl-2OG (**4**),[20] and 4-hydroxy-2OG (**5**),[21] were tested to compare the reaction outcomes with those reported for PK2 PsEFE.[6b] ^{1}H NMR analyses (700 MHz) reveal that the *glycinea* and 1449B PsEFEs catalysed formation of a diacid and an alcohol product from all the four tested 2OG derivatives (Figure 3.4, Supporting Figures 3.S4-3.S9), as also observed for PK2 PsEFE.[6] The estimated

diacid:alcohol ratios of the reactions of the 1449B and *glycinea* PsEFEs with 2OG derivatives were similar within experimental error and did not differ substantially from those reported for PK2 PsEFE (Figure 3.4).[6b] As observed with 2OG (Figure 3.3), the specific activities of the three PsEFEs with 2OG derivatives decreased in the order: PK2 PsEFE > 1449B PsEFE > *glycinea* PsEFE (Table 3.1). Note that the concentration of *glycinea* PsEFE in the ^{1}H NMR assays with the 2OG derivatives was 3-fold that of PK2 and 1449B PsEFE to enable robust product formation.

Table 3.1. Specific activities and product ratios of the PK2, 1449B, and *glycinea* PsEFEs with 2OG and 2OG derivatives.

2OG derivative	[a]Specific activities [nmol·min^{-1}·mg^{-1}]			[b]Ethylene:succinic acid ratios (for 2OG) or [c]diacid:alcohol ratios		
	PK2 PsEFE	1449B PsEFE	*glycinea* PsEFE	PK2 PsEFE	1449B PsEFE	*glycinea* PsEFE
[b]2OG	357 ± 26	136 ± 4	17 ± 6	1.0 ± 0.1	1.5 ± 0.1	1.2 ± 0.1
[c]2OA	52 ± 1	23 ± 14	2 ± 1	1.2 ± 0.1	1.3 ± 0.1	1.2 ± 0.1
[c]4-Methyl-2OG (**3**)	72 ± 4	46 ± 7	3 ± 1	1.9 ± 0.3	1.3 ± 0.3	1.3 ± 0.2
[c]3-Methyl-2OG (**4**)	62 ± 1	36 ± 3	8 ± 1	1.2 ± 0.1	1.4 ± 0.1	0.9 ± 0.1
[c]4-Hydroxy-2OG (**5**)	69 ± 2	10 ± 4	3 ± 2	2.2 ± 0.3	2.6 ± 0.9	1.4 ± 0.4

a) Specific activities (nmol·min^{-1}·mg^{-1}) based on 2OG/2OG derivative depletion after ~10 min incubations with PsEFE. b) Ethylene:succinic acid ratios ~90 min post incubation with PsEFE. c) Diacid:alcohol ratios ~90 min post incubation with PsEFE. ^{1}H NMR assay conditions: 400 μM 2OG or 2OG derivative, 500 μM *L*-arginine, 500 μM *L*-ascorbic acid, 50 μM Fe(II), 800 μM TMSP-d_4, and PsEFE (using: 2 μM PK2, 5 μM 1449B, and 10 μM *glycinea* PsEFE with 2OG; or using: 10 μM PK2 or 1449B and 30 μM *glycinea* PsEFE with 2OG derivatives) in buffer (50 mM sodium phosphate, pH 7.4, 10% $_{v/v}$ D_2O). Results are means of independent triplicates (n = 3; mean ± SD).




Figure 3.4. The *glycinea* and 1449B PsEFEs catalyse oxidation of 2OG derivatives. Outcomes of the reactions of the *glycinea* and 1449B PsEFEs with: (**a**) 2-oxoadipate (2OA, **2**), (**b**) 4-methyl-2OG (**3**), (**c**) 3-methyl-2OG (**4**), and (**d**) 4-hydroxy-2OG (**5**), compared to those reported for PK2 PsEFE.[6b] Conditions: 400 μM 2OG derivative, 500 μM *L*-arginine, 500 μM *L*-ascorbic acid, 50 μM Fe(II), 800 μM TMSP-d_4, and PsEFE (PK2[6b] and 1449B PsEFE: 10 μM; *glycinea* PsEFE: 30 μM) in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). Results are means of independent triplicates (n = 3; mean ± SD). Note that the 2OG derivatives **3**, **4**, and **5** were used as racemic mixtures; there is potential for stereoselectivity in their reactions, including that one enantiomer may be a PsEFE substrate and the other an inhibitor and/or that one enantiomer favours a particular PsEFE reaction pathway.

3.5.4 Crystallography

1449B PsEFE was crystallized in the presence of 2OG and Mn(II), substituting for catalytically active Fe(II). 1449B PsEFE crystallized in the P43 space group (1.65 Å resolution; Figure 3.5 and Supporting Table 3.S2), whereas PK2 PsEFE crystallized in different space groups.[9,12b,22]

The 1449B PsEFE:Mn:2OG complex crystal structure was solved by molecular replacement using a reported PK2 PsEFE structure as a search model (PDB ID: 5MOF[9a]). Superimposition of the 1449B PsEFE:Mn:2OG structure and a reported PK2 PsEFE:Mn:2OG structure (PDB ID: 5MOF[9a]) reveal a similar PsEFE fold (RMSD: 0.415 Å) (Supporting Figure 3.S10) and active site geometry, in accord with the ability of 1449B PsEFE to catalyse conversion of 2OG to ethylene (Figure 3.3).




Figure 3.5. 2OG adopts an inverted binding mode in complex with 1449B PsEFE and Mn. (**a**) Locations of 1449B PsEFE residues in the 1449B PsEFE:Mn:2OG crystal structure (PDB ID: 9SCF) that differ from PK2 PsEFE (in red; Figure 3.2); the corresponding PK2 PsEFE residues are in black parentheses; the 1449B PsEFE DSBH is in cyan. (**b**) Representative Polder omit electron density map contoured to 3σ around 2OG in the 1449B PsEFE:Mn:2OG structure. 2OG coordinates to Mn and interacts with the R171 guanidinium group with its C5 carboxylate; it interacts with the R277 guanidinium group via its C1 carboxylate and C2 ketone. (**c**) Active site view of a PK2 PsEFE:Mn:Cl:2OG structure (PDB ID: 5MOF[9a]). (**d**) Superimposition of active site views from the 1449B PsEFE:Mn:2OG (PDB ID: 9SCF) and PK2 PsEFE:Mn:2OG:Cl (PDB ID: 5MOF[9a]) structures reveal different 2OG metal ion binding modes. (**e**) Active site view of a reported PK2 PsEFE:Mn:2OG structure (PDB ID: 5V2X[9b]). (**f** and **g**) Superimposition of views from the 1449B PsEFE:Mn:2OG (PDB ID: 9SCF) and PK2 PsEFE:Mn:2OG (PDB ID: 5V2X[9b]) structures. Note that 2OG adopts two binding modes in the PK2 PsEFE:Mn:2OG structure, with Binding Mode A shown in: (**f**) and

Binding Mode B shown in: (**g**). Binding Mode A and B differ from that observed in the 1449B PsEFE:Mn:2OG structure. (**h**) Superimposition of active site views from the 1449B PsEFE:Mn:2OG (PDB ID: 9SCF) and PK2 PsEFE:Mn:2OG:*L*-arginine (PK2 PsEFE: orange; Mn: violet; carbon-backbone of 2OG: turquoise; carbon-backbone of *L*-arginine: salmon; PDB ID: 5V2Y[9b]) structures imply *L*-arginine binding affects the 2OG conformation.

Analysis of electron density maps for the 1449B PsEFE:Mn:2OG structure reveals both 2OG and Mn bound at the active site with 2OG adopting an inverted metal ion binding mode compared to that typically observed for 2OG.[5b,8-9] In this structure, the 2-oxoacid group is positioned to interact with the guanidinium group of R277 via its C1 carboxylate and C2 ketone groups (2.5 and 3.0 Å, respectively) rather than with the metal ion, as typically observed.[5b,8] Mn is instead complexed by one of the two 2OG C5 carboxylate oxygen atoms (2.2 Å), whereas the other 2OG C5 carboxylate oxygen atom is positioned to interact with the guanidinium group of R171 (2.8 Å) (Figure 3.5). No evidence for the typical 2OG binding mode was apparent, but the presence of low levels of this cannot be excluded. As anticipated, Mn is coordinated by the side chain imidazole rings of H189 and H268 (2.2 Å), the side chain carboxylate of D191 (2.2 Å),[7a] and the water molecules W1 and W2 (1.6 and 2.0 Å, respectively) (Figure 3.5b).

The two Mn-bound water molecules W1 and W2 are part of a water network which extends across the active site (Figure 3.6a). The water network links the 2OG C1 and C5 carboxylate groups, the D191 side chain carboxylate, and the Y192 and Y318 side chain alcohols (Y192 is involved in binding of the *L*-arginine substrate, as observed in the PK2 PsEFE:Mn:2OG:*L*-arginine (PDB ID: 5V2Y[9b]) structure) with Q37, T241, V246, Q200, and T248, with some of latter residues being distal to the 1449B PsEFE active site. The positions of water molecules in and proximal to the active site is similar, but not identical to those observed in reported PK2 PsEFE structures (Figure 3.6d).[9] Note that in the PK2 PsEFE:Mn:Cl:2OG structure (PDB ID: 5MOF[9a]) a chloride ion was modelled at the 2OG binding site (~30% occupancy), binding of

which is possibly stabilized by interactions with water molecules (Figure 3.5c-d).[9a] PsEFE turnover assays are typically performed in the absence of high chloride concentrations,[3k,4,6,9,12a,12b,13,22-23] though chloride ions from the enzyme sample are present and do not ablate activity.[6b,23]

The 2OG binding mode in the 1449B PsEFE:Mn:2OG structure was similar, but not identical, to the atypical inverted 2OG binding mode (Binding Mode B; Figure 3.5g) observed in a reported PK2 PsEFE:Mn:2OG structure (PDB ID: 5V2X[9b]), along with the typical 2OG binding mode (Binding Mode A; Figure 3.5f). Notably, in our structure and in the reported Binding Mode B, 2OG adopts a C2/C5 *anti*-periplanar conformation (Figure 3.5b and g), whereas it adopts a C2/C5 *gauche* conformation in reported Binding Mode A (Figure 3.5f). The PK2 PsEFE:Mn:Cl:2OG (PDB ID: 5MOF[9a]) and PK2 PsEFE:Mn:2OG:*L*-arginine (PDB ID: 5V2Y[9b]) structures reveal a substantially different 2OG binding mode, in which it coordinates to Mn via its 2-oxoacid group, as typically observed in 2OG oxygenase crystal structures in complex with 2OG (Figure 3.5h).[5b,8] Notably, the 2OG binding mode in the 1449B PsEFE:Mn:2OG structure also differed from that observed in a reported *Penicillium digitatum* EFE structure in complex with Mn and 2OG (PDB ID: 9EIS[13b]), in which only the 2OG C1 carboxylate group, but not the 2OG C2 ketone group, was modelled in a position to form a salt-bridge with the guanidinium group of R336, the R277-equivalent residue in the *Penicillium digitatum* EFE.[13b]




Figure 3.6. An extensive water network stabilises 2OG binding in PsEFE:Mn:2OG complex crystal structures in the absence of *L*-arginine. (**a**) Multiple water molecules are within hydrogen bonding distance (< 3.2 Å) of 2OG in the 1449B PsEFE:Mn:2OG structure, apparently stabilising 2OG binding. (**b** and **c**) Close up views of the waters in proximity of: (**b**) the 2OG C1 carboxylate and (**c**) the 2OG C5 carboxylate in the 1449B PsEFE:Mn:2OG structure. (**d**) A water network similar to that at the 1449B PsEFE active site was observed in the PK2 PsEFE:Mn:2OG structure (PDB ID: 5V2X[9b]). (**e**) The PK2 PsEFE:Mn:2OG:*L*-arginine structure (PDB ID: 5V2Y[9b]) reveals that the presence of *L*-arginine at the PsEFE active site alters the water network.

3.6 Discussion

Analysis of specific activities indicates that the *glycinea* PsEFE and the 1449B PsEFEs are less efficient in catalysing 2OG depletion and ethylene production compared to PK2 PsEFE (Table 3.1), in accord with reported studies showing that the R326S PK2 PsEFE variation reduced levels of ethylene production in cell extracts of different *Pseudomonas* species (Supporting Table 3.S1).[11b] The *glycinea* PsEFE S326 residue is located on the surface of the protein relatively close to the C-terminal region, which has been shown to affect 2OG oxygenases catalysis, at least in some cases, in a manner involving conformational changes.[24]

Notably, although the R326S PK2 PsEFE variation is conserved amongst several *Pseudomonas* pathovars,[9b,11b] work with intact cells implies that these *Pseudomonas* pathovars produce variable levels of ethylene, despite bearing PsEFEs with identical sequences.[25] Results with cell extracts suggested that the 1449B PsEFE is less efficient in producing ethylene than the *glycinea* PsEFE;[11b] by contrast, our work with isolated enzymes indicates that the 1449B PsEFE is more efficient in catalysing ethylene production than the *glycinea* PsEFE (Figure 3.3; Table 3.1). Thus, the combined results suggest that ethylene production in *Pseudomonas* pathovars does not only depend on the PsEFE sequence, but also on other factors.

The *glycinea* and PK2 PsEFEs produced ethylene and succinic acid in approximately equal ratios under the tested conditions (Figure 3.3). By contrast, 1449B PsEFE apparently favoured ethylene over succinic acid production (Figure 3.3), possibly reflecting the lower sequence similarity of 1449B and PK2 PsEFE (Figure 3.2). Although functional and structural studies on 2OG oxygenases have revealed that the C-terminus of some 2OG oxygenases is proximal to the active site and can affect catalysis,[24] our work supports reported results showing that the 13 C-terminal PK2 PsEFE residues are not required for ethylene production and *L*-arginine hydroxylation (Figure 3.3).[9b] The combined results thus suggest that one or more of the 28 additional residues distinguishing the 1449B and PK2 PsEFEs affect the ethylene:succinic acid

ratio, which is reported to be ~1:1 for the PK2 PsEFE variant lacking the 13 C-terminal residues (as for wildtype PK2 PsEFE),[6b,9b] but which is ~3:2 for 1449B PsEFE (Figure 3.3). As many of the 28 residues distinguishing the 1449B and PK2 PsEFEs are not at the active site (Figure 3.5), they may influence second coordination sphere interactions, possibly involving the active site water network (Figure 3.6).[26]

The observation that the PsEFE sequence affects the ethylene:succinic acid ratio is precedented by work on PsEFE-related enzymes, including the *Penicillium digitatum* EFE which features ~55% sequence similarity with PK2 PsEFE (Supporting Figure 3.S1) and which also favours ethylene over succinic acid production.[13] Thus, 1449B PsEFE-related EFEs might exist, or might be evolved, which exclusively catalyse ethylene production from 2OG, but not C5 hydroxylation of *L*-arginine. Such EFEs are attractive from biocatalysis perspectives to enable sustainable ethylene production from renewable resources.[27]

The 1449B PsEFE:Mn:2OG structure reveals an atypical inverted 2OG metal ion binding mode in which the 2OG 2-oxoacid group is positioned to coordinate with the guanidinium group of R277 rather than with Mn, which is instead complexed by the 2OG C5 carboxylate (Figure 3.5). This observation is precedented by similar, but not identical 2OG binding modes in reported PK2 PsEFE:Mn:2OG and *Penicillium digitatum* EFE:Mn:2OG structures (Figure 3.5).[9b,13b] It is unclear whether this atypical binding mode of 2OG in complex with PsEFE is relevant in cells, including because the crystal structures reported to date suggest that 2OG only adopts these inverted modes in complex with PK2 PsEFE and catalytically inert Mn(II), but not Fe(II), and in the absence of *L*-arginine (Figure 3.5h).[9b,12b]

It is possible that the atypical 2OG metal ion binding modes observed with PK2 PsEFE, 1449B PsEFE, and *Penicillium digitatum* EFE are of *in vivo* relevance.[9b,13b] Some Fe(II)-dependent oxygenases are proposed to act as non-Fe transition metal ion sensors as a result of Fe(II)

displacement and consequent activity loss.[28] It is possible that catalytically inactive 2OG binding modes at the PsEFE active site may be involved in metal ion sensing and/or in enabling different reactions; in the latter case, it is notable that structurally distinct dioxygenases are catalytically active using either Fe(II) or other divalent transition metal ions.[29]

2OG not only adopts atypical metal ion binding modes in complex with PsEFE and Mn, but also atypical (*i.e.*, non *anti*-periplanar) conformations, *i.e.*, with its C2/C5 groups being positioned *gauche* (Figure 3.5f).[9b] Atypical 2OG conformations are also observed in complex with 2OG oxygenases other than PsEFE. For example, 2OG was observed to adopt a C2/C5 *gauche* conformation with the 2OG C5 carboxylate group complexing Mn via a water molecule in crystal structures with disease-related active site variants of human aspartate/asparagine-β-hydroxylase (AspH), whereas it adopted a C2/C5 antiperiplanar conformation in complex with wildtype AspH.[30] The oncometabolite *D*-2-hydoxyglutarate (2HG) has been observed to adopt a C2/C5 *gauche* conformation in complex with the JmjC N^{ε}-methyllysine demethylase 5B (KDM5B) and Mn, *i.e.*, with the 2HG C5 carboxylate group being positioned to interact with the hydroxy group of 2HG.[31] Notably, oxaloacetate has been observed in an inverted metal ion binding mode in complex with KDM5B and Mn.[31] Although oxaloacetate lacks one of the two 2OG methylene groups, the combined observations indicate that 2OG binding modes and conformations may be flexible at 2OG oxygenase active site, possibly in a manner affecting catalysis.

Given that the *glycinea*, 1449B, and PK2 PsEFEs can employ naturally occurring 2OG derivatives as cosubstrates to sustain *L*-arginine hydroxylation and alcohol formation, but not ethylene production (Figure 3.4),[6] it is possible that the 2OG derivative substitution pattern affects the binding mode and/or conformation of the 2OG derivatives at the PsEFE active site, as observed in the case of AspH,[30a] potentially in a manner affecting reaction outcomes. Although we obtained crystal structures of 1449B PsEFE in complex with Mn and 4-methyl-

2OG or 3-methyl-2OG to test this proposal, their resolutions (2.05 Å and 1.85 Å, respectively) were insufficient to unambiguously assign the precise 2OG derivative binding modes at the active site.

An extensive water network was observed in the active site of the 1449B PsEFE:Mn:2OG structure, this network likely stabilises 2OG binding and may affect reaction outcomes (Figure 3.6). Some of the 1449B PsEFE residues involved in stabilising the water network at the active site are also likely involved in binding *L*-arginine (Supporting Table 3.S1).[9b] All EFE crystal structures[9,13b] and ACCO structures[15b] also feature active site water networks, suggesting such networks may be an important general characteristic of ethylene-producing enzymes of the 2OG superfamily. It is therefore of interest to investigate the effects of variations in residues which are involved in stabilising the water network on PsEFE fold, 2OG binding mode and conformation, catalytic efficiency, and reaction outcome.

3.7 Conclusion

The combined spectroscopic and crystallographic evidence demonstrates that both the *glycinea* and 1449B PsEFEs have capacity to produce ethylene from 2OG (Figure 3.3), however, apparently less efficiently than PK2 PsEFE and, at least in the case of 1449B PsEFE, with an altered product ratio, favouring ethylene over succinic acid formation. The results provide further evidence for variations in the binding modes of 2OG and 2OG derivatives at the active sites of 2OG oxygenases.

3.8 Experimental Procedures

Protein production. Plasmid DNA encoding for *glycinea* PsEFE (*i.e.*, the R326S PK2 PsEFE variation) was obtained by standard site-directed mutagenesis of our reported PK2 PsEFE construct[9a] (Supporting Table 3.S3). *Glycinea* PsEFE was produced and purified as reported for PK2 PsEFE.[6b,9a] Plasmid DNA encoding for 1449B PsEFE in a pet-28a(+)-TEV vector

backbone was codon optimised for expression in *E. coli* cells and prepared by GenScript Biotech. 1449B PsEFE was produced and purified as reported for PK2 PsEFE;[6b,9a] however, the Tobacco Etch Virus (TEV) protease was used to remove the N-terminal His_6-tag of 1449B PsEFE instead of the SENP2 protease. Thus, His_6-tagged 1449B PsEFE was incubated overnight with TEV protease (~100:1 ratio) in cleavage buffer (50 mM Tris, 100 mM NaCl, 10%$_{v/v}$ glycerol, pH 7.8) at 4 ˚C, followed by reverse Ni(II) affinity chromatography purification and size exclusion chromatography as reported for PK2 PsEFE.[6b,9a] *Glycinea* and 1449B PsEFE were >90% pure by SDS-PAGE analysis and had the anticipated masses (Supporting Figure 3.S2).

2OG derivatives. 2-Oxoglutarate (2OG), 2-oxoadipate (2OA), and racemic 4-hydroxy-2OG (**3**) were from Sigma-Aldrich. Racemic 4-methyl-2OG (**1**) and 3-methyl-2OG (**6**) were synthesised as reported.[30b]

^{1}H NMR assays. ^{1}H NMR assays were performed as reported,[6b] typically using: 2OG or 2OG derivative (400 μM), *L*-arginine (500 μM), *L*-ascorbic acid (500 μM), Fe(II) (50 μM), TMSP-d_4 (800 μM), and a PsEFE (10 μM for 1449B PsEFE and 30 μM for *glycinea* PsEFE) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O), if not specified otherwise. Freshly thawed *glycinea* and 1449B PsEFE aliquots were used for all experiments. Assays were performed using a BrukerAVIII 700 MHz NMR spectrometer with a 5-mm inverse triple-resonance-inverse cryoprobe. The water peak in ^{1}H NMR experiments was suppressed using the perfect echo-modified WATERGATE solvent suppression method.[32]

Crystallography. Full-length 1449B PsEFE (11 mg/mL) was incubated with $MnCl_2$ (3 mM), and 2OG (10 mM) in buffer (50 mM Tris, pH 7.5) at 4 °C for 15 min and centrifuged using a MicroCL 21R centrifuge (Thermo Fisher Scientific; 18,800 × *g*, 4 °C, 1 min). Crystallisations were performed using the sitting drop vapor diffusion method (3 μL drops) at 17 °C using 2:1,

1:1, and 1:2 well solution (μL): protein solution (μL) ratios. Conditions were: 0.03 M citric acid, 0.07 M Bis-Tris propane/pH 7.6, 20%$_{w/v}$ polyethylene glycol (PEG) 3350. Crystals were cryo-protected with 20-25%$_{v/v}$ glycerol prior to flash-freezing with liquid N_2.

Diffraction data for the 1449B PsEFE single crystals were collected at the Diamond Light Source (beamline: I03). The data were indexed, integrated, and scaled using the Xia2 pipeline.[33] Structures were solved by molecular replacement with Phaser,[34] using a reported PK2 PsEFE structure as a search model (PDB ID: 5MOF[9a]). Structures were iteratively refined using PHENIX[35] and manual model building using COOT.[36]

3.9 Acknowledgements

This research was funded in part by the Wellcome Trust (106244/Z/14/Z). For the purpose of open access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission. We thank Cancer Research UK (C8717/A18245) and the Biotechnology and Biological Sciences Research Council (BB/J003018/1 and BB/R000344/1) for funding and Prof. Dr. C. Redfield for helpful discussions.

3.10 Author Contributions

Y.S. and S.D. produced enzymes with help of E.S.Z.C.; Y.S., S.D., and E.S.Z.C. performed assays; Y.S., S.D., and E.S.Z.C. performed crystallizations, M.D.A. solved and refined the structure with help from Y.S.; S.D. and L.B. conceived the project together with Y.S. and C.J.S.; L.B., Z.Z., and C.J.S. supervised the research; all authors analysed data; L.B., S.D. and C.J.S. wrote the manuscript with help from all authors. We thank the Diamond Light Source (beamline I03) and staff for allocation of beam time and support.

3.11 References

[1] a) B. M. Binder, *J. Biol. Chem.* **2020**, *295*, 7710–7725; b) X. Wang, H. Wen, A. Suprun, H. Zhu, *Plants* **2025**, *14*, 309; c) N. Papon, B. M. Binder, *Trends Microbiol.* **2019**, *27*, 193–196; d) M. Ravanbakhsh, R. Sasidharan, L. A. C. J. Voesenek, G. A. Kowalchuk, A. Jousset, *Microbiome* **2018**, *6*, 52; e) B. Bidon, S. Kabbara, V. Courdavault, G. Glévarec, A. Oudin, F. Héricourt, S. Carpin, L. Spíchal, B. M. Binder, J. M. Cock, N. Papon, *Cells* **2020**, *9*, 2526; f) I. Camehl, I. Sherameti, Y. Venus, G. Bethke, A. Varma, J. Lee, R. Oelmüller, *New Phytol.* **2010**, *185*, 1062–1073; g) S. Gu, L. Xie, Q. Guan, X. Sheng, Y. Fang, X. Wang, *Int. J. Food Microbiol.* **2024**, *418*, 110729.

[2] M. Houben, B. Van de Poel, *Front. Plant Sci.* **2019**, *10*, 695.

[3] a) C. C. Lee, Y. Hu, M. W. Ribbe, *Science* **2010**, *329*, 642–642; b) Y. Hu, C. C. Lee, M. W. Ribbe, *Science* **2011**, *333*, 753–755; c) C. C. Lee, Y. Hu, M. W. Ribbe, *Angew. Chem. Int. Ed.* **2011**, *50*, 5545–5547; d) Y. Hu, C. C. Lee, M. Grosch, J. B. Solomon, W. Weigand, M. W. Ribbe, *Chem. Rev.* **2023**, *123*, 5755–5797; e) L. C. Seefeldt, Z.-Y. Yang, D. A. Lukoyanov, D. F. Harris, D. R. Dean, S. Raugei, B. M. Hoffman, *Chem. Rev.* **2020**, *120*, 5082–5106; f) B. M. Barney, R. Y. Igarashi, P. C. Dos Santos, D. R. Dean, L. C. Seefeldt, *J. Biol. Chem.* **2004**, *279*, 53621–53624; g) J. A. North, A. B. Narrowe, W. Xiong, K. M. Byerly, G. Zhao, S. J. Young, S. Murali, J. A. Wildenthal, W. R. Cannon, K. C. Wrighton, R. L. Hettich, F. R. Tabita, *Science* **2020**, *369*, 1094–1098; h) A. Lago-Maciel, J. C. Soares, J. Zarzycki, C. J. Buchanan, T. Reif-Trauttmansdorff, F. V. Schmidt, S. Lometto, N. Paczia, J. M. Schuller, D. F. Hansen, G. T. Heller, S. Prinz, G. K. A. Hochberg, A. J. Pierik, J. G. Rebelein, *bioRxiv* **2024**, https://doi.org/10.1101/2024.10.19.619033; i) J. A. North, A. R. Miller, J. A. Wildenthal, S. J. Young, F. R. Tabita, *Proc. Natl. Acad. Sci. USA* **2017**, *114*, E10455–E10464; j) R. P. Hausinger, S. B. J. S. Rifayee, M. G. Thomas, S. Chatterjee, J. Hu, C. Z. Christov, *RSC Chem. Biol.* **2023**, *4*, 635–646; k) Y. Cui, Y. Jiang, M. Xiao, M. Z. Munir, S. Riaz, F. Rasul, M. Daroch, *Int. J. Mol. Sci.* **2022**, *23*, 4500; l) K. Nagahama, T. Ogawa, T. Fujii, M. Tazaki, S. Tanase, Y. Morino, H. Fukuda, *J. Gen. Microbiol.* **1991**, *137*, 2281–2286; m) H. Fukuda, T. Ogawa, K. Ishihara, T. Fujii, K. Nagahama, T. Omata, Y. Inoue, S. Tanase, Y. Morino, *Biochem. Biophys. Res. Commun.* **1992**, *188*, 826–832; n) M. Goto, Y. Ishida, Y. Takikawa, H. Hyodo, *Plant Cell Physiol.* **1985**, *26*, 141–150.

[4] a) K. Nagahama, T. Ogawa, T. Fujii, M. Tazaki, M. Goto, H. Fukuda, *J. Gen. Microbiol.* **1991**, *137*, 1641–1646; b) S. Martinez, R. P. Hausinger, *Biochemistry* **2016**, *55*, 5989–5999.

[5] a) H. Fukuda, T. Ogawa, M. Tazaki, K. Nagahama, T. Fujii, S. Tanase, Y. Morino, *Biochem. Biophys. Res. Commun.* **1992**, *188*, 483–489; b) S. Martinez, R. P. Hausinger, *J. Biol. Chem.* **2015**, *290*, 20702–20711.

[6] a) R. A. Copeland, S. Zhou, I. Schaperdoth, T. K. C. Shoda, J. M. Bollinger, C. Krebs, *Science* **2021**, *373*, 1489–1493; b) S. Dhingra, Z. Zhang, C. T. Lohans, L. Brewitz, C. J. Schofield, *J. Biol. Chem.* **2024**, *300*, 107546.

[7] a) S. Kal, L. Que, *J. Biol. Inorg. Chem.* **2017**, *22*, 339–365; b) W. Aik, M. A. McDonough, A. Thalhammer, R. Chowdhury, C. J. Schofield, *Curr. Opin. Struct. Biol.* **2012**, *22*, 691–700.

[8] a) M. A. McDonough, C. Loenarz, R. Chowdhury, I. J. Clifton, C. J. Schofield, *Curr. Opin. Struct. Biol.* **2010**, *20*, 659–672; b) *2-Oxoglutarate-dependent oxygenases* (Ed.: R. P. Hausinger, C. J. Schofield), The Royal Society of Chemistry, Cambridge, United Kingdom, **2015**.

[9] a) Z. Zhang, T. J. Smart, H. Choi, F. Hardy, C. T. Lohans, M. I. Abboud, M. S. W. Richardson, R. S. Paton, M. A. McDonough, C. J. Schofield, *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 4667–4672; b) S. Martinez, M. Fellner, C. Q. Herr, A. Ritchie, J. Hu, R. P. Hausinger, *J. Am. Chem. Soc.* **2017**, *139*, 11980–11988.

[10] K. Watanabe, K. Nagahama, M. Sato, *Phytopathology* **1998**, *88*, 1205–1209.
[11] a) H. Weingart, B. Volksch, *Appl. Environ. Microbiol.* **1997**, *63*, 156–161; b) H. Weingart, B. Völksch, M. S. Ullrich, *Phytopathology* **1999**, *89*, 360–365; c) M. Sato, K. Watanabe, M. Yazawa, Y. Takikawa, K. Nishiyama, *Phytopathology* **1997**, *87*, 1192–1196.
[12] a) E. J. Burke, R. A. Copeland, Y. Dixit, C. Krebs, J. M. Bollinger, Jr., *J. Am. Chem. Soc.* **2024**, *146*, 1977–1983; b) R. A. Copeland, K. M. Davis, T. K. C. Shoda, E. J. Blaesi, A. K. Boal, C. Krebs, J. M. Bollinger, Jr., *J. Am. Chem. Soc.* **2021**, *143*, 2293–2303; c) N. Johansson, K. O. Persson, C. Larsson, J. Norbeck, *BMC Biochem.* **2014**, *15*, 22; d) K. Nagahama, K. Yoshino, M. Matsuoka, S. Tanase, T. Ogawa, H. Fukuda, *J. Ferment. Bioeng.* **1998**, *85*, 255–258.
[13] a) S. Chatterjee, J. A. Rankin, M. A. Farrugia, B. J. Delaney, N. S. Pascual, J. VanAntwerp, D. R. Woldring, J. Hu, R. P. Hausinger, *Biochemistry* **2025**; b) S. Chatterjee, J. A. Rankin, M. A. Farrugia, S. B. J S Rifayee, C. Z. Christov, J. Hu, R. P. Hausinger, *Biochemistry* **2025**.
[14] a) H. Fukuda, T. Fujii, T. Ogawa, *Agric. Biol. Chem.* **1986**, *50*, 977–981; b) H. Fukuda, H. Kitajima, T. Fujii, M. Tazaki, T. Ogawa, *FEMS Microbiol. Lett.* **1989**, *59*, 1–5.
[15] a) Z. Zhang, J.-S. Ren, I. J. Clifton, C. J. Schofield, *Chem. Biol.* **2004**, *11*, 1383–1394; b) X. Sun, Y. Li, W. He, C. Ji, P. Xia, Y. Wang, S. Du, H. Li, N. Raikhel, J. Xiao, H. Guo, *Nat. Commun.* **2017**, *8*, 15758.
[16] X. Robert, P. Gouet, *Nucleic Acids Res.* **2014**, *42*, W320–W324.
[17] a) A. J. Copper, A. G. Redfield, *J. Biol. Chem.* **1975**, *250*, 527–532; b) T. S. Viswanathan, R. E. Johnson, H. F. Fisher, *Biochemistry* **1982**, *21*, 339–345.
[18] a) K. Danhauser, S. W. Sauer, T. B. Haack, T. Wieland, C. Staufner, E. Graf, J. Zschocke, Tim M. Strom, T. Traub, Jürgen G. Okun, T. Meitinger, Georg F. Hoffmann, H. Prokisch, S. Kölker, *Am. J. Hum. Genet.* **2012**, *91*, 1082–1087; b) D. E. Matthews, *J. Nutr.* **2020**, *150*, 2548S–2555S.
[19] a) R. Díaz, H. Gallart-Ayala, J. V. Sancho, O. Nuñez, T. Zamora, C. P. B. Martins, F. Hernández, S. Hernández-Cassou, J. Saurina, A. Checa, *J. Chromatogr. A* **2016**, *1433*, 90–97; b) D. Mukherjee, M. M. Laloraya, *Biochem. Physiol. Pflanz.* **1974**, *166*, 429–436; c) A. I. Virtanen, A.-M. Berg, *Acta Chem. Scand.* **1955**, *9*, 553–554.
[20] A. L. Wilkins, Y. Lu, S.-T. Tan, *J. Agric. Food Chem.* **1995**, *43*, 3021–3025.
[21] D. Siegel, A. C. Meinema, H. Permentier, G. Hopfgartner, R. Bischoff, *Anal. Chem.* **2014**, *86*, 5089–5100.
[22] S. Chatterjee, M. Fellner, J. Rankin, M. G. Thomas, S. B. J S Rifayee, C. Z. Christov, J. Hu, R. P. Hausinger, *Biochemistry* **2024**, *63*, 1038–1050.
[23] S. Dhingra, Y. Sun, Z. Zhang, C. J. Schofield, L. Brewitz, manuscript under review.
[24] a) M. Sami, T. J. N. Brown, P. L. Roach, C. J. Schofield, J. E. Baldwin, *FEBS Lett.* **1997**, *405*, 191–194; b) H. S. Chin, J. Sim, K. I. Seah, T. S. Sim, *FEMS Microbiol. Lett.* **2003**, *218*, 251–257; c) S. J. Lipscomb, H.-J. Lee, M. Mukherji, J. E. Baldwin, C. J. Schofield, M. D. Lloyd, *Eur. J. Biochem.* **2002**, *269*, 2735–2739; d) M. D. Lloyd, S. J. Lipscomb, K. S. Hewitson, C. M. H. Hensgens, J. E. Baldwin, C. J. Schofield, *J. Biol. Chem.* **2004**, *279*, 15420–15426; e) H.-J. Lee, M. D. Lloyd, K. Harlos, I. J. Clifton, J. E. Baldwin, C. J. Schofield, *J. Mol. Biol.* **2001**, *308*, 937–948.
[25] K. Ishihara, M. Matsuoka, T. Ogawa, H. Fukuda, *J. Ferment. Bioeng.* **1996**, *82*, 509–511.
[26] a) S. S. Chaturvedi, R. Ramanan, J. Hu, R. P. Hausinger, C. Z. Christov, *ACS Catal.* **2021**, *11*, 1578–1592; b) M. G. Thomas, S. B. Jaber Sathik Rifayee, C. Z. Christov, *ACS Catal.* **2024**, *14*, 18550–18569.
[27] a) J. Wilson, S. Gering, J. Pinard, R. Lucas, B. R. Briggs, *Biotechnol. Biofuels* **2018**, *11*, 234; b) C. Eckert, W. Xu, W. Xiong, S. Lynch, J. Ungerer, L. Tao, R. Gill, P.-C. Maness, J. Yu, *Biotechnol. Biofuels* **2014**, *7*, 33.

[28] a) H. Chen, M. Costa, *BioMetals* **2009**, *22*, 191–196; b) N. R. Rose, M. A. McDonough, O. N. F. King, A. Kawamura, C. J. Schofield, *Chem. Soc. Rev.* **2011**, *40*, 4364–4397; c) L. Brewitz, A. Tumber, I. Pfeffer, M. A. McDonough, C. J. Schofield, *Sci. Rep.* **2020**, *10*, 8650; d) A. Brasnett, I. Pfeffer, L. Brewitz, R. Chowdhury, Y. Nakashima, A. Tumber, M. A. McDonough, C. J. Schofield, *Angew. Chem. Int. Ed.* **2021**, *60*, 14657–14663.

[29] a) J. P. Emerson, E. G. Kovaleva, E. R. Farquhar, J. D. Lipscomb, L. Que, *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 7347–7352; b) A. R. Deshpande, T. C. Pochapsky, D. Ringe, *Chem. Rev.* **2017**, *117*, 10474–10501.

[30] a) C. X. Hou, A. Brasnett, P. Rabe, C. J. Schofield, L. Brewitz, *J. Biol. Chem.*, accepted for publication; b) L. Brewitz, Y. Nakashima, C. J. Schofield, *Chem. Sci.* **2021**, *12*, 1327–1342; c) I. Pfeffer, L. Brewitz, T. Krojer, S. A. Jensen, G. T. Kochan, N. J. Kershaw, K. S. Hewitson, L. A. McNeill, H. Kramer, M. Münzel, R. J. Hopkinson, U. Oppermann, P. A. Handford, M. A. McDonough, C. J. Schofield, *Nat. Commun.* **2019**, *10*, 4910.

[31] H. Tarhonskaya, R. P. Nowak, C. Johansson, A. Szykowska, A. Tumber, R. L. Hancock, P. Lang, E. Flashman, U. Oppermann, C. J. Schofield, A. Kawamura, *J. Mol. Biol.* **2017**, *429*, 2895–2906.

[32] R. W. Adams, C. M. Holroyd, J. A. Aguilar, M. Nilsson, G. A. Morris, *Chem. Commun.* **2013**, *49*, 358–360.

[33] a) G. Winter, *J. Appl. Cryst.* **2010**, *43*, 186–190; b) G. Winter, J. Beilsten-Edmands, N. Devenish, M. Gerstel, R. J. Gildea, D. McDonagh, E. Pascal, D. G. Waterman, B. H. Williams, G. Evans, *Protein Sci.* **2022**, *31*, 232–250.


[34] A. J. McCoy, R. W. Grosse-Kunstleve, P. D. Adams, M. D. Winn, L. C. Storoni, R. J. Read, *J. Appl. Cryst.* **2007**, *40*, 658–674.

[35] D. Liebschner, P. V. Afonine, M. L. Baker, G. Bunkóczi, V. B. Chen, T. I. Croll, B. Hintze, L.-W. Hung, S. Jain, A. J. McCoy, N. W. Moriarty, R. D. Oeffner, B. K. Poon, M. G. Prisant, R. J. Read, J. S. Richardson, D. C. Richardson, M. D. Sammito, O. V. Sobolev, D. H. Stockwell, T. C. Terwilliger, A. G. Urzhumtsev, L. L. Videau, C. J. Williams, P. D. Adams, *Acta Cryst.* **2019**, *D75*, 861–877.


[36] P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, *Acta Cryst. D* **2010**, *66*, 486–501.

3.12 Supporting Information


3.12.1 Supporting Figures




Supporting Figure 3.S1. Sequence comparison of functionally and structurally characterised ethylene/succinate-forming enzymes (EFEs) and 1-aminocyclopropane-1-carboxylate oxidases (ACCOs). *Pseudomonas savastanoi* pv. *phaseolicola* PK2 EFE (Uniprot ID: P32021): PK2 PsEFE; *Pseudomonas savastanoi* pv. *glycinea* EFE (Uniprot ID: Q7BS32): *glycinea* PsEFE; *Pseudomonas savastanoi* pv. 1449B EFE (Uniprot ID: Q9Z3T0): 1449B PsEFE; *Penicillium digitatum* EFE (Uniprot ID: K9GJX7): PdEFE; *Petunia hybrida* ACCO1 (Uniprot ID: Q08506): PhACCO1; *Arabidopsis thaliana* ACCO1 (Uniprot ID: Q9ZUN4): AtACCO1. α-Helices and β-strands of PK2 PsEFE are numbered based on a PK2 PsEFE structure (PDB ID: 5MOF[1]); β-turns: TT. Cyan triangles: β-strands forming the DSBH fold of PK2 PsEFE; green circles: the PK2 PsEFE HXD…H Fe(II)-binding motif. Identical residues are in red; residues of similar polarity are in yellow. The Figure 3.S1 was generated using ESPript 3.0[2].




Supporting Figure 3.S2. Purity of the *glycinea* and 1449B PsEFEs. (**a**) SDS-PAGE analysis (using a 4-12% Novex™ 330 Tris-glycine gel; Thermo Fisher) of the isolated recombinant *glycinea* and 1449B PsEFEs indicate >90% purity. The PageRuler™ Plus Prestained Protein Ladder (Thermo Fisher) was used. (**b**) Mass spectrometric analyses of PsEFEs (3 µM) in buffer (25 mM Tris, pH 7.8) were performed using an iFunnel Agilent 6550 accurate mass quadrupole time-of-flight (Q-TOF) mass spectrometer operated in the positive ionization mode for: (**i**) 1449B PsEFE (calculated mass: 38548 Da; observed mass: 38547 Da) and (**ii**) *glycinea* PsEFE (calculated mass: 39301 Da; observed mass: 39301 Da).




Supporting Figure 3.S3. 1449B PsEFE and *glycinea* PsEFE do not catalyse efficient oxidation of 2OG in the absence of *L*-arginine. (**a-d**) ^{1}H NMR analyses (~1.5 to ~4.0 ppm and ~5.0 to ~7.5 ppm) of reactions of 2OG ~6 h post incubation with: (**a**) 1449B PsEFE and (**b**) *glycinea* PsEFE; (**c**) Reported ^{1}H NMR spectrum for a ~6 h incubation of 2OG with PK2 PsEFE.[3] Conditions: 400 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and appropriate concentrations of PsEFE (2 µM for PK2;[3] 5 µM for1449B; 10 µM for *glycinea*) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). The peaks at ~2.1 and ~2.3 ppm (dashed circles) correspond to the hydrated ketone form of 2OG.[4] The peaks at ~3.6 and ~3.7 ppm in (**a-c**) are from glycerol; the singlet at ~3.7 ppm in (**a** and **b**) corresponds to Tris.




Supporting Figure 3.S4 *Glycinea* PsEFE catalyses oxidation of 2OG and 2OG derivatives in the presence of *L*-arginine. Reaction of the *glycinea* PsEFE with: (**a**) 2OG, (**b**) 2OA, (**c**) 4-methyl-2OG (**3**),[5] (**d**) 3-methyl-2OG (**4**),[5] and (**e**) 4-hydroxy-2OG (**5**). Conditions: 400 μM 2OG/2OG derivative, 500 μM *L*-arginine, 500 μM *L*-ascorbate, 50 μM Fe(II), 800 μM TMSP-d_4, and *glycinea* PsEFE (30 μM with 2OG derivatives; 10 μM with 2OG) in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Reaction times are normalized to the end of the first ^{1}H NMR acquisition following *glycinea* PsEFE addition to the reaction mixture (t = 0 min), by which time varied levels of turnover were manifest. Results are means of independent triplicates (n = 3; mean ± standard deviation, SD). Note that the 2OG derivatives **3**, **4**, and **5** were used as racemic mixtures; there is potential for stereoselectivity, including preferred conversion (including for a particular pathway) / inhibition for one enantiomer.




Supporting Figure 3.S5. 1449B PsEFE catalyses oxidation of 2OG and 2OG derivatives in the presence of *L*-arginine. Reaction of the 1449B PsEFE with: (**a**) 2OG, (**b**) 2OA, (**c**) 4-methyl-2OG (**3**),[5] (**d**) 3-methyl-2OG (**4**),[5] and (**e**) 4-hydroxy-2OG (**5**). Conditions: 400 μM 2OG/2OG derivative, 500 μM *L*-arginine, 500 μM *L*-ascorbate, 50 μM Fe(II), 800 μM TMSP-d_4, and 1449B PsEFE (10 μM with 2OG derivatives; 5 μM with 2OG) in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Reaction times are normalized to the end of the first ^{1}H NMR acquisition following 1449B PsEFE addition to the reaction mixture (t = 0 min), by which time varied levels of turnover were manifest. Results are means of independent triplicates (n = 3; mean ± SD). Note that the 2OG derivatives **3**, **4**, and **5** were used as racemic mixtures; there is potential for stereoselectivity, including preferred conversion (including for a particular pathway) / inhibition for one enantiomer.




Supporting Figure 3.S6. ^{1}H NMR analysis of the reactions of the 1449B, *glycinea*, and PK2 PsEFEs with 2OA. (a-b) Reported ^{1}H NMR spectra (700 MHz; ~0.8 to ~4.4 ppm and ~5.4 to ~8.0 ppm) of authentic samples of: (**a**) **2b** and (**b**) **2a**.[6] **(c-e)** ^{1}H NMR spectra of 2OA reaction mixtures, ~90 min post addition of: (**c**) 1449B PsEFE and (**d**) *glycinea* PsEFE, showing conversion of 2OA to **2a** and **2b**; (**e**) Reported ^{1}H NMR spectrum for a ~90 min incubation of 2OA with PK2 PsEFE.[6] (**f**) **Reported** ^{1}H NMR spectrum of 2OA under standard conditions in the absence of PsEFE.[6] Conditions: 400 µM 2OA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and appropriate concentrations of PsEFEs (10 µM for PK2 PsEFE[6] and 1449B PsEFE; 30 µM for *glycinea* PsEFE) in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**c** and **e**) are from glycerol; the singlet at ~3.7 ppm in (**c-e**) corresponds to Tris. Representative spectra from independent triplicates are shown.




Supporting Figure 3.S7. ^{1}H NMR analysis of the reactions of the 1449B, *glycinea*, and PK2 PsEFEs with 4-methyl-2OG (3). (a-b) Reported ^{1}H NMR spectra (700 MHz; ~0.8 to ~4.4 ppm and ~5.4 to ~8.0 ppm) of authentic samples of: (**a**) **3b** and (**b**) **3a**, the latter in the presence of PK2 PsEFE.[6] **(c-e)** ^{1}H NMR spectra of 4-methyl-2OG (**3**) reaction mixtures, ~90 min post addition of: (**c**) 1449B PsEFE and (**d**) *glycinea* PsEFE, showing conversion of **3** to **3a** and **3b**; (**e**) Reported ^{1}H NMR spectrum for a ~90 min incubation of **3** with PK2 PsEFE.[6] (**f**) Reported ^{1}H NMR spectrum of **3** under standard conditions in the absence of PsEFE.[6] Conditions: 400 µM **3**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and appropriate concentrations of PsEFEs (10 µM for PK2 PsEFE[6] and 1449B PsEFE; 30 µM for *glycinea* PsEFE) in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**b**, **c**, and **e**) are from glycerol; the singlet at ~3.7 ppm in (**b-e**) corresponds to Tris. Representative spectra from independent triplicates are shown.



Supporting Figure 3.S8. ^{1}H NMR analysis of the reactions of the 1449B, *glycinea*, and PK2 PsEFEs with 3-methyl-2OG (4). (a-b) Reported ^{1}H NMR spectra (700 MHz; ~0.8 to ~4.4 ppm and ~5.4 to ~8.0 ppm) of authentic samples of: (**a**) **4b** and (**b**) **3a**, the latter in the presence of PK2 PsEFE.[6] **(c-e)** ^{1}H NMR spectra of 3-methyl-2OG (**4**) reaction mixtures, ~90 min post addition of: (**c**) 1449B PsEFE and (**d**) *glycinea* PsEFE, showing conversion of **4** to **3a** and **4b**; (**e**) Reported ^{1}H NMR spectrum for a ~90 min incubation of **4** with PK2 PsEFE.[6] (**f**) Reported ^{1}H NMR spectrum of **4** under standard conditions in the absence of PsEFE.[6] Conditions: 400 µM **4**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and appropriate concentrations of PsEFEs (10 µM for PK2 PsEFE[6] and 1449B PsEFE; 30 µM for *glycinea* PsEFE) in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**b**, **c**, and **e**) are from glycerol; the singlet at ~3.7 ppm in (**b-e**) corresponds to Tris. Representative spectra from independent triplicates are shown.



Supporting Figure 3.S9. ^{1}H NMR analysis of the reactions of the 1449B, *glycinea*, and PK2 PsEFEs with 4-hydroxy-2OG (5). (a-b) Reported ^{1}H NMR spectra (700 MHz; ~0.8 to ~4.4 ppm and ~5.4 to ~8.0 ppm) of authentic samples of: (**a**) **5b** and (**b**) **5a**.[6] **(c-e)** ^{1}H NMR spectra of 4-hydroxy-2OG (**5**) reaction mixtures, ~90 min post addition of: (**c**) 1449B PsEFE and (**d**) *glycinea* PsEFE, showing conversion of **5** to **5a** and **5b**; (**e**) Reported ^{1}H NMR spectrum for a ~90 min incubation of **5** with PK2 PsEFE.[6] (**f**) Reported ^{1}H NMR spectrum of **5** under standard conditions in the absence of PsEFE.[6] Conditions: 400 µM **5**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and appropriate concentrations of PsEFEs (10 µM for PK2 PsEFE[6] and 1449B PsEFE; 30 µM for *glycinea* PsEFE) in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**c** and **e**) are from glycerol; the singlet at ~3.7 ppm in (**c-f**) corresponds to Tris. Representative spectra from independent triplicates are shown.



Supporting Figure 3.S10. Comparison of views from crystal structures of 1449B and PK2 PsEFE in complex with 2OG and Mn. Green: 1449B PsEFE; lavender: carbon-backbone of 2OG; light grey: Mn; red: oxygen; blue: nitrogen. (**a**) Overview of the 1449B PsEFE:Mn:2OG crystal structure. The core DSBH fold is in cyan; (**b**) superimposition of views from the 1449B PsEFE:Mn:2OG and PK2 PsEFE:Mn:2OG (PK2 PsEFE: orange; Mn: yellow; carbon-backbone of 2OG: pink; PDB ID: 5MOF[1]) crystal structures reveal similar, but not identical, PsEFE conformations (Cα RMSD ~ 0.415 Å). Note that the C-terminal region of 1449B PsEFE is truncated; (**c**) the 1449B PsEFE active site is shaped by mostly hydrophobic residues; (**d**) superimposition of active site views from the 1449B PsEFE:Mn:2OG and PK2 PsEFE:Mn:2OG (PK2 PsEFE: orange; Mn: yellow; carbon-backbone of 2OG: pink; PDB ID: 5MOF[1]) crystal structures reveal near identical conformation of amino acid sidechains; note, however, that conformations of 2OG differ (Figure 3.5).

3.12.2 Supporting Tables

Supporting Table 3.S1. Effects of single amino acid residue variations on the ethylene-producing capacity of wildtype (wt) PK2 PsEFE. Note that only the effects of some of the shown variations in the PK2 PsEFE sequence were validated using isolated recombinant PK2 PsEFE variants,[1,7] whereas other studies used intact cells (denoted: *)[8] or cell extracts (denoted: #),[9] a difference complicating comparison including because cells may produce ethylene via pathways not involving PK2 PsEFE.

Variation from wt PK2 PsEFE	Proposed role of residue in wt PK2 PsEFE	Ethylene production activity	References
L22M*	Surface residue	**Yes** (no effect compared to wt PK2 PsEFE)	[8]
L73K/R	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[7a]
S81R/Y	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[7a]
E84A/Q	*L*-arginine binding	**No**	[1,7a]
E84D	*L*-arginine binding	**Trace**	[1]
T86S/V	*L*-arginine binding	**Reduced** (compared to wt PK2 PsEFE)	[1]
H116Q	Structural (interior/surface)	**Reduced** (compared to wt PK2 PsEFE)	[7e]
H168Q	Structural (interior/surface)	**Reduced** (compared to wt PK2 PsEFE)	[7e]
H169Q	Structural (interior)	**Reduced** (compared to wt PK2 PsEFE)	[7e]
R171A	*L*-arginine binding	**No**	[7a]
R171K	*L*-arginine binding	**No**	[1]
V172T*	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[8]
F175Y	2OG binding	**Reduced** (compared to wt PK2 PsEFE)	[1]
R184A	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[7a]
H189A*	Metal-binding	**No**	[8]
H189Q	Metal-binding	**No**	[7e]
D191A*	Metal-binding	**No**	[8]
D191E	Metal-binding	**Reduced** (compared to wt PK2 PsEFE)	[7b,7d]
Y192F	*L*-arginine binding	**Reduced** (compared to wt PK2 PsEFE)	[1]
Y192R/W	*L*-arginine binding	**Reduced** (compared to wt PK2 PsEFE)	[7a]
V196R	2OG binding (hydrophobic)	**Reduced** (compared to wt PK2 PsEFE)	[7a]
A198V	2OG binding (hydrophobic)	**Reduced** (compared to wt PK2 PsEFE)	[7a]
A199G*	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[8]
L206V	2OG binding (hydrophobic)	**Reduced** (compared to wt PK2 PsEFE)	[7c]
E213A	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[7a]
E215A	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[7a]
H233A*	Surface residue	**Enhanced** (compared to wt PK2 PsEFE)	[8]
H233Q	Surface residue	**No**	[7e]
E235D*	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[8]
R236S*	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[8]

Variation from wt PK2 PsEFE	Proposed role of residue in wt PK2 PsEFE	Ethylene production activity	References
I254M*	Surface residue	**Yes** (no effect compared to wt PK2 PsEFE)	[8]
H268A*	Metal-binding	**No**	[8]
H268Q	Metal-binding	**Reduced** (compared to wt PK2 PsEFE)	[7e]
V270T	2OG binding (hydrophobic)	**Reduced** (compared to wt PK2 PsEFE)	[1]
R277K	2OG binding	**Yes** (no effect compared to wt PK2 PsEFE)	[7b]
F278Y*	Surface residue	**Yes** (no effect compared to wt PK2 PsEFE)	[8]
C280F	Surface/possible interaction with M54	**Reduced** (compared to wt PK2 PsEFE)	[1]
A281R	2OG binding (hydrophobic)	**Reduced** (compared to wt PK2 PsEFE)	[7a]
A281V	2OG binding (hydrophobic)	**Reduced** (compared to wt PK2 PsEFE)	[7a]
F283A/R/V/Y	2OG binding (hydrophobic)	**Reduced** (compared to wt PK2 PsEFE)	[7a]
H284Q	Structural (interior)	**Reduced** (compared to wt PK2 PsEFE)	[7e]
E285A	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[7a]
E285Q	Structural (interaction with Y306)	**Reduced** (compared to wt PK2 PsEFE)	[7a]
I304N*	Surface residue	**No**	[8]
H305Q	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[7e]
H309Q	Structural (interior)	**Reduced** (compared to wt PK2 PsEFE)	[7e]
Y306A/F	Interaction with Y192 and F283	**Reduced** (compared to wt PK2 PsEFE)	[7a]
F310R	Interaction with Y192 and F283	**Reduced** (compared to wt PK2 PsEFE)	[7a]
F310W	Interaction with Y192 and F284	**Yes** (no effect compared to wt PK2 PsEFE)	[7a]
R316A/K	*L*-arginine binding	**Reduced** (compared to wt PK2 PsEFE)	[1]
C317A	*L*-arginine binding	**Reduced** (compared to wt PK2 PsEFE)	[1]
C317N	*L*-arginine binding	**Reduced** (compared to wt PK2 PsEFE)	[7a]
C317S	*L*-arginine binding	**Reduced** (compared to wt PK2 PsEFE)	[1]
Y318F	*L*-arginine binding	**Reduced** (compared to wt PK2 PsEFE)	[1]
I322V*	Surface residue	**Enhanced** (compared to wt PK2 PsEFE)	[8]
R326S[#]	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[9]
H335Q	Surface residue	**Reduced** (compared to wt PK2 PsEFE)	[7e]

Supporting Table 3.S2. Data collection and refinement statistics for the 1449B PsEFE:Mn:2OG complex.

1449B PsEFE:Mn:2OG	
PDB ID	**9SCF**
Data collection	
Space group	P 43
Cell dimensions	
a, *b*, *c* (Å)	56.063, 56.063, 100.988
α, β, γ (°)	90, 90, 90
Resolution (Å)	56.08 - 1.65 (1.68 - 1.65)
R_{merge}	0.099 (1.514)
I / sI	10.7 (1.1)
CC (1/2)	0.999 (0.867)
Completeness (%)	99.8 (95.6)
Redundancy	11.5 (11.5)
Refinement	
No. unique reflections	37156 (1921)
R_{work} / R_{free}	0.172 (0.307) / 0.198 (0.324)
No. atoms	
Protein	2688
Ligand/ion	11
Water	282
B-factors	
Protein	39.92
Ligand/ion	36.25
Water	42.34
R.m.s. deviations	
Bond lengths (Å)	0.010
Bond angles (°)	1.06

*Values in parentheses are for highest-resolution shell.

Supporting Table 3.S3. Primer pairs used for site-directed mutagenesis of PK2 PsEFE to give *glycinea* PsEFE. Substitutions are highlighted in red. PCR reactions to generate the plasmid encoding *glycinea* PsEFE (PK2 PsEFE bearing the R326S mutation) utilized the primer pair shown below. A standard hot-start PCR protocol was employed for site-directed mutagenesis of PsEFE: an annealing temperature of 60 °C was used, the elongation time was set to 5 min.

PsEFE variant		Primers pairs (5'-3')	Length (nt)	GC content (%)	Annealing temperature (°C)
R326S	Fwd	CACCACCAAA**AGC**ATCCACAAGG	23	52.0	66.0
	Rev	ATACGATCCGGATAACAACG	20	45.0	62.0

3.12.3 Supporting Information References

[1] Z. Zhang, T. J. Smart, H. Choi, F. Hardy, C. T. Lohans, M. I. Abboud, M. S. W. Richardson, R. S. Paton, M. A. McDonough, C. J. Schofield, *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 4667–4672.

[2] X. Robert, P. Gouet, *Nucleic Acids Res.* **2014**, *42*, W320–W324.

[3] S. Dhingra, Y. Sun, Z. Zhang, C. J. Schofield, L. Brewitz, manuscript under review.

[4] a) A. J. Copper, A. G. Redfield, *J. Biol. Chem.* **1975**, *250*, 527–532; b) T. S. Viswanathan, R. E. Johnson, H. F. Fisher, *Biochemistry* **1982**, *21*, 339–345.

[5] L. Brewitz, Y. Nakashima, C. J. Schofield, *Chem. Sci.* **2021**, *12*, 1327–1342.

[6] S. Dhingra, Z. Zhang, C. T. Lohans, L. Brewitz, C. J. Schofield, *J. Biol. Chem.* **2024**, *300*, 107546.

[7] a) S. Martinez, M. Fellner, C. Q. Herr, A. Ritchie, J. Hu, R. P. Hausinger, *J. Am. Chem. Soc.* **2017**, *139*, 11980–11988; b) R. A. Copeland, S. Zhou, I. Schaperdoth, T. K. C. Shoda, J. M. Bollinger, C. Krebs, *Science* **2021**, *373*, 1489–1493; c) E. J. Burke, R. A. Copeland, Y. Dixit, C. Krebs, J. M. Bollinger, Jr., *J. Am. Chem. Soc.* **2024**, *146*, 1977–1983; d) R. A. Copeland, K. M. Davis, T. K. C. Shoda, E. J. Blaesi, A. K. Boal, C. Krebs, J. M. Bollinger, Jr., *J. Am. Chem. Soc.* **2021**, *143*, 2293–2303; e) K. Nagahama, K. Yoshino, M. Matsuoka, S. Tanase, T. Ogawa, H. Fukuda, *J. Ferment. Bioeng.* **1998**, *85*, 255–258.


[8] N. Johansson, K. O. Persson, C. Larsson, J. Norbeck, *BMC Biochem.* **2014**, *15*, 22.

[9] H. Weingart, B. Völksch, M. S. Ullrich, *Phytopathology* **1999**, *89*, 360–365.

3.13 Statement of Authorship

Title of Paper	**Structural and functional validation of *Pseudomonas savastanoi* ethylene forming enzymes reveals flexibility in 2-oxoglutarate binding mode and conformation.**
Publication Status	□ Published □ Accepted for Publication X Submitted for Publication □ Unpublished and unsubmitted work written in a manuscript style
Publication Details	Sun, Y.; Dhingra, S.; Allen, M. D.; Cheng, E. S. Z.; Zhang, Z.; Schofield, C. J.; Brewitz, L. Structural and functional validation of *Pseudomonas savastanoi* ethylene forming enzymes reveals flexibility in 2-oxoglutarate binding mode and conformation. **Manuscript under review in** *ChemBioChem*.

Student Confirmation

Student Name:	Siddhant Dhingra		
Contribution to the Paper	The experimental work in this manuscript was performed by myself and Mr. Yihong Sun and Mr. Ethan Sip under my laboratory supervision. I conducted all the bioinformatic analyses, NMR data analysis and visualisation described here. Dr. Mark Allen solved and refined the 1449B crystal structure. The manuscript was prepared by myself and Dr. Lennart Brewitz under the guidance of Prof. Christopher J. Schofield. Dr. Zhihong Zhang provided the wt PsEFE plasmid and the initial training for protein expression and purification. Dr. Lennart Brewitz, Prof. Christopher J. Schofield, and I jointly conceptualised the project.		
Signature 		Date	05-10-2025

Supervisor Confirmation

By signing the Statement of Authorship, you are certifying that the candidate made a substantial contribution to the publication, and that the description described above is accurate.

Supervisor name and title: Prof. Christopher J. Schofield		
Supervisor comments I'm happy to state that the candidate made the major and a very substantial contribution to the publication, and that the description described above is accurate.		
Signature C. J. Schofield	Date	25th September 2025

CHAPTER 4

MANUSCRIPT III

Expanding the alkene forming capability of the *Pseudomonas savastanoi* ethylene-forming enzyme

Contents

4.1 Overview **208**

4.2 Table of Contents Graphic **209**

4.3 Abstract **209**

4.4 Introduction **210**

4.5 Results **214**

4.5.1 Reactivity of L206 PsEFE variants with 2OG **214**

4.5.2 L206 PsEFE variants catalyse oxidation of C4-substituted 2OG derivatives **216**

4.5.2 L206 PsEFE variants catalyse alkene formation from C4-substituted 2OG derivatives **219**

4.6 Discussion **222**

4.7 Conclusions **224**

4.8 Experimental Section **225**

4.9 Acknowledgements **226**

4.10 Author Contributions **226**

4.11 References **227**

4.12 Supporting Information **230**

4.12.1 Supporting Figures **230**

4.12.2 Supporting Tables **248**

4.12.3 Supporting Information References **248**

4.13 Statement of Authorship **249**

4.1 Overview

Mutagenesis studies on different ethylene-forming enzymes, including ACCO, and nitrogenases, have shown that these enzymes can be engineered for producing alkenes other than ethylene (Section 1.8).[8e, 8f, 29, 30] The broad cosubstrate-scope of *Pseudomonas savastanoi* ethylene-forming enzyme, PsEFE, and the dependence of its product partitioning on the steric bulk of the C4-substituent of 2OG derivatives, as described in Chapter 2, suggests that PsEFE might be engineered for producing higher alkenes using site-directed mutagenesis.


This chapter discusses qualitative biochemical studies on PsEFE variants bearing bulk-diminishing mutations at the Leu206 residue, aimed at enabling productive binding of C4-substituted 2OG derivatives. ^{1}H NMR assays and product analysis with authentic standards show that L206 PsEFE variants are capable of producing higher alkenes; L206V PsEFE produces propylene and acetaldehyde from 4-methyl-2OG and 4-hydroxy-2OG respectively; L206A catalyses butylene formation from 4-ethyl-2OG. Collectively, our results highlight the potential of PsEFE for engineering biocatalysts capable of producing alkenes.

At the time of writing this thesis, Manuscript III has been submitted to *ChemCatChem* and is under review.

Dhingra, S.; Sun, Y.; Zhang, Z.; Schofield, C. J.; Brewitz, L. Expanding the alkene forming capability of the *Pseudomonas savastanoi* ethylene-forming enzyme. **Manuscript under review in** *ChemCatChem*.

Each author's contribution is described in the Author Contributions (Section 4.10). A Statement of Authorship is appended at the end of the chapter (Section 4.13).

4.2 Table of Contents Graphic



Bump-and-hole studies on the *Pseudomonas savastanoi* ethylene-forming enzyme (PsEFE) enable formation of low levels of alkene products other than ethylene from naturally occurring 2-oxoglutarate derivatives. The results show that plants and/or microorganisms have potential for enzyme-catalysed production of propylene and 1-butylene, potentially to exert distinctive signalling effects.

4.3 Abstract

Ethylene is an established signalling molecule in plants and other organisms, however, the biosynthesis and biological roles of gaseous alkenes other than ethylene are less well defined. The *Pseudomonas savastanoi* ethylene/succinate-forming enzyme (PsEFE) catalyses ethylene production from 2-oxoglutarate (2OG), though does not catalyse formation of alkenes from C4 alkyl- or hydroxyl-substituted 2OG derivatives. Here we report studies on the reactivity of L206 PsEFE variants with C4-substituted 2OG derivatives. Spectroscopic evidence reveals that L206V and L206A PsEFE react with C4-substituted 2OG derivatives to give diacid and alcohol products, similarly to wildtype (wt) PsEFE. Importantly, L206V PsEFE, but not L206A or L206G PsEFE, catalysed production of relatively low levels of acetaldehyde and propylene from the natural metabolites 4-hydroxy-2OG and 4-methyl-2OG, respectively. By contrast,

L206A PsEFE, but not L206V and L206G PsEFE, catalysed formation of low levels of 1-butylene from 4-ethyl-2OG. Together with studies from others, the combined results indicate the potential of bump-and-hole studies to modify the substrate and product selectivities of PsEFE reactions, provided that the PsEFE variant:2OG derivative pairs are matched. The results suggest that wildtype 2OG oxygenases other than PsEFE may catalyse production of gaseous alkenes other than ethylene.

4.4 Introduction

2-Oxoglutarate (2OG) and Fe(II) dependent oxygenases catalyse 2-electron oxidations of their substrates, a process typically coupled to 2OG oxidation, producing succinic acid and CO_2.[1] 2OG oxygenases share a characteristic distorted double-stranded β-helix (DSBH) fold which, together with other conserved secondary structure elements, they use to form Fe(II) and 2OG binding sites; there is more variation in the modes of prime substrate binding.[2] 2OG oxygenases have important functions *inter alia* in plant, fungal and bacterial ethylene production.[3]

Ethylene has validated signalling roles in plant development,[4] whereas its functions in bacteria and fungi are less well defined.[5] In seeding plants, 1-aminocyclopropane-1-carboxylate (ACC) oxidase (ACCO) catalyses production of ethylene from ACC.[3b,6] Although ACCO belongs to the 2OG oxygenase superfamily, it does not require 2OG as a cosubstrate for productive ACC turnover.[3,7]

Bacteria and fungi employ enzymes other than ACCOs to catalyse ethylene production,[3a,8] including ethylene/succinate-forming enzymes (EFEs) which catalyse fragmentation of 2OG to give ethylene and CO_2.[9] The EFE of *Pseudomonas savastanoi* (previously: *P. syringae*) pv. *phaseolicola* PK2 (*i.e.*, PsEFE)[9f,9g] not only catalyses oxidative fragmentation of 2OG to give ethylene,[9f,10] but also catalyses low levels of 2OG fragmentation to give 3-hydroxypropionic

acid and CO_2.[9b,9e,11] Both of these PsEFE reactions require *L*-arginine to proceed efficiently.[11-12] PsEFE also employs 2OG as a cosubstrate in a more typical 2OG oxygenase reaction involving C5 hydroxylation of *L*-arginine to give succinic acid, CO_2, H_2O, guanidine, and *L*-Δ^1-pyrroline-5-carboxylic acid (P5C), the later likely produced via *L*-glutamate-5-semialdehyde formed via fragmentation of nascent 5-hydroxyarginine (Figure 4.1a).[10,12a] PsEFE also catalyses oxidation of 2OG to give succinic acid and CO_2 uncoupled from *L*-arginine substrate oxidation, though not efficiently.[9b,9e]




Figure 4.1. The wildtype PsEFE substrate scope. (**a**) Wildtype (wt) PsEFE reacts with 2OG to give ethylene and succinic acid as major products, with formation of succinic acid being coupled to C5 oxidation of *L*-arginine to give guanidine and P5C;[9f,10] note that low levels of an alcohol product have also been observed.[9b,9e,11] (**b**) wt PsEFE reacts with C3/C4-alkyl/hydroxy-substituted 2OG derivatives to give a diacid (coupled to P5C formation) and an alcohol, but not an alkene product.[13] (**c**) View from a wt PsEFE:Mn:2OG:arginine crystal structure (with Mn substituting for catalytically active Fe(II); PDB ID: 5V2Y[14]) showing that the methyl groups of L206 are proximate with the 2OG C4 methylene group (dotted red lines) suggesting that reducing steric bulk at residue-206 may affect the conformation of C4-substituted 2OG derivatives at the active site. Polar interactions with 2OG and Mn are shown by dotted black lines.

Recent work on the (co)substrate scope of wildtype (wt) PsEFE shows that it can employ certain C3- and/or C4-substituted 2OG derivatives to catalyse *L*-arginine oxidation in the absence of 2OG to give diacid products (Figure 4.1b).[13] This property is reminiscent of the cosubstrate flexibility manifested by at least some other 2OG oxygenases.[15] Additionally, wt PsEFE was observed to catalyse formation of an alcohol product with some, but not all, of the tested 2OG derivatives, with the 2OG substitution pattern regulating the reaction outcomes (Figure 4.1b).[13] By contrast, wt PsEFE did not catalyse formation of alkene products from any of the tested C3- and/or C4-substituted 2OG diacid derivatives with the exception of fluoroethylene from 4-fluoro-2OG.[13a]

Analysis of PsEFE:2OG complex crystal structures reveals that side chains of hydrophobic amino acid residues (*e.g.*, L173, F175, L206, V270, A279) help shape the 2OG binding site, likely stabilising 2OG binding via hydrophobic interactions with the 2OG C3 and C4 methylene groups (Figure 4.1, Supporting Figure 4.S1),[14,16] with the L206 side chain methyl groups being proximate with the 2OG C4 carbon (distances: 4.1 and 4.8 Å) (Figure 4.1). Altering enzyme active site geometries using site-directed mutagenesis to reduce/increase steric bulk proximate to the substrate is an established approach to modify reaction outcomes of wt enzymes with (co)substrate derivatives (*i.e.*, the bump-and-hole strategy),[17] including for 2OG oxygenases.[18] Recent studies with L206V PsEFE have revealed its potential to catalyse propylene formation from 4-methyl-2OG.[11] However, the reactivity of L206V PsEFE with C4-substituted 2OG derivatives other than 4-methyl-2OG was not investigated in this work.[11]

Here we report on the reactivity of L206V, L206A, and L206G PsEFE with five C4-substituted 2OG derivatives.[15c] The results reveal that, in addition to the reported propylene formation from 4-methyl-2OG,[11] L206V PsEFE catalyses formation of acetaldehyde from the natural metabolite[19] 4-hydroxy-2OG. Notably, L206A PsEFE, but not L206V or L206G PsEFE,

catalyse 1-butylene formation from 4-ethyl-2OG. The combined results indicate that active site mutation studies with PsEFE have potential to enable formation of novel products, including 1-butylene, from non-native substrates, provided that the 2OG derivative : PsEFE variant pairs are matched.

4.5 Results

4.5.1 Reactivity of L206 PsEFE variants with 2OG

The L206V, L206A, and L206G PsEFE variants (UNIPROT ID: P32021; EC:1.13.12.19)[9f,9g] were produced in >90% purity, as judged by SDS-PAGE and MS analyses (Supporting Figure 4.S2), to investigate whether altering the steric bulk at residue-206 can affect reaction outcomes with C4-substituted 2OG derivatives. Initially, the reactivities of the three isolated recombinant L206 PsEFE variants were investigated with 2OG as a (co)substrate together with *L*-arginine using a reported ^{1}H NMR-based assay.[13b] These assays enable quantification of 2OG/2OG derivative consumption and product formation by comparing signals to that of an internal standard (*i.e.*, 3-(trimethylsilyl)-2,2,3,3-tetradeuteropropionic acid,[20] TMSP-d_4), and product identification by comparison with authentic standards.

^{1}H NMR analysis (700 MHz) of reactions of L206V, L206A, and L206G PsEFE with 2OG and *L*-arginine reveals that all of these PsEFE variants were less active than wt PsEFE: the efficiency of PsEFE-variants, as measured by depletion of the 2OG peak, decreased with decreasing steric bulk at residue-206 in the order: L206 (wt) > V206 > A206 > G206, with L206G PsEFE being inactive under the tested conditions (Figure 4.2). Note that analysis of wt, L206V, L206A, and L206G PsEFE by circular dichroism (CD) spectroscopy indicates that substituting L206 did not affect the overall PsEFE fold (Supporting Figure 4.S3). There was no evidence accrued for formation of substantial amounts of 2OG-derived products other than ethylene and succinic acid (Supporting Figure 4.S4), including 3-hydroxypropionic acid, *i.e.*,

the low level alcohol product of the wt PsEFE reaction with 2OG.[9b,13a] This observation contrasts with the reported detection of 3-hydroxypropionic acid from reactions of L206V PsEFE with 2OG by infrared spectroscopy and LCMS assays, possibly reflecting differences in assay method sensitivity and/or assay conditions.[11]

The estimated succinic acid : ethylene ratio following reaction of L206V and L206A PsEFE with 2OG was similar to that reported for wt PsEFE (~1:1) (Figure 4.2).[13b] P5C formation was not quantified, because the abundance of its characteristic singlet (δ ~ 7.70 ppm) apparently varied over time.[13b] Experiments in the absence of *L*-arginine indicated that L206V and L206A PsEFE, like wt PsEFE, require *L*-arginine for, at least, efficient 2OG oxidation (Supporting Figures 4.S5).




Figure 4.2. Substitutions at PsEFE residue-206 influence reactivity with 2OG. Reactions of: (**a**) L206V PsEFE, (**b**) L206A PsEFE, and (**c**) L206G PsEFE with 2OG. (**d**) Outcomes of L206 PsEFE variant reactions with 2OG compared to that reported for wt PsEFE.[13b] Ethylene formation was estimated by subtracting the normalised integral of succinic acid (obtained via comparison with TMSP-d_4) from that of 2OG. The time-scales were calibrated to the end of the acquisition of the first ^{1}H NMR experiment following PsEFE addition to the reaction mixture (t = 0 min), by which time low levels of conversion were manifest. Results are means of independent triplicates (n = 3; mean ± standard deviation, SD). Conditions: 400 µM 2OG, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and L206 PsEFE variant (10 µM) or wt PsEFE (2 µM)[13b] in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O).

4.5.2 L206 PsEFE variants catalyse oxidation of C4-substituted 2OG derivatives

Having validated that L206V and L206A PsEFE can catalyse 2OG oxidation, we next investigated the ability of five representative C4-substituted 2OG derivatives (all as racemates) to sustain catalysis by L206V, L206A, and L206G PsEFE. ^{1}H NMR analysis (700 MHz) reveals that L206V and L206A PsEFE reacted with 4-methyl-2OG (**1**), 4,4-dimethyl-2OG (**2**), 4-hydroxy-2OG (**3**), 4-ethyl-2OG (**4**), and 4-propyl-2OG (**5**) to give both the corresponding

diacid and alcohol products (Figure 4.3, Supporting Figures 4.S6-4.S10), as reported for wt PsEFE.[13] Note that the stereochemistry of the products was not assigned. By contrast, L206G PsEFE did not catalyse conversion of any of the 2OG derivatives **1-5** within 2 h (Figure 4.3, Supporting Figures S6-S10). Experiments in the absence of *L*-arginine indicated that the three L206 PsEFE variants require *L*-arginine for, at least, efficient oxidation of C4-substituted 2OG derivatives (Supporting Figures 4.S13-4.S15).




Figure 4.3. L206 PsEFE variants employ C4-substituted 2OG derivatives as (co)substrates. Outcomes of the reaction of L206 PsEFE variants with: (**a**) 4-methyl-2OG (**1**), (**b**) 4,4-dimethyl-2OG (**2**), (**c**) 4-hydroxy-2OG (**3**), (**d**) 4-ethyl-2OG (**4**), (**e**) 4-propyl-2OG (**5**), and (**f**) 3-methyl-2OG (**6**) compared to that reported for wt PsEFE.[13b] Results are means of independent triplicates (n = 3; mean ± SD); ^{1}H NMR spectra are shown in Supporting Figures 4.S6-4.S11. Conditions: 400 µM 2OG derivative, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Note that the stereochemistry of the products was not assigned.

The results indicate that, in general, L206V and L206A PsEFE reacted less efficiently with the five C4-substituted 2OG derivatives **1-5** than wt PsEFE (Figure 4.3).[13b] Notable exceptions

were the reactions of L206V PsEFE with racemic 4-methyl-2OG (**1**) and 4-propyl-2OG (**5**), which were equally or more efficient than those reported with wt PsEFE,[13b] giving 2-methylsuccinic acid (**1a**)/β-hydroxyisobutyric acid (**1b**) and 2-propylsuccinic acid (**5a**)/2-(hydroxymethyl)pentanoic acid (**5b**), respectively.

Analysis of the diacid : alcohol ratios for the reactions of L206V and L206A PsEFE with 2OG derivatives **1**-**5** reveals that the nature of the L206 variant did not substantially affect product ratios. The diacid : alcohol ratios for the reactions of L206V and L206A PsEFE with **1**-**5** matched those reported for wt PsEFE (*i.e.*, ~1:1 for **2**; ~2:1 for **3**; ~2:1 for **4**; ~3:2 for **5**); the exception was the product ratios obtained with **1**, which were ~1:1 for L206V and L206A PsEFE, but ~2:1 for wt PsEFE (Figure 4.3).[13b]

4.5.2 L206 PsEFE variants catalyse alkene formation from C4-substituted 2OG derivatives

Importantly, ^{1}H NMR analyses of the L206V and L206A PsEFE reactions with 2OG derivatives **1**-**5** not only revealed diacid and alcohol formation, but also the formation of relatively low levels of alkene products from the 4-methyl-2OG (**1**), 4-hydroxy-2OG (**3**), and 4-ethyl-2OG (**4**) 2OG derivatives (Supporting Figures 4.S16-4.S18). In accord with reported work,[11] L206V PsEFE catalysed formation of propylene from **1**, as supported by direct comparison with an authentic sample of propylene (Figure 4.4). Note that alkene formation was not directly quantified by ^{1}H NMR, because alkenes may evaporate from the reaction mixture; in some cases, overlapping peaks (*e.g.*, the peaks of the propylene methyl group and the C4 methylene group of *L*-arginine; Figure 4.4) further complicated quantification of alkene formation. The alkene levels were thus approximated by subtracting the normalised integral of the diacid and alcohol products (obtained via comparison with TMSP-d_4) from that of **1** because no substantial amounts of additional products derived from **1** were detected and because diacid formation appeared to be mostly coupled to *L*-arginine oxidation.[13] By contrast

with L206V PsEFE, L206A and L206G PsEFE (like wt PsEFE[13b]) did not catalyse formation of propylene from **1**, possibly reflecting their substantially reduced reactivity with **1** compared to that of L206V PsEFE.




Figure 4.4. L206V PsEFE catalyses production of propylene from 4-methyl-2OG (1). (**a-d**) ^{1}H NMR analysis (~0.5 to ~4.0 ppm and ~5.5 to ~8.5 ppm) for ~90 min incubations of 4-methyl-2OG (**1**) with: (**a**) L206G PsEFE, (**b**) L206A PsEFE, and (**c**) L206V PsEFE. (**d**) Reported ^{1}H NMR analysis of ~90 min incubation of **1** with wt PsEFE.[13b] (**e**) Reported ^{1}H NMR spectrum of **1** under standard conditions in the absence of PsEFE.[13b] (**f**) ^{1}H NMR spectrum of an authentic sample of propylene under assay conditions. Note that the C1 alkene protons of propylene were not observed, likely because they coincide with the HOD peak. (**g**) ^{1}H NMR spectrum of an authentic sample of propylene in buffer. (**h**) ^{1}H NMR spectra (~5.8 to ~6.0 ppm and ~1.6 to ~1.8 ppm) showing that L206V PsEFE catalysed propylene production from **1**. Conditions: 400 μM **1**, 500 μM *L*-arginine, 500 μM *L*-ascorbic acid, 50 μM Fe(II), 800 μM TMSP-d_4, and 10 μM PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris buffer. Note that the stereochemistry of the products was not assigned. Representative spectra of independent triplicates are shown.

Like wt PsEFE,[13b] L206V, L206A, and L206G PsEFE did not catalyse formation of propylene from 2-oxoadipate (2OA) and 3-methyl-2OG (**6**), the regioisomer of **1**, within 2 h (Figure 4.3; Supporting Figures 4.S11-4.12). None of the L206 PsEFE variants catalysed isobutene

production from 4,4-dimethyl-2OG (**2**) within 2 h (Supporting Figure 4.S6), in accord with the reported reactivity of wt PsEFE with **2**.[13b]

L206V PsEFE not only catalysed propylene formation from 4-methyl-2OG (**1**), but also catalysed production of relatively low levels of acetaldehyde from 4-hydroxy-2OG (**3**); note that we observed no evidence for formation of pyruvate via the possible retroaldol reaction of **3** (Supporting Figure 4.S8). Neither L206A, L206G, nor wt PsEFE[13b] catalysed acetaldehyde formation from **3** (Supporting Figure 4.S7). By contrast, analysis of the reaction of L206A PsEFE, but not of L206V, L206G, and wt PsEFE,[13b] showed evidence for formation of relatively low levels of 1-butylene from 4-ethyl-2OG (**4**), as manifest by peaks for the methyl group of 1-butylene (Supporting Figure 4.S9). The results suggest that the C4 ethyl group of **4** requires more space in proximity to the PsEFE residue-206 relative to the methyl group of **1** in order to adopt a conformation enabling alkene formation. Consistent with this proposal, none of the L206 PsEFE variants catalysed formation of 1-pentene from 4-propyl-2OG (**5**) (Supporting Figure 4.S10)

4.6 Discussion

Studies have indicated that plants and/or microorganisms emit gaseous alkenes other than ethylene, however, it is unclear whether all of these alkenes are produced in an enzyme-catalysed manner.[21] The effects of these alkenes on plant signalling can differ from those of ethylene.[22] Pioneering work with fruit extracts on the substrate scope of ACCOs revealed the possibility of ACCO-catalysed propylene and 1-butylene production from naturally occurring C-alkylated ACC derivatives.[23] By contrast, studies on the substrate scope of wt PsEFE have indicated that it does not catalyse propylene production from 2-oxoadipate, 4-methyl-2OG (**1**), or 3-methyl-2OG (**6**),[12b,13-14] although wt PsEFE catalysed production of fluoroethylene from

4-fluoro-2OG,[13a] a reaction likely lacking biological significance as 4-fluoro-2OG is not a naturally occurring cellular metabolite.

The results presented here show that decreasing steric bulk at PsEFE residue-206 can enable the capacity to catalyse production of alkenes, *i.e.*, propylene from the natural product[24] **1** and 1-butylene from 4-ethyl-2OG (**4**), as well as acetaldehyde from the metabolite[19] 4-hydroxy-2OG (**3**) (Figure 4.3), which is formed in cells as an intermediate of 4-hydroxyproline catabolism.[25] They confirm the recently reported ability of L206V PsEFE to catalyse propylene production from **1**.[11] Thus, the biocatalytic potential of PsEFE variants extends beyond catalysing production of ethylene, diacids, and alcohols[13,26] to production of alkenes other than ethylene and aldehydes which are commodities.[26b,27] L206V and L206A PsEFE likely catalyse alkene formation from C4-substituted 2OG derivatives via a radical Grob-type mechanism similar to that proposed for the reaction of wt PsEFE with 2OG.[11,13a,28] The reaction of L206V PsEFE with **3** might involve transient production of vinyl alcohol giving acetaldehyde following tautomerisation.

The observations: (i) that L206V PsEFE, but not L206A, L206G, and wt PsEFE,[13b] catalysed propylene formation from **1**, (ii) that the tested L206 PsEFE variants did not catalyse formation of propylene and isobutylene from **6** and 4,4-dimethyl-2OG (**2**), respectively, and (iii) that L206A PsEFE, but not L206V, L206G, and wt PsEFE,[13b] catalyse formation of 1-butene from **4** suggest matching of specific 2OG derivative:PsEFE variant pairs is required to induce novel PsEFE reactivity (Figure 4.3). Thus, there is potential for design of PsEFE variants which catalyse alkene and/or acetaldehyde formation more efficiently than the L206 PsEFE variants tested in this work and for bump-and-hole studies with PsEFE to enable formation of (novel) products from 2OG derivatives by varying PsEFE residues other than 206.

Aside from the work on wt ACCO, the ability of substitutions in the PsEFE active site to induce capacity for production of alkenes from C4-substituted 2OG derivatives is precedented by work on nitrogenases which catalyse ethylene production from acetylene in their wildtype form,[8e,8f,29] as substitutions in the active site of nitrogenases enabled production of propylene from propyne.[30] The combined evidence highlight the importance of second coordination sphere residues in influencing reactivity of mechanistically unrelated ethylene-producing nitrogenases[31] and PsEFE,[32] as is also the case for typical 2OG oxygenases.[33] Thus, mutations beyond bump-and-hole type studies reported here and elsewhere will likely be required to optimise formation of specific PsEFE products including alkenes. Given that the alkene products can be measured by headspace analysis, a forced-evolution approach[34] (taking into account substrate stereochemistry) may be particularly productive.

The combined results on the reactivity of the L206 PsEFE variants and the observation that disease-associated variations in the active sites of certain human 2OG oxygenases can alter their reactivity with 2-oxoacids[35] raises the question as to whether 2OG oxygenase(s) can catalyse alkene formation from naturally abundant C3- and/or C4-substituted 2OG derivatives in their wildtype form. To our knowledge, L206V/A (equivalent) PsEFE variants have not yet been identified in microbes.

4.7 Conclusions

The results reveal L206V and L206A PsEFE have capacity to catalyse production of alkenes other than ethylene, including propylene and 1-butylene, from 2OG-derived substrates, including from natural products. L206V PsEFE also catalysed low levels of acetaldehyde formation from the abundant metabolite 4-hydroxy-2OG (**3**). The combined results further highlight the remarkable versatility of the PsEFE reaction with respect to 2-oxoacid oxidation. They suggest that plants and/or microorganisms may have potential to produce gaseous alkenes

other than ethylene in an enzyme-catalysed manner, possibly to exert signalling effects distinct from those of ethylene.

4.8 Experimental Section

Protein production. Plasmid DNA encoding for the L206V, L206A, and L206G PsEFE variants was obtained from our reported wt PsEFE construct[16a] by standard site-directed mutagenesis (Supporting Table S1). The L206V, L206A, and L206G PsEFE variants were produced and purified as reported for wt PsEFE.[13b,16a] The L206 PsEFE variants were >90% pure by SDS-PAGE analysis and had the anticipated masses (Supporting Figure 4.S2). They were stored at −70 ºC, at ~20-25 mg/mL concentrations.

2OG derivatives. 2-Oxoglutarate (2OG), 2-oxoadipate (2OA), and racemic 4-hydroxy-2OG (**3**) were from Sigma-Aldrich. 4-Methyl-2OG (**1**), 4,4-dimethyl-2OG (**2**), 4-ethyl-2OG (**4**), 4-propyl-2OG (**5**), and 3-methyl-2OG (**6**) were synthesised as reported;[15c] chiral 2OG derivatives were prepared as racemic mixtures.

^{1}H NMR assays. ^{1}H NMR assays were performed as reported,[13b] typically using: 2OG derivative (400 μM), *L*-arginine (500 μM), *L*-ascorbic acid (500 μM), Fe(II) (50 μM), TMSP-d_4 (800 μM), and L206 PsEFE variant (10 μM) in buffer (50 mM sodium phosphate, pH 7.4, 10%$_{v/v}$ D_2O), if not specified otherwise in the Figure legends. Freshly thawed aliquots of L206 PsEFE variants were used for all experiments. A Bruker AVIII 700 MHz NMR spectrometer with a 5-mm inverse triple-resonance-inverse cryoprobe was used for analysis. The water peak in ^{1}H NMR experiments was suppressed using the perfect echo-modified WATERGATE solvent suppression method.[36]

4.9 Acknowledgements

This research was funded in part by the Wellcome Trust (106244/Z/14/Z). For the purpose of open access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission. We thank the Biotechnology and Biological Sciences Research Council (BB/J003018/1 and BB/R000344/1) for funding and Prof. Dr. C. Redfield, Dr. C. Mycroft, and L. Harrington for helpful discussions.

4.10 Author Contributions

S.D. and Y.S. produced enzymes; S.D. and Y.S. performed assays; all authors analysed data; L.B. conceived the project together with S.D.; L.B., Z.Z., and C.J.S. supervised the research; L.B., S.D. and C.J.S. wrote the manuscript with help from all authors.

4.11 References

[1] *2-Oxoglutarate-dependent oxygenases* (Ed.: R. P. Hausinger, C. J. Schofield), The Royal Society of Chemistry, Cambridge, United Kingdom, **2015**.

[2] a) S. Kal, L. Que, *J. Biol. Inorg. Chem.* **2017**, *22*, 339–365; b) W. Aik, M. A. McDonough, A. Thalhammer, R. Chowdhury, C. J. Schofield, *Curr. Opin. Struct. Biol.* **2012**, *22*, 691–700.

[3] a) R. P. Hausinger, S. B. J. S. Rifayee, M. G. Thomas, S. Chatterjee, J. Hu, C. Z. Christov, *RSC Chem. Biol.* **2023**, *4*, 635–646; b) M. Houben, B. Van de Poel, *Front. Plant Sci.* **2019**, *10*, 695.

[4] a) B. M. Binder, *J. Biol. Chem.* **2020**, *295*, 7710–7725; b) X. Wang, H. Wen, A. Suprun, H. Zhu, *Plants* **2025**, *14*, 309; c) B. Van de Poel, E. D. Cooper, C. F. Delwiche, C. Chang, in *Ethylene in Plants* (Ed.: C.-K. Wen), Springer Netherlands, Dordrecht, **2015**, pp. 109–134; d) B. Van de Poel, J. de Vries, *Environ. Exp. Bot.* **2023**, *214*, 105456.

[5] a) M. Ravanbakhsh, R. Sasidharan, L. A. C. J. Voesenek, G. A. Kowalchuk, A. Jousset, *Microbiome* **2018**, *6*, 52; b) N. Papon, B. M. Binder, *Trends Microbiol.* **2019**, *27*, 193–196; c) B. Bidon, S. Kabbara, V. Courdavault, G. Glévarec, A. Oudin, F. Héricourt, S. Carpin, L. Spíchal, B. M. Binder, J. M. Cock, N. Papon, *Cells* **2020**, *9*, 2526; d) I. Camehl, I. Sherameti, Y. Venus, G. Bethke, A. Varma, J. Lee, R. Oelmüller, *New Phytol.* **2010**, *185*, 1062–1073; e) V. Chagué, in *Handbook of Hydrocarbon and Lipid Microbiology* (Ed.: K. N. Timmis), Springer Berlin Heidelberg, Berlin, Heidelberg, **2010**, pp. 3011–3020; f) S. Gu, L. Xie, Q. Guan, X. Sheng, Y. Fang, X. Wang, *Int. J. Food Microbiol.* **2024**, *418*, 110729.

[6] a) A. J. Hamilton, G. W. Lycett, D. Grierson, *Nature* **1990**, *346*, 284–287; b) Z. Zhang, J.-S. Ren, I. J. Clifton, C. J. Schofield, *Chem. Biol.* **2004**, *11*, 1383–1394.

[7] J. G. Dong, J. C. Fernández-Maculet, S. F. Yang, *Proc. Natl. Acad. Sci. USA* **1992**, *89*, 9789–9793.

[8] a) C. C. Lee, Y. Hu, M. W. Ribbe, *Science* **2010**, *329*, 642–642; b) Y. Hu, C. C. Lee, M. W. Ribbe, *Science* **2011**, *333*, 753–755; c) C. C. Lee, Y. Hu, M. W. Ribbe, *Angew. Chem. Int. Ed.* **2011**, *50*, 5545–5547; d) Y. Hu, C. C. Lee, M. Grosch, J. B. Solomon, W. Weigand, M. W. Ribbe, *Chem. Rev.* **2023**, *123*, 5755–5797; e) L. C. Seefeldt, Z.-Y. Yang, D. A. Lukoyanov, D. F. Harris, D. R. Dean, S. Raugei, B. M. Hoffman, *Chem. Rev.* **2020**, *120*, 5082–5106; f) B. M. Barney, R. Y. Igarashi, P. C. Dos Santos, D. R. Dean, L. C. Seefeldt, *J. Biol. Chem.* **2004**, *279*, 53621–53624; g) J. A. North, A. R. Miller, J. A. Wildenthal, S. J. Young, F. R. Tabita, *Proc. Natl. Acad. Sci. USA* **2017**, *114*, E10455–E10464; h) J. A. North, A. B. Narrowe, W. Xiong, K. M. Byerly, G. Zhao, S. J. Young, S. Murali, J. A. Wildenthal, W. R. Cannon, K. C. Wrighton, R. L. Hettich, F. R. Tabita, *Science* **2020**, *369*, 1094–1098; i) A. Lago-Maciel, J. C. Soares, J. Zarzycki, C. J. Buchanan, T. Reif-Trauttmansdorff, F. V. Schmidt, S. Lometto, N. Paczia, J. M. Schuller, D. F. Hansen, G. T. Heller, S. Prinz, G. K. A. Hochberg, A. J. Pierik, J. G. Rebelein, *bioRxiv* **2024**, 2024.2010.2019.619033.

[9] a) Y. Cui, Y. Jiang, M. Xiao, M. Z. Munir, S. Riaz, F. Rasul, M. Daroch, *Int. J. Mol. Sci.* **2022**, *23*, 4500; b) S. Chatterjee, J. A. Rankin, M. A. Farrugia, S. B. J S Rifayee, C. Z. Christov, J. Hu, R. P. Hausinger, *Biochemistry* **2025**, *64*, 2054–2067; c) H. Fukuda, T. Fujii, T. Ogawa, *Agric. Biol. Chem.* **1986**, *50*, 977–981; d) T. Hottiger, T. Boller, *Arch. Microbiol.* **1991**, *157*, 18–22; e) S. Chatterjee, J. A. Rankin, M. A. Farrugia, B. J. Delaney, N. S. Pascual, J. VanAntwerp, D. R. Woldring, J. Hu, R. P. Hausinger, *Biochemistry* **2025**, *64*, 3432–3445; f) K. Nagahama, T. Ogawa, T. Fujii, M. Tazaki, S. Tanase, Y. Morino, H. Fukuda, *J. Gen. Microbiol.* **1991**, *137*, 2281–2286; g) H. Fukuda, T. Ogawa, K. Ishihara, T. Fujii, K. Nagahama, T. Omata, Y. Inoue, S. Tanase, Y. Morino, *Biochem. Biophys. Res. Commun.* **1992**, *188*, 826–832; h) N. Johansson, K. O. Persson, C. Larsson, J. Norbeck, *BMC Biochem.* **2014**, *15*, 22.

[10] H. Fukuda, T. Ogawa, M. Tazaki, K. Nagahama, T. Fujii, S. Tanase, Y. Morino, *Biochem. Biophys. Res. Commun.* **1992**, *188*, 483–489.

[11] E. J. Burke, R. A. Copeland, Y. Dixit, C. Krebs, J. M. Bollinger, Jr., *J. Am. Chem. Soc.* **2024**, *146*, 1977–1983.

[12] a) K. Nagahama, T. Ogawa, T. Fujii, M. Tazaki, M. Goto, H. Fukuda, *J. Gen. Microbiol.* **1991**, *137*, 1641–1646; b) S. Martinez, R. P. Hausinger, *Biochemistry* **2016**, *55*, 5989–5999.

[13] a) R. A. Copeland, S. Zhou, I. Schaperdoth, T. K. C. Shoda, J. M. Bollinger, C. Krebs, *Science* **2021**, *373*, 1489–1493; b) S. Dhingra, Z. Zhang, C. T. Lohans, L. Brewitz, C. J. Schofield, *J. Biol. Chem.* **2024**, *300*, 107546.

[14] S. Martinez, M. Fellner, C. Q. Herr, A. Ritchie, J. Hu, R. P. Hausinger, *J. Am. Chem. Soc.* **2017**, *139*, 11980–11988.

[15] a) K. Majamaa, H. M. Hanauske-Abel, V. Günzler, K. I. Kivirikko, *Eur. J. Biochem.* **1984**, *138*, 239–245; b) C. J. Cunliffe, T. J. Franklin, N. J. Hales, G. B. Hill, *J. Med. Chem.* **1992**, *35*, 2652–2658; c) L. Brewitz, Y. Nakashima, C. J. Schofield, *Chem. Sci.* **2021**, *12*, 1327–1342; d) Y. Nakashima, L. Brewitz, A. Tumber, E. Salah, C. J. Schofield, *Nat. Commun.* **2021**, *12*, 6478; e) A. Tumber, E. Salah, L. Brewitz, T. Corner, C. J. Schofield, *RSC Chem. Biol.* **2023**, *4*, 399–413; f) G. Fiorini, S. A. Marshall, W. D. Figg, W. K. Myers, L. Brewitz, C. J. Schofield, *Sci. Rep.* **2024**, *14*, 26162.

[16] a) Z. Zhang, T. J. Smart, H. Choi, F. Hardy, C. T. Lohans, M. I. Abboud, M. S. W. Richardson, R. S. Paton, M. A. McDonough, C. J. Schofield, *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 4667–4672; b) R. A. Copeland, K. M. Davis, T. K. C. Shoda, E. J. Blaesi, A. K. Boal, C. Krebs, J. M. Bollinger, Jr., *J. Am. Chem. Soc.* **2021**, *143*, 2293–2303; c) S. Chatterjee, M. Fellner, J. Rankin, M. G. Thomas, S. B. J S Rifayee, C. Z. Christov, J. Hu, R. P. Hausinger, *Biochemistry* **2024**, *63*, 1038–1050.

[17] K. Islam, *Cell Chem. Biol.* **2018**, *25*, 1171–1184.

[18] M. Breski, D. Dey, S. Obringer, B. Sudhamalla, K. Islam, *J. Am. Chem. Soc.* **2016**, *138*, 13505–13508.

[19] D. Siegel, A. C. Meinema, H. Permentier, G. Hopfgartner, R. Bischoff, *Anal. Chem.* **2014**, *86*, 5089–5100.

[20] L. Pohl, M. Eckle, *Angew. Chem. Int. Ed.* **1969**, *8*, 381.

[21] a) H. Fukuda, T. Fujii, T. Ogawa, *Agric. Biol. Chem.* **1984**, *48*, 1679–1682; b) M. A. K. Khalil, R. A. Rasmussen, *J. Air Waste Manag. Assoc.* **1992**, *42*, 810–813; c) T. D. Sharkey, S. Yeh, *Annu. Rev. Plant Biol.* **2001**, *52*, 407–436.

[22] a) A. E. H. W. Crocker, P. W. Zimmerman, *Contrib. Boyce Thompson Inst.* **1935**, *7*, 231–248; b) E. J. McMurchie, W. B. McGlasson, I. L. Eaks, *Nature* **1972**, *237*, 235–236; c) G. Stavroulakis, E. Sfakiotakis, *Postharvest Biol. Technol.* **1997**, *10*, 189–194; d) T. Suzuki, T. Yazawa, N. Morishita, A. Maruyama, H. Fuse, *Microbes Environ.* **2019**, *34*, 33–42.

[23] a) N. E. Hoffman, S. F. Yang, A. Ichihara, S. Sakamura, *Plant Physiol.* **1982**, *70*, 195–199; b) M. C. Pirrung, G. M. McGeehan, *J. Org. Chem.* **1986**, *51*, 2103–2106; c) M. C. Pirrung, *Acc. Chem. Res.* **1999**, *32*, 711–718; d) J. E. Baldwin, R. M. Adlington, G. A. Lajoie, B. J. Rawlings, *J. Chem. Soc., Chem. Commun.* **1985**, 1496–1498.

[24] a) R. Díaz, H. Gallart-Ayala, J. V. Sancho, O. Nuñez, T. Zamora, C. P. B. Martins, F. Hernández, S. Hernández-Cassou, J. Saurina, A. Checa, *J. Chromatogr. A* **2016**, *1433*, 90–97; b) D. Mukherjee, M. M. Laloraya, *Biochem. Physiol. Pflanz.* **1974**, *166*, 429–436; c) A. I. Virtanen, A.-M. Berg, *Acta Chem. Scand.* **1955**, *9*, 553–554.

[25] a) T. J. Riedel, L. C. Johnson, J. Knight, R. R. Hantgan, R. P. Holmes, W. T. Lowther, *PLOS ONE* **2011**, *6*, e26021; b) T. J. Riedel, J. Knight, M. S. Murray, D. S. Milliner, R. P. Holmes, W. T. Lowther, *Biochim. Biophys. Acta* **2012**, *1822*, 1544–1552.

[26] a) C. Eckert, W. Xu, W. Xiong, S. Lynch, J. Ungerer, L. Tao, R. Gill, P.-C. Maness, J. Yu, *Biotechnol. Biofuels* **2014**, *7*, 33; b) J. Wilson, S. Gering, J. Pinard, R. Lucas, B. R. Briggs, *Biotechnol. Biofuels* **2018**, *11*, 234.

[27] a) R. Jira, *Angew. Chem. Int. Ed.* **2009**, *48*, 9034–9037; b) J. A. Keith, P. M. Henry, *Angew. Chem. Int. Ed.* **2009**, *48*, 9038–9049; c) V. Kazimírová, M. Rebroš, *Int. J. Mol. Sci.* **2021**, *22*, 4949.

[28] a) S. B. Jaber Sathik Rifayee, M. G. Thomas, C. Christov, *Chem. Sci.* **2025**, *16*, 7667–7684; b) C. Wang, E. R. Sayfutyarova, *J. Phys. Chem. B* **2025**, *129*, 4335–4349.

[29] C. C. Lee, Y. Hu, M. W. Ribbe, *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 9209–9214.

[30] a) S. M. Mayer, W. G. Niehaus, D. R. Dean, *J. Chem. Soc., Dalton Trans.* **2002**, 802–807; b) P. C. Dos Santos, S. M. Mayer, B. M. Barney, L. C. Seefeldt, D. R. Dean, *J. Inorg. Biochem.* **2007**, *101*, 1642–1648.

[31] S. T. Stripp, B. R. Duffus, V. Fourmond, C. Léger, S. Leimkühler, S. Hirota, Y. Hu, A. Jasniewski, H. Ogata, M. W. Ribbe, *Chem. Rev.* **2022**, *122*, 11900–11973.

[32] a) S. S. Chaturvedi, R. Ramanan, J. Hu, R. P. Hausinger, C. Z. Christov, *ACS Catal.* **2021**, *11*, 1578–1592; b) M. G. Thomas, S. B. Jaber Sathik Rifayee, C. Z. Christov, *ACS Catal.* **2024**, *14*, 18550–18569.

[33] a) M. G. Thomas, S. B. Jaber Sathik Rifayee, S. S. Chaturvedi, K. R. Gorantla, W. White, J. Wildey, C. J. Schofield, C. Z. Christov, *Inorg. Chem.* **2024**, *63*, 10737–10755; b) M. E. Neugebauer, E. N. Kissman, J. A. Marchand, J. G. Pelton, N. A. Sambold, D. C. Millar, M. C. Y. Chang, *Nat. Chem. Biol.* **2022**, *18*, 171–179; c) A. Papadopoulou, J. Meierhofer, F. Meyer, T. Hayashi, S. Schneider, E. Sager, R. Buller, *ChemCatChem* **2021**, *13*, 3914–3919; d) S. S. Chaturvedi, S. B. Jaber Sathik Rifayee, S. O. Waheed, J. Wildey, C. Warner, C. J. Schofield, T. G. Karabencheva-Christova, C. Z. Christov, *JACS Au* **2022**, *2*, 2169–2186.


[34] a) C. Zeymer, D. Hilvert, *Annu. Rev. Biochem.* **2018**, *87*, 131–157; b) Y. Wang, P. Xue, M. Cao, T. Yu, S. T. Lane, H. Zhao, *Chem. Rev.* **2021**, *121*, 12384–12444.

[35] a) M. Mukherji, N. J. Kershaw, C. H. MacKinnon, I. J. Clifton, A. S. Wierzbicki, C. J. Schofield, M. D. Lloyd, *Chem. Commun.* **2001**, 972–973; b) C. X. Hou, A. Brasnett, P. Rabe, C. J. Schofield, L. Brewitz, *J. Biol. Chem.*, accepted for publication.


[36] R. W. Adams, C. M. Holroyd, J. A. Aguilar, M. Nilsson, G. A. Morris, *Chem. Commun.* **2013**, *49*, 358–360.

4.12 Supporting Information


4.12.1 Supporting Figures




Supporting Figure 4.S1. Interactions of 2OG within the wt PsEFE active site. LIGPLOT[1] diagram based on a reported wt PsEFE:Mn:2OG crystal structure (PDB ID: 5MOF[2]) showing the interactions of the methylene groups of 2OG with hydrophobic residues, including L206, in the active site of wt PsEFE.




Supporting Figure 4.S2. Purity of the L206 PsEFE variants. (**a**) SDS-PAGE analysis (using a 4-12% Novex™ 330 Tris-glycine gel; Thermo Fisher) of isolated recombinant L206 PsEFE variants indicate >95% purity. The PageRuler™ Plus Prestained Protein Ladder (Thermo Fisher) was used. (**b**) Mass spectrometric analyses of L206 PsEFE variants (3 μM) in buffer (25 mM Tris, pH 7.8) were performed using an iFunnel Agilent 6550 accurate mass quadrupole time-of-flight (Q-TOF) mass spectrometer operated in the positive ionization mode for: (**i**) L206G PsEFE (calculated mass: 39314 Da; observed mass: 39316 Da), (**ii**) L206A PsEFE (calculated mass: 39328 Da; observed mass: 39329 Da), and (**iii**) L206V PsEFE (calculated mass: 39356 Da; observed mass: 39357 Da).




Supporting Figure 4.S3. Analysis of the effects of L206 substitutions on the PsEFE fold using circular dichroism (CD) spectroscopy. CD spectra of wt PsEFE and L206 PsEFE variants showing: (**a**) raw ellipticity and (**b**) mean residue ellipticity (MRE). The results indicate that the PsEFE fold is conserved in wt PsEFE and the three L206 PsEFE variants; note that minor variations in intensities are likely due to differences in protein concentration.[3] Data were normalised to the PsEFE concentration for computing the MRE and smoothened using SELCON3[4] (using DichroWeb[5]). Conditions: 0.1 mg/mL wt PsEFE/L206 PsEFE variant in buffer (20 mM sodium phosphate, pH 7.4).




Supporting Figure 4.S4. L206A and L206V PsEFE catalyse conversion of *L*-arginine in the presence of 2OG. (**a-d**) ^{1}H NMR analyses (~0.5 to ~4.0 ppm and ~5.5 to ~8.5 ppm) for ~90 min incubations of 2OG with: (**a**) L206G PsEFE, (**b**) L206A PsEFE, and (**c**) L206V PsEFE. (**d**) Reported ^{1}H NMR spectrum for a ~90 min incubation of 2OG with wt PsEFE.[6] (**e**) Reported ^{1}H NMR spectrum of 2OG under standard conditions in the absence of PsEFE.[6] Conditions: 400 µM 2OG, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE[6]/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a-c**) are from glycerol; the peak at ~3.7 ppm corresponds to Tris. The peaks at ~2.1 and ~2.3 ppm (dashed circles) correspond to the hydrated ketone form of 2OG.[7] Representative spectra from independent triplicates are shown.




Supporting Figure 4.S5. L206 PsEFE variants do not catalyse efficient oxidation of 2OG in the absence of *L*-arginine. (**a-d**) ^{1}H NMR analyses (~2.3 to ~4.2 ppm and ~5.1 to ~5.9 ppm) of reactions of 2OG ~6 h post incubation with: (**a**) L206G PsEFE, (**b**) L206A PsEFE, (**c**) L206V PsEFE, and (**d**) wt PsEFE. (**e**) ^{1}H NMR spectrum of 2OG in buffer. Conditions: 400 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a-c**) are from glycerol; the peak at ~3.7 ppm in (**a-d**) corresponds to Tris.




Supporting Figure 4.S6. L206A and L206V PsEFE catalyse conversion of *L*-arginine in the presence of 4,4-dimethyl-2OG (2). (**a-d**) ^{1}H NMR analyses (~0.5 to ~4.0 ppm and ~5.5 to ~8.5 ppm) for ~90 min incubations of 4,4-dimethyl-2OG (**2**) with: (**a**) L206G PsEFE, (**b**) L206A PsEFE, and (**c**) L206V PsEFE. (**d**) Reported ^{1}H NMR spectrum for a ~90 min incubation of **2** with wt PsEFE.[6] (**e**) Reported ^{1}H NMR spectrum of **2** under standard conditions in the absence of PsEFE.[6] (**f**) ^{1}H NMR spectrum of an authentic sample of isobutylene (15% solution in THF) under standard assay conditions in the presence of wt PsEFE; note that the alkene protons of isobutylene were not observed, likely because they coincide with the HOD peak. (**g**) ^{1}H NMR spectrum of an authentic sample of isobutylene (15% solution in THF) in buffer. Conditions: 400 µM **2**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE[6]/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a-c**) are from glycerol; the peak at ~3.7 ppm corresponds to Tris. The broad signals at ~0.5 to ~1.5 ppm and ~6.5 to ~7.5 ppm are from the PsEFE sample, which was used in a relatively high concentration. Representative spectra from independent triplicates are shown.




Supporting Figure 4.S7. L206V PsEFE catalyses production of low levels of acetaldehyde from 4-hydroxy-2OG (3). (**a-d**) ^{1}H NMR analyses (~0.5 to ~4.0 ppm and ~5.5 to ~10.0 ppm) for ~90 min incubations of 4-hydroxy-2OG (**3**) with: (**a**) L206G PsEFE, (**b**) L206A PsEFE, and (**c**) L206V PsEFE. (**d**) Reported ^{1}H NMR spectrum for a ~90 min incubation of **3** with wt PsEFE.[6] (**e**) Reported ^{1}H NMR spectrum of **3** under standard conditions in the absence of PsEFE.[6] (**f**) ^{1}H NMR spectrum of an authentic sample of acetaldehyde under standard assay conditions in the presence of wt PsEFE. (**g**) ^{1}H NMR spectrum of an authentic sample of acetaldehyde in buffer. (**h**) ^{1}H NMR spectra (~2.2 to ~2.3 ppm and ~9.6 to ~9.8 ppm) showing that L206V PsEFE catalysed production of relatively low levels of acetaldehyde from **3**. Note that the C2 proton of malic acid (**3a**) was not observed, likely because it coincides with the HOD peak. The stereochemistry of the malic acid (**3a**) and glyceric acid (**3b**) products was not assigned. Conditions: 400 µM **3**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE[6]/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a-d**) are from glycerol; the peak at ~3.7 ppm corresponds to Tris. The broad signals at ~0.5 to ~1.5 ppm and ~6.5 to ~7.5 ppm are from the PsEFE sample, which was used in a relatively high concentration. Note that an unassigned peak was observed at ~2.25 ppm (singlet), the intensity of which apparently did not alter during the reaction. The authentic sample of acetaldehyde in buffer showed peaks corresponding to the hydrated aldehyde form of acetaldehyde (black asterisk, ~1.3 ppm).[8] Representative spectra from independent triplicates are shown.




Supporting Figure 4.S8. No evidence for formation of the 4-hydroxy-2OG (3) retro-aldol product pyruvic acid was accrued under standard assay conditions. (**a**) ^{1}H NMR analysis (~0.4 to ~3.4 ppm) of a ~90 min incubation of 4-hydroxy-2OG (**3**) with L206V PsEFE. (**b**) Reported ^{1}H NMR spectrum of **3** under standard conditions in the absence of PsEFE.[6] (**c**) Reported ^{1}H NMR spectrum of an authentic sample of citramalic acid in buffer.[6] (**d**) Reported ^{1}H NMR spectrum of an authentic sample of parapyruvic acid in buffer.[6] (**e**) Reported ^{1}H NMR spectrum of an authentic sample of pyruvic acid in buffer.[6] Conditions: 400 μM **3**, 500 μM *L*-arginine, 500 μM *L*-ascorbic acid, 50 μM Fe(II), 800 μM TMSP-d_4, and 10 μM L206V PsEFE in buffer (50 mM sodium phosphate, pH 7.4, 10% $_{v/v}$ D_2O). The broad signals in (**a**) at ~0.5 to ~1.5 ppm are from the PsEFE sample, which was used in a relatively high concentration. The authentic sample of pyruvic acid in buffer showed a peak corresponding to the hydrated form of pyruvic acid (dashed star, ~1.48 ppm).[9]




Supporting Figure 4.S9. L206A PsEFE catalyses production of low levels of 1-butylene from 4-ethyl-2OG (4). (**a-d**) ^{1}H NMR analyses (~0.5 to ~4.0 ppm and ~5.5 to ~8.5 ppm) for ~90 min incubations of **4** with: (**a**) L206G PsEFE, (**b**) L206A PsEFE, and (**c**) L206V PsEFE. (**d**) Reported ^{1}H NMR spectrum for a ~90 min incubation of **4** with wt PsEFE.[6] (**e**) Reported ^{1}H NMR spectrum of **4** under standard conditions in the absence of PsEFE.[6] (**f**) ^{1}H NMR spectrum of an authentic sample of 1-butylene under standard assay conditions in the presence of wt PsEFE; note that the C1 alkene protons of 1-butylene were not observed, likely because they coincide with the HOD peak. (**g**) ^{1}H NMR spectrum of an authentic sample of 1-butylene in buffer. **h**) ^{1}H NMR spectra (~0.8 to ~1.0 ppm, ~1.9 to ~2.1 ppm, and ~5.9 to ~6.0 ppm) indicating that L206A PsEFE may catalyse formation of low levels of 1-butylene from **4**; note, however, that peaks corresponding to the methylene and alkene protons of 1-butylene were not observed, possibly due to the low abundance of the alkene product. The stereochemistry of the 2-ethylsuccinic acid (**4a**) and 2-(hydroxymethyl)butanoic acid (**4b**) products was not assigned. Conditions: 400 µM **4**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE[6]/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a-d**) are from glycerol; the peak at ~3.7 ppm corresponds to Tris. Representative spectra from independent triplicates are shown.




Supporting Figure 4.S10. L206A and L206V PsEFE catalyse conversion of *L*-arginine in the presence of 4-propyl-2OG (5). (**a-d**) ^{1}H NMR analyses (~0.5 to ~4.0 ppm and ~5.5 to ~8.5 ppm) for ~90 min incubations of **5** with: (**a**) L206G PsEFE, (**b**) L206A PsEFE, and (**c**) L206V PsEFE. (**d**) Reported ^{1}H NMR spectrum for a ~90 min incubation of **5** with wt PsEFE.[6] (**e**) Reported ^{1}H NMR spectrum of **5** under standard conditions in the absence of PsEFE.[6] Note that the stereochemistry of the 2-propylsuccinic acid (**5a**) and 2-(hydroxymethyl)pentanoic acid (**5b**) products was not assigned. Conditions: 400 μM **5**, 500 μM *L*-arginine, 500 μM *L*-ascorbic acid, 50 μM Fe(II), 800 μM TMSP-d_4, and 10 μM wt PsEFE[6]/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a-d**) are from glycerol; the peak at ~3.7 ppm corresponds to Tris. Representative spectra from independent triplicates are shown.




Supporting Figure 4.S11. L206A and L206V PsEFE catalyse conversion of *L*-arginine in the presence of 3-methyl-2OG (6). (**a-d**) ^{1}H NMR analyses (~0.5 to ~4.0 ppm and ~5.5 to ~8.5 ppm) for ~90 min incubations of **6** with: (**a**) L206G PsEFE, (**b**) L206A PsEFE, and (**c**) L206V PsEFE, indicating that these L206 PsEFE variants do not catalyse formation of propylene from **6**. (**d**) Reported ^{1}H NMR spectrum for a ~90 min incubation of **6** with wt PsEFE.[6] (**e**) Reported ^{1}H NMR spectrum of **6** under standard conditions in the absence of PsEFE.[6] (**f**) ^{1}H NMR spectrum of an authentic sample of propylene under standard assay conditions in the presence of wt PsEFE; note that the C1 alkene protons of propylene were not observed, likely because they coincide with the HOD peak. (**g**) ^{1}H NMR spectrum of an authentic sample of propylene in buffer. Note that the stereochemistry of the 2-methylsuccinic acid (**1a**) and 3-hydroxybutyric acid (**6b**) products was not assigned. Conditions: 400 μM **6**, 500 μM *L*-arginine, 500 μM *L*-ascorbic acid, 50 μM Fe(II), 800 μM TMSP-d_4, and 10 μM wt PsEFE[6]/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm are from glycerol; the peak at ~3.7 ppm corresponds to Tris. The broad signals at ~0.5 to ~1.5 ppm and ~6.5 to ~7.5 ppm are from the PsEFE sample, which was used in a relatively high concentration. Representative spectra from independent triplicates are shown.




Supporting Figure 4.S12. L206A and L206V PsEFE catalyse conversion of *L*-arginine in the presence of 2OA. (**a-d**) ^{1}H NMR analyses (~0.5 to ~4.0 ppm and ~5.5 to ~8.5 ppm) for ~90 min incubations of 2OA with: (**a**) L206G PsEFE, (**b**) L206A PsEFE, and (**c**) L206V PsEFE, indicating that these L206 PsEFE variants do not catalyse formation of propylene from 2OA. (**d**) Reported ^{1}H NMR spectrum for a ~90 min incubation of 2OA with wt PsEFE.[6] (**e**) Reported ^{1}H NMR spectrum of 2OA under standard conditions in the absence of PsEFE.[6] (**f**) ^{1}H NMR spectrum of an authentic sample of propylene under standard assay conditions in the presence of wt PsEFE; note that the C1 alkene protons of propylene were not observed, likely because they coincide with the HOD peak. (**g**) ^{1}H NMR spectrum of an authentic sample of propylene in buffer. Conditions: 400 µM 2OA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE[6]/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm are from glycerol; the peak at ~3.7 ppm corresponds to Tris. The broad signals at ~0.5 to ~1.5 ppm and ~6.5 to ~7.5 ppm are from the PsEFE sample, which was used in a relatively high concentration. Representative spectra of independent triplicates are shown.




Supporting Figure 4.S13. L206 PsEFE variants do not catalyse efficient oxidation of 4-methyl-2OG (1) in the absence of *L*-arginine. (**a-d**) ^{1}H NMR analyses (~2.3 to ~4.2 ppm and ~5.1 to ~5.9 ppm) of reactions of **1** ~8 h post incubation with: (**a**) L206G PsEFE, (**b**) L206A PsEFE, (**c**) L206V PsEFE, and (**d**) wt PsEFE. Note that the results reveal low levels of oxidation of **1** in the absence of *L*-arginine; low levels to 2OG turnover were also observed (Supporting Figure 4.S5). (**e**) ^{1}H NMR spectrum of **1** in buffer. Note that the stereochemistry of the 2-methylsuccinic acid (**1a**) product was not assigned. Conditions: 400 µM **1**, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a-c**) are from glycerol; the peak at ~3.7 ppm in (**a-d**) corresponds to Tris.




Supporting Figure 4.S14. L206 PsEFE variants do not catalyse efficient oxidation of 4-hydroxy-2OG (3) in the absence of *L*-arginine. (**a-d**) ^{1}H NMR analyses (~2.3 to ~4.2 ppm and ~5.1 to ~5.9 ppm) of reactions of **3** ~8 h post incubation with: (**a**) L206G PsEFE, (**b**) L206A PsEFE, (**c**) L206V PsEFE, and (**d**) wt PsEFE. (**e**) ^{1}H NMR spectrum of **3** in buffer. Conditions: 400 µM **3**, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a-c**) are from glycerol; the peak at ~3.7 ppm in (**a-d**) corresponds to Tris.




Supporting Figure 4.S15. L206 PsEFE variants do not catalyse efficient oxidation of 4-ethyl-2OG (4) in the absence of *L*-arginine. (**a-d**) ^{1}H NMR analyses (~2.3 to ~4.2 ppm and ~5.1 to ~5.9 ppm) of reactions of **4** ~8 h post incubation with: (**a**) L206G PsEFE, (**b**) L206A PsEFE, (**c**) L206V PsEFE, and (**d**) wt PsEFE. (**e**) ^{1}H NMR spectrum of **4** in buffer. Conditions: 400 µM **4**, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a-c**) are from glycerol; the peak at ~3.7 ppm in (**a-d**) corresponds to Tris.



Supporting Figure 4.S16. Substitutions at PsEFE residue-206 influence reactivity with 4-methyl-2OG (1). Reactions of: (**a**) L206V PsEFE, (**b**) L206A PsEFE, and (**c**) L206G PsEFE with 4-methyl-2OG (**1**). (**d**) Outcomes of L206 PsEFE variant reactions with **1** compared to that reported for wt PsEFE.[6] Propylene formation was indirectly estimated by subtracting the normalized integrals of 2-methylsuccinic acid and β-hydroxyisobutyric acid (obtained via comparison with TMSP-d_4) from that of **1**; note, substantial amounts of reaction products derived from **1** other than 2-methylsuccinic acid, β-hydroxyisobutyric acid, or propylene were not apparent in the corresponding spectra (Figure 4.4). The time-scales were calibrated to the end of the acquisition of the first ^{1}H NMR experiment following PsEFE addition to the reaction mixture (t = 0 min), by which time low levels of conversion were manifest. Results are means of independent triplicates (n = 3; mean ± standard deviation, SD). Conditions: 400 μM **1**, 500 μM *L*-arginine, 500 μM *L*-ascorbic acid, 50 μM Fe(II), 800 μM TMSP-d$_4$, and 10 μM wt PsEFE[6]/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O).



Supporting Figure 4.S17. Substitutions at PsEFE residue-206 influence reactivity with 4-hydroxy-2OG (3). Reactions of: (**a**) L206V PsEFE, (**b**) L206A PsEFE, and (**c**) L206G PsEFE with 4-hydroxy-2OG (**3**). (**d**) Outcomes of L206 PsEFE variant reactions with **3** compared to that reported for wt PsEFE.[6] The time-scales were calibrated to the end of the acquisition of the first ^{1}H NMR experiment following PsEFE addition to the reaction mixture (t = 0 min), by which time low levels of conversion were manifest. Results are means of independent triplicates (n = 3; mean ± SD). Conditions: 400 µM **3**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d$_4$, and 10 µM wt PsEFE[6]/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O).



Supporting Figure 4.S18. Substitutions at PsEFE residue-206 influence reactivity with 4-ethyl-2OG (4). Reactions of: (**a**) L206V PsEFE, (**b**) L206A PsEFE, and (**c**) L206G PsEFE with 4-ethyl-2OG (**4**). (**d**) Outcomes of L206 PsEFE variant reactions with **4** compared to that reported for wt PsEFE.[6] 1-Butylene formation was indirectly estimated by subtracting the normalized integrals of 2-ethylsuccinic acid and 2-(hydroxymethyl)butanoic acid (obtained via comparison with TMSP-d_4) from that of **4**; note, substantial amounts of reaction products derived from **4** other than 2-ethylsuccinic acid, 2-(hydroxymethyl)butanoic acid, or 1-butylene were not apparent in the corresponding spectra (Supporting Figure 4.S9). The time-scales were calibrated to the end of the acquisition of the first ^{1}H NMR experiment following PsEFE addition to the reaction mixture (t = 0 min), by which time low levels of conversion were manifest. Results are means of independent triplicates (n = 3; mean ± SD). Conditions: 400 μM **4**, 500 μM *L*-arginine, 500 μM *L*-ascorbic acid, 50 μM Fe(II), 800 μM TMSP-d$_4$, and 10 μM wt PsEFE[6]/L206 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.4, 10% v/v D_2O).

4.12.2 Supporting Tables

Supporting Table 4.S1. Primer pairs used for site-directed mutagenesis of wt PsEFE to give the L206 PsEFE variants. Substitutions are highlighted in red. PCR reactions to generate plasmids encoding the L206 PsEFE variants utilized the same forward primer (L206V PsEFE Fwd). A standard hot-start PCR protocol was employed for site-directed mutagenesis of PsEFE: an annealing temperature of 60 °C was used to prevent non-specific primer annealing and the elongation time was set to 5 min to promote amplification of the entire construct (~6.5 kb).

PsEFE variant	Primers pairs (5'-3')		Length (nt)	GC content (%)	Annealing temperature (°C)
L206V	Fwd	TATATTCGTCCGCCTGTTGA	20	45.00	63.0
	Rev	**TAC**ACCACCAACATCATCCTGT	22	45.45	64.0
L206A	Rev	**TGC**ACCACCAACATCATCCTGT	22	50.00	69.8
L206G	Rev	**TCC**ACCACCAACATCATCCTGT	22	50.00	68.7

4.12.3 Supporting Information References

[1] A. C. Wallace, R. A. Laskowski, J. M. Thornton, *Protein Eng.* **1995**, *8*, 127–134.

[2] Z. Zhang, T. J. Smart, H. Choi, F. Hardy, C. T. Lohans, M. I. Abboud, M. S. W. Richardson, R. S. Paton, M. A. McDonough, C. J. Schofield, *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 4667–4672.

[3] N. J. Greenfield, *Nat. Protoc.* **2006**, *1*, 2876–2890.

[4] a) N. Sreerama, R. W. Woody, *Anal. Biochem.* **1993**, *209*, 32–44; b) N. Sreerama, S. Y. U. Venyaminov, R. W. Woody, *Protein Sci.* **1999**, *8*, 370–380.

[5] A. J. Miles, S. G. Ramalli, B. A. Wallace, *Protein Sci.* **2022**, *31*, 37–46.

[6] S. Dhingra, Z. Zhang, C. T. Lohans, L. Brewitz, C. J. Schofield, *J. Biol. Chem.* **2024**, *300*, 107546.

[7] a) A. J. Copper, A. G. Redfield, *J. Biol. Chem.* **1975**, *250*, 527–532; b) T. S. Viswanathan, R. E. Johnson, H. F. Fisher, *Biochemistry* **1982**, *21*, 339–345.

[8] G. Socrates, *J. Org. Chem.* **1969**, *34*, 2958–2961.

[9] A. Lopalco, J. Douglas, N. Denora, V. J. Stella, *J. Pharm. Sci.* **2016**, *105*, 664–672.

4.13 Statement of Authorship

Title of Paper	**Expanding the alkene forming capability of the *Pseudomonas savastanoi* ethylene-forming enzyme.**
Publication Status	□ Published □ Accepted for Publication X Submitted for Publication □ Unpublished and unsubmitted work written in a manuscript style
Publication Details	Dhingra, S.; Sun, Y.; Zhang, Z.; Schofield, C. J.; Brewitz, L. Expanding the alkene forming capability of the *Pseudomonas savastanoi* ethylene-forming enzyme. **Manuscript under review in** *ChemCatChem*.

Student Confirmation

Student Name:	Siddhant Dhingra		
Contribution to the Paper	The experimental work in this manuscript was performed by myself and Mr. Yihong Sun, under my laboratory supervision (I cloned and purified wt and L206V PsEFE; Mr. Yihong Sun purified L206A and L206G PsEFE; both of us performed NMR assays). I conducted all the data analysis and data visualisation described here. The manuscript was prepared by myself and Dr. Lennart Brewitz under the guidance of Prof. Christopher J. Schofield. Dr. Zhihong Zhang provided the wt PsEFE plasmid and the initial training for protein expression and purification. Dr. Lennart Brewitz, Prof. Christopher J. Schofield, and I jointly conceptualised the project.		
Signature 		Date	05-10-2025

Supervisor Confirmation

By signing the Statement of Authorship, you are certifying that the candidate made a substantial contribution to the publication, and that the description described above is accurate.

Supervisor name and title: Prof. Christopher J. Schofield		
Supervisor comments I'm happy to state that the candidate made the major and a very substantial contribution to the publication, and that the description described above is accurate.		
Signature C. J. Schofield	Date	25th September 2025

CHAPTER 5

MANUSCRIPT IV

Oxygenase-catalyzed production of ethylene from the methionine transamination product α-keto-γ-methylthiobutyric acid

Contents

5.1 Overview 252

5.2 Abstract 253

5.3 Introduction 253

5.4 Results 257

5.4.1 Hydrophobic 2-oxoacids with a single carboxylate are PsEFE (co)substrates. 257

5.4.2 The structure of hydrophobic 2-oxoacids affects PsEFE catalysis. 260

5.4.3 The structure of the hydrophobic 2-oxoacid (co)substrate affects the product ratio 263

5.4.4 PsEFE efficiency and reaction outcomes with hydrophobic 2-oxoacids vary with pH 264

5.4.5 2OG is a more efficient PsEFE (co)substrate than hydrophobic 2-oxoacids with a single carboxylate 264

5.4.6 R277 PsEFE variants employ hydrophobic 2-oxoacids as (co)substrates 266

5.4.7 R277 PsEFE variants catalyze ethylene formation from KMBA 268

5.5 Discussion 272

5.6 Methods 280

5.7 Acknowledgements 281

5.8 Author Contributions 282

5.9 References 282

5.10 Supporting Figures 288

5.11 Supporting Tables 372

5.12 Supporting Information References 374

5.13 Statement of Authorship 375

5.1 Overview

Expanding on its (co)substrate scope, PsEFE was found to accept hydrophobic 2-oxoacids lacking the distal carboxylate group that interacts with the Arg277 residue, such as, α-keto-γ-methylthiobutyric acid (KMBA), as alternative (co)substrates. Notably, R277 PsEFE variants did not accept 2OG as a (co)substrate but retained activity with hydrophobic 2-oxoacids, including KMBA. In this chapter, KMBA was validated as a direct ethylene precursor, consistent with pioneering work suggesting its involvement in ethylene biosynthesis in plants and microbes. R277 PsEFE variants catalysed ethylene formation from KMBA alongside its corresponding acid and alcohol products, with ethylene formation apparently linked to methanesulphonic acid (MSA) formation. Isotope-labelling studies probing sulphur oxidation suggested incorporation of a full O_2 molecule into MSA. This study also extends the repertoire of small-molecule ethylene precursors to include the ethyl-variant of KMBA, α-keto-γ-ethylthiobutyric acid (KEBA).

Broadly, these results suggest that hydrophobic 2-oxoacids may influence the activity of PsEFE and, potentially, may influence other 2OG-dependent oxygenases. Such modulation may have context-dependent roles varying with substrate availability. However, further studies are needed to clarify their physiological significance.

At the time of writing this thesis, Manuscript IV is being prepared for submission and is written in the style of a *Nature Communications* paper.

Dhingra, S.; Harrington, L; Sun, Y.; Legge, I.; Zhang, Z.; McCullagh, J. S.; Schofield, C. J.; Brewitz, L. Oxygenase-catalyzed production of ethylene from the methionine transamination product α-keto-γ-methylthiobutyric acid. Manuscript in preparation.

Each author's contribution is described in the Author Contributions (Section 5.8). A Statement of Authorship is appended at the end of the chapter (Section 5.13).

5.2 Abstract

Prior to discovery of ethylene biosynthesis in plants via Fe(II)-dependent 1-aminocyclopropane-1-carboxylate (ACC) oxidase catalysis, pioneering studies demonstrated ethylene production from the methionine transamination product α-keto-γ-methylthiobutyric acid (KMBA); however, isolated enzymes catalyzing KMBA oxidation giving ethylene have not been reported to date. *Pseudomonas savastanoi* employs an Fe(II)-oxygenase (PsEFE) to produce ethylene from the diacid 2-oxoglutarate (2OG). We show that PsEFE reacts with hydrophobic 2-oxoacids with a single carboxylic acid group, including KMBA, to give an acid and, in some cases, an alcohol product. Importantly, PsEFE active site variants employing hydrophobic 2-oxoacids, but not 2OG, as (co)substrates, also catalyze KMBA oxidation giving ethylene and methanesulfonic acid. These results, identifying a new mode of reactivity for Fe(II)/2OG-dependent oxygenases, show that these oxygenases can employ 2-oxoacids with a single carboxylic acid as efficient (co)substrates, and support the proposal that KMBA is a direct precursor for ethylene production.

5.3 Introduction

Ethylene is an important commodity chemical which is produced on an annual scale of >150 million metric tons from non-renewable sources;[1, 2, 3] ethylene is a precursor for manufacture of polymer plastics and base chemicals.[4, 5, 6] Ethylene is also a phytohormone with pleiotropic regulatory functions in plants, including in fruit ripening, senescence, and development.[7] The Fe(II)-dependent 1-aminocyclopropane-1-carboxylate (ACC) oxidase (ACCO) catalyzes the final step in ethylene biosynthesis in seeding plants, *i.e.*, the fragmentation of ACC to give ethylene (Figure 5.1a).[8] By contrast, bacteria employ nitrogenases,[9, 10, 11, 12, 13, 14] nitrogenase-like enzymes,[15, 16, 17] and Fe(II) and 2-oxoglutarate (2OG) dependent ethylene/succinate-forming enzymes (EFEs) to catalyze ethylene production (Figure 5.1).[7, 18] Pioneering work demonstrated that bacteria, ferns, fungi, and other organisms can produce ethylene from the

methionine transamination product α-keto-γ-methylthiobutyric acid (KMBA; 2-oxo-4-methylthiobutyric acid).[19, 20, 21, 22, 23, 24, 25] To our knowledge, however, the proposed enzymatic conversion of KMBA to ethylene has not yet been biochemically validated by characterization of isolated enzymes catalyzing this reaction.[21, 25, 26]

Like ACCO, the EFE of the plant pathogen *Pseudomonas savastanoi* (previously: *P. syringae*) pv. *phaseolicola* PK2 (PsEFE) belongs to the 2-oxoglutarate (2OG) oxygenase structural superfamily.[27, 28] These enzymes share a conserved distorted double-stranded β-helix (DSBH) core fold that supports binding of Fe(II) and 2OG with additional secondary structure elements.[29] 2OG oxygenases catalyze a variety of 2 electron oxidations of their substrates, normally coupled to the oxidative decarboxylation of the diacid 2OG to give succinate and CO_2.[30] Although ACCO does not employ 2OG as a cosubstrate, it belongs to the 2OG oxygenase structural superfamily and has a similar active site as PsEFE, including an HXD...H triad of residues that coordinate the single active site Fe(II) ion (Figure 5.1).[31]




Figure 5.1. Representative enzymatic ethylene forming pathways. (**a**) ACCO catalyzes ethylene formation from ACC.[8, 32] *L*-Ascorbate and bicarbonate promote ACCO catalysis by incompletely understood processes;[7, 8] (**b**) the nitrogenase-like enzyme, methylthio-alkane reductase (Mar), catalyzes anaerobic ethylene formation from (2-methylthio)ethanol (methionol, **1b**);[15, 16, 17] (**c**) PsEFE catalyzes: (i) ethylene formation from 2OG,[27, 28] (ii) *L*-arginine C5 hydroxylation (5-hydroxyarginine fragments to guanidine and P5C) coupled to succinate/diacid formation, and (iii) alcohol formation from most C3/C4-substituted 2OG derivatives;[33, 34] (**d**) view from a PsEFE:Fe:NOG:arginine structure (PDB ID: 5LUN[35]) showing that the catalytically inert 2OG analogue *N*-oxalylglycine (NOG) adopts two conformations with its glycine carboxylate engaging in a salt-bridge interaction with the PsEFE R277 guanidinium group with either one (yellow) or two (pink) of its carboxylate oxygen atoms; (**e**) view from a *Petunia hybrida* ACCO:Ni:ACC structure (PDB ID: 5TCV), with Ni(II) replacing catalytically active Fe(II).

PsEFE employs 2OG as a substrate to give ethylene and CO_2 in a reaction requiring *L*-arginine (Figure 5.1c).[7, 36, 37] PsEFE also employs 2OG as a cosubstrate to catalyze C5 hydroxylation of *L*-arginine to give succinate, CO_2, and 5-hydroxyarginine, the latter of which fragments to give guanidine, water, and *L*-Δ^1-pyrroline-5-carboxylic acid (P5C) (Figure 5.1c).[36, 37] Recent studies have shown that PsEFE can employ C3- and/or C4-substituted 2OG diacid derivatives to sustain arginine oxidation in the absence of 2OG, at least *in vitro*.[33, 34] However, wildtype (wt) PsEFE did not catalyze formation of alkene products from the tested 2OG diacid derivatives, with the exception of 4-fluoro-2OG,[33] but catalyzed formation of an alcohol product from some, but not all, of the tested 2OG diacid derivatives (Figure 5.1c).[33, 34] Thus, the 2OG derivative substitution pattern can dictate the outcome of the PsEFE reaction.[33, 34]

Analysis of PsEFE:*L*-arginine:2OG structures reveals that side chains of hydrophobic amino acids (*e.g.*, L173, F175, L206, V270, A279) shape the 2OG binding site, possibly contributing to stabilizing 2OG binding via hydrophobic interactions with its two methylene groups (Figure 5.1).[38, 39] Notably, the conformation of 2OG at the PsEFE active site can vary, with one or two of the 2OG C5 carboxylate oxygen atoms forming a salt-bridge interaction with the guanidinium group of R277 (Figure 5.1). These observations suggest that hydrophobic 2-oxoacids bearing only a single carboxylic acid moiety, which cannot form a salt-bridge with R277, may bind in the 2OG binding pocket as mediated by stabilizing hydrophobic interactions within the PsEFE active site, possibly in a manner that supports productive catalysis. However, a study on the reactivity of wt PsEFE with hydrophobic 2-oxoacids (*e.g.*, 2-oxovalerate and 2-oxobutyrate) suggest that they do not support catalysis.[40]

Here we show that KMBA and other naturally occurring hydrophobic 2-oxoacids, all lacking the diacid motif of 2OG, are (co)substrates of PsEFE. These hydrophobic 2-oxoacids efficiently sustained *L*-arginine hydroxylation to give the corresponding acid products. PsEFE also catalyzed conversion of some, but not all, the tested hydrophobic 2-oxoacids to the

corresponding alcohol products. Although evidence for alkene formation from the reaction of wt PsEFE with hydrophobic 2-oxoacids was not accrued, R277 PsEFE active site variants catalyzed conversion of KMBA to ethylene. This observation supports the proposal that KMBA can serve as a direct precursor for the enzyme-catalyzed formation of ethylene in plants and bacteria.

5.4 Results

5.4.1 Hydrophobic 2-oxoacids with a single carboxylate are PsEFE (co)substrates.

The (co)substrate scope of isolated recombinant wt PsEFE (UNIPROT ID: P32021; EC:1.13.12.19)[27, 28] was investigated using a set of 20 hydrophobic 2-oxoacids which, unlike the diacid 2OG, only bear a single carboxylate group. Reactions were analyzed using a reported ^{1}H NMR assay,[34] which enables measurement of substrate depletion and product formation by comparison with an internal standard (*i.e.*, 3-(trimethylsilyl)-2,2,3,3-tetradeuteropropionic acid,[41] TMSP-d_4), and characterization of reaction products by comparison with authentic standards. Assays were initially performed overnight to estimate levels of substrate depletion and product formation, followed by time-course studies for active hydrophobic 2-oxoacids; separate assays with 2OG served as controls.

^{1}H NMR analysis (700 MHz) of the overnight reactions reveals that wt PsEFE did not catalyze conversion of any of the 20 tested hydrophobic 2-oxoacids, including KMBA, to alkene products, at least under the tested conditions (Supporting Figures 5.S1-5.S47). Reactions in the absence of PsEFE provided evidence for formation of low levels of ethylene from KMBA (see below). The analyses show that 15 of the 20 hydrophobic 2-oxoacids sustained PsEFE-catalyzed hydroxylation of *L*-arginine to give the corresponding decarboxylated carboxylic acid products and P5C, including naturally occurring 2-oxoacids derived from transamination of proteinogenic α-amino acids such as KMBA (Figure 5.2; Supporting Figures 5.S1-5.S47).

P5C formation was not quantified, because the intensity of its characteristic signal (singlet, δ ~ 7.70 ppm) varied over time.[34] Control experiments showed that the reaction of PsEFE with hydrophobic 2-oxoacids was slow in the absence of *L*-arginine (Supporting Figure 5.S48); however, signal overlap hampered quantification of *L*-arginine depletion.




Figure 5.2. The reactivity of wt PsEFE with hydrophobic 2-oxoacids with a single carboxylic acid. (**a**) wt PsEFE-catalyzed reactions of hydrophobic 2-oxoacids bearing a single carboxylic acid. (**b** - **n**) ^{1}H NMR analyses (700 MHz) of the PsEFE-catalyzed reaction with: (**b**) 2OG, (**c**) KMBA (**1**), (**d**) 4-methyl-2-oxovalerate (**2**), (**e**) 3-methyl-2-oxovalerate (**3**), (**f**) 3-methyl-2-oxobutyrate (**4**), (**g**) 2-oxobutyrate (**5**), (**h**) 2-oxovalerate (**6**), (**i**) 2-oxocaproate (**7**), (**j**) 2-cyclohexyl-2-oxoacetate (**8**), (**k**) 3-cyclohexylpyruvate (**9**), (**l**) 4-cyclohexyl-2-oxobutyrate (**10**), (**m**) 4,4-dimethyl-2-oxovalerate (**11**), and (**n**) 2-oxo-4-phenylbutanoate (**12**). (**o**) Reactions of PsEFE with hydrophobic 2-oxoacids with a single carboxylic acid for which conversion was not quantified. (**p**) Hydrophobic 2-oxoacids which did not react with PsEFE (30 µM) with overnight incubation. Conditions: 400 µM hydrophobic 2-oxoacid, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TSP-d_4, and wt PsEFE (2 µM for 2OG; 5 µM for KMBA; 10 µM for **2-12**; 30 µM for **13-20**) in buffer (50 mM sodium phosphate, pH

7.0, 10%$_{v/v}$ D_2O) at 25˚C. Reaction times are normalized to the end of the first ^{1}H NMR acquisition following PsEFE addition to the reaction mixture (t = 0 min), by which time variable levels of activity were manifest. Results are means of independent triplicates (n = 3; mean ± standard deviation, SD). Representative ^{1}H NMR spectra are shown in Supporting Figures 5.S1-5.S47.

PsEFE also catalyzed conversion of 13 of the 15 hydrophobic 2-oxoacids to give the corresponding alcohol products (Figure 5.2; Supporting Figures 5.S1-5.S47). Although PsEFE-catalyzed alcohol formation was apparently not coupled to *L*-arginine depletion/P5C formation, it did not proceed in the absence of *L*-arginine (Supporting Figure 5.S48). Thus, the reaction of PsEFE with hydrophobic 2-oxoacids is reminiscent of its reaction with some C3- and/or C4-substituted 2OG diacid derivatives, which gave both the corresponding decarboxylated diacid and alcohol products in the presence of *L*-arginine.[33, 34]

5.4.2 The structure of hydrophobic 2-oxoacids affects PsEFE catalysis.

^{1}H NMR analyses reveal that the structure of the linear hydrophobic 2-oxoacids affects the catalytic efficiency of PsEFE (Figure 5.2): Incrementally increasing the chain length/hydrophobicity of 2-oxobutyric acid (**5**) by addition of methylene groups to give 2-oxovaleric acid (**6**) and 2-oxocaproic acid (**7**) correlated with increased efficiency (Supporting Table 5.S2). KMBA, a more hydrophobic C5 sulfur derivative of **7**, reacted even more efficiently than **7**. Note that previous work has reported that **5** and **6** do not sustain PsEFE catalysis, possibly reflecting differences in assays and/or conditions.[40] Specific activities were calculated for wt PsEFE by analysing 2-oxoacid depletion ~10 min post incubation with PsEFE (Supporting Table 5.S2).

The structure of branched hydrophobic 2-oxoacids also affects PsEFE efficiency (Figure 5.2, Supporting Table 5.S2): PsEFE catalyzed depletion of β-branched 3-cyclohexylpyruvic acid

(**9**) more efficiently than of α-branched 2-cyclohexyl-2-oxoacetic acid (**8**) and γ-branched 4-cyclohexyl-2-oxobutyric acid (**10**); **9** was also more efficient in sustaining catalysis than the linear 2-oxoacid **7** (Figures 2 and 3). Substituting the cyclohexyl group of **9** (A value:[42] ~2.15 kcal mol^{-1}) for a sterically bulkier *tert*-butyl group (A value:[42] ~4.2 kcal mol^{-1}) reduced activity, while its substitution for an aryl group reduced activity to an extent that required the use of 3-fold higher PsEFE concentrations to detect turnover (Figure 5.2): PsEFE employed 3-phenylpyruvic acid (PPA, **13**), 3-(4-hydroxyphenyl)pyruvic acid (HPPA, **14**), and 3-(4-methoxyphenyl)pyruvic acid (**15**) as (co)substrates, but not the structurally related 2-(4-nitrophenyl)-2-oxoacetic acid (**16**), 2-(furan-2-yl)-2-oxoacetic acid (**17**), and 3-(4-chlorophenyl)pyruvic acid (**18**), within limits of detection (Figure 5.2). However, turnover of PPA, HPPA, and **15** could not be quantified due to contamination of the starting materials with the acid product or signal overlap with reaction products or different tautomeric forms (Supporting Figures 5.S39-5.S47).




Figure 5.3. The reaction of wt PsEFE with 3-cyclohexyl-2-oxopyruvic acid (9). PsEFE uses 3-cyclohexyl-2-oxopropanoic acid (**9**) to catalyze conversion of *L*-arginine to P5C to give 2-cyclohexylacetic acid (**9a**) and cyclohexylmethanol (**9b**), the latter in a reaction likely uncoupled from *L*-arginine oxidation. (**a**-**f**) ^{1}H NMR analyses (~0.6 to ~3.6 ppm) of: (**a**) reaction of PsEFE and **9** in the presence of an authentic sample of **9b**, (**b**) an authentic sample of **9b**; (**c**) reaction of PsEFE and **9** in the presence of an authentic sample of **9a**, (**d**) an authentic sample of **9a**; (**e**) reaction of PsEFE and **9** ~75 min post addition of PsEFE; (**f**) reaction mixture containing **9**, but without PsEFE. Conditions: 400 µM **9**, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM phosphate, pH 7.0, 10%$_{v/v}$ D_2O) at 25˚C. Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.

5.4.3 The structure of the hydrophobic 2-oxoacid (co)substrate affects the product ratio

The results reveal that the structure of hydrophobic 2-oxoacids not only affects the efficiency of PsEFE catalysis, but also the reaction outcome, *i.e.*, the acid:alcohol ratio. Acid:alcohol ratios were in the 2:1 range for PsEFE reactions with linear (*e.g.*, KMBA) or α-branched (*e.g.*, **3**) hydrophobic 2-oxoacids (Figure 5.4). By contrast, PsEFE reactions with β-branched hydrophobic 2-oxoacids favored acid formation to a greater extent: For example, the estimated acid:alcohol ratio increased from ~2:1 with **6** to ~4.7:1 with both **2** and **11**, and from ~1.8:1 with **8** to ~18:1 with **9** (Figures 5.3 and 5.4). The estimated acid:alcohol ratio decreased from ~18:1 with **9** to ~11:1 with **10**, indicating that acid formation is more favorable for β-branched hydrophobic 2-oxoacids than for γ-branched hydrophobic 2-oxoacids.




Figure 5.4. The structure of the hydrophobic 2-oxoacid (co)substrate governs the outcome of the PsEFE reaction. wt PsEFE reacted with hydrophobic 2-oxoacids to give varying acid:alcohol ratios. Results are means of independent triplicates (n = 3; mean ± SD).

wt PsEFE reacted with PPA and HPPA to exclusively give the acid products, within limits of detection (Supporting Figures 5.S39-5.S44). However, the observations that PsEFE-catalyzed turnover of PPA and HPPA was low and that PsEFE catalyzed alcohol formation from the

PPA/HPPA derivative **15** (Supporting Figures 5.S45-5.S47), suggest that product ratios obtained with relatively inefficient (co)substrates should be interpreted with care. The observation that the structure of the hydrophobic 2-oxoacid (co)substrates affects the PsEFE product ratio is precedented by the reported effects of the structure of 2OG diacid derivatives on the PsEFE reaction outcome.[34]

5.4.4 PsEFE efficiency and reaction outcomes with hydrophobic 2-oxoacids vary with pH

Both the turnover levels and the ethylene:succinate ratio of the PsEFE reaction with 2OG vary with the pH.[34] We therefore investigated the effect of pH on the outcome of the PsEFE reaction with the efficient (co)substrate KMBA (Figure 5.2). The results reveal that PsEFE catalyzed KMBA turnover most effectively at pH ~6.5-7.4, whereas no conversion was observed at pH <5.5 and at pH >9.0 (Supporting Figure 5.S49). Notably, the 3-(methylthio)propionic acid (**1a**):2-(methylthio)ethanol (**1b**) product ratio was ~2:1 at pH >7.0, but ~3:1 at pH 6.0 (Supporting Figure 5.S49), indicating that the reaction conditions can affect the outcome of the PsEFE reaction with hydrophobic 2-oxoacids.

5.4.5 2OG is a more efficient PsEFE (co)substrate than hydrophobic 2-oxoacids with a single carboxylate

Competition assays were performed with both 2OG and KMBA being present in the same reaction mixture to assess effects of the interaction between 2OG and R277 on PsEFE catalysis. Analysis of (co)substrate consumption in reaction mixtures containing equimolar amounts of KMBA and 2OG reveal that 2OG was a more efficient PsEFE (co)substrate than KMBA (Figure 5.5a-c). Notably, the presence of both KMBA and 2OG in the same reaction mixture reduced (co)substrate consumption compared to reactions with a single 2-oxoacid, indicating that KMBA inhibited 2OG turnover and *vice versa* (Figure 5.5a-c). Incrementally increasing the 2OG:KMBA ratio from 1:1 to 1:15 further reduced 2OG depletion, with no 2OG being consumed within ~1 h at 2OG:KMBA ratios >1:10, while levels of KMBA depletion did not

improve. The competition results obtained with 2OG and **9** were similar to those obtained with 2OG and KMBA (Figure 5.5d-f).

Competition assays were also performed with either 2OG or KMBA in the presence of equimolar amounts of the 2OG-competitive, broad spectrum 2OG oxygenase inhibitor, *N*-oxalylglycine (NOG) (Figure 5.1)[35] to inform on the potential of NOG to inhibit PsEFE. PsEFE catalysis was observed with 2OG in the presence of NOG, but not KMBA (Supporting Figure 5.S50), possibly reflecting higher binding affinity of PsEFE for 2OG.




Figure 5.5. wt PsEFE reacts more efficiently with 2OG than with KMBA and 3-cyclohexylpyruvic acid. Time-dependent PsEFE-catalyzed consumption of 2OG, KMBA, and 3-cyclohexylpyruvic acid (**9**) was monitored by ^{1}H NMR (700 MHz): (**a**) 2OG conversion in the presence of incrementally increasing concentrations of KMBA (0-15 equivalents); (**b**) KMBA consumption at varied KMBA concentrations in the presence of 2OG; (**c**) PsEFE assays with 2OG, KMBA or 2OG:KMBA (1:1); (**d**) 2OG conversion in the presence of **9** (0, 1 or 2 equivalents); (**e**) **9** consumption at varied concentrations of **9** in the presence of 2OG; (**f**) PsEFE assays with 2OG, **9** or 2OG:**9** (1:1) mixtures. Reaction times are normalized to the end of the first ^{1}H NMR acquisition following PsEFE addition to the reaction mixture (t = 0 min), by which time variable levels of activity were manifest. Reaction conditions: 2OG, KMBA, or **9** (as indicated), 500 μM *L*-arginine, 500 μM *L*-ascorbate, 50 μM Fe(II), 800 μM TMSP-d_4, and 5 μM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Results are means of independent duplicates (n = 2; mean ± SD).

5.4.6 R277 PsEFE variants employ hydrophobic 2-oxoacids as (co)substrates

The observation that the 2OG diacid structure is not an essential (co)substrate requirement for productive PsEFE catalysis *in vitro* (Figures 5.2-5) suggested that PsEFE variants lacking the R277 guanidinium group may employ hydrophobic 2-oxoacids as (co)substrates. We therefore

produced the R277K, R277F, R277L, and R277A PsEFE variants to test this proposal (Supporting Figures 5.S51-5.S52). The results reveal that R277K, R277F, R277L, and R277A PsEFE do not catalyze conversion of 2OG into ethylene and succinate, even at higher enzyme concentrations (30 μM) and extended incubation times (>10 h) (Supporting Figure 5.S53), although circular dichroism spectroscopy indicated that they share a similar fold as wt PsEFE (Supporting Figure 5.S54). Note that the R277 PsEFE variants also did not react with 2OG derivatives (Supporting Figures 5.S55-5.S57). By contrast, 15 of the 20 tested hydrophobic 2-oxoacids, including KMBA, sustained catalysis by the R277 PsEFE variants in the presence of *L*-arginine (Figure 5.6 and Supporting Figures 5.S58-5.S69).

The nature of the hydrophobic 2-oxoacid substituent and of the R277 substitution affected the relative efficiencies and product outcomes of the reactions of R277K, R277F, R277L, R277A, and wt PsEFE with hydrophobic 2-oxoacids (Supporting Table 5.S2). For example, R277L PsEFE typically catalyzed conversion of hydrophobic 2-oxoacids less efficiently than the other R277 PsEFE variants, possibly reflecting the steric bulk of the leucine side chain (Figure 5.6). Notably, wt PsEFE and R277F PsEFE oxidized **10** substantially more efficiently than R277K, R277A, or R277L PsEFE (Figure 5.6j). wt PsEFE and R277F PsEFE reacted with **10** to give similar acid:alcohol ratios (~10:1), whereas the reaction of R277K PsEFE with **10** apparently produced relatively more alcohol (acid:alcohol ratio ~ 5:1; Supporting Figure 5.S67). These observations are consistent with the calculated specific activities of R277 PsEFE variants for hydrophobic 2-oxoacids (Supporting Table 5.S2).




Figure 5.6. The reactivity of R277 PsEFE variants with hydrophobic 2-oxoacids. ^{1}H NMR analysis (700 MHz) of R277A (orange diamonds), R277F (green triangles), R277K (dark blue inverse triangles), and R277L (pink circles) PsEFE-catalyzed depletion of hydrophobic 2-oxoacids; wt PsEFE was used as a control (light blue boxes): (**a**) KMBA (**1**), (**b**) 4-methyl-2-oxovaleric acid (**2**), (**c**) 3-methyl-2-oxovaleric acid (**3**), (**d**) 3-methyl-2-oxobutyric acid (**4**), (**e**) 2-oxobutyric acid (**5**), (**f**) 2-oxovaleric acid (**6**), (**g**) 2-oxocaproaic acid (**7**), (**h**) 2-cyclohexyl-2-oxoacetic acid (**8**), (**i**) 3-cyclohexylpyruvic acid (**9**), (**j**) 4-cyclohexyl-2-oxobutyric acid (**10**), (**k**) 4,4-dimethyl-2-oxovaleric acid (**11**), and (**l**) 2-oxo-4-phenylbutyric acid (**12**). Conditions: 400 µM 2-oxoacid, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TSP-d_4, 10 µM R277 PsEFE variant in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O) at 25˚C. Reaction times are normalized to the end of the first ^{1}H NMR acquisition following PsEFE addition to the reaction mixture (t = 0 min), by which time variable levels of activity were manifest. Results are means of independent triplicates (for KMBA) or duplicates (± SD). Representative ^{1}H NMR spectra are shown in Supporting Figures 5.S70-5.S72.

5.4.7 R277 PsEFE variants catalyze ethylene formation from KMBA

Notably, the reactions of R277K, R277F, R277L, and R277A PsEFE with KMBA not only gave the acid (**1a**) and alcohol (**1b**) products as previously observed with wt PsEFE (Figure

5.2), but also produced ethylene and methanesulfonic acid (MSA; **1c**) (Figure 5.7; Supporting Figures 5.S70-5.S73). Although ethylene formation could not be quantified, likely due to its evaporation from the reaction mixture, the ^{1}H NMR results imply that ethylene was continuously formed (Supporting Figure 5.S73). Ethylene formation is likely not a result of reactions with co-purified 2OG or 2OG contaminations given that R277 PsEFE variants did not use 2OG as a (co)substrate and no evidence for 2OG or succinate was observed by ^{1}H NMR (Supporting Figure 5.S53).

The capacity of the R277 PsEFE variants to produce ethylene from KMBA apparently varied and was highest for R277K PsEFE under the tested conditions (Supporting Figure 5.S73). This observation may reflect the higher reactivity of R277K PsEFE with KMBA compared to the other R277 PsEFE variants, in accord with the specific activities of R277 PsEFE variants with KMBA (Supporting Table 5.S2). The rates of KMBA depletion decreased in the order: wt PsEFE ~ R277K PsEFE > R277A PsEFE > R277F PsEFE > R277L PsEFE (Figure 5.6a). R277K PsEFE, but not wt PsEFE, also catalyzed ethylene production from the ethionine[43] transamination product α-keto-γ-ethylthiobutyric acid (KEBA) (Supporting Figures 5.S74-5.S77).




Figure 5.7. R277 PsEFE variants catalyze ethylene formation from KMBA. (**a**) R277 PsEFE variants catalyze the reaction of KMBA (**1**) to give ethylene and methanesulfonic acid (MSA, **1c**) together with the 3-(methylthio)propionic acid (**1a**) and 2-(methylthio)ethanol (**1b**); (**b**-**e**) representative ^{1}H NMR spectra (~2.0 to ~5.5 ppm) of the R277K PsEFE reaction with KMBA: (**b**) ~75 min; (**c**) ~50 min; (**d**) ~20 min; and (**e**) ~5 min post addition of R277K PsEFE to the mixture. (**f**) ^{1}H NMR spectrum of the mixture in the absence of R277K PsEFE. Note, the multiplet signals at ~2.5 ppm and ~2.06 ppm correspond to the hydrated ketone form of KMBA (Supporting Figure 5.S2); (**g**-**h**) representative ^{1}H NMR spectra (~5.42 to ~5.46 ppm) showing the R277K PsEFE-catalyzed conversion of KMBA to ethylene in 5 min increments over 80 min in: (**g**) the absence of catalase and (**h**) the presence of catalase (1 µM). Conditions: 400 µM KMBA, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TSP-d_4, and 5 µM R277K PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O) at 25˚C. The peaks at ~3.6 and ~3.7 ppm in (**b**-**e**) are from glycerol; the singlet at ~3.7 ppm in (**b**-**e**) corresponds to Tris. Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.

MSA formation could not be quantified by ^{1}H NMR analysis, because of its low concentration (Figure 5.7). However, its low abundance and the estimated MSA:**1a** and MSA:**1b** ratios suggest that MSA is formed through the same reaction pathway that gives ethylene. There was

no evidence for the formation of methanesulfinic acid, methanesulfenic acid or methanethiol as per comparison with authentic standards. We cannot, however, rule out the possibility that these or related MSA derivatives are transient intermediates. Ion chromatography coupled to mass spectrometry (IC-MS)[44] confirmed MSA formation from KMBA with R277K PsEFE, but not wt PsEFE (Supporting Figure 5.S81). Note that, the low levels of MSA produced during R277K PsEFE catalysis limited the absolute quantification of MSA because of non-linearity of the calibration curve at low concentrations.

We observed very low levels of ethylene formation in reactions of wt PsEFE with KMBA after >1 h and prior to addition of R277 PsEFE variants to reaction mixtures containing KMBA (Figure 5.7g). These observations are consistent with non-enzymatic ethylene production from KMBA, possibly mediated by reactive oxygen species.[45] Thus, catalase was added to the reaction mixture of R277K PsEFE with KMBA to degrade hydrogen peroxide. The results reveal that, in the presence of catalase, ethylene was produced only in the presence of the R277K PsEFE variant, but not in its absence; this supports the proposal that ethylene formation from KMBA is enzyme-catalyzed (Figure 5.7h; Supporting Figures 5.S78-5.S80).

IC-MS-based[44] isotope-labelling studies with $^{18}O_2$ and $H_2{}^{18}O$ were carried out to probe the source of oxygen in MSA produced during R277K PsEFE-catalyzed KMBA turnover. Nearly complete conversion of KMBA was observed with both wt and R277K PsEFE (under $^{16}O_2$ and $^{18}O_2$ atmosphere), as evidenced by the levels of 3-(methylthio)propanoate (**1a**) formed in the reaction mixture; note that the levels of **1a** formed are in accord with the amount of **1a** produced in the ^{1}H NMR assays (Supporting Figures 5.S57 and 5.S81). In reactions under $^{18}O_2$ atmosphere, MS analyses showed nearly exclusive incorporation of a single ^{18}O label into **1a** (Supporting Figure 5.S81), consistent with the prior mechanistic studies on PsEFE which report single ^{18}O incorporation into the PsEFE-catalyzed decarboxylated carboxylic acid product.[33] Relative comparison of MSA isotopologues formed with R277K PsEFE under $^{18}O_2$ atmosphere

showed ~50% abundance of di- and tri-^{18}O-labelled MSA in ~1:1 ratio (Supporting Figure 5.S82). This observation suggests incorporation of one full $^{18}O_2$ molecule into MSA during ethylene formation. Reactions conducted in $H_2{}^{18}O$ did not yield detectable conversion of KMBA and warrant further investigation. However, it is possible, that commercial $H_2{}^{18}O$ may contain trace levels of transition metal ions,[46] which are inhibitors of 2OG oxygenases.[30] Note that, low levels of MSA were also observed in the non-enzymatic reaction of KMBA, consistent with ^{1}H NMR assays (Supporting Figure 5.S82).

5.5 Discussion

The combined results extend the (co)substrate scope of wt PsEFE to hydrophobic 2-oxoacids with a single carboxylic acid group, including the methionine transamination product KMBA (Figures 5.2 and 5.8). The reactions of wt PsEFE with hydrophobic 2-oxoacids gave, at least in most cases, both an acid and an alcohol product, but without alkene production. The relative catalytic efficiencies and product ratios are dependent *inter alia* on the 2-oxoacid substitution pattern, the reaction conditions, and the PsEFE active site geometry (Figures 5.2, 5.4, 5.6; Supporting Figure 5.S49; Supporting Table S2). Thus, the reactivity of wt PsEFE with hydrophobic 2-oxoacids is reminiscent of that with C3- and/or C4-substituted 2OG diacid derivatives which afforded a diacid and an alcohol product in varied ratios, but, at least in most cases, no alkene product.[33, 34]

Although pioneering work on the (co)substrate scope of wt PsEFE suggested that hydrophobic 2-oxoacids may not support catalysis,[40] our results clearly show that at least some of the tested hydrophobic 2-oxoacids are relatively good (co)substrates of wt PsEFE (Figure 5.2), albeit less efficient than 2OG (Figure 5.5). This observation is precedented by work on the (co)substrate scopes of 2OG oxygenases other than PsEFE, the results of which imply that the reactivity of 2OG oxygenases with hydrophobic 2-oxoacids varies on a case-dependent basis.[47, 48, 49, 50, 51,

[52] The combined results further support the possibility that 2OG oxygenases may employ 2-oxoacids other than 2OG in cells.[33, 34, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60] Given the abundance of hydrophobic 2-oxoacids with a single carboxylic acid in cells,[61] the results may be biologically relevant, both in terms of promoting or hindering '2OG' oxygenase catalysis.

It is notable that HPPA (**14**) is a substrate of wt PsEFE because it is also the substrate for 4-hydroxyphenylpyruvate dioxygenase (HPPD) which catalyzes conversion of HPPA to homogentisic acid during tyrosine metabolism; mutations to the *HPD* gene cause the diseases tyrosinemia type III and hawkinsinuria.[62, 63] Like the 2OG oxygenases, HPPD is a non-heme Fe(II)-oxygenase, albeit one from a different structural family which catalyzes a different type of oxidation. Interestingly, HPPD from rat liver also accepts 2-oxoacids other than HPPA as substrates, including KMBA which was observed to give a single enantiomer of the oxidatively decarboxylated sulfoxide ($CH_3S(O)CH_2CH_2CO_2H$),[64, 65] a product we did not observe in PsEFE / KMBA incubations by ^{1}H NMR analysis (Figure 5.7; Supporting Figure 5.S1).




Figure 5.8. Summary of the reactions of wt PsEFE and PsEFE active site variants with 2-oxoacids. (**a**) The reaction of wt PsEFE and L206V PsEFE with 2OG and 2OG diacid derivatives to give diacid, alcohol, and, in some cases, alkene products.[33, 34, 66] (**b**) The reactivity of wt PsEFE and R277 PsEFE variants with hydrophobic 2-oxoacids to give acid, alcohol, and, in some cases, alkene products.

It is possible that PsEFE and, by implication, other 2OG oxygenases have context-dependent functions which may vary in response to (co)substrate availability, nutrition, pH, and other factors. PsEFE/2OG oxygenases can directly alter cellular amino acid concentrations through catalyzing their likely irreversible conversion into (di)acid, alcohol, and alkene products

(Figure 5.8), affecting amino acid flux, the 2-oxoacid pool, and availability of metabolites/signaling molecules. Hydrophobic 2-oxoacids may affect PsEFE/2OG oxygenase catalysis in bacterial cells, because of their relatively high abundance[61] and because they can also inhibit turnover (Figure 5.5). Such modulation of activity may: (i) help in adaptation to environmental stress, (ii) affect the activities of 2OG oxygenases with respect to 'prime' substrate oxidation, and/or (iii) enable cross-talk between metabolic pathways.

Importantly, our results demonstrate that the natural product KMBA and KEBA, the transamination product of naturally occurring ethionine[43] which may likely be a natural product itself,[67, 68] are direct substrates of R277 PsEFE variants for production of ethylene (Figure 5.7). To our knowledge, KEBA has not previously been shown to be a direct ethylene precursor, whereas the corresponding amino acid ethionine has been reported to give ethylene under light in the absence of enzymes.[69, 70] R277 PsEFE variants, but not wt PsEFE, catalyzed conversion of KMBA and KEBA into ethylene (Figures 5.2 and 5.7). By contrast, wt PsEFE, but not R277 PsEFE variants, catalyzed conversion of 2OG into ethylene (Figure 5.2; Supporting Figure 5.S53), indicating that the altered active site geometry of the R277 PsEFE variants governs the reaction outcome. This proposal is precedented by work on nitrogenases showing that active site substitutions enable ethylene production from CO,[71] and by work on L206V PsEFE showing that the L206V variant catalyzes propylene formation from 4-methyl-2OG.[66]

Levels of PsEFE-catalyzed ethylene formation from KMBA can likely be increased by further modifications either at the active site or beyond. It is possible that currently uncharacterized 2OG oxygenase(s) in their wildtype form may catalyze ethylene production from KMBA. Indeed, KMBA is a reported precursor of ethylene in some bacteria[20, 21, 22, 72] and other organisms[21, 23, 24, 25]. Enzymes mechanistically distinct from those of the 2OG oxygenase structural superfamily may also have capacity to catalyze conversion of KMBA into ethylene.[26]

The PsEFE mechanism is the subject of ongoing investigations;[33, 73, 74, 75, 76, 77, 78] however, the importance of the hydrophobic 2OG pocket of PsEFE for enabling ethylene formation from 2OG is apparent.[38, 73, 76] The observation that an increase in the hydrophobicity of linear hydrophobic 2-oxoacids correlated with higher PsEFE (co)substrate activities (*i.e.*, KMBA > **7** > **6** > **5**; Figure 5.2) implies that the hydrophobic PsEFE active site also helps in stabilizing binding of hydrophobic 2-oxoacids with a single carboxylate group to the 2OG pocket. Turnover of hydrophobic 2-oxoacids to acid and alcohol products may proceed at the PsEFE active site via a radical mechanism in a similar manner to that proposed for 2OG and 2OG diacid derivatives to give the same product types.[33, 73]

Different mechanistic scenarios can rationalize the R277 PsEFE-catalyzed ethylene and sulfonic acid formation from KMBA and KEBA. Our results collectively support a scenario in which KMBA/KEBA oxidation occurs at the active site of R277 PsEFE variants in an enzyme-catalyzed manner. The observations that R277 PsEFE variants did not catalyze ethylene formation from 2-oxobutyric acid (**5**) and from the KMBA C5 methylene derivative **7** suggest an essential role of the KMBA γ-S atom for ethylene formation (Figure 5.6). It is unlikely that ethylene formation from KMBA involves *S*-methyl bond fission to give methyl radicals,[79] because MS and ^{1}H NMR analyses did not provide evidence for PsEFE methylation or methanol formation (Supporting Figures 5.S70-5.S72), instead for MSA formation (Figure 5.7). However, the precise sequence of the PsEFE-catalyzed conversion of KMBA to ethylene is unclear, in particular whether sulfur oxidation proceeds or follows KMBA fission to give ethylene (Figure 5.7). Attempts to synthesize KMBA sulfoxide/sulfone as mechanistic probes have been unsuccessful, possibly reflecting their intrinsic instability.[22, 23]

Assuming that ethylene and MSA are produced via the same pathway in a 1:1 ratio as implied by ^{1}H NMR analyses (Figure 5.7), two molecules of O_2 are required for the reaction. Thus, it is possible that the reaction of KMBA and O_2 initially gives ethylene and a PsEFE:Fe(II):OSMe

complex which undergoes reaction with a second O_2 molecule to give MSA, potentially in a manner related to the mechanism of Fe(II)-dependent cysteine dioxygenase.[80, 81] This hypothesis is further supported by MS-based isotope labelling studies on R277K PsEFE-catalyzed KMBA turnover where one full $^{18}O_2$ molecule is apparently incorporated into MSA (Supporting Figure 5.S82). However, we cannot rule out other possibilities, including that methanethiol is transiently formed and undergoes efficient enzymatic or non-enzymatic oxidation to MSA. The PsEFE-catalyzed oxidation of KMBA and KEBA to ethylene is the subject of ongoing investigations.

The four R277 PsEFE active site variants characterized in this work employed hydrophobic 2-oxoacids as (co)substrates, but did not catalyze 2OG oxidation (Figure 5.6). To our knowledge, R277 (equivalent) PsEFE variants have not yet been identified in microbes. However, disease-associated variations in human phytanoyl-CoA 2-hydroxylase (PAHX) and aspartate/asparagine-β-hydroxylase (AspH) occur at R277 PsEFE-equivalent arginine residues forming salt-bridge interactions with the 2OG C5 carboxylate: *i.e.*, R275 in PAHX[82] and R735 in AspH. The reactivity of KMBA with R277 PsEFE variants differs from that with wt PsEFE, an observation precedented by the (co)substrate scopes of R275W PAHX and R735W AspH which do not employ 2OG as a (co)substrate *in vitro*, but hydrophobic 2-oxoacids lacking a second carboxylate.[83] The combined results show that the salt-bridge interaction between 2OG and R277 of wt PsEFE is a requirement for, at least efficient, productive PsEFE catalysis with 2OG and 2OG diacid derivatives (Supporting Figures 5.S53 and 5.S55-5.S57), but not for productive reaction with hydrophobic 2-oxoacids.

The observations that R277 PsEFE (Figure 5.6), R275W PAHX,[83] and R735W AspH variants (unpublished work, CJS group) are catalytically active in the presence of hydrophobic 2-oxoacids suggest that 2OG oxygenase active site geometries required for productive catalysis may be more diverse than has been perceived. This proposal is supported by the identification

of 2OG oxygenases with Fe(II) binding motives diverging from the canonical HXD/E…H triad present in 2OG protein hydroxylases,[31, 84, 85, 86] including catalytically competent 2OG protein hydroxylase variants with two[87, 88, 89] or one protein-bound Fe(II) ligand.[89, 90] Recent evidence implies that the DSBH-fold supporting the active site geometry of 2OG oxygenases is also not essential for catalysis.[91]

The observation that KMBA serves as a direct precursor for enzyme-catalyzed ethylene formation is of interest from evolutionary perspectives. KMBA is a Yang cycle intermediate and a precursor in the biosynthesis of ACC, suggesting the possibility that ethylene signaling mechanisms may have evolved from the same methionine-related origin.[92, 93] Interestingly, the methionine/KMBA-derived metabolite (2-methylthio)ethanol (methionol; **1b**)[94] is the substrate of a bacterial nitrogenase-like enzyme catalyzing ethylene formation under anaerobic and sulfur-limiting conditions.[15, 16, 17] Thus, ethylene formation mechanisms in aerobic and anaerobic bacteria may have evolved from a common ancestor given that methionol is also produced by the reaction of wt PsEFE and PsEFE variants with KMBA (Figure 5.2).

The observation that enzymes can catalyze ethylene production from KMBA is of interest from sustainable ethylene production perspectives.[3, 95] Although biocatalysis is an attractive method to enable ethylene production from renewable resources,[2, 12, 96] efforts to engineer wt PsEFE into organisms to introduce capacity for ethylene production are compromised by the relatively broad (co)substrate scope of PsEFE and its bifurcating reaction pathway affording product mixtures (Figure 5.8).[1, 2, 97] Our observations that PsEFE (co)substrates can be tuned to exclusively give the acid product and that the (co)substrate scope of wt PsEFE differs from that of PsEFE active site variants (Figures 5.2 and 5.6) indicate the possibility of tailoring PsEFE for selective ethylene production from 2OG, KMBA and/or related molecules. Note that PsEFE is also of interest for biocatalysis applications beyond ethylene production, because it catalyzes formation of industrially-relevant acid and alcohol products.[98]

KMBA, 3-cyclohexylpyruvic acid (**9**), and, by implication, other hydrophobic 2-oxoacids can impede PsEFE-catalyzed 2OG turnover in a concentration-dependent manner (Figure 5.5). This observation is of interest from an inhibitor development perspective given that human 2OG oxygenases are validated drug targets.[99, 100] Hydrophobic 2-oxoacids may serve as scaffolds for 2OG oxygenase inhibitors as they do not interact with polar active site residues coordinating the 2OG C5 carboxylate, thereby possibly enhancing inhibition selectivity. This proposal is supported by observations that derivatives of 2OG-mimics such as pyridine-2,4-dicarboxylic acid[101, 102, 103] and *N*-oxalylglycine[104], and C3- and/or C4-substituted 2OG diacid derivatives[54, 55, 56, 105, 106] are (partially) selective 2OG oxygenase inhibitors.

In summary, the combined results reveal the capacity of PsEFE to employ hydrophobic 2-oxoacids with a single carboxylate group as (co)substrates for formation of acid and alcohol products (Figures 5.2 and 5.6). Importantly, our work shows that enzymes of the 2OG oxygenase superfamily do not only employ 2OG and ACC as direct precursors for ethylene formation, but also KMBA and KEBA (Figure 5.7), suggesting that they also have capacity to catalyze conversion of other suitably functionalized naturally occurring molecules to ethylene and/or other gaseous alkenes with potential functions in signaling.

5.6 Methods

Protein production. Plasmid DNA encoding for R277K, R277F, R277L, and R277A PsEFE was obtained from our reported wt PsEFE construct[35] by standard site-directed mutagenesis (Supporting Table 5.S1). The R277 PsEFE variants were produced and purified as reported for wt PsEFE;[34, 35] they were >90% pure by SDS-PAGE analysis (Supporting Figure 5.S51), had the anticipated masses (Supporting Figure 5.S52), and shared a similar fold with wt PsEFE (Supporting Figure 5.S54). They were stored at −70 ºC, at ~20-25 mg/mL concentrations.

2-Oxoacids. Hydrophobic 2-oxoacids with a single carboxylate, 2-oxoglutarate (2OG), and 2-oxoadipate (2OA) were from commercial sources (mostly Sigma-Aldrich) and were used as received. Racemic 3-methyl-2OG and 4-methyl-2OG were synthesised as reported.[54]

^{1}H NMR assays. ^{1}H NMR assays were performed as reported,[34] typically using: hydrophobic 2-oxoacids (400 μM), *L*-arginine (500 μM), *L*-ascorbic acid (500 μM), Fe(II) (50 μM), TMSP-d_4 (800 μM), and R277 PsEFE variant (10 μM or, in some cases, 30 μM) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O), if not specified otherwise in the Figure 5.legends. Freshly thawed aliquots of R277 PsEFE variants were used for all experiments. A Bruker AVIII 700 MHz NMR spectrometer with a 5-mm inverse triple-resonance-inverse cryoprobe was used for analysis. The water peak in ^{1}H NMR experiments was suppressed using the perfect echo-modified WATERGATE solvent suppression method.[107]

IC-MS analysis. Anion-exchange chromatography-MS analysis was performed as reported,[44] using KMBA (400 μM), *L*-arginine (500 μM), *L*-ascorbic acid (500 μM), Fe(II) (50 μM), and wt/R277K PsEFE (5 μM) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Samples were incubated for ~ 6 h after addition of PsEFE and then filtered using 10 kDa molecular weight cut-off filters. A 5 μL partial loop injection was used for all analyses. Chromatographic separations were performed using a Dionex ICS-5000+ high-pressure ion chromatography

system (Dionex, Sunnyvale, CA, USA) coupled to a Q-Exactive HF hybrid quadrupole-Orbitrap mass spectrometer (Thermo Scientific, San Jose, CA, USA). A Thermo IonPac AS11-HC (2 × 250 mm, 4 μm) column was used at 30 °C with an aqueous hydroxide ion gradient at a flow rate of 0.25 mL/min with the following steps: 0 min, 5 mM; 1 min, 5 mM; 15 min, 60 mM; 25 min, 100 mM; 30 min, 100 mM; 30.1 min, 5 mM; 37 min, 5 mM. A continuously regenerated trap column (CR-ATC 500, 2 mm) was used to remove ionic contaminants from the eluent and ion suppression was achieved using a Dionex AERS 500e 2 mm (Dionex, Sunnyvale, CA, USA) in external water mode with a flow rate of 0.5 mL/min. The mass spectrometer was equipped with a HESI II probe in negative ion mode with source parameters set as follows: sheath gas flow rate, 60; auxiliary gas flow rate, 20; sweep gas flow rate, 0; spray voltage, -3.2 kV; capillary temperature, 320 °C; S-lens RF level, 70 and heater temperature 350 °C. MS scan parameters were set as follows: microscans, 2; resolution, 70000; AGC target, 1 × 10 6 ions; maximum IT, 120 ms and scan range, 60 900 m/z. MS/MS scan parameters were set as follows: micro- scans, 1; resolution, 17500; AGC target, 1 × 10 5 ions; maximum IT, 250 ms; loop count, 10; MSX count, 1; isolation window, 2.0 m/z; collision energy, 35; minimum AGC target, 5 × 10 3 ions; apex trigger 1–15 s; charge exclusion, 3–8, >8 and dynamic exclusion, 20.0 s.

5.7 Acknowledgements

This research was funded in part by the Wellcome Trust (106244/Z/14/Z). For the purpose of open access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission. We thank Cancer Research UK (C8717/A18245) and the Biotechnology and Biological Sciences Research Council (BB/J003018/1 and BB/R000344/1) for funding and Prof. Dr. C. Redfield for helpful discussions.

5.8 Author Contributions

S.D., L.H., and Y.S. performed assays; S.D., L.H., and Z.Z. and produced enzymes; L.B. conceived the project together with S.D. and C.J.S.; S.D., I. L., and J. S. M. performed the MS analyses; L. B. and C.J.S. supervised the research; all authors analyzed data; L.B., S.D. and C.J.S. wrote the manuscript with help from all authors.

5.9 References

1. Wilson J, Gering S, Pinard J, Lucas R, Briggs BR. Bio-production of gaseous alkenes: ethylene, isoprene, isobutene. *Biotechnol Biofuels* **11**, 234 (2018).
2. Eckert C, *et al.* Ethylene-forming enzyme and bioethylene production. *Biotechnol Biofuels* **7**, 33 (2014).
3. Gao Y, *et al.* Recent Advances in Intensified Ethylene Production—A Review. *ACS Catalysis* **9**, 8592–8621 (2019).
4. Egbuhuzor MO, Ogbobe P, Umunnakwe C. Polyethylenes: A Vital Recyclable Polymer. In: *Waste Material Recycling in the Circular Economy - Challenges and Developments* (ed Achilias DS). IntechOpen (2022).
5. Liang J, *et al.* Security analysis of ethylene glycol production pathways. *Chem Eng Res Des* **214**, 65–77 (2025).
6. Wu L, Moteki T, Gokhale Amit A, Flaherty David W, Toste FD. Production of Fuels and Chemicals from Biomass: Condensation Reactions and Beyond. *Chem* **1**, 32–58 (2016).
7. Hausinger RP, Rifayee SBJS, Thomas MG, Chatterjee S, Hu J, Christov CZ. Biological formation of ethylene. *RSC Chem Biol* **4**, 635–646 (2023).
8. Houben M, Van de Poel B. 1-Aminocyclopropane-1-carboxylic acid oxidase (ACO): the enzyme that makes the plant hormone ethylene. *Front Plant Sci* **10**, 695 (2019).
9. Lee CC, Hu Y, Ribbe MW. Vanadium nitrogenase reduces CO. *Science* **329**, 642–642 (2010).
10. Hu Y, Lee CC, Ribbe MW. Extending the Carbon Chain: Hydrocarbon Formation Catalyzed by Vanadium/Molybdenum Nitrogenases. *Science* **333**, 753–755 (2011).
11. Lee CC, Hu Y, Ribbe MW. Tracing the hydrogen source of hydrocarbons formed by Vanadium Nitrogenase. *Angew Chem Int Ed* **50**, 5545–5547 (2011).
12. Hu Y, Lee CC, Grosch M, Solomon JB, Weigand W, Ribbe MW. Enzymatic Fischer–Tropsch-type reactions. *Chem Rev* **123**, 5755–5797 (2023).
13. Seefeldt LC, *et al.* Reduction of substrates by nitrogenases. *Chem Rev* **120**, 5082–5106 (2020).
14. Barney BM, Igarashi RY, Dos Santos PC, Dean DR, Seefeldt LC. Substrate interaction at an iron-sulfur face of the FeMo-cofactor during nitrogenase catalysis. *J Biol Chem* **279**, 53621–53624 (2004).
15. North JA, *et al.* A nitrogenase-like enzyme system catalyzes methionine, ethylene, and methane biogenesis. *Science* **369**, 1094–1098 (2020).
16. Lago-Maciel A, *et al.* Methylthio-alkane reductases use nitrogenase metalloclusters for carbon-sulfur bond cleavage. *bioRxiv*, 2024.2010.2019.619033 (2024).

17. North JA, Miller AR, Wildenthal JA, Young SJ, Tabita FR. Microbial pathway for anaerobic 5′-methylthioadenosine metabolism coupled to ethylene formation. *Proc Natl Acad Sci USA* **114**, E10455–E10464 (2017).
18. Cui Y, *et al.* Discovery of Five New Ethylene-Forming Enzymes for Clean Production of Ethylene in E. coli (2022).
19. Osborne DJ, Walters J, Milborrow BV, Norville A, Stange LMC. Evidence for a non-ACC ethylene biosynthesis pathway in lower plants. *Phytochemistry* **42**, 51–60 (1996).
20. Shipston BN, Bunch AW. The physiology of L-methionine catabolism to the secondary metabolite ethylene by *Escherichia coli*. *J Gen Microbiol* **135**, 1489–1497 (1989).
21. Mansouri S, Bunch AW. Bacterial Ethylene Synthesis from 2-Oxo-4-thiobutyric Acid and from Methionine. *J Gen Microbiol* **135**, 2819–2827 (1989).
22. Billington DC, Golding BT, Primrose SB. Biosynthesis of ethylene from methionine. Isolation of the putative intermediate 4-methylthio-2-oxobutanoate from culture fluids of bacteria and fungi. *Biochem J* **182**, 827–836 (1979).
23. Chagué V, Elad Y, Barakat R, Tudzynski P, Sharon A. Ethylene biosynthesis in Botrytis cinerea. *FEMS Microbiol Ecol* **40**, 143–149 (2002).
24. Gu S, Xie L, Guan Q, Sheng X, Fang Y, Wang X. Effect of ethylene production by four pathogenic fungi on the postharvest diseases of green pepper (Capsicum annuum L.). *Int J Food Microbiol* **418**, 110729 (2024).
25. Fukuda H, Takahashi M, Fujii T, Tazaki M, Ogawa T. An NADH: Fe(III) EDTA oxidoreductase from Cryptococcus albidus: an enzyme involved in ethylene production in vivo? *FEMS Microbiol Lett* **60**, 107–111 (1989).
26. Ogawa T, Takahashi M, Fujii T, Tazaki M, Fukuda H. The role of NADH:Fe(III)EDTA oxidoreductase in ethylene formation from 2-keto-4-methylthiobutyrate. *J Ferment Bioeng* **69**, 287–291 (1990).
27. Nagahama K, *et al.* Purification and properties of an ethylene-forming enzyme from Pseudomonas syringae pv. phaseolicola PK2. *J Gen Microbiol* **137**, 2281–2286 (1991).
28. Fukuda H, *et al.* Molecular cloning in Escherichia coli, expression, and nucleotide sequence of the gene for the ethylene-forming enzyme of Pseudomonas syringae pv. phaseolicola PK2. *Biochem Biophys Res Commun* **188**, 826–832 (1992).
29. Aik W, McDonough MA, Thalhammer A, Chowdhury R, Schofield CJ. Role of the jelly-roll fold in substrate binding by 2-oxoglutarate oxygenases. *Curr Opin Struct Biol* **22**, 691–700 (2012).
30. *2-Oxoglutarate-dependent oxygenases*. The Royal Society of Chemistry (2015).
31. Kal S, Que L. Dioxygen activation by nonheme iron enzymes with the 2-His-1-carboxylate facial triad that generate high-valent oxoiron oxidants. *J Biol Inorg Chem* **22**, 339–365 (2017).
32. Bleecker AB, Kende H. Ethylene: a gaseous signal molecule in plants. *Annu Rev Cell Dev Biol* **16**, 1–18 (2000).
33. Copeland RA, Zhou S, Schaperdoth I, Shoda TKC, Bollinger JM, Krebs C. Hybrid radical-polar pathway for excision of ethylene from 2-oxoglutarate by an iron oxygenase. *Science* **373**, 1489–1493 (2021).
34. Dhingra S, Zhang Z, Lohans CT, Brewitz L, Schofield CJ. Substitution of 2-oxoglutarate alters reaction outcomes of the Pseudomonas savastanoi ethylene-forming enzyme. *J Biol Chem* **300**, 107546 (2024).
35. Zhang Z, *et al.* Structural and stereoelectronic insights into oxygenase-catalyzed formation of ethylene from 2-oxoglutarate. *Proc Natl Acad Sci USA* **114**, 4667–4672 (2017).
36. Fukuda H, *et al.* Two reactions are simultaneously catalyzed by a single enzyme: The arginine-dependent simultaneous formation of two products, ethylene and succinate, from 2-

oxoglutarate by an enzyme from Pseudomonas syringae. *Biochem Biophys Res Commun* **188**, 483–489 (1992).
37. Nagahama K, Ogawa T, Fujii T, Tazaki M, Goto M, Fukuda H. L-Arginine is essential for the formation in vitro of ethylene by an extract of Pseudomonas syringae. *J Gen Microbiol* **137**, 1641–1646 (1991).
38. Martinez S, Fellner M, Herr CQ, Ritchie A, Hu J, Hausinger RP. Structures and mechanisms of the non-heme Fe(II)- and 2-oxoglutarate-dependent ethylene-forming enzyme: substrate binding creates a twist. *J Am Chem Soc* **139**, 11980–11988 (2017).
39. Chatterjee S, *et al.* Structural, spectroscopic, and computational insights from canavanine-bound and two catalytically compromised variants of the ethylene-forming enzyme. *Biochemistry* **63**, 1038–1050 (2024).
40. Martinez S, Hausinger RP. Biochemical and spectroscopic characterization of the non-heme Fe(II)- and 2-oxoglutarate-dependent ethylene-forming enzyme from pseudomonas syringae pv. phaseolicola PK2. *Biochemistry* **55**, 5989–5999 (2016).
41. Pohl L, Eckle M. Sodium 3-trimethylsilyltetradeuteriopropionate, a new water-soluble standard for ^{1}H-NMR. *Angew Chem Int Ed* **8**, 381 (1969).
42. *Stereochemistry of Organic Compounds*. Wiley (1993).
43. Fischer NS, Steinhaus M. Identification of an Important Odorant Precursor in Durian: First Evidence of Ethionine in Plants. *J Agric Food Chem* **68**, 10397–10402 (2020).
44. Williams R, *et al.* Metabolomics using Anion-Exchange Chromatography Mass Spectrometry for the Analysis of Cells, Tissues and Biofluids. *Nat Protoc*, (2025).
45. Primrose SB. Evaluation of the Role of Methional, 2-Keto-4-methylthiobutyric Acid and Peroxidase in Ethylene Formation by Escherichia coli. *Microbiol* **98**, 519–528 (1977).
46. Gillies JM, Najim N, Zweit J. Analysis of Metal Radioisotope Impurities Generated in [^{18}O]H_2O During the Cyclotron Production of Fluorine-18. *Appl Radiat Isot* **64**, 431–434 (2006).
47. Ng SF, Hanauske-Abel HM, Englard S. Cosubstrate binding site of Pseudomonas sp. AK1 gamma-butyrobetaine hydroxylase. Interactions with structural analogs of alpha-ketoglutarate. *J Biol Chem* **266**, 1526–1533 (1991).
48. White AK, Metcalf WW. Isolation and biochemical characterization of hypophosphite/ 2-oxoglutarate dioxygenase: a novel phosphorus-oxidizing enzyme from Pseudomonas Stutzeri WM88. *J Biol Chem* **277**, 38262–38271 (2002).
49. Kahnert A, Kertesz MA. Characterization of a sulfur-regulated oxygenative alkylsulfatase from Pseudomonas putida S-313. *J Biol Chem* **275**, 31661–31667 (2000).
50. Fukumori F, Hausinger RP. Purification and characterization of 2,4-dichlorophenoxyacetate/alpha-ketoglutarate dioxygenase. *J Biol Chem* **268**, 24311–24317 (1993).
51. Eichhorn E, van der Ploeg JR, Kertesz MA, Leisinger T. Characterization of α-ketoglutarate-dependent taurine dioxygenase from Escherichia coli. *J Biol Chem* **272**, 23031–23036 (1997).
52. Müller TA, Fleischmann T, Meer JRvd, Kohler H-PE. Purification and characterization of two enantioselective α-ketoglutarate-dependent dioxygenases, RdpA and SdpA, from *Sphingomonas herbicidovorans* MH. *Appl Environ Microbiol* **72**, 4853–4861 (2006).
53. Breski M, Dey D, Obringer S, Sudhamalla B, Islam K. Engineering biological C–H functionalization leads to allele-specific regulation of histone demethylases. *J Am Chem Soc* **138**, 13505–13508 (2016).
54. Brewitz L, Nakashima Y, Schofield CJ. Synthesis of 2-oxoglutarate derivatives and their evaluation as cosubstrates and inhibitors of human aspartate/asparagine-β-hydroxylase. *Chem Sci* **12**, 1327–1342 (2021).

55. Nakashima Y, Brewitz L, Tumber A, Salah E, Schofield CJ. 2-Oxoglutarate derivatives can selectively enhance or inhibit the activity of human oxygenases. *Nat Commun* **12**, 6478 (2021).
56. Tumber A, Salah E, Brewitz L, Corner T, Schofield CJ. Kinetic and inhibition studies on human Jumonji-C (JmjC) domain-containing protein 5. *RSC Chem Biol* **4**, 399–413 (2023).
57. Tiwari AD, Guan Y, Grabowski DR, Maciejewski JP, Jha BK, Phillips JG. SAR insights into TET2 catalytic domain inhibition: Synthesis of 2-hydroxy-4-methylene-pentanedicarboxylates. *Bioorg Med Chem* **39**, 116141 (2021).
58. Fiorini G, Marshall SA, Figg WD, Myers WK, Brewitz L, Schofield CJ. Human prolyl hydroxylase domain 2 reacts with O2 and 2-oxoglutarate to enable formation of inactive Fe(III).2OG.hypoxia-inducible-factor α complexes. *Sci Rep* **14**, 26162 (2024).
59. Majamaa K, Hanauske-Abel HM, Günzler V, Kivirikko KI. The 2-oxoglutarate binding site of prolyl 4-hydroxylase. *Eur J Biochem* **138**, 239–245 (1984).
60. Cunliffe CJ, Franklin TJ, Hales NJ, Hill GB. Novel inhibitors of prolyl 4-hydroxylase. 3. Inhibition by the substrate analog N-oxaloglycine and its derivatives. *J Med Chem* **35**, 2652–2658 (1992).
61. Bröer S, Gauthier-Coles G. Amino acid homeostasis in mammalian cells with a focus on amino acid fransport. *J Nutr* **152**, 16–28 (2022).
62. Moran GR. 4-Hydroxyphenylpyruvate dioxygenase. *Arch Biochem Biophys* **433**, 117–128 (2005).
63. Moran GR. 4-Hydroxyphenylpyruvate dioxygenase and hydroxymandelate synthase: Exemplars of the α-keto acid dependent oxygenases. *Arch Biochem Biophys* **544**, 58–68 (2014).
64. Adlington RM, Baldwin JE, Crouch NP, Lee M-H, MacKinnon CH. Identification and stereochemistry of the product of 4-HPPD catalyzed oxidation of the ketoacid of methionine. *Bioorg Med Chem Lett* **6**, 2003–2006 (1996).
65. Crouch NP, Adlington RM, Baldwin JE, Lee MH, MacKinnon CH. A mechanistic rationalisation for the substrate specificity of recombinant mammalian 4-hydroxyphenylpyruvate dioxygenase (4-HPPD). *Tetrahedron* **53**, 6993–7010 (1997).
66. Burke EJ, Copeland RA, Dixit Y, Krebs C, Bollinger JM, Jr. Steric perturbation of the Grob-like final step of ethylene-forming enzyme enables 3-hydroxypropionate and propylene production. *J Am Chem Soc* **146**, 1977–1983 (2024).
67. Ikeda T, Konishi Y, Ichihara A. Transaminase of branched chain amino acids: XI. leucine (methionine) transaminase of rat liver mitochondria. *Biochim Biophys Acta* **445**, 622–631 (1976).
68. Steele RD, Benevenga NJ. Identification of a transaminative pathway for ethionine catabolism. *Cancer Res* **39**, 3935–3941 (1979).
69. Shimokawa K, Kasai Z. Ethylene Formation from Ethyl Moiety of Ethionine. *Science* **156**, 1362–1363 (1967).
70. Yang SF. Photosensitized conversion of ethionine to ethylene by flavin mononucleotide. *Photochem Photobiol* **12**, 419–422 (1970).
71. Yang Z-Y, Dean DR, Seefeldt LC. Molybdenum nitrogenase catalyzes the reduction and coupling of CO to form hydrocarbons. *J Biol Chem* **286**, 19417–19421 (2011).
72. de Zélicourt A, *et al.* Ethylene induced plant stress tolerance by Enterobacter sp. SA187 is mediated by 2-keto-4-methylthiobutyric acid production. *PLOS Genet* **14**, e1007273 (2018).
73. Jaber Sathik Rifayee SB, Thomas MG, Christov C. Revealing the nature of the second branch point in the catalytic mechanism of the Fe(II)/2OG-dependent ethylene forming enzyme. *Chem Sci* **16**, 7667–7684 (2025).
74. Thomas MG, Jaber Sathik Rifayee SB, Christov CZ. How do variants of residues in the first coordination sphere, second coordination sphere, and remote areas influence the catalytic

mechanism of non-Heme Fe(II)/2-oxoglutarate dependent ethylene-forming enzyme? *ACS Catal* **14**, 18550–18569 (2024).
75. Wang C, Sayfutyarova ER. Diverging reaction pathways and key intermediates in ethylene forming enzyme. *J Phys Chem B* **129**, 4335–4349 (2025).
76. Chaturvedi SS, Ramanan R, Hu J, Hausinger RP, Christov CZ. Atomic and electronic structure determinants distinguish between ethylene formation and L-arginine hydroxylation reaction mechanisms in the ethylene-forming enzyme. *ACS Catal* **11**, 1578–1592 (2021).
77. Xue J, Lu J, Lai W. Mechanistic insights into a non-heme 2-oxoglutarate-dependent ethylene-forming enzyme: selectivity of ethylene-formation versusl-Arg hydroxylation. *Phys Chem Chem Phys* **21**, 9957–9968 (2019).
78. Yeh CCG, Ghafoor S, Satpathy JK, Mokkawes T, Sastri CV, de Visser SP. Cluster Model Study into the Catalytic Mechanism of α-Ketoglutarate Biodegradation by the Ethylene-Forming Enzyme Reveals Structural Differences with Nonheme Iron Hydroxylases. *ACS Catal* **12**, 3923–3937 (2022).
79. Althoff F, *et al.* Abiotic methanogenesis from organosulphur compounds under ambient conditions. *Nat Commun* **5**, 4205 (2014).
80. Joseph CA, Maroney MJ. Cysteine dioxygenase: structure and mechanism. *Chem Commun*, 3338–3349 (2007).
81. Stipanuk MH, Simmons CR, Andrew Karplus P, Dominy JE. Thiol dioxygenases: unique families of cupin proteins. *Amino Acids* **41**, 91–102 (2011).
82. McDonough MA, Kavanagh KL, Butler D, Searls T, Oppermann U, Schofield CJ. Structure of Human Phytanoyl-CoA 2-Hydroxylase Identifies Molecular Mechanisms of Refsum Disease. *J Biol Chem* **280**, 41101–41110 (2005).
83. Mukherji M, *et al.* 'Chemical co-substrate rescue' of phytanoyl-CoA 2-hydroxylase mutants causing Refsum's Disease. *Chem Commun*, 972–973 (2001).
84. Horton JR, Upadhyay AK, Hashimoto H, Zhang X, Cheng X. Structural basis for human PHF2 jumonji domain interaction with metal ions. *J Mol Biol* **406**, 1–8 (2011).
85. Liu L, Kim H, Casta A, Kobayashi Y, Shapiro LS, Christiano AM. Hairless is a histone H3K9 demethylase. *FASEB J* **28**, 1534–1542 (2014).
86. Pfeffer I, *et al.* Aspartate/asparagine-β-hydroxylase crystal structures reveal an unexpected epidermal growth factor-like domain substrate disulfide pattern. *Nat Commun* **10**, 4910 (2019).
87. Hewitson KS, *et al.* Evidence that two enzyme-derived histidine ligands are sufficient for iron binding and catalysis by factor inhibiting HIF (FIH). *J Biol Chem* **283**, 25971–25978 (2008).
88. Hangasky JA, Taabazuing CY, Martin CB, Eron SJ, Knapp MJ. The facial triad in the α-ketoglutarate dependent oxygenase FIH: A role for sterics in linking substrate binding to O2 activation. *J Inorg Biochem* **166**, 26–33 (2017).
89. Brasnett A, *et al.* Human oxygenase variants employing a single protein Fe(II) ligand are catalytically active. *Angew Chem Int Ed* **60**, 14657–14663 (2021).
90. Zou Q, *et al.* Hydroxylase Activity of ASPH Promotes Hepatocellular Carcinoma Metastasis Through Epithelial-to-Mesenchymal Transition Pathway. *EBioMedicine* **31**, 287–298 (2018).
91. Morishita Y, *et al.* Fused radical SAM and αKG-HExxH domain proteins contain a distinct structural fold and catalyse cyclophane formation and β-hydroxylation. *Nat Chem* **16**, 1882–1893 (2024).
92. Adams DO, Yang SF. Methionine Metabolism in Apple Tissue: Implication of S-Adenosylmethionine as an Intermediate in the Conversion of Methionine to Ethylene 1. *Plant Physiol* **60**, 892–896 (1977).

93. Adams DO, Yang SF. Ethylene biosynthesis: Identification of 1-aminocyclopropane-1-carboxylic acid as an intermediate in the conversion of methionine to ethylene. *Proc Natl Acad Sci USA* **76**, 170–174 (1979).
94. Perpète P, *et al.* Methionine catabolism in Saccharomyces cerevisiae. *FEMS Yeast Res* **6**, 48–56 (2006).
95. Ren T, Patel M, Blok K. Olefins from conventional and heavy feedstocks: Energy use in steam cracking and alternative processes. *Energy* **31**, 425–451 (2006).
96. Pichler CM, *et al.* Bio-Electrocatalytic Conversion of Food Waste to Ethylene via Succinic Acid as the Central Intermediate. *ACS Catal* **12**, 13360–13371 (2022).
97. Digiacomo F, *et al.* Ethylene-producing bacteria that ripen fruit. *ACS Synth Biol* **3**, 935–938 (2014).
98. Wang J, Lin M, Xu M, Yang S-T. Anaerobic Fermentation for Production of Carboxylic Acids as Bulk Chemicals from Renewable Biomass. In: *Anaerobes in Biotechnology* (eds Hatti-Kaul R, Mamo G, Mattiasson B). Springer International Publishing (2016).
99. Joharapurkar AA, Pandya VB, Patel VJ, Desai RC, Jain MR. Prolyl Hydroxylase inhibitors: a breakthrough in the therapy of anemia associated with chronic diseases. *J Med Chem* **61**, 6964–6982 (2018).
100. Locatelli F, Minutolo R, De Nicola L, Del Vecchio L. Evolving strategies in the treatment of anaemia in chronic kidney disease: the HIF-prolyl hydroxylase inhibitors. *Drugs* **82**, 1565–1589 (2022).
101. Brewitz L, Tumber A, Thalhammer A, Salah E, Christensen KE, Schofield CJ. Synthesis of novel pyridine-carboxylates as small-molecule inhibitors of human aspartate/asparagine-β-hydroxylase. *ChemMedChem* **15**, 1139–1149 (2020).
102. Brewitz L, Nakashima Y, Tumber A, Salah E, Schofield CJ. Fluorinated derivatives of pyridine-2,4-dicarboxylate are potent inhibitors of human 2-oxoglutarate dependent oxygenases. *J Fluor Chem* **247**, 109804 (2021).
103. Brewitz L, *et al.* 5-Substituted Pyridine-2,4-dicarboxylate Derivatives Have Potential for Selective Inhibition of Human Jumonji-C Domain-Containing Protein 5. *J Med Chem* **66**, 10849–10865 (2023).
104. McDonough MA, *et al.* Selective inhibition of factor inhibiting hypoxia-inducible factor. *J Am Chem Soc* **127**, 7680–7681 (2005).
105. Windsor P, Chatterjee S, Damodaran AR, Bhagi-Damodaran A. 2-oxoglutarate analogue-based biomolecular tools for exploring structure-activity relationships in non-heme iron enzymes. *ChemBioChem* **n/a**, e202500177 (2025).
106. Liu X, *et al.* Natural and synthetic 2-oxoglutarate derivatives are substrates for oncogenic variants of human isocitrate dehydrogenase 1 and 2. *J Biol Chem* **299**, 102873 (2023).
107. Adams RW, Holroyd CM, Aguilar JA, Nilsson M, Morris GA. “Perfecting” WATERGATE: clean proton NMR spectra from aqueous solution. *Chem Commun* **49**, 358–360 (2013).

5.10 Supporting Figures


^{1}H NMR assays (700 MHz or 600 MHz) were performed using a Bruker AVIII 600 instrument equipped with a Prodigy nitrogen broadband cryoprobe or using a Bruker AVIII 700 MHz NMR spectrometer equipped with a 5-mm inverse triple-resonance-inverse cryoprobe.


For ^{1}H NMR experiments, water suppression at ~4.7 ppm was achieved using perfect echo-modified WATERGATE experiments (1). 3-(Trimethylsilyl)-2,2,3,3-tetradeuteropropionic acid,(2) TMSP-d_4) was used as the internal standard (0 ppm) to enable substrate and product quantification via comparison of peak integrals with that of TMSP-d_4. The stereochemistry of P5C was not assigned; note that P5C is in equilibrium with *L*-glutamate-5-semialdehyde(3, 4) and may epimerize under the assay conditions.

Symbols are used to annotate signals in the ^{1}H NMR spectra as shown below:


Annotation	Molecule(s)
squares	*L*-ascorbic acid (LAA)
pentagons	*L*-arginine (Arg)
inverted triangles	*L*-Δ^1-pyrroline-5-carboxylic acid (P5C)
circles	2-oxoacids
diamonds/triangles	PsEFE reaction products

Colors are used to distinguish between chemically distinct protons, as shown below.







Supporting Figure 5.S1. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of KMBA. (**a**) ^{1}H NMR analysis of the reaction of wt PsEFE with KMBA ~75 min post incubation with wt PsEFE reveals formation of pyrroline-5-carboxylic acid (P5C). Low levels of ethylene are formed likely in a non-enzymatic manner; (**b** and **c**) ^{1}H NMR spectra of KMBA under standard assay conditions in the absence of wt PsEFE: (**b**) ~75 min and (**c**) ~5 min post addition of KMBA to the reaction mixture. The results indicate low levels of ethylene are formed from KMBA in a non-enzymatic manner (Supporting Figure 5.S78). Conditions: 400 µM KMBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Peaks corresponding to the hydrated ketone form of KMBA are annotated with dashed circles (Supporting Figure 5.S2). Signals at ~3.6 and ~3.7 ppm correspond to glycerol. Representative spectra from independent triplicates are shown.




Supporting Figure 5.S2. KMBA is present as ketone and hydrated ketone forms in aqueous buffer. (**a**) ^{1}H NMR (700 MHz) spectrum (~1.9 to ~4.2 ppm) of KMBA in buffer shows characteristic signals corresponding to both the ketone form (closed circles) and low levels of the hydrated ketone form (dashed circles) of KMBA; (**b**) ^{13}C NMR (175 MHz) analysis of KMBA supports its presence in ketone (~205 ppm; red B) and hydrated ketone forms (gem-diol, ~95 ppm; orange B); (**c** and **d**) 2D NMR analysis of KMBA using: (**c**) ^{1}H,^{1}H-COSY and (**d**) ^{1}H-^{13}C HMBC supports the assignment of signals corresponding to KMBA in its ketone or hydrated ketone form. Conditions: ~30 mM KMBA in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O).




Supporting Figure 5.S3. wt PsEFE catalyzed time-dependent oxidation of KMBA. (**a**-**c**) ^{1}H NMR analysis (~2.0 to ~3.9 ppm) of a reaction of wt PsEFE with KMBA: (**a**) ~25 min, (**b**) ~15 min, and (**c**) ~5 min post addition of wt PsEFE to the reaction mixture. (**d**) ^{1}H NMR spectrum of the reaction mixture containing KMBA in the absence of PsEFE. Conditions: 400 µM KMBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a**-**c**) are from glycerol. Peaks corresponding to the hydrated ketone form of KMBA are annotated with dashed circles (Supporting Figure 5.S2). Note that wt PsEFE catalyzed P5C formation in the presence of KMBA (Supporting Figure 5.S1). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S4. ^{1}H NMR analysis of KMBA and authentic samples of the acid (1a) and alcohol (1b) products formed by PsEFE-catalyzed oxidation of KMBA. (**a**-**c**) ^{1}H NMR spectra (~2.0 to ~3.9 ppm) of: (**a**) the alcohol product, 2-(methylthio)ethanol (**1b**), (**b**) the acid product, 3-(methylthio)propionic acid (**1a**), and (**c**) KMBA in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Peaks corresponding to the hydrated form of KMBA are annotated with dashed circles (Supporting Figure 5.S2).




Supporting Figure 5.S5. Analysis of the reaction of wt PsEFE with KMBA. ^{1}H NMR analysis (~2.0 to ~3.9 ppm) of: (**a**) a reaction of wt PsEFE and KMBA in the presence of an authentic sample of 2-(methylthio)ethanol (**1b**), (**b**) ^{1}H NMR spectrum of an authentic sample of **1b** under standard assay conditions in the presence of wt PsEFE; (**c**) a reaction of wt PsEFE and KMBA in the presence of an authentic sample of 3-(methylthio)propionic acid (**1a**), (**d**) ^{1}H NMR spectrum of an authentic sample of **1a** under standard assay conditions in the presence of wt PsEFE; (**e**) ^{1}H NMR spectrum of a reaction of wt PsEFE and KMBA after ~25 min incubation; (**f**) ^{1}H NMR spectrum of KMBA under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM KMBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM wt PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The singlet at ~3.7 ppm in (**a**-**d**) corresponds to Tris. Peaks corresponding to the hydrated form of KMBA are annotated with dashed circles (Supporting Figure 5.S2). Note, wt PsEFE catalyzed P5C formation in the presence of KMBA, but not ethylene production (Supporting Figure 5.S1). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S6. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of 4-methyl-2-oxopentanoic acid (2). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 4-methyl-2-oxopentanoic acid (**2**) after ~75 min incubation reveals formation of P5C; (**b**) ^{1}H NMR spectrum of a mixture containing **2** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **2**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S7. ^{1}H NMR analysis of 4-methyl-2-oxopentanoic acid (2) and authentic samples of the acid (2a) and alcohol (2b) products formed by wt PsEFE-catalyzed oxidation of 2. (**a**-**c**) ^{1}H NMR spectra (~0.5 to ~3.5 ppm) of: (**a**) the alcohol product, 2-methylpropan-1-ol (**2b**), (**b**) the acid product, 3-methylbutanoic acid (**2a**), and (**c**) 4-methyl-2-oxopentanoic acid (**2**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O).




Supporting Figure 5.S8. Analysis of the reaction of wt PsEFE with 4-methyl-2-oxopentanoic acid (2). ^{1}H NMR analysis (~0.8 to ~3.8 ppm) of: (**a**) a reaction of wt PsEFE and 4-methyl-2-oxopentanoic acid (**2**) in the presence of an authentic sample of 2-methylpropan-1-ol (**2b**), (**b**) authentic sample of **2b**; (**c**) a reaction of wt PsEFE and **2** in the presence of an authentic sample of 3-methylbutanoic acid (**2a**), (**d**) authentic sample of **2a**; (**e**) a reaction of wt PsEFE and **2** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **2** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **2**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**c**) are from glycerol; the singlet at ~3.7 ppm in (**a**) and (**c**) corresponds to Tris. Note, wt PsEFE catalyzed P5C formation in the presence of **2** (Supporting Figure 5.S6). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S9. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of 3-methyl-2-oxopentanoic acid (3). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 3-methyl-2-oxopentanoic acid (**3**) after ~75 min incubation reveals formation of P5C; (**b**) ^{1}H NMR spectrum of **3** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **3**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-*d*$_4$, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol; low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed in (**a**) and (**b**).(5) Representative spectra from independent triplicates are shown.




Supporting Figure 5.S10. ^{1}H NMR analysis of 3-methyl-2-oxopentanoic acid (3) and authentic samples of the acid (3a) and alcohol (3b) products formed by wt PsEFE-catalyzed oxidation of 3. (**a**-**c**) ^{1}H NMR spectra (~0.7 to ~3.9 ppm) of: (**a**) the alcohol product, butan-2-ol (**3b**), (**b**) the acid product, 2-methylbutanoic acid (**3a**), and (**c**) 3-methyl-2-oxopentanoic acid (**3**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O).




Supporting Figure 5.S11. Analysis of the reaction of wt PsEFE with 3-methyl-2-oxopentanoic acid (3). ^{1}H NMR analysis (~0.8 to ~3.9 ppm) of: (**a**) a reaction of wt PsEFE and 3-methyl-2-oxopentanoic acid (**3**) in the presence of an authentic sample of butan-2-ol (**3b**), (**b**) authentic sample of **3b**; (**c**) a reaction of wt PsEFE and **3** in the presence of an authentic sample of 2-methylbutanoic acid (**3a**), (**d**) authentic sample of **3a**; (**e**) a reaction of wt PsEFE and **3** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **3** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **3**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm in (**a**) and (**c**) corresponds to Tris. Note, wt PsEFE catalyzed P5C formation in the presence of **3** (Supporting Figure 5.S9). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S12. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of 3-methyl-2-oxobutanoic acid (4). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 3-methyl-2-oxobutanoic acid (**4**) after ~75 min incubation reveals formation of P5C; (**b**) ^{1}H NMR spectrum of **4** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **4**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-*d*$_4$, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol; low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed.(5) The broad peaks at ~0.5 to ~1.5 ppm in (**a**) are likely from the PsEFE sample, which was used in a relatively high concentration. Representative spectra from independent triplicates are shown.




Supporting Figure 5.S13. ^{1}H NMR analysis of 3-methyl-2-oxobutanoic acid (4) and authentic samples of the acid (4a) and alcohol (4b) products formed by wt PsEFE-catalyzed oxidation of 4. (**a**-**c**) ^{1}H NMR spectra (~0.8 to ~4.1 ppm) of: (**a**) the alcohol product, propan-2-ol (**4b**), (**b**) the acid product, isobutyric acid (**4a**), and (**c**) 3-methyl-2-oxobutanoic acid (**3**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O).




Supporting Figure 5.S14. Analysis of the reaction of wt PsEFE with 3-methyl-2-oxobutanoic acid (4). ^{1}H NMR analysis (~0.8 to ~4.2 ppm) of: (**a**) a reaction of wt PsEFE and 3-methyl-2-oxobutanoic acid (**4**) in the presence of an authentic sample of propan-2-ol (**4b**), (**b**) authentic sample of **4b**; (**c**) a reaction of wt PsEFE and **4** in the presence of an authentic sample of isobutyric acid (**4a**), (**d**) authentic sample of **4a**; (**e**) a reaction of wt PsEFE and **4** after ~75 min incubation; (**f**) a reaction of wt PsEFE and **4** after ~5 min incubation. Conditions: 400 µM **4**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm in (**a**) and (**c**) corresponds to Tris. The broad peaks at ~0.5 to ~1.5 ppm in (**e**) are likely from the PsEFE sample, which was used in a relatively high concentration. Note, wt PsEFE catalyzed P5C formation in the presence of **4** (Supporting Figure 5.S12). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S15. wt PsEFE catalyzes conversion of *L*-arginine to low levels of P5C in the presence of 2-oxobutanoic acid (5). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 2-oxobutanoic acid (**5**) after ~75 min incubation reveals formation of low levels of P5C; (**b**) ^{1}H NMR spectrum of **5** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **5**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol; low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed.(5) The broad peaks at ~0.5 to ~1.5 ppm in (**a**) are likely from the PsEFE sample, which was used in a relatively high concentration. Representative spectra from independent triplicates are shown. Note that **5** contained low levels of **5a**.




Supporting Figure 5.S16. ^{1}H NMR analysis of 2-oxobutanoic acid (5) and authentic samples of the acid (5a) and alcohol (5b) products formed by wt PsEFE-catalyzed oxidation of 5. (**a**-**c**) ^{1}H NMR spectra (~1.0 to ~3.8 ppm) of: (**a**) the alcohol product, ethanol (**5b**), (**b**) the acid product, propanoic acid (**5a**), and (**c**) 2-oxobutanoic acid (**5**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Note that **5** contained low levels of **5a**.




Supporting Figure 5.S17. Analysis of the reaction of wt PsEFE with 2-oxobutanoic acid (5). ^{1}H NMR analysis (~1.0 to ~3.8 ppm) of: (**a**) a reaction of wt PsEFE and 2-oxobutanoic acid (**5**) in the presence of an authentic sample of ethanol (**5b**), (**b**) authentic sample of **5b**; (**c**) a reaction of wt PsEFE and **5** in the presence of an authentic sample of propanoic acid (**5a**), (**d**) authentic sample of **5a**; (**e**) a reaction of wt PsEFE and **5** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **5** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **5**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a**,**c**,**e**) are from glycerol; the singlet at ~3.7 ppm in (**a**,**c**,**e**) corresponds to Tris. Note that **5** contained low levels of **5a** and that wt PsEFE catalyzed P5C formation in the presence of **5** (Supporting Figure 5.S15). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S18. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of 2-oxovaleric acid (6). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 2-oxovaleric acid (**6**) after ~75 min incubation reveals formation of P5C; (**b**) ^{1}H NMR spectrum of **6** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **6**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol; low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed in (**a**).(5) Representative spectra from independent triplicates are shown.




Supporting Figure 5.S19. ^{1}H NMR analysis of 2-oxovaleric acid (6) and authentic samples of the acid (6a) and alcohol (6b) products formed by wt PsEFE-catalyzed oxidation of 6. (**a**-**c**) ^{1}H NMR spectra (~0.6 to ~3.7 ppm) of: (**a**) the alcohol product, propanol (**6b**), (**b**) the acid product, butanoic acid (**6a**), and (**c**) 2-oxovaleric acid (**6**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O).




Supporting Figure 5.S20. Analysis of the reaction of wt PsEFE with 2-oxovaleric acid (6). ^{1}H NMR analysis (~0.8 to ~3.6 ppm) of: (**a**) a reaction of wt PsEFE and 2-oxovaleric acid (**6**) in the presence of an authentic sample of propanol (**6b**), (**b**) authentic sample of **6b**; (**c**) a reaction of wt PsEFE and **6** in the presence of an authentic sample of butanoic acid (**6a**), (**d**) authentic sample of **6a**; (**e**) a reaction of wt PsEFE and **6** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **6** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **6**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM wt PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Note that wt PsEFE catalyzed P5C formation in the presence of **6** (Supporting Figure 5.S18). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S21. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of 2-oxocaproic acid (7). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 2-oxocaproic acid (**7**) after ~75 min incubation reveals formation of low levels of P5C; (**b**) ^{1}H NMR spectrum of **7** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **7**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol; the singlet at ~3.7 ppm in (**a**) corresponds to Tris. The broad peaks at ~0.5 to ~1.5 ppm in (**a**) are likely from the PsEFE sample, which was used in a relatively high concentration. Representative spectra from independent triplicates are shown.




Supporting Figure 5.S22. ^{1}H NMR analysis of 2-oxocaproic acid (7) and authentic samples of the acid (7a) and alcohol (7b) products formed by wt PsEFE-catalyzed oxidation of 7. (**a**-**c**) ^{1}H NMR spectra (~0.7 to ~3.7 ppm) of: (**a**) the alcohol product, butanol (**7b**), (**b**) the acid product, pentanoic acid (**7a**), and (**c**) 2-oxocaproic acid (**7**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O).




Supporting Figure 5.S23. Analysis of the reaction of wt PsEFE with 2-oxocaproic acid (7). ^{1}H NMR analysis (~0.8 to ~3.7 ppm) of: (**a**) a reaction of wt PsEFE and 2-oxocaproic acid (**7**) in the presence of an authentic sample of butanol (**7b**), (**b**) authentic sample of **7b**; (**c**) a reaction of wt PsEFE and **7** in the presence of an authentic sample of pentanoic acid (**7a**), (**d**) authentic sample of **7a**; (**e**) a reaction of wt PsEFE and **7** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **7** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **7**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM wt PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Note that wt PsEFE catalyzed P5C formation in the presence of **7** (Supporting Figure 5.S21). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S24. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of 2-cyclohexyl-2-oxoacetic acid (8). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 2-cyclohexyl-2-oxoacetic acid (**8**) after ~75 min incubation reveals formation of P5C; (**b**) ^{1}H NMR spectrum of **8** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **8**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-*d*$_4$, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual cyclohexyl protons, except for those at the β position of **8**, the α position of **8a**, and the C1 position of **8b**. The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol. The broad peaks at ~0.5 to ~1.5 ppm in (**a**) are likely from the PsEFE sample, which was used in a relatively high concentration. Representative spectra from independent triplicates are shown.




Supporting Figure 5.S25. ^{1}H NMR analysis of 2-cyclohexyl-2-oxoacetic acid (8) and authentic samples of the acid (8a) and alcohol (8b) products formed by wt PsEFE-catalyzed oxidation of 8. (**a-c**) ^{1}H NMR spectra (~1.0 to ~4.0 ppm) of: (**a**) the alcohol product, cyclohexanol (**8b**), (**b**) the acid product, cyclohexanecarboxylic acid (**8a**), and (**c**) 2-cyclohexyl-2-oxoacetic acid (**8**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual cyclohexyl protons, except for those at the β position of **8**, the α position of **8a**, and the C1 position of **8b**.




Supporting Figure 5.S26. Analysis of the reaction of wt PsEFE with 2-cyclohexyl-2-oxoacetic acid (8). ^{1}H NMR analysis (~1.0 to ~3.9 ppm) of: (**a**) a reaction of wt PsEFE and 2-cyclohexyl-2-oxoacetic acid (**8**) in the presence of an authentic sample of cyclohexanol (**8b**), (**b**) authentic sample of **8b**; (**c**) a reaction of wt PsEFE and **8** in the presence of an authentic sample of cyclohexanecarboxylic acid (**8a**), (**d**) authentic sample of **8a**; (**e**) a reaction of wt PsEFE and **8** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **8** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **8**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM wt PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual cyclohexyl protons, except for those at the β position of **8**, the α position of **8a**, and the C1 position of **8b**. The peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Note that wt PsEFE catalyzed P5C formation in the presence of **8** (Supporting Figure 5.S24). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S27. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of 3-cyclohexyl-2-oxopropanoic acid (9). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 3-cyclohexyl-2-oxopropanoic acid (**9**) after ~75 min incubation reveals formation of P5C; (**b**) ^{1}H NMR spectrum of **9** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **9**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual cyclohexyl protons, except for those at the β and γ positions of **9**, the α position of **9a**, and the C1 position of **9b**. The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol; low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed in (**b**).(5) Representative spectra from independent triplicates are shown.




Supporting Figure 5.S28. ^{1}H NMR analysis of 3-cyclohexyl-2-oxopropanoic acid (9) and authentic samples of the acid (9a) and alcohol (9b) products formed by wt PsEFE-catalyzed oxidation of 9. (**a**-**c**) ^{1}H NMR spectra (~0.6 to ~3.6 ppm) of: (**a**) the alcohol product, cyclohexylmethanol (**9b**), (**b**) the acid product, 2-cyclohexylacetic acid (**9a**), and (**c**) 3-cyclohexyl-2-oxopropanoic acid (**9**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual cyclohexyl protons, except for those at the β and γ positions of **9**, the α position of **9a**, and the C1 position of **9b**.




Supporting Figure 5.S29. Analysis of the reaction of wt PsEFE with 3-cyclohexyl-2-oxopropanoic acid (9). ^{1}H NMR analysis (~0.6 to ~3.6 ppm) of: (**a**) a reaction of wt PsEFE and 3-cyclohexyl-2-oxopropanoic acid (**9**) in the presence of an authentic sample of cyclohexylmethanol (**9b**), (**b**) authentic sample of **9b**; (**c**) a reaction of wt PsEFE and **9** in the presence of an authentic sample of 2-cyclohexylacetic acid (**9a**), (**d**) authentic sample of **9a**; (**e**) a reaction of wt PsEFE and **9** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **9** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **9**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM wt PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual cyclohexyl protons, except for those at the β and γ positions of **9**, the α position of **9a**, and the C1 position of **9b**. Note that wt PsEFE catalyzed P5C formation in the presence of **9** (Supporting Figure 5.S27). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S30. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of 4-cyclohexyl-2-oxobutanoic acid (10). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 4-cyclohexyl-2-oxobutanoic acid (**10**) after ~75 min incubation reveals formation of P5C; (**b**) ^{1}H NMR spectrum of **10** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **10**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual cyclohexyl protons, except for those at the β and γ positions of **10**, the α and β positions of **10a**, and the C1 and C2 positions of **10b**. The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol; low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed in (**b**).(5) Representative spectra from independent triplicates are shown.




Supporting Figure 5.S31. ^{1}H NMR analysis of 4-cyclohexyl-2-oxobutanoic acid (10) and authentic samples of the acid (10a) and alcohol (10b) products formed by wt PsEFE-catalyzed oxidation of 10. (**a**-**c**) ^{1}H NMR spectra (~0.7 to ~4.0 ppm) of: (**a**) the alcohol product, 2-cyclohexylethan-1-ol (**10b**), (**b**) the acid product, 3-cyclohexylpropanoic acid (**10a**), and (**c**) 4-cyclohexyl-2-oxobutanoic acid (**10**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual cyclohexyl protons, except for those at the β and γ positions of **10**, the α and β positions of **10a**, and the C1 and C2 positions of **10b**.




Supporting Figure 5.S32. Analysis of the reaction of wt PsEFE with 4-cyclohexyl-2-oxobutanoic acid (10). ^{1}H NMR analysis (~0.7 to ~4.0 ppm) of: (**a**) a reaction of wt PsEFE and 4-cyclohexyl-2-oxobutanoic acid (**10**) in the presence of an authentic sample of 2-cyclohexylethan-1-ol (**10b**), (**b**) authentic sample of **10b**; (**c**) a reaction of wt PsEFE and **10** in the presence of an authentic sample of 3-cyclohexylpropanoic acid (**10a**), (**d**) authentic sample of **10a**; (**e**) a reaction of wt PsEFE and **10** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **10** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **10**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM wt PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual cyclohexyl protons, except for those at the β and γ positions of **10**, the α and β positions of **10a**, and the C1 and C2 positions of **10b**. Note that wt PsEFE catalyzed P5C formation in the presence of **10** (Supporting Figure 5.S30). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S33. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of 4,4-dimethyl-2-oxopentanoic acid (11). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 4,4-dimethyl-2-oxopentanoic acid (**11**) after ~75 min incubation reveals formation of P5C; (**b**) ^{1}H NMR spectrum of **11** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **11**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol; low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed.(5) Representative spectra from independent triplicates are shown.




Supporting Figure 5.S34. ^{1}H NMR analysis of 4,4-dimethyl-2-oxopentanoic acid (11) and authentic samples of the acid (11a) and alcohol (11b) products formed by wt PsEFE-catalyzed oxidation of 11. (**a**-**c**) ^{1}H NMR spectra (~0.7 to ~3.5 ppm) of: (**a**) the alcohol product, neopentanol (**11b**), (**b**) the acid product, 3,3-dimethylbutanoic acid (**11a**), and (**c**) 4,4-dimethyl-2-oxopentanoic acid (**11**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O).




Supporting Figure 5.S35. Analysis of the reaction of wt PsEFE with 4,4-dimethyl-2-oxopentanoic acid (11). ^{1}H NMR analysis (~0.7 to ~3.5 ppm) of: (**a**) a reaction of wt PsEFE and 4,4-dimethyl-2-oxopentanoic acid (**11**) in the presence of an authentic sample of neopentanol (**11b**), (**b**) authentic sample of **11b**; (**c**) a reaction of wt PsEFE and **11** in the presence of an authentic sample of 3,3-dimethylbutanoic acid (**11a**), (**d**) authentic sample of **11a**; (**e**) a reaction of wt PsEFE and **11** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **11** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **11**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM wt PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Note that wt PsEFE catalyzed P5C formation in the presence of **11** (Supporting Figure 5.S33). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S36. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of 2-oxo-4-phenylbutanoic acid (12). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 2-oxo-4-phenylbutanoic acid (**12**) after ~75 min incubation reveals formation of P5C; (**b**) ^{1}H NMR spectrum of **12** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **12**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual phenyl protons. Peaks corresponding to the hydrated ketone form of **12** are annotated with dashed circles; the assignment of the peaks corresponding to the hydrated form of **12** was based on that for the hydrated ketone forms of KMBA (Supporting Figure 5.S2) and 2OG.(6, 7). Low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed in (**b**).(5) The broad peaks at ~0.5 to ~1.5 ppm in (**a**) are likely from the PsEFE sample, which was used in a relatively high concentration. Representative spectra from independent triplicates are shown.




Supporting Figure 5.S37. ^{1}H NMR analysis of 2-oxo-4-phenylbutanoic acid (12) and authentic samples of the acid (12a) and alcohol (12b) products formed by wt PsEFE-catalyzed oxidation of 12. (**a**-**c**) ^{1}H NMR spectra (~2.2 to ~7.6 ppm) of: (**a**) the alcohol product, 2-phenylethanol (**12b**), (**b**) the acid product, 3-phenylpropanoic acid (**12a**), and (**c**) 2-oxo-4-phenylbutanoic acid (**12**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual phenyl protons. Peaks corresponding to the hydrated ketone form of **12** are annotated with dashed circles; the assignment of the peak corresponding to the hydrated form of **12** was based on that for the hydrated ketone forms of KMBA (Supporting Figure 5.S2) and 2OG.(6, 7).




Supporting Figure 5.S38. Analysis of the reaction of wt PsEFE with 2-oxo-4-phenylbutanoic acid (12). ^{1}H NMR analysis (~2.2 to ~7.8 ppm) of: (**a**) a reaction of wt PsEFE and 2-oxo-4-phenylbutanoic acid (**12**) in the presence of an authentic sample of 2-phenylethanol (**12b**), (**b**) authentic sample of **12b**; (**c**) a reaction of wt PsEFE and **12** in the presence of an authentic sample of 3-phenylpropanoic acid (**12a**), (**d**) authentic sample of **12a**; (**e**) a reaction of wt PsEFE and **12** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **12** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **12**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM wt PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Peaks corresponding to the hydrated ketone form of **12** are annotated with dashed circles; the assignment of the peaks corresponding to the hydrated form of **12** was based on that for the hydrated ketone forms of KMBA (Supporting Figure 5.S2) and 2OG.(6, 7). The peaks at ~3.6 and ~3.7 ppm in (**c**) are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. ^{1}H NMR signals were not assigned to individual phenyl protons. Note that wt PsEFE catalyzed P5C formation in the presence of **12** (Supporting Figure 5.S36). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S39. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of phenylpyruvic acid (PPA, 13). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with phenylpyruvic acid (PPA, **13**) after ~75 min incubation reveals formation of P5C. The concentration of wt PsEFE was increased 3-fold with respect to the standard assay conditions (30 µM), likely resulting in decreased signal-to-noise ratio due to increased levels of peaks/contaminants from the protein sample; (**b**) ^{1}H NMR spectrum of **13** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **13**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 30 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual phenyl protons and that commercially-sourced **13** contained low amounts of **13a**. The methylene protons of **13b** were not observed, likely because they coincide with the HOD peak. The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol. The broad peaks at ~0.5 to ~1.5 ppm in (**a**) are likely from the PsEFE sample, which was used in a relatively high concentration. Representative spectra from independent triplicates are shown.



Supporting Figure 5.S40. ^{1}H NMR analysis of phenylpyruvic acid (PPA, 13) and authentic samples of the acid (13a) and alcohol (13b) products formed by wt PsEFE-catalyzed oxidation of 13. (**a**-**c**) ^{1}H NMR spectra (~2.9 to ~4.2 ppm and ~7.0 to ~7.6 ppm) of: (**a**) the alcohol product, benzyl alcohol (**13b**), (**b**) the acid product, 2-phenylacetic acid (**13a**), and (**c**) phenylpyruvic acid (PPA, **13**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual phenyl protons and that commercially-sourced **13** contained low amounts of **13a**. The methylene protons of **13b** were not observed, likely because they coincide with the HOD peak.




Supporting Figure 5.S41. Analysis of the reaction of wt PsEFE with phenylpyruvic acid (PPA, 13). ^{1}H NMR analysis (~3.0 to ~4.2 ppm and ~7.0 to ~7.6 ppm) of: (**a**) a reaction of wt PsEFE and phenylpyruvic acid (PPA, **13**) in the presence of an authentic sample of benzyl alcohol (**13b**), (**b**) authentic sample of **13b**; (**c**) a reaction of wt PsEFE and **13** in the presence of an authentic sample of 2-phenylacetic acid (**13a**), (**d**) authentic sample of **13a**; (**e**) a reaction of wt PsEFE and **13** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **13** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **13**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 30 µM wt PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual phenyl protons. Commercially-sourced **13** contained low amounts of **13a**; thus, conversion was not quantified. The methylene protons of **13b** were not observed, likely because they coincide with the HOD peak. The peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Note that wt PsEFE catalyzed P5C formation in the presence of **13** (Supporting Figure 5.S39). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S42. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of 4-hydroxyphenylpyruvic acid (HPPA, 14). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 4-hydroxyphenylpyruvic acid (HPPA, **14**) after ~75 min incubation reveals formation of P5C. The concentration of wt PsEFE was increased 3-fold with respect to the standard assay conditions (30 µM), likely resulting in decreased signal-to-noise ratio due to increased levels of peaks/contaminants from the protein sample; (**b**) ^{1}H NMR spectrum of **14** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **14**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 30 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual phenyl protons. Peaks corresponding to the hydrated ketone form of **14** are annotated with dashed circles. The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol; low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed in (**b**).(5) The broad peaks at ~0.5 to ~1.5 ppm in (**a**) are likely from the PsEFE sample, which was used in a relatively high concentration. Representative spectra from independent triplicates are shown.


.




Supporting Figure 5.S43. ^{1}H NMR analysis of 4-hydroxyphenylpyruvic acid (HPPA, 14) and authentic samples of the acid (14a) and alcohol (14b) products formed by wt PsEFE-catalyzed oxidation of 14. (**a**-**c**) ^{1}H NMR spectra (~3.0 to ~4.6 ppm and ~6.5 to ~7.4 ppm) of: (**a**) the alcohol product, 4-hydroxybenzyl alcohol (**14b**), (**b**) the acid product, 2-(4-hydroxyphenyl)acetic acid (**14a**), and (**c**) 4-hydroxyphenylpyruvic acid (HPPA, **14**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The methylene protons of **14b** were not observed, likely because they coincide with the HOD peak. ^{1}H NMR signals were not assigned to individual phenyl protons. Peaks corresponding to the hydrated ketone form of **14** are annotated with dashed circles.




Supporting Figure 5.S44. Analysis of the reaction of wt PsEFE with 4-hydroxyphenylpyruvic acid (HPPA, 14). ^{1}H NMR analysis (~3.0 to ~4.0 ppm and ~6.5 to ~7.4 ppm) of: (**a**) a reaction of wt PsEFE and 4-hydroxyphenylpyruvic acid (HPPA, **14**) in the presence of an authentic sample of 4-hydroxybenzyl alcohol (**14b**), (**b**) authentic sample of **14b**; (**c**) a reaction of wt PsEFE and **14** in the presence of an authentic sample of 2-(4-hydroxyphenyl)acetic acid (**14a**), (**d**) authentic sample of **14a**; (**e**) a reaction of wt PsEFE and **14** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **14** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **14**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 30 µM wt PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual phenyl protons. Commercially-sourced **14** contained low amounts of **14a**; thus, conversion was not quantified. The peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. The methylene protons of **14b** were not observed, likely because they coincide with the HOD peak. Note that wt PsEFE catalyzed P5C formation in the presence of **14** (Supporting Figure 5.S42). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S45. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of 4-methoxyphenylpyruvic acid (15). (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with 4-methoxyphenylpyruvic acid (**15**) after ~75 min incubation reveals formation of low levels of P5C. The concentration of wt PsEFE was increased 3-fold with respect to the standard assay conditions to 30 µM, likely resulting in decreased signal-to-noise ratio due to increased levels of peaks/contaminants from the protein sample; (**b**) ^{1}H NMR spectrum of **15** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **15**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 30 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual phenyl protons. The methylene protons of **15b** were not observed, likely because they coincide with the HOD peak. The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol; the singlet at ~3.7 ppm in (**a**) corresponds to Tris. Low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed.(5) The broad peaks at ~0.5 to ~1.5 ppm in (**a**) are likely from the PsEFE sample, which was used in a relatively high concentration. Representative spectra from independent triplicates are shown.




Supporting Figure 5.S46. ^{1}H NMR analysis of 4-methoxyphenylpyruvic acid (15) and authentic samples of the acid (15a) and alcohol (15b) products formed by wt PsEFE-catalyzed oxidation of 15. (**a**-**c**) ^{1}H NMR spectra (~3.0 to ~4.6 ppm and ~6.6 to ~7.6 ppm) of: (**a**) the alcohol product, 4-methoxybenzyl alcohol (**15b**), (**b**) the acid product, 2-(4-methoxyphenyl)acetic acid (**15a**), and (**c**) 4-methoxyphenylpyruvic acid (**15**) in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual phenyl protons. The methylene protons of **15b** were not observed, likely because they coincide with the HOD peak.




Supporting Figure 5.S47. Analysis of the reaction of wt PsEFE with 4-methoxyphenylpyruvic acid (15). ^{1}H NMR analysis (~3.0 to ~ 4.6 ppm and ~6.6 to ~7.4 ppm) of: (**a**) a reaction of wt PsEFE and 4-methoxyphenylpyruvic acid (**15**) in the presence of an authentic sample of 4-methoxybenzyl alcohol (**15b**), (**b**) authentic sample of **15b**; (**c**) a reaction of wt PsEFE and **15** in the presence of an authentic sample of 2-(4-methoxyphenyl)acetic acid (**15a**), (**d**) authentic sample of **15a**; (**e**) a reaction of wt PsEFE and **15** after ~75 min incubation; (**f**) ^{1}H NMR spectrum of **15** under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM **15**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 30 µM wt PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). ^{1}H NMR signals were not assigned to individual phenyl protons. The methylene protons of **15b** were not observed, likely because they coincide with the HOD peak. The peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Note that wt PsEFE catalyzed P5C formation in the presence of **15** (Supporting Figure 5.S45). Representative spectra from independent triplicates are shown.




Supporting Figure 5.48. wt PsEFE and R277K PsEFE require *L*-arginine for efficient KMBA turnover. (**a-b**) ^{1}H NMR analysis of a reaction containing KMBA: (**a**) ~10 h and (**b**) ~5 min post addition of R277K PsEFE to the mixture in the absence of *L*-arginine. (**c-d**) ^{1}H NMR analysis of a reaction containing KMBA: (**c**) ~10 h and (**d**) ~5 min post addition of wt PsEFE to the mixture in the absence of *L*-arginine. Conditions: 400 µM KMBA, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE or R277K PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Peaks corresponding to the hydrated ketone form of KMBA are annotated with dashed circles (Supporting Figure 5.S2); the singlet at ~3.7 ppm corresponds to Tris. Note that wt PsEFE catalyzed P5C formation in the presence of KMBA (Supporting Figure 5.S1).




Supporting Figure 5.S49. Effects of pH on the reaction of PsEFE with KMBA. ^{1}H NMR (700 MHz) analysis of the PsEFE-catalyzed oxidation of KMBA (**1**) at variable pH values implies that the pH affects the estimated 3-(methylthio)propionic acid (**1a**):2-(methylthio)ethanol (**1b**) product ratios. The estimated **1a**:**1b** ratios were: (**a**) 3.2 ± 0.01:1 at pH 6.0, (**b**) 2.3 ± 0.1:1 at pH 6.5, (**c**) 1.9 ± 0.1:1 at pH 7.0, (**d**) 1.8 ± 0.1:1 at pH 7.4, (**e**) 2.0 ± 0.01:1 at pH 8.0, and (**f**) 2.1 ± 0.1:1 at pH 8.5. Note that no conversion was observed at pH <5.5 (in: 50 mM sodium acetate, 10%$_{v/v}$ D_2O) or at pH 9.0, under otherwise standard assay conditions. (**g**) Estimated **1a**:**1b** ratios at pH 6.0 – 8.5 after ~1 h incubations. Reaction times are normalized to the end of the first ^{1}H NMR acquisition following PsEFE addition to the reaction mixture (t = 0 min), by which time variable levels of turnover were manifest. Conditions: 400 µM KMBA, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, and 5 µM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O at the indicated pH) at 25 °C; note that full conversion of KMBA was observed when incubating reaction mixtures overnight or at increased PsEFE concentrations. Data are means of two independent duplicates (n = 2; mean ± SD).




Supporting Figure 5.S50. 2OG is a more efficient (co)substrate of wt PsEFE than KMBA in the presence of the broad-spectrum 2OG oxygenase inhibitor *N*-oxalylglycine (NOG). (**a**) wt PsEFE catalyzes conversion of 2OG in the presence of equimolar amounts of NOG (orange inverse triangles), however, substantially less efficiently than in the absence of NOG (blue circles). (**b**) wt PsEFE catalyzes conversion of KMBA in the absence of NOG (green boxes), but, at least, not efficiently in the presence of equimolar amounts of NOG (purple triangles). Conditions: 400 µM 2OG or KMBA, 400 µM NOG (if appropriate), 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 5 µM wt PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Reaction times are normalized to the end of the first ^{1}H NMR acquisition following wt PsEFE addition to the reaction mixture (t = 0 min), by which time variable levels of turnover were manifest. Note that wt PsEFE and NOG were incubated for 5 min before addition to the reaction mixture. Results are the mean of independent duplicates (n = 2; mean ± SD); graphs were plotted using GraphPad Prism 10.3.0 using non-linear regression analysis and a single-phase decay model.




Supporting Figure 5.S51. Mass spectrometric analysis of isolated recombinant R277 PsEFE variants. MS analyses of R277 PsEFE variants (3 μM) in buffer (25 mM Tris, pH 7.8) were performed using an iFunnel Agilent 6550 accurate mass quadrupole time-of-flight (Q-TOF) mass spectrometer operated in the positive ionization mode for: (**a**) R277A PsEFE (calculated mass: 39285 Da; observed mass: 39285 Da), (**b**) R277K PsEFE (calculated mass: 39342 Da; observed mass: 39342 Da), (**c**) R277L PsEFE (calculated mass: 39327 Da; observed mass: 39330 Da), and (**d**) R277F PsEFE (calculated mass: 39361 Da; observed mass: 39364 Da). Raw spectra were deconvoluted using the Agilent MassHunter workstation B.08.00 software.




Supporting Figure 5.S52. SDS-PAGE gel analysis of isolated recombinant R277 PsEFE variants. The PageRuler™ Plus Prestained Protein Ladder (Thermo Fisher) was used. Proteins were separated on a 4-12% Novex™ 330 Tris-glycine gel (Thermo Fisher) according to the manufacturer's protocol. The combined results indicate that the R277 PsEFE variants were obtained in >90% purity.




Supporting Figure 5.S53. The reactivity of R277 PsEFE variants with 2OG. ^{1}H NMR analysis (700 MHz) reveals that none of the tested R277 PsEFE variants catalyze 2OG oxidation under standard assay conditions, *i.e.*, no evidence for ethylene or P5C formation was accrued. (**a**) wt PsEFE (light blue circles) catalyzes the conversion of 2OG to succinate and ethylene, whereas R277A (orange diamonds), R277F (green triangles), R277K (dark blue inverse triangles), and R277L (pink circles) PsEFE did not catalyze 2OG oxidation. This observation contrasts with the reported ability of R277K PsEFE to employ 2OG when using LC-MS-based assays,(8) possibly as a result of different analytical and/or assay conditions. It is possible that the N^{ε}-amino group of K277 may interact with the 2OG C5 carboxylate, thus, stabilizing 2OG binding to the R277K PsEFE active site and enabling productive catalysis. (**b** and **c**) Representative ^{1}H NMR spectra of a reaction mixture containing 2OG: (**b**) ~10 h post addition of R277K PsEFE and (**c**) prior to addition of R277K PsEFE reveals no substantial levels of 2OG turnover. Reaction times are normalized to the end of the first ^{1}H NMR acquisition following PsEFE addition to the reaction mixture (t = 0 min), by which time variable levels of activity were manifest. Conditions: 400 µM 2OG, 500 µM *L*-arginine, 500 µM *L*-ascorbate, 50 µM Fe(II), 800 µM TMSP-d_4, and 2 µM wt PsEFE or 30 µM R277 PsEFE variants in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. The peaks at ~3.6 and ~3.7 ppm in (**b**) are from glycerol; the singlet at ~3.7 ppm in (**b**) corresponds to Tris. Low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed.(5) The peaks at ~2.1 and ~2.3 ppm (dashed circles) correspond to the hydrated ketone form of 2OG.(6, 7) The broad peaks at ~0.5 to ~1.5 ppm in (**b**) are likely from the PsEFE sample, which was used in a relatively high concentration. Representative spectra from independent triplicates are shown.




Supporting Figure 5.S54. R277K PsEFE does not catalyze oxidation of 2-oxoadipic acid (2OA). (**a**-**b**) ^{1}H NMR analysis of a reaction mixture containing 2-oxoadipic acid (2OA): (**a**) ~10 h and (**b**) ~5 min post addition of R277K PsEFE to the reaction mixture. (**c**) ^{1}H NMR analysis of a reaction mixture containing 2OA in the absence of R277K PsEFE. The results reveal R277K PsEFE and, by implication, the other tested R277 PsEFE variants do not employ 2OA as a cosubstrate. Conditions: 400 µM 2OA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 30 µM R277K PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. The peaks at ~3.6 and ~3.7 ppm in (**a**) and (**b**) are from glycerol; the singlet at ~3.7 ppm in (**a**) and (**b**) corresponds to Tris. Low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed.(5) The broad peaks at ~0.5 to ~1.5 ppm in (**a**) and (**b**) are likely from the PsEFE sample, which was used in a relatively high concentration. Representative spectra from independent duplicates are shown.




Supporting Figure 5.S55. R277K PsEFE does not catalyze oxidation of 4-methyl-2OG (21). (**a-b**) ^{1}H NMR analysis of a reaction mixture containing 4-methyl-2OG (**21**): (**a**) ~10 h and (**b**) ~5 min post addition of R277K PsEFE to the reaction mixture. (**c**) ^{1}H NMR analysis of a reaction mixture containing **21** in the absence of R277K PsEFE. The results reveal R277K PsEFE and, by implication, the other tested R277 PsEFE variants do not employ **21** as a cosubstrate. Conditions: 400 µM **21**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 30 µM R277K PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. The peaks at ~3.6 and ~3.7 ppm in (**a**) and (**b**) are from glycerol; the singlet at ~3.7 ppm in (**a**) and (**b**) corresponds to Tris; low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed.(5) Representative spectra from independent duplicates are shown.




Supporting Figure 5.S56. R277K PsEFE does not catalyze oxidation of 3-methyl-2OG (22). (**a-b**) ^{1}H NMR analysis of a reaction mixture containing 3-methyl-2OG (**22**): (**a**) ~10 h and (**b**) ~5 min post addition of R277K PsEFE to the reaction mixture. (**c**) ^{1}H NMR analysis of a reaction mixture containing **22** in the absence of R277K PsEFE. The results reveal R277K PsEFE and, by implication, the other tested R277 PsEFE variants do not employ **22** as a cosubstrate. Conditions: 400 µM **22**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 30 µM R277K PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. The peaks at ~3.6 and ~3.7 ppm in (**a**) and (**b**) are from glycerol; the singlet at ~3.7 ppm in (**a**) and (**b**) corresponds to Tris; low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed.(5) Representative spectra from independent duplicates are shown.




Supporting Figure 5.S57. R277 PsEFE variants catalyze oxidation of KMBA. (**a**) Reaction outcomes of R277 PsEFE variants with KMBA (400 µM): R277 PsEFE variants catalyzed conversion of KMBA to give ethylene, 3-(methylthio)propionic acid (**1a**), and 2-(methylthio)ethanol (**1b**). wt PsEFE did not catalyze ethylene formation from KMBA under identical conditions. (**b**) Estimated **1a**:**1b** ratios: 2.1 ± 0.2:1 (wt PsEFE), 1.4 ± 0.1:1 (R277A PsEFE), 1.3 ± 0.1:1 (R277F PsEFE), 1.4 ± 0.1:1 (R277K PsEFE), and 1.4 ± 0.1:1 (R277L PsEFE). Conditions: 400 µM KMBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 5 µM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Note that full conversion of KMBA was observed after overnight incubations or when increasing PsEFE concentrations. Data are means of independent triplicates (n = 3; mean ± SD). The corresponding ^{1}H NMR spectra are shown in Supporting Figures 5.S1-5.S5 and 5.S68-5.S70.




Supporting Figure 5.S58. R277 PsEFE variants catalyze oxidation of 4-methyl-2-oxopentanoic acid (2). (**a**) Reaction outcomes of R277 PsEFE variants with **2** (400 µM) after ~90 min incubations: R277 PsEFE variants catalyzed conversion of **2** to give 3-methylbutanoic acid (**2a**) and 2-methylpropan-1-ol (**2b**), in accord with wt PsEFE catalysis. Evidence for formation of an alkene product was not accrued. (**b**) Estimated **2a**:**2b** ratios after ~90 min incubations: 4.8 ± 0.2:1 (wt PsEFE), 3.1 ± 0.1:1 (R277A PsEFE), 3.2 ± 0.1:1 (R277F PsEFE), 3.1 ± 0.1:1 (R277K PsEFE), and 3.3 ± 0.1:1 (R277L PsEFE). Conditions: 400 µM **2**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Data are means of independent triplicates (n = 3; mean ± SD) for wt PsEFE and of independent duplicates (n = 2; mean ± SD) for R277 PsEFE variants. The corresponding ^{1}H NMR spectra are shown in Supporting Figures 5.S6-5.S8.




Supporting Figure 5.S59. R277 PsEFE variants catalyze oxidation of 3-methyl-2-oxopentanoic acid (3). (**a**) Reaction outcomes of R277 PsEFE variants with **3** (400 μM) after ~90 min incubations: R277 PsEFE variants catalyzed conversion of **3** to give 2-methylbutanoic acid (**3a**) and butan-2-ol (**3b**), in accord with wt PsEFE catalysis. Evidence for formation of an alkene product was not accrued. (**b**) Estimated **3a**:**3b** ratios after ~90 min incubations: 2.1 ± 0.1:1 (wt PsEFE), 2.9 ± 0.1:1 (R277A PsEFE), 2.5 ± 0.1:1 (R277F PsEFE), 2.2 ± 0.2:1 (R277K PsEFE), and 2.7 ± 0.2:1 (R277L PsEFE). Conditions: 400 μM **3**, 500 μM *L*-arginine, 500 μM *L*-ascorbic acid, 50 μM Fe(II), 800 μM TMSP-d_4, and 10 μM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Data are means of independent triplicates (n = 3; mean ± SD) for wt PsEFE and of independent duplicates (n = 2; mean ± SD) for R277 PsEFE variants. The corresponding ^{1}H NMR spectra are shown in Supporting Figures 5.S9-5.S11.




Supporting Figure 5.S60. R277 PsEFE variants catalyze oxidation of 3-methyl-2-oxobutanoic acid (4). (**a**) Reaction outcomes of R277 PsEFE variants with **4** (400 µM) after ~90 min incubations: R277 PsEFE variants catalyzed conversion of **4** to give isobutyric acid (**4a**) and isopropanol (**4b**), in accord with wt PsEFE catalysis. Evidence for formation of an alkene product was not accrued. (**b**) Estimated **4a**:**4b** ratios after ~90 min incubations: 2.3 ± 0.1:1 (wt PsEFE), 2.2 ± 0.1:1 (R277A PsEFE), 2.7 ± 0.3:1 (R277F PsEFE), 2.4 ± 0.3:1 (R277K PsEFE), and 2.0 ± 0.4:1 (R277L PsEFE). Conditions: 400 µM **4**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Data are means of independent triplicates (n = 3; mean ± SD) for wt PsEFE and of independent duplicates (n = 2; mean ± SD) for R277 PsEFE variants. The corresponding ^{1}H NMR spectra are shown in Supporting Figures 5.S12-5.S14.




Supporting Figure 5.S61. R277 PsEFE variants catalyze oxidation of 2-oxobutanoic acid (5). (**a**) Reaction outcomes of R277 PsEFE variants with **5** (400 µM) after ~90 min incubations: R277 PsEFE variants catalyzed conversion of **5** to give propanoic acid (**4a**) and ethanol (**4b**) product, in accord with wt PsEFE catalysis. Evidence for formation of an alkene product was not accrued. (**b**) Estimated **5a**:**5b** ratios after ~90 min incubations: 1.8 ± 0.3:1 (wt PsEFE), 2.0 ± 0.2:1 (R277A PsEFE), 2.1 ± 0.1:1 (R277F PsEFE), 1.6 ± 0.1:1 (R277K PsEFE), and 0.9 ± 0.1:1 (R277L PsEFE). Conditions: 400 µM **5**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Data are means of independent triplicates (n = 3; mean ± SD) for wt PsEFE and of independent duplicates (n = 2; mean ± SD) for R277 PsEFE variants. The corresponding ^{1}H NMR spectra are shown in Supporting Figures 5.S15-5.S17.




Supporting Figure 5.S62. R277 PsEFE variants catalyze oxidation of 2-oxovaleric acid (6). (**a**) Reaction outcomes of R277 PsEFE variants with **6** (400 µM) after ~90 min incubations: R277 PsEFE variants catalyzed conversion of **6** to give butanoic acid (**6a**) and propanol (**6b**), in accord with wt PsEFE catalysis. Evidence for formation of an alkene product was not accrued. (**b**) Estimated **6a**:**6b** ratios after ~90 min incubations: 2.4 ± 0.4:1 (wt PsEFE), 1.6 ± 0.1:1 (R277A PsEFE), 1.9 ± 0.5:1 (R277F PsEFE), 1.3 ± 0.1:1 (R277K PsEFE), and 1.9 ± 0.3:1 (R277L PsEFE). Conditions: 400 µM **6**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Data are means of independent triplicates (n = 3; mean ± SD) for wt PsEFE and of independent duplicates (n = 2; mean ± SD) for R277 PsEFE variants. The corresponding ^{1}H NMR spectra are shown in Supporting Figures 5.S18-5.S20.




Supporting Figure 5.S63. R277 PsEFE variants catalyze oxidation of 2-oxocaproic acid (7). (**a**) Reaction outcomes of R277 PsEFE variants with **7** (400 µM) after ~90 min incubations: R277 PsEFE variants catalyzed conversion of **7** to give pentanoic acid (**7a**) and butanol (**7b**), in accord with wt PsEFE catalysis. Evidence for formation of an alkene product was not accrued. (**b**) Estimated **7a**:**7b** ratios after ~90 min incubations: 2.8 ± 0.2:1 (wt PsEFE), 2.5 ± 0.3:1 (R277A PsEFE), 2.2 ± 0.1:1 (R277F PsEFE), 2.2 ± 0.1:1 (R277K PsEFE), and 1.5 ± 0.1:1 (R277L PsEFE). Conditions: 400 µM **7**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Data are means of independent triplicates (n = 3; mean ± SD) for wt PsEFE and of independent duplicates (n = 2; mean ± SD) for R277 PsEFE variants. The corresponding ^{1}H NMR spectra are shown in Supporting Figures 5.S21-5.S23.




Supporting Figure 5.S64. R277 PsEFE variants catalyze oxidation of 2-cyclohexyl-2-oxoacetic acid (8). (**a**) Reaction outcomes of R277 PsEFE variants with **8** (400 μM) after ~90 min incubations: R277 PsEFE variants catalyzed conversion of **8** to give cyclohexanecarboxylic acid (**8a**) and cyclohexanol (**8b**), in accord with wt PsEFE catalysis. Evidence for formation of an alkene product was not accrued. (**b**) Estimated **8a**:**8b** ratios after ~90 min incubations: 1.9 ± 0.2:1 (wt PsEFE), 1.7 ± 0.1:1 (R277A PsEFE), 1.6 ± 0.1:1 (R277F PsEFE), 1.8 ± 0.2:1 (R277K PsEFE), and 1.4 ± 0.1:1 (R277L PsEFE). Conditions: 400 μM **8**, 500 μM *L*-arginine, 500 μM *L*-ascorbic acid, 50 μM Fe(II), 800 μM TMSP-d_4, and 10 μM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Data are means of independent triplicates (n = 3; mean ± SD) for wt PsEFE and of independent duplicates (n = 2; mean ± SD) for R277 PsEFE variants. The corresponding ^{1}H NMR spectra are shown in Supporting Figures 5.S24-5.S26.




Supporting Figure 5.S65. R277 PsEFE variants catalyze oxidation of 3-cyclohexyl-2-oxopropanoic acid (9). (**a**) Reaction outcomes of R277 PsEFE variants with **9** (400 µM) after ~90 min incubations: R277 PsEFE variants catalyzed conversion of **9** to give 2-cyclohexylacetic acid (**9a**) and cyclohexylmethanol (**9b**), in accord with wt PsEFE catalysis. Evidence for formation of an alkene product was not accrued. (**b**) Estimated **9a**:**9b** ratios after ~90 min incubations: 18 ± 0.3:1 (wt PsEFE), 26 ± 1:1 (R277A PsEFE), 19 ± 1:1 (R277F PsEFE), 20 ± 2:1 (R277K PsEFE), and 20 ± 4:1 (R277L PsEFE). Conditions: 400 µM **9**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Data are means of independent triplicates (n = 3; mean ± SD) for wt PsEFE and of independent duplicates (n = 2; mean ± SD) for R277 PsEFE variants. The corresponding ^{1}H NMR spectra are shown in Supporting Figures 5.S27-5.S29.




Supporting Figure 5.S66. R277 PsEFE variants catalyze oxidation of 4-cyclohexyl-2-oxobutanoic acid (10). (**a**) Reaction outcomes of R277 PsEFE variants with **10** (400 µM) after ~90 min incubations: R277 PsEFE variants catalyzed conversion of **10** to give 3-cyclohexylpropionic acid (**10a**) and cyclohexylethan-1-ol (**10b**), in accord with wt PsEFE catalysis. Evidence for formation of an alkene product was not accrued. (**b**) Estimated **10a**:**10b** ratios after ~90 min incubations: 11 ± 1:1 (wt PsEFE), 9.7 ± 1.3:1 (R277F PsEFE), and 4.6 ± 0.7:1 (R277K PsEFE). Conditions: 400 µM **10**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Note that the **10a**:**10b** ratios could not be computed for R277A PsEFE and R277L PsEFE, because the relatively low turnover of **10** prevented efficient alcohol detection. Data are means of independent triplicates (n = 3; mean ± SD) for wt PsEFE and of independent duplicates (n = 2; mean ± SD) for R277 PsEFE variants. The corresponding ^{1}H NMR spectra are shown in Supporting Figures 5.S30-5.S32.




Supporting Figure 5.S67. R277 PsEFE variants catalyze oxidation of 4,4-dimethyl-2-oxopentanoic acid (11). (**a**) Reaction outcomes of R277 PsEFE variants with **11** (400 µM) after ~90 min incubations: R277 PsEFE variants catalyzed conversion of **11** to give 3,3-dimethylbutanoic acid (**11a**) and neopentanol (**11b**), in accord with wt PsEFE catalysis. Evidence for formation of an alkene product was not accrued. (**b**) Estimated **11a**:**11b** ratios after ~90 min incubations: 4.8 ± 0.1:1 (wt PsEFE), 4.9 ± 0.4:1 (R277A PsEFE), 4.4 ± 0.2:1 (R277F PsEFE), 5.2 ± 0.4:1 (R277K PsEFE), and 3.5 ± 0.4:1 (R277L PsEFE). Conditions: 400 µM **11**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Data are means of independent triplicates (n = 3; mean ± SD) for wt PsEFE and of independent duplicates (n = 2; mean ± SD) for R277 PsEFE variants. The corresponding ^{1}H NMR spectra are shown in Supporting Figures 5.S33-5.S35.




Supporting Figure 5.S68. R277 PsEFE variants catalyze oxidation of 2-oxo-4-phenylbutanoic acid (12). (**a**) Reaction outcomes of R277 PsEFE variants with **12** (400 μM) after ~90 min incubations: R277 PsEFE variants catalyzed conversion of **12** to give 3-phenylpropanoic acid (**12a**) and 2-phenylethanol (**12b**), in accord with wt PsEFE catalysis. Evidence for formation of an alkene product was not accrued. (**b**) Estimated **12a**:**12b** ratios after ~90 min incubations: 2.8 ± 1.0:1 (wt PsEFE), 1.0 ± 0.1:1 (R277A PsEFE), 1.1 ± 0.1:1 (R277F PsEFE), 1.7 ± 0.2:1 (R277K PsEFE), and 0.9 ± 0.2:1 (R277L PsEFE). Conditions: 400 μM **12**, 500 μM *L*-arginine, 500 μM *L*-ascorbic acid, 50 μM Fe(II), 800 μM TMSP-d_4, and 10 μM PsEFE in buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O) at 25 °C. Data are means of independent triplicates (n = 3; mean ± SD) for wt PsEFE and of independent duplicates (n = 2; mean ± SD) for R277 PsEFE variants. The corresponding ^{1}H NMR spectra are shown in Supporting Figures 5.S36-5.S38.




Supporting Figure 5.S69. Representative ^{1}H NMR spectra with R277K PsEFE showing that R277 PsEFE variants catalyze ethylene formation from KMBA. (**a**) ^{1}H NMR analysis of a R277K PsEFE reaction with KMBA after ~75 min incubation reveals formation of P5C and ethylene; (**b-c**) ^{1}H NMR spectra of KMBA under standard assay conditions in the absence of PsEFE: (**b**) ~75 min and (**c**) ~5 min after the addition of all assay components. The results indicate low levels of ethylene are formed in a non-enzymatic manner (see below). Conditions: 400 µM KMBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM R277K PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol; the singlet at ~3.7 ppm in (**a**) corresponds to Tris; low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed in (**b**).(5) Peaks corresponding to the hydrated ketone form of KMBA are annotated with dashed circles (Supporting Figure 5.S2). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S70. R277K PsEFE catalyzed time-dependent oxidation of KMBA. (**a**-**c**) Representative ^{1}H NMR spectra (~2.0 to ~3.9 ppm and ~5.4 to ~5.5 ppm) monitoring the reaction of R277K PsEFE with KMBA: (**a**) ~75 min, (**b**) ~50 min, (**c**) ~20 min, and (**d**) ~5 min post addition of R277K PsEFE to the reaction mixture. (**e**) ^{1}H NMR spectrum of the reaction mixture in the absence of R277K PsEFE. Conditions: 400 µM KMBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM R277K PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a**-**d**) are from glycerol; the singlet at ~3.7 ppm in (**a**-**d**) corresponds to Tris. Peaks corresponding to the hydrated ketone form of KMBA are annotated with dashed circles (Supporting Figure 5.S2). Note that R277K PsEFE catalyzed P5C formation in the presence of KMBA (Supporting Figure 5.S68). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S71. Analysis of the reaction of R277K PsEFE with KMBA. ^{1}H NMR analysis (~2.0 to ~3.9 ppm) of: (**a**) a reaction of R277K PsEFE and KMBA in the presence of an authentic sample of methanesulfonic acid (**1c**), (**b**) authentic sample of **1c**; (**c**) a reaction of R277K PsEFE and KMBA in the presence of an authentic sample of 2-(methylthio)ethanol (**1b**), (**d**) authentic sample of **1b**; (**e**) a reaction of R277K PsEFE and KMBA in the presence of an authentic sample of 3-(methylthio)propionic acid (**1a**), (**f**) authentic sample of **1a**; (**g**) a reaction of R277K PsEFE and KMBA ~75 min post addition of R277K PsEFE; (**h**) ^{1}H NMR spectrum of KMBA under standard assay conditions in the absence of R277K PsEFE. Conditions: 400 µM KMBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM R277K PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a**-**g**) are from glycerol; the singlet at ~3.7 ppm in (**a**-**g**) corresponds to Tris. Peaks corresponding to the hydrated ketone form of KMBA are annotated with dashed circles (Supporting Figure 5.S2). Note that R277K PsEFE catalyzed P5C formation in the presence of KMBA (Supporting Figure 5.S68). Representative spectra from independent triplicates are shown.




Supporting Figure 5.S72. R277 PsEFE variants catalyze time-dependent ethylene formation from KMBA. ^{1}H NMR analyses of the reactions of: (**a**) R277A PsEFE, **(b**) R277F PsEFE, (**c**) R277K PsEFE, and (**d**) R277L PsEFE with KMBA to give 3-(methylthio)propionic acid (**1a**), 2-(methylthio)ethanol (**1b**), and ethylene. Conditions: 400 µM KMBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, 10 µM PsEFE variant in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O) at 25˚C. Reaction times are normalized to the end of the first ^{1}H NMR acquisition following PsEFE variant addition to the reaction mixture (t = 0 min), by which time variable levels of turnover were manifest. Results are means of independent triplicates (n = 3; mean ± SD); graphs were plotted using non-linear regression analysis and a single-phase decay model (GraphPad Prism 10.3.0).




Supporting Figure 5.S73. wt PsEFE catalyzes conversion of *L*-arginine to P5C in the presence of KEBA. (**a**) ^{1}H NMR analysis of a reaction of wt PsEFE with KEBA after ~75 min incubation reveals P5C formation; (**b**) ^{1}H NMR spectrum of KEBA under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM KEBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM wt PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**a**) are from glycerol; low levels of dehydroascorbic acid (~4.1 ppm and ~4.3 ppm) were observed in (**b**).(5) the assignment of the peaks corresponding to the hydrated form of KEBA was based on that for the hydrated ketone forms of KMBA (Supporting Figure 5.S2) and 2OG.(6, 7) Representative spectra from independent duplicates are shown.




Supporting Figure 5.S74. ^{1}H NMR analysis of KEBA and authentic samples of the acid (21a) and alcohol (22b) products formed by PsEFE catalyzed oxidation of KEBA. (**a**-**c**) ^{1}H NMR spectra (~0.9 to ~4.0 ppm) of: (**a**) the alcohol product, 2-(ethylthio)ethanol (**21b**), (**b**) the acid product, 3-(ethylthio)propionic acid (**21a**), and (**c**) KEBA in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Peaks corresponding to the hydrated form of KEBA (~2.1 and ~2.5 ppm) are annotated with dashed circles; the assignment of the peaks corresponding to the hydrated form of KEBA was based on that for the hydrated ketone forms of KMBA (Supporting Figure 5.S2) and 2OG.(6, 7)




Supporting Figure 5.S75. Analysis of the reaction of wt PsEFE with KEBA. ^{1}H NMR analysis (~0.9 to ~3.8 ppm) of: (**a**) a reaction of wt PsEFE and KEBA in the presence of an authentic sample of 2-(ethylthio)ethanol (**21b**), (**b**) authentic sample of **21b**; (**c**) a reaction of wt PsEFE and KEBA in the presence of an authentic sample of 3-(ethylthio)propionic acid (**21a**), (**d**) authentic sample of **21a**; (**e**) a reaction of wt PsEFE and KEBA after ~75 min incubation; (**f**) ^{1}H NMR spectrum of KEBA under standard assay conditions in the absence of wt PsEFE. Conditions: 400 µM KEBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM wt PsEFE, and, if appropriate, 2 mM of the appropriate authentic standard in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Peaks corresponding to the hydrated form of KEBA (~2.1 and ~2.5 ppm) are annotated with dashed circles; the assignment of the peaks corresponding to the hydrated form of KEBA was based on that for the hydrated ketone forms of KMBA (Supporting Figure 5.S2) and 2OG.(6, 7) Note that wt PsEFE catalyzed P5C formation in the presence of KEBA (Supporting Figure 5.S72). Representative spectra from independent duplicates are shown.




Supporting Figure 5.S76. R277 PsEFE variants catalyze ethylene formation from KEBA. (**a**) R277 PsEFE variants catalyze conversion of KEBA to give ethylene together with 3-(ethylthio)propionic acid (**21a**) and 2-(ethylthio)ethanol (**21b**). (**b-h**) ^{1}H NMR analysis (~1.0 to ~6.0 ppm and ~7.0 to ~8.0 ppm) of: (**b**) a reaction of R277K PsEFE and KEBA in the presence of an authentic sample of **21b**, (**c**) authentic sample of **21b**; (**d**) a reaction of R277K PsEFE and KEBA in the presence of an authentic sample of **21a**, (**e**) authentic sample of **21a**; (**f**) a reaction of R277K PsEFE and KEBA after ~75 min incubation; (**g**) a reaction of R277K PsEFE and KEBA after ~25 min incubation; (**h**) ^{1}H NMR spectrum of KEBA under standard assay conditions in the absence of R277K PsEFE. (**i**) ^{1}H NMR spectra (~5.42 to ~5.46 ppm) showing the R277K PsEFE-catalyzed conversion of KEBA to ethylene in 5 min increments over 80 min; note that ethylene formation cannot be quantified under the assay conditions. Conditions: 400 µM KEBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and 10 µM R277K PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The peaks at ~3.6 and ~3.7 ppm in (**d**,**f**,**g**) are from glycerol; the singlet at ~3.7 ppm in (**b**,**d**,**f**,**g**) corresponds to Tris. Peaks corresponding to the hydrated ketone form of KEBA (~2.1 and ~2.5 ppm) are annotated with dashed circles; the assignment of the peaks corresponding to the hydrated form of KEBA was based on that for the hydrated ketone forms of KMBA (Supporting Figure 5.S2) and 2OG.(6, 7) Representative spectra from independent duplicates are shown.




Supporting Figure 5.S77. R277K PsEFE catalyzed ethylene formation from KMBA in the presence of catalase. (**a**-**d**) ^{1}H NMR spectra (~2.0 to ~5.5 ppm) monitoring R277K PsEFE-catalyzed ethylene formation from KMBA in the presence of catalase: (**a**) ~75 min, (**b**) ~50 min, (**c**) ~20 min, and (**d**) ~5 min post addition of R277K PsEFE and catalase to the reaction mixture. (**e**) ^{1}H NMR spectrum of KMBA in the absence of PsEFE and catalase. (**f**) ^{1}H NMR spectrum of catalase in buffer. Conditions: 400 µM KMBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, 10 µM PsEFE and 1 µM catalase in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Lyophilized catalase from bovine liver was from Sigma-Aldrich; peaks from catalase and *L*-ornithine, the product of the catalase reaction with *L*-arginine, are annotated by an asterisk (*). The peaks at ~3.6 and ~3.7 ppm in (**a**-**d**) are from glycerol; the singlet at ~3.7 ppm in (**a**-**d**) corresponds to Tris. Peaks corresponding to the hydrated ketone form of KMBA are annotated with dashed circles (Supporting Figure 5.S2). Note that R277K PsEFE catalyzed P5C formation in the presence of KMBA (Supporting Figure 5.S68).




Supporting Figure 5.S78. Ethylene is produced non-enzymatically from KMBA in the presence of ammonium iron (II) sulfate hexahydrate. (**a**-**c**) ^{1}H NMR analysis (~2.0 to ~5.5 ppm) reveals low levels of non-enzymatic production of ethylene from KMBA in the presence of Fe(II): (**a**) ^{1}H NMR spectrum of KMBA in buffer; (**b**) non-enzymatic ethylene formation from KMBA requires Fe(II), but not *L*-arginine and/or *L*-ascorbic acid; (**c**) *L*-ascorbic acid enhances the non-enzymatic ethylene formation in the presence of Fe(II). (**d**) Catalase inhibits non-enzymatic production of KMBA from ethylene in the presence of Fe(II). Notably, the reaction of wt PsEFE with KMBA did not produce substantial amounts of ethylene although identical conditions were employed (Supporting Figure 5.S1), possibly because wt PsEFE reacts relatively fast with KMBA, whereas the non-enzymatic ethylene formation from KMBA may be relatively slower. Conditions: 400 µM KMBA, 800 µM TMSP-d_4 and, if appropriate, 50 µM Fe(II), 500 µM *L*-ascorbic acid, and 1 µM catalase in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Lyophilized catalase from bovine liver was from Sigma-Aldrich; peaks from catalase are annotated by an asterisk (*). Peaks corresponding to the hydrated ketone form of KMBA are annotated with dashed circles (Supporting Figure 5.S2).




Supporting Figure 5.S79. Commercially-sourced catalase catalyzes *L*-ornithine production from *L*-arginine. (**a**) ^{1}H NMR (700 MHz) spectrum (~1.5 to ~3.9 ppm and ~5.4 to ~6.0 ppm) of *L*-ornithine in the presence of catalase. (**b**) ^{1}H NMR spectrum of *L*-ornithine. (**c-d**) ^{1}H NMR analysis of the reaction of catalase with *L*-arginine: (**c**) ~90 min and (**d**) ~5 min post addition of catalase to the mixture. It is possible that the commercial catalase contains trace amounts of arginase.(9) (**e**) ^{1}H NMR spectrum of catalase in buffer. Conditions: 400 µM *L*-arginine/*L*-ornithine, 800 µM TMSP-d_4 and, if appropriate, 1 µM catalase in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Lyophilized catalase from bovine liver was from Sigma-Aldrich; signals originating from catalase are annotated by an asterisk (*).




Supporting Figure 5.S80. Analysis of R277 PsEFE variants using Circular dichroism (CD) spectroscopy. CD spectra of wt and R277 PsEFE variants showing: **a**) raw ellipticity (millidegrees) and **b**) mean residue ellipticity (MRE, Θ) are indicative of folded proteins.(10) No major differences between the CD spectra of the R277 PsEFE variants were observed, suggesting that R277 substitutions do not substantially affect the PsEFE fold; variations in intensities are likely due to differences in protein concentration.(10) Data were normalised to the PsEFE concentration for computing the mean residue ellipticity (MRE) and smoothened using SELCON3(11, 12) (using DichroWeb(13)). Conditions: 0.1 mg/mL wt PsEFE/R277 PsEFE variant in buffer (20 mM sodium phosphate, pH 7.4).



Supporting Figure 5.S81. PsEFE catalysis incorporates a single ^{18}O label in 3-(methylthio)propanoate (1a), the oxidative decarboxylation product of KMBA. (**a**) wt and R277K PsEFE catalyzed turnover of KMBA under $^{18}O_2$ atmosphere produces singly ^{18}O-labeled 3-(methylthio)propanoate (**1a**). The levels of **1a** formed in the reaction are in accord with the results of ^{1}H NMR assays showing complete conversion of KMBA (Supporting Figure 5.S57). (**b**) Stacked bar plots showing the distribution of ^{18}O-labeled and unlabeled **1a** species formed under $^{16}O_2$ and $^{18}O_2$ atmospheres. Unlabeled **1a**: blue; singly ^{18}O-labeled **1a**: orange. (**c**-**d**) Representative extracted mass spectra of **1a** species formed under $^{16}O_2$ (black) and $^{18}O_2$ (red) atmospheres from KMBA with: (**c**) wt PsEFE and (**d**) R277K PsEFE. (**e**-**h**) Extracted ion chromatograms (retention times ~ 6-9 min) of ^{18}O-labeled and unlabeled **1a** species from KMBA turnover, showing overlays of wt PsEFE (green), R277K PsEFE (blue), and no-PsEFE control (red): (**e**) **1a** under $^{16}O_2$, (**f**) ^{18}O-labeled **1a** under $^{16}O_2$, (**g**) **1a** under $^{18}O_2$, and (**h**) ^{18}O-labeled **1a** under $^{18}O_2$. Conditions: 400 μM KMBA, 500 μM *L*-arginine, 500 μM *L*-ascorbic acid, 50 μM Fe(II), and 5 μM wt/R277K PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). The absolute levels of **1a** were quantified using a calibration curve constructed from authentic samples of **1a** under identical reaction conditions (Supporting Figure 5.S83).



Supporting Figure 5.S82. PsEFE-catalysed KMBA turnover produces methanesulfonic acid (MSA) incorporating a full O_2 molecule. (**a**-**f**) Extracted ion chromatograms (retention times ~ 6-9 min) of ^{18}O-labeled and unlabeled MSA species from KMBA turnover, showing overlays of wt PsEFE (green), R277K PsEFE (blue), and no-PsEFE control (red): (**a**) MSA under $^{16}O_2$, (**b**) singly ^{18}O-labeled MSA under $^{16}O_2$, (**c**) MSA under $^{18}O_2$, (**d**) singly ^{18}O-labeled MSA under $^{18}O_2$, (**e**) doubly ^{18}O-labeled MSA under $^{18}O_2$, and (**f**) triply ^{18}O-labeled MSA under $^{18}O_2$. (**g**) Representative extracted mass spectra of MSA species formed in the reaction of KMBA with R277K PsEFE under $^{16}O_2$ (black) and $^{18}O_2$ (red) atmospheres. (**h**) Stacked bar plots showing the distribution of singly-, doubly-, and triply-^{18}O-labeled and unlabeled MSA species formed under $^{16}O_2$ and $^{18}O_2$ atmospheres. Unlabeled MSA: blue; singly ^{18}O-labeled MSA: orange; doubly ^{18}O-labeled MSA: green; triply ^{18}O-labeled MSA: violet. Conditions: 400 µM KMBA, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), and 5 µM wt/R277K PsEFE in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O). Note that, the low levels of MSA produced during R277K PsEFE catalysis limited the absolute quantification of MSA because of non-linearity of the calibration curve at low concentrations.



Supporting Figure 5.S83. Calibration curve of 3-(methylthio)propanoic acid (1a) showing peak intensity as a function of concentration. The calibration demonstrates linearity across the measured range under the reaction conditions ($R^{2.}$ = 0.99). Conditions: 0.05-60 µg/mL **1a**, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, and 50 µM Fe(II), in buffer (50 mM sodium phosphate, pH 7.0, 10%$_{v/v}$ D_2O).

5.11 Supporting Tables

Supporting Table 5.S1. Primer pairs used for site-directed mutagenesis of wt PsEFE to give R277 PsEFE variants. Substitutions are highlighted in bold red. PCR reactions to generate plasmids encoding the R277 PsEFE variants utilized the same reverse primer (R277A PsEFE rev). A standard hot-start PCR protocol was employed for site-directed mutagenesis of PsEFE: an annealing temperature of 64 °C was used to prevent non-specific primer annealing and the elongation time was set to 5 min to ensure amplification of the entire construct (~6.5 kb).

PsEFE variant	Primers pairs (5'-3')		Length (nt)	GC content (%)	Annealing temperature (°C)
R277A	Fwd	AATACCCGTGAA**GCA**TTTGCCTGTGC	26	50.0	74.3
	Rev	CAGTTTAACTTTATGCGGTGTGCTCAGCAG	30	46.7	73.3
R277F	Fwd	GAATACCCGTGAA**TTC**TTTGCCTGTGCC	28	50.0	75.0
R277K	Fwd	AATACCCGTGAA**AAG**TTTGCCTTGCC	27	48.2	73.9
R277L	Fwd	GAATACCCGTGAAC**T**TTTTGCCTGTGCC	28	50.0	75.0

Supporting Table 5.S2. Specific activities of the wt and R277 PsEFE variants with 2OG, KMBA, and other 2-oxoacids accepted as alternative PsEFE (co)substrates.

2-oxoacid	[a] Specific activity [nmol·min^{-1}·mg^{-1}]				
	wt PsEFE	**R277A PsEFE**	**R277F PsEFE**	**R277K PsEFE**	**R277L PsEFE**
2OG	357 ± 26	-	-	-	-
KMBA	163 ± 6	113 ± 14	80 ± 28	184 ± 8	66 ± 9
2	81 ± 3	73 ± 4	62 ± 12	88 ± 2	18 ± 10
3	40 ± 3	20 ± 4	12 ± 5	36 ± 2	13 ± 2
4	23 ± 5	46 ± 2	27 ± 3	34 ± 4	22 ± 3
5	2 ± 2	1 ± 0	1 ± 0	9 ± 3	1 ± 0
6	37 ± 12	23 ± 4	72 ± 13	51 ± 1	16 ± 5
7	74 ± 2	32 ± 4	67 ±4	72 ± 2	34 ± 1
8	84 ± 1	95 ± 1	95 ± 1	81 ± 1	74 ± 4
9	84 ± 3	76 ± 2	77 ± 2	78 ± 4	28 ± 1
10	52 ± 10	19 ± 3	40 ± 2	13 ± 4	13 ± 2
11	11 ± 5	25 ± 6	12 ± 5	21 ± 6	13 ± 2
12	41 ± 4	14 ± 8	17 ± 1	27 ± 1	7 ± 2

a) Specific activities (nmol·min^{-1}·mg^{-1}) based on 2OG/2-oxoacid depletion after ~10 min incubations with PsEFE. ^{1}H NMR assay conditions: 400 µM 2OG or 2OG derivative, 500 µM *L*-arginine, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TMSP-d_4, and PsEFE (using 2 µM wt PsEFE with 2OG; 5 µM wt PsEFE or 10 µM R277 PsEFE variant with KMBA; 10 µM wt PsEFE or R277 PsEFE variant with **2-12**) in buffer (50 mM sodium phosphate, pH 7.0, 10% $_{v/v}$ D_2O).


5.12 Supporting Information References

1. Adams, R. W., Holroyd, C. M., Aguilar, J. A., Nilsson, M., and Morris, G. A. (2013) "Perfecting" WATERGATE: clean proton NMR spectra from aqueous solution. *Chem. Commun.* **49**, 358–360
2. Pohl, L., and Eckle, M. (1969) Sodium 3-trimethylsilyltetradeuteriopropionate, a new water-soluble standard for ^{1}H-NMR. *Angew. Chem. Int. Ed.* **8**, 381
3. Ducho, C., Hamed, R. B., Batchelar, E. T., Sorensen, J. L., Odell, B., and Schofield, C. J. (2009) Synthesis of regio- and stereoselectively deuterium-labelled derivatives of l-glutamate semialdehyde for studies on carbapenem biosynthesis. *Org. Biomol. Chem.* **7**, 2770–2779
4. Hamed, R. B., Henry, L., Gomez-Castellanos, J. R., Mecinović, J., Ducho, C., Sorensen, J. L., *et al.* (2012) Crotonase catalysis enables flexible production of functionalized prolines and carbapenams. *J. Am. Chem. Soc.* **134**, 471–479
5. Dhingra, S., Zhang, Z., Lohans, C. T., Brewitz, L., and Schofield, C. J. (2024) Substitution of 2-oxoglutarate alters reaction outcomes of the Pseudomonas savastanoi ethylene-forming enzyme. *J. Biol. Chem.* **300**, 107546
6. Copper, A. J., and Redfield, A. G. (1975) Proton magnetic resonance studies of alpha-keto acids. *J. Biol. Chem.* **250**, 527–532
7. Viswanathan, T. S., Johnson, R. E., and Fisher, H. F. (1982) α-Ketoglutaric acid: solution structure and the active form for reductive amination by bovine liver glutamate dehydrogenase. *Biochemistry* **21**, 339–345
8. Copeland, R. A., Zhou, S., Schaperdoth, I., Shoda, T. K. C., Bollinger, J. M., and Krebs, C. (2021) Hybrid radical-polar pathway for excision of ethylene from 2-oxoglutarate by an iron oxygenase. *Science* **373**, 1489–1493
9. Tian, Y., Xing, Y., Magliozzo, R., Yu, K., Bloom, B. R., and Chan, J. (2000) A Commercial preparation of catalase inhibits nitric oxide production by activated murine macrophages: role of arginase. *Infect. Immun.* **68**, 3015–3018
10. Greenfield, N. J. (2006) Using circular dichroism spectra to estimate protein secondary structure. *Nat. Protoc.* **1**, 2876–2890
11. Sreerama, N., and Woody, R. W. (1993) A self-consistent method for the analysis of protein secondary structure from circular dichroism. *Anal. Biochem.* **209**, 32–44
12. Sreerama, N., Venyaminov, S. Y. U., and Woody, R. W. (1999) Estimation of the number of α-helical and β-strand segments in proteins using circular dichroism spectroscopy. *Protein Sci.* **8**, 370–380
13. Miles, A. J., Ramalli, S. G., and Wallace, B. A. (2022) DichroWeb, a website for calculating protein secondary structure from circular dichroism spectroscopic data. *Protein Sci.* **31**, 37–46

5.13 Statement of Authorship

Title of Paper	**Oxygenase-catalyzed production of ethylene from the methionine transamination product α-keto-γ-methylthiobutyric acid.**
Publication Status	□ Published □ Accepted for Publication □ Submitted for Publication X Unpublished and unsubmitted work written in a manuscript style
Publication Details	Dhingra, S.; Harrington, L; Sun, Y.; Legge, I.; Zhang, Z.; McCullagh, J. S.; Schofield, C. J.; Brewitz, L. Oxygenase-catalyzed production of ethylene from the methionine transamination product α - keto - γ - methylthiobutyric acid. **Manuscript in preparation.** Written in Nat. Commun. style.

Student Confirmation

Student Name:	Siddhant Dhingra		
Contribution to the Paper	The experimental work in this manuscript was performed by myself and Mr. Luke Harrington, under my laboratory supervision; I cloned and purified R277A/K PsEFE, Mr. L. Harrington cloned and purified R277F/L PsEFE, both of us performed the NMR assays. Mr. Yihong Sun performed initial assays for exploring non-enzymatic ethylene formation. Prof. James S. McCullagh and Ms. Isabel Legge assisted with MS-characterisation and analysis of the KMBA reaction. I conducted all the data analysis and data visualisation described here. The manuscript was prepared by Dr. Lennart Brewitz and myself under the guidance of Prof. Christopher J. Schofield. Dr. Zhihong Zhang provided the wt PsEFE plasmid and the initial training for protein expression and purification. Dr. Lennart Brewitz, Prof. Christopher J. Schofield, and I jointly conceptualised the project.		
Signature 		Date	05-10-2025

Supervisor Confirmation

By signing the Statement of Authorship, you are certifying that the candidate made a substantial contribution to the publication, and that the description described above is accurate.

Supervisor name and title: Prof. Christopher J. Schofield		
Supervisor comments I'm happy to state that the candidate made the major and a very substantial contribution to the publication, and that the description described above is accurate.		
Signature C. J. Schofield	Date	25th September 2025

CHAPTER 6

MANUSCRIPT V

Crystallographic and biochemical characterisation of HIS1 and OsHSL1 inform on strategies to evade resistance-associated β-triketone herbicide metabolism

Contents

6.1 Overview 378
6.2 Abstract 379
6.3 Introduction 379
6.4 Results 382
6.4.1 ^{1}H NMR assays are suitable for monitoring HIS1 catalysis 382
6.4.2 Characterisation of the HIS1 reaction products. 385
6.4.3 The substrate scopes of HIS1 and OsHSL1 differ. 386
6.4.4 Crystallography. 389
6.4.5 HIS1 and OsHSL1 crystal structures inform on potential substrate binding sites. 393
6.5 Discussion 396
6.6 Experimental Procedures 400
6.7 Acknowledgements 403
6.8 Author Contributions 403
6.9 References 404
6.10 Supporting Figures 408
6.11 Supporting Tables 430
6.12 Supporting Information References 433
6.13 Statement of Authorship 435

6.1 Overview

Herbicide resistance in crops is a major challenge in agriculture. Enzyme-mediated metabolic resistance is a key mechanism by which plants evade herbicidal action. The rice enzyme, HPPD-Inhibitor Sensitive I, HIS1, a 2-oxoglutarate (2OG)-dependent oxygenase, has been identified as conferring resistance to β-triketone herbicides. Related oxygenases such as *Oryza sativa* HIS1-like enzyme, OsHSL1, share high sequence similarity with HIS1 but differ in their substrate selectivity, providing an opportunity to explore the structural and biochemical basis of xenobiotic metabolism by plant 2OG oxygenases.

This chapter reports structural and biochemical characterisation of HIS1 and OsHSL1. Structural analyses revealed highly similar features of typical 2OG-dependent oxygenases, yet differences in active-site geometry and bound ligands support distinct substrate preferences. Functional studies demonstrate that HIS1 efficiently catalyses oxidation of β-triketone herbicides, while OsHSL1 displays narrower selectivity. Multiple crystallographic studies inform on structural differences between HIS1 and OsHSL1. These results advance the understanding of herbicide metabolism by plant 2OG oxygenases and highlight their potential for use in engineering of transgenic crops and/or potent inhibitors/herbicides.

At the time of writing this thesis, Manuscript V is being prepared for submission and is written in the style of a *Journal of Biological Chemistry* paper.

Dhingra, S.; Sharma, S.; Farcas, I.M.; Alshref, F. M.; Allen, M. D.; Redfield, C.; Schofield, C. J.; Brewitz, L. Crystallographic and biochemical characterisation of HIS1 and OsHSL1 inform on strategies to evade resistance-associated β-triketone herbicide metabolism. **Manuscript in preparation.**

Each author's contribution is described in the Author Contributions (Section 6.8). A Statement of Authorship is appended at the end of the chapter (Section 6.13).

6.2 Abstract

4-Hydroxyphenylpyruvate dioxygenase (HPPD) is an essential enzyme in tyrosine metabolism and a validated agrochemical target. β-Triketone herbicides (βTHs) are potent HPPD inhibitors which are widely used for weed-control *e.g.*, in rice farming. The Fe(II) and 2-oxoglutarate (2OG)-dependent oxygenase HPPD-inhibitor sensitive 1 (HIS1) catalyses hydroxylation of a broad spectrum of βTHs, conferring resistance. By contrast, the HIS1 ortholog *Oryza sativa* HIS1-like enzyme 1 (OsHSL1) is reported to have a narrower substrate scope than HIS1 with which it shares ~87% sequence identity. Here we report ^{1}H NMR assays suitable for quantification of HIS1/OsHSL1 catalysis and product characterisation. The results indicate that both HIS1 and OsHSL1 catalyse βTH oxidation at the cyclohexanedione C5 position and confirm the relatively larger HIS1 substrate scope compared to that of OsHSL1. Importantly, the βTH fenquinotrione appears to be a less efficient HIS1 substrate than other βTHs. High-resolution crystal structures of both full-length HIS1 and OsHSL1 each in complex with 2OG and Mn(II) inform on active-site differences that could contribute to the relatively reduced OsHSL1 substrate scope. The combined results improve understanding of herbicide metabolism in crops and inform on strategies for development of improved βTHs to evade HIS1/OsHSL1 metabolism.

6.3 Introduction

Weeds compromise crop productivity worldwide because they compete with crops for water, light, and nutrients, which can result in reduced crop yields.[1,2] Several classes of widely used small-molecule herbicides have been developed to control weed growth,[3-10] including the β-triketone herbicides (βTHs) which comprise mesotrione (**1**),[10-12] sulcotrione (**2**),[10,12] tembotrione (**3**),[10,13] tefuryltrione (**4**),[3,10] nitisinone (**5**),[10,12] fenquinotrione (**6**),[10,14] and benzobicyclon (BBC)[10,15] (Figure 6.1). βTHs work by inhibiting catalysis of 4-

hydroxyphenylpyruvic acid dioxygenase (HPPD), a non-haem Fe(II)-dependent oxygenase conserved across several kingdoms of life.[16-19]

HPPD is an essential enzyme in eukaryotic tyrosine catabolism, catalysing oxidation of the tyrosine transamination product 4-hydroxyphenylpyruvic acid (HPPA) to give homogentisic acid (HGA).[16-19] In plants, HPPD is essential in part because HGA serves as a precursor for the biosynthesis of both plastoquinone and tocopherol (vitamin E).[19] Plastoquinone is a cofactor for carotenoid biosynthesis and mediates protection of chlorophyll in plants,[19,20] and tocopherol provides protection against oxidative stress and regulates phytohormonal balance.[21] Inhibition of HPPD depletes plastoquinone and tocopherol, resulting in chlorosis, disruption of photosynthesis, and, ultimately, plant death (Figure 6.1).[16-18,22,23]

βTHs and other classes of HPPD inhibitors (Figure 6.1), including isoxazoles and pyrazoles,[24,25] are widely employed in agriculture, including during rice and maize farming.[3-8] However, their efficacy is threatened by the emergence of βTH-resistant weeds, as also observed for herbicides targeting enzymes other than HPPD.[26]

Herbicide metabolism is a common resistance mechanism in plants; it is mediated by different classes of enzymes, including cytochrome P450 monooxygenases, glutathione-*S*-transferases, and non-haem Fe(II)/2OG-dependent oxygenases.[27,28] The 2OG oxygenase superfamily catalyses 2-electron oxidation reactions of DNA, RNA, lipids, proteins, and small-molecules, coupled to the oxidative decarboxylation of 2OG to give succinic acid.[29] 2OG oxygenases share a conserved double-stranded β-helix (DSBH) core fold which typically bears a HXD/E…H triad of ligands to bind Fe(II), that serves as a cofactor for the oxidation reaction.[29]




Figure 6.1. HIS1 and HIS1-like enzymes confer βTH resistance. (**a**) HPPD catalysis.[16-19] (**b**) Reported classes of HPPD inhibitors[24,25]: β-triketones (blue), isoxazoles (pink), and pyrazolones (green). (**c**) *Oryza sativa* HIS1, and HIS1-like enzymes (HSLs), confer resistance to βTHs by catalysing βTH metabolism of *e.g.*, benzobicyclon-OH (BBC-OH) and mesotrione (**1**).[3-7] (**d**) A reported ZmHSL1A:Mn(II):2OG:**1** complex crystal structure (PDB ID: 8X74[8]) informs on binding of **1** at the active site.[8]

Pioneering studies identified 2OG oxygenases in *Oryza sativa* (rice) that catalyse oxidation of βTHs, *i.e.*, HPPD Inhibitor Sensitive 1 (HIS1)[6] and six phylogenetically related HIS1-like enzymes (HSLs) (Supporting Figure 6.S1-6.S2).[3-7] Related HSLs have subsequently been identified by sequence similarity searches in species other than rice, including, wheat (*Triticum aestivum*), barley (*Hordeum vulgare*), sorghum (*Sorghum bicolor*) and maize (*Zea mays*) (Supporting Figure 6.S1-6.S2).[3-7]

Structural and functional studies have shown that isolated HIS1 and HSLs catalyse hydroxylation of βTHs at the C5 position of the cyclohexanedione ring,[9] producing metabolites

(Figure 6.1c) with markedly reduced HPPD inhibitory activity.[3-9] Notably, HIS1 catalyses hydroxylation of a broad range of βTHs,[3-9] whereas the substrate scope (and efficiency) of OsHSL1 is substantially narrower than that of HIS1,[3-7] possibly reflecting differences in active-site architecture, as implied by mutation studies on OsHSL1 active-site residues.[8] Crystal structures of HIS1 in its apo-form have been reported,[9,30] however, at relatively poor resolution (3.7 Å and 2.4 Å) using truncated instead of full-length protein and/or HIS1 variants.[9] HSL enzymes have also been crystallographically characterised, including *e.g.*, ZmHSL1A, for which a crystal structure in complex with mesotrione (**1**) is reported (Figure 6.1d).[8]

Here we report ^{1}H NMR-based assays of isolated HIS1 and OsHSL1 enabling investigations on the substrate scopes and catalytic efficiencies. The results revealed that the βTH fenquinotrione (**6**) is not an efficient substrate of either HIS1 or OsHSL1, informing on strategies to design improved HPPD inhibitors. High resolution crystal structure of both isolated full-length HIS1 and OsHSL1 inform on the molecular basis for the observed differences in substrate scopes and on their active site geometry, including metal ion and 2OG binding.

6.4 Results

6.4.1 ^{1}H NMR assays are suitable for monitoring HIS1 catalysis

Reported HIS1 and OsHSL1 assays employ HPLC coupled to mass spectrometry (MS)-based assays,[6,8,9,30-32] while the reaction product of HIS1 with mesotrione (**1**) has been characterised using ^{1}H NMR.[9] We envisioned ^{1}H NMR-based assays could be used to directly monitor the reaction of HIS1/OsHSL1 with βTHs and related compounds, because they enable quantification of substrate depletion and product formation using 3-(trimethylsilyl)-2,2,3,3-tetradeuteropropionic acid (TMSP-d_4) as an internal standard and because they are suitable for product characterisation.[33]

Full-length isolated HIS1 and OsHSL1 were recombinantly produced in *E. coli* cells and purified (>90% by SDS-PAGE and MS analyses; Supporting Figure 6.S3). Conditions previously used in ^{1}H NMR turnover assays of 2OG oxygenases other than HIS1/OsHSL1[33] served as a starting point for optimisation using the reported HIS1 substrate mesotrione (**1**).

The ^{1}H NMR assay results reveal that HIS1 was more active under mildly acidic conditions (pH 6.0) than basic conditions (pH 8.0), and that it required *L*-ascorbate, Fe(II), and 2OG for catalysis (Supporting Figure 6.S4). The optimised reaction conditions are: HIS1 or OsHSL1 (20 μM), substrate (400 μM), 2OG (800 μM), Fe(II) (50 μM), and *L*-ascorbic acid (500 μM) in 50 mM sodium phosphate buffer (pH 6.0, 10%$_{v/v}$ D_2O) at 25 °C. Notably, in the absence of a βTH substrate, HIS1 and OsHSL1 catalysed oxidation of ~30% of 2OG to succinic acid within 90 min uncoupled from substrate oxidation. Under the optimised conditions, HIS1 catalysed ~50% depletion of mesotrione (**1**) within 30 min with initial formation of a single product which was slowly converted into a second product (Figure 6.2).




Figure 6.2. HIS1 catalyses hydroxylation of mesotrione (1) in the presence of 2OG. (**a**) HIS1 catalyses C5 hydroxylation of **1** in the presence of 2OG to give 5-hydroxymesotrione (**1a**); small peaks suggesting the formation of a second product **1b** are observed after ~6 h incubation of **1** with HIS1 in an open Eppendorf. Note that 100% conversion of **1** was observed in the open Eppendorf reaction. (**b**) The relative orientation of the hydroxyl group was assigned as equatorial based on the analysis of coupling constants. The C5 proton of **1a** was not observed, likely because it coincides with the HOD peak; however, the splitting pattern of the C4/C6 proton signals implies that these protons couple with the C5 proton as observed for other βTHs (Supporting Figures 6.S5 and 6.S8). (**c-e**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of HIS1 and **1**: (**c**) ~6 h, (**d**) ~90 min, and (**e**) ~5 min post addition of HIS1. (**f**) ^{1}H NMR spectrum of a reaction mixture in the absence of HIS1. (**g**) ^{1}H NMR spectrum of an authentic sample of **1** in buffer. Conditions: 400 µM **1**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM HIS1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. For *J*-values of **1a** see: Supporting Table 6.S2. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed squares).[34-37] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Both **1** and **1a** were assigned as the

C2 *exo*-enol tautomer as implied by a crystal structure of **1** and ^{1}H NMR analysis of **1** and **5**;[8,38-40] note, however, that the corresponding C2 *endo*-enol tautomer may also form.[9] The second reaction product **1b** was assigned by comparison with structurally related 2-acylphloroglucinol derivatives showing a characteristic singlet for C4/C6-protons at ~5.5 to ~6.5 ppm.[41-43] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.

6.4.2 Characterisation of the HIS1 reaction products.

^{1}H NMR was employed to characterise product(s) of the HIS1 reaction with βTHs. The results reveal that HIS1 catalyses C5-hydroxylation of the βTH cyclohexane-1,3-diketone group (Figure 6.2; Table 6.1; Supporting Table 6.S2), in accord with the reported assignment of the mesotrione-derived reaction product.[9]

Oxidation at the C5 position of the cyclohexyl ring of **1** gave a symmetrically substituted cyclohexane-1,3-diketone with the four C4 and C6 protons showing peaks at ~2.65 ppm (J = 16.7, 4.0 Hz) and ~2.40 ppm (J = 16.6, 7.0 Hz) due to conformational bias imposed by the cyclohexane ring (Figure 6.2; Supporting Table 6.S2).[44]

The relatively large coupling constants of ~16.5 Hz are characteristic of geminal couplings in methylene groups,[45] while the vicinal coupling of ~7.0 Hz indicates a *trans*-diaxial coupling of the C4/C6 methylene protons with the C5 proton; the relatively smaller coupling constant of ~4.0 Hz indicates an axial-equatorial coupling between the C5 proton and the C4/C6 methylene protons.[45] The combined analysis of the vicinal coupling constants thus informs on the relative orientation of the hydroxyl group in the mesotrione-derived product (**1a**),[45] suggesting that the C5 proton adopts an axial and the C5 hydroxyl group an equatorial orientation. Similar coupling patterns were observed for the other tested βTHs (Supporting Table 6.S2).

After ~ 20 min incubation of HIS1 with mesotrione (**1**), signals corresponding to a new species were observed: the new species showed a characteristic singlet at ~5.6 ppm, but lacked the C4/C6 multiplets of **1a**, an observation consistent with the formation of a second product during HIS1 catalysis in the reported MS-based assays (Figure 6.2).[8,9,32] The low abundance and

extensive peak broadening of the second product complicated its characterisation using ^{1}H NMR; the peaks corresponding to the second product were assigned by comparison with reported ^{1}H NMR data of structurally related 2-acylphloroglucinol derivatives having the characteristic peak for C4/C6-protons at ~5.5-6.5 ppm.[41-43] Thus, **1b** was tentatively assigned as the acylphloroglucinol derivative of mesotrione (**1**). Note that peaks corresponding to a second product of **1** were not observed in reported ^{1}H NMR studies.[9] No evidence for the formation of other products was accrued under the assay conditions.

Reported NMR studies on βTHs,[39] including mesotrione (**1**) and nitisinone (**5**), small molecule crystal structure of the sodium salt of **1**,[40] and crystal structures of HSLs in complex with βTHs,[8,38] suggest that the βTHs are predominantly present as the enol tautomer. Accordingly, our ^{1}H NMR spectra did not show peaks corresponding to the C2 proton of the triketone form of βTHs (Figure 6.2, Supporting Figures 6.S5-6.S13). Note that, however, βTHs, may interconvert between different keto-enol forms in a substrate, time, and/or condition-dependent manner.[9,39,40]

6.4.3 The substrate scopes of HIS1 and OsHSL1 differ.

The optimised ^{1}H NMR assays were employed to investigate the substrate scope of HIS1 and OsHSL1. Reported HPPD-targeting agrochemicals were tested as HIS1 substrates, including βTHs[24,25] (*i.e.*, mesotrione (**1**)[10-12], sulcotrione (**2**)[10,12], tembotrione (**3**)[10,13], tefuryltrione (**4**)[3,10], nitisinone (**5**)[10,12], fenquinotrione (**6**)[10,14]), the isoxazole isoxaflutole (**7**),[24,25] and the pyrazole topramezone (**8**)[24,25] (Figure 6.1). The plant growth regulators prohexadione (**9**)[46-48] and trinexapac-ethyl (**10**)[48,49] were also tested for activity (Figure 6.3), because of their structural similarities to βTHs. Note that **9** and **10** lack an aryl group capable of engaging in π,π-stacking interactions, a key feature associated with the interaction of βTHs with HPPD.[24,25]

The results reveal HIS1-catalysed substrate depletion of **1-6** and **10** with formation of the corresponding C5 hydroxylated products (Figure 6.3, Supporting Figures 6.S5-6.S13), consistent with reported results using MS-based assays;[6,8,9,30-32] note the reaction of HIS1 with fenquinotrione (**6**) has, to our knowledge, not previously been reported. Analyses of specific activities and substrate turnover profiles reveals relatively lower activity of HIS1 with βTHs **3**, **4**, and, in particular, **6** compared to **1**, **2**, and **5**, suggesting that substitutions at the C3 position of the mesotrione aryl ring reduce HIS1 activity (Figure 6.3, Table 6.1). Relatively low levels of the corresponding second products of **1**-**5** were observed ~6 h post incubation with HIS1, in accord with the literature (Supporting Figures 6.S5-6.S13).[6,8,9,30-32] Interestingly, HIS1-catalysed production of the second product was more efficient with **1-4** than with **5** (Figure 6.3), whereas HIS1 did not catalyse formation of any product other than **6a** from **6** under the assay conditions, within limits of detection (Figure 6.3; Supporting Figure 6.S9). Note that the relatively low abundance and/or peak broadening may perturb quantification of the second product at earlier time points of the ^{1}H NMR assays.

By contrast to **1-6** and **10**, HIS1 did not catalyse oxidation of **7** and **8**, implying that the triketone functionality of the βTHs is required for efficient HIS1 catalysis (Supporting Figures 6.S10-6.S11, Figure 6.3). Consistent with this proposal, and reported studies,[32] HIS1 catalysed hydroxylation of triketone **10**, although it bears a carboxyethyl ester substituent at the C5 position (Figure 6.3). By contrast, HIS1 did not catalyse hydroxylation of triketone **9** within limits of detection (Supporting Figures 6.S12), consistent with reported studies proposing that HIS1 is inhibited by **9**, despite being structurally related to **10** (Figure 6.3).[32] In the presence of **9**, the HIS1-catalysed oxidation of 2OG to succinic acid uncoupled from triketone oxidation was substantially lower than for the other triketones tested, possibly reflecting the reported inhibitory effect of **9** (Figure 6.3j).[32]




Figure 6.3. HIS1 catalyses oxidation of most but not all commercially used HPPD inhibitors. Reported HPPD inhibitors[24,25] (**1-8**) and structurally similar triketones (**9-10**) were tested as HIS1 substrates. (**a**) HIS1-catalysed reactions of triketones bearing a cyclohexanedione moiety, including βTHs. (**b-k**) ^{1}H NMR analyses (600 MHz) of the HIS1-catalysed reaction of: (**b**) mesotrione (**1**), (**c**) sulcotrione (**2**), (**d**) tembotrione (**3**), (**e**) tefuryltrione (**4**), (**f**) nitisinone (**5**), (**g**) fenquinotrione (**6**), (**h**) isoxaflutole (**7**), (**i**) topramezone (**8**), (**j**) prohexadione (**9**), and (**k**) trinexapac ethyl (**10**). (**l**) ^{1}H NMR analyses of the reactions of HIS1 with **1-10** after ~6 h incubation. Substrate depletion for **4** could not be directly quantified due to overlapping peaks of **4** and **4a**. Conditions: 400 µM substrate, 800 µM 2OG,

500 μM *L*-ascorbic acid, 50 μM Fe(II), 800 μM TSP-d_4, and 20 μM HIS1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Reaction times are normalized to the end of the first ^{1}H NMR acquisition following HIS1 addition to the reaction mixture (t = 0 min), by which time variable levels of conversion were manifest. Results are means of independent triplicates (n = 3; mean ± standard deviation, SD) for time-course profiles and duplicates for open-Eppendorf reactions (n = 2; mean ± SD). Representative ^{1}H NMR spectra are shown in Figure 6.2 and Supporting Figures 6.S5-6.S13.

^{1}H NMR assays with OsHSL1 reveal that OsHSL1 only employed βTH **4** as a substrate, but, unlike HIS1, not any of the other tested βTHs (Supporting Figures 6.S14-6.S19), consistent with reported results from assays with transgenic plants.[32] However, this observation is in contrast with studies on the substrate scope of isolated OsHSL1 showing that mesotrione (**1**) and topramezone (**8**) were substrates.[8] This difference might reflect the different reaction conditions employed in the ^{1}H NMR and MS-based assays and/or the intrinsic detection limitations of the ^{1}H NMR-based assays. Note that analysis of 2OG depletion revealed relatively constant levels of OsHSL1-catalysed oxidation of 2OG to succinic acid uncoupled from triketone oxidation (Supporting Figures 6.S15-6.S19).

6.4.4 Crystallography.

HIS1 and OsHSL1 were crystallised in the presence of 2OG and Mn(II), substituting for catalytically active Fe(II) to inform on the differences in substrate scope. HIS1 crystallised in the C 1 2 1 space group (1.75 Å resolution; Figure 6.5 and Supporting Table 6.S1) and OsHSL1 in the P 21 21 21 space group (2.10 Å; Figure 6.5 and Supporting Table 6.S1). By contrast, reported HIS1 crystal structures were in the I 2 2 2[30] and P41[9] space groups and a reported OsHSL1 variant crystal structure in the C 1 2 1 space group[8]. The HIS1:Mn:2OG and OsHSL1:Mn:2OG complex crystal structures were solved by molecular replacement using an AlphaFold[50] model structure of HIS1 from *Oryza sativa* as a search model (PDB accession code: AF_AFQ6ES21F1).

The results reveal the characteristic distorted DSBH core fold for both HIS1 and OsHSL1, with the overall fold of HIS1 and OsHSL1 being highly similar (RMSD: 0.55 Å), consistent with a sequence identity of ~87% (Figure 6.4). The HIS1 and OsHSL1 folds comprise 9 α-helices and 14 β-strands, eight of which *i.e.*, β7- β14**,** constitute the conserved distorted DSBH core fold (Supporting Figure 6.S20). The *N*-terminal region, having six α-helices (α1-α6), forms a belt around the DSBH core, stabilises the catalytic pocket. The *C*-terminal α-helices, α8 and α9, form a lid near the active-site bearing the Fe(II) binding residues (*vide infra*) (Supporting Figure 6.S20). The HIS1 fold in the HIS1:Mn:2OG structure is similar to that in the reported HIS1 apo[30] and HIS1:Fe[9] crystal structures (RMSD = 0.58 and 1.16 Å, respectively) (Figure 6.4d and e).

A defining feature of the IPNS/DAOCS 2OG oxygenase subfamilies,[51] the insert between strands IV (β10) and V (β11) of the DSBH core is constituted of a short turn (loop) in both HIS1 and OsHSL1. Structural comparison of HIS1 and OsHSL1 using the DALI server[52] reveals similarities with 2OG oxygenases of the IPNS/DAOCS subfamily,[51] including 1-aminocyclopropane-1-carboxylic acid oxidase (ACCO) and Isopenicillin *N* synthase (IPNS), which both do not employ 2OG as a co-substrate.[51]

Analysis of electron density maps for the HIS1:Mn:2OG (PDB ID: 8S7B; 1.75 Å) and OsHSL1:Mn:2OG (PDB ID: 9GI5; 2.35 Å) structures reveals that both 2OG and Mn were bound at the active site (Figure 6.4). Interestingly, a polyethylene glycol-8 (PEG8) molecule from the precipitant co-crystallised in complex with HIS1 (*vide infra*). Comparison of our HIS1:Mn:2OG structure with the reported HIS1 apo structure (PDB ID: 8Y4U; 2.40 Å resolution; RMSD = 0.58 Å)[30] and HIS1:Fe(III) complex structure (PDB ID: 8ETN; 3.16 Å resolution; RMSD = 1.16 Å)[9] reveals similar overall folds, but different conformations of the side chains of residues directly involved in metal ion and/or 2OG binding (Figure 6.4).

The electron density maps reveal that Mn(II) is complexed by an HXD…H triad of ligands in both the HIS1:Mn:2OG and OsHSL1:Mn:2OG structures, *i.e.*, H226 (2.1 Å), H283 (2.2 Å), and D228 (2.2 Å) for HIS1 and H225 (2.3 Å), H282 (2.3 Å), and D227 (2.2 Å) for OsHSL1 (Figure 6.4). 2OG chelated to Mn(II) via its C1 carboxylate and C2 keto groups (2.1 and 2.1 Å for HIS1; 2.3 and 2.2 Å for OsHSL1). A water molecule bound to Mn(II), completing its octahedral coordination sphere (2.3 Å for HIS1; 2.7 Å for OsHSL1). The 2OG C5 carboxylate group was positioned to interact with the guanidinium group of R293 of HIS1 (2.9 and 2.8 Å) and R292 of OsHSL1 (2.8 and 2.8 Å), respectively. A tyrosine residue was also positioned to interact with the 2OG C5 carboxylate (Y211 for HIS1: 2.6 Å; Y210 for OsHSL1: 2.6 Å).




Figure 6.4. HIS1 and OsHSL1 share a similar overall fold and Fe(II) binding site. (**a-b**) Representative Polder omit[53] electron density maps (green mesh) contoured to 5σ around 2OG in: (**a**) the HIS1:Mn:2OG structure (PDB ID: 8S7B) and (**b**) the OsHSL1:Mn:2OG structure (PDB ID: 9GI5) reveal similar 2OG and metal ion binding sites. (**c**) Superimposition of overall and active site views of the HIS1:Mn:2OG (PDB ID: 8S7B) and OsHSL1:Mn:2OG (PDB ID: 9GI5) structures reveal similar overall folds and active site geometries (RMSD = 0.55 Å). Note that a PEG8 molecule crystallised at the HIS1 active site. (**d** and **e**) Superimposition of overall and active site views of the HIS1:Mn:2OG (PDB ID: 8S7B) structure and: (**d**) the reported HIS1 apo (PDB ID: 8Y4U[30]) structure (RMSD = 0.58 Å), and (**e**) the reported HIS1:Fe (PDB ID: 8ETN[9]) structures (RMSD = 1.16 Å) revealing different conformations of residues directly involved in metal ion and/or 2OG binding. HIS1: pink; OsHSL1: lavender; DSBH core fold: cyan; Mn: magenta spheres; 2OG: green. Representative Polder omit[53] electron density maps contoured to 5σ around 2OG are shown as green mesh.

6.4.5 HIS1 and OsHSL1 crystal structures inform on potential substrate binding sites.

Superimposition of the HIS1:Mn:2OG and OsHSL1:Mn:2OG structures reveals conserved active site geometries, including residues forming the HXD…H triad of Fe(II) ligands and residues interacting with the 2OG C5 carboxylate (Figure 6.4). Although HIS1/OsHSL1 structures in complex with a βTH substrate were not obtained, the PEG8 molecule, which co-crystallised with HIS1, potentially informs on the substrate binding site (Figure 6.5a). The side chains of R139, H141, R143, F205, L299, F336, and F339 are in proximity of the PEG8 molecule (Figure 6.5a).

The superimposition of the HIS1:Mn:2OG and OsHSL1:Mn:2OG structures reveals that multiple residues of those in proximity of the PEG8 molecule are not conserved between HIS1 and OsHSL1, *e.g.*, H141, F205, L299, and F336 in HIS1 are substituted for F140, L204, F298, and I335 in OsHSL1 (Figure 6.4c), the side chains of which differ in steric bulk and/or polarity. These differences could rationalise the different substrate scopes of HIS1 and OSHSL1. This proposal is supported by analysis of the superimposed HIS1:Mn:2OG and the ZmHSL1A:Mn:2OG:mesotrione (**1**) structures (PDB ID: 8X74[8]; RMSD = 0.72 Å) which reveals that **1** and PEG8 bind at similar locations in the active site (Figure 6.5a and b). Note that **1** is bound in a likely non-productive orientation at the ZmHSL1A active site, with its cyclohexane-1,3-dione ring facing away from Mn, consistent with the reported inability of ZmHSL1A to catalyse hydroxylation of **1**.[8]

Notably, HIS1 and ZmHSL1A residues in proximity of PEG8 and **1**, respectively, are not conserved, *e.g.*, H141, F205, L299, and F336 in HIS1 are substituted for Q142, I207, F301, and R338 in ZmHSL1A, possibly indicating why the substrate scopes of HIS1 and ZmHSL1A differ.[8] Superimposition of the OsHSL1:Mn:2OG structure with the reported ZmHSL1A:Mn:2OG:mesotrione structure (PDB ID: 8X74;[8] RMSD = 0.78 Å) reveals that

residues involved in mesotrione binding are conserved to a higher degree than the case with HIS1, consistent with a similar substrate scope for OsHSL1 and ZmHSL1A (Figure 6.5b).




Figure 6.5. PEG binds to HIS1 at the potential substrate binding site. (**a**) Representative Polder omit electron density maps contoured to 5σ around PEG8 and 2OG in the HIS1:Mn:2OG structure (PDB ID: 8S7B). (**b-c**) Overlay views of a reported crystal structure of ZmHSL1A in complex with Mn(II), 2OG, and mesotrione (**1**) (PDB ID: 8X74[8]) with: (**b**) OsHSL1 (PDB ID: 9GI5; RMSD = 0.78 Å) and (**c**) HIS1 (PDB ID: 8S7B; RMSD = 0.72 Å). Pink residues: HIS1; lavender residues: OsHSL1; yellow residues: ZmHSL1A; the DSBH core fold: cyan. Mn(II): magenta spheres; 2OG: green.

Table 6.1. Specific activities of HIS1 and OsHSL1 with the HPPD inhibitors **1-8** and the plant growth regulators **9** and **10**.

Substrate	[a]Specific activities [nmol·min^{-1}·mg^{-1}]	
	HIS1	**OsHSL1**
1	10 ± 3	n.r.
2	12 ± 2	n.r.
3	3 ± 1	n.r.
4	5 ± 3	6 ± 2
5	13 ± 4	n.r.
6	4 ± 1	n.r.
7	n.r.	n.r.
8	n.r.	n.r.
9	n.r.	n.r.
10	16 ± 3	n.r.

a) Specific activities (nmol·min^{-1}·mg^{-1}) based on formation of the C5-hydroxylated product after ~20 min incubations with HIS1 and ~90 min incubations with OsHSL1 (Figure 6.3). Conditions: 400 µM substrate, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM HIS1 or 20 µM OsHSL1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. n.r.: no reaction. Results are means of independent triplicates (n = 3; mean ± SD).

6.5 Discussion

Plant herbicide resistance mechanisms can be broadly categorised into target-site resistance (TSR) and non-target-site resistance (NTSR).[27] TSR arises from mutations or overexpression of the herbicide-target enzyme and consequently reduces herbicide sensitivity.[27,54] By contrast, NTSR involves metabolic detoxification or reduced uptake of the herbicide. The identification of HIS1[6] and phylogenetically related HSLs in rice and other crops (Supporting Figure 6.S1-6.S2)[3-7], which all catalyse oxidation of βTHs, suggests that the efficacy of HPPD-targeting

βTH inhibitors may be threatened given that similar 2OG oxygenases have been identified in weeds.[6,8,31,32]

We report ^{1}H NMR-based assays to monitor HIS1/OsHSL1 catalysis which are suitable to rank substrates and which enable product characterisation (Table 6.1; Figure 6.3); they complement reported MS-based assays.[6,8,9,30-32] The combined results are overall consistent with literature data, showing that catalysis of isolated HIS1/OsHSL1 is relatively inefficient and that the substrate scope of HIS1 is substantially broader than that of OsHSL1 (Table 6.1). The ^{1}H NMR assays support the assignment of the HIS1/OsHSL1 reaction products as the C5 hydroxylated βTHs.

Notably, the second product of the HIS1-catalysed reaction with **1-5** shows characteristic ^{1}H NMR signals of acyl substituted phloroglucinols.[41-43] This second product may be formed as a consequence of a second-hydroxylation catalysed by HIS1 at the C5 position of the first product to give the corresponding C5-dihydroxylated product which loses water to produce the corresponding ketone that can tautomerise in solution to give the corresponding substituted 1,3,5-benzenetriols (phloroglucinols) **1b-5b** (Figure 6.2, Supporting Figure 6.S4-6.S8). These observations are in accord with reported MS-based studies showing the formation of a second product with the mass corresponding to the ketone product.[32] Note, however, we did not accrue direct evidence for a second hydroxylation reaction.

Importantly, our results with **3** and **4** show that substitutions at the C3 position of the mesotrione aryl ring reduce HIS1 activity. Consistently, formation of the corresponding second product has not been observed for fenquinotrione (**6**) (Figure 6.3) suggesting that **6** might serve as a scaffold to develop improved β-triketone HPPD inhibitors which are not substrates of HIS1 and/or HSL-like enzymes. The AtHPPD:Mn:fenquinotrione complex structure suggests that the HPPD fenquinotrione binding site is sufficiently spacious to accommodate substituents at

the fenquinotrione C5 position (*e.g.*, fluorine atoms or methyl groups),[38] suggesting that fenquinotrione derivatives can be designed that are not HIS1/HSL substrates but retain efficient HPPD inhibition.

Our work supports the reported observation that prohexadione (**9**) is not a substrate of HIS1/OsHSL1, but instead inhibits catalysis[32] by showing that the oxidation of 2OG to succinic acid decoupled from substrate oxidation is lower in the presence of **9** (Figure 6.3). Thus, triketone **9** may serve as a scaffold for the development of HIS1 and/or HSLs inhibitors to rescue the activity of βTHs. Such HIS1/HSL inhibitors may be used in combination with βTHs to suppress the βTH metabolism.

The plant growth regulators **9** and **10** inhibit gibberellin biosynthesis by targeting GA oxidases.[32,55] The observation that HIS1 is inhibited by **9** but can metabolise **10** highlights that HSLs and GA oxidase activity differs, suggesting that selective inhibitors/substrates can be designed (Figure 6.3). Since, trinexapac ethyl (**10**) regulates GA biosynthesis, HIS1-mediated hydroxylation of **10** suggests its potential involvement in plant hormone homeostasis.

Together with reported crystallographic and mutational studies,[6,8,31,32] our HIS1:Mn:2OG and OsHSL1:Mn:2OG structures provide insights into substrate binding. The observation that a PEG8 molecule binds to the HIS1 active site at a position at which mesotrione binds in ZmHSL1A suggests conserved substrate binding sites amongst HIS1 and HSLs (Figure 6.5). Notably, however, the HIS1/ZmHSL1A residues in proximity of PEG8/mesotrione differ, *e.g.*, H141, F205, L299, and F336 in HIS1, possibly indicating why the substrate scope of HIS1 and HSLs also differs.

The serendipitous binding of PEG8 at the HIS1 active site suggests that polyketide-derived metabolites may be endogenous substrates of HIS1/HSLs. Since plants frequently encounter polyketide-type phytotoxic allelochemicals from microbes and other plants,[42,56] it is possible

that HIS1/HSLs may have originally evolved for their detoxification. Note that the widely used βTHs are derived from the phytotoxic plant metabolite, leptospermone.[11,57]

Transgenic expression of HIS1/HSLs in *O. sativa* and *A. thaliana* has shown broad-spectrum resistance against multiple βTHs without compromising plant growth, consistent with their ability to catalyse the oxidation of βTHs and the reduced herbicidal efficacy of the oxidised metabolites.[3-7] The ability to rationally modify the active sites of HIS1 and HSLs expands the horizon for engineering and tuning herbicide tolerance in crops enabling more efficient weed control via increasing herbicide selectivity. The design of HSLs which metabolise a specific βTH can also potentially limit the risk of resistance evolution in weeds, as seeds encoding for different HSL proteins may be used annually.

The distinct activities and roles of HIS1 and OsHSL1 raise questions about their physiological substrates and functions; they apparently differ in the lifecycle of the plant. HIS1, localised in leaves, is linked to the early stages of plant growth, while OsHSL1 is expressed in roots later during plant development.[6] It is possible that HIS1 and OsHSL1 may have evolved to guard different exposure windows during the plant lifecycle.

6.6 Experimental Procedures

Chemicals. All chemicals used in this work were from commercial sources (Sigma Aldrich Inc., Fluorochem Ltd., Bionet Key Organics, Medchem Express, Tokyo Chemical Industry Co. Ltd.) and were used without further purification.

HIS1 and OsHSL1 production and purification. Pet28a-(+)-TEV plasmids encoding for HIS1 (UNIPROT ID: Q6ES21) or OsHSL1 (UNIPROT ID: Q8H620), codon optimised for expression in *E. coli* cells, were obtained from GenScript Biotech. The plasmids were transformed into competent T7 Express LysY competent *E. coli* cells (New England Biolabs). Cells were cultured in 2YT media supplemented in the presence of 30 μg/mL kanamycin at 37 °C at 180 rpm, until HIS1/OsHSL1 expression was induced at an OD_{600} ~0.6-0.8 by addition of isopropyl-β-D-thiogalactopyranoside (final concentration: 0.5 mM). Cells were incubated at 28 °C for 4 h with agitation at 180 rpm, before being harvested by centrifugation (11,395 x g, 15 min, 4 °C); cell pellets were stored at –80 °C.

For purification, the cell pellet was resuspended in ice-cold buffer (50 mM Tris, pH 8.0, 500 mM NaCl, 20 mM imidazole) in the presence of DNAase I (bovine pancreas, Sigma-Aldrich) and lysed at 4 °C at 20 kPSI using a high-pressure cell disrupter (Continuous Flow Cell Disruptor CF1; Constant Systems Ltd) followed by centrifugation (58545 x g, 20 min, 4 °C). The supernatant was purified at 4 °C using Ni(II)-affinity chromatography (HisTrap HP column, GE Healthcare) and an ÄKTA Pure chromatography system (GE Healthcare) with an elution gradient of increasing imidazole concentration (from 20 mM to 500 mM imidazole) in buffer (50 mM Tris, pH 8.0, 500 mM NaCl). Fractions containing the protein of interest (determined by SDS-PAGE analysis) were combined, concentrated using Amicon Ultra centrifugal filters (3082 x g, 4 °C), and buffer exchanged using Amicon Ultra centrifugal filters (3082 x g, 4 °C) to remove imidazole (25 mM Tris, pH 8.0, 500 mM NaCl, 10%$_{v/v}$ glycerol). The *N*-terminal His_6 tag was cleaved overnight using the Tobacco Etch Virus protease (TEV

protease) at 4 °C; and the resultant mixture was purified using Ni(II)-affinity chromatography (50 mM Tris, pH 8.0, 500 mM NaCl, 20 mM imidazole). The flow-through was collected, concentrated, and purified using size-exclusion chromatography (300 mL HiLoad 26/60 Superdex 200 pg column; 1 mL/min; ÄKTA Pure chromatography system) and gel filtration buffer (25 mM Tris, pH 8.0, 100 mM NaCl). Eluted fractions containing >95% pure HIS1/OsHSL1 (determined by SDS-PAGE analysis) were pooled, then concentrated using Amicon Ultra centrifugal filters (3082 x g, 4 °C) and aliquoted. The aliquots were flash frozen in liquid N_2 and stored at –80 °C. The protein concentration was determined using a NanoDrop One microvolume UV-Vis spectrophotometer (Thermo Scientific); fresh HIS1/OsHSL1 aliquots were used for all biochemical experiments.

^{1}H NMR assays. Aqueous solutions of 2OG (80 mM), *L*-ascorbic acid (LAA; 50 mM), and ferrous ammonium sulphate (FAS; 100 mM) were prepared on the day the assay was performed. Substrate solutions (100 mM) were prepared in DMSO-d_6 because of typically limited water solubility and stored at –20 °C. These solutions and a solution (100 mM) of (trimethylsilyl)propionic-2,2,3,3-d_4 acid sodium salt (TMSP-d_4), which was used as an internal standard, were diluted using buffer (50 mM sodium phosphate, 10%$_{v/v}$ D_2O, pH 6.0) to give: 2OG (8 mM), LAA (5 mM), FAS (0.5 mM), and TMSP-d_4 (8 mM) stock solutions for direct use in the ^{1}H NMR assays.

^{1}H NMR assays were performed in a reaction volume of 180 µL with the following final concentrations of assay components, added strictly in the following order: 500 µM *L*-ascorbic acid, 800 µM 2OG, 400 µM substrate, 50 µM Fe(II), 800 µM TMSP-*d4,* and 20 µM HIS1/OsHSL1. The assay mixture was carefully mixed with a glass Pasteur pipette, transferred to a 3 mm Bruker NMR tube, and hand centrifuged. Assays were performed using a Bruker AVIII 600 or a Bruker AVIII 700 MHz NMR spectrometer equipped with a 5-mm inverse

triple-resonance-inverse cryoprobe. 2D NMR characterisation was run on a Bruker AVIII 750 NMR spectrometer equipped with a 5-mm inverse triple-resonance-inverse cryoprobe.

The NMR tubes were quickly transferred to the NMR spectrometer. The probe was locked to D_2O (10%$_{v/v}$ in water), followed by auto-tuning. The sample was then shimmed, receiver gain was adjusted, and consecutive acquisitions of ^{1}H NMR spectra were started. The variable delay was set to zero and the number of scans per acquisition was set to 64 scans per acquisition to ensure a sufficient signal-to-noise ratio; the relaxation delay was minimized to reduce the time-gap between acquisitions. For ^{1}H NMR experiments, water suppression was achieved using perfect echo-modified WATERGATE experiments.[58] Assays were performed in independent triplicates.

Spectra were automatically phased and baseline corrected. Integral precision was attenuated by zero filling which adjusted the size of the spectra to the number of data points used. Spectra were integrated using MestReNova x86_64 (Mestrelab research) and integrals calibrated and scaled to the internal standard TMSP-d_4. Data was plotted using GraphPad Prism 9.

Crystallography. Full-length HIS1 and OsHSL1 (10 mg/mL) were incubated with $MnCl_2$ (4 mM) and 2OG (10 mM) in buffer (50 mM Tris, pH 7.5) at 4 °C for 30 min and centrifuged using a MicroCL 21R centrifuge (Thermo Fisher Scientific; 18,800 × *g*, 4 °C, 1 min). Crystallisation was carried out using sitting drop vapor diffusion method (300 nL drops) at 17 °C using 2:1, 1:1, and 1:2 well solution (µL): protein solution (µL) ratios in 96-3-subwell low profile SWISSSCI plates (Molecular Dimensions), using a Mosquito crystallisation robot (SPT Labtech). HIS1 conditions: 0.15 M magnesium acetate tetrahydrate, 0.1 M sodium citrate, pH 5.6, and 20% $_{v/v}$ PEG Smear Broad. OsHSL1 conditions: 0.1 M carboxylic acids, 1.0 M imidazole, MES hydrate, pH 6.5, 30% $_{v/v}$ glycerol, and PEG 4000. Crystals were cryo-protected with 15% $_{v/v}$ glycerol prior to flash-freezing with liquid N_2.

Diffraction data for HIS1 and OsHSL1 single crystals were collected at the Diamond Light Source (beamline: I03). The data were indexed, integrated, and scaled using the Xia2 pipeline.[59,60] Structures were solved by molecular replacement with Phaser,[61] using an AlphaFold[50] model structure of HIS1 from *Oryza sativa* as a search model (Accession code: AF_AFQ6ES21F1). Structures were iteratively refined using PHENIX[62] and manual model building using COOT.[63]

6.7 Acknowledgements

This research was funded in part by the Wellcome Trust (106244/Z/14/Z). For the purpose of open access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission. We thank the Biotechnology and Biological Sciences Research Council (BB/J003018/1 and BB/R000344/1) for funding. We thank Dr. C. Mycroft, Dr. A. Tumber, and Ms. Y. Wang for helpful discussions.

6.8 Author Contributions

S.D. and I.M.F. produced enzymes; S.D., S.S., and F.M.A. performed assays; S.D. and C.R. characterised reaction products; S.D. and I.M.F. crystallised the proteins with the help of F.M.A.; M.D.A. solved the crystal structures; C.J.S, L.B., and S.D. conceived the project; L.B. and C.J.S. supervised the research; all authors analysed data; S.D., L.B., and C.J.S. wrote the manuscript with help from all authors.

6.9 References

1 Paul, S. K., Mazumder, S. & Naidu, R. Herbicidal weed management practices: History and future prospects of nanotechnology in an eco-friendly crop production system. *Heliyon* **10**, e26527 (2024). https://doi.org/https://doi.org/10.1016/j.heliyon.2024.e26527

2 Kubiak, A., Wolna-Maruwka, A., Niewiadomska, A. & Pilarska, A. A. The Problem of Weed Infestation of Agricultural Plantations vs. the Assumptions of the European Biodiversity Strategy. *Agronomy* **12**, 1808 (2022).

3 Aung, K. M. *et al.* Analyzing the response of rice to tefuryltrione herbicide: Haplotype variation and evolutionary dynamics of the HIS1 gene. *Environmental Research* **252**, 118839 (2024). https://doi.org/https://doi.org/10.1016/j.envres.2024.118839

4 Wu, J. *et al.* Directed evolution of HIS1-Like proteins to enhance herbicide metabolism in crops. *Advanced Agrochem* (2025). https://doi.org/https://doi.org/10.1016/j.aac.2025.05.005

5 Takamura, N. *et al.* Catalytic promiscuity of rice 2-oxoglutarate/Fe(II)-dependent dioxygenases supports xenobiotic metabolism. *Plant Physiol* **187**, 816–828 (2021). https://doi.org/10.1093/plphys/kiab293

6 Maeda, H. *et al.* A rice gene that confers broad-spectrum resistance to β-triketone herbicides. *Science* **365**, 393–396 (2019). https://doi.org/doi:10.1126/science.aax0379

7 Lv, Q. *et al.* Exploring Natural Allelic Variations of the β-Triketone Herbicide Resistance Gene HIS1 for Application in indica Rice and Particularly in Two-Line Hybrid Rice. *Rice (N Y)* **14**, 7 (2021). https://doi.org/10.1186/s12284-020-00448-7

8 Dong, J. *et al.* An Artificially Evolved Gene for Herbicide-resistant Rice Breeding. *Proc. Natl. Acad. Sci. U. S. A.* **121**, e2407285121 (2024). https://doi.org/doi:10.1073/pnas.2407285121

9 Duff, S. M. G. *et al.* Structural and Functional Characterization of Triketone Dioxygenase from *Oryza Sativa*. *Biochem. Biophys. Res. Commun.* **1868**, 130504 (2024). https://doi.org/https://doi.org/10.1016/j.bbagen.2023.130504

10 Kraehmer, H. *et al.* Herbicides as Weed Control Agents: State of the Art: II. Recent Achievements. *Plant Physiol.* **166**, 1132–1148 (2014). https://doi.org/10.1104/pp.114.241992

11 Mitchell, G. *et al.* Mesotrione: A New Selective Herbicide for Use in Maize. *Pest Manag. Sci.* **57**, 120–128 (2001). https://doi.org/https://doi.org/10.1002/1526-4998(200102)57:2<120::AID-PS254>3.0.CO;2-E

12 Lin, H.-Y. *et al.* Molecular Insights into the Mechanism of 4-Hydroxyphenylpyruvate dioxygenase Inhibition: Enzyme Kinetics, X-ray Crystallography and Computational Simulations. *FEBS J.* **286**, 975–990 (2019). https://doi.org/https://doi.org/10.1111/febs.14747

13 Küpper, A. *et al.* Tembotrione detoxification in 4-hydroxyphenylpyruvate dioxygenase (HPPD) inhibitor-resistant Palmer amaranth (Amaranthus palmeri S. Wats.). *Pest Manag. Sci.* **74**, 2325–2334 (2018). https://doi.org/10.1002/ps.4786

14 Nagamatsu, A., Ueda, K., Tamai, R., Tani, S. & Yamamoto, S. Development of a Rice Herbicide, Fenquinotrione. *J Pestic. Sci.* **47**, 139–145 (2022). https://doi.org/10.1584/jpestics.J22-02

15 Brabham, C., Norsworthy, J. K., Sha, X., Varanasi, V. K. & González-Torralva, F. Benzobicyclon efficacy is affected by plant growth stage, HPPD Inhibitor Sensitive 1 (HIS1) expression and zygosity in weedy rice (Oryza sativa). *Weed Science* **70**, 328–334 (2022). https://doi.org/10.1017/wsc.2022.17

16 Santucci, A., Bernardini, G., Braconi, D., Petricci, E. & Manetti, F. 4-Hydroxyphenylpyruvate Dioxygenase and Its Inhibition in Plants and Animals: Small Molecules as Herbicides and

Agents for the Treatment of Human Inherited Diseases. *J Med Chem* **60**, 4101–4125 (2017). https://doi.org/10.1021/acs.jmedchem.6b01395

17 Dayan, F. E. *et al.* β-Triketone Inhibitors of Plant p-Hydroxyphenylpyruvate Dioxygenase: Modeling and Comparative Molecular Field Analysis of Their Interactions. *Journal of Agricultural and Food Chemistry* **57**, 5194–5200 (2009). https://doi.org/10.1021/jf9005593

18 Dayan, F. E. *et al.* p-Hydroxyphenylpyruvate dioxygenase is a herbicidal target site for beta-triketones from Leptospermum scoparium. *Phytochemistry* **68**, 2004–2014 (2007). https://doi.org/10.1016/j.phytochem.2007.01.026

19 Moran, G. R. 4-Hydroxyphenylpyruvate dioxygenase. *Arch Biochem Biophys* **433**, 117–128 (2005). https://doi.org/10.1016/j.abb.2004.08.015

20 Havaux, M. Plastoquinone In and Beyond Photosynthesis. *Trends Plant Sci* **25**, 1252–1265 (2020). https://doi.org/10.1016/j.tplants.2020.06.011

21 Hasanuzzaman, M., Nahar, K. & Fujita, M.

22 Jhala, A. J. *et al.* 4-Hydroxyphenylpyruvate dioxygenase (HPPD)-inhibiting herbicides: past, present, and future. *Weed Technology* **37**, 1–14 (2023). https://doi.org/10.1017/wet.2022.79

23 Ahrens, H. *et al.* 4-Hydroxyphenylpyruvate dioxygenase inhibitors in combination with safeners: solutions for modern and sustainable agriculture. *Angew Chem Int Ed Engl* **52**, 9388–9398 (2013). https://doi.org/10.1002/anie.201302365

24 Lin, H.-Y., Dong, J., Dong, J., Yang, W.-C. & Yang, G.-F. Insights into 4-hydroxyphenylpyruvate dioxygenase-inhibitor interactions from comparative structural biology. *Trends in Biochemical Sciences* **48**, 568–584 (2023). https://doi.org/https://doi.org/10.1016/j.tibs.2023.02.006

25 Lin, H. Y. *et al.* Crystal Structure of 4-Hydroxyphenylpyruvate Dioxygenase in Complex with Substrate Reveals a New Starting Point for Herbicide Discovery. *Research (Wash D C)* **2019**, 2602414 (2019). https://doi.org/10.34133/2019/2602414

26 Paiman, Nurul Fasehah Ismail, S. & Shah Abdullah, A. Recent Developments of Weed Management in Rice Fields. *Reviews in Agricultural Science* **8**, 343–353 (2020). https://doi.org/10.7831/ras.8.0_343

27 Gaines, T. A. *et al.* Mechanisms of evolved herbicide resistance. *J Biol Chem* **295**, 10307–10330 (2020). https://doi.org/10.1074/jbc.REV120.013572

28 Jugulam, M. & Shyam, C. Non-Target-Site Resistance to Herbicides: Recent Developments. *Plants (Basel)* **8** (2019). https://doi.org/10.3390/plants8100417

29 *2-Oxoglutarate-Dependent Oxygenases*. (The Royal Society of Chemistry, 2015).

30 Wang, N. *et al.* Crystal structure of HPPD Inhibitor Sensitive Protein from *Oryza sativa*. *Biochem. Biophys. Res. Commun.* **704**, 149672 (2024). https://doi.org/https://doi.org/10.1016/j.bbrc.2024.149672

31 Wu, J. *et al.* Directed Evolution of HIS1-Like Proteins to Enhance Herbicide Metabolism in Crops. *Adv. Agrochem.* **in press** (2025). https://doi.org/https://doi.org/10.1016/j.aac.2025.05.005

32 Takamura, N. *et al.* Catalytic Promiscuity of Rice 2-Oxoglutarate/Fe(II)-dependent Dioxygenases Supports Xenobiotic Metabolism. *Plant Physiol.* **187**, 816–828 (2021). https://doi.org/10.1093/plphys/kiab293

33 Dhingra, S., Zhang, Z., Lohans, C. T., Brewitz, L. & Schofield, C. J. Substitution of 2-Oxoglutarate Alters Reaction Outcomes of the *Pseudomonas savastanoi* Ethylene-forming Enzyme. *J. Biol. Chem.* **300**, 107546 (2024). https://doi.org/https://doi.org/10.1016/j.jbc.2024.107546

34 Copper, A. J. & Redfield, A. G. Proton magnetic resonance studies of alpha-keto acids. *J. Biol. Chem.* **250**, 527–532 (1975). https://doi.org/https://doi.org/10.1016/S0021-9258(19)41928-5

35 Viswanathan, T. S., Johnson, R. E. & Fisher, H. F. α-Ketoglutaric acid: solution structure and the active form for reductive amination by bovine liver glutamate dehydrogenase. *Biochemistry* **21**, 339–345 (1982). https://doi.org/10.1021/bi00531a022

36 Sun, Y. D., S.; Allen, M. D.; Cheng, E. S. Z.; Zhang, Z.; Schofield, C. J.; Brewitz, L. . Structural and Functional Validation of *Pseudomonas savastanoi* Ethylene Forming Enzymes Reveals Flexibility in 2-Oxoglutarate Binding Mode and Conformation. . *Manuscript submitted.* (2025).

37 Dhingra, S. S., Y.; Zhang, Z.; Schofield, C. J.; Brewitz, L. . Expanding the Alkene Forming Capability of the *Pseudomonas savastanoi* Ethylene-forming Enzyme. . *Manuscript submitted.* (2025).

38 Alshref, F. *et al.* Structural and Inhibition studies on *Arabidopsis thaliana* HPPD. *Manuscript in preparation* (2025).

39 Kavana, M. & Moran, G. R. Interaction of (4-Hydroxyphenyl)pyruvate Dioxygenase with the Specific Inhibitor 2-[2-Nitro-4-(trifluoromethyl)benzoyl]-1,3-cyclohexanedione. *Biochemistry* **42**, 10238–10245 (2003). https://doi.org/10.1021/bi034658b

40 Bereziuk, O., Gubina, K., Trush, V. & Ovchynnikov, V. Crystal Structure of the Sodium Salt of Mesotrione: a Triketone Herbicide. *Acta Crystallogr. E.* **80**, 296–299 (2024). https://doi.org/doi:10.1107/S2056989024001439

41 Meepagala, K. M. & Schrader, K. K. Antibacterial Activity of Acylglucinol Derivatives against Flavobacterium columnare. *J. Agric. Chem. Environ.* **3**, 7 (2014). https://doi.org/10.4236/jacen.2014.33011

42 Popoola, O. K., Marnewick, J. L., Rautenbach, F., Iwuoha, E. I. & Hussein, A. A. Acylphloroglucinol Derivatives from the South African *Helichrysum niveum* and Their Biological Activities. *Molecules* **20**, 17309–17324 (2015). https://doi.org/10.3390/molecules200917309

43 Rahman, M. M., Shiu, W. K. P., Gibbons, S. & Malkinson, J. P. Total synthesis of acylphloroglucinols and their antibacterial activities against clinical isolates of multi-drug resistant (MDR) and methicillin-resistant strains of *Staphylococcus aureus*. *Eur. J. Med. Chem.* **155**, 255–262 (2018). https://doi.org/https://doi.org/10.1016/j.ejmech.2018.05.038

44 Katritzky, A. R., Ramsden, C. A., Joule, J. A. & Zhdankin, V. V. in *Handbook of Heterocyclic Chemistry (Third Edition)* (eds Alan R. Katritzky, Christopher A. Ramsden, John A. Joule, & Viktor V. Zhdankin) 37–86 (Elsevier, 2010).

45 Balci, M. in *Basic 1H- and 13C-NMR Spectroscopy* (ed Metin Balci) 87–133 (Elsevier Science, 2005).

46 Diwan, G. *et al.* Effect of New Plant Growth Regulator Prohexadione- Ca on Horticultural Crops. *AMA* **53**, 9671 (2022).

47 Li, D. *et al.* Prohexadione-calcium Improves Grape Quality by Regulating Endogenous Hormones, Sugar and Acid Metabolism and Related Enzyme Activities in Grape Berries. *BMC Plant Biol.* **24**, 122 (2024). https://doi.org/10.1186/s12870-024-04803-4

48 Aftab, T. & Naeem B.Sc, M. S. (Elsevier Science & Technology, 2021).

49 Trethewey, J. A. K., Rolston, M. P., McCloy, B. L. & Chynoweth, R. J. The Plant growth Regulator, Trinexapac-ethyl, Increases Seed Yield in Annual Ryegrass (*Lolium multiflorum Lam.*). *N.Z. J. Agri. Res.* **59**, 113–121 (2016). https://doi.org/10.1080/00288233.2015.1134590

50 Abramson, J. *et al.* Accurate Structure Prediction of Biomolecular Interactions with AlphaFold 3. *Nature* **630**, 493–500 (2024). https://doi.org/10.1038/s41586-024-07487-w

51 Aik, W. S. *et al.* in *2-Oxoglutarate-Dependent Oxygenases* (eds Christopher Schofield & Robert Hausinger) 0 (The Royal Society of Chemistry, 2015).

52 Holm, L., Laiho, A., Törönen, P. & Salgado, M. DALI Shines a Light on Remote Homologs: One Hundred Discoveries. *Protein Sci.* **32**, e4519 (2023). https://doi.org/https://doi.org/10.1002/pro.4519

53 Liebschner, D. *et al.* Polder Maps: Improving OMIT Maps by Excluding Bulk Solvent. *Acta Crystallogr. D. Struct. Biol.* **73**, 148–157 (2017). https://doi.org/10.1107/s2059798316018210

54 Murphy, B. P. & Tranel, P. J. Target-Site Mutations Conferring Herbicide Resistance. *Plants* **8**, 382 (2019).

55 Rademacher, W. Growth Retardants: Effects on Gibberellin Biosynthesis and Other Metabolic Pathways. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **51**, 501–531 (2000). https://doi.org/10.1146/annurev.arplant.51.1.501

56 Iqbal, S. *et al.* Allelochemicals Degradation and Multifarious Plant Growth Promoting Potential of Two *Bacillus* spp.: Insights Into Genomic Potential and Abiotic Stress Alleviation. *Chemosphere* **373**, 144191 (2025). https://doi.org/https://doi.org/10.1016/j.chemosphere.2025.144191

57 Owens, D. K., Nanayakkara, N. P. & Dayan, F. E. In Planta Mechanism of Action of Leptospermone: Impact of its Physico-chemical Properties on Uptake, Translocation, and Metabolism. *J. Chem. Ecol.* **39**, 262–270 (2013). https://doi.org/10.1007/s10886-013-0237-8

58 Adams, R. W., Holroyd, C. M., Aguilar, J. A., Nilsson, M. & Morris, G. A. "Perfecting" WATERGATE: clean proton NMR spectra from aqueous solution. *Chem. Commun.* **49**, 358–360 (2013). https://doi.org/10.1039/C2CC37579F

59 Winter, G. xia2: an expert system for macromolecular crystallography data reduction. *J. Appl. Cryst.* **43**, 186–190 (2010). https://doi.org/doi:10.1107/S0021889809045701


60 Winter, G. *et al.* DIALS as a toolkit. *Protein Sci.* **31**, 232–250 (2022). https://doi.org/https://doi.org/10.1002/pro.4224

61 McCoy, A. J. *et al.* Phaser crystallographic software. *J. Appl. Cryst.* **40**, 658–674 (2007). https://doi.org/doi:10.1107/S0021889807021206

62 Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in *Phenix*. *Acta Cryst.* **D75**, 861–877 (2019). https://doi.org/doi:10.1107/S2059798319011471

63 Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta Cryst. D* **66**, 486–501 (2010). https://doi.org/doi:10.1107/S0907444910007493


6.10 Supporting Figures



Supporting Figure 6.S1. Sequence comparison of HIS1 and HIS1-like enzymes (HSLs). The sequence of HIS1 (UNIPROT ID: Q6ES21) is compared to that of: *Oryza sativa* HIS1-like 1 (OsHSL1; UNIPROT ID: Q8H620), *Oryza sativa* HIS1-like 2 (OsHSL2; UNIPROT ID: Q8H618), *Oryza sativa* HIS1-like 4 (OsHSL4; UNIPROT ID: A0A7G2L474), *Oryza sativa* HIS1-like 5 (OsHSL5; UNIPROT ID: UPI001862E832), *Oryza sativa* HIS1-like 6 (OsHSL6; UNIPROT ID: A0A7G2L586), *Zea mays* HIS1-like 1A (ZmHSL1A; UNIPROT ID: B6U9L0), *Zea mays* HIS1-like 1B (ZmHSL1B; UNIPROT

ID: B6SI41), *Sorghum bicolor* HIS1-like 1A (SbHSL1A; UNIPROT ID: UPI0001A8917D), *Triticum aestivum* HIS1-like 1 (TaHSL1; UNIPROT ID: A0A9R1HWI4), *Triticum aestivum* HIS1-like 2 (TaHSL2; UNIPROT ID: UPI0012A9436F), *Hordeum vulgare* HIS1-like 1A (HvHSL1A; UNIPROT ID: Q7XJ26), and *Hordeum vulgare* HIS1-like 6A (HvHSL6A; UNIPROT ID: UPI0012A81F00). α-Helices and β-strands of HIS1 are numbered based on a HIS1 structure (PDB ID: 8S7B); β-turns: TT. The HXD…H Fe(II)-binding motif: green circles. Identical residues: red; residues of similar polarity: yellow. The Figure 6.was generated using ESPript 3.0[1].


a)




b)

	OsHSL5	OsHSL6	HvHSl6A	OsHSL2	TaHSL2	OsHSL4	ZmHSL1A	ZmHSL1B	SbHSL1A	HIS1	OsHSL1	TaHSL1	HvHSL1A
OsHSL5	100	50	46	48	46	45	48	46	45	44	48	43	44
OsHSL6	50	100	69	54	49	52	52	49	50	50	51	53	53
HvHSl6A	46	69	100	53	50	52	54	50	49	53	54	55	55
OsHSL2	48	54	53	100	67	54	58	57	56	61	61	57	59
TaHSL2	46	49	50	67	100	51	56	54	53	58	56	56	57
OsHSL4	45	52	52	54	51	100	58	52	52	53	52	54	55
ZmHSL1A	48	52	54	58	56	58	100	66	67	64	63	63	64
ZmHSL1B	46	49	50	57	54	52	66	100	89	61	59	58	62
SbHSL1A	45	50	49	56	53	52	67	89	100	60	59	59	61
HIS1	44	50	53	61	58	53	64	61	60	100	87	63	64
OsHSL1	48	51	54	61	56	52	63	59	59	87	100	64	63
TaHSL1	43	53	55	57	56	54	63	58	59	63	64	100	87
HvHSL1A	44	53	55	59	57	55	64	62	61	64	63	87	100


Supporting Figure 6.S2. Phylogenetic relationship and percentage sequence identity between HIS1 and HSLs. (**a**) The sequence alignment-based phylogram reveals distinct HIS1/HSL clades: the HIS1/OsHSL1 cluster, the ZmHSL1A/SbHSL1A cluster, and a divergent group containing OsHSL2–6 and HSLs from wheat (TaHSL2) and barley (HvHSL6A).[4-6] The phylogenetic tree was created using EMBL-EBI Simple Phylogeny[2] and visualised using Interactive Tree of Life version 7.0[3]. (**b**) Percentage sequence identity of: HIS1 and HSLs from rice (OsHSL1-6), maize (ZmHSL1A and ZmHSL1B), wheat (TaHSL1 and TaHSL2), Indian millet (SbHSL1A), and barley (HvHSL1A and HvHSL6A). The ZmHSL1B/SbHSL1A, HIS1/OsHSL1, and TaHSL1/HvHSL1A pairs share the highest sequence identity (>85%). Note that the HSLs are reported to have variable substrate scopes and catalytic efficiencies.[4-6] Sequence alignments are given in: Supporting Figure 6.S1.




Supporting Figure 6.S3. Purity of HIS1 and OsHSL1. (**a-b**) Mass spectrometric analyses of HIS1 and OsHSL1 (3 μM) in buffer (25 mM Tris, pH 7.8) were performed using an iFunnel Agilent 6550 accurate mass quadrupole time-of-flight (Q-TOF) mass spectrometer operated in the positive ionization mode for: (**a**) HIS1 (calculated mass: 40601 Da; observed mass: 40602 Da) and (**b**) OsHSL1 (calculated mass: 40418 Da; observed mass: 40417 Da). (**c**) SDS-PAGE analysis of isolated recombinant HIS1 and OsHSL1 indicate >95% purity. The PageRuler™ Plus Prestained Protein Ladder (Thermo Fisher) was used as a reference. Proteins were separated using a 4-12% NovexTM 330 Tris-glycine gel (Thermo Fisher) according to the manufacturer's protocol.




Supporting Figure 6.S4. pH dependence of the HIS1-catalysed hydroxylation of mesotrione (1). (**a-c**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of HIS1 and **1**, ~6 h post addition of HIS1 at pH: (**a**) 6.0, (**b**) 7.4, and (**c**) 8.0. (**d**) ^{1}H NMR spectrum of a reaction mixture in the absence of HIS1. Conditions: 400 µM **1**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM HIS1 in buffer (50 mM sodium phosphate, pH 6.0-8.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that the C5 proton of **1a** was not observed, likely because it coincides with the HOD peak; however, the splitting pattern of the C4/C6 proton signals implies that these protons couple with the C5 proton, as observed for other βTHs (Supporting Figures 6.S5 and 6.S8). For *J*-values of **1a** see: Supporting Table 6.S2. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Both **1** and **1a** were assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analyses of **1** and **5**;[5,11-13] however, **1** and/or **1a** could, in part, be present as the corresponding C2 *endo*-enol tautomers.[14] The peaks corresponding to the second reaction product **1b** were assigned by comparison with ^{1}H NMR spectra reported for structurally related 2-acylphloroglucinol derivatives, showing a characteristic singlet for C4/C6-protons at ~5.5 to ~6.5 ppm.[15-17] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S5. HIS1 catalyses hydroxylation of sulcotrione (2) in the presence of 2OG. (**a-c**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of HIS1 and **2**: (**a**) ~6 h, (**b**) ~90 min, and (**c**) ~5 min post addition of HIS1. (**d**) ^{1}H NMR spectrum of a reaction mixture in the absence of HIS1. (**e**) ^{1}H NMR spectrum of an authentic sample of **2** in buffer. Conditions: 400 µM **2**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM HIS1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. For *J*-values of **2a** see: Supporting Table 6.S2. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). The peaks at ~4.1 and ~4.2 ppm are from dehydroascorbate (annotated with: *).[18] Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Note that peaks at ~6.3 and ~7.0 ppm appear to increase with time but were not assigned due to their relatively low abundance. Both **2** and **2a** were assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analyses of **1** and **5**;[5,11-13] however, **2** and/or **2a** could, in part, be present as the corresponding C2 *endo*-enol tautomers.[14] The peaks corresponding to the second reaction product **2b** were assigned by comparison with ^{1}H NMR spectra reported for structurally related 2-acylphloroglucinol derivatives, showing a characteristic singlet for C4/C6-protons at ~5.5 to ~6.5 ppm.[15-17] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S6. HIS1 catalyses hydroxylation of tembotrione (3) in the presence of 2OG. (**a-c**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of HIS1 and **3**: (**a**) ~6 h, (**b**) ~90 min, and (**c**) ~5 min post addition of HIS1. (**d**) ^{1}H NMR spectrum of a reaction mixture in the absence of HIS1. (**e**) ^{1}H NMR spectrum of an authentic sample of **3** in buffer. Conditions: 400 µM **3**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM HIS1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that the C5 proton of **3a** was not observed, likely because it coincides with the HOD peak; however, the splitting pattern of the C4/C6 proton signals implies that these protons couple with the C5 proton, as observed for other βTHs (Supporting Figures 6.S5 and 6.S8). For *J*-values of **3a** see: Supporting Table 6.S2. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Note that the peak at ~7.0 ppm appears to increase with time but was not assigned due to its low abundance. Both **3** and **3a** were assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analyses of **1** and **5**;[5,11-13] however, **3** and/or **3a** could, in part, be present as the corresponding C2 *endo*-enol tautomers.[14] The peaks corresponding to the second reaction product **3b** were assigned by comparison with ^{1}H NMR spectra reported for structurally related 2-acylphloroglucinol derivatives, showing a characteristic singlet for C4/C6-protons at ~5.5 to ~6.5 ppm.[15-17] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S7. HIS1 catalyses hydroxylation of tefuryltrione (4) in the presence of 2OG. (**a-c**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of HIS1 and **4**: (**a**) ~6 h, (**b**) ~90 min, and (**c**) ~5 min post addition of HIS1. (**d**) ^{1}H NMR spectrum of a reaction mixture in the absence of HIS1. (**e**) ^{1}H NMR spectrum of an authentic sample of **4** in buffer. Conditions: 400 µM **4**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM HIS1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that the C5 proton of **4a** was not observed, likely because it coincides with the HOD peak; however, the splitting pattern of the C4/C6 proton signals implies that these protons couple with the C5 proton, as observed for other βTHs (Supporting Figures 6.S5 and 6.S8). For *J*-values of **4a** see: Supporting Table 6.S2. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Both **4** and **4a** were assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analyses of **1** and **5**;[5,11-13] however, **4** and/or **4a** could, in part, be present as the corresponding C2 *endo*-enol tautomers.[14] The peaks corresponding to the second reaction product **4b** were assigned by comparison with ^{1}H NMR spectra reported for structurally related 2-acylphloroglucinol derivatives, showing a characteristic singlet for C4/C6-protons at ~5.5 to ~6.5 ppm.[15-17] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S8. HIS1 catalyses hydroxylation of nitisinone (5) in the presence of 2OG. (**a-c**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of HIS1 and **5**: (**a**) ~6 h, (**b**) ~90 min, and (**c**) ~5 min post addition of HIS1. (**d**) ^{1}H NMR spectrum of a reaction mixture in the absence of HIS1. (**e**) ^{1}H NMR spectrum of an authentic sample of **5** in buffer. Conditions: 400 µM **5**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM HIS1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. For *J*-values of **5a** see: Supporting Table 6.S2. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). The peaks at ~4.1 and ~4.2 ppm are from dehydroascorbate (annotated with: *).[18] Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Both **5** and **5a** were assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analyses of **1** and **5**;[5,11-13] however, **5** and/or **5a** could, in part, be present as the corresponding C2 *endo*-enol tautomers.[14] The peaks corresponding to the second reaction product **5b** were assigned by comparison with ^{1}H NMR spectra reported for structurally related 2-acylphloroglucinol derivatives, showing a characteristic singlet for C4/C6-protons at ~5.5 to ~6.5 ppm.[15-17] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S9. HIS1 catalyses hydroxylation of fenquinotrione (6) in the presence of 2OG. (**a-c**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of HIS1 and **6**: (**a**) ~6 h, (**b**) ~90 min, and (**c**) ~5 min post addition of HIS1. (**d**) ^{1}H NMR spectrum of a reaction mixture in the absence of HIS1. (**e**) ^{1}H NMR spectrum of an authentic sample of **6** in buffer. Conditions: 400 µM **6**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM HIS1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that the C5 proton of **6a** was not observed, likely because it coincides with the HOD peak; however, the splitting pattern of the C4/C6 proton signals implies that these protons couple with the C5 proton, as observed for other βTHs (Supporting Figures 6.S5 and 6.S8). For *J*-values of **6a** see: Supporting Table 6.S2. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Note that the aromatic and methoxy peaks of **6a** were not assigned because of peak broadening and overlapping with substrate peaks. **6** is assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analyses of **1** and **5**;[5,11-13] however, **6** could, in part, be present as the corresponding C2 *endo*-enol tautomer.[14] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S10. HIS1 does not catalyse oxidation of isoxaflutole (7) in the presence of 2OG. (**a-c**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of HIS1 and **5**: (**a**) ~6 h, (**b**) ~90 min, and (**c**) ~5 min post addition of HIS1. (**d**) ^{1}H NMR spectrum of a reaction mixture in the absence of HIS1. (**e**) ^{1}H NMR spectrum of an authentic sample of **7** in buffer. Conditions: 400 µM **7**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM HIS1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that turnover of 2OG to succinic acid uncoupled from the oxidation of **7** was observed. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. The peaks at ~4.1 and ~4.2 ppm are from dehydroascorbate (annotated with: *).[18] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S11. HIS1 does not catalyse oxidation of topramezone (8) in the presence of 2OG. (**a-c**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of HIS1 and **8**: (**a**) ~6 h, (**b**) ~90 min, and (**c**) ~5 min post addition of HIS1. (**d**) ^{1}H NMR spectrum of a reaction mixture in the absence of HIS1. (**e**) ^{1}H NMR spectrum of an authentic sample of **8** in buffer. Conditions: 400 µM **8**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM HIS1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that turnover of 2OG to succinic acid uncoupled from the oxidation of **7** was observed. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S12. HIS1 does not catalyse oxidation of prohexadione (9) in the presence of 2OG. (**a-c**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~7.0 ppm) of a reaction of HIS1 and **9**: (**a**) ~6 h, (**b**) ~90 min, and (**c**) ~5 min post addition of HIS1. (**d**) ^{1}H NMR spectrum of a reaction mixture in the absence of HIS1. (**e**) ^{1}H NMR spectrum of an authentic sample of **9** in buffer. Conditions: 400 µM **9**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM HIS1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that turnover of 2OG to succinic acid uncoupled from the oxidation of **7** was observed. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. **9** is assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analyses of **1** and **5**;[5,11-13] however, **9** could, in part, be present as the corresponding C2 *endo*-enol tautomer.[14] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S13. HIS1 catalyses hydroxylation of trinexapac ethyl (10) in the presence of 2OG. (**a-c**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of HIS1 and **10**: (**a**) ~6 h, (**b**) ~90 min, and (**c**) ~5 min post addition of HIS1. (**d**) ^{1}H NMR spectrum of a reaction mixture in the absence of HIS1. (**e**) ^{1}H NMR spectrum of an authentic sample of **10** in buffer. Conditions: 400 µM **10**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM HIS1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. For *J*-values of **10a** see: Supporting Table 6.S2. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG[7-10] (dashed boxes). The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Both **10** and **10a** were assigned as the C2 *exo*-enol tautomer, as implied by crystallographic analysis of **1** and ^{1}H NMR analyses of **1** and **5**,[5,11-13] however, **10** and/or **10a** could, in part, be present as the corresponding C2 *endo*-enol tautomers.[14] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S14. OsHSL1 catalyses hydroxylation of tefuryltrione (4) in the presence of 2OG. (**a-b**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of OsHSL1 and **4**: (**a**) ~6 h, (**b**) ~5 min post addition of OsHSL1. (**c**) ^{1}H NMR spectrum of a reaction mixture in the absence of OsHSL1. (**d**) ^{1}H NMR spectrum of an authentic sample of **4** in buffer. Conditions: 400 µM **4**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM OsHSL1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that the C5 proton of **4a** was not observed, likely because it coincides with the HOD peak; however, the splitting pattern of the C4/C6 proton signals implies that these protons couple with the C5 proton as observed for other βTHs (Supporting Figures 6.S5 and 6.S8). For *J*-values of **4a** see: Supporting Table 6.S2. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). The peaks at ~4.1 and ~4.2 ppm are from dehydroascorbate (annotated with: *).[18] Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. Note that the peak at ~7.6 ppm appears to increase with time but was not assigned due to its relatively low abundance. Both **4** and **4a** were assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analyses of **1** and **5**;[5,11-13] however, **4** and/or **4a** could, in part, be present as the corresponding C2 *endo*-enol tautomers.[14] The peaks corresponding to the second reaction product **4b** were assigned by comparison with ^{1}H NMR spectra reported for structurally related 2-acylphloroglucinol derivatives, showing a characteristic singlet for C4/C6-protons at ~5.5 to ~6.5 ppm.[15-17] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S15. OsHSL1 does not catalyse oxidation of mesotrione (1) in the presence of 2OG. (**a-b**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of OsHSL1 and **1**: (**a**) ~6 h, (**b**) ~5 min post addition of OsHSL1. (**c**) ^{1}H NMR spectrum of a reaction mixture in the absence of OsHSL1. (**d**) ^{1}H NMR spectrum of an authentic sample of **1** in buffer. Conditions: 400 µM **1**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM OsHSL1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that turnover of 2OG to succinic acid uncoupled from the oxidation of **1** was observed. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. **1** is assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analyses of **1** and **5**;[5,11-13] however, **1** could, in part, be present as the corresponding C2 *endo*-enol tautomer .[14] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S16. OsHSL1 does not catalyse oxidation of sulcotrione (2) in the presence of 2OG. (**a-b**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of OsHSL1 and **2**: (**a**) ~6 h, (**b**) ~5 min post addition of OsHSL1. (**c**) ^{1}H NMR spectrum of a reaction mixture in the absence of OsHSL1. (**d**) ^{1}H NMR spectrum of an authentic sample of **2** in buffer. Conditions: 400 µM **2**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM OsHSL1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that turnover of 2OG to succinic acid uncoupled from the oxidation of **2** was observed. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. **2** is assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analyses of **1** and **5**;[5,11-13] however, **2** could, in part, be present as the corresponding C2 *endo*-enol tautomer.[14] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S17. OsHSL1 does not catalyse oxidation of tembotrione (3) in the presence of 2OG. (**a-b**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of OsHSL1 and **3**: (**a**) ~6 h, (**b**) ~5 min post addition of OsHSL1. (**c**) ^{1}H NMR spectrum of a reaction mixture in the absence of OsHSL1. (**d**) ^{1}H NMR spectrum of an authentic sample of **3** in buffer. Conditions: 400 µM **3**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM OsHSL1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that turnover of 2OG to succinic acid uncoupled from the oxidation of **3** was observed. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). The peaks at ~4.1 and ~4.2 ppm are from dehydroascorbate (annotated with: *).[18] Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. **3** is assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analysis of **1** and **5**;[5,11-13] however, **3** could, in part, be present as the corresponding C2 *endo*-enol tautomer.[14] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.




Supporting Figure 6.S18. OsHSL1 does not catalyse oxidation of nitisinone (5) in the presence of 2OG. (**a-b**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of OsHSL1 and **5**: (**a**) ~6 h, (**b**) ~5 min post addition of OsHSL1. (**c**) ^{1}H NMR spectrum of a reaction mixture in the absence of OsHSL1. (**d**) ^{1}H NMR spectrum of an authentic sample of **5** in buffer. Conditions: 400 µM **5**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM OsHSL1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that turnover of 2OG to succinic acid uncoupled from the oxidation of **5** was observed. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. The peaks at ~4.1 and ~4.2 ppm are from dehydroascorbate (annotated with: *).[18] **5** is assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analysis of **1** and **5**;[5,11-13] however, **5** could, in part, be present as the corresponding C2 *endo*-enol tautomer.[14] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.



Supporting Figure 6.S19. OsHSL1 does not catalyse oxidation of fenquinotrione (6) in the presence of 2OG. (**a-b**) ^{1}H NMR analysis (~1.5 to ~4.0 ppm and ~5.5 to ~9.0 ppm) of a reaction of OsHSL1 and **6**: (**a**) ~6 h, (**b**) ~5 min post addition of OsHSL1. (**c**) ^{1}H NMR spectrum of a reaction mixture in the absence of OsHSL1. (**d**) ^{1}H NMR spectrum of an authentic sample of **6** in buffer. Conditions: 400 µM **6**, 800 µM 2OG, 500 µM *L*-ascorbic acid, 50 µM Fe(II), 800 µM TSP-d_4, and 20 µM OsHSL1 in buffer (50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O) at 25˚C. Note that turnover of 2OG to succinic acid uncoupled from the oxidation of **6** was observed. Signals at ~2.2 ppm and ~2.0 ppm correspond to the hydrated ketone form of 2OG (dashed boxes).[7-10] The peak at ~2.6 ppm is from DMSO-d_6 (annotated with: *). The peaks at ~4.1 and ~4.2 ppm are from dehydroascorbate (annotated with: *).[18] Peaks at ~3.6 and ~3.7 ppm are from glycerol; the singlet at ~3.7 ppm corresponds to Tris. **6** is assigned as the C2 *exo*-enol form, consistent with a small-molecule crystal structure of **1** and ^{1}H NMR analysis of **1** and **5**;[5,11-13] however, **6** could, in part, be present as the corresponding C2 *endo*-enol tautomer.[14] Representative ^{1}H NMR spectra (700 MHz) of independent triplicates are shown.



Supporting Figure 6.S20. HIS1 and OsHSL1 share features with the IPNS/ACCO subfamily of 2OG oxygenases. (**a-b**) The overall fold of: (**a**) HIS1 and (**b**) OsHSL1 comprises 9 α-helices and 14 β-strands of which eight β-strands (I-VIII) form the conserved distorted DSBH core fold. Non-DSBH β-strands: yellow; α-helices: red; loops: green. β-Strands and α-helices are sequentially numbered.[1] (**c-d**) Superimposition of views from a HIS1:Mn:2OG structure (PDB ID: 8S7B) with that from: (**c**) a IPNS:Fe:HCV structure (PDB ID: 3ZKU; RMSD = 1.89 Å; ~22% sequence identity) and (**d**) an ACCO:Ni:ACC structure (PDB ID: 5TCV; RMSD = 1.48 Å; ~28% sequence identity) reveals conserved structural features around the DSBH core fold, suggesting HIS1/OsHSL1 may be related to the IPNS/ACCO subfamily of 2OG oxygenases.[19]



Supporting Figure 6.S21. The L204, F298, and I335 residues of OsHSL1 affect its reactivity with βTHs. (**a-b**) Superimposition of views from a L204F/F298L/I335F OsHSL1 variant:Co:2OG:ACHD structure (PDB ID: 8X6Q[5]) with that from: (**a**) a HIS1:Mn:2OG structure (PDB ID: 8S7B; RMSD: 0.511 Å) and (**b**) a OsHSL1:Mn:2OG structure (PDB ID: 9GI5; RMSD: 0.60 Å). The OsHSL1 variant in which the L204, F298, and I335 were substituted for the equivalent HIS1 residues is reported to manifest an improved catalytic efficiency and broader βTH substrate scope similar to that of HIS1,[5] suggesting that these residues are involved in substrate binding.





6.11 Supporting Tables




Supporting Table 6.S1. Data collection and refinement statistics for the HIS1:Mn:2OG and OsHSL1:Mn:2OG complex crystal structures.

	HIS1:Mn:2OG	OsHSL1:Mn:2OG
PDB ID	**8S7B**	**9GI5**
Data collection		
Space group	C 1 2 1	P 21 21 21
Cell dimensions		
a, *b*, *c* (Å)	68.33, 67.43, 83.18	89.67, 102.59, 156.06
α, β, γ (°)	90.00, 92.09, 90.00	90.00, 90.00, 90.00
Resolution (Å)	47.98 - 1.75 (1.78 - 1.75)	51.30 – 2.35 (2.41 - 2.35)
R_{merge}	0.060 (0.895)	0.147 (1.529)
I / sI	18.5 (1.1)	12.9 (1.7)
CC (1/2)	0.999 (0.804)	0.999 (0.688)
Completeness (%)	99.8 (98.0)	99.9 (99.8)
Redundancy	7.0 (6.1)	13.7 (13.1)
Refinement		
No. unique reflections	38097 (1983)	60662 (3007)
R_{work} / R_{free}	0.175 / 0.203	0.198 / 0.233
No. atoms		
Protein	3128	11683
Ligand/ion	11	78
Water	252	226
B-factors		
Protein	39.92	68.97
Ligand/ion	36.25	55.24
Water	42.34	48.62
R.m.s. deviations		
Bond lengths (Å)	0.04	0.04
Bond angles (°)	0.61	0.69

*Values in parentheses are for highest-resolution shell.

Supporting Table 6.S2. Assignment of the C5-hydroxylated HIS1 reaction products with triketones based on ^{1}H NMR analysis.

C5-Hydroxylated product of HIS1 catalysis	**Peak assignments** using ^{1}H NMR (600 MHz, 50 mM sodium phosphate, pH 6.0, 10%$_{v/v}$ D_2O)
 **1a***	δ: 8.73 (d, *J* = 1.8 Hz, 1H, **f**), 8.28 (dd, *J* = 8.0, 1.8 Hz, 1H, **e**), 7.54 (d, *J* = 8.0 Hz, 1H, **d**), 3.37 (s, 3H, **g**), 2.75 (dd, *J* = 16.7, 3.9 Hz, 2H, **a**), 2.50 ppm (dd, *J* = 16.6, 6.6 Hz, 2H, **b**).
 **2a**	δ: 8.02 (s, 1H, **f**), 7.89 (dd, *J* = 8.1, 1.7 Hz, 1H, **e**), 7.43 (d, *J* = 8.1 Hz, 1H, **d**), 4.60 (m, 1H, **c**), 3.31 (s, 3H, **g**), 2.66 (dd, *J* = 16.7 Hz, 4.0 Hz, 2H, **a**), 2.46 ppm (dd, *J* = 16.6, 6.7 Hz, 2H, **b**).
 **3a***	δ: 8.05 (d, *J* = 8.1 Hz, 1H, **e**), 7.42 (d, *J* = 8.0 Hz, 1H, **d**), 5.33 (s, 2H, **g**), 4.22 (t, *J* = 9.0 Hz, 2H, **h**), 3.38 (s, 3H, **f**), 2.76 (dd, *J* = 16.4, 4.0 Hz, 2H, **a**), 2.51 ppm (dd, *J* = 16.4, 6.7 Hz, 2H, **b**).
 **4a***	δ: 8.04 (d, *J* = 8.2 Hz, 1H, **e**), 7.38 (d, *J* = 8.1 Hz, 1H, **d**), 5.17 (s, 2H, **g**), 4.31 – 4.29 (m, 1H, **i**), 3.87 – 3.82 (m, 2H, **j**), 3.41 (s, 3H, **f**), 2.75 (dd, *J* = 16.5, 4.0 Hz, 2H, **a**), 2.51 (dd, *J* = 16.5, 6.6 Hz, 2H, **b**), 2.12 – 1.93 (m, 1H, **k**), 1.91 (m, 2H, **k**), 1.68 – 1.56 ppm (m, 1H, **k**). The peaks corresponding to **h** of **6a** could not be assigned because of peak broadening and overlapping with peaks of other assay components.

 **5a**	δ: 8.52 (s, 1H, **f**), 8.05 (d, *J* = 7.9 Hz, 1H, **e**), 7.44 (d, *J* = 7.9 Hz, 1H, **d**), 4.61 (m, 1H, **c**), 2.74 (dd, *J* = 16.4, 4.0 Hz, 2H, **a**), 2.49 ppm (dd, J = 16.4, 6.7 Hz, 2H, **b**).
 **6a***	δ: 2.79 (dd, *J* = 16.0, 4.1 Hz, 2H, **a**), 2.54 ppm (dd, *J* = 16.3, 6.6 Hz, 2H, **b**). The aromatic and methoxy peaks of **6a** were not assigned because of peak broadening and overlapping with substrate peaks.
 **10a**	δ: 4.26 (m, 2H, **b**), 3.00 (d, *J* = 16.3, 2H, **c**), 2.60 (d, *J* = 16.4, 2H, **c**), 2.33 (m, 1H, **d**), 1.28 (t, *J* = 7.3 Hz, 3H, **a**), 0.97 ppm (d, *J* = 8.1 Hz, 4H, **e**).

*The C5 proton of **1a, 3a, 4a,** and **6a** was not observed, likely because it coincides with the HOD peak; however, the splitting pattern of the C4/C6 proton signals implies that these protons couple with the C5 proton as observed for other βTHs (Supporting Figures S5 and S8).

6.12 Supporting Information References

1 Robert, X. & Gouet, P. Deciphering Key Features in Protein Structures with the New ENDscript Server. *Nucleic Acids Res.* **42**, W320–W324 (2014). https://doi.org/10.1093/nar/gku316

2 Madeira, F. *et al.* The EMBL-EBI Job Dispatcher Sequence Analysis Tools Framework in 2024. *Nucleic acids Res.* **52**, W521–W525 (2024). https://doi.org/10.1093/nar/gkae241

3 Letunic, I. & Bork, P. Interactive Tree of Life (iTOL) v6: Recent Updates to the Phylogenetic Tree Display and Annotation Tool. *Nucleic Acids Res.* **52**, W78–W82 (2024). https://doi.org/10.1093/nar/gkae268

4 Maeda, H. *et al.* A Rice Gene that Confers Broad-spectrum Resistance to β-Triketone Herbicides. *Science* **365**, 393–396 (2019). https://doi.org/doi:10.1126/science.aax0379

5 Dong, J. *et al.* An Artificially Evolved Gene for Herbicide-resistant Rice Breeding. *Proc. Natl. Acad. Sci. U. S. A.* **121**, e2407285121 (2024). https://doi.org/doi:10.1073/pnas.2407285121

6 Wu, J. *et al.* Directed Evolution of HIS1-Like Proteins to Enhance Herbicide Metabolism in Crops. *Adv. Agrochem.* **in press** (2025). https://doi.org/https://doi.org/10.1016/j.aac.2025.05.005

7 Viswanathan, T. S., Johnson, R. E. & Fisher, H. F. Alpha-Ketoglutaric Acid: Solution Structure and the Active Form for Reductive Amination by Bovine Liver Glutamate Dehydrogenase. *Biochemistry* **21**, 339–345 (1982). https://doi.org/10.1021/bi00531a022

8 Copper, A. J. & Redfield, A. G. Proton Magnetic Resonance Studies of Alpha-keto Acids. *J. Biol. Chem.* **250**, 527–532 (1975).

9 Sun, Y. D., S.; Allen, M. D.; Cheng, E. S. Z.; Zhang, Z.; Schofield, C. J.; Brewitz, L. . Structural and Functional Validation of *Pseudomonas savastanoi* Ethylene Forming Enzymes Reveals Flexibility in 2-Oxoglutarate Binding Mode and Conformation. . *Manuscript submitted.* (2025).

10 Dhingra, S. S., Y.; Zhang, Z.; Schofield, C. J.; Brewitz, L. . Expanding the Alkene Forming Capability of the *Pseudomonas savastanoi* Ethylene-forming Enzyme. . *Manuscript submitted.* (2025).

11 Alshref, F. *et al.* Structural and Inhibition studies on *Arabidopsis thaliana* HPPD. *Manuscript in preparation* (2025).

12 Kavana, M. & Moran, G. R. Interaction of (4-Hydroxyphenyl)pyruvate Dioxygenase with the Specific Inhibitor 2-[2-Nitro-4-(trifluoromethyl)benzoyl]-1,3-cyclohexanedione. *Biochemistry* **42**, 10238–10245 (2003). https://doi.org/10.1021/bi034658b

13 Bereziuk, O., Gubina, K., Trush, V. & Ovchynnikov, V. Crystal Structure of the Sodium Salt of Mesotrione: a Triketone Herbicide. *Acta Crystallogr. E.* **80**, 296–299 (2024). https://doi.org/doi:10.1107/S2056989024001439

14 Duff, S. M. G. *et al.* Structural and Functional Characterization of Triketone Dioxygenase from *Oryza Sativa*. *Biochem. Biophys. Res. Commun.* **1868**, 130504 (2024). https://doi.org/https://doi.org/10.1016/j.bbagen.2023.130504

15 Meepagala, K. M. & Schrader, K. K. Antibacterial Activity of Acylglucinol Derivatives against Flavobacterium columnare. *J. Agric. Chem. Environ.* **3**, 7 (2014). https://doi.org/10.4236/jacen.2014.33011

16 Popoola, O. K., Marnewick, J. L., Rautenbach, F., Iwuoha, E. I. & Hussein, A. A. Acylphloroglucinol Derivatives from the South African *Helichrysum niveum* and Their Biological Activities. *Molecules* **20**, 17309–17324 (2015). https://doi.org/10.3390/molecules200917309

17 Rahman, M. M., Shiu, W. K. P., Gibbons, S. & Malkinson, J. P. Total synthesis of acylphloroglucinols and their antibacterial activities against clinical isolates of multi-drug

resistant (MDR) and methicillin-resistant strains of *Staphylococcus aureus*. *Eur. J. Med. Chem.* **155**, 255–262 (2018). https://doi.org/https://doi.org/10.1016/j.ejmech.2018.05.038


18 Dhingra, S., Zhang, Z., Lohans, C. T., Brewitz, L. & Schofield, C. J. Substitution of 2-Oxoglutarate Alters Reaction Outcomes of the *Pseudomonas savastanoi* Ethylene-forming Enzyme. *J. Biol. Chem.* **300**, 107546 (2024). https://doi.org/https://doi.org/10.1016/j.jbc.2024.107546

19 Aik, W. S. *et al.* in *2-Oxoglutarate-Dependent Oxygenases* (eds Christopher Schofield & Robert Hausinger) 0 (The Royal Society of Chemistry, 2015).

6.13 Statement of Authorship

Title of Paper	**Crystallographic and Biochemical Characterisation of HIS1 and OsHSL1 Inform on Strategies to Evade Resistance-Associated β-Triketone Herbicide Metabolism**
Publication Status	□ Published □ Accepted for Publication □Submitted for Publication X Unpublished and unsubmitted work written in a manuscript style
Publication Details	Dhingra, S.; Sharma, S.; Farcas, I.M.; Alshref, F. M.; Allen, M. D.; Redfield, C.; Schofield, C. J.; Brewitz, L. Crystallographic and biochemical characterisation of HIS1 and OsHSL1 inform on strategies to evade resistance-associated β-triketone herbicide metabolism. *Manuscript in preparation.*

Student Confirmation

Student Name:	Siddhant Dhingra		
Contribution to the Paper	I conducted all the protein purification and NMR assays described here alongside Ms. Sakshi Sharma and Ms. Ilinca Farcas, under my laboratory supervision. Ms. Ilinca Farcas and I crystallised HIS1 and OsHSL1 with help from Mr. Faisal Alshref. Dr. Mark Allen solved the crystal structures. I conducted all the data analysis, and data visualisation described here. Prof. Christina Redfield helped with the analysis of the NMR spectra. The manuscript was prepared by myself and Dr. Lennart Brewitz under the guidance of Prof. Christopher J. Schofield. Dr. Lennart Brewitz, myself, and Prof. Christopher J. Schofield conceptualised the project.		
Signature 		Date	05-10-2025

Supervisor Confirmation

By signing the Statement of Authorship, you are certifying that the candidate made a substantial contribution to the publication, and that the description described above is accurate.

Supervisor name and title: Prof. Christopher J. Schofield		
Supervisor comments I'm happy to state that the candidate made the major and a very substantial contribution to the publication, and that the description described above is accurate.		
Signature C. J. Schofield	Date	25th September 2025

CHAPTER 7

CONCLUSIONS AND FUTURE PERSPECTIVES

Contents

7.1 Catalytic versatility of 2OG oxygenases **438**

7.2 PsEFE: bifurcating catalysis, substrate scope, and engineering new reactivity **440**

7.2.1 Future directions **442**

7.3 KMBA and KEBA as ethylene precursors **442**

7.3.1 Future directions **447**

7.4 HIS1 and OsHSL1: catalytic promiscuity - a herbicide resistance challenge **447**

7.4.1 Future directions **448**

7.5 Concluding Remarks **449**

7.6 References **449**

7.1 Catalytic versatility of 2OG oxygenases

Fe(II)/2OG-dependent oxygenases are one of the most versatile enzyme superfamilies found in nature.[1] Despite having conserved features, such as, a distorted DSBH core fold and the Fe(II)-binding facial triad, they catalyse a diverse range of oxidative reactions. They have roles in primary metabolism, signalling, and natural product biosynthesis (Section 1.6.1).[1-3]

2OG oxygenases are being increasingly explored for their industrial applications, because of their diverse catalytic abilities and high regio- and stereoselectivity[4] (Figure 7.1). Prolyl hydroxylases are already employed as biocatalysts by companies, such as Kyowa Hakko Kogyo[5] and API Corp[6], for the large-scale manufacturing of hydroxyproline[5] or hydroxypipecolic acid derivatives,[6] which are important building blocks for pharmaceuticals[4]. 2OG-dependent amino acid hydroxylases, including leucine dioxygenase, GriE, have been engineered for producing hydroxylated amino acids;[4] GriE was optimised for producing hydroxyleucine, an intermediate for the synthesis of antibiotics, such as, manzacidin C[7] and cavinafungin[8]. Clavaminate synthase and deacetoxycephalosporin synthase (DAOCS) are also widely studied 2OG oxygenases involved in the biosynthesis of clavulanic acid, a β-lactamase inhibitor, and cephalosporins, respectively.[4,9,10] The catalytic plasticity of 2OG oxygenases is, therefore, a source of opportunity, in the form of biocatalytic innovation (Chapter 2-5), but may also pose challenges to human intervention in the form of resistance in agriculture (Chapter 6) and medicine.




Figure 7.1. The catalytic versatility of 2OG oxygenases can be harnessed for biocatalytic applications, including functionalisation of amino acids and synthesis of antibiotics.[4] (**a**) Kyowa Hakko Kogyo Ltd. has patented several prolyl hydroxylase-based processes for the synthesis of 3-hydroxy-*L*-proline and 4-hydroxy-*L*-proline; cis-P3H1 converts *L*-proline into *cis*-3-hydroxy-*L*-proline and SmP4H and DaP4H produce cis- and trans-4-hydroxy-*L*-proline respectively.[4,5] (**b**) API Corp. has used a prolyl hydroxylase X5PH to produce cis-5-hydroxypipecolic acid from pipecolic acid; note that cis-P3H1, SmP4H, and DaP4H have also been engineered for the large-scale synthesis of hydroxypipecolic acid derivatives.[4,6] (**c**) GriE is used for the chemo-enzymatic synthesis of hydroxyleucine, a precursor to antibiotics, such as, manzacidin C[7] and cavinafungin[4,8]. (**d**) Clavulanic acid, a commercially available β-lactamase inhibitor, can be produced by a trifunctional 2OG oxygenase, clavaminate synthase (CAS), from *Streptomyces clavuligerus*.[4,9,10] (**e**) β-Lactam antibiotic biosynthesis employs 2OG oxygenases; the ring expansion of penicillin N, is mediated by deacetoxycephalosporin synthase (DAOCS), a key step in the biosynthesis of cephalosporins.[4] Efforts to engineer DAOCS variants, that can accept the more abundantly available penicillin G as a substrate, have shown promise for improving the efficiency for large-scale cephalosporin production.[11]

The work described in this thesis concerned investigations on two complementary aspects of the catalytic versatility of 2OG oxygenases, one from bacteria (PsEFE) and one from plants (HIS1). PsEFE, the *Pseudomonas savastanoi* pv. *phaseolicola* PK2 ethylene-forming enzyme, catalyses a unique bifurcating reaction of 2OG producing both ethylene and succinate in the presence of *L*-arginine (Section 1.7).[12-19] PsEFE was investigated because of its interesting mechanism and its potential for biocatalysis. Work on HIS1 and OsHSL1, rice enzymes conferring resistance to β-triketone herbicides,[20-23] aimed to explore the molecular basis of herbicide resistance to guide herbicide and transgenic crop development (Chapter 6). Together, these studies further highlight the catalytic promiscuity of 2OG oxygenases; in PsEFE, it represents an opportunity for biocatalytic production of relevant chemicals, including ethylene, while in HIS1/OsHSL1, it highlights emerging challenges in crop protection and food security.

7.2 PsEFE: bifurcating catalysis, substrate scope, and engineering new reactivity

To date, PsEFE catalysed conversion of 2OG to ethylene is unusual to biology; its chemistry partitions between two competing pathways: fragmentation of 2OG to ethylene and CO_2 or the oxidative decarboxylation of 2OG to succinate, coupled to the hydroxylation of *L*-arginine to P5C and guanidine.[12-19] The ratio between these outcomes is not only sensitive to the reaction conditions (Figure 2.2) but also varies amongst ethylene-forming enzymes from different strains (Figure 3.3). Notably, other members of the 2OG oxygenase superfamily, such as deacetoxycephalosporin C synthase (DAOCS), also show bifurcating reactivity.[9] DAOCS catalyses the transformation of penicillin N to deacetoxycephalosporin C (DAOC), following ring expansion; the reaction is proposed to bifurcate at the penultimate step, producing 3β-hydroxy cepham, as a side product.[9]

A novel reaction pathway replacing the alkene-forming reaction, unobserved for 2OG, was discovered for the reaction of PsEFE with substituted 2OG derivatives by competitive studies published during my DPhil studies.[24] This involves the formation of a substituted 3-hydroxypropionate derivative (alcohol) and succinate product (diacid) in the presence of *L*-arginine (Figure 1.14).[15,24] Systematic substrate scope studies with >30 C3- and C4-substituted 2OG derivatives, discussed in Chapter 2, showed that steric and electronic features of the 2-oxoacid (co)substrate can influence catalytic efficiency and product partitioning between competing reaction pathways.[7] Bulky or cyclic substituents were found to suppress the alcohol product formation and favour diacid/P5C formation (Figure 7.2, Figure 2.8); highlighting the substrate-dependent tunability of PsEFE catalysis.

Importantly, several naturally-occurring 2OG derivatives, such as, 3-methyl-2OG and 4-hydroxy-2OG, were identified as alternative PsEFE substrates,[15,24] highlighting its potential involvement in other metabolic pathways beyond ethylene formation in bacteria. For example, 3-hydroxybutyrate, formed from 3-methyl-2OG, is a known metabolite in *Pseudomonas* involved in polyester biosynthesis.[25] It is possible that other naturally-occurring 2OG derivatives may be the preferred (co)substrates of some 2OG oxygenases or that competition between 2OG and 2OG derivatives regulates catalysis in cells.[26]

Work on PsEFE orthologues from other *P. savastanoi* strains, *glycinea* and 1449B, highlighted the conservation of *efe* across different strains (Chapter 3). *Glycinea* PsEFE bears a single surface-residue mutation (R326S) while 1449B PsEFE has 28 substitutions, distal from the active site, and 14 deletions at the C-terminus of the PK2 PsEFE (Chapter 3). Both the *glycinea* and 1449B PsEFE were found to be less efficient than PK2 PsEFE in catalysing ethylene formation, at least under the tested conditions, but differed in the ratios of ethylene and succinate produced. 1449B PsEFE shifted the reaction flux towards ethylene formation while *glycinea* PsEFE produced ethylene and succinate from 2OG in approximately equal amounts, similar to as observed for PK2 PsEFE. This product partitioning shows that subtle sequence variations, despite being distal from the active-site, can bias reaction partitioning.

A crystal structure of 1449B PsEFE in complex with Mn(II) and 2OG displayed an atypical inverted binding mode of 2OG and a conserved extensive water network, similar to PK2 PsEFE (Chapter 3). These results, together with reported work on the *Penicillium digitatum* ethylene-forming enzyme,[27] show that naturally-occurring or evolved EFE variants might preferentially catalyse ethylene formation over arginine hydroxylation, an attractive avenue for biocatalysis.

Building on the substrate scope studies on PK2 PsEFE, mutagenesis at Leu206 offered a promising avenue for engineering new PsEFE variants capable of producing higher alkenes from substituted 2OG derivatives.[28] (Chapter 4). The L206V and L206A 'bump and hole' variants can accommodate the additional steric bulk of the C4-substituted 2OG derivatives and catalyse the formation of higher alkene products, including propylene, 1-butylene, and acetaldehyde from 4-methyl-2OG, 4-ethyl-2OG, and 4-hydroxy-2OG respectively (Section 4.3). This work forms one of the first studies[28] demonstrating that PsEFE can be engineered to produce higher alkenes beyond ethylene (Figure 7.2), similar to ACC oxidase[29-33], the ethylene-forming enzyme in plants. This work mirrors the mutagenesis studies aimed at engineering

nitrogenases capable of producing higher hydrocarbons from alkynes (Section 1.5).[34,35] A summary of PsEFE and PsEFE variants-catalysed reactions is shown in Figure 7.2.

7.2.1 Future directions

The results described in Chapters 2-4 strengthen the case for further developing PsEFE for biocatalysis. Structural studies of alternative substrates and PsEFE variants may be useful for expanding the mechanistic understanding of PsEFE catalysis, specifically for understanding how radical intermediates are stabilised and reaction partitioning is governed.[36] Directed evolution on PsEFE may help identify variants with improved efficiency and selectivity, such as for engineering 'ethylene-only' enzymes for sustainable ethylene production. PsEFE may provide a versatile and engineerable scaffold for biocatalysis, capable of enabling the integration of engineered PsEFE variants into microbial systems for the large-scale production of industrially and medicinally relevant small molecules, following optimisation of its catalytic efficiency and product partitioning.

7.3 KMBA and KEBA as ethylene precursors

Apart from 2OG and ACC, other small molecules have also been implicated in ethylene production (Figure 7.3). β-Alanine has been reported to produce ethylene in cellular extracts containing ATP/CoA/PLP/Mg^{2+},[37] although the precise chemistry of this conversion remains ambiguous and may come from non-enzymatic fragmentation of β-alanine. Beyond metabolites, acetylene can also be non-enzymatically reduced to ethylene using transition-metal catalysts;[38] Mo- and V-dependent nitrogenases are also known to catalyse ethylene production from acetylene.[39] KMBA has also been shown to produce ethylene in bacteria and fungi;[40-42] however, pioneering studies have shown that KMBA oxidation to ethylene may occur non-enzymatically in the presence of peroxide or metal catalysts (Section 1.4).[43] However, no enzyme capable of directly converting KMBA to ethylene has yet been characterised.[18] Notably, KMBA has been mentioned to produce ethylene widely in biological literature over the years,[40-42,44] but the biochemical characterisation of this reaction has not been studied in the recent years.

An important outcome of the work described in this thesis was the demonstration that PsEFE and its variants can accept hydrophobic 2-oxoacids, lacking the distal carboxylate moiety of 2OG/2OG derivatives, as alternative (co)substrates. wt PsEFE catalyses the conversion of 2-

oxoacids, including KMBA, to their acid and alcohol products similar to 2OG derivatives (Section 5.2). The uptake of hydrophobic 2-oxoacids by PsEFE suggests that 2OG oxygenases may have context-dependent functions where they may utilise the cellular 2-oxoacid pool for signalling and/or secondary metabolism, because of the relatively high abundance of hydrophobic 2-oxoacids in cells.[45] Other 2-oxoacid dependent dioxygenases, like HPPD, accept hydrophobic 2-oxoacids as their substrates.[46,47]

R277 PsEFE variants unlocked a new reactivity, the direct conversion of KMBA into ethylene, showing for the first-time the validation of KMBA as a direct ethylene precursor (Figure 7.3, Figure 5.7). Additionally, the work also identified the *S*-ethyl substituted KMBA, KEBA, as a novel small-molecule ethylene precursor, extending the repertoire of ethylene precursors in nature.




Figure 7.2. Summary of the PsEFE- and PsEFE variant-catalysed reactions of: a) 2OG and 2OG derivatives and b) hydrophobic 2-oxoacids. (a) wt PsEFE and L206 PsEFE variants catalyse the oxidation of 2OG and C3- or C4-substituted 2OG derivatives, in the presence of *L*-arginine, to give diacid, alcohol, and in some cases ethylene or other alkene products; the alkene products and their corresponding precursor and enzyme-variant pair are highlighted in blue. **(b)** wt PsEFE and R277 PsEFE variants catalyse the oxidation of hydrophobic 2-oxoacids, in the presence of *L*-arginine, to give an acid and alcohol product; in case of KMBA and KEBA, R277 PsEFE variants catalyse ethylene and methanesulphonic acid or ethanesulphonic acid formation respectively.

Disruption of the R277 salt-bridge in PsEFE, which interacts with the C5 carboxylate of 2OG in R277A/F/K/L PsEFE, ablates PsEFE activity with 2OG and its derivatives. Notably, the

designed R277 PsEFE variants are still able to productively bind to hydrophobic 2-oxoacids, producing both acid and alcohol products, reminiscent of the activity of wt PsEFE with 2-oxoacids and 2OG derivatives. However, only KMBA and KEBA were found to manifest ethylene formation with the R277 PsEFE variants, indicating the importance of the terminal *S* atom for reactivity. The diverging reactivity of R277 PsEFE variants from that of wt PsEFE parallels with the catalytic activity of human oxygenases having disease-associated mutations at equivalent positions.[48,49] Note that, other sulphur containing molecules, such as the reaction products of KMBA with PsEFE, (2-(methylthio)ethanol and 3-(methylthio)propanoic acid), did not produce ethylene under the reaction conditions.




Figure 7.3. Reported and newly validated small-molecule precursors (shown in blue) that can be enzymatically converted to ethylene. These substrates illustrate different biochemical solutions to ethylene biosynthesis employing varied enzymes (shown in red), organisms, and metabolic contexts (shown in black); small molecules which can produce ethylene non-enzymatically are shown in a dashed box. **(a)** 2OG is converted to ethylene in *Pseudomonas* spp. and *Penicillium* spp. by ethylene-forming enzymes in the presence of *L*-arginine;[12-19,27] nitrogenase-like enzymes of the MarBHDK gene cluster in *R. rubrum* produce ethylene in the absence of oxygen using 2-(methylthio)ethanol;[50-52] Mo- and V-dependent nitrogenases can

convert acetylene to ethylene;[39,51,53] higher plants produce ethylene from ACC using ACC-oxidase;[18,54,55] R277A/F/K/L PsEFE variants can catalyse ethylene formation from KEBA and KMBA in the presence of *L*-arginine. Note that, there is also the potential of non-enzymatic ethylene production from KMBA[43] and acetylene[38] (highlighted in a dashed box). (**b**) β-Alanine has been reported as an ethylene precursor in subcellular extracts of bean cotyledons; the conversion of β-alanine may also proceed non-enzymatically via malonatesemialdehyde and acrylate.[37]

From an evolutionary perspective, ethylene formation from KMBA draws a link between the ethylene production machinery in plants and bacteria. KMBA is a natural intermediate of the Yang cycle, which links methionine metabolism to ethylene production via ACC in plants (Figure 1.3). The evidence that KMBA can itself be enzymatically converted to ethylene suggests that ethylene biosynthesis may have stemmed from methionine or methionine-derived intermediates and later evolved to ACC, a specialised ethylene precursor. The convergence of aerobic microbial EFE reactivity with KMBA and the anaerobic nitrogenase-like enzymes acting on methionol, one of the PsEFE reaction products of KMBA, to produce ethylene, suggests common ancestry (Figure 7.4).




Figure 7.4. Methionine (red) is the central building block for the biosynthesis of known ethylene precursors (blue) including via the Yang cycle. Note that, KMBA can be oxidised non-enzymatically to produce ethylene.[43] The complete Yang cycle is shown and discussed in Figure 1.3.

7.3.1 Future directions

KMBA and KEBA, potentially, offer a renewable, methionine-derived route to ethylene. The wasteful reaction flux to acid/alcohol formation could be minimised by enzyme engineering, using directed evolution, for sustainable ethylene biosynthesis. Screening putative ethylene-forming enzymes across different biological kingdoms, including microbes and plants, may reveal additional enzymes capable of naturally catalysing this reaction; this would also help in understanding the position of PsEFE within a broader evolutionary landscape. Comparative studies with other oxygenases that utilise hydrophobic 2-oxoacids, such as HPPD, as their natural substrates, can help elucidate the catalytic promiscuity and its mechanism across the superfamily.

7.4 HIS1 and OsHSL1: catalytic promiscuity - a herbicide resistance challenge

In contrast to the potential of PsEFE as a biocatalyst, 2OG oxygenases, including HIS1 and OsHSL1, highlight the potential challenges posed by catalytic promiscuity that can undermine the efficacy of agrochemicals.[20] HIS1 and OsHSL1 confer resistance to β-triketone herbicides, such as benzobicyclon and mesotrione, which are reported HPPD inhibitors and widely employed in agriculture for weed management.[20,21,56-59] The detoxified metabolites of HIS1 and OsHSL1 have reduced HPPD inhibition and, consequently, reduced efficacy.[21] Notably, other 2OG oxygenases, like RdpA and SdpA, found in *Sphingobium herbicidovorans* and *Delftia acidovorans*, are also implicated in herbicide resistance; RdpA and SdpA oxidise herbicides (*S*)- and (*R*)-dichlorprop respectively, rendering them inactive.[60]

Despite ~87% sequence identity, HIS1 and OsHSL1 display divergent reaction profiles with triketone herbicides (Section 6.1);[20-23] HIS1 has a significantly broader substrate scope than OsHSL1 suggesting that it can manifest herbicide resistance to a broad range of herbicides in rice and potentially be passed on across species as a resistance mechanism.




Figure 7.5. Reactions catalysed by 2OG-dependent oxygenases identified to be involved in herbicide degradation.[20,56-61] **a)** RdpA and SdpA, 2OG oxygenases found in *Sphingobium herbicidovorans* and *Delftia acidovorans*, facilitate the oxidative cleavage of (*R,S*)-dichloroprop.[60] **b) H**PPD **I**nhibitor **S**ensitive 1 or HIS1 confers resistance to βTHs in *Oryza sativa*; benzobicyclon (prodrug) is metabolised to benzobicyclon-OH (BBC-OH), a reported HPPD inhibitor, which is deactivated by HIS1-catalysed hydroxylation.[20,56-59]

The structural work discussed in Section 6.4.4 identifies residues that are indispensable for the catalytic activity of HIS1. Coupled with complementary independent studies published during the course of my DPhil on the reverse engineering of OsHSL1 to encode for these residues, supports this hypothesis.[20,56-59] This underscores the need for detailed structural and biochemical analyses of HIS1-like proteins. These findings can inform on rational strategies for crop development and engineering herbicide resistance traits into transgenic crops. The structure-activity relationships drawn from the work can also guide the design of herbicides resilient to HIS1-mediated inactivation, helping in the management of the emergence of resistance in weeds.

7.4.1 Future directions

The structural and biochemical information on HIS1/OsHSL1 catalysis provides a foundation for HIS1/OsHSL1 inhibitor development and for designing next-generation herbicides that are resilient to degradation by herbicide resistance enzymes. Expanding these studies to other HSL-like proteins, across crop and weed species, will be important for understanding the evolution of resistance patterns, developing strategies to harness or surpass them, and maintaining herbicide efficacy. The resistance can also be deployed to stack resistance traits selectively and minimise cross-resistance by allowing for the sustainable and selective application of

herbicides. This would have implications not only in the development of transgenic crops but also, broadly, in ensuring food security.

7.5 Concluding Remarks

The research described here has advanced the biochemical understanding and applied perspectives on bacterial and plant 2OG oxygenases. PsEFE has been shown to be an engineerable scaffold with a broad substrate scope, substrate or variant-dependent tuning of product partitioning, and the capacity to produce alkene products other than ethylene from suitable precursors. The discovery that enzyme-catalysed ethylene production from KMBA and KEBA is possible, connects microbial ethylene-forming machinery to methionine metabolism in plants and offers evolutionary insights into ethylene biosynthesis. Structural studies on herbicide resistance enzymes, HIS1 and OsHSL1, illustrate how subtle structural differences govern catalysis influencing herbicide metabolism, impacting modern agriculture.

Together, the results described in this thesis show that from a human perspective that the catalytic promiscuity of 2OG oxygenases can be a 'double-edged sword'. It can be harnessed for biotechnology and biocatalysis, as may be in the case of PsEFE as has been previously done for prolyl hydroxylases,[5,6] but also presents challenges in the form of resistance, as with HIS1 and OsHSL1. This shows the importance of fundamental enzymology for not only unlocking opportunities but also for addressing global challenges in sustainability and food security.

7.6 References

1 *2-Oxoglutarate-Dependent Oxygenases*. (The Royal Society of Chemistry, 2015).

2 Clifton, I. J. *et al.* Structural Studies on 2-Oxoglutarate Oxygenases and Related Double-stranded β-helix Fold Proteins. *J. Inorg. Biochem.* **100**, 644–669 (2006). https://doi.org/https://doi.org/10.1016/j.jinorgbio.2006.01.024

3 Martinez, S. & Hausinger, R. P. Catalytic Mechanisms of Fe(II)- and 2-Oxoglutarate-dependent Oxygenases. *J. Biol. Chem.* **290**, 20702–20711 (2015). https://doi.org/10.1074/jbc.R115.648691

4 Peters, C. & Buller, R. M. Industrial Application of 2-Oxoglutarate-Dependent Oxygenases. *Catalysts* **9**, 221 (2019).

5 Ozaki, A. *et al.* (Google Patents, 1999).

6 Miyake, R. & Dekishima, Y. (Google Patents, 2018).

7 Zwick, C. R., 3rd & Renata, H. Evolution of Biocatalytic and Chemocatalytic C-H Functionalization Strategy in the Synthesis of Manzacidin C. *J. Org. Chem.* **83**, 7407–7415 (2018). https://doi.org/10.1021/acs.joc.8b00248

8 Zwick, C. R. & Renata, H. A One-pot Chemoenzymatic Synthesis of (2S, 4R)-4-Methylproline Enables the First Total Synthesis of Antiviral Llipopeptide Cavinafungin B. *Tetrahedron* **74**, 6469–6473 (2018). https://doi.org/https://doi.org/10.1016/j.tet.2018.09.046

9 Foster, F. *Substrate Analogue Studies on Cephalosporin Biosynthesis Enzymes*, University of Oxford, (2022).

10 Ser, H. L. *et al.* Fermentation Conditions that Affect Clavulanic Acid Production in *Streptomyces clavuligerus*: A Systematic Review. *Front. Microbiol.* **7**, 522 (2016). https://doi.org/10.3389/fmicb.2016.00522

11 Ji, J. *et al.* Iterative Combinatorial Mutagenesis as an Effective Strategy for Generation of Deacetoxycephalosporin C Synthase with Improved Activity Toward Penicillin G. *Appl. Environ. Microbiol.* **78**, 7809–7812 (2012). https://doi.org/10.1128/aem.02122-12

12 Johansson, N., Persson, K. O., Larsson, C. & Norbeck, J. Comparative Sequence Analysis and Mutagenesis of Ethylene Forming Enzyme (EFE) 2-Oxoglutarate/Fe(II)-dependent Dioxygenase Homologs. *BMC Biochem.* **15**, 22–29 (2014). https://doi.org/10.1186/1471-2091-15-22

13 Xue, J., Lu, J. & Lai, W. Mechanistic Insights into a Non-heme 2-Oxoglutarate-dependent Ethylene-forming Enzyme: Selectivity of Ethylene-formation versus L-Arg Hydroxylation. *Phys. Chem. Chem. Phys.* **21**, 9957–9968 (2019). https://doi.org/10.1039/C9CP00794F

14 Cui, Y. *et al.* Discovery of Five New Ethylene-Forming Enzymes for Clean Production of Ethylene in *E. coli*. *Int. J. Mol. Sci.* **23**, 4500–4515 (2022).

15 Dhingra, S., Zhang, Z., Lohans, C. T., Brewitz, L. & Schofield, C. J. Substitution of 2-Oxoglutarate Alters Reaction Outcomes of the *Pseudomonas savastanoi* Ethylene-forming Enzyme. *J. Biol. Chem.* **300**, 107546 (2024). https://doi.org/https://doi.org/10.1016/j.jbc.2024.107546

16 Martinez, S. & Hausinger, R. P. Biochemical and Spectroscopic Characterization of the Non-Heme Fe(II)- and 2-Oxoglutarate-Dependent Ethylene-Forming Enzyme from *Pseudomonas syringae* pv. *phaseolicola* PK2. *Biochemistry* **55**, 5989–5999 (2016). https://doi.org/10.1021/acs.biochem.6b00890

17 Zhang, Z. *et al.* Structural and Stereoelectronic Insights into Oxygenase-catalyzed Formation of Ethylene from 2-Oxoglutarate. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 4667–4672 (2017). https://doi.org/10.1073/pnas.1617760114

18 Hausinger, R. P. *et al.* Biological Formation of Ethylene. *RSC Chem. Biol.* **4**, 635–646 (2023). https://doi.org/10.1039/D3CB00066D

19 Walewska, A., Szewczyk, A. & Koprowski, P. Gas Signaling Molecules and Mitochondrial Potassium Channels. *Int. J. Mol. Sci.* **19**, 3227–3247 (2018). https://doi.org/10.3390/ijms19103227

20 Maeda, H. *et al.* A Rice Gene that Confers Broad-spectrum Resistance to β-Triketone Herbicides. *Science* **365**, 393–396 (2019). https://doi.org/doi:10.1126/science.aax0379

21 Dong, J. *et al.* An Artificially Evolved Gene for Herbicide-resistant Rice Breeding. *Proc. Natl. Acad. Sci. U. S. A.* **121**, e2407285121 (2024). https://doi.org/doi:10.1073/pnas.2407285121

22 Wu, J. *et al.* Directed Evolution of HIS1-Like Proteins to Enhance Herbicide Metabolism in Crops. *Adv. Agrochem.* **in press** (2025). https://doi.org/https://doi.org/10.1016/j.aac.2025.05.005

23 Takamura, N. *et al.* Catalytic Promiscuity of Rice 2-Oxoglutarate/Fe(II)-dependent Dioxygenases Supports Xenobiotic Metabolism. *Plant Physiol.* **187**, 816–828 (2021). https://doi.org/10.1093/plphys/kiab293

24 Copeland, R. A. *et al.* Hybrid Radical-polar Pathway for Excision of Ethylene from 2-Oxoglutarate by an Iron Oxygenase. *Science* **373**, 1489–1493 (2021). https://doi.org/doi:10.1126/science.abj4290

25 Verlinden, R. A., Hill, D. J., Kenward, M. A., Williams, C. D. & Radecka, I. Bacterial synthesis of biodegradable polyhydroxyalkanoates. *J Appl Microbiol* **102**, 1437–1449 (2007). https://doi.org/10.1111/j.1365-2672.2007.03335.x

26 Tiwari, A. D. *et al.* SAR Insights into TET2 Catalytic Domain Inhibition: Synthesis of 2-Hydroxy-4-Methylene-pentanedicarboxylates. *Bioorg. Med. Chem.* **39**, 116141 (2021). https://doi.org/https://doi.org/10.1016/j.bmc.2021.116141

27 Chatterjee, S. *et al.* Biochemical, Structural, and Conformational Characterization of a Fungal Ethylene-Forming Enzyme. *Biochemistry* **64**, 2054–2067 (2025). https://doi.org/10.1021/acs.biochem.5c00038

28 Burke, E. J., Copeland, R. A., Dixit, Y., Krebs, C. & Bollinger, J. M., Jr. Steric Perturbation of the Grob-like Final Step of Ethylene-Forming Enzyme Enables 3-Hydroxypropionate and Propylene Production. *J. Am. Chem. Soc.* **146**, 1977–1983 (2024). https://doi.org/10.1021/jacs.3c09733

29 Khan, S., Alvi, A. F., Saify, S., Iqbal, N. & Khan, N. A. The Ethylene Biosynthetic Enzymes, 1-Aminocyclopropane-1-Carboxylate (ACC) Synthase (ACS) and ACC Oxidase (ACO): The Less Explored Players in Abiotic Stress Tolerance. *Biomolecules* **14**, 90–123 (2024). https://doi.org/10.3390/biom14010090

30 Dilley, D. R. *et al.* 1-Aminocyclopropane-1-carboxylic Acid Oxidase Reaction Mechanism and Putative Post-translational Activities of the ACCO Protein. *AoB Plants* **5**, plt031 (2013). https://doi.org/10.1093/aobpla/plt031

31 McGarvey, D. J. & Christoffersen, R. E. Characterization and Kinetic Parameters of Ethylene-forming Enzyme from Avocado Fruit. *J. Biol. Chem.* **267**, 5964–5967 (1992). https://doi.org/https://doi.org/10.1016/S0021-9258(18)42649-X

32 Fernández-Maculet, J. C. & Yang, S. F. Extraction and Partial Characterization of the Ethylene-forming Enzyme from Apple Fruit. *Plant Physiol.* **99**, 751–754 (1992). https://doi.org/10.1104/pp.99.2.751

33 Zhang, Z., Schofield, C. J., Baldwin, J. E., Thomas, P. & John, P. Expression, Purification and Characterization of 1-Aminocyclopropane-1-carboxylate Oxidase from Tomato in *Escherichia coli*. *Biochem. J.* **307 (Pt 1)**, 77–85 (1995). https://doi.org/10.1042/bj3070077

34 Mayer, S. M., Niehaus, W. G. & Dean, D. R. Reduction of short chain alkynes by a nitrogenase α-70Ala-substituted MoFe protein. *J. Chem. Soc., Dalton Trans.*, 802–807 (2002). https://doi.org/10.1039/B107336B

35 Dos Santos, P. C., Mayer, S. M., Barney, B. M., Seefeldt, L. C. & Dean, D. R. Alkyne substrate interaction within the nitrogenase MoFe protein. *J. Inorg. Biochem.* **101**, 1642–1648 (2007). https://doi.org/https://doi.org/10.1016/j.jinorgbio.2007.05.007

36 Jaber Sathik Rifayee, S. B., Thomas, M. G. & Christov, C. Z. Revealing the Nature of the Second Branch Point in the Catalytic Mechanism of the Fe(II)/2OG-dependent Ethylene Forming Enzyme. *Chem. Sci.* **16**, 7667–7684 (2025). https://doi.org/10.1039/D4SC08378D

37 Stinson, R. A. & Spencer, M. beta-Alanine as an Ethylene Precursor. Investigations Towards Preparation, and Properties, of a Soluble Enzyme System From a Subcellular Particulate Fraction of Bean Cotyledons. *Plant Physiol.* **44**, 1217–1226 (1969). https://doi.org/10.1104/pp.44.9.1217

38 Xie, K. *et al.* Catalysts for Selective Hydrogenation of Acetylene: A Review. *Mater. Today Catal.* **3**, 100029 (2023). https://doi.org/https://doi.org/10.1016/j.mtcata.2023.100029

39 Seefeldt, L. C. *et al.* Reduction of Substrates by Nitrogenases. *Chem. Rev.* **120**, 5082–5106 (2020). https://doi.org/10.1021/acs.chemrev.9b00556

40 Chagué, V., Elad, Y., Barakat, R., Tudzynski, P. & Sharon, A. Ethylene Biosynthesis in *Botrytis cinerea*. *FEMS Microbiol. Ecol.* **40**, 143–149 (2002). https://doi.org/10.1111/j.1574-6941.2002.tb00946.x

41 Fukuda, H., Takahashi, M., Fujii, T., Tazaki, M. & Ogawa, T. An NADH : Fe(III):EDTA Oxidoreductase from *Cryptococcus albidus:* an Enzyme involved in Ethylene Production *in vivo*? *FEMS Microbiol. Lett.* **60**, 107–111 (1989).

42 Yang, J. *et al.* An Insight on the Ethylene Biosynthetic Pathway of Two Major Fruit Postharvest Pathogens with different Host Specificity: P*enicillium digitatum* and *Penicillium expansum*. *Eur. J. Plant Pathol.* **149**, 575–585 (2017). https://doi.org/10.1007/s10658-017-1205-x

43 Fukuda, H., Fujii, T. & Ogawa, T. Preparation of a Cell-free Ethylene-forming System from *Penicillium digitatum*. *Agric. Biol. Chem.* **50**, 977–981 (1986). https://doi.org/10.1271/bbb1961.50.977

44 Ballester, A. R. & González-Candelas, L. EFE-Mediated Ethylene Synthesis Is the Major Pathway in the Citrus Postharvest Pathogen *Penicillium digitatum* during Fruit Infection. *J. Fungi* **6**, 175–192 (2020). https://doi.org/10.3390/jof6030175

45 Bröer, S. & Gauthier-Coles, G. Amino Acid Homeostasis in Mammalian Cells with a Focus on Amino Acid Transport. *J Nutr.* **152**, 16–28 (2022). https://doi.org/https://doi.org/10.1093/jn/nxab342

46 Crouch, N. P., Adlington, R. M., Baldwin, J. E., Lee, M. H. & MacKinnon, C. H. A Mechanistic Rationalisation for the Substrate Specificity of Recombinant Mammalian 4-Hydroxyphenylpyruvate Dioxygenase (4-HPPD). *Tetrahedron* **53**, 6993–7010 (1997). https://doi.org/https://doi.org/10.1016/S0040-4020(97)00398-0

47 Adlington, R. M., Baldwin, J. E., Crouch, N. P., Lee, M.-H. & MacKinnon, C. H. Identification and Stereochemistry of the Product of 4-HPPD Catalyzed Oxidation of the Ketoacid of Methionine. *Bioorg. Med. Chem. Lett.* **6**, 2003–2006 (1996). https://doi.org/https://doi.org/10.1016/0960-894X(96)00362-9

48 Pfeffer, I. *et al.* Aspartate/asparagine-β-hydroxylase crystal structures reveal an unexpected epidermal growth factor-like domain substrate disulfide pattern. *Nature Communications* **10**, 4910 (2019). https://doi.org/10.1038/s41467-019-12711-7

49 McDonough, M. A. *et al.* Structure of human phytanoyl-CoA 2-hydroxylase identifies molecular mechanisms of Refsum disease. *J Biol Chem* **280**, 41101–41110 (2005). https://doi.org/10.1074/jbc.M507528200

50 Lago-Maciel, A. *et al.* Methylthio-alkane Reductases use Nitrogenase Metalloclusters for Carbon-sulfur Bond Cleavage. *bioRxiv* (2024). https://doi.org/10.1101/2024.10.19.619033

51 North, J. A. *et al.* A Nitrogenase-like Enzyme System Catalyzes Methionine, Ethylene, and Methane Biogenesis. *Science* **369**, 1094–1098 (2020). https://doi.org/10.1126/science.abb6310

52 North, J. A., Miller, A. R., Wildenthal, J. A., Young, S. J. & Tabita, F. R. Microbial Pathway for Anaerobic 5'-Methylthioadenosine Metabolism Coupled to Ethylene Formation. *Proc. Natl. Acad. Sci. U. S. A.* **114**, e10455–e10464 (2017). https://doi.org/10.1073/pnas.1711625114

53 Lee, C. C., Hu, Y. & Ribbe, M. W. Vanadium Nitrogenase Reduces CO. *Science* **329**, 642–642 (2010). https://doi.org/doi:10.1126/science.1191455

54 Houben, M. & Van de Poel, B. 1-Aminocyclopropane-1-Carboxylic Acid Oxidase (ACO): The Enzyme That Makes the Plant Hormone Ethylene. *Front. Plant. Sci.* **10**, 695 (2019). https://doi.org/10.3389/fpls.2019.00695

55 Wang, K. L., Li, H. & Ecker, J. R. Ethylene biosynthesis and signaling networks. *Plant Cell* **14 Suppl**, S131–151 (2002). https://doi.org/10.1105/tpc.001768

56 Aung, K. M. *et al.* Analyzing the response of rice to tefuryltrione herbicide: Haplotype variation and evolutionary dynamics of the HIS1 gene. *Environmental Research* **252**, 118839 (2024). https://doi.org/https://doi.org/10.1016/j.envres.2024.118839

57 Wu, J. *et al.* Directed evolution of HIS1-Like proteins to enhance herbicide metabolism in crops. *Advanced Agrochem* (2025). https://doi.org/https://doi.org/10.1016/j.aac.2025.05.005

58 Takamura, N. *et al.* Catalytic promiscuity of rice 2-oxoglutarate/Fe(II)-dependent dioxygenases supports xenobiotic metabolism. *Plant Physiol* **187**, 816–828 (2021). https://doi.org/10.1093/plphys/kiab293

59 Lv, Q. *et al.* Exploring Natural Allelic Variations of the β-Triketone Herbicide Resistance Gene HIS1 for Application in indica Rice and Particularly in Two-Line Hybrid Rice. *Rice (N Y)* **14**, 7 (2021). https://doi.org/10.1186/s12284-020-00448-7

60 Müller, T. A., Fleischmann, T., Meer, J. R. v. d. & Kohler, H.-P. E. Purification and Characterization of Two Enantioselective α-Ketoglutarate-Dependent Dioxygenases, RdpA and SdpA, from Sphingomonas herbicidovorans MH. *Applied and Environmental Microbiology* **72**, 4853–4861 (2006). https://doi.org/doi:10.1128/AEM.02758-05

61 Gaines, T. A. *et al.* Mechanisms of evolved herbicide resistance. *J Biol Chem* **295**, 10307–10330 (2020). https://doi.org/10.1074/jbc.REV120.013572