

Full Title: Understanding the regulation of α -globin gene and its implication on the treatment of β -thalassemia

Authors' names: Sachith Mettananda^{1,2}, Richard J Gibbons¹ and Douglas R Higgs^{1,3}

Authors' affiliations:

¹ Medical Research Council (MRC) Molecular Hematology Unit, Weatherall Institute of Molecular Medicine, University of Oxford, Oxford, United Kingdom

² Department of Pediatrics, Faculty of Medicine, University of Kelaniya, Sri Lanka

³ Oxford National Institute for Health Research Biomedical Research Centre, Blood Theme, Oxford University Hospital, Oxford, United Kingdom

Corresponding author's contact information:

Name: Douglas R Higgs

Address: MRC Molecular Hematology Unit, Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Headington, Oxford OX3 9DS, UK.

Email: doug.higgs@imm.ox.ac.uk

Telephone: +44 1865 222393

Fax: +44 1865 222424

Short Title: α -globin regulation and β -thalassemia

Key words: α -globin; β -thalassemia; gene regulation; epigenetic drug targeting; genome editing

Abstract

Over the past three decades, a vast amount of new information has been uncovered describing how the globin genes are regulated. This knowledge has provided significant insights into the general understanding of the regulation of human genes. It is now known that molecular defects within and around the α - and β -globin genes as well as in the distant regulatory elements can cause thalassemia. Unbalanced production of globin chains due to defective synthesis of one and the continued, unopposed synthesis of another is the central causative factor in the cellular pathology and pathophysiology of thalassemia. A large body of clinical, genetic and experimental evidence suggests that altering globin chain imbalance by reducing the production of α -globin synthesis ameliorates the disease severity in patients with β -thalassemia. With the development of new genetic based therapeutic tools that have a potential to decrease the expression of a selected gene in a tissue-specific manner, the possibility of decreasing expression of the α -globin gene to improve the clinical severity of β -thalassemia could become a reality.

Introduction

Hemoglobin is the highly specialized oxygen carrier molecule in human red blood cells (RBC). It is composed of two α - and two β -like globin chains and four heme groups each having an iron (Fe^{2+}) ion as a cofactor. The α - and β -globin gene clusters are two most extensively studied regions of the human genome and the genes in both loci demonstrate well-coordinated, tissue- and developmental-stage-specific patterns of expression. Over the past three decades, a vast amount of new information has been uncovered on the precise regulation of these genes, which includes *cis*-acting regulatory elements (enhancers), *trans*-acting transcription factors and epigenetic mechanisms mediated via chromatin modifications. Extensive knowledge on how these genes are switched 'on' and 'off' has provided significant insights into the general understanding of the regulation of human genes¹.

Thalassemia is caused by impaired synthesis of either α - or β -globin chains. Reduced or absent production of β -globin chains, associated with nearly 300 genetic defects in and around the β -globin gene, result in β -thalassemia²; the most clinically significant form of thalassemia. Despite being one of the first molecular diseases identified and the pathophysiology characterized, the management of β -thalassemia is still far from optimal, limited to supportive therapy in the majority of patients³. It has been known for many decades that natural reductions of α -globin output in the form of co-inherited α -thalassemia ameliorate the disease phenotype in patients with β -thalassemia⁴. However,

except for a limited number of studies, the usefulness of this pathway for treatment of β -thalassemia has not been explored.

During the past few years, genetic therapies have made tremendous progress and generated considerable excitement in the field of translational medicine. Thanks to this, the scope of understanding of regulation of genes has transformed from merely academic exercises to have wider applications including ability to treat diseases. The range of potential genetic therapies now spread far beyond standard gene therapy approaches to methods which fine-tune the expression of specific genes in a tissue-specific manner^{5, 6}.

In this review, we first summarize our current knowledge on regulation of the α -globin genes. Then we present the clinical, genetic and experimental evidence describing how silencing of the α -globin genes ameliorates the clinical phenotype of β -thalassemia. Finally, with the help of new genetic-based therapeutic tools, we propose various pathways which could be utilized to selectively silence the α -globin gene during erythropoiesis.

Regulation of α -globin

Distal enhancers of α -globin

The human α -globin gene cluster is located on the short arm of chromosome 16 (16p13.3) very close (~ 150 kb) to the telomere. The *cis*-acting regulatory network of this gene cluster has been well characterised and it is now known that expression of the α -globin genes is controlled by distant enhancers located 10-48 kilo bases (kb) upstream of the genes (Figure 1). Distal enhancer regulators of α -globin were first identified in the early 1990s, following a report of a family with α -thalassemia due to a large (62kb) deletion upstream of the α -globin locus, which leaves the α -globin genes intact^{7, 8}. A more recent comparative sequence analysis of 22 species identified a conserved syntenic region of ~ 135 kb containing the α -globin cluster and all of the *cis*-acting regulatory elements including multispecies conserved sequences (MCS) required for full tissue- and developmental-stage specific expression of α -globin⁹. Mapping the positions of DNase1 hypersensitive sites to the MCSs showed that in humans, there are four erythroid-specific distant enhancers situated 10kb (MCS-R4), 33kb (MCS-R3), 40kb (MCS-R2) and 48kb (MCS-R1) upstream of the gene locus within the conserved syntenic region. These elements enhance human α -globin expression^{9, 10}.

The four upstream enhancers vary in their capacity to enhance α -globin expression. A variety of experiments suggested that MCS-R2 (previously known as HS-40) is the most critical regulatory element capable of enhancing α -globin expression on its own^{8, 11}. Multispecies analysis revealed that MCS-R2 is the most highly conserved regulatory

element and it could be detected in all mammals⁹. In a humanised mouse model, in which the mouse α -globin locus was replaced by a 117kb segment of DNA containing the human α -globin gene cluster and regulatory elements, the deletion of a 1.1kb segment covering MCS-R2 in the humanised chromosome resulted in a reduction of human α -globin expression to very low levels¹². Similarly, all naturally occurring human deletions which cause α -thalassemia by removing the upstream regulatory elements include MCS-R2¹³. A rare patient homozygous for a naturally occurring deletion limited to MCS-R2 region developed α -thalassemia with haemoglobin H disease¹⁴. These observations confirm that MCS-R2 is crucial for the expression of the human α -globin genes. Of the remaining enhancer like elements only MCS-R1 has been shown to have significant activity *in vivo*.

Epigenetic regulation of α -globin

The human α -globin gene cluster is located in a gene dense, early replicating and open chromatin region of the genome surrounded by a number of housekeeping genes which are expressed in all cell types¹. DNA at this locus is substantially unmethylated and the promoters of all α -like globin genes are associated with unmethylated cytosine-guanine dinucleotide (CpG) rich islands. Because the human α -globin locus is in an open chromatin environment, the genes are actively silenced through a transcriptional repressive protein complex known as polycomb repressive complex (PRC) 2 in non-erythroid cells, in which α -globin is not expressed¹⁵. The transcriptional repressor action of PRC2 is associated with a histone methyltransferase (EZH2) which increases trimethylation of histone H3 lysine 27 (H3K27me3), a chromatin signature associated

with gene silencing¹⁶. The mechanism involved in the binding of PRC2 complex to the target genomic sites is not fully understood but clearly depends on the unmethylated CpG island promoters¹⁷. In non-erythroid cells in humans, PRC2 is recruited to the α -globin promoter and the chromatin is modified by H3K27me3 and this is associated with transcriptional silencing¹⁵.

In humanised mouse erythroid cells, when the α -globin gene is activated, PRC2 is displaced and the H3K27me3 chromatin mark is erased through both passive and active mechanisms. The active mechanism of demethylation of H3K27me3 is thought to be mediated via histone lysine demethylase (KDM) 6B (also known as Jumonji domain-containing protein 3 [JMJD3])¹⁸. In addition, as cells differentiate into erythroid cells a dramatic increase in the levels of the H3K4me3 chromatin modification which signals transcriptional activation is also seen at the promoter of the α -globin genes¹⁹. This is believed to be mediated through the action of Set/MLL histone methyltransferases. The net effect of these chromatin modifications and the recruitment of transcription factors (TF) is the activation of transcription of the α -globin genes to produce large amounts of globin mRNA and proteins required to make hemoglobin.

Transcription factors controlling expression of α -globin

During human erythroid differentiation the key TFs, GATA-binding factor 1 (GATA1), Kruppel-like factor-1 (KLF1), stem cell leukaemia (SCL), transcription factor E2-alpha (E2A), Lim-only 2 (LMO2) and LIM domain binding protein 1 (Ldb1) are recruited to the four upstream enhancers of the α -globin genes. In addition, another erythroid TF,

nuclear factor-erythroid 2 (NF-E2) is bound to two of the enhancers; MCS-R2 and to a lesser extent MCS-R1²⁰. Finally, general transcription factors together with RNA polymerase II (PolII) (collectively referred to as the preinitiation complex) are recruited to the MCS-R1 and MCS-R2 enhancers and then subsequently to the α -globin promoters thereby initiating transcription. During this process a physical interaction between enhancers and promoters are thought to occur through the mechanism of chromatin looping^{21, 22}.

Contrasting regulation α - and β -globin expression

Despite being expressed in the same cell type, in a similar developmental-stage-specific manner, the regulation of β -globin demonstrates significant differences to that of α -globin. The human β -globin gene is located on chromosome 11 in a relatively gene sparse, late replicating region of the genome which, in non-erythroid cells, is in a closed heterochromatic environment^{1, 23}. The promoter of β -globin is methylated and not associated with CpG islands. Therefore, in contrast to α -globin, in non-erythroid cells the promoter of β -globin is not bound by PRC2 and does not have the repressive chromatin mark H3K27me3^{15, 24}. Instead, silencing of the β -globin gene is mediated through DNA methylation and co-localisation with nuclear heterochromatin¹.

In erythroid cells when the β -globin is activated, in a process similar to α -globin, the distant enhancers (referred to as the β -locus control region) and the gene promoters are bound by erythroid specific TFs which include (but are not limited to) GATA1, NF-E2, KLF1 and SCL^{25, 26}. Similar to α -globin, the interaction of the enhancers and promoters

are believed to be maintained by chromosomal looping²⁷. The promoter is marked by the H3K4me3 active chromatin modification. However, in contrast to α -globin, the repressive chromatin modification H3K27me3 is not present at the β -locus and, histone demethylases have not been shown to play a role in the activation of β -globin¹⁵.

Reduction of α -globin as a therapeutic option for β -thalassemia

Role of free α -globin chains in the molecular pathology of β -thalassemia

Unbalanced production of α - and β -globin chains within erythroid cells is the main causative factor leading to hemolysis and ineffective erythropoiesis which are the two important contributing factors to anemia in thalassemia^{4, 28}. During every stage of normal development, production of α -like and β -like globin chains are closely balanced to prevent an accumulation of free globin chains which damage the RBCs. The deleterious effects of small amounts of free α -globin chains are diminished by the erythroid specific molecular chaperone, α -hemoglobin stabilizing protein (AHSP). AHSP specifically binds to α -globin and stabilizes free α -chains by promoting protein folding and resistance to protease digestion thereby protecting erythroid cells from toxic effects of free α -globin²⁹.

In patients with β -thalassemia, impaired production of β -globin chains results in an excess of free α -globin chains. The levels of free α -globin chains exceed the capacity of AHSP to buffer their effect. Once this capacity is exceeded, highly unstable free α -globin molecules undergo auto-oxidation forming α -hemichromes (α -globin monomers which contain oxidized ferric iron) and reactive oxygen species (ROS) to trigger a cascade of events leading to hemolysis and ineffective erythropoiesis (Figure 2)^{30, 31}.

Co-inheritance of α - and β -thalassemia

Since the initial descriptions by Fessas³², and Kan and Nathan³³, co-inheritance of α -thalassemia has been predicted to ameliorate the severity of β -thalassemia. A number of family, cohort and case-control studies have reported that the co-inheritance of a deletion of two α -globin genes ($--/\alpha\alpha$ or $-\alpha/-\alpha$) is associated with a milder clinical phenotype in most patients with β -thalassemia while deletion of a single α -globin gene ($-\alpha/\alpha\alpha$) is beneficial in patients other than those with the most severe reduction in β -globin synthesis (β^0/β^0 genotype)³⁴⁻⁴⁰. Furthermore, a recent study showed that the presence of a mutated α -globin allele provides an additive effect to induction of γ -globin expression, the other major disease modifier of β -thalassemia⁴¹.

Co-inheritance of α -thalassemia appears to have its most pronounced effect on patients with HbE β -thalassemia, the most common sub-type of β -thalassemia worldwide. Importantly, co-inheritance of even a single α -globin gene deletion ($-\alpha/\alpha\alpha$) is able to produce beneficial effects in these patients. In large scale prospective studies, all patients with co-inherited α -thalassemia displayed a milder phenotype, older age at presentation, smaller splenic and hepatic sizes, normal physical and sexual maturation and significantly less transfusion requirement^{39, 42-46}. Furthermore, in HbE β -thalassemia patient cohorts, the frequency of α -thalassemia is significantly lower than in the normal matched population. Similarly it is extremely rare to find a patient with HbE β -thalassemia and two α -globin gene deletions ($--/\alpha\alpha$ or $-\alpha/-\alpha$) requiring transfusion treatment^{42, 44}. These observations suggest that individuals with HbE β -thalassemia,

who co-inherit only two, rather than the normal four, α -globin genes have a very mild phenotype and rarely come to medical attention.

Effect of excess α -globin chains on β -thalassemia

Contrary to the beneficial effect demonstrated by co-inheritance of α and β -thalassemia, inheritance of excess α -globin genes in the presence of β -thalassemia worsens disease severity⁴⁷⁻⁴⁹. Patients with heterozygous β -thalassemia mutations, who would otherwise have a β -thalassemia minor phenotype, develop a thalassemia intermedia phenotype when they inherit more than four functional copies of the α -globin gene. This is because the inheritance of excess α -globin genes in these patients increases the amount of free α -globin within RBCs and worsens the α -like/ β -like globin chain imbalance. Increase in free α -globin chains exacerbates the premature erythroid cell destruction and worsens the severity of anaemia. This observation further emphasizes that the number of functional α -globin genes has clear and direct effects on the clinical severity of patients with β -thalassemia.

Evidence from mouse models

Amelioration of β -thalassemia by reducing α -globin has also been recapitulated in the β -globin knockout (β -KO) mouse model. Voon et al generated a double heterozygous mouse knocked-out for both α - and β -globin genes ($\alpha^{++/-}$; $\beta^{+/-}$) and compared the hematological parameters to those of the β -KO mice ($\alpha^{++/++}$; $\beta^{+/-}$)⁵⁰. Interestingly, all the observed parameters including hemoglobin level, hematocrit, red cell distribution width, reticulocyte number and the percentage of erythroid progenitors in the bone

marrow and spleen were corrected to near normal values in the double heterozygous mouse. This observation further emphasises that a better balanced globin chain synthesis by co-inheritance of α -thalassemia might make profound improvements to the β -thalassemia phenotype and provides an experimental proof that α -globin is a suitable target to develop therapies.

Desired level of reduction of α -globin

The clinical, genetic and experimental evidence discussed in the preceding sections strongly supports the idea that reducing α -globin expression in patients with β -thalassemia is a rational pathway to develop new therapies. Reducing α -globin output to levels comparable to a single or two α -globin gene deletion should be clinically beneficial to patients with β -thalassemia. Among patients with α -thalassemia, single and two α -globin gene deletions correspond to 75% and 50% α -globin levels respectively compared to normal controls. Therefore, these levels could be considered as rational therapeutic aims in β -thalassemia^{13, 51}. Although, the data on coinheritance of β -thalassemia and three α -globin gene deletions are sparse, reductions down to 25% of the normal levels (as seen in patients with α -thalassemia due to three α -globin gene deletions) do not result in significant morbidity due to anemia and can be considered as a safe tolerable lower limit.

Selective silencing of α -globin expression: therapeutic pathways

The development of new drugs targeting specific epigenetic pathways and the advent of genome engineering using programmable, sequence-specific endonucleases have generated potentials to specifically regulate the expression of a single gene. Similarly, current understanding of regulation of α - and β -globin genes in erythroid cells have uncovered pathways to consider new strategies to selectively control α -globin expression to the appropriate degree to be useful for patients with β -thalassemia. Several pathways warrant careful consideration. Post-transcriptional gene silencing through RNA interference (RNAi) is one such plausible avenue which has been explored, to some extent, in the past. Pharmacological targeting of the contrasting epigenetic regulation of α - and β -globin genes and selective silencing of α -globin using genome engineering strategies are the other pathways which have potentials for clinical application.

RNA interference strategy

RNAi mediated through small double stranded RNAs (small interfering RNA [siRNA] and short hairpin RNA [shRNA]) can produce efficient and specific post-transcriptional gene silencing^{52, 53}. Attempts to reduce α -globin expression using this technique have been reported in the past. Up to 50% knockdown of α -globin gene expression using siRNA targeting the α -globin in β -thalassemia murine primary erythroid cells, demonstrated significant phenotypic improvement by reducing the levels of intracellular ROS⁵⁴. Transgenic β -thalassemia mice which produced 20%-35% less α -globin,

generated by microinjecting shRNA targeting α -globin into single-cell embryos, demonstrated sustained phenotypic improvement in the RBCs with less poikilocytosis, fewer target cells, increased hemoglobin values and red cell counts, and low reticulocyte counts⁵⁵. However, direct injection of siRNA plasmids targeting α -globin mRNA into the tail veins of the β -thalassemia mice did not produce improvements in red cell counts and haemoglobin levels⁵⁶.

Pharmacological strategy through epigenetic drug targeting

Small molecules which alter the chromatin environment by selectively inhibiting epigenetic enzymes have become popular as new therapeutics in recent years⁶. Although, most are indicated and trialled for various forms of cancers, their usefulness in the treatment of non-malignant conditions are being investigated. In fact, the results from the very first clinical trial which tested the effect of a histone deacetylase inhibitor for sickle cell disease were recently published⁵⁷. As outlined earlier, the chromatin environment and silencing mechanisms of the α - and β -globin clusters have very clear differences. This distinction may be very useful in the development of therapeutic strategies and it is likely that drugs or small molecules which act by altering the chromatin environment will have a differential effect on α - and β -globin expression.

Genome editing strategy

Genome editing using sequence-specific programmable artificially engineered nucleases has provided a realistic approach for the treatment of many human genetic diseases including the hemoglobinopathies. These nucleases create double strand

breaks at specific chosen locations in the genome, which when repaired, create mutations at the targeted sites and, depending on their position, result in either loss of gene function or disruption of non-coding regulatory sequences. The zinc-finger nuclease (ZFN) was the first such technology to be developed. Although, this generated lots of promise, many drawbacks including cytotoxicity, low-targeting efficiency, difficulties in engineering proteins for each target and limitations in target sites (one per ~100bp) precluded its widespread use⁵⁸. The second group, transcription activator-like effector nucleases (TALENs), were popular as they dramatically increased the availability of target sites allowing almost any sequence to be targeted. However, the construction of TALENs is challenging and time-consuming⁵⁹. Both ZFNs and TALENs use DNA-binding proteins to tether endonuclease catalytic domains to specific genomic loci. By contrast, in the recently developed RNA-guided CRISPR–Cas9 (clustered, regularly interspaced, short palindromic repeat-CRISPR associated system) system, the endonuclease Cas9 is guided by small RNAs through Watson-Crick base pairing with target DNA⁶⁰. Therefore it represents a system that is markedly easier to design, highly specific, efficient and well-suited for high-throughput and multiplexed gene editing for a variety of cell types and organisms⁶¹.

These genome editing approaches can be used to disrupt the expression of the α -globin gene. This can be done by directly targeting the genes or through targeting regulatory elements of the genes. If this strategy is successful, translation into clinical practice would broadly involve three steps. Firstly, the hematopoietic stem cells (HSC) will be harvested from patients with β -thalassemia after mobilisation using granulocyte colony-

stimulating factor. Then the harvested HSCs will be edited *ex vivo* using programmable nucleases and transplanted back to the same patient after myeloablation.

Conclusions

Advances in the understanding of regulation of α - and β -globin genes have demonstrated clear and striking differences in silencing mechanisms of these two gene clusters. Pathways are thus uncovered for selective silencing of α -globin expression to levels beneficial to patients with β -thalassemia. Novel genetic therapeutic tools now offer the potential to translate these theoretical mechanisms into practical realities. In this way, we are approaching the dawn of realising the transformation of a long standing clinical observation to an effective therapy in β -thalassemia.

Acknowledgements

This work was supported by grants to D.R.H. by the UK Medical Research Council and the NIHR Oxford Biomedical Research Centre. S.M. is a Commonwealth Scholar, funded by the UK government.

References

1. Higgs, D. R., *et al.* 2005. Understanding alpha-globin gene regulation: Aiming to improve the management of thalassemia. *Annals of the New York Academy of Sciences*. 1054: 92-102.
2. Thein, S. L. 2013. The molecular basis of beta-thalassemia. *Cold Spring Harbor perspectives in medicine*. 3: a011700.
3. Higgs, D. R., J. D. Engel & G. Stamatoyannopoulos. 2012. Thalassaemia. *Lancet*. 379: 373-383.
4. Mettananda, S., R. J. Gibbons & D. R. Higgs. 2015. alpha-Globin as a molecular target in the treatment of beta-thalassemia. *Blood*. 125: 3694-3701.
5. Cox, D. B., R. J. Platt & F. Zhang. 2015. Therapeutic genome editing: prospects and challenges. *Nature medicine*. 21: 121-131.
6. Helin, K. & D. Dhanak. 2013. Chromatin proteins and modifications as drug targets. *Nature*. 502: 480-488.
7. Hatton, C. S., *et al.* 1990. Alpha-thalassemia caused by a large (62 kb) deletion upstream of the human alpha globin gene cluster. *Blood*. 76: 221-227.
8. Higgs, D. R., *et al.* 1990. A major positive regulatory region located far upstream of the human alpha-globin gene locus. *Genes & development*. 4: 1588-1601.
9. Hughes, J. R., *et al.* 2005. Annotation of cis-regulatory elements by identification, subclassification, and functional assessment of multispecies conserved sequences. *Proceedings of the National Academy of Sciences of the United States of America*. 102: 9830-9835.
10. Higgs, D. R. & W. G. Wood. 2008. Long-range regulation of alpha globin gene expression during erythropoiesis. *Current opinion in hematology*. 15: 176-183.
11. Sharpe, J. A., *et al.* 1992. Analysis of the human alpha globin upstream regulatory element (HS-40) in transgenic mice. *The EMBO journal*. 11: 4565-4572.
12. Wallace, H. A. C., *et al.* 2007. Manipulating the mouse genome to engineer precise functional syntenic replacements with human sequence. *Cell*. 128: 197-209.
13. Hartevelde, C. L. & D. R. Higgs. 2010. Alpha-thalassaemia. *Orphanet journal of rare diseases*. 5: 13.
14. Coelho, A., *et al.* 2010. Novel large deletions in the human alpha-globin gene cluster: Clarifying the HS-40 long-range regulatory role in the native chromosome environment. *Blood cells, molecules & diseases*. 45: 147-153.
15. Garrick, D., *et al.* 2008. The role of the polycomb complex in silencing alpha-globin gene expression in nonerythroid cells. *Blood*. 112: 3889-3899.
16. Simon, J. A. & R. E. Kingston. 2013. Occupying chromatin: Polycomb mechanisms for getting to genomic targets, stopping transcriptional traffic, and staying put. *Molecular cell*. 49: 808-824.
17. Lynch, M. D., *et al.* 2012. An interspecies analysis reveals a key role for unmethylated CpG dinucleotides in vertebrate Polycomb complex recruitment. *The EMBO journal*. 31: 317-329.
18. Vernimmen, D., *et al.* 2011. Polycomb eviction as a new distant enhancer function. *Genes & development*. 25: 1583-1588.
19. De Gobbi, M., *et al.* 2011. Generation of bivalent chromatin domains during cell fate decisions. *Epigenetics & chromatin*. 4: 9.

20. De Gobbi, M., *et al.* 2007. Tissue-specific histone modification and transcription factor binding in alpha globin gene expression. *Blood*. 110: 4503-4510.
21. Vernimmen, D., *et al.* 2009. Chromosome looping at the human alpha-globin locus is mediated via the major upstream regulatory element (HS -40). *Blood*. 114: 4253-4260.
22. Vernimmen, D. 2014. Uncovering Enhancer Functions Using the alpha-Globin Locus. *PLoS genetics*. 10: e1004668.
23. Higgs, D. R., J. A. Sharpe & W. G. Wood. 1998. Understanding alpha globin gene expression: a step towards effective gene therapy. *Seminars in hematology*. 35: 93-104.
24. Chang, K. H., *et al.* 2013. Epigenetic modifications and chromosome conformations of the beta globin locus throughout development. *Stem cell reviews*. 9: 397-407.
25. Cantor, A. B. & S. H. Orkin. 2002. Transcriptional regulation of erythropoiesis: an affair involving multiple partners. *Oncogene*. 21: 3368-3376.
26. Kim, A. & A. Dean. 2012. Chromatin loop formation in the beta-globin locus and its role in globin gene transcription. *Molecules and cells*. 34: 1-5.
27. Palstra, R. J., *et al.* 2003. The beta-globin nuclear compartment in development and erythroid differentiation. *Nature genetics*. 35: 190-194.
28. Ribeil, J. A., *et al.* 2013. Ineffective erythropoiesis in beta -thalassemia. *TheScientificWorldJournal*. 2013: 394295.
29. Yu, X., *et al.* 2007. An erythroid chaperone that facilitates folding of alpha-globin subunits for hemoglobin synthesis. *The Journal of clinical investigation*. 117: 1856-1865.
30. Voon, H. P. & J. Vadolas. 2008. Controlling alpha-globin: a review of alpha-globin expression and its impact on beta-thalassemia. *Haematologica*. 93: 1868-1876.
31. Shinar, E. & E. A. Rachmilewitz. 1993. Haemoglobinopathies and red cell membrane function. *Bailliere's clinical haematology*. 6: 357-369.
32. Fessas, P., G. Stamatoyannopoulos & A. Karaklis. 1961. Hereditary persistence of foetal haemoglobin and its combination with alpha and beta-thalassaemia. *In* 8th Congress of European Society of Haematology: 302. Karger, Basel, Vienna.
33. Kan, Y. W. & D. G. Nathan. 1970. Mild thalassemia: the result of interactions of alpha and beta thalassemia genes. *The Journal of clinical investigation*. 49: 635-642.
34. Wainscoat, J. S., *et al.* 1983. The molecular basis for the clinical diversity of beta thalassaemia in Cypriots. *Lancet*. 1: 1235-1237.
35. Thein, S. L., *et al.* 1988. The molecular basis of thalassaemia major and thalassaemia intermedia in Asian Indians: application to prenatal diagnosis. *Br J Haematol*. 70: 225-231.
36. Galanello, R., *et al.* 1989. Molecular analysis of beta zero-thalassemia intermedia in Sardinia. *Blood*. 74: 823-827.
37. Gringras, P., *et al.* 1994. Effect of alpha thalassaemia trait and enhanced gamma chain production on disease severity in beta thalassaemia major and intermedia. *Archives of disease in childhood*. 70: 30-34.
38. Ho, P. J., *et al.* 1998. Beta-thalassaemia intermedia: is it possible consistently to predict phenotype from genotype? *Br J Haematol*. 100: 70-78.
39. Winichagoon, P., *et al.* 2000. Genetic factors affecting clinical severity in beta-thalassemia syndromes. *Journal of pediatric hematology/oncology*. 22: 573-580.

40. Camaschella, C., *et al.* 1995. Genetic interactions in thalassemia intermedia: analysis of beta-mutations, alpha-genotype, gamma-promoters, and beta-LCR hypersensitive sites 2 and 4 in Italian patients. *American journal of hematology*. 48: 82-87.
41. Badens, C., *et al.* 2011. Variants in genetic modifiers of beta-thalassemia can help to predict the major or intermedia type of the disease. *Haematologica*. 96: 1712-1714.
42. Sripichai, O., *et al.* 2008. Coinheritance of the different copy numbers of alpha-globin gene modifies severity of beta-thalassemia/Hb E disease. *Annals of hematology*. 87: 375-379.
43. Premawardhena, A., *et al.* 2005. Haemoglobin E beta thalassaemia in Sri Lanka. *Lancet*. 366: 1467-1470.
44. Suresh, S., *et al.* 2013. Alpha thalassaemia and extended alpha globin genes in Sri Lanka. *Blood cells, molecules & diseases*. 50: 93-98.
45. Bandyopadhyay, S., *et al.* 2001. Variable severity of beta-thalassemia patients of eastern India: effect of alpha-thalassemia and xmnI polymorphism. *Clinical and experimental medicine*. 1: 155-159.
46. Sharma, V. & R. Saxena. 2009. Effect of alpha-gene numbers on phenotype of HbE/beta thalassemia patients. *Annals of hematology*. 88: 1035-1036.
47. Premawardhena, A., *et al.* 2005. A novel molecular basis for beta thalassemia intermedia poses new questions about its pathophysiology. *Blood*. 106: 3251-3255.
48. Sollaino, M. C., *et al.* 2009. Association of alpha globin gene quadruplication and heterozygous beta thalassemia in patients with thalassemia intermedia. *Haematologica*. 94: 1445-1448.
49. Origa, R., *et al.* 2013. alpha-Globin Gene Quadruplication and Heterozygous beta-Thalassemia: A Not So Rare Cause of Thalassemia Intermedia. *Acta haematologica*. 131: 162-164.
50. Voon, H. P., H. Warden & J. Vadolas. 2007. Co-inheritance of alpha- and beta-thalassaemia in mice ameliorates thalassaemic phenotype. *Blood cells, molecules & diseases*. 39: 184-188.
51. Chaisue, C., *et al.* 2007. alpha/beta-Globin mRNA ratio determination by multiplex quantitative real-time reverse transcription-polymerase chain reaction as an indicator of globin gene function. *Clinical biochemistry*. 40: 1373-1377.
52. Hannon, G. J. 2002. RNA interference. *Nature*. 418: 244-251.
53. Gavrillov, K. & W. M. Saltzman. 2012. Therapeutic siRNA: principles, challenges, and strategies. *The Yale journal of biology and medicine*. 85: 187-200.
54. Voon, H. P., H. Warden & J. Vadolas. 2008. siRNA-mediated reduction of alpha-globin results in phenotypic improvements in beta-thalassemic cells. *Haematologica*. 93: 1238-1242.
55. Xie, S. Y., *et al.* 2007. Restoration of the balanced alpha/beta-globin gene expression in beta654-thalassemia mice using combined RNAi and antisense RNA approach. *Human molecular genetics*. 16: 2616-2625.
56. Xie, S. Y., *et al.* 2011. Correction of beta654-thalassaemia mice using direct intravenous injection of siRNA and antisense RNA vectors. *International journal of hematology*. 93: 301-310.

57. Okam, M. M., *et al.* 2015. Phase 1/2 trial of vorinostat in patients with sickle cell disease who have not benefitted from hydroxyurea. *Blood*. 125: 3668-3669.
58. Kim, H. & J. S. Kim. 2014. A guide to genome engineering with programmable nucleases. *Nature reviews. Genetics*. 15: 321-334.
59. Cheng, J. K. & H. S. Alper. 2014. The genome editing toolbox: a spectrum of approaches for targeted modification. *Current opinion in biotechnology*. 30C: 87-94.
60. Garneau, J. E., *et al.* 2010. The CRISPR/Cas bacterial immune system cleaves bacteriophage and plasmid DNA. *Nature*. 468: 67-71.
61. Ran, F. A., *et al.* 2013. Genome engineering using the CRISPR-Cas9 system. *Nature protocols*. 8: 2281-2308.

Figure legends




Figure 1- Schematic representation of the α -globin gene cluster (top) and the MCS-R2 enhancer element (bottom). The main globin genes in the α -globin gene cluster are ζ (embryonic) and $\alpha 1$ and $\alpha 2$ (adult). The four upstream regulatory elements of the α -locus situated 10-48kb upstream to the genes are known as MCS-R1 to MCS-R4. The most critical MCS-R2 enhancer has binding sites for erythroid specific TFs GATA1 and NF-E2.




Figure 2- Pathophysiology of β -thalassemia. Absent or reduced β -globin production leads to an unbalanced excess of α -globin chains, which then trigger cascade of events through the generation of ROS, resulting in hemolysis of mature red blood cells and destruction of immature erythroid precursors in the bone marrow (ineffective erythropoiesis). The resultant anemia increases the intestinal iron absorption which together with transfusion therapy leads to body iron overload and subsequent organ dysfunction.