

Study References [posted as supplied by authors]

w1. GlaxoSmithKline. Study: NAI30028. 2006 Jan 4 (on line). Available from: URL: <http://www.gsk-clinicalstudyregister.com/files/pdf/3304.pdf>. Accessed July 1, 2009.

w2. Hedrick JA, Barzilai A, Behre U, et al. Zanamivir for treatment of symptomatic influenza A and B infection in children five to twelve years of age: a randomized controlled trial. *Pediatr Infect Dis J* 2000; **19**: 410–7. Additional data: GlaxoSmithKline. Study: NAI30009. 2005 June 4. Available from: URL: <http://www.gsk-clinicalstudyregister.com/files/pdf/3298.pdf>. Accessed July 1, 2009.

w3. Whitley RJ, Hayden FG, Reisinger KS, et al. Oral oseltamivir treatment of influenza in children. *Pediatr Infect Dis J*. 2001; **20**: 127–33. Erratum in: *Pediatr Infect Dis J* 2001; **20**: 421.

w4. Johnston SL, Ferrero F, Garcia ML, Dutkowski R. Oral oseltamivir improves pulmonary function and reduces exacerbation frequency for influenza-infected children with asthma. *Pediatr Infect Dis J* 2005; **24**: 225–32.

w5. Hayden FG, Gubareva LV, Monto AS, et al. Inhaled zanamivir for the prevention of influenza in families. Zanamivir Family Study Group. *N Engl J Med* 2000; **343**: 1282–9. Additional data: GlaxoSmithKline. Study: NAI30010. 2005 July 12 Available from: URL: <http://www.gsk-clinicalstudyregister.com/files/pdf/3300.pdf>. Accessed July 1, 2009. Further unpublished data supplied by GlaxoSmithKline.

w6. Monto AS, Pichichero ME, Blanckenberg SJ, et al. Zanamivir prophylaxis: an effective strategy for the prevention of influenza types A and B within households. *J Infect Dis* 2002; **186**: 1582–8. Additional data from: Roche. Study: NAI30031. 2005 June 14. Available from: URL: <http://www.gsk-clinicalstudyregister.com/files/pdf/3371.pdf>. Accessed July 1, 2009. Further unpublished data supplied by GlaxoSmithKline

w7. Hayden FG, Belshe R, Villanueva C, et al. Management of influenza in households: a prospective, randomized comparison of oseltamivir treatment with or without postexposure prophylaxis. *J Infect Dis* 2004; **189**: 440–9.

W8. Winther B, Hayden FG, Whitley R, et al. Oral oseltamivir reduces the risk of developing acute otitis media following influenza infection in children. In: 40th Interscience Conference on Antimicrobial Agents and Chemotherapy, Toronto. 2000