

Remission in PsA – where are we now?

Laura C. Coates,^{1†} Philip G. Conaghan,¹ Maria Antonietta D’Agostino,² Maarten De Wit,³ Oliver FitzGerald,⁴ Tore K Kvien,⁵ Rik Lories,⁶ Philip Mease,⁷ Peter Nash,⁸ Georg Schett,⁹ Enrique R. Soriano,¹⁰ Paul Emery¹

1. Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, and National Institute for Health Research (NIHR) Leeds Biomedical Research Centre, Leeds, UK
2. APHP, Hôpital Ambroise Paré, Rheumatology Department, 92100 Boulogne-Billancourt; INSERM U1173, Laboratoire d’Excellence INFLAMEX, UFR Simone Veil, Versailles-Saint-Quentin University, 78180 Saint-Quentin en Yvelines; FRANCE
3. Department of Medical Humanities, VU University Medical Centre, Amsterdam, The Netherlands
4. Department of Rheumatology, St Vincent’s University Hospital, and Conway Institute for Biomolecular Research, University College Dublin, Ireland
5. Department of Rheumatology, Diakonhjemmet Hospital, Oslo, Norway
6. Skeletal Biology and Engineering Research Center, KU Leuven, Leuven, Belgium; Division of Rheumatology, UZ Leuven, Leuven, Belgium.
7. University of Washington School of Medicine/ Swedish Medical Center, Seattle, USA
8. Department of Medicine, University of Queensland, Brisbane, Queensland, Australia
9. Department of Internal Medicine 3 and Institute for Clinical Immunology, Friedrich Alexander University of Erlangen-Nuremberg, Erlangen, Germany
10. Rheumatology Unit, Internal Medical Services, Hospital Italiano de Buenos Aires, Buenos Aires, Argentina

†Current address: Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Oxford, UK.

Keywords: PsA, remission, Treat to target, tight control, disease activity, imaging, biomarkers, Doppler ultrasound, MRI

Running title: PsA remission review

Corresponding author:

L.C. Coates, MB, BS, MRCP, PhD

[Current full postal address and email]

Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences,

Botnar Research Centre

University of Oxford

Windmill Road

Oxford, OX3 7LD

UK

Telephone - +44 1865 737545

Abstract

Advances in treatments and treatment strategies for psoriatic arthritis (PsA) have led to many patients responding well to management of their disease, and targeting remission as a treatment goal is now a possibility. Treat to target (T2T) is a strategy aimed at maximizing benefit, irrespective of the type of medication used, by monitoring disease activity and using it to guide therapy. The measurement of response to treatment has been the subject of wide discussions among experts for some time, and many instruments exist. Comparisons of the different measures and their different strengths and weaknesses, is ongoing. The impact of modern imaging techniques on monitoring disease progression is also evolving, and advanced techniques using both magnetic resonance imaging (MRI) and ultrasound (US) have the potential to improve management of PsA through identification of risk factors for poor prognosis as well as accurate assessment of inflammation and damage, including subclinical disease. Increased understanding of the pathways that drive the pathogenesis of PsA will be key to identifying specific biomarkers for the disease and developing effective treatment strategies. Targets for response, considerations for use of a T2T strategy in PsA, different imaging techniques, and serological aspects of remission are all discussed in this review, and areas for further research are identified.

Introduction

The treatment goals for patients with psoriatic arthritis (PsA) are control of disease activity, improvement of physical function and quality of life, and prevention of structural damage to joints [1-3]. In the last few years, advances in pharmacological treatment of PsA, particularly the introduction of biologic therapies, have enabled excellent responses to be achieved in many patients [4]. However, PsA is a heterogeneous disease and measuring its response to treatment, both in the clinic and in clinical trials, has been the subject of wide debate.

Advances in treatment strategies for rheumatic diseases have also occurred. Treat to target (T2T) is aimed at maximizing benefit, irrespective of the type of medication used, by monitoring disease activity using the best current measures and remission criteria [5]. The Tight Control of disease activity in rheumatoid arthritis (RA), (TICORA) [6] study showed that escalating therapy in a T2T strategy could improve outcomes in RA. The study investigated an intensive treatment strategy consisting of frequent, objective assessment of patients, intensive use of intra-articular steroid injections if needed, and a structured protocol for the escalation of treatment in patients with active disease despite treatment. The targets in RA have become more stringent over time, related to a greater ability to achieve remission as new, better treatments are developed [7].

Evidence for T2T in PsA only began to emerge in 2013, and many treatments and outcome measures have been ‘borrowed’ from RA. There has been little agreement on what target(s) for response should be used in PsA [8], and a literature review by the European League Against Rheumatism (EULAR) showed that there were few relevant studies on T2T in PsA [8].

The use of magnetic resonance imaging (MRI), ultrasound (US), and computed tomography in the study of PsA has permitted a better understanding of the various pathologies of the PsA phenotypes. These sensitive imaging techniques have highlighted the high frequency of subclinical inflammation and added insights into the persistence of inflammation and structural damage after therapy [9-12].

This review provides an overview of the current status of targeting remission in PsA, including a focus on areas that need more research. It resulted from a consensus meeting with an expert panel of clinicians involved in PsA routine management and research in February 2016.

Considerations in Applying Treat to Target (T2T) in PsA Clinical Practice

Specific aspects and challenges of remission in PsA

In order to assess remission, it must first be defined. Remission implies that at a minimum, the inflammatory disease process will be controlled such that the patient has no symptoms and no long-term functional or structural joint consequences [13]. Even in clinical manifestations, PsA is a multifaceted disease with varied rheumatological and dermatological presentations. Beyond this, PsA not only has clinical manifestations, but is also characterized by structural and immunologic changes. Therefore PsA remission may encompass more than remission of the clinical signs and symptoms of musculoskeletal and skin disease.

Core domains for assessment of PsA were defined by Outcome Measures in Rheumatology (OMERACT) in 2006 [14] and updated in 2016 [15, 16], and a core set of domains criteria for minimal disease activity (MDA) in PsA have also been defined [7, 17]. Ideally, the target for remission should be feasible for clinical use and, as PsA is a heterogeneous condition, should include assessment of all key different domains. As yet, there are no reliable serum markers of PsA disease activity.

Another major factor affecting quality of life for PsA patients is comorbidity, and this aspect needs to be considered when setting realistic expectations of disease remission. A large proportion of PsA patients have comorbidities, which are often under-recognized and undertreated, which may influence treatment, prognosis and outcomes; they include cardiovascular disease, obesity, metabolic syndrome, depression, uveitis and cancer [18, 19]. One study has found that 42% of PsA patients have three or more comorbidities; however, the incremental effects of comorbidities on quality of life relate more to the type rather than the number of comorbidities [20]. Targeted treatment is therefore an important concept in achieving patient-defined remission.

Patient perspectives on disease activity, treatment and remission in PsA

Patients with PsA and their physicians may view the disease differently, and there is a discrepancy between patient and physician assessment of joint activity [21]. An analysis of 565 patients found that patients scored their disease worse than physicians, with the discordance greater for joints than for skin parameters. Similar discrepancy is well-documented in RA [22], but has been less well-studied in PsA.

Patient education in PsA is often not optimal and PsA patients are less empowered than those with RA [23]. However, a recent study showed that the difference between patients and physician global assessment of disease activity as well as the difference between tender and swollen joint count were associated with a reduced risk of achieving remission, both in PsA and RA [24].

Results of the Multinational Assessment of Psoriasis and Psoriatic Arthritis Survey [25] showed that 59% of surveyed PsA patients were receiving no treatment or only topical treatment. This is partly due to low expectations on the part of patients that dermatologists or rheumatologists will be able to offer effective treatments.

For patients, the impact of disease on quality of life and function is important. Although individuals may have very different expectations of how their disease is managed, aspects of disease and treatment that are important to patients are not adequately covered by the self-report measures (both patient reported outcomes [PROs] and existing composite scores) most often used in PsA patients [26]. These include the impact of environmental factors, societal attitudes towards individuals with psoriasis (PsO) or PsA, the increased feeling of isolation from social activities, and treatment burden, resulting in, for instance, lack of leisure time. Expectations are an important factor in disease management. For example, there is evidence that RA patients consider remission more as a feeling of returning to normality, rather than an absence or reduction of symptoms [27]. Treatment clearly impacts quality of life for PsA patients, and there is evidence that treatment early in the course of the disease (< 2 years disease duration) led to greater improvements in arthritis scores and quality of life measures compared with those treated after having the disease for more than 2 years [28].

Clinical remission in PsA: How to measure it

Comparison of different instruments used to assess disease activity and measure outcomes

The composite measures used to assess disease activity are compared in Table 1, built by author consensus while writing the paper. These composite measures combine individual measures of disease activity into a single score and, while this may be a more efficient approach than comparing across individual scores, the ability to distinguish between changes in disease activity in individual clinical features may be lost [29]. Different outcome measures may be used in clinical practice from those used in clinical trials although, if being used to guide treatment decisions in a trial, these measures must be feasible. A joint count assessing 68 joints for tenderness and 66 joints for swelling is employed in virtually all randomized controlled trials (RCTs) to constitute the primary outcome measure and is endorsed by OMERACT [14]. However, the Disease Activity Score 28 (DAS28) developed for RA [30] is an often-used measure of disease activity and remission in PsA in clinical practice around the world and is a secondary measure used in RCTs. However, it was not developed for PsA patients, it is purely a measure of joint inflammation and confined to 28 joints and it does not assess disease in common domains of PsA involvement, i.e., distal interphalangeal joints, feet or ankles, skin, and nails.

Work is ongoing to compare different measures. The GRACE (GRAppa Composite Exercise) project aimed to develop new composite measures in PsA and compare them with existing indices [31]. The new indices included the psoriatic arthritis disease activity score (PASDAS) and the GRAPPA composite index, which uses the arithmetic mean of desirability functions. These have been compared with existing indices such the Composite Psoriatic arthritis Disease Activity Index (CPDAI), Disease Activity for PSoriatic Arthritis (DAPSA), and DAS28.

A recent study in patients with active PsA demonstrated that different remission criteria provide different results [32], while the performance of six composite activity indices was compared in a real-world study [33]; all six showed good discriminant capacity, but the proportions of patients classified in the disease activity levels differed and, in particular, the rate of patients in remission was clearly different among the indices. Of note, none of the existing composite measures, including MDA, capture the original (2006) [14] nor the updated (2016) [15, 16] PsA core set.

Is Treat to Target applicable in PsA?

The preferred 'target' (state) of a T2T approach is remission or inactive disease as the primary goal and low disease activity or MDA as the secondary goal.

The TICOPA (Tight Control of Psoriatic Arthritis) study [34] has recently shown that treating to target by escalating therapy, with a greater use of combination disease-modifying anti-rheumatic drugs (DMARDs) and biologics in the tight control arm of the study significantly improves joint outcomes for newly diagnosed patients (Fig. 1) [34]. In the Standard Care arm patients were reviewed every 12 weeks in a general rheumatology outpatient clinic supervised by a consultant rheumatologist. No formal measures of disease activity were used to guide treatment decisions and there was no restriction on prescribing. By contrast, in the Tight Control arm patients were seen every 4 weeks by the study physician and treated according to a predefined treatment protocol. At each visit, patients were assessed for MDA criteria. Those not achieving MDA had their treatment escalated to the maximum dose according to the protocol. Patients achieving the MDA criteria continued on their current therapy.

Patients who received tight control treatment did experience more treatment-related adverse and serious adverse events than those receiving standard care, reporting more colds, nausea, fatigue and gastrointestinal upsets than those in the control arm (only partly explained by more frequent visits and recording of adverse events). However, despite larger doses of methotrexate in the tight control arm, liver enzyme abnormalities were similar in both arms. Patients in the tight control arm also required 27% more tumor necrosis factor inhibitor (TNFi) usage compared with those on standard care.

Patients in the TICOPA study were selected for early disease, and current T2T concepts may be more appropriate for newly diagnosed patients, and may be more difficult to apply in patients with longer disease duration with relatively more damage. This damage may affect the optimal primary target as patients with longstanding disease may be unable to meet these stringent criteria.

TICOPA is the first trial of strategy in PsA and further strategy trials are needed to weigh effectiveness against safety, since adverse events were also higher in the tight control arm of the TICOPA study compared with the standard of care (SOC) arm.

In two additional studies, a delay in diagnosis and intervention by 6 months demonstrated an impact on structural damage and long-term functional outcomes [35-37]. Data from the Swedish Early Psoriatic Arthritis Register (SwePsA) [38] suggest that a shorter time between onset of symptoms and diagnosis is associated with better clinical outcomes at 5 years. It therefore appears that, as is the case with RA, early intervention combined with a tight control strategy is important to prevent irreversible damage.

Insights from Modern Imaging

Much less information is available on the use of ultrasound (US) and magnetic resonance imaging (MRI) in PsA compared with RA, and imaging outcomes for remission in PsA are still evolving. What is clear is that MRI and US have the potential to improve PsA management [39]. Both techniques offer capability for assessing both inflammation and damage, with MRI enabling visualization of the spine in axial disease. Both may evaluate peripheral joints, with US being more patient friendly while providing multiple joint examinations in real time, though it is unable to visualize intra-bone pathology (osteitis). MRI can evaluate only one joint or a joint area during one session, and may be less acceptable to patients due to the enclosed nature of the technique.

Although there are no typical US patterns characterizing PsA synovitis, with the exception of possibly more intense intra-articular vascularization seen in inflamed tissue, US has demonstrated good accuracy in assessing synovitis in PsA [10, 39-42]. In addition, the presence of US-detected synovitis has been shown to be associated with long-term radiographic erosion progression and poor outcomes [23]. Recently, Ficjan et al. [11] in a prospective and longitudinal study, developed an US composite score for the assessment of inflammatory and structural lesions in PsA, which demonstrates good metric properties including good sensitivity to change. US has also shown to be of added value in assessing enthesitis and dactylitis. US can also be used for visualizing structural changes and inflammatory activity at the psoriatic skin and nail level; thickening of both the epidermis and dermis is the most constant US pathologic finding in psoriatic plaques, whereas the hypoechoic band in the upper dermis is associated with Power Doppler (PD) activity (an expression of neoangiogenesis) and is particularly detectable in the active stages of the disease [43, 44].

Recommendations on imaging in spondyloarthritis (SpA) have been proposed by EULAR, including use of X-rays, US or MRI [45]. In axial SpA, the recommendation is for disease activity to be monitored with MRI of the sacroiliac (SI) joints and/or the spine, whereas conventional radiography should be used for long-term monitoring of structural damage. Similarly, for peripheral SpA, the recommendation is for US and MRI to be considered when monitoring disease activity (particularly synovitis and enthesitis), and conventional radiography is recommended to monitor structural damage.

The EULAR recommendations reflect the benefits of advanced imaging in assessing inflammation rather than assessing damage on X-rays, which has previously been an issue for trials conducted over short periods of time and trials that are not placebo controlled, where there is little radiographic structural progression. The recommendations and recent evidence from clinical trials suggest that the field could be moving towards a time when X-rays are of limited value for imaging in SpA clinical studies.

Using imaging to monitor disease activity

Multiple studies have shown that MRI and US can detect inflammatory and structural lesions [46] and identify risk factors for poor prognosis in PsA [39, 47]. In terms of quantifying change, most US composite scores have been developed for the assessment of inflammatory and structural lesions in PsA (in terms of quantifying change), and they have demonstrated construct validity, sensitivity to change, reliability and feasibility [11]. The OMERACT PsA MRI Score (PsAMRIS) has similarly demonstrated good performance metrics [11, 48]. Several studies have now demonstrated the use of imaging to monitor disease activity and therapeutic response. A study of more than 300 SpA patients being treated with TNFi showed that PD US is a reliable method to monitor therapeutic response by measuring enthesitis [49], while US had a pivotal role in differential diagnosis and treatment monitoring in a patient with early PsA undergoing an aggressive tight control treatment program and being monitored by US [50]. Similarly MRI has demonstrated responsiveness in PsA clinical studies [51].

Imaging of subclinical disease and remission

In line with the concept of subclinical disease first described in RA (inflammation detected by modern imaging but not examination), studies have found discrepancies between modern imaging and clinical findings, uncovering issues with accurate detection and clinical assessment of inflammation [9] and enthesitis, tenosynovitis or perisynovitis (i.e., extracapsular inflammation) in PsA patients in clinical remission [52]. In a study of newly diagnosed PsA comparing clinical examination with US in 49 patients, three-quarters were found to have sub-clinical synovitis, most frequently in the wrist and knee (Fig. 2) [53]. In patients on treatment, subclinical synovitis has been detected using US in patients classified as being in remission (as defined by MDA or DAS28) [9, 52]. There is some evidence that US detected synovitis might predict short-term flares in PsA patients in remission. However, it is not clear how important these US-detected manifestations really are and whether a T2T approach based on imaging would be superior to one based on clinical assessments. Some studies have shown that US can detect inflammatory and structural lesions and identify risk factors for poor prognosis in PsA [47]. Most of the studies have found discrepancies between US and clinical findings, uncovering issues with accurate detection and assessment of inflammation [9] and enthesitis, tenosynovitis or perisynovitis in PsA patients in clinical remission [52].

Enthesitis is another key, but often underestimated, feature of PsA, and therefore assessment of enthesitis with imaging is important, particularly as clinical measurements are often unreliable. Enthesitis may be predictive of flares, can predict clinical outcome, and can be present, although at a lower level, in remission or low disease activity states [54, 55]. A number of studies have been published supporting the validity of US in the assessment of entheses [56-60]. A recent study in newly diagnosed PsA found that three-quarters had sub-clinical synovitis, most frequently in the wrist and knee (Fig. 2).

Using contrast-enhanced US to detect persistent joint inflammation among patients in clinical remission showed that this technique is sufficiently sensitive to identify the presence of synovitis and thereby monitor remission [61]. Although there are limited data, there is some evidence that US detected synovitis might predict short-term flares in PsA patients in remission [62]. While Power Doppler US assessment may have an important role in monitoring treatment, its use at every clinic visit may not be feasible due to expertise required, time and financial constraints [63, 64]. However its use at specific time points where accurate assessment of inflammation is critical (e.g. evaluation of true remission state) may add value to usual care. Further developments such as whole-body MRI could be an additional tool for use in clinical decision making, allowing the assessment of disease activity in axial and peripheral sites, and improving the detection of inflammatory changes in PsA in locations that are difficult to assess clinically [65]. Again, feasibility is an important consideration given the equipment required and the costs associated with scanning.

Serological and Immunological Aspects of Remission in PsA

Two hypotheses have been formulated for the pathogenesis of PsA: firstly, that PsA is a classic autoimmune disease, or alternatively, that it begins with microtrauma at the enthesis, which then initiates innate immune events. [66]. A better understanding of the key pathologic pathways that drive progression from skin to bone involvement is needed in order to develop more effective treatment strategies.

Several studies on the origins of PsA have revealed signs of subclinical synovitis and enthesitis by MRI and US examination in the joints of patients who have psoriasis but not PsA [67-70], although the significance of these findings is not clear. Enthesitis has also been documented in healthy controls [59], and in patients with psoriasis without arthritis [71]. Psoriasis patients also have a greater risk of developing enthesophytes than healthy controls [72]. There is also evidence to suggest that skin-bone interactions are triggered by IL-17, and interleukin-17 (IL-17) overexpression in mice with chronic skin inflammation induces bone loss through inhibition of osteoblast-mediated bone formation [73]. Finally, recent data show that body mass index (BMI) may also have an effect on the development of enthesitis, with overweight patients having less chance of fulfilling MDA criteria for tender enthesial points [74].

There may be differences in the pathologies of the various phenotypes of PsA, in terms of presence of certain cytokines/immune cells in synovitis and enthesitis; for example, T-cell concentration changes or abnormalities in early disease may be predictive of progression and/or response to therapy [75]. Genetic factors, such as IL-23R polymorphisms, may also predispose to exaggerated cytokine production and a hyperproliferative response, which can combine with mechanical stress factors into clinically apparent skin disease and clinically unapparent enthesial proliferation [76].

Biochemical markers of inflammation

The concept of immunological remission in PsA is only beginning to be understood. Standard biomarkers of inflammation are not particularly helpful in judging inflammatory disease activity in PsA. Unlike the situation in RA, there are few established biomarkers for immunological pathology in PsA. As the IL-17 pathway is integral in psoriasis and psoriatic disease [77], the IL-17–IL-23 pathway may provide more reliable markers for PsA in future and recently, changes in CD3+ T-cell expression in PsA synovium have been shown to correlate with clinical response to treatment [75]. Biomarkers are under review as part of the OMERACT/GRAPPA initiative and several new biomarkers for PsA have been proposed, including calprotectin, SAA and MRP, although none has been extensively validated to date. In the future, newer approaches such as proteomics may reveal better biomarkers of disease activity for PsA.

Conclusions

Advances in the treatment of PsA, particularly the introduction of biologic therapies, have allowed the disease to be controlled in many patients; however, measuring response to treatment in PsA patients is widely debated, partly caused by the heterogeneity of the disease. Changes have also occurred in treatment strategies for rheumatic diseases and the development of Treat to Target approaches have led to a change in the established treatment paradigm.

Both MRI and US techniques have the potential to improve PsA management, and imaging outcomes for remission in PsA are still evolving. The concept of immunological remission in PsA is only just beginning to be discussed and biomarkers for the disease are yet to be fully identified.

While remission is the ultimate goal for PsA patients and their physicians, questions on what exactly we should aim to achieve still remain; this review has examined the current status of targeting remission in PsA, with a focus on areas that need more research.

Figures





FIG 1: Proportion of patients achieving an ACR response at 48 weeks in TICOPA (tight control vs SOC) [34]



411 *FIG 2: Subclinical synovitis in 49 patients with early PsA [53].*

412 *A: US positive, clinical exam negative; B: US negative, clinical exam positive.*

413 © 2014 The Authors. Arthritis Care & Research is published by Wiley Periodicals, Inc. on behalf of the American

414 College of Rheumatology.

Instrument:	MDA [7, 29, 55]	PASDAS [31]	CPDAI [78, 79]	DAPSA [80, 81]	RAPID 3 [82]	PsAID* [83]	DAS28 [30]
Developed for PsA?	Yes	Yes	Yes	Yes	No - Generic	Yes	No - RA
Approx time for patient to perform	2-5 mins	2-5 mins	5-10 mins	1-2 mins	2-5 mins	2-5 mins	1-2 mins
Approx time for assessor to perform	5-10 min	5-10 min	5-10 min	5-10 min	1 min	1 min	3-5 min
Complex calculation required	No	Yes	No	No	No	No	Yes
Continuous measure of disease activity	No	Yes	Yes	Yes	Yes	Impact not activity*	Yes
Measures peripheral arthritis	Yes	Yes	Yes	Yes	No	Impact*	Yes
Measures enthesitis	Yes	Yes	Yes	No	No	Impact*	No
Measures skin disease	Yes	Within global only	Yes	No	No	Impact*	No
Sensitive to change in PsA	Yes	Yes	Yes	Yes	Yes	Yes	Polyarticular only
Additional comments		Requires SF-36 and CRP	Development not evidence based	Requires CRP Cutoffs based on physician opinion only, peripheral arthritis only	No physician exam	Impact measure rather than activity	Peripheral arthritis only and measures only 28 joints.

TABLE 1. Comparison of features of clinical remission instruments

416

- 417 • PsAID measures the impact of the disease on the patient rather than disease activity but identifies impact of the disease in many
418 domains including MSK and skin.

419

420

421 **Acknowledgements**

422 The authors thank Tina Patrick, PhD, of Novartis Ireland Ltd for providing medical writing support with
423 the drafting of this manuscript, which was funded by Novartis in accordance with Good Publication
424 Practice (GPP3) guidelines (<http://www.ismpp.org/gpp3>).

425 **Conflict of interest statement**

426 LCC: Advisory boards or consultancies or speaker fees for Abbvie, BMS, Celgene, Janssen, Lilly, MSD,
427 Novartis, Pfizer, Sun Pharma, UCB

428 PGC: Advisory boards or consultancies for Abbvie, BMS, Eli Lilly, Novartis, Pfizer, Roche

429 MADA: Advisory boards or consultancies or speaker's fees for Abbvie, BMS, Novartis, Roche

430 MdW: No conflicts to disclose

431 OF: Advisory boards, grants or consultancies for Abbvie, BMS, Celgene, Eli Lilly, Novartis, Pfizer, Janssen,
432 UCB.

433 TK: Tore K Kvien has received fees for speaking and/or consulting from AbbVie, Biogen, BMS, Boehringer
434 Ingelheim, Celgene, Celltrion, Eli Lilly, Epirus, Hospira, Merck-Serono, MSD, Mundipharma, Novartis,
435 Oktal, Orion Pharma, Hospira/Pfizer, Roche, Sandoz and UCB and received research funding to
436 Diakonhjemmet Hospital from AbbVie, BMS, MSD, Pfizer, Roche and UCB.

437 RL: Leuven R&D, KU Leuven's Knowledge and Technology Transfer Office, has received consultancy and
438 speaker's fees from Abbvie, Celgene, Janssen, Novartis and Pfizer and research support from Boehringer
439 Ingelheim, Celgene and Pfizer on behalf of Rik Lories.

440 PM: Research grants, consultation and/or speaker honoraria: Abbvie, Amgen, BMS, Celgene, Janssen,
441 Lilly, Novartis, Pfizer, SUN, UCB.

442 PN: Consultancy – AbbVie, Amgen, BMS, Celgene, Lilly, Hospira, MSD, Pfizer, Janssen, UCB, Novartis,
443 Roche; Novartis Sanofi; Speaker bureau – AbbVie, Amgen, BMS, Celgene, Lilly, Hospira, MSD, Pfizer,
444 Janssen, UCB, Novartis, Roche; Research funding – AbbVie, Amgen, BMS, Celgene, Lilly, Hospira, MSD,
445 Pfizer, Janssen, UCB, Novartis, Roche, Sanofi.

446 GS: No conflicts to disclose

447 ERS: ER Soriano has received fees for speaking and/or consulting from AbbVie, BMS, Janssen, Novartis,
448 Pfizer, Roche, and UCB and received research funding to Hospital Italiano de Buenos Aires from AbbVie,
449 BMS, Janssen, Novartis, Pfizer, Roche and UCB.

450 PE: Paul Emery has undertaken clinical trials and provided expert advice to Pfizer, MSD, Abbvie, BMS,
451 UCB, Roche, Novartis, Samsung, Sandoz and Lilly

Funding

PGC and PE are supported in part by the National Institute for Health Research (NIHR) Leeds Biomedical Research Centre. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

References

- 1 Coates LC, Kavanaugh A, Mease PJ, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis 2015 Treatment Recommendations for Psoriatic Arthritis. *Arthritis & rheumatology* (Hoboken, N.J.) 2016;68(5):1060-71.
- 2 Gossec L, Smolen JS, Ramiro S, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2015 update. *Annals of the rheumatic diseases* 2016;75(3):499-510.
- 3 Smolen JS, Braun J, Dougados M, et al. Treating spondyloarthritis, including ankylosing spondylitis and psoriatic arthritis, to target: recommendations of an international task force. *Annals of the rheumatic diseases* 2014;73(1):6-16.
- 4 Gottlieb A, Korman NJ, Gordon KB, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 2. Psoriatic arthritis: overview and guidelines of care for treatment with an emphasis on the biologics. *Journal of the American Academy of Dermatology* 2008;58(5):851-64.
- 5 Smolen JS. Treat-to-target as an approach in inflammatory arthritis. *Current opinion in rheumatology* 2016;28(3):297-302.
- 6 Grigor C, Capell H, Stirling A, et al. Effect of a treatment strategy of tight control for rheumatoid arthritis (the TICORA study): a single-blind randomised controlled trial. *Lancet* 2004;364(9430):263-9.
- 7 Coates LC, Fransen J, Helliwell PS. Defining minimal disease activity in psoriatic arthritis: a proposed objective target for treatment. *Annals of the rheumatic diseases* 2010;69(1):48-53.
- 8 Schoels MM, Braun J, Dougados M, et al. Treating axial and peripheral spondyloarthritis, including psoriatic arthritis, to target: results of a systematic literature search to support an international treat-to-target recommendation in spondyloarthritis. *Annals of the rheumatic diseases* 2014;73(1):238-42.
- 9 Michelsen B, Diamantopoulos AP, Hammer HB, Soldal DM, Kavanaugh A, Haugeberg G. Ultrasonographic evaluation in psoriatic arthritis is of major importance in evaluating disease activity. *Annals of the rheumatic diseases* 2016;75(12):2108-13.
- 10 Wiell C, Szkudlarek M, Hasselquist M, et al. Ultrasonography, magnetic resonance imaging, radiography, and clinical assessment of inflammatory and destructive changes in fingers and toes of patients with psoriatic arthritis. *Arthritis research & therapy* 2007;9(6):R119.
- 11 Ficjan A, Husic R, Gretler J, et al. Ultrasound composite scores for the assessment of inflammatory and structural pathologies in Psoriatic Arthritis (PsASon-Score). *Arthritis research & therapy* 2014;16(5):476.
- 12 Curran SA, FitzGerald OM, Costello PJ, et al. Nucleotide sequencing of psoriatic arthritis tissue before and during methotrexate administration reveals a complex inflammatory T cell infiltrate with very few clones exhibiting features that suggest they drive the inflammatory process by recognizing autoantigens. *Journal of immunology* 2004;172(3):1935-44.
- 13 Kavanaugh A, Fransen J. Defining remission in psoriatic arthritis. *Clinical and experimental rheumatology* 2006;24(6 Suppl 43):S-83-7.

- 14 Gladman DD, Mease PJ, Strand V, et al. Consensus on a core set of domains for psoriatic arthritis. *The Journal of rheumatology* 2007;34(5):1167-70.
- 15 Orbai AM, Mease PJ, de Wit M, et al. Report of the GRAPPA-OMERACT Psoriatic Arthritis Working Group from the GRAPPA 2015 Annual Meeting. *The Journal of rheumatology* 2016;43(5):965-9.
- 16 Orbai AM, de Wit M, Mease P, et al. International patient and physician consensus on a psoriatic arthritis core outcome set for clinical trials. *Annals of the rheumatic diseases* 2016.
- 17 Coates LC, Helliwell PS. Defining Low Disease Activity States in Psoriatic Arthritis using Novel Composite Disease Instruments. *The Journal of rheumatology* 2016;43(2):371-5.
- 18 Haroon M, FitzGerald O. Psoriatic arthritis: complexities, comorbidities and implications for the clinic. *Expert Rev Clin Immunol* 2016;12(4):405-16.
- 19 Haroon M, Gallagher P, Heffernan E, FitzGerald O. High prevalence of metabolic syndrome and of insulin resistance in psoriatic arthritis is associated with the severity of underlying disease. *The Journal of rheumatology* 2014;41(7):1357-65.
- 20 Husted JA, Thavaneswaran A, Chandran V, Gladman DD. Incremental effects of comorbidity on quality of life in patients with psoriatic arthritis. *The Journal of rheumatology* 2013;40(8):1349-56.
- 21 Eder L, Thavaneswaran A, Chandran V, Cook R, Gladman DD. Factors explaining the discrepancy between physician and patient global assessment of joint and skin disease activity in psoriatic arthritis patients. *Arthritis care & research* 2015;67(2):264-72.
- 22 Barton JL, Imboden J, Graf J, Glidden D, Yelin EH, Schillinger D. Patient-physician discordance in assessments of global disease severity in rheumatoid arthritis. *Arthritis care & research* 2010;62(6):857-64.
- 23 Helliwell P, Coates L, Chandran V, et al. Qualifying unmet needs and improving standards of care in psoriatic arthritis. *Arthritis Care Res (Hoboken)* 2014;66(12):1759-66.
- 24 Michelsen B, Kristianslund EK, Hammer HB, et al. Discordance between tender and swollen joint count as well as patient's and evaluator's global assessment may reduce likelihood of remission in patients with rheumatoid arthritis and psoriatic arthritis: data from the prospective multicentre NOR-DMARD study. *Ann Rheum Dis* 2016.
- 25 Lebwohl MG, Bachelez H, Barker J, et al. Patient perspectives in the management of psoriasis: results from the population-based Multinational Assessment of Psoriasis and Psoriatic Arthritis Survey. *J Am Acad Dermatol* 2014;70(5):871-81 e1-30.
- 26 Stamm TA, Nell V, Mathis M, et al. Concepts important to patients with psoriatic arthritis are not adequately covered by standard measures of functioning. *Arthritis and rheumatism* 2007;57(3):487-94.
- 27 van Tuyl LH, Hewlett S, Sadlonova M, et al. The patient perspective on remission in rheumatoid arthritis: 'You've got limits, but you're back to being you again'. *Annals of the rheumatic diseases* 2015;74(6):1004-10.
- 28 Kirkham B, de Vlam K, Li W, et al. Early treatment of psoriatic arthritis is associated with improved patient-reported outcomes: findings from the etanercept PRESTA trial. *Clinical and experimental rheumatology* 2015;33(1):11-9.
- 29 Soriano ER. Defining remission in psoriatic arthritis: are we getting closer? *The Journal of rheumatology* 2015;42(6):907-8.
- 30 van Gestel AM, Haagsma CJ, van Riel PL. Validation of rheumatoid arthritis improvement criteria that include simplified joint counts. *Arthritis and rheumatism* 1998;41(10):1845-50.
- 31 Helliwell PS, FitzGerald O, Fransen J, et al. The development of candidate composite disease activity and responder indices for psoriatic arthritis (GRACE project). *Annals of the rheumatic diseases* 2013;72(6):986-91.
- 32 Van den Bosch F, Kavanaugh A, Kron M, Kupper H, Mease PJ. Clinical remission in patients with active psoriatic arthritis treated with adalimumab and correlations in joint and skin manifestations. *The Journal of rheumatology* 2015;42(6):952-9.

- 33 Salaffi F, Ciapetti A, Carotti M, Gasparini S, Gutierrez M. Disease activity in psoriatic arthritis: comparison of the discriminative capacity and construct validity of six composite indices in a real world. *BioMed research international* 2014;2014:528105.
- 34 Coates LC, Moverley AR, McParland L, et al. Effect of tight control of inflammation in early psoriatic arthritis (TICOPA): a UK multicentre, open-label, randomised controlled trial. *Lancet* 2015;386(10012):2489-98.
- 35 Haroon M, Gallagher P, FitzGerald O. Diagnostic delay of more than 6 months contributes to poor radiographic and functional outcome in psoriatic arthritis. *Ann Rheum Dis* 2015;74(6):1045-50.
- 36 Tillett W, Jadon D, Shaddick G, et al. Smoking and delay to diagnosis are associated with poorer functional outcome in psoriatic arthritis. *Annals of the Rheumatic Diseases* 2013;72(8):1358-61.
- 37 Gladman DD, Thavaneswaran A, Chandran V, Cook RJ. Do patients with psoriatic arthritis who present early fare better than those presenting later in the disease? *Annals of the rheumatic diseases* 2011;70(12):2152-4.
- 38 Theander E, Husmark T, Alenius GM, et al. Early psoriatic arthritis: short symptom duration, male gender and preserved physical functioning at presentation predict favourable outcome at 5-year follow-up. Results from the Swedish Early Psoriatic Arthritis Register (SwePsA). *Annals of the rheumatic diseases* 2014;73(2):407-13.
- 39 Coates LC, Hodgson R, Conaghan PG, Freeston JE. MRI and ultrasonography for diagnosis and monitoring of psoriatic arthritis. *Best Pract Res Clin Rheumatol* 2012;26(6):805-22.
- 40 Gutierrez M, Filippucci E, De Angelis R, Filosa G, Kane D, Grassi W. A sonographic spectrum of psoriatic arthritis: "the five targets". *Clin Rheumatol* 2010;29(2):133-42.
- 41 De Simone C, Caldarola G, D'Agostino M, et al. Usefulness of ultrasound imaging in detecting psoriatic arthritis of fingers and toes in patients with psoriasis. *Clin Dev Immunol* 2011;2011:390726.
- 42 Sankowski AJ, Lebkowska UM, Cwikla J, Walecka I, Walecki J. The comparison of efficacy of different imaging techniques (conventional radiography, ultrasonography, magnetic resonance) in assessment of wrist joints and metacarpophalangeal joints in patients with psoriatic arthritis. *Pol J Radiol* 2013;78(1):18-29.
- 43 Gutierrez M, De Angelis R, Bernardini ML, et al. Clinical, power Doppler sonography and histological assessment of the psoriatic plaque: short-term monitoring in patients treated with etanercept. *Br J Dermatol* 2011;164(1):33-7.
- 44 Gutierrez M, Wortsman X, Filippucci E, De Angelis R, Filosa G, Grassi W. High-frequency sonography in the evaluation of psoriasis: nail and skin involvement. *J Ultrasound Med* 2009;28(11):1569-74.
- 45 Mandl P, Navarro-Compan V, Terslev L, et al. EULAR recommendations for the use of imaging in the diagnosis and management of spondyloarthritis in clinical practice. *Ann Rheum Dis* 2015;74(7):1327-39.
- 46 Kaeley GS. Review of the use of ultrasound for the diagnosis and monitoring of enthesitis in psoriatic arthritis. *Current rheumatology reports* 2011;13(4):338-45.
- 47 El Miedany Y, El Gaafary M, Youssef S, Ahmed I, Nasr A. Tailored approach to early psoriatic arthritis patients: clinical and ultrasonographic predictors for structural joint damage. *Clinical rheumatology* 2015;34(2):307-13.
- 48 Boyesen P, McQueen FM, Gandjbakhch F, et al. The OMERACT Psoriatic Arthritis Magnetic Resonance Imaging Score (PsAMRIS) is reliable and sensitive to change: results from an OMERACT workshop. *J Rheumatol* 2011;38(9):2034-8.
- 49 Naredo E, Batlle-Gualda E, Garcia-Vivar ML, et al. Power Doppler ultrasonography assessment of entheses in spondyloarthropathies: response to therapy of enthesal abnormalities. *The Journal of rheumatology* 2010;37(10):2110-7.

- 50 Bandinelli F, Bonacci E, Matucci-Cerinic M. Ultrasound-integrated tight control in early psoriatic arthritis during adalimumab treatment. *Clinical and experimental rheumatology* 2013;31(3):440-2.
- 51 Glinatsi D, Bird P, Gandjbakhch F, et al. Validation of the OMERACT Psoriatic Arthritis Magnetic Resonance Imaging Score (PsAMRIS) for the Hand and Foot in a Randomized Placebo-controlled Trial. *J Rheumatol* 2015;42(12):2473-9.
- 52 Husic R, Gretler J, Felber A, et al. Disparity between ultrasound and clinical findings in psoriatic arthritis. *Annals of the rheumatic diseases* 2014;73(8):1529-36.
- 53 Freeston JE, Coates LC, Nam JL, et al. Is there subclinical synovitis in early psoriatic arthritis? A clinical comparison with gray-scale and power Doppler ultrasound. *Arthritis care & research* 2014;66(3):432-9.
- 54 Perrotta FM, Astorri D, Zappia M, Reginelli A, Brunese L, Lubrano E. An ultrasonographic study of enthesitis in early psoriatic arthritis patients naive to traditional and biologic DMARDs treatment. *Rheumatol Int* 2016;36(11):1579-83.
- 55 Marin J, Acosta-Felquer ML, Ferreyra-Garrot L, Ruta S, Rosa J, Soriano ER. Patients with Psoriatic Arthritis Fulfilling the Minimal Disease Activity Criteria Do Not Have Swollen and Tender Joints, but Have Active Skin. *The Journal of rheumatology* 2016;43(5):907-10.
- 56 Filippucci E, Aydin SZ, Karadag O, et al. Reliability of high-resolution ultrasonography in the assessment of Achilles tendon enthesopathy in seronegative spondyloarthropathies. *Ann Rheum Dis* 2009;68(12):1850-5.
- 57 de Miguel E, Munoz-Fernandez S, Castillo C, Cobo-Ibanez T, Martin-Mola E. Diagnostic accuracy of enthesitis ultrasound in the diagnosis of early spondyloarthritis. *Ann Rheum Dis* 2011;70(3):434-9.
- 58 D'Agostino MA, Aegerter P, Bechara K, et al. How to diagnose spondyloarthritis early? Accuracy of peripheral enthesitis detection by power Doppler ultrasonography. *Ann Rheum Dis* 2011;70(8):1433-40.
- 59 D'Agostino MA, Said-Nahal R, Hacquard-Bouder C, Brasseur JL, Dougados M, Breban M. Assessment of peripheral enthesitis in the spondylarthropathies by ultrasonography combined with power Doppler: a cross-sectional study. *Arthritis and rheumatism* 2003;48(2):523-33.
- 60 Wang CH, Feng Y, Ren Z, et al. Performance of ultrasound to monitor Achilles enthesitis in patients with ankylosing spondylitis during TNF- α antagonist therapy. *Clin Rheumatol* 2015;34(6):1073-8.
- 61 Bonifati C, Elia F, Graceffa D, et al. Clinical and Contrast-Enhanced Ultrasound Echography Outcomes in Psoriatic Arthritis Patients after One Year of Continuous Therapy with Anti-TNF Drugs. *ISRN Dermatol* 2014;2014:932721.
- 62 Marin JA-F, M.L.; Ferreyra-Garrot, L.; Catay, E.; Rosa, J.; Ruta, S.; Sabelli, M.; Garcia-Monaco, R.; Soriano, E.R. Utility of Power Doppler Ultrasound-Detected Synovitis for the Prediction of Short Term Flare in Psoriatic Arthritis Patients in Clinical Remission [abstract]. *Arthritis Rheumatol* 2015; 67 ((suppl 10)).
- 63 Diaz-Torne C, Moragues C, Toniolo E, et al. Impact of ultrasonography on treatment decision in rheumatoid arthritis: the IMPULSAR study. *Rheumatol Int* 2017.
- 64 Naredo E, Rodriguez M, Campos C, et al. Validity, reproducibility, and responsiveness of a twelve-joint simplified power doppler ultrasonographic assessment of joint inflammation in rheumatoid arthritis. *Arthritis and rheumatism* 2008;59(4):515-22.
- 65 Weckbach S, Schewe S, Michaely HJ, Steffinger D, Reiser MF, Glaser C. Whole-body MR imaging in psoriatic arthritis: additional value for therapeutic decision making. *Eur J Radiol* 2011;77(1):149-55.
- 66 FitzGerald O, Haroon M, Giles JT, Winchester R. Concepts of pathogenesis in psoriatic arthritis: genotype determines clinical phenotype. *Arthritis research & therapy* 2015;17:115.
- 67 Naredo E, Moller I, de Miguel E, et al. High prevalence of ultrasonographic synovitis and enthesopathy in patients with psoriasis without psoriatic arthritis: a prospective case-control study. *Rheumatology (Oxford, England)* 2011;50(10):1838-48.

- 68 Gisondi P, Tinazzi I, El-Dalati G, et al. Lower limb enthesopathy in patients with psoriasis without clinical signs of arthropathy: a hospital-based case-control study. *Annals of the rheumatic diseases* 2008;67(1):26-30.
- 69 Offidani A, Cellini A, Valeri G, Giovagnoni A. Subclinical joint involvement in psoriasis: magnetic resonance imaging and X-ray findings. *Acta Derm Venereol* 1998;78(6):463-5.
- 70 Faustini F, Simon D, Oliveira I, et al. Subclinical joint inflammation in patients with psoriasis without concomitant psoriatic arthritis: a cross-sectional and longitudinal analysis. *Annals of the rheumatic diseases* 2016;75(12):2068-74.
- 71 Moshrif A, Mosallam A, Mohamed EE, Gouda W, Doma M. Subclinical enthesopathy in patients with psoriasis and its association with other disease parameters: a power Doppler ultrasonographic study. *Eur J Rheumatol* 2017;4(1):24-8.
- 72 Simon D, Faustini F, Kleyer A, et al. Analysis of periarticular bone changes in patients with cutaneous psoriasis without associated psoriatic arthritis. *Annals of the rheumatic diseases* 2016;75(4):660-6.
- 73 Uluckan O, Wagner EF. Role of IL-17A signalling in psoriasis and associated bone loss. *Clinical and experimental rheumatology* 2016;34(4 Suppl 98):17-20.
- 74 Eder L, Thavaneswaran A, Chandran V, Cook RJ, Gladman DD. Obesity is associated with a lower probability of achieving sustained minimal disease activity state among patients with psoriatic arthritis. *Annals of the rheumatic diseases* 2015;74(5):813-7.
- 75 Pontifex EK, Gerlag DM, Gogarty M, et al. Change in CD3 positive T-cell expression in psoriatic arthritis synovium correlates with change in DAS28 and magnetic resonance imaging synovitis scores following initiation of biologic therapy--a single centre, open-label study. *Arthritis Res Ther* 2011;13(1):R7.
- 76 Cargill M, Schrodli SJ, Chang M, et al. A large-scale genetic association study confirms IL12B and leads to the identification of IL23R as psoriasis-risk genes. *Am J Hum Genet* 2007;80(2):273-90.
- 77 Durham LE, Kirkham BW, Taams LS. Contribution of the IL-17 Pathway to Psoriasis and Psoriatic Arthritis. *Current rheumatology reports* 2015;17(8):55.
- 78 Ritchlin CT, Kavanaugh A, Gladman DD, et al. Treatment recommendations for psoriatic arthritis. *Annals of the rheumatic diseases* 2009;68(9):1387-94.
- 79 Mumtaz A, Gallagher P, Kirby B, et al. Development of a preliminary composite disease activity index in psoriatic arthritis. *Annals of the rheumatic diseases* 2011;70(2):272-7.
- 80 Schoels MM, Aletaha D, Alasti F, Smolen JS. Disease activity in psoriatic arthritis (PsA): defining remission and treatment success using the DAPSA score. *Annals of the rheumatic diseases* 2016;75(5):811-8.
- 81 Aletaha D, Alasti F, Smolen JS. Disease activity states of the DAPSA, a psoriatic arthritis specific instrument, are valid against functional status and structural progression. *Annals of the rheumatic diseases* 2016.
- 82 Pincus T, Bergman MJ, Yazici Y. RAPID3-an index of physical function, pain, and global status as "vital signs" to improve care for people with chronic rheumatic diseases. *Bulletin of the NYU hospital for joint diseases* 2009;67(2):211-25.
- 83 Gossec L, de Wit M, Kiltz U, et al. A patient-derived and patient-reported outcome measure for assessing psoriatic arthritis: elaboration and preliminary validation of the Psoriatic Arthritis Impact of Disease (PsAID) questionnaire, a 13-country EULAR initiative. *Ann Rheum Dis* 2014;73(6):1012-9.