

15th Water-Rock Interaction International Symposium, WRI-15

## Tritium Tracers of Rapid Surface Water Ingression into Arsenic-Bearing Aquifers in the Lower Mekong Basin, Cambodia

Laura A Richards<sup>a,1</sup>, Jürgen Sültenfuß<sup>b</sup>, Christopher J Ballentine<sup>c</sup>, Daniel Magnone<sup>a</sup>, Bart E van Dongen<sup>a</sup>, Chansopheaktra Sovann<sup>d</sup>, David A Polya<sup>a</sup>

<sup>a</sup> School of Earth, Atmospheric and Environmental Sciences and Williamson Research Centre for Molecular Environmental Science, University of Manchester, Manchester M13 9PL, UK

<sup>b</sup> Institute of Environmental Physics, University of Bremen, Bremen 28359, Germany

<sup>c</sup> Department of Earth Sciences, University of Oxford, Oxford OX1 3AN, UK

<sup>d</sup> Department of Environmental Sciences, Royal University of Phnom Penh; Cambodia

---

### Abstract

Arsenic (As) contamination of groundwaters in South and Southeast Asia is a major threat to public health in these areas. Understanding the source and age of the groundwaters is critically important to understanding the controls on As mobilization in these aquifers. Using tritium ( $^3\text{H}$ ) and noble gas (He and Ne) signatures, model groundwater ages and dominant hydrological controls were identified in a transect oriented broadly parallel to inferred groundwater flowpaths in Kandal Province, Cambodia in the lower Mekong Basin. Apparent  $^3\text{H}$ - $^3\text{He}$  ages showed that most groundwaters are modern (< 55 years), indicating relatively fast recharge even in the absence of large-scale groundwater abstraction. The age-depth relationship indicates a strong vertical component of groundwater flow and allows for recharge rates to be estimated. Vertical and horizontal flow velocities are heterogeneous and site-specific. The conceptual framework will be used to better understand As mobilization and subsequent transport with these and similar aquifers.

© 2017 Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of the organizing committee of WRI-15

**Keywords:** tritium, arsenic, groundwater age, environmental pollutants, Cambodia

---

### 1. Introduction

Millions of people in South/Southeast Asia chronically consume groundwater containing dangerous concentrations of naturally-occurring  $\text{As}^{1-4}$ . As release in shallow aquifers in this region is widely attributed to the

---

\* Corresponding author. Tel.: +44 (0)161 306 0361; fax: +44(0)161 306 9361.

E-mail address: [laura.richards@manchester.ac.uk](mailto:laura.richards@manchester.ac.uk)

reductive dissolution of As-bearing Fe(III) minerals<sup>5</sup>, driven by metal reducing bacteria and bioavailable organic matter<sup>2,5-7</sup>. The nature of the organic matter implicated in As mobilization is generally thought to derive, in some proportion, from (i) plant-derived organic matter internal to aquifers<sup>8,9</sup>; (ii) external, modern surface-derived organic matter, largely from ponds, rivers and rice paddies<sup>10-18</sup>; and/or (iii) petroleum-derived hydrocarbons<sup>7,19,20</sup>. Debates surrounding the sub-surface location where As mobilization takes place and the subsequent controls on As mobility<sup>10,14,15,17, 21-25</sup> are intrinsically linked to the type and amount of the organic matter implicated in As release<sup>7-10,14,15,19,20,26</sup>. Determining the relative importance of these various inputs is essential in predicting future changes in As hazard<sup>10,15,27</sup>. The aim is to characterize the  $^3\text{H}$  and noble gas (He and Ne) isotopic signature of As-affected groundwater to (i) determine groundwater age; (ii) examine dominant hydrological controls; and (iii) assess local heterogeneity, which ultimately may affect As mobilization and transport in these aquifers.

## 2. Methods and Materials

The field sites are located in Kien Svay, Kandal Province, Cambodia, an area heavily affected by groundwater As<sup>11,12,14,15,19,27-30</sup> and representative of pre-development conditions. Groundwater samples (6 - 45 m deep) were collected from flushed and developed wells<sup>31</sup> pre- (May – June 2014) and post-monsoon (Nov – Dec 2014). In-situ measurements of basic groundwater parameters and subsequent analysis of geochemical composition was conducted<sup>31</sup>. Duplicate samples for  $^3\text{H}$  analysis were collected in 1 L argon filled amber glass bottles. Duplicate samples for He and Ne analysis were collected in flushed soft copper tubes. Analysis of  $^3\text{H}$ , He and Ne isotopes was conducted at the “Helis – Helium Isotopes Studies Bremen” Noble Gas Laboratory (Institute of Environmental Physics, University of Bremen, Germany)<sup>32</sup>, with apparent  $^3\text{H}$ - $^3\text{He}$  ages derived using previous methods<sup>33</sup>.

## 3. Results and Discussion

### 3.1. $^3\text{H}$ Concentration of Groundwater and Surface Water

$^3\text{H}$  measurements of groundwaters, river water and rain range from 0.04 to 3.6 TU (Fig. 1A). Relatively high pre-monsoon  $^3\text{H}$  in the Mekong suggests an additional and variable upstream source for  $^3\text{H}$ , likely from higher altitude Himalayan precipitation<sup>34</sup>. Concentrations of ‘stable tritium’ (the sum of  $^3\text{H}$  and  $^3\text{He}_{\text{in}}$ ) are similar to the input function for Bangkok precipitation (Fig. 1B)<sup>35</sup>. The similarity in independent datasets provides verification of  $^3\text{H}$  sampling and analysis. For all samples (except  $\geq 30$  m depth at LR09), the ‘stable tritium’ follows the  $^3\text{H}$  input function for Bangkok closely, indicating that no old  $^3\text{H}$ -free water is admixed with young groundwater at most sites.




Fig. 1. (A)  $^3\text{H}$  for pre-/post-monsoon groundwaters, Mekong River and rain.  $^3\text{H}$  was  $> 0.02$  TU in all groundwater samples; (B) ‘Stable tritium’ (the sum of  $^3\text{H}$  and  $^3\text{He}_{\text{in}}$ ) values and the input function for Bangkok precipitation (WMO Code 4845500<sup>35</sup>) against infiltration date.

### 3.2. Apparent $^3\text{H}$ - $^3\text{He}$ Ages of Groundwater and Surface Water

Apparent  $^3\text{H}$ - $^3\text{He}$  ages show distinct spatial patterns (Fig. 2A). Ages are mostly modern (e.g.  $< 55$  years in age), with the exception of two sites (LR09-30 and LR01-45). Age increases with depth at every site, with the exception of site LR09. For the same depth there is age variation on the order of 10 – 20 years. The spatial variation of apparent ages allows for estimates of groundwater flow velocities. The overall increase in  $^3\text{H}$ - $^3\text{He}$  age with depth (Fig. 2B) has an overall gradient representing mean vertical flow velocity of  $0.5 \pm 0.1 \text{ m.yr}^{-1}$  ( $t(24) = 5.62$ ,  $p = 8.7 \cdot 10^{-6}$  for ages  $< 55$  years). Depth profiles at specific sites show localized and heterogeneous vertical flow velocities. The site-specific variability gives rise to the identification of relatively rapid recharge zones.




Fig. 2. (A) Apparent  $^3\text{H}$ - $^3\text{He}$  age (in years, 2014 basis) with increasing perpendicular distance from the Bassac River. Blue, labelled ages are pre-monsoon (orange post-monsoon). Dashed lines indicate ages > 55 years. The contour represents grain size (lighter colors are sandier); (B) Apparent  $^3\text{H}$ - $^3\text{He}$  age versus depth.

### 3.3. Implications on Arsenic Release

As concentrations are typically lower in very young waters and increase in deeper, older waters (**Fig. 3A**). This relationship allows for the calculation of an overall As loading rate of  $6.3 \pm 2.6 \mu\text{g}\cdot\text{L}^{-1}\cdot\text{yr}^{-1}$  ( $t(24) = 2.40$ ,  $p = 0.0246$ ). As loading is heterogeneous and site-specific, and is attributed in part to changes in lithology and the likely presence of bio-available carbon. Depth profiles allow for examination of the association of As with  $^3\text{H}$ , noble gases, Eh and dissolved organic carbon (DOC) (**Fig. 3B**). In addition to the general correlation of As with depth and  $^3\text{H}$ - $^3\text{He}$  age, and the inverse As/Eh correlation,  $^4\text{He}_{\text{rad}}$  increases with depth which is consistent with age.




Fig. 3. (A) As versus apparent  $^3\text{H}$ - $^3\text{He}$  age (basis 2014). Samples in the dashed circle are > 55 years old; (B) Depth profiles for site LR01.

## 4. Conclusions

The  $^3\text{H}$  composition of As-bearing groundwater was used to determine groundwater ages along groundwater flowpaths. Apparent  $^3\text{H}$ - $^3\text{He}$  ages showed that most groundwaters are modern and younger than previously considered. Consistency of ‘stable tritium’ with the input function for Bangkok precipitation provides independent verification of analysis. A strong depth-age relationship suggests the dominance of a vertical hydrological control. Ages can be used to estimate flow velocities and heterogeneity. Such hydrological controls provide the conceptual framework for an improved groundwater model and may affect As mobilization/transport with the aquifer.

## Acknowledgements

This research was funded by a NERC Standard Research Grant (NE/J023833/1) to DP, BvD and CB, a NERC PhD studentship (NE/L501591/1) to DM, and a Leverhulme Trust Early Career Fellowship (ECF2015-657) to LR. We are very grateful to: Chivuth Kong, Pheary Meas and Yut Yann (all Royal University of Agriculture, Phnom Penh), Chhenggunn Aing, Zongta Sang and Teyden Sok (all Royal University of Phnom Penh), Helen Downie (The University of Manchester) and Lee Chambers (The University of Lancaster) for fieldwork contributions. The local drilling team and landowners are thanked for project support. Ann Hall, Marc Hall, Lori Frees, Lori Allen and Dina Kuy (Resources Development International – Cambodia) are thanked for logistical support.

## References

1. Smedley PL, Kinniburgh DG. A review of the source, behaviour and distribution of arsenic in natural waters. *Applied Geochemistry*. 2002;17(5):517-68.
2. Charlet L, Polya DA. Arsenic in shallow, reducing groundwaters in Southern Asia: An environmental health disaster. *Elements*. 2006;2:91-6.
3. World Health Organization. Guidelines for Drinking-water Quality: Fourth Edition. World Health Organization, editor. Geneva 2011.
4. Ravenscroft P, Brammer H, Richards K. Arsenic pollution - A global synthesis. Chichester: Wiley-Blackwell; 2009. 588 p.
5. Islam FS, Gault AG, Boothman C, Polya DA, Charnock JM, Chatterjee D, et al. Role of metal-reducing bacteria in arsenic release from Bengal delta sediments. *Nature*. 2004;430:68-71.
6. Postma D, Larsen F, Minh Hue NT, Duc MT, Viet PH, Nhan PQ, et al. Arsenic in groundwater of the Red River floodplain, Vietnam: Controlling geochemical processes and reactive transport modeling. *Geochimica et Cosmochimica Acta*. 2007;71(21):5054-71.
7. Rowland HAL, Boothman C, Pancost RD, Gault AG, Polya DA, Lloyd JR. The role of indigenous microorganisms in the biodegradation of naturally occurring petroleum, the reduction of iron, and the mobilization of arsenite from west bengal aquifer sediments. *Journal of Environmental Quality*. 2009;38(4):1598 - 607.
8. Nickson RT, McArthur JM, Burgess WG, Ahmed KM, Ravenscroft P, Rahman M. Arsenic poisoning of Bangladesh groundwater. *Nature*. 1998;395:338.
9. McArthur JM, Banerjee DM, Hudson-Edwards KA, Mishra R, Purohit R, Ravenscroft P, et al. Natural organic matter in sedimentary basins and its relation to arsenic in anoxic ground water: the example of West Bengal and its worldwide implications. *Applied Geochemistry*. 2004;19(8):1255-93.
10. Harvey CF, Swartz CH, Badruzzaman ABM, Keon-Blute N, Yu W, Ashraf Ali M, et al. Arsenic mobility and groundwater extraction in Bangladesh. *Science*. 2002;298:1602-6.
11. Polizzotto ML, Kocar BD, Benner SG, Sampson M, Fendorf S. Near-surface wetland sediments as a source of arsenic release to ground water in Asia. *Nature*. 2008;454:505-8.
12. Kocar BD, Polizzotto ML, Benner SG, Ying SC, Ung M, Ouch K, et al. Integrated biogeochemical and hydrologic processes driving arsenic release from shallow sediments to groundwaters of the Mekong delta. *Applied Geochemistry*. 2008;23(11):3059-71.
13. Neumann RB, Ashfaq KN, Badruzzaman ABM, Ali MA, Shoemaker JK, Harvey CF. Anthropogenic influences on groundwater arsenic concentrations in Bangladesh. *Nature Geoscience*. 2010;3(1):46-52.
14. Lawson M, Polya DA, Boyce AJ, Bryant C, Mondal D, Shantz A, et al. Pond-derived organic carbon driving changes in arsenic hazard found in Asian groundwaters. *Environmental Science and Technology*. 2013;47:7085 - 94.
15. Lawson M, Polya DA, Boyce AJ, Bryant C, Ballentine CJ. Tracing organic matter composition and distribution and its role on arsenic release in shallow Cambodian groundwaters. *Geochimica et Cosmochimica Acta*. 2016;178:160 - 77.
16. Sengupta S, McArthur J, Sarkar A, Leng MJ, Ravenscroft P, Howarth RJ, et al. Do Ponds Cause Arsenic-Pollution of Groundwater in the Bengal Basin? An Answer from West Bengal. *Environmental Science & Technology*. 2008;42:5156 - 64.
17. McArthur JM, Ravenscroft P, Sracek O. Aquifer arsenic source. *Nature Geoscience*. 2011;4(10):655 - 6.
18. Mailloux BJ, Trembath-Reichert E, Cheung J, Watson M, Stute M, Freyer GA, et al. Advection of surface-derived organic carbon fuels microbial reduction in Bangladesh groundwater. *PNAS*. 2013;110(13):5331 - 5.
19. van Dongen B, Rowland HAL, Gault AG, Polya DA, Bryant C, Pancost RD. Hopane, sterane and n-alkane distributions in shallow sediments hosting high arsenic groundwaters in Cambodia. *Applied Geochemistry*. 2008;23:3047 - 58.
20. Rowland HAL, Pederick RL, Polya DA, Pancost RD, Van Dongen BE, Gault AG, et al. The control of organic matter on microbially mediated iron reduction and arsenic release in shallow alluvial aquifers, Cambodia. *Geobiology*. 2007;5:281-92.
21. Neumann RB, Asfaq KN, Badruzzaman BM, Ali MA, Shoemaker JK, Harvey CF. Aquifer arsenic source reply. *Nature Geoscience*. 2011.
22. Datta S, Neal AW, Mohajerin TJ, Ocheltree T, Rosenheim BE, White CD, et al. Perennial ponds are not an important source of water or dissolved organic matter to groundwaters with high arsenic concentrations in West Bengal, India. *Geophysical Research Letters*. 2011;38(L20404).
23. Schaefer MV, Ying SC, Benner SG, Duan Y, Wang Y, Fendorf S. Aquifer Arsenic Cycling Induced by Seasonal Hydrologic Changes within the Yangtze River Basin. *Environmental Science and Technology*. 2016;DOI: 10.1021/acs.est.5b04986.
24. Stuckey JW, Schaefer MV, Kocar BD, Benner SG, Fendorf S. Arsenic release metabolically limited to permanently water-saturated soil in Mekong Delta. *Nature Geoscience*. 2016;9:70 - 6.
25. McArthur JM, Banerjee DM, Sengupta S, Ravenscroft P, Klump S, Sarkar A, et al. Migration of As, and  $^3\text{H}/^3\text{He}$  ages, in groundwater from West Bengal: Implications for monitoring. *Water Research*. 2010;44(14):4171-85.
26. Fendorf S, Michael HA, van Geen A. Spatial and temporal variations of groundwater arsenic in south and southeast Asia. *Science*. 2010;328:1123-7.
27. Polya DA, Charlet L. Rising arsenic risk? *Nature Geoscience*. 2009;2(6):383-4.
28. Polya DA, Gault AG, Diebe N, Feldman P, Rosenboom JW, Gilligan E, et al. Arsenic hazard in shallow Cambodian groundwaters. *Mineralogical Magazine*. 2005;69(5):807-23.
29. Polya DA, Charlet L. Arsenic in Shallow, Reducing Groundwaters in Southern Asia: An Environmental Health Disaster. *Elements*. 2006;2(2).
30. Benner SG, Polizzotto ML, Kocar BD, Ganguly S, Phan K, Ouch K, et al. Groundwater flow in an arsenic-contaminated aquifer, Mekong Delta, Cambodia. *Applied Geochemistry*. 2008;23(11):3072-87.
31. Richards LA, Magnone D, van Dongen BE, Ballentine CJ, Polya DA. Use of Lithium Tracers to Quantify Drilling Fluid Contamination for Groundwater Monitoring in Southeast Asia. *Applied Geochemistry*. 2015;63:190 - 202.
32. Sültenfuß J, Roether W, Rhein M. The Bremen mass spectrometric facility for the measurement of helium isotopes, neon, and tritium in water. *Isotopes in Environmental and Health Studies*. 2009;45(2):83 - 95.
33. Sültenfuß J, Purtscher R, Führböter JF. Age structure and recharge conditions of a coastal aquifer (northern Germany) investigated with  $^{39}\text{Ar}$ ,  $^{14}\text{C}$ ,  $^3\text{H}$ , He isotopes and Ne. *Hydrogeology Journal*. 2011;19:221 - 36.
34. Schell WR, Sauzay G, Payne BR. World Distribution of Environmental Tritium. Physical Behaviour of Radioactive Contaminants in the Atmosphere. Vienna: International Atomic Energy Agency; 1974. p. 388.
35. IAEA/WMO. Global Network of Isotopes in Precipitation. The GNIP Database. (<http://www.iaea.org/water>, accessed 15 Feb 2016). 2015.