

OPEN

**The Pediatric Infectious Disease Journal Publish Ahead of Print**

**DOI: 10.1097/INF.0000000000001242**

**The Epidemiology of Hand, Foot and Mouth Disease in Asia: A Systematic Review and Analysis**

Wee Ming Koh<sup>1</sup>, MSc, Tiffany Bogich<sup>2</sup>, PhD, Karen Siegel<sup>1</sup>, MPH, Jing Jin<sup>1</sup>, MSc, Elizabeth Y. Chong<sup>3</sup>, PhD, Chong Yew Tan<sup>4</sup>, BSc, Mark IC Chen<sup>1,5</sup>, MBBS, PhD, Peter Horby<sup>6</sup>, MBBS, MSc, PhD, and Alex R. Cook<sup>1,5,7</sup>, PhD

**Affiliations:**

1 Saw Swee Hock School of Public Health, National University of Singapore and National University Health System, Singapore

2 Standard Analytics, New York, US

3 Rollins School of Public Health, Emory University, US

4 Duke-NUS Graduate Medical School Singapore

5 Communicable Disease Centre, Tan Tock Seng Hospital, Singapore

6 Nuffield Department of Medicine, University of Oxford, UK

7 Yale-NUS College, National University of Singapore

**Author for correspondence:** Dr Alex R Cook

Saw Swee Hock School of Public Health

Tahir Foundation Building

12 Science Drive 2

National University of Singapore

Singapore 117549

alex.richard.cook@gmail.com

**Abbreviated title:** Epidemiology of HFMD in Asia: A Systematic Review

**Running head:** Epidemiology of HFMD in Asia

**Funding source:** The work was funded by Singapore's Ministry of Health Health Services Research grant number HSRG12MAY023, Communicable Disease Public Health Research Grant number CDPHRG12NOV021, the Centre for Infectious Disease Epidemiology and Research, the Ministry of Education Tier 1 grant and the President's Graduate Fellowship to WMK. The funders had no role in the decision to publish.

**Financial Disclosure:** Tiffany Bogich is employed by commercial company, Standard Analytics. The remaining authors have no financial relationships relevant to this article to disclose.

**Conflict of Interest:** The authors have no conflicts of interest to disclose.

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially.

## **Abstract**

**Context:** Hand, foot and mouth disease (HFMD) is a widespread pediatric disease caused primarily by human enterovirus 71 (EV71) and coxsackievirus A16 (CA16).

**Objective:** This study reports a systematic review of the epidemiology of HFMD in Asia.

**Data Sources:** PubMed, Web of Science and Google Scholar were searched up to December 2014.

**Study Selection:** Two reviewers independently assessed studies for epidemiologic and serologic information about prevalence and incidence of HFMD against predetermined inclusion/exclusion criteria.

**Data extraction:** Two reviewers extracted answers for eight specific research questions on HFMD epidemiology. The results are checked by three others.

**Results:** HFMD is found to be seasonal in temperate Asia with a summer peak, and in subtropical Asia, with spring and fall peaks, but not in tropical Asia; evidence of a climatic role was identified for temperate Japan. Risk factors for HFMD include hygiene, age, gender and social contacts, but most studies were underpowered to adjust rigorously for confounding variables. Both community-level and school-level transmission have been implicated but their relative importance for HFMD is inconclusive. Epidemiologic indices are poorly understood: No supporting quantitative evidence was found for the incubation period of EV71; the symptomatic rate of EV71/CA16 infection was 10% to 71% in four studies; while the basic reproduction number was between 1.1 to 5.5 in three studies. The uncertainty in these estimates inhibits their use for further analysis.

**Limitations:** Diversity of study designs complicates attempts to identify features of HFMD epidemiology.

**Conclusions:** Knowledge on HFMD remains insufficient to guide interventions such as the incorporation of an EV71 vaccine in pediatric vaccination schedules. Research is urgently needed to fill these gaps.

**Key Words:** Hand, foot and mouth disease (HFMD), EV71, CA16, Epidemiology

ACCEPTED

## Introduction

Hand foot and mouth disease (HFMD) has become an endemic childhood disease in East and South East Asia. Its main etiologic agents are Human Enterovirus 71 (EV-A71) and Coxsackievirus 16 (CV-A16). Although usually mild—with symptoms limited to  $>38^{\circ}\text{C}$  fever, malaise, rashes on the volar regions of the hands and feet, herpangina and difficulty eating and drinking—more rarely, infection can lead to complications of the nervous or cardiopulmonary systems. Such cases can result in long term sequelae such as cognitive and motor disorders<sup>1,2</sup> or death, usually from pulmonary edema or brainstem encephalitis<sup>3</sup>. Although complications are rare, the number of children being infected in high incidence countries such as China ( $\sim 2.7\text{M}$  cases in 2014<sup>3</sup>) means the death toll can be substantial (384 deaths in China in 2014<sup>3</sup>). The EV-A71 virus seems to be responsible for more severe outcomes, while CV-A16 and other Coxsackieviruses, such as CV-A2, CV-A6 and CV-A10, usually present milder symptoms that resolve within a few weeks<sup>4-6</sup>.

There are nearly 25 years of literature from Asia that describes the epidemiology of HFMD, drawing on pediatric cohorts, national surveillance systems, outbreak investigations and clinical data, and from disparate countries that span stages of economic development and with climates that range from tropical to temperate. This diversity complicates attempts to identify general features of HFMD epidemiology and conceals gaps in the body of knowledge of this important pediatric disease.

The objective of this paper is to provide a robust systematic review of the epidemiology of HFMD that informs public health policy making about HFMD epidemics. The review covers three major areas: (i) history and seasonality of HFMD, and the efforts in predictive modeling; (ii) risk factors for infection, to guide control; and (iii) global epidemiologic parameters, such as the incubation period and basic reproduction number, which may determine the effectiveness of control policies.

## Methods

### Search strategy and selection criteria

Using a combination of search terms, including “Hand foot and mouth disease”, “Hand foot and mouth”, “HFMD”, “Enterovirus”, “Enterovirus 71”, “EV71”, “Coxsackie A16”, “CA16”, “CVA16”, we searched PubMed, Thomson Reuters Web of Science and Google Scholar to identify 1305, 1255 and 100 articles respectively.

Eligibility criteria were articles that: (i) were published in peer reviewed journals from January 1957 to Dec 2014; (ii) were studies with epidemiologic and/or serologic information (quantitative/qualitative) about incidence, prevalence of HFMD; and/or (iii) contained information about factors associated with prevalence and incidence and/or (iv) employed statistical models to derive the above. Articles not in English, not related to HFMD, or HFMD articles that did not cover epidemiologic or clinical factors were excluded.

Two independent readers examined each of the 407 abstracts to determine if specific research questions were answered. The 8 specific research questions are as follows: i) What time of the year do HFMD outbreaks occur, and with what seasonal factors are outbreaks associated? ii) How long have EV-A71 and CV-A16 been circulating in Asia? iii) What age groups are at higher risk of infection? iv) What risk factors are associated with infection and severe outcomes? v) Where do infections predominantly occur (home or school)? vi) What is the incubation period? vii) What proportion of infections are symptomatic? viii) What is the basic reproduction number for HFMD by virus? An article was retained as long as both readers indicated that it answered at least one specific research question and was discarded if both readers agreed that no questions were answered. A third independent reader arbitrated when there was a disparity between the original readers.

The two original readers each read the full text of half of the articles to identify answers to the questions. A second pair of independent readers read the articles again. Finally, the first

author compiled all answers to the specific questions and compared the extracted answers to the original text. Relevant references from these papers were included in the analysis, in particular to identify non-English and early references. In total, information from 242 papers was compiled and 108 papers were used in data synthesis.

Hourly weather data were downloaded from the Weather Underground and aggregated at a weekly scale. Incidence data from Tokyo, Hong Kong, Taiwan and Singapore were extracted from routine surveillance data published by government agencies (the National Institute of Infectious Diseases, Japan<sup>7</sup>, the Department of Health, Hong Kong<sup>8</sup>, the Taiwan National Infectious Disease Statistics System<sup>9</sup>, and the Ministry of Health, Singapore).

Non-tabular data were extracted from figures using Plot Digitizer<sup>10</sup>. Data on weather and incidence were analyzed using a time series model. Symptomatic proportions were pooled by aggregating denominators and numerators. Other analyses used standard statistical methods and were conducted using R<sup>11</sup>.

## **Results**

### **Timing and seasonality of HFMD outbreaks**

Outbreaks of HFMD do not occur uniformly throughout the year across Asia. In Fukuoka, Japan, for example, weekly numbers of HFMD cases have been found to increase with average temperature and humidity, especially among younger children<sup>18</sup>. By digitizing incidence data from publications on Japan<sup>5,18,22,23</sup> and North China<sup>24-29</sup> (**Figure 1**), we observe that May through July are the months with highest incidence in temperate regions of Asia. However, this relationship is less clear for tropical and subtropical Asia. The extracted data on Southwest China<sup>24,30</sup>, South China<sup>2,24,31,32</sup>, Hong Kong<sup>33,34</sup> and Taiwan<sup>35-37</sup> shows that outbreaks typically happen in late spring and fall. No distinct pattern is obvious for tropical regions as seen from data in Thailand<sup>38-40</sup>, Vietnam<sup>41,42</sup>, Malaysia<sup>43</sup> and Singapore<sup>44-47</sup>, where outbreaks occur sporadically throughout the year, although models have been developed for

Singapore (~1° North) that show a positive statistical relationship between maximum daily temperature above 32°C with HFMD incidence in the subsequent 1-2 weeks<sup>46</sup>.

To assess how general the relationship between climate and transmissibility of HFMD was, we took incidence data from Tokyo, Hong Kong, Taiwan, and Singapore (**Figure 2**), i.e. spanning temperate, sub-tropical and tropical latitudes, and fitted time series models to them. After controlling for contagion via autoregression terms, the effect of meteorologic factors was weak: a small positive increase in transmissibility with rising absolute humidity/temperature during the current week in Tokyo and Singapore. There was no evidence for temperature and humidity in having the same effect in Hong Kong or Taiwan, although rising relative humidity seems to decrease transmissibility in Singapore.

The earliest recorded cases of HFMD in Asia are from Japan (1967)<sup>39</sup>, Singapore (1970)<sup>40</sup>, Taiwan (1980)<sup>41</sup> and Shanghai, China (1981)<sup>42</sup>. Since then, outbreaks have been reported in many parts of Asia, including mainland China<sup>18,19,22,23,42-59</sup>, Korea<sup>60-62</sup>, Japan<sup>13,63-76</sup>, Taiwan<sup>6,75,77-80</sup>, Hong Kong<sup>26,27,81</sup>, India<sup>82-87</sup>, Thailand<sup>30,32,88</sup>, Vietnam<sup>33</sup>, Malaysia<sup>35,75,89-93</sup>, Singapore<sup>4,94</sup> and Brunei<sup>95</sup>, as summarized in (**Top**: incidence, temperature, absolute and relative humidity. Top panels indicate incidence (data in gray, mean and 95% interval in black) for the time period Jan 2001-Mar 2012 (Tokyo), Jan 2001-Dec 2009 (Hong Kong, Peoples' Republic of China), Jan 2001-Dec 2011 (Taiwan, Republic of China), Jan 2001-Jan 2012 (Singapore). Middle and bottom panels show mean daily temperature, absolute and relative humidity at Tokyo Narita, Hong Kong International, Taipei Taoyuan International and Singapore Changi airports, downloaded from the Weather Underground. **Bottom**: Coefficients of meteorological variables at 0 to 2 week time lags in autoregressive models of z-scored HFMD case counts to facilitate comparison between locations. Each city is analyzed separately using a model in which HFMD incidence in week  $t$  is (auto)regressed on incidence in weeks  $t-1$  up to  $t-3$  (using the Akaike information criterion to select the order of the

autoregression component) and, independently, on each meteorological variable. Weather parameters are not regressed together in a single model due to collinearity. The effect of weather on HFMD incidence can be seen by coefficient mean (points) and 95% confidence intervals (lines), colored red if statistically significant at the 5% level.

**Figure 3).** These reported outbreaks are unlikely to reflect the true first outbreaks of HFMD, as serologic studies provide evidence that by the time surveillance systems were established, EV-A71 and CV-A16 were already endemic in many of these countries. Early serologic tests conducted in Japan in 1970 show evidence of EV-A71 and CV-A16 circulation<sup>96</sup>. Serum taken in the late 1990s in Singapore, before the start of surveillance in 2000, shows that around 50% children and 44% cord blood, indicating maternal infection, had already seroconverted to EV-A71<sup>97</sup>. Blood samples from Taiwan from 1989 to 1997 show 3% to 11% EV-A71 incidence per year, and up to 68% of children<sup>98</sup> had serologic evidence of EV-A71 infection before the large HFMD outbreak of 1997. Similarly, although China has reported millions of HFMD cases since the beginning of the HFMD surveillance program in 2008, evidence from Anhui<sup>54</sup> shows high seroprevalence of up to 74.6% in older children before the 2008 outbreaks. Retrospective seroepidemiologic tests from blood serum collected in 2005<sup>99</sup> also show that China had positive rates of 32.0% to EV-A71 and 43.4% to CV-A16, indicating that outbreaks happened earlier but were simply not reported in the literature.

### **Risk Factors**

Risk factors for infection are depicted in **(Figure 4)** and summarized below.

**Hygiene:** Evidence from Qiaosi, China<sup>100</sup> indicates the importance of hygiene for protection against HFMD infection. Children who always wash their hands before meals are about 50 times less likely to contract HFMD, while those whose caregivers wash their hands before feeding are about 25 times less likely. Additional protective habits include washing of hands after play, washing of hands more than 4 times per day, using soap, and not sucking fingers.

A study in Korea<sup>101</sup> revealed that drinking unboiled water (3.34 OR [1.59-6.99]), a change in water quality such as color, taste, smell, presence of precipitation or floating materials (6.93 OR [2.17-22.15]), using communal toilets/toilets outside the house (2.77 OR [1.14-6.74]), and eating outside the home (37.0 OR [5.1-269.5]) were risk factors for HFMD.

**Rural vs urban areas:** All papers<sup>58,102-104</sup> that compared urban with rural areas agreed that the latter conferred a higher risk for HFMD. However, this might be confounded by socioeconomic status and hygiene practices.

**Sex:** While most papers show that being male is a risk factor for both mild<sup>4,16,17,23,25,32,36,43,58,46,88,104,105</sup> and severe<sup>20,59,103,106</sup> HFMD (OR ranges between 1.2 to 2), surprisingly, serologic evidence does not support this finding: a study from Singapore<sup>107</sup> shows marginal evidence that females are more likely to have seroconverted to EV-A71 (OR 0.79 [0.61-1.01]) while a Taiwanese<sup>102</sup> study shows no statistically significant differences (OR 0.94 [0.76-1.16]). Taken together, these suggest that infection rates are comparable, but that boys are more likely to develop symptoms, more involved in propagation of outbreaks or more likely to be brought for medical care than girls.

**Other:** A case-control study in Xi'an, China<sup>103</sup>, found that breastfeeding may lower the risk of developing severe HFMD (aOR 0.57 [0.33-0.98]), even though breastfeeding does not apparently lower the chance of being infected by EV-A71 (OR 1.1 [0.93-1.3])<sup>102</sup>. It further found that patients with a history of Epstein - Barr virus are at greater risk of contracting severe, rather than mild, HFMD (aOR 2.6 [1.5-4.4]). A spatial-temporal model of Guangdong<sup>23</sup> showed that sunshine could be protective against HFMD. This is agreed by a matched case-control study of preschoolers in Beijing<sup>48</sup> which showed that UV radiation in classrooms is associated with lower HFMD attack rate (p-value 0.027), and recommended installing UV lamps to sterilize unoccupied classrooms. These findings are however,

inconsistent with the seasonal nature of HFMD, where outbreaks in temperate countries tend to occur in summer, when sunlight and UV exposure is strongest.

### **Age distribution of HFMD cases**

The age distribution of HFMD cases in Asia, compiled from a variety of sources including surveillance and cohort data, is summarized in **Figure 5**. Data from China<sup>16,18,20,22,23,43,56–59,100,105,106,108–110</sup> and Taiwan<sup>5,6,79,111–114</sup> are particularly abundant. Other sources include Hong Kong<sup>26,27</sup>, India<sup>82,86</sup>, Japan<sup>13,15</sup>, Korea<sup>61,101</sup>, Malaysia<sup>90,115</sup>, Singapore<sup>4,36,94</sup>, Thailand<sup>31,32</sup> and Vietnam<sup>33</sup>.

The symptomatic HFMD incidence rate varies widely even within the narrow 0 to 6 year age-band. The greatest proportion of cases occur at ages 1 (18.8% [17.4–20.2%]) and 2 (17.9% [16.6–19.2%]). By the age of formal schooling, from 6 in most Asian countries, the proportion is substantially lower (8.7% [7.9–9.5%]). Overall, 82.6% (82.2–82.9%) of all cases occur before age 6. The lower rate during the first year of life could be due to lack of contact with other children or to presence of maternal antibodies<sup>97</sup>.

### **Community versus school as medium for infection**

The literature is ambiguous about the importance of locations for transmission. Four studies showed that contact with a case, particularly a household member, is as or more significant a risk factor than preschool attendance<sup>30,94,102,111</sup>. An early study in Singapore observed 60 families with secondary cases and found the secondary attack rate amongst children below 12 years old to be 77%<sup>94</sup>. Similarly, in a large seroepidemiologic study of EV-A71 in Taiwanese children<sup>102</sup>, multivariate analysis showed attendance at a preschool imparted a similar magnitude of risk as contact with a case (adjusted ORs of 1.6 [1.2–2.1] and 1.8 [1.3–2.5], respectively), as well as a strong concordance (84%) between seropositivity in younger and older siblings.

Also, a number of studies showed that a higher percentage of diagnoses occurred among children who did not attend a nursery or preschool<sup>46,58</sup>. Liu et al<sup>56</sup> note that about half of symptomatic cases in Nanchang, China, are among children under 3, the age at which preschooling starts in China.

Conversely, some studies suggest that preschool attendance is a key risk factor<sup>4,116</sup>. For example, a seroepidemiologic study in 1996–7 in Singapore showed that seropositivity to EV-A71 increases rapidly from age 2 to age 5<sup>97</sup>, when attendance at childcare or preschool is the norm. Also, a case-control study in Japan<sup>116</sup> showed that preschool attendance was associated with increased risk of severe disease.

Other studies suggest that both locations are important. In Shanghai, China,<sup>106</sup> there was a marked shift from 2007 to 2008 in the proportion of cases among children in preschools (from 59% to 37%) with a concurrent shift from local to migrant children, suggesting that the importance of routes of transmission can vary over time within the same locale. A case-control study from Zhejiang<sup>100</sup> showed that while attending preschool is a risk factor (2.1 OR), other factors such as contact with neighbours (11 OR), going to hospital (20 OR), and going to parties (31 OR) impart greater risk. Yet, a Korean case-control study<sup>101</sup>, found no significant relationship between infection and school attendance or household size.

Overall, the evidence points to both home and school environments contributing to transmission, but the relative importance of these venues remains murky.

### **Incubation period**

Several papers describe the incubation period (**Figure 5**), though it is striking that the majority do not provide a source to justify the claimed period. These unsupported claims vary substantially from paper to paper, from the incubation period ‘is’ 3–6 days<sup>117</sup>, or 3–7 days<sup>82</sup>, ‘is usually 3–4 days, but can be ... 10 days or more’<sup>41</sup>, or ‘is usually 3 to 5 days (range, 2 to 12 days)’<sup>114</sup>, is ‘typically’ 3–7 days<sup>118</sup> or 3–5 days<sup>56</sup>, ranges from 5 to 7 days<sup>49,104</sup> or 3 to 7

days<sup>115</sup>, and the ‘usual period’ is 3 to 5 days ‘with longest period of 7 days’<sup>119</sup>. Only a few provide evidence to justify the claim: one reports<sup>101</sup> that the incubation period is usually 3 to 7 days, citing a US CDC factsheet on aseptic meningitis. Another cites<sup>12</sup> an early study from Singapore<sup>94</sup> that presented the median and range for the serial interval (3 days [1–7]), not the incubation period. Another early study<sup>120</sup> states that the incubation period is “said to be” 3 to 5 days, but notes that this is inconsistent with the serial interval observed in the study. It appears that there is no empirical support whatsoever for any distribution of incubation periods.

### **Symptomatic proportion**

Although several studies report that the asymptomatic rate of EV-A71 infection is high, few studies report data (**Table 1**). Two studies, from Taiwan and Shanghai, tested sera for evidence of EV-A71 infection and asked patients or their families to recall past HFMD infection, deriving estimates of 29%<sup>121</sup> and 10%<sup>109</sup> of symptomatic infection respectively. Some HFMD cases may have been caused by other enteroviruses, biasing these estimates upwards, while some may have been diagnosed as another viral illness or forgotten, biasing them downwards.

Two additional studies in Taiwan found much higher symptomatic infection rates. The first<sup>111</sup> recruited symptomatic cases suspected of having EV-A71 infection, and took throat and rectal swabs or stool samples, of cases and their household members with repeat telephone interviews. Excluding the 94 symptomatic index cases, 68% of confirmed infections in the household were symptomatic (88% of infected children and 47% of infected adults). A second study<sup>122</sup> prospectively followed a cohort of neonates over 3 years, taking repeat sera, requesting that parents report suspected HFMD and giving reminders during HFMD epidemics. This study found that 71% of serologically confirmed infections were symptomatic, though the sample size is only 28.

The discrepancy between these two pairs of papers is substantial, undoubtedly due to differences in methodology. An overall estimate, combining the four studies, is 36% (33%–39%) but given the large discrepancy between studies this estimate does not appear reliable. The latter pair of studies are prospective, thereby circumventing recall bias, and thus appear to provide a more accurate description of the epidemiology of enterovirus infection.

### **Basic reproduction number for HFMD by virus**

Only three papers have sought to estimate the reproduction number for HFMD or the viruses that cause it. One<sup>16</sup> estimates what they call the ‘local effective reproduction number’ in China—meaning using the average number of secondary cases from a randomly selected index to estimate the cases that would be caused in a fully susceptible population (note, this is substantially different from the effective reproduction number<sup>123</sup> in a partially susceptible population)—using a sophisticated Poisson regression model that incorporated infection from the environment, the prefecture and neighboring prefectures. This model did not, however, account for the accumulation of herd immunity, and required arbitrary assignment of the infectious period, so the estimated local effective reproduction number of 1.1 to 1.2 during peak periods may be biased.

A second paper<sup>119</sup> used a method from Choi and Pak<sup>124</sup> to estimate the basic reproduction number to be 5.5 (IQR 4.2 to 6.5) for EV-A71 and 2.5 (IQR 2.0 to 3.7) for CV-A16. These estimates are likely inaccurate because the method assumes (i) a known generation time distribution, labelled incubation period in the paper; (ii) a completely immunonaïve population, though applied to groups of individuals for whom past exposure was highly plausible; and (iii) an early exponential growth period, despite being applied to complete outbreak data.

The third paper<sup>125</sup> attempted to estimate the reproduction number using a SEIQRS (Susceptible, exposed, infectious, quarantined, recovered) simulation model, and obtained an

estimate of 1.1 for the years 2009 to 2012 in China. However, the model used 10% of China's population as the initial susceptible population but did not conduct a sensitivity analysis on this vital parameter.

## **Discussion**

Despite the substantial number of papers on HFMD, this systematic review shows that many fundamental questions about EV-A71 and CV-A16 persist. Both viruses occur year round in tropical Asia, but are epidemic in the summer in North-East Asia. A role for temperature or humidity therefore seems plausible<sup>14,23,126–128</sup>, although given the relative lack of seasonality in equatorial Asia, it is not clear whether prediction of outbreaks is possible there. In Japan, summer temperatures peak after HFMD incidence does, suggesting correlation but not temporality, and that it may not be possible to provide early warning of impending epidemics. This also differs from other human enteric viruses including Poliovirus 1 (also an Enterovirus), Hepatitis A, and Adenovirus that have been shown to survive longer on colder surfaces<sup>129</sup>.

Urashima et al (2003)<sup>14</sup> claimed that enteroviruses experience a more rapid virus decline during dry seasons than during wet seasons, which could explain the seasonality. This result is supported by Wang et al (2011)<sup>130</sup>, where they showed that precipitation patterns has the most similar structure as HFMD incidence, more so than other meteorologic variables, albeit with only 11 months of data.

While any causal relationship between climate and HFMD is unknown, speculations include a lower HFMD incidence due to decreased social contact during temperate zones' winter<sup>12,131</sup>. In contrast, increased social contacts during winter have been speculated to facilitate spread of other droplet borne diseases, such as influenza<sup>132</sup>, which are epidemic in winter. Given the unknowns surrounding this issue, further research is clearly required to

ascertain whether meteorologic factors or seasonal social contact patterns is an adequate explanation for the seasonality of HFMD.

The next step to analyzing the dynamics of HFMD seasonality is likely to involve social and environmental factors – another under-researched area for this pediatric disease. For instance, the literature is unclear on the relative importance of school versus community transmission, with evidence to support both, yet knowledge of where HFMD most often is transmitted is important as school closure policies are employed to control outbreaks in some countries. Further, the environment of schools in Asia may vary widely, and attributes such as hygiene practices should be characterized and quantified to allow more definitive results and conclusions in future studies.

Even without being able to determine the relative importance of school versus community transmission, the effectiveness of school closure to prevent large scale HFMD outbreaks is questionable, as the interruption to social networks cannot be enforced while children are out of school. Additionally, while we know little about the infectiousness of asymptomatic cases of HFMD, the proportion of infections that are asymptomatic is substantial, and so even quite modest school closure attack rate thresholds, such as Singapore's 25%<sup>133</sup>, corresponds to a possible majority of students being infected prior to the trigger for closure being met. Further, EV-A71 can be found in fecal samples for up to 54 days after infection<sup>134</sup>, and thus continue to be shed after a school is closed, disinfected, and reopened.

Studies on risk factors were rare, and we identified only 3 papers that describe risk factors for hygiene and contact patterns, making a meta-analysis of risk factors unfeasible. These typically were only powered to provide unadjusted effect sizes, and so provide evidence of correlation, not causation. One interesting finding was the apparent protective effect of a caregiver "always washing" their hands. This suggests that adult to child transmission might be important, even if adults are mostly asymptomatic with EV-A71 and CV-A16, but may

reflect confounding with general hygiene. Future work may elicit hygiene factors at the preschool level and relate these to attack rates.

A recently developed EV-A71 vaccine has undergone phase 3 trials in China<sup>135–137</sup>. To determine the cost-effectiveness of incorporating the vaccine in pediatric vaccination schedules, or of other interventions such as school closure or isolation, would require epidemiologic models that account for the protective effects of herd immunity. However, this review indicates that vital parameters for such models remain unknown. The asymptomatic rate and relative infectiousness of asymptomatic cases are both poorly known, while estimates of the incubation period, although commonly cited as 3 to 5 days, appear to be based solely on expert opinion. Most importantly, estimates of the basic reproduction number range widely from 1.1 to 5.5. This uncertainty prohibits utilitarian estimation of the necessary vaccine coverage to prevent epidemics of EV-A71.

To reconcile the differences between the disparate estimates, the age distributions of the samples need to be considered. As shown in this review, symptomatic HFMD incidence rates differ greatly even between ages 0 to 6, and thus studies conducted predominantly on preschoolers may derive higher estimates of  $R_0$  compared to studies in older children.

Accordingly, future studies on HFMD should use narrower age bands and also state the distribution clearly to allow adjustments or standardization.

Two final omissions from the literature are quantitative estimates of the impact of infection on complications, child and caregiver absenteeism, and costings of complications, and qualitative evidence on the impact of infection and enforced isolation on families and schools. Given the promising direct effects of the EV-A71 vaccine, and the huge public health impacts of HFMD in East and South East Asia, research is urgently needed to fill these gaps.

The research questions in this systematic review were generally answered only by a limited number of papers, with substantial differences in their study design, and thus most data were not synthesized through meta-analysis. More research to assess risk factors and measure key epidemiologic parameters is needed. We were also unable to trace the earliest cases of HFMD in Asia as our scope only covers published material on outbreaks, which leads us back to 1967 in Japan. Finally, we limited the scope of our study to exclude virologic characteristics or molecular epidemiology, which have been well reviewed elsewhere<sup>118,139–141,142</sup>, and clinical manifestations of EV-A71 and CV-A16<sup>118,138,139,142,143</sup>. A recent review of the case-fatality rate has recently been published<sup>144</sup>, as has a review of the epidemiology in Taiwan<sup>145</sup>.

### **Acknowledgments**

We thank Alex S Leow, Vincent J Pang for screening some of the papers and Zhao Xiahong for processing the meteorologic data.

## References

1. Huang M-C, Wang S-M, Hsu Y-W, Lin H-C, Chi C-Y, Liu C-C. Long-term cognitive and motor deficits after enterovirus 71 brainstem encephalitis in children. *Pediatrics*. 2006;118(6):e1785-e1788. doi:10.1542/peds.2006-1547.
2. Chang L-Y, Huang L-M, Gau SS-F, et al. Neurodevelopment and cognition in children after enterovirus 71 infection. *N Engl J Med*. 2007;356(12):1226-1234. doi:10.1056/NEJMoa065954.
3. World Health Organization Western Pacific Region. Hand, Foot, and Mouth Disease Situation Update. December 2014. [http://www.wpro.who.int/emerging\\_diseases/hfmd\\_biweekly\\_29dec2014.pdf?ua=1](http://www.wpro.who.int/emerging_diseases/hfmd_biweekly_29dec2014.pdf?ua=1). Accessed March 11, 2015.
4. Ang LW, Koh BK, Chan KP, Chua LT, James L, Goh KT. Epidemiology and control of hand, foot and mouth disease in Singapore, 2001-2007. *Ann Acad Med Singap*. 2009;38(2):106-112.
5. Chen S-P, Huang Y-C, Li W-C, et al. Comparison of clinical features between coxsackievirus A2 and enterovirus 71 during the enterovirus outbreak in Taiwan, 2008: a children's hospital experience. *J Microbiol Immunol Infect*. 2010;43(2):99-104.
6. Lo S-H, Huang Y-C, Huang C-G, et al. Clinical and epidemiologic features of Coxsackievirus A6 infection in children in northern Taiwan between 2004 and 2009. *J Microbiol Immunol Infect*. 2011;44(4):252-257. doi:10.1016/j.jmii.2011.01.031.
7. National Institute of Infectious Diseases, Japan. Infectious Disease Weekly Report, Surveillance Data Tables. <http://www0.niid.go.jp/niid/index-e.html>.
8. Centre for Health Protection, Department of Health, Hong Kong. Updated Situation of Hand, Foot and Mouth Disease (HFMD) and Enterovirus Infection 2010. August 2010.

- [http://www.chp.gov.hk/files/pdf/updated\\_situation\\_of\\_hand\\_foot\\_and\\_mouth\\_disease\\_hfmd\\_r.pdf](http://www.chp.gov.hk/files/pdf/updated_situation_of_hand_foot_and_mouth_disease_hfmd_r.pdf). Accessed November 5, 2015.
9. Taipei (Taiwan): Taiwan Centers for Disease Control. Taiwan National Infectious Disease Statistics System. Taiwan National Infectious Disease Statistics System. <http://nidss.cdc.gov.tw/en/>. Published April 15, 2014.
  10. Huwaldt J. *Plot Digitizer*. <http://sourceforge.net/projects/plotdigitizer/>.
  11. R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.; 2013.
  12. Onozuka D, Hashizume M. The influence of temperature and humidity on the incidence of hand, foot, and mouth disease in Japan. *Sci Total Environ*. 2011;410-411:119-125. doi:10.1016/j.scitotenv.2011.09.055.
  13. Miwa C, M. Ohtani, H. Watanabe, J. Fujimoto, K. Kinbara, H. Tanaka. Epidemic of hand, foot and mouth disease in Gifu Prefecture in 1978. *Jpn J Med Sci Biol*. 1980;33(3):167-180.
  14. Urashima M, Shindo N, Okabe N. Seasonal models of herpangina and hand-foot-mouth disease to simulate annual fluctuations in urban warming in Tokyo. *Jpn J Infect Dis*. 2003;56(2):48-53.
  15. Momoki ST. Surveillance of enterovirus infections in Yokohama city from 2004 to 2008. *Jpn J Infect Dis*. 2009;62(6):471-473.
  16. Wang Y, Feng Z, Yang Y, et al. Hand, Foot, and Mouth Disease in China: Patterns of Spread and Transmissibility. *Epidemiology*. 2011;22(6):781-792. doi:10.1097/EDE.0b013e318231d67a.
  17. Yan L, Li X, Yu Y, et al. Distribution and risk factors of hand, foot, and mouth disease in Changchun, northeastern China. *Chin Sci Bull*. 2014;59(5-6):533-538. doi:10.1007/s11434-013-0069-5.

18. Liu Y, Wang X, Liu Y, et al. Detecting Spatial-Temporal Clusters of HFMD from 2007 to 2011 in Shandong Province, China. Khudyakov YE, ed. *PLoS One*. 2013;8(5):e63447. doi:10.1371/journal.pone.0063447.
19. Cao Z, Zeng D, Wang Q, Zheng X, Wang F. An epidemiological analysis of the Beijing 2008 Hand-Foot-Mouth epidemic. *Chin Sci Bull*. 2010;55(12):1142-1149. doi:10.1007/s11434-010-0144-0.
20. Zhu D, Zhao X-Y, Yao Y, et al. A new factor influencing pathogen detection by molecular assay in children with both mild and severe hand, foot, and mouth disease. *Diagn Microbiol Infect Dis*. 2013;76(2):162-167. doi:10.1016/j.diagmicrobio.2013.02.011.
21. Tian H, Zhang Y, Sun Q, et al. Prevalence of Multiple Enteroviruses Associated with Hand, Foot, and Mouth Disease in Shijiazhuang City, Hebei Province, China: Outbreaks of Coxsackieviruses A10 and B3. Riley S, ed. *PLoS One*. 2014;9(1):e84233. doi:10.1371/journal.pone.0084233.
22. Xu W, Jiang L, Thammawijaya P, Thamthitiwat S. Hand, Foot and Mouth Disease in Yunnan Province, China, 2008-2010. *Asia Pac J Public Health*. December 2011. doi:10.1177/1010539511430523.
23. Deng T, Huang Y, Yu S, et al. Spatial-Temporal Clusters and Risk Factors of Hand, Foot, and Mouth Disease at the District Level in Guangdong Province, China. Viboud C, ed. *PLoS One*. 2013;8(2):e56943. doi:10.1371/journal.pone.0056943.
24. Di B, Zhang Y, Xie H, et al. Circulation of Coxsackievirus A6 in hand-foot-mouth disease in Guangzhou, 2010-2012. *Viol J*. 2014;11:157.
25. Wang Y, Sun L, Xiao W, Chen L, Wang X, Pan D. Epidemiology and clinical characteristics of hand foot, and mouth disease in a Shenzhen sentinel hospital from 2009 to 2011. *BMC Infect Dis*. 2013;13(1):539.

26. Ma E, Chan KC, Cheng P, Wong C, Chuang SK. The enterovirus 71 epidemic in 2008—public health implications for Hong Kong. *Int J Infect Dis.* 2010;14(9):e775-e780. doi:10.1016/j.ijid.2010.02.2265.
27. Ma E, Lam T, Chan KC, Wong C, Chuang SK. Changing epidemiology of hand, foot, and mouth disease in Hong Kong, 2001–2009. *Jpn J Infect Dis.* 2010;63(6):422-426.
28. Lee T-C, Guo H-R, Su H-JJ, Yang Y-C, Chang H-L, Chen K-T. Diseases caused by enterovirus 71 infection. *Pediatr Infect Dis J.* 2009;28(10):904-910. doi:http://dx.doi.org/10.1097/inf.0b013e3181a41d63.
29. Yang T-T, Huang L-M, Lu C-Y, et al. Clinical features and factors of unfavorable outcomes for non-polio enterovirus infection of the central nervous system in northern Taiwan, 1994-2003. *J Microbiol Immunol Infect.* 2005;38(6):417-424.
30. Chatproedprai S, Theanboonlers A, Korkong S, Thongmee C, Wananukul S, Poovorawan Y. Clinical and molecular characterization of hand-foot-and-mouth disease in Thailand, 2008-2009. *Jpn J Infect Dis.* 2010;63(4):229-233.
31. Linsuwanon P, Puenpa J, Huang S-W, et al. Epidemiology and seroepidemiology of human enterovirus 71 among Thai populations. *J Biomed Sci.* 2014;21:16.
32. Samphutthanon R, Tripathi N, Ninsawat S, Duboz R. Spatio-Temporal Distribution and Hotspots of Hand, Foot and Mouth Disease (HFMD) in Northern Thailand. *Int J Environ Res Public Health.* 2013;11(1):312-336. doi:10.3390/ijerph110100312.
33. Tu PV, Thao NTT, Perera D, et al. Epidemiologic and virologic investigation of hand, foot, and mouth disease, southern Vietnam, 2005. *Emerg Infect Dis.* 2007;13(11):1733-1741. doi:10.3201/eid1311.070632.
34. Nguyen NT, Pham HV, Hoang CQ, et al. Epidemiological and clinical characteristics of children who died from hand, foot and mouth disease in Vietnam, 2011. *BMC Infect Dis.* 2014;14(1):341.

35. Podin Y, Gias EL, Ong F, et al. Sentinel surveillance for human enterovirus 71 in Sarawak, Malaysia: lessons from the first 7 years. *BMC Public Health*. 2006;6(1):180.
36. Chan KP, Goh KT, Chong CY, Teo ES, Lau G, Ling AE. Epidemic Hand, Foot and Mouth Disease Caused by Human Enterovirus 71, Singapore. *Emerg Infect Dis*. 2003;9(1):78-85. doi:10.3201/eid1301.020112.
37. Hii YL, Rocklöv J, Ng N. Short term effects of weather on hand, foot and mouth disease. *PLoS One*. 2011;6(2):e16796.
38. Wu Y, Yeo A, Phoon MC, et al. The largest outbreak of hand; foot and mouth disease in Singapore in 2008: The role of enterovirus 71 and coxsackievirus A strains. *Int J Infect Dis*. 2010;14(12):e1076-e1081. doi:10.1016/j.ijid.2010.07.006.
39. Sawada K, Nakamura Y, Ito S. On the hand, foot and mouth disease prevailed in Ohmiya City in summer, 1967. *Nihon Shonika Gakkai Zasshi*. 1968;74(1594).
40. Chan MCK, Wong HB. Hand-foot and mouth disease. *J Singapore Paediatr Soc*. 1973;15:31-34.
41. Chang L-Y. Enterovirus 71 in Taiwan. *Pediatr Neonatol*. 2008;49(4):103-112.
42. Li C. The prevalence of Hand, Foot and Mouth Disease. *J Pediatr Pharm*. 2008;14:64-66.
43. Sun L, Zheng H, Zheng H, et al. An enterovirus 71 epidemic in Guangdong Province of China, 2008: epidemiological, clinical, and virogenic manifestations. *Jpn J Infect Dis*. 2011;64(1):13-18.
44. Zhang Y, Tan X-J, Wang H-Y, et al. An outbreak of hand, foot, and mouth disease associated with subgenotype C4 of human enterovirus 71 in Shandong, China. *J Clin Virol*. 2009;44(4):262-267. doi:10.1016/j.jcv.2009.02.002.

45. Ding N-Z, Wang X-M, Sun S-W, Song Q, Li S-N, He C-Q. Appearance of mosaic enterovirus 71 in the 2008 outbreak of China. *Virus Res.* 2009;145(1):157-161. doi:10.1016/j.virusres.2009.06.006.
46. Wang J, Cao Z, Zeng DD, Wang Q, Wang X, Qian H. Epidemiological Analysis, Detection, and Comparison of Space-Time Patterns of Beijing Hand-Foot-Mouth Disease (2008–2012). Rong L, ed. *PLoS One.* 2014;9(3):e92745. doi:10.1371/journal.pone.0092745.
47. Wang X, Wu X, Jia L, et al. Estimating the number of hand, foot and mouth disease amongst children aged under-five in Beijing during 2012, based on a telephone survey of healthcare seeking behavior. *BMC Infect Dis.* 2014;14(1):437.
48. Wu X, Sun Y, Lin C, et al. A case-control study to identify environmental risk factors for hand, foot, and mouth disease outbreaks in Beijing. *Jpn J Infect Dis.* 2014;67(2):95-99. doi:10.7883/yoken.67.95.
49. De W, Changwen K, Wei L, et al. A large outbreak of hand, foot, and mouth disease caused by EV71 and CAV16 in Guangdong, China, 2009. *Arch Virol.* 2011;156(6):945-953. doi:10.1007/s00705-011-0929-8.
50. Li W, Yi L, Su J, et al. Seroprevalence of Human Enterovirus 71 and Coxsackievirus A16 in Guangdong, China, in Pre- and Post-2010 HFMD Epidemic Period. Borrow R, ed. *PLoS One.* 2013;8(12):e80515. doi:10.1371/journal.pone.0080515.
51. Xu W, Liu C, Yan L, et al. Distribution of enteroviruses in hospitalized children with hand, foot and mouth disease and relationship between pathogens and nervous system complications. *Virol J.* 2012;9(1):8.
52. Zhang Y, Zhu Z, Yang W, et al. An emerging recombinant human enterovirus 71 responsible for the 2008 outbreak of hand foot and mouth disease in Fuyang city of China. *Virol J.* 2010;7(1):94.

53. Bingjun T, Yoshida H, Yan W, et al. Molecular typing and epidemiology of non-polio enteroviruses isolated from Yunnan Province, the People's Republic of China. *J Med Virol*. 2008;80(4):670-679. doi:10.1002/jmv.21122.
54. Yu H, Wang M, Chang H, et al. Prevalence of antibodies against enterovirus 71 in children from Lu'an City in Central China. *Jpn J Infect Dis*. 2011;64(6):528-532.
55. Cardoso MJ, Perera D, Brown BA, et al. Molecular epidemiology of human enterovirus 71 strains and recent outbreaks in the Asia-Pacific region: comparative analysis of the VP1 and VP4 genes. *Emerg Infect Dis*. 2003;9(4):461-468. doi:10.3201/eid0904.020395.
56. Liu MY, Liu W, Luo J, et al. Characterization of an Outbreak of Hand, Foot, and Mouth Disease in Nanchang, China in 2010. Zheng JC, ed. *PLoS One*. 2011;6(9):e25287. doi:10.1371/journal.pone.0025287.
57. Zhu Q, Hao Y, Ma J, Yu S, Wang Y. Surveillance of hand, foot, and mouth disease in mainland China (2008–2009). *Biomed Env Sci*. 2011;24(4):349-356. doi:10.3967/0895-3988.2011.04.005.
58. Mao L-X, Wu B, Bao W-X, et al. Epidemiology of hand, foot, and mouth disease and genotype characterization of Enterovirus 71 in Jiangsu, China. *J Clin Virol*. 2010;49(2):100-104. doi:10.1016/j.jcv.2010.07.009.
59. He S-J, Han J-F, Ding X-X, Wang Y-D, Qin C-F. Characterization of enterovirus 71 and coxsackievirus A16 isolated in hand, foot, and mouth disease patients in Guangdong, 2010. *Int J Infect Dis*. 2013;17(11):e1025-e1030. doi:10.1016/j.ijid.2013.04.003.
60. Jee YM, Cheon D-S, Kim K, et al. Genetic analysis of the VP1 region of Human enterovirus 71 strains isolated in Korea during 2000. *Arch Virol*. 2003;148(9):1735-1746. doi:10.1007/s00705-003-0133-6.

61. Baek K, Yeo S, Lee B, et al. Epidemics of enterovirus infection in Chungnam Korea, 2008 and 2009. *Viol J*. 2011;8(1):297. doi:10.1186/1743-422X-8-297.
62. Kim SJ, Kim J-H, Kang J-H, et al. Risk Factors for Neurologic Complications of Hand, Foot and Mouth Disease in the Republic of Korea, 2009. *J Korean Med Sci*. 2013;28(1):120. doi:10.3346/jkms.2013.28.1.120.
63. Gobara F, Itagaki A, Ito Y, Saito K, Katsumoto T, Kimura G. Properties of Virus Isolated from an Epidemic of Hand-Foot-and-Mouth Disease in 1973 in the City of Matsue. *Microbiol Immunol*. 1977;21(4):207-217.
64. Hagiwara A, Tagaya I, Yoneyama T. Epidemic of hand, foot and mouth disease associated with enterovirus 71 infection. *Intervirology*. 1978;9(1):60-63.
65. Tagaya I, Moritsugu Y. Epidemic of Hand, Foot and Mouth Disease in Japan. *Jpn J Med Sci Biol*. 1973;26(3):143-147. doi:10.7883/yoken1952.26.143.
66. Fujimiya Y, Oyama S, Numazaki Y, Ishida N. Serologic Relationship of Coxsackie A16 Viruses from the Epidemic of Hand-Foot-and-Mouth Disease in Japan, 1970, to the Prototype Strain. *Jpn J Microbiol*. 1974;18(5):379-384. doi:10.1111/j.1348-0421.1974.tb00824.x.
67. Itagaki A, Gobara F, Ito Y, Saito K. A Survey on Enterovirus Infections in Matsue City, Shimane Prefecture in the Period from May, 1969 to March, 1971. *J Jpn Assoc Infect Dis*. 1972;46(8):280-289. doi:10.11150/kansenshogakuzasshi1970.46.280.
68. Tagaya I, Tachibana K. Epidemic of Hand, Foot and Mouth Disease in Japan, 1972-1973: Difference in Epidemiologic and Virologic Features from the Previous One. *Jpn J Med Sci Biol*. 1975;28(4):231-234. doi:10.7883/yoken1952.28.231.
69. Gobara F, Itagaki A, Ito Y, Saito K, Nishino Y. A Survey on Enterovirus Infection in Matsue City, Shimane Prefecture in the Period from April to October, 1973. *J Jpn Assoc Infect Dis*. 1975;49(7):282-287. doi:10.11150/kansenshogakuzasshi1970.49.282.

70. Ishimaru Y, Nakano S, Yamaoka K, Takami S. Outbreaks of hand, foot, and mouth disease by enterovirus 71. High incidence of complication disorders of central nervous system. *Arch Dis Child*. 1980;55(8):583-588.
71. Tagaya I, Takayama R, Hagiwara A. A Large-Scale Epidemic of Hand, Foot and Mouth Disease Associated with Enterovirus 71 Infection in Japan in 1978. *Jpn J Med Sci Biol*. 1981;34(3):191-196. doi:10.7883/yoken1952.34.191.
72. Hosoya M, Kawasaki Y, Sato M, et al. Genetic Diversity of Coxsackievirus A16 Associated with Hand, Foot, and Mouth Disease Epidemics in Japan from 1983 to 2003. *J Clin Microbiol*. 2007;45(1):112-120. doi:10.1128/JCM.00718-06.
73. Mizuta K, Abiko C, Murata T, et al. Frequent Importation of Enterovirus 71 from Surrounding Countries into the Local Community of Yamagata, Japan, between 1998 and 2003. *J Clin Microbiol*. 2005;43(12):6171-6175. doi:10.1128/JCM.43.12.6171-6175.2005.
74. Komatsu H, Shimizu Y, Takeuchi Y, Ishiko H, Takada H. Outbreak of severe neurologic involvement associated with enterovirus 71 infection. *Pediatr Neurol*. 1999;20(1):17-23.
75. Bible JM, Pantelidis P, Chan PKS, Tong CYW. Genetic evolution of enterovirus 71: epidemiological and pathological implications. *Rev Med Virol*. 2007;17(6):371-379. doi:10.1002/rmv.538.
76. Fujimoto T, Chikahira M, Yoshida S, et al. Outbreak of central nervous system disease associated with hand, foot, and mouth disease in Japan during the summer of 2000: detection and molecular epidemiology of enterovirus 71. *Microbiol Immunol*. 2002;46(9):621-627.
77. Ho M, Chen E-R, Hsu K-H, et al. An epidemic of enterovirus 71 infection in Taiwan. *N Engl J Med*. 1999;341(13):929-935.

78. Wang J-R, Tuan Y-C, Tsai H-P, Yan J-J, Liu C-C, Su I-J. Change of Major Genotype of Enterovirus 71 in Outbreaks of Hand-Foot-and-Mouth Disease in Taiwan between 1998 and 2000. *J Clin Microbiol.* 2002;40(1):10-15. doi:10.1128/JCM.40.1.10-15.2002.
79. Liu C-C, Tseng H-W, Wang S-M, Wang J-R, Su I-J. An outbreak of enterovirus 71 infection in Taiwan, 1998: epidemiologic and clinical manifestations. *J Clin Virol.* 2000;17(1):23-30.
80. Wang J-R, Tsai H-P, Chen P-F, et al. An outbreak of enterovirus 71 infection in Taiwan, 1998. II. Laboratory diagnosis and genetic analysis. *J Clin Virol.* 2000;17(2):91-99.
81. Samuda G, Chang W-K, Yeung C-Y, Tang P-S. Monoplegia caused by Enterovirus 71: an outbreak in Hong Kong. *Pediatr Infect Dis J.* 1987;6(2):206-208.
82. Sasidharan CK, Sugathan P, Agarwal R, Khare S, Lal S, Paniker CJ. Hand-foot-and-mouth disease in Calicut. *Indian J Pediatr.* 2005;72(1):17-21.
83. Beig FK, Malik A, Rizvi M, Acharya D, Khare S. Etiology and clinico-epidemiological profile of acute viral encephalitis in children of western Uttar Pradesh, India. *Int J Infect Dis.* 2010;14(2):e141-e146. doi:10.1016/j.ijid.2009.03.035.
84. Saoji VA, others. Hand, foot and mouth disease in Nagpur. *Indian J Dermatol Venereol Leprol.* 2008;74(2):133.
85. Sarma N, Sarkar A, Mukherjee A, Ghosh A, Dhar S, Malakar R. Epidemic of hand, foot and mouth disease in West Bengal, India in August, 2007: a multicentric study. *Indian J Dermatol.* 2009;54(1):26.
86. Kar BR, Dwibedi B, Kar SK. An outbreak of hand, foot and mouth disease in Bhubaneswar, Odisha. *Indian Pediatr.* 2013;50(1):139-142.

87. Kashyap S, Verma GK. Hand-foot-mouth Disease: Outbreak in Shimla. *Indian Pediatr.* 2014;51(2):155-155.
88. Puenpa J, Mauleekoonphairoj J, Linsuwanon P, et al. Prevalence and Characterization of Enterovirus Infections among Pediatric Patients with Hand Foot Mouth Disease, Herpangina and Influenza Like Illness in Thailand, 2012. *PLoS One.* 2014;9(6):e98888.
89. Cardoso MJ, Krishnan S, Tio PH, Perera D, Wong SC. Isolation of subgenus B adenovirus during a fatal outbreak of enterovirus 71-associated hand, foot, and mouth disease in Sibu, Sarawak. *Lancet.* 1999;354(9183):987-991. doi:10.1016/S0140-6736(98)11032-2.
90. Hooi PS, Chua BH, Lee CS, Lam SK, Chua KB. Hand, foot and mouth disease: University Malaya Medical Centre experience. *Med J Malaysia.* 2002;57(1):88-91.
91. Lum LC, Wong KT, Lam SK, et al. Fatal enterovirus 71 encephalomyelitis. *J Pediatr.* 1998;133(6):795-798.
92. Shekhar K, Lye MS, Norlijah O, et al. Deaths in children during an outbreak of hand, foot and mouth disease in Peninsular Malaysia—clinical and pathological characteristics. *Med J Malaysia.* 2005;60(3):297-304.
93. Ooi M, Wong S, Mohan A, et al. Identification and validation of clinical predictors for the risk of neurological involvement in children with hand, foot, and mouth disease in Sarawak. *BMC Infect Dis.* 2009;9(1):3. doi:10.1186/1471-2334-9-3.
94. Goh KT, Doraisingam S, Tan JL, Lim GN, Chew SE. An outbreak of hand, foot, and mouth disease in Singapore. *Bull World Health Organ.* 1982;60(6):965.
95. AbuBakar S, Sam C, Yusof J, et al. Enterovirus 71 Outbreak, Brunei. *Emerg Infect Dis.* 2009;15(1):79-82. doi:10.3201/eid1501.080264.

96. Sudo T, Morita M. [Hand, foot and mouth disease]. *Nihon Rinsho Jpn J Clin Med.* 1971;29(4):1137-1142.
97. Ooi E-E, Phoon M-C, Ishak B, Chan S-H. Seroepidemiology of human enterovirus 71, Singapore. *Emerg Infect Dis.* 2002;8(9):995-997.
98. Lu C-Y, Lee C-Y, Kao C-L, et al. Incidence and case-fatality rates resulting from the 1998 enterovirus 71 outbreak in Taiwan. *J Med Virol.* 2002;67(2):217-223. doi:10.1002/jmv.2210.
99. Zhu Z, Zhu S, Guo X, et al. Retrospective seroepidemiology indicated that human enterovirus 71 and coxsackievirus A16 circulated widely in central and southern China before large-scale outbreaks from 2008. *Viol J.* 2010;7(1):300-305.
100. Ruan F, Yang T, Ma H, et al. Risk Factors for Hand, Foot, and Mouth Disease and Herpangina and the Preventive Effect of Hand-washing. *Pediatrics.* 2011;127(4):e898-e904. doi:10.1542/peds.2010-1497.
101. Park SK, Park B, Ki M, et al. Transmission of Seasonal Outbreak of Childhood Enteroviral Aseptic Meningitis and Hand-foot-mouth Disease. *J Korean Med Sci.* 2010;25(5):677. doi:10.3346/jkms.2010.25.5.677.
102. Chang L-Y, King C-C, Hsu K-H, et al. Risk Factors of Enterovirus 71 Infection and Associated Hand, Foot, and Mouth Disease/Herpangina in Children During an Epidemic in Taiwan. *Pediatrics.* 2002;109(6):e88-e88. doi:10.1542/peds.109.6.e88.
103. Li Y, Dang S, Deng H, et al. Breastfeeding, previous Epstein-Barr virus infection, Enterovirus 71 infection, and rural residence are associated with the severity of hand, foot, and mouth disease. *Eur J Pediatr.* 2013;172(5):661-666. doi:10.1007/s00431-013-1939-1.

104. Qiaoyun F, Xiongfei J, lihuan L, Angao X. Epidemiology and etiological characteristics of hand, foot and mouth disease in Huizhou City between 2008 and 2011. *Arch Virol.* 2013;158(4):895-899. doi:10.1007/s00705-012-1566-6.
105. Li J, Fu Y, Xu A, Zhou Z, Wang W. A Spatial-Temporal ARMA Model of the Incidence of Hand, Foot, and Mouth Disease in Wenzhou, China. *Abstr Appl Anal.* 2014;2014:1-9. doi:10.1155/2014/238724.
106. Zeng M, Li Y-F, Wang X-H, et al. Epidemiology of hand, foot, and mouth disease in children in Shanghai 2007–2010. *Epidemiol Infect.* 2012;140(06):1122-1130. doi:10.1017/S0950268811001622.
107. Ang L-W, Phoon M-C, Wu Y, Cutter J, James L, Chow VT. The changing seroepidemiology of enterovirus 71 infection among children and adolescents in Singapore. *BMC Infect Dis.* 2011;11:270. doi:10.1186/1471-2334-11-270.
108. Yan X-F, Gao S, Xia J-F, Ye R, Yu H, Long J-E. Epidemic characteristics of hand, foot, and mouth disease in Shanghai from 2009 to 2010: Enterovirus 71 subgenotype C4 as the primary causative agent and a high incidence of mixed infections with coxsackievirus A16. *Scand J Infect Dis.* 2012;44(4):297-305. doi:10.3109/00365548.2011.634433.
109. Zeng M, El Khatib NF, Tu S, et al. Seroepidemiology of Enterovirus 71 infection prior to the 2011 season in children in Shanghai. *J Clin Virol.* 2012;53(4):285-289. doi:10.1016/j.jcv.2011.12.025.
110. Zhang J, Sun J, Chang Z, Zhang W, Wang Z, Feng Z. Characterization of hand, foot, and mouth disease in China between 2008 and 2009. *Biomed Env Sci.* 2011;24(3):214-221. doi:10.3967/0895-3988.2011.03.002.

111. Chang L-Y, Tsao K-C, Hsia S-H, et al. Transmission and clinical features of enterovirus 71 infections in household contacts in Taiwan. *Jama*. 2004;291(2):222-227.
112. Chen K-T, Chang H-L, Wang S-T, Cheng Y-T, Yang J-Y. Epidemiologic Features of Hand-Foot-Mouth Disease and Herpangina Caused by Enterovirus 71 in Taiwan, 1998-2005. *Pediatrics*. 2007;120(2):e244-e252. doi:10.1542/peds.2006-3331.
113. Chen S-C, Chang H-L, Yan T-R, Cheng Y-T, Chen K-T. An eight-year study of epidemiologic features of enterovirus 71 infection in Taiwan. *Am J Trop Med Hyg*. 2007;77(1):188-191.
114. Yen F-B, Chang L-Y, Kao C-L, et al. Coxsackieviruses infection in northern Taiwan—epidemiology and clinical characteristics. *J Microbiol Immunol Infect*. 2009;42(1):38-46.
115. Chan Y-F, Sam I-C, Wee K-L, Abubakar S. Enterovirus 71 in Malaysia: A decade later. *Neurol Asia*. 2011;16(1):1-15.
116. Suzuki Y, Taya K, Nakashima K, et al. Risk factors for severe hand foot and mouth disease. *Pediatr Int*. 2010;52(2):203-207. doi:10.1111/j.1442-200X.2009.02937.x.
117. Frydenberg A, Starr M. Hand, foot and mouth disease. *Aust Fam Physician*. 2003;32(8):594-595.
118. Wong SSY, Yip CCY, Lau SKP, Yuen KY. Human enterovirus 71 and hand, foot and mouth disease. *Epidemiol Infect*. 2010;138(08):1071-1089. doi:10.1017/S0950268809991555.
119. Ma E, Fung C, Yip SH, Wong C, Chuang SK, Tsang T. Estimation of the basic reproduction number of enterovirus 71 and coxsackievirus A16 in hand, foot, and mouth disease outbreaks. *Pediatr Infect Dis J*. 2011;30(8):675-679.
120. Dale MH. Hand, foot and mouth disease—1973. *Br J Gen Pr*. 1975;25(150):35-38.

121. Chang L-Y, Lin T-Y, Hsu K-H, et al. Clinical features and risk factors of pulmonary oedema after enterovirus-71-related hand, foot, and mouth disease. *The Lancet*. 1999;354(9191):1682-1686.
122. Lee M-S, Chiang P-S, Luo S-T, et al. Incidence Rates of Enterovirus 71 Infections in Young Children during a Nationwide Epidemic in Taiwan, 2008–09. Walson JL, ed. *PLoS Negl Trop Dis*. 2012;6(2):e1476. doi:10.1371/journal.pntd.0001476.
123. Wallinga J, Teunis P. Different epidemic curves for severe acute respiratory syndrome reveal similar impacts of control measures. *Am J Epidemiol*. 2004;160(6):509-516. doi:10.1093/aje/kwh255.
124. Choi BCK, Pak AWP. A simple approximate mathematical model to predict the number of severe acute respiratory syndrome cases and deaths. *J Epidemiol Community Health*. 2003;57(10):831-835.
125. Li Y, Zhang J, Zhang X. Modeling and Preventive Measures of Hand, Foot and Mouth Disease (HFMD) in China. *Int J Environ Res Public Health*. 2014;11(3):3108-3117. doi:10.3390/ijerph110303108.
126. Chang H-L, Chio C-P, Su H-J, et al. The Association between Enterovirus 71 Infections and Meteorological Parameters in Taiwan. Yang Y, ed. *PLoS ONE*. 2012;7(10):e46845. doi:10.1371/journal.pone.0046845.
127. Ma E, Lam T, Wong C, Chuang SK. Is hand, foot and mouth disease associated with meteorological parameters? *Epidemiol Infect*. 2010;138(12):1779-1788. doi:10.1017/S0950268810002256.
128. Huang Y, Deng T, Yu S, et al. Effect of meteorological variables on the incidence of hand, foot, and mouth disease in children: a time-series analysis in Guangzhou, China. *BMC Infect Dis*. 2013;13(1):134.

129. Abad FX, Pinto RM, Bosch A. Survival of enteric viruses on environmental fomites. *Appl Environ Microbiol.* 1994;60(10):3704-3710.
130. Wang JF, Guo Y-S, Christakos G, et al. Hand, foot and mouth disease: spatiotemporal transmission and climate. *Int J Health Geogr.* 2011;10(25):1-10.
131. Bélanger M, Gray-Donald K, O'loughlin J, Paradis G, Hanley J. Influence of Weather Conditions and Season on Physical Activity in Adolescents. *Ann Epidemiol.* 2009;19(3):180-186. doi:10.1016/j.annepidem.2008.12.008.
132. Lofgren E, Fefferman NH, Naumov YN, Gorski J, Naumova EN. Influenza seasonality: underlying causes and modeling theories. *J Virol.* 2007;81(11):5429-5436.
133. Singapore Ministry of Health. Hand, foot & Mouth Disease: Updates. [https://www.moh.gov.sg/content/moh\\_web/home/diseases\\_and\\_conditions/h/hand\\_foot\\_mouth\\_disease.html](https://www.moh.gov.sg/content/moh_web/home/diseases_and_conditions/h/hand_foot_mouth_disease.html).
134. Li J, Lin C, Qu M, et al. Excretion of enterovirus 71 in persons infected with hand, foot and mouth disease. *Virol J.* 2013;10(31):422X - 10.
135. Zhu F-C, Meng F-Y, Li J-X, et al. Efficacy, safety, and immunology of an inactivated alum-adjutant enterovirus 71 vaccine in children in China: a multicentre, randomised, double-blind, placebo-controlled, phase 3 trial. *The Lancet.* 2013;381(9882):2024-2032.
136. Li R, Liu L, Mo Z, et al. An Inactivated Enterovirus 71 Vaccine in Healthy Children. *N Engl J Med.* 2014;370(9):829-837. doi:10.1056/NEJMoa1303224.
137. Zhu F, Xu W, Xia J, et al. Efficacy, Safety, and Immunogenicity of an Enterovirus 71 Vaccine in China. *N Engl J Med.* 2014;370(9):818-828. doi:10.1056/NEJMoa1304923.
138. Lee T-C, Guo H-R, Su H-JJ, Yang Y-C, Chang H-L, Chen K-T. Diseases Caused by Enterovirus 71 Infection. *Pediatr Infect Dis J.* 2009;28(10):904-910. doi:http://dx.doi.org/10.1097/inf.0b013e3181a41d63.

139. Solomon T, Lewthwaite P, Perera D, Cardosa MJ, McMinn P, Ooi MH. Virology, epidemiology, pathogenesis, and control of enterovirus 71. *Lancet Infect Dis*. 2010;10(11):778-790.
140. Tong CW, Bible JM. Global epidemiology of enterovirus 71. *Future Virol*. 2009;4(5):501-510.
141. Yip CCY, Lau SKP, Woo PCY, Yuen K-Y. Human enterovirus 71 epidemics: what's next? *Emerg Health Threats J*. 2013;6(0). doi:10.3402/ehth.v6i0.19780.
142. Ooi MH, Wong SC, Lewthwaite P, Cardosa MJ, Solomon T. Clinical features, diagnosis, and management of enterovirus 71. *Lancet Neurol*. 2010;9(11):1097-1105.
143. Fang Y, Wang S, Zhang L, et al. Risk factors of severe hand, foot and mouth disease: A meta-analysis. *Scand J Infect Dis*. 2014;46(7):515-522. doi:10.3109/00365548.2014.907929.
144. Zhao YY, Jin H, Zhang XF, Wang B. Case-fatality of hand, foot and mouth disease associated with EV71: a systematic review and meta-analysis. *Epidemiol Infect*. 2015;143(14):3094-3102. doi:10.1017/S095026881500028X.
145. Chia M-Y, Chiang P-S, Chung W-Y, Luo S-T, Lee M-S. Epidemiology of Enterovirus 71 Infections in Taiwan. *Pediatr Neonatol*. 2014;55(4):243-249. doi:10.1016/j.pedneo.2013.07.007.

Figure captions

**Figure 1: Temporal patterns of HFMD outbreaks in Asia, by latitude.** Left: Plot Digitizer is used to convert charts into numbers. White boxes are the months where HFMD cases fall below the year's median. The remaining cells are then shaded into 4 darker shades by octiles. The regions of China were based on Wang et al's classification<sup>16</sup> (C standing for central). The regions are arranged by latitude. South China, Hong Kong and Taiwan have subtropical climates. Areas further north are temperate, while the South East Asian regions are tropical. Right: The coefficient of variation is the ratio of the standard deviation to its mean, and the proportion of cases in top 3 months is the proportion of cases of the 3 months with highest incidence to the annual incidence. Points represent one year per region. The lines are obtained from ordinary least squares regression with Latitude as the independent variable, and show how clearly defined epidemics become the further north from the equator.

**Figure 2: Temporal incidence of HFMD or enterovirus and climatic factors for four Asian cities spanning temperate, subtropical and tropical latitudes.** Top: incidence, temperature, absolute and relative humidity. Top panels indicate incidence (data in gray, mean and 95% interval in black) for the time period Jan 2001-Mar 2012 (Tokyo), Jan 2001-Dec 2009 (Hong Kong, Peoples' Republic of China), Jan 2001-Dec 2011 (Taiwan, Republic of China), Jan 2001-Jan 2012 (Singapore). Middle and bottom panels show mean daily temperature, absolute and relative humidity at Tokyo Narita, Hong Kong International, Taipei Taoyuan International and Singapore Changi airports, downloaded from the Weather Underground. Bottom: Coefficients of meteorological variables at 0 to 2 week time lags in autoregressive models of z-scored HFMD case counts to facilitate comparison between locations. Each city is analyzed separately using a model in which HFMD incidence in week  $t$  is (auto)regressed on incidence in weeks  $t - 1$  up to  $t - 3$  (using the Akaike information criterion to select the order of the autoregression component) and, independently, on each

meteorological variable. Weather parameters are not regressed together in a single model due to collinearity. The effect of weather on HFMD incidence can be seen by coefficient mean (points) and 95% confidence intervals (lines), colored red if statistically significant at the 5% level.

**Figure 3: Historical establishment of HFMD in Asia.** Dots represent a reported outbreak in that year, with the main causal agent written below. Boxes with right arrow indicate endemicity of HFMD, evidenced by repeated reporting of HFMD. Boxes with left arrow indicate seroepidemiologic evidence that the pathogen is already in existence, even if no significant outbreaks were documented previously. Triangles indicate the point where data on HFMD started to be collected systematically, for example, through government surveillance. The length of the left arrows are arbitrary as there is no way to know how long has HFMD been circulating prior to the tests.

**Figure 4: Excerpts from studies on risk factors.** Ruan et al's case-control study in Qiaosi<sup>100</sup>, China, was conducted by taking 273 diagnosed HFMD/Herpangina as cases (age 6 years or younger), and 273 stratified random sample as controls. Park et al's case-control study uses hospital cases of enteroviral Aseptic Meningitis (n = 205) and HFMD (n=116), and non-enteroviral disease controls (n = 170). Their case-crossover design uses only cases. 1-7 days prior to admission was set as the hazard period, and 22-28 days prior to admission was set as the non-hazard period. Both studies gather data via questionnaires. White points indicate non-adjusted odds ratios (ORs), black, adjusted (aOR). For the effect of sex (bottom right), confidence intervals are omitted from complete case notifications, and alternatingly colored gray and white for visual distinction.

**Figure 5: HFMD cases by age and estimates of incubation period.** Left: Each line indicates a unique data set (total 79 lines). Distributions within age ranges were assumed to be constant. The black dots are average proportion for that age (with 95% CI). Right:

Reported incubation periods for HFMD, by year of publication and provision of evidence to support claimed period. Lines indicate that the incubation period “is X days”. Gray bars indicate that the incubation period is “usually” or “typically” X days. Gray bars with lines shows an extended interval “can be up to X days”. The point indicates a median. **Notes:** (1) provides information within the paper which is inconsistent this estimate. (2) uses generation interval distribution as a proxy for incubation period. (3) Cites a US CDC factsheet on aseptic meningitis, which in turn provides no supporting evidence. (4) Cites Goh et al<sup>94</sup>.

ACCEPTED

## Tables


**Table 1: Estimates of symptomatic proportion from the literature.**

Location	Ref	Year	Inclusion Criteria	Number Symptomatic	Number Infected	Percent Symptomatic (95% CI)
Taiwan	<sup>121</sup>	1998	Stratified sampling. Infection determined using serology vs EV-A71 and recall of previous infection history	140	484	28.9 (24.9-33.0)
Shanghai	<sup>109</sup>	2010 - 2011	Routine blood samples and recall of previous infection history	12	122	9.8 (4.6 – 15.1)
Taiwan	<sup>111</sup>	2001 - 2002	Household members of hospital symptomatic cases were swabbed for EV-A71	119 (Excl 94 symptomatic cases)	176 (Excl 94 symptomatic cases)	67.6 (60.7-74.5)
Taiwan	<sup>122</sup>	2006 - 2009	Prospective cohort of neonates with repeat serology (EV-A71) and swabs taken upon subsequent illness	20	28	71.4 (54.7-88.2)
Total				291	810	35.9


						(32.6-39.2)
--	--	--	--	--	--	-------------

ACCEPTED


**Figure 1.**




**Figure 2.**




**Figure 3.**



**Figure 4.**



**Figure 5.**



## Appendix

### 1 Timing and seasonality of HFMD outbreaks

An autoregressive model was employed to investigate the effect of meteorological variables after correcting for contagion via autoregression.

A lag 2 model can be specified as follows:

$$\begin{aligned} \text{HFMD}_t = & c + A_1 \text{HFMD}_{t-1} + A_2 \text{HFMD}_{t-2} + B_0 \text{WEA}_t \\ & + B_1 \text{WEA}_{t-1} + B_2 \text{WEA}_{t-2} + \epsilon_t \end{aligned}$$

The A coefficients are the coefficients for the AR terms, while the B coefficients represent how a change in weather is correlated with changes in HFMD incidence. The number of lag terms are determined by the Akaike information criterion values of the regression models.

This same model was used for 4 countries – Japan (lag 2), Hong Kong (lag 3), Taiwan (lag 3) and Singapore (lag 2), for 3 meteorological parameters – temperature, absolute humidity and relative humidity. As this model carries autocorrelated terms, we used Generalized Least Squares (GLS) for model fitting. Coefficients from the fitted models are presented in tables A1 to A3.

Table A1: Coefficients for autoregressive model using temperature as predictor. The outcome variable is the number of reported HFMD cases per sentinel per week (Japan), the number of reported HFMD cases per GP per week (Hong Kong), the number of reported severe enterovirus cases per week (Taiwan), and the number of reported HFMD cases per week (Singapore). To facilitate comparability, the incidence measures were standardized to have mean 0 and variance 1.

Location	Lag (weeks)	Autoregression term (95%CI)	p	Temperature (°C) coefficient (95%CI)	p
Japan	0	-	-	0.016 (0.008, 0.024)	<0.001
	1	1.517 (1.452, 1.582)	<0.001	-0.009 (-0.018, 0.001)	0.07
	2	-0.601 (-0.665, -0.536)	<0.001	-0.001 (-0.009, 0.006)	0.723
Hong Kong	0	-	-	0.012 (-0.007, 0.031)	0.204
	1	0.756 (0.678, 0.835)	<0.001	-0.009 (-0.031, 0.013)	0.415
	2	-0.079 (-0.178, 0.02)	0.116	-0.01 (-0.031, 0.012)	0.388
Taiwan	0	-	-	0.014 (-0.003, 0.032)	0.108
	1	-0.181 (-0.26, -0.102)	<0.001	0.011 (-0.007, 0.029)	0.222
	2	0.062 (-0.019, 0.142)	0.132	-0.003 (-0.02, 0.015)	0.779
	3	0.192 (0.113, 0.271)	<0.001	0 (-0.018, 0.017)	0.968

Singapore	0	-	-	0.049 (0.017, 0.081)	0.003
	1	1.42 (1.349, 1.491)	<0.001	-0.041 (-0.082, 0.001)	0.054
	2	-0.458 (-0.53, -0.387)	<0.001	0.01 (-0.023, 0.042)	0.564

Table A2: Coefficients for autoregressive model using relative humidity as predictor. The outcome variable is the same as the temperature model.

Location	Lag (weeks)	Autoregression term (95%CI)	p	RH (°C) coefficient (95%CI)	p
Japan	0	-	-	-0.001 (-0.003, 0.001)	0.363
	1	1.567 (1.503, 1.631)	<0.001	0.002 (-0.001, 0.004)	0.186
	2	-0.634 (-0.699, -0.569)	<0.001	0.001 (-0.001, 0.003)	0.231
Hong Kong	0	-	-	0.002 (-0.004, 0.007)	0.578
	1	0.942 (0.864, 1.021)	<0.001	-0.002 (-0.008, 0.004)	0.483
	2	-0.242 (-0.349, -0.135)	<0.001	-0.003 (-0.009, 0.004)	0.413
Taiwan	0	-	-	0.001 (-0.005, 0.007)	0.72
	1	-0.178 (-0.257, -0.1)	<0.001	0.001 (-0.005, 0.007)	0.755

	2	0.062 (-0.018, 0.142)	0.13	0.003 (-0.004, 0.009)	0.426
	3	0.19 (0.111, 0.269)	<0.001	-0.001 (-0.006, 0.005)	0.849
Singapore	0	-	-	-0.009 (-0.016, -0.003)	0.004
	1	1.43 (1.36, 1.501)	<0.001	0.008 (0, 0.016)	0.055
	2	-0.469 (-0.539, -0.398)	<0.001	-0.003 (-0.01, 0.003)	0.349

Table A3: Coefficients for autoregressive model using absolute humidity as predictor. The outcome variable is the same as the temperature model. coefficients for autoregressive model using temperature as predictor.

Location	Lag (weeks)	Autoregression term (95%CI)	p	AH (°C) coefficient (95%CI)	p
Japan	0	-	-	0.021 (0.013, 0.029)	<0.001
	1	0.847 (0.766, 0.928)	<0.001	0.003 (-0.005, 0.011)	0.484
	2	-0.141 (-0.221, -0.061)	0.001	0.005 (-0.003, 0.013)	0.228
Hong Kong	0	-	-	0.013 (-0.005, 0.031)	0.153
	1	0.781 (0.703, 0.86)	<0.001	-0.012 (-0.033, 0.01)	0.285
	2	-0.102 (-0.202, -0.001)	0.047	-0.004 (-0.026, 0.017)	0.703
	3	0.195 (0.117, 0.274)	<0.001	0.013 (-0.005, 0.031)	0.156

Taiwan	0	-	-	0.013 (-0.002, 0.029)	0.096
	1	-0.18 (-0.259, -0.101)	<0.001	0.011 (-0.004, 0.027)	0.158
	2	0.064 (-0.016, 0.144)	0.117	-0.002 (-0.018, 0.014)	0.814
	3	0.194 (0.115, 0.273)	<0.001	0.001 (-0.015, 0.017)	0.893
Singapore	0	-	-	0.034 (0.012, 0.056)	0.003
	1	1.42 (1.349, 1.492)	<0.001	-0.029 (-0.058, 0)	0.047
	2	-0.459 (-0.53, -0.388)	<0.001	0.007 (-0.015, 0.03)	0.536

The AR coefficients are generally statistically significant across models.

Significant AR terms indicate that incidence is highly autocorrelated, which is expected as the contagious nature of HFMD is the primary driver of temporal patterns of incidence. The primary parameters of interest are the coefficients of the meteorological variables, which represent how much temperature/humidity affects the z-score of incidence, after controlling for contagion. The results are tabulated in Tables A1 to A3, summarised in Figure 1 and the paper itself.

Incidence data is obtained from various sources, summarised in Table A4. For Japan and Hong Kong, “Cases per sentinel” and “Cases per consultation” are used instead of the actual number of notified cases because these data are from voluntary sentinel reporting. Thus, actual notified cases will increase with an increase of GP sentinel participation rate.

Table A4: Data source for incidence.

Country	Data	Source
Singapore	Notified Cases (low underreporting, as notification is mandatory and laws are strict)	Ministry of Health, Singapore
Japan	Notified Cases, Cases per sentinel (not mandatory reporting)	National Institute of Infectious Diseases, Japan (Released weekly)
Hong Kong	Notified Cases per 1000 GP consultations (Not mandatory reporting)	Department of Health (Website and digitized from historical documents)
Taiwan	Enterovirus with complications	Taiwan National Infectious Disease Statistics System

## 2 Odds Ratio from Figure 3

Paper	Type	Location	Details	OR
97	Serology	Singapore	1200 serum samples, aged 1-17 years. Children were split into 3 equal groups of age 1-6, 7-12 and 13-17	0.788, 0.608, 1.019
87	Serology	Taiwan	1800 children between 6 mth to 6 years old	0.941, 0.763, 1.159
91	Cases	Whole China	2008, 2009 reported cases for entire China (almost 1million cases). Controlled for population male/female ratio of entire population. 632.84m girls, 667.20m boys	1.658 (1.648, 1.667) 1.605 (1.599, 1.611)
33	Cases	Jiangsu, China	2008, 2009 reported cases for Jiangsu, zhenjiang. 6324 HFMD cases. Controlled for population male/female	1.543 (1.424,1.673) 1.300 (1.218,1.387)
20	Cases	Beijing, China	157k cases in 2008 to 2012, each OR represents a particular year. Not controlled for population male/female	1.568, 1.535, 1.517, 1.493, 1.494
34	Cases	Guangdong, China	48876 cases in 2008. Do not have denominator for population male/female.	1.85
35	Cases	Guangdong, China	Incidence ratios for 2008,9,10,11. Total of 641k cases.	1.84, 1.81, 1.74, 1.68
89	Cases	Huizhou	42012 cases from 2008-2011. Incidence ratio	1.65
92	Cases	Wenzhou	103k cases from 2010-2012. Not controlled for male/female ratio.	1.639, 1.691, 1.633
93	Cases	Shenzhen	Total 12132 reported cases for 2009, 2010 and 2011. Not controlled for male/female ratio	1.851, 1.697, 1.854
94	Cases	Changchun	17464 cases reported from 2008-2011. Not controlled for male/female ratio in population.	1.480
5	Cases	Singapore	Incidence ratios for 2001-2007. All cases in Singapore.	1.328, 1.420, 1.607, 1.309, 1.268, 1.236, 1.214
90	Cases	Singapore	year 2000. All cases reported to MOH	1.700
70	Cases	Thailand	Reported cases in 2012.	1.496
71	Cases	Thailand	Reported cases from 2003 to 2012	1.212, 1.297, 1.321, 1.323, 1.330, 1.320, 1.341, 1.360,

			28058 hospital cases -> not reported as case because the other data above are all surveillance data, not hospital. 1.165 is the incidence ratio of severe cases, 473/17206 boys and 257/10852 girls	1.372, 1.429
95	Hospital	Shanghai, China	Xian Jiaotong University and Xian children hospital Apr-Oct 2011. 116 (83m, 33f) severe cases of HFMD. 318 hospital cases (211m, 107f)	1.1653 (0.999, 1.359)
88	Severe	Xian, China	Beijing Youan hospital June-Oct 2010, 233 (158m, 75f) severe, 1104 total (667m, 437f)	1.454 (0.8867, 2.3844)
96	Severe	Beijing, China	Zhujiang hospital Mar to Dec 2010. 542 children diagnosed with HFMD. CNS: 34m, 13f Total: 349m, 193f	1.498 (1.103, 2.035)
36	Central Nervous System	Guangdong, China		1.495 (0.769, 2.906)

### 3 Data source for Figure 5 (left)

Page	Date	Data source	Size	Type	Location	Found in:
30	Apr 7 to May 11, 2010	9th People's Hospital of Nanchang	109	Hospital cases	Nanchang, China	Page 5, Figure 4 (A)
31	May 2008 to Dec 2009	Weekly Reports to China CDC	1065000	Surveillance	Mainland China	Figure 1 and 2
32	2007-2011	China Information System for Disease Control and Prevention	421488	Surveillance	Shandong, China	Table 1
33	May 2008 to Oct 2009.	Reported cases to Jiangsu CDC	6324	Surveillance	Jiangsu, China	Figure 3
34	2008	Reported cases to Guangdong HFMD web-based surveillance system (871 clinics)	48876	Surveillance	Guangdong, China	Figure 2
35	2008-2011	Guangdong surveillance data	641318	Surveillance	Guangdong, China	Table 1, proportion
36	2010	Laboratory samples	542	Laboratory	Guangdong, China	Figure 1 and 2
37	2008-2010	Reported cases under Yunnan HFMD web-based surveillance system (871 clinics)	75109	Surveillance	Yunnan, China	Table 2
85	Apr 30 to June 26, 2008	All 6mth to 6yr cases from Qiaosi, Zhejiang	273	Case-control	Zhejiang, China	Table 1
91	2008-2009	China surveillance data	1500000	Surveillance	Whole China	Table 1
92	2010-2012	Reported cases to Wenzhou CDC	103671	Surveillance	Wenzhou, China	Table 1
95	2007-2010	Cases from Children's Hospital of Fudan University	28058	Hospital cases	Shanghai, China	Table 1
96	Jun-Oct 2010	Cases from Beijing Youan Hospital	1104	Hospital cases	Beijing, China	Figure 2
98	Jan 2009-Dec 2010	Cases from Children's Hospital of Fudan University	3208	Hospital cases	Shanghai, China	Figure 3
99	2011	Cases from Children's Hospital of Fudan	8020	Hospital cases	Shanghai, China	Table 1

		University				
100	May 2008 to April 2009	China surveillance data	765220	Surveillance	China	Figure 4
6	2008	Laboratory confirmed cases from Chang Gung Children's Hospital	280	Hospital cases	Taoyuan, Taiwan	Table 1
7	Jan 2004-Dec 2009	Laboratory confirmed CA6 cases from Chang Gung Memorial Hospital	229	Hospital cases	Taoyuan, Taiwan	Figure 3
58	Apr-Dec 1998	Laboratory confirmed EV71 cases from Taiwan MOH passive surveillance	119	Surveillance	Tainan, Chiayi, Taiwan	Figure 3
101	Feb 2001-Aug 2002	Chang Gung Children's Hospital, ages 0-40	256	Cohort study	Taiwan	Table 2
102	Mar 1998-Dec 2005	Taiwan surveillance data, ages 0-15	8000	Surveillance	Taiwan	Figure 4
103	Mar 1998 - Dec 2005	Taiwan surveillance data, severe cases	1584	Surveillance	Taiwan	Figure 2
104	Jan 1999 to Dec 2006.	Coxsackievirus confirmed cases from National Taiwan University Hospital	457	Hospital cases	Taiwan	Figure 3
60	2008	Laboratory confirmed EV71 cases from voluntary reporting to Public Health Laboratory of the Dept of Health	98	Surveillance	Hong Kong	Figure 2
61	2001-2009	HK GP-based sentinel surveillance and Public Health Laboratory of the Dept of Health	3512	Surveillance	Hong Kong	Figure 4
63	Oct 2003-Feb 2004	Cases from 1 outpatient clinic	81	Clinical cases	Calicut, India	Page 2, Results
67	Sep 2009 - Nov 2009	Hospitals and community in urban areas	78	Clinical cases	Bhubaneswar, Odisha, India	Table 1
49	1978	Cases from Gifu Prefectural Hospital	108	Hospital cases	Gifu Prefecture, Japan	Figure 2
105	2004-2008	Survey	166	Survey	Yokohama City, Japan	Table 3
39	2008-2009	Enterovirus-positive cases	1214	Survey	Chungnam, Korea	Figure 2

86	2002-2003	HFMD cases from 3 general hospitals	116	Hospital Case-control study	Seoul, Gyeongju, Pohang Korea	Table 1
75	May 1997 to June 2001	HFMD cases investigated in the University Malaya Medical Centre	467	Laboratory	Malaysia	Figure 1
106	1997-2008	EV71 and CA16 confirmed cases	145	Laboratory	Malaysia	Page 7, Host Factors
5	2001-2007	All Singapore cases, GP reporting and laboratory cases	83970	Surveillance	Singapore	Table 1
79	Sep - Dec 1981	Notified cases	270	Clinical cases	Singapore	Table 1
90	Early Sep 2000 to Mar 2001	Notifications to the Ministry of the Environment	175	Clinical cases	Singapore	Table 1
71	Jan 2003 - Nov 2012	Cases reported to Ministry of Public Health	20281	Surveillance	Thailand	Table 1
107	2008-2013	Laboratory tested cases from King Chulalongkorn Memorial Hospital	1182	Hospital cases	Bangkok, Thailand	Figure 5
72	2005	Paediatric hospital in HCM City	764	Hospital cases	Ho Chi Minh City, Vietnam	Figure 2B

#### 4 PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
<b>TITLE</b>			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	Page 1
<b>ABSTRACT</b>			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria; participants; and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	Page 3
<b>INTRODUCTION</b>			
Rationale	3	Describe the rationale for the review in the context of what is already known.	Page 5
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	Page 5 and Page 19 (Table 1)
<b>METHODS</b>			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	Attached as supporting document
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	Page 7
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	Page 7
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Page 7
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	Page 7 and Page 19 (Table 1)
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	7

Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	Data sought answers questions stated in page 19 (Table 1)
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	Most studies are not synthesized due to small sample size. Potential biases are discussed throughout the paper.
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	Risk factors: OR, page 10-11
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., $I^2$ ) for each meta-analysis.	Epidemic patterns: page 21 Climate patterns: page 22

Page 1 of 2

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	Cumulative evidence that might be biased were not synthesized.
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	NA
<b>RESULTS</b>			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	Attached as supporting document
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	Details are found in Figures and as supporting doc
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	Studies that might be biased are identified throughout the review.
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals,	Synthesis of results was avoided when i) data is too

		ideally with a forest plot.	sparse; or ii) studies are too different.
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	For these studies, we have summarised the key results into figures for comparison.
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	Attached as supporting document. (Gender and Age data synthesis)
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	Cumulative evidence that might be biased were not synthesized.
<b>DISCUSSION</b>			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	Page 19
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	Page 19
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	Page 19
<b>FUNDING</b>			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	Page 1

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 6(6): e1000097. doi:10.1371/journal.pmed.1000097

For more information, visit: [www.prisma-statement.org](http://www.prisma-statement.org).

## 5 PRISMA 2009 Flow Diagram

