

Chikungunya and Zika Virus Cases Detected against a Backdrop of Endemic Dengue Transmission in Vietnam

Nguyen Than Ha Quyen,¹ Duong Thi Hue Kien,¹ Maia Rabaa,^{1,2} Nguyen Minh Tuan,³ Tran Thuy Vi,¹ Le Van Tan,¹ Nguyen Thanh Hung,³ Ha Manh Tuan,⁴ Ta Van Tram,⁵ Nguyen Le Da Ha,⁶ Han Khoi Quang,⁷ Nguyen Quoc Doanh,⁸ Nguyen Van Vinh Chau,⁹ Bridget Wills,^{1,2} and Cameron P. Simmons^{1,2,10*}

¹Oxford University Clinical Research Unit, Wellcome Trust Major Overseas Programme, Hospital for Tropical Diseases, Ho Chi Minh City, Vietnam; ²Nuffield Department of Medicine, Centre for Tropical Medicine, University of Oxford, Oxford, United Kingdom; ³Children's Hospital No. 1, Ho Chi Minh, Vietnam; ⁴Children's Hospital No. 2, Ho Chi Minh, Vietnam; ⁵Tien Giang Hospital, My Tho, Vietnam; ⁶Dong Nai Children Hospital, Bien Hoa, Vietnam; ⁷Binh Duong Hospital, Thu Dau Mot, Vietnam; ⁸Long An Hospital, Tan An, Vietnam; ⁹Hospital for Tropical Diseases, Ho Chi Minh, Vietnam; ¹⁰Department of Microbiology and Immunology at the Doherty Institute, University of Melbourne, Parkville, Australia

Abstract. Between 2010 and 2014, four chikungunya and two Zika virus infections were identified among 8,105 febrile children in southern Vietnam. Zika viruses were linked to French Polynesian strains, chikungunya to Cambodian strains. Against a backdrop of endemic dengue transmission, chikungunya and Zika present an additional arboviral disease burden in Vietnam.

INTRODUCTION

The emergence of the Zika virus (ZIKV) in the Western Pacific and Latin America, followed by the World Health Organization (WHO) declaration of a global public health emergency, has brought renewed attention to this and other epidemic arboviral infections. ZIKV is a flavivirus, whereas chikungunya virus (CHIKV) is an alphavirus. Both viruses are transmitted to humans by *Aedes* spp. mosquitoes. Clinical signs and symptoms of CHIKV and ZIKV infections are similar to those of dengue, particularly in the first few days of illness when the symptoms are non-specific, including but not limited to fever, arthralgia, maculopapular rash, muscle and joint pain, malaise, and headache.¹

There are three genotypes of CHIKV: Asian, West African, and East/Central/South African (ECSA). The Asian genotype is thought to have circulated in southeast Asia for many decades without ever posing a perceivable public health problem. More recently, one lineage of the CHIKV ECSA genotype, the Indian Ocean Lineage (IOL), emerged from east Africa and seeded large epidemics in the Indian Ocean region, and later in Europe and Asia.^{2–5} Sporadic cases of illness caused by the CHIKV IOL were first detected in Cambodia in 2011.⁵ In neighboring Vietnam, no virologically confirmed cases of CHIKV infection have been described.

ZIKV has also evolved into three distinct genotypes of similar geographic distribution: West African (Nigerian cluster), East African (MR766 prototype cluster), and Asian.⁶ ZIKV was first detected in humans in 1954 but has recently emerged as a global threat to public health^{7,8} because it can be an infectious teratogen to the unborn fetus and an occasional cause of Guillain-Barré syndrome in adults.^{9,10} Since 2015, 67 countries have reported outbreaks (WHO situation report, October 27, 2016; <http://www.who.int/emergencies/zika-virus/situation-report/27-october-2016/>). Prior to 2016, only sporadic ZIKV infections have been

reported in southeast Asia; sequence analysis shows that historical southeast Asian viruses cluster within the Asian ZIKV lineage.^{7,11} In Vietnam, the first two confirmed cases of locally acquired ZIKV infections were reported in southern Vietnam in April 2016 (<http://www.who.int/csr/don/12-april-2016-zika-viet-nam/en/>).

Against a backdrop of seasonally hyperendemic dengue transmission, this retrospective study sought to understand the burden and trajectory of CHIKV and ZIKV infections in Vietnam from 2010 to 2014.

THE STUDY

This was a retrospective diagnostic study. The study population was children from 1 to 15 years of age with fever of less than 72 hours and clinical symptoms that suggested to the attending physician that dengue could be a possible diagnosis. Study participants were enrolled at the outpatient clinics of seven hospitals in southern Vietnam between October 2010 and December 2014. The detailed inclusion and exclusion criteria and study procedures are described elsewhere.¹² Reverse transcription polymerase chain reaction (RT-PCR) tests for ZIKV¹³ and CHIKV¹⁴ were performed on stored frozen plasma samples from study participants that were tested negative in a dengue virus (DENV) RT-PCR assay that has been described extensively elsewhere.¹⁵ Virus surveillance was only carried out on stored specimens from patients who had consented to having their samples used for future research. The study protocol was reviewed and approved by the institutional review boards of the seven hospitals and the Oxford Tropical Research Ethical Committee (OxTREC 592-16).

RESULTS

During the study period, 8,105 participants were enrolled and 2,203 dengue cases identified on the basis of direct virological confirmation (non-structural protein 1 or RT-PCR positive) or IgM seroconversion (Panbio IgM capture enzyme-linked immunosorbent assay) as described previously.¹² Among all DENV PCR-negative samples (6,037), including those showing serological evidence of recent DENV

* Address correspondence to Cameron P. Simmons, Department of Microbiology and Immunology at the Doherty Institute, University of Melbourne, Parkville, Victoria 3010, Australia. E-mail: csimmons@unimelb.edu.au

infection, a total of 5,617 frozen acute plasma samples (collected within 72 hours of fever onset) were of adequate sample volume for CHIKV and ZIKV testing. Of the 5,617 samples tested, four CHIKV (0.07%) and two ZIKV (0.035%) cases were identified.

Clinical manifestations. All four CHIKV cases were detected between August and November 2012—one case lived in Ho Chi Minh City and three in Binh Duong Province (Figure 1). Three of the infections were relatively mild, with fever, rash and mild to moderate systemic symptoms





FIGURE 1. Residential location of the study patients and study hospitals. Locations of the chikungunya and Zika cases are shown with black circles and stars, respectively; hospitals are shown as black crosses. Two chikungunya cases were located approximately 20 km from the border with Cambodia.



reported. One case, a 13-year-old boy, was briefly hospitalized. All patients made unremarkable recoveries.

The two ZIKV infections were identified in a 12-year-old from Long An Province, and a 4-year-old from Cu Chi District, Ho Chi Minh City, in January and June 2013, respectively. The 12-year-old was briefly hospitalized. Both children recovered fully.

Viral genetics. Nucleotide sequences of the 1,743-nucleotide (nt) CHIKV E1 gene with partial 3' untranslated region and 1,512-nt ZIKV E gene were determined using Miseq Illumina (San Diego, CA) techniques. Sequence assembly and alignment of these sequences to global databases was performed using CLC Workbench 9.1 (Qiagen, Redwood City, CA). The four Vietnamese CHIKV E1 sequences (GenBank Accession KY131437, KY131438, KY131439, and KY131440) shared 99.77–100% pairwise nucleotide identity and carried the E1-A226V mutation reported to increase infectiousness for *Aedes albopictus*.¹⁶ Phylogenetic analysis indicated that the four Vietnamese samples were of the Indian Ocean Outbreak lineage within the ECSA genotype and clustered closely with 2011 Cambodian sequences (99.71–99.94% identity) (Figure 2A). Notably, two of the Vietnamese cases in Binh Duong Province resided approximately 20 km from the Vietnam–Cambodian border.

The two ZIKV E gene sequences captured in Vietnam in 2013 (GenBank Accession KY131441 and KY131442) differed by a single nucleotide. There was strong bootstrap support to place these sequences among other Asian lineage ZIKV sampled from southeast Asia and the Western Pacific, with a close phylogenetic link to a sequence obtained from French Polynesia in late 2013 (Figure 2B). The similarity of the two Vietnamese sequences coupled with their spatial and temporal spread (42 km and 5 months apart), suggests that ZIKV may have circulated at very low levels or caused mild enough illness to be largely undetected within southern Vietnam in 2013. Alternatively, the viruses could represent multiple isolated importations of ZIKV from another population. As this investigation was conducted retrospectively and travel histories were not collected at enrollment, we cannot definitively determine whether the viruses detected in this study were locally acquired or travel related. Importantly, these sequences do not appear to be phylogenetically linked to the only other available ZIKV sequence from Vietnam, sampled in 2016 (GenBank accession KX216633), which suggests that the two 2013 viruses characterized in this study are unlikely to represent ancestral viruses to those causing more recent infections within Vietnam.

CONCLUSIONS

These data suggest that CHIKV and ZIKV have recently circulated, or been imported, into southern Vietnam. The detection of Zika cases in 2013 is concerning for the potential consequences of local transmission for pregnant women. Greater surveillance is required to understand the trajectory of the incidence of these arboviral diseases in Vietnam, and underscores the importance of performing differential diagnosis in populations where multiple arboviral pathogens may cocirculate.

Received December 14, 2016. Accepted for publication January 21, 2017.

Published online April 10, 2017.

Acknowledgments: We are indebted to Tran Tan Thanh for his support with setting up the ZIKV PCR.

Financial support: This work was funded by the Wellcome Trust (United Kingdom).

Disclaimer: The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' addresses: Nguyen Than Ha Quyen, Duong Thi Hue Kien, Nguyen Minh Tuan, and Tran Thuy Vi, Dengue, Oxford University Clinical Research Unit, Wellcome Trust Major Overseas Programme, Hospital for Tropical Diseases, Ho Chi Minh, Vietnam, E-mails: quyennt@oucru.org, kiendth@oucru.org, tuannm@oucru.org, and vitt@oucru.org. Maia Rabaa and Le Van Tan, Enteric Infections, Oxford University Clinical Research Unit, Wellcome Trust Major Overseas Programme, Hospital for Tropical Diseases, Ho Chi Minh, Vietnam, E-mails: mrabaa@oucru.org and tanlv@oucru.org. Nguyen Thanh Hung, Department of Dengue Hemorrhagic Fever, Children's Hospital No. 1, Ho Chi Minh, Vietnam, E-mail: drthanhhung@gmail.com. Ha Manh Tuan, General Planning, Children's Hospital 2, Ho Chi Minh, Vietnam, E-mail: manhtuanzu@yahoo.com. Ta Van Tram, Paediatric Department, Tien Giang Provincial Hospital, My Tho, Vietnam, E-mail: tavantram@gmail.com. Nguyen Le Da Ha, Pediatrics, Dong Nai Children Hospital, Bien Hoa, Vietnam, E-mail: daha_nl@yahoo.com. Han Khoi Quang, Pediatrics, Binh Duong Hospital Thu Dau Mot, Vietnam, E-mail: hkquang@gmail.com. Nguyen Quoc Doanh, Pediatrics, Long An Hospital, Tan An, Vietnam, E-mail: nguyenguocdoanh1964@gmail.com. Nguyen Van Vinh Chau, Hospital for Tropical Diseases, Ho Chi Minh, Vietnam, E-mail: chaunvv@oucru.org. Bridget Wills, Dengue, Oxford University Clinical Research Unit, Wellcome Trust Major Overseas Programme, Hospital for Tropical Diseases, Ho Chi Minh, Vietnam, and Nuffield Department of Medicine, Centre for Tropical Medicine, University of Oxford, Oxford, United Kingdom, E-mail: bwills@oucru.org. Cameron P. Simmons, Dengue, Oxford University Clinical Research Unit, Wellcome Trust Major Overseas Programme, Hospital for Tropical Diseases, Ho Chi Minh, Vietnam, and Microbiology and Immunology, University of Melbourne, Parkville, Australia, E-mail: csimmons@oucru.org.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

REFERENCES

1. Grossi-Soyster EN, LaBeaud AD, 2017. Clinical aspects of Zika virus. *Curr Opin Pediatr* 29: 102–106.
2. Hapuarachchi HC, Bandara KB, Sumanadasa SD, Hapugoda MD, Lai YL, Lee KS, Tan LK, Lin RT, Ng LF, Bucht G, Abeyewickreme W, Ng LC, 2010. Re-emergence of chikungunya virus in southeast Asia: virological evidence from Sri Lanka and Singapore. *J Gen Virol* 91: 1067–1076.
3. Sam IC, Chan YF, Chan SY, Loong SK, Chin HK, Hooi PS, Ganeswari R, Abubakar S, 2009. Chikungunya virus of Asian and Central/East African genotypes in Malaysia. *J Clin Virol* 46: 180–183.
4. Tun MM, Thant KZ, Inoue S, Nabeshima T, Aoki K, Kyaw AK, Myint T, Tar T, Maung KT, Hayasaka D, Morita K, 2014. Detection of east/central/south African genotype of chikungunya virus in Myanmar, 2010. *Emerg Infect Dis* 20: 1378–1381.
5. Duong V, Andries AC, Ngan C, Sok T, Richner B, Asgari-Jirhandeh N, Bjorge S, Huy R, Ly S, Laurent D, Hok B, Roces MC, Ong S, Char MC, Deubel V, Tarantola A, Buchy P, 2012. Reemergence of chikungunya virus in Cambodia. *Emerg Infect Dis* 18: 2066–2069.
6. Musso D, Cao-Lormeau VM, Gubler DJ, 2015. Zika virus: following the path of dengue and chikungunya? *Lancet* 386: 243–244.
7. Haddow AD, Schuh AJ, Yasuda CY, Kasper MR, Heang V, Huy R, Guzman H, Tesh RB, Weaver SC, 2012. Genetic characterization of Zika virus strains: geographic expansion of the Asian lineage. *PLoS Negl Trop Dis* 6: e1477.

8. Faye O, Freire CC, Iamarino A, Faye O, de Oliveira JV, Diallo M, Zanotto PM, Sall AA, 2014. Molecular evolution of Zika virus during its emergence in the 20th century. *PLoS Negl Trop Dis* 8: e2636.
9. Calvet G, Aguiar RS, Melo AS, Sampaio SA, de Filippis I, Fabri A, Araujo ES, de Sequeira PC, de Mendonca MC, de Oliveira L, Tschoeke DA, Schrago CG, Thompson FL, Brasil P, Dos Santos FB, Nogueira RM, Tanuri A, de Filippis AM, 2016. Detection and sequencing of Zika virus from amniotic fluid of fetuses with microcephaly in Brazil: a case study. *Lancet Infect Dis* 16: 653–660.
10. Cao-Lormeau VM, Blake A, Mons S, Lastere S, Roche C, Vanhomwegen J, Dub T, Baudouin L, Teissier A, Larre P, Vial AL, Decam C, Choumet V, Halstead SK, Willison HJ, Musset L, Manuguerra JC, Despres P, Fournier E, Mallet HP, Musso D, Fontanet A, Neil J, Ghawche F, 2016. Guillain-Barre syndrome outbreak associated with Zika virus infection in French Polynesia: a case-control study. *Lancet* 387: 1531–1539.
11. Ellison DW, Ladner JT, Buathong R, Alera MT, Wiley MR, Hermann L, Rutvisuttinunt W, Klungthong C, Chinnawirotpisan P, Manasatienkij W, Melendrez MC, Maljkovic Berry I, Thaisomboonsuk B, Ong-Ajchaowlerd P, Kaneechit W, Velasco JM, Tac-An IA, Villa D, Lago CB, Roque VG Jr, Plipat T, Nisalak A, Srikiatkachorn A, Fernandez S, Yoon IK, Haddow AD, Palacios GF, Jarman RG, Macareo LR, 2016. Complete genome sequences of Zika virus strains isolated from the blood of patients in Thailand in 2014 and the Philippines in 2012. *Genome Announc* 4: e00359-16.
12. Tuan NM, Nhan HT, Chau NV, Hung NT, Tuan HM, Tram TV, Ha Nle D, Loi P, Quang HK, Kien DT, Hubbard S, Chau TN, Wills B, Wolbers M, Simmons CP, 2015. Sensitivity and specificity of a novel classifier for the early diagnosis of dengue. *PLoS Negl Trop Dis* 9: e0003638.
13. Waggoner JJ, Pinsky BA, 2016. Zika virus: diagnostics for an emerging pandemic threat. *J Clin Microbiol* 54: 860–867.
14. Pastorino B, Bessaud M, Grandadam M, Murri S, Tolou HJ, Peyrefitte CN, 2005. Development of a TaqMan RT-PCR assay without RNA extraction step for the detection and quantification of African chikungunya viruses. *J Virol Methods* 124: 65–71.
15. Hue KD, Tuan TV, Thi HT, Bich CT, Anh HH, Wills BA, Simmons CP, 2011. Validation of an internally controlled one-step real-time multiplex RT-PCR assay for the detection and quantitation of dengue virus RNA in plasma. *J Virol Methods* 177: 168–173.
16. Tssetsarkin KA, Vanlandingham DL, McGee CE, Higgs S, 2007. A single mutation in chikungunya virus affects vector specificity and epidemic potential. *PLoS Pathog* 3: e201.