

Studying glycobiology at the single molecule level

Abirami Lakshminarayanan, Mylène Richard and Benjamin G. Davis*

Department of Chemistry, Chemistry Research Laboratory, University of Oxford, Oxford, U.K.

*e-mail: Ben.Davis@chem.ox.ac.uk

Abstract | Attempts to elucidate the roles of carbohydrate-associated structures in biology have led to the distinct field of glycobiology research. The focus of this field has been in understanding the evolution, biosynthesis and interactions of glycans, both individually and as components of larger bio-molecules. However, as most approaches for studying glycans (including mass spectrometry and various binding assays) use ensemble measurements, they lack the precision required to uncover the roles of glycoconjugates, which are often heterogeneous, in biomolecular processes. Single molecule techniques can examine individual events within challenging mixtures and they are beginning to be applied to glycobiology. For example, single molecule force spectroscopy (SMFS) by atomic force microscopy (AFM) has enabled the molecular interactions of sugars to be studied, single molecule fluorescence microscopy and spectroscopy have led to insight into the role of sugars in biological processes and nanopores have revealed key interactions between polysaccharides and their transporters. Thus, single molecule technology is becoming a valuable tool in glycoscience.

[H1] Introduction

Glycobiology explores a variety of biomolecules, including conjugates containing carbohydrate moieties and those that are discrete mono-, oligo- and polysaccharides. The roles of such sugars and glycoconjugates (that is, carbohydrates that are covalently linked to another chemical moiety) in cell biology are varied. For example, these moieties are involved in recognition¹, structure, modulation and signalling mechanisms^{2,3} (**Fig. 1**). These functions highlight glycosylation as one of the most important and diverse post-translation modifications that is at the heart of fine-tuning biological mechanisms. Indeed, there has been widespread interest among researchers in understanding and modulating glycobiology. [Au: OK?] However, there has been a lack of progress in the field owing not only to its intrinsic complexity and diversity but also, as we discuss here, due to a lack of requisite tools and techniques to study glycoconjugates^{2,4,5}. [Au: OK?]

The structural heterogeneity of glycoconjugates, which arises from branching, interlinking and different anomeric stereochemistries, has made it difficult to de-lineate individual structures and their roles in cellular pathways^{5,6}. The study of glycobiology would therefore benefit from approaches that allow: the elucidation of precise structures within dynamic mixtures, and the study of single events (that are mediated by broad populations of biomolecules) within diverse mixed interactions⁷⁻⁹. Single molecule methods can, in principle, address these goals by providing discrete analysis of molecules, and the events in which they are involved, in the context of the biological spectrum; this motivates the use of such methods in the study of glycans, where dynamic mixtures dominate.

In this Review we suggest that, although single molecule methods are currently used rarely in glycoscience, they might enable the discrete structural characterization of glycans and the elucidation of the mechanisms underlying sugar-mediated events. We choose key examples, from within and outside glycobiology, to illustrate the power of single molecule methods and acknowledge that these methods cannot address all of the issues in the field of glycobiology. Rather than collating all of the methods that can be used to study glycobiology, this Review highlights single molecule strategies that do, or have the potential to, allow a discrete understanding of the structure, function and dynamics of glycans and it represents our personal perspective on future directions for advancing the field.

[H1] Major challenges in glycobiology

[H2] *Structure determination of glycoconjugates.* Seemingly minor structural details, such as the anomeric stereochemistry of, or type of glycosidic linkage in, a glycoconjugate can influence the biochemical processes in which glycoconjugates are involved in to varying degrees. Thus, elucidating the exact structural details of glycoconjugates is an important starting point when studying the mechanisms underlying their biological roles, and for developing system-specific models or analogues that test glycoconjugate-driven hypotheses. Advances in carbohydrate nuclear magnetic resonance (NMR) spectroscopy^{10,11} and mass spectrometry¹²⁻¹⁴ have enabled formidable progress in this direction, yet neither technique is routinely used by all in the biological community. Challenges remain in readily determining the structure and position of glycans in glycoproteins at a sensitivity that allows the proteome-wide survey of glycoproteins (that is, of the glycoproteome) in primary tissue samples.^{15,16} Strategies that can distinguish between the isobaric species that are ubiquitous in glycoscience, but more rarely plague proteomics, show promise in enabling us to elucidate the full ‘primary sequence’ of glycoproteins; such sequence data was once restricted to a report of the ratios of hexose (Hex) to N-acetylhexosamine (HexNAc) to higher sugar types.^{13,17-20} Additionally, methods for determining the ‘shape’ of glycoproteins – that is, for determining secondary and higher order structural motifs²¹⁻²⁵ – are valuable given the relative rarity of glycoconjugates in current structural databases. Thus ‘sugar sequencing’ and comprehensive structure–function correlations still remain incompletely solved issues but promising methods are now touching on useful strategies.

[H2] *Understanding glycan dynamics and glycoconjugate modulation.* Glycoconjugates not only provide a stable matrix for biomolecules, but they are also involved in transferring information during extracellular communication and through signalling networks⁶. Thus, understanding the population dynamics of glycans in such glycoconjugates, and how these dynamics change in response to various stimuli, will enable a deeper understanding of the mechanisms underscoring information transfer and may inform medical diagnostics in the clinic.

Understanding the population dynamics of glycans requires advanced methods for synthesising and assembling glycans that can be used as key standards and probes, as well as methods for the ready labelling, sample enrichment and rapid and repeatable detection of metabolite glycans in minute quantities within mixtures; many current, widely-used methods

may not achieve the levels of efficiency or sensitivity needed. Therefore, assays and technologies that can determine, *in situ*, the functional moiety of the glycoconjugate, if it is present or partitionable, are needed. [Au: Sentence OK now?] Recent advances of relevance include the continuing development of automated glycan synthesis²⁶, the introduction of sensitive labels (for example, fluorophores) into molecules [Au: OK?] by suitably selective reactions²⁷, chemoenzymatic strategies²⁸ for altering cellular glycans to study their function²⁹ as well as the increased efficiency of, and alternative strategies in, glycosylation methods.³⁰ These efforts should help us understand the roles of sugars in disease and dysfunction³¹⁻³³, perhaps by correlating sugar biomarkers with pathology. Indeed, fluorescent imaging and the use of fluorescence resonance energy transfer (FRET), even using methods that have not been optimized for the detection of glycoconjugates, [Au: OK?] have provided valuable insight into processes, including the clustering of glycolipids on the cell surface^{34,35}, that inform us about glycan function in certain settings. However, despite bioinformatic analyses to identify the most biologically relevant sugars³⁶, the high number of sugar permutations is a barrier that needs to be acknowledged^{36,37} [Au: Your edit deleted ref 37; reinstate here? Or please cite it elsewhere, renumbering the references if needed.] when considering the synthesis and sequencing of such sugar biomarkers.

[H2] Understanding multivalency. The role of multivalency is a primary consideration in glycobiology. Carbohydrate–protein interactions are key to many biological processes but involve often weak binding affinities, a seeming contradiction at first inspection. In the limited number of systems that have been studied, nature appears to use differing modes of multivalency. To compensate for their weak binding affinities, multiple copies of ligands and receptors are suggested to cooperate to bind sequentially or simultaneously, providing enhanced binding avidities that convert small affinity interactions to physiologically relevant levels. This phenomenon has been termed the ‘cluster glycoside effect’^{38,39} and is thought to be integral to the study of many carbohydrate–mediated interactions in glycobiology. Studying this ‘effect’ requires a continued and improved understanding of models and modes of multivalency,⁴⁰ as well as the development of appropriate techniques.

Glycan arrays have been developed to qualitatively and semi-quantitatively probe interactions that involve carbohydrates in various multivalent contexts, including as biomarkers for various diseases^{41,42}. Along with surface plasmon resonance (SPR)⁴³, isothermal titration calorimetry (ITC)⁴⁴, and mass spectrometry^{12,13}, glycan arrays are now

semi-routine tools for more detailed but situation-dependent quantification of host–ligand multivalent interactions from which seminal bulk models of multivalency have emerged.^{39,44-}

⁴⁶ Further, techniques for studying multivalent interactions and delineating the pathways, stochasticity and collective modes of such binding will improve our understanding of many bioprocesses. These bioprocesses include the attachment of pathogenic bacteria to receptors and signalling induced by small molecules and cytokines. We feel that the development of such techniques for studying multivalent interactions could prove invaluable for analysing suggested cluster glycoside effects on a case-by-case basis rather than in a generalized manner. [Au:OK?] Indeed, it is unclear whether multivalent and/or cooperative and anticooperative effects are routinely or occasionally present in glycoscience and more regular observation of relevant molecular events will allow us to understand the ‘if, how and when’ of the multivalency of sugars.

[H1] Studying single molecule glycobiology

The intricacies of glycobiology have necessitated the development of various bulk techniques that have become indispensable for the study and characterization of the structures and interactions of carbohydrates; some of these techniques are relevant to single-molecule methods for studying glycobiology and we will highlight these here. Conventional bulk techniques (often also referred to as ensemble techniques) provide the deterministic outcome of a given process, for example, through a comparison between the bound and unbound states of an analyte. However, two potentially confounding aspects of ensembles in glycobiology, where single-molecule methods could prove particularly useful, are in heterogeneity and collective effects.

Furthermore, kinetics obtained in ensembles can be complex and difficult to analyse.^{46,47} Single molecule studies could reveal information regarding the underlying dynamics of a system and the precise roles of individual components within mixtures to complement ensemble studies, which themselves may simply show the existence of a global interaction. Moreover, spatial and temporal resolution provides unique mechanistic information or different facets of a molecular mechanism that give a common outcome, which cannot be resolved by ensemble analysis (see **Fig. 2**).⁴⁸ These facets may provide further biological insight to broad ‘effects’.

The elucidation of [Au:OK?] binding mechanisms by new methods could be important for the understanding, exploitation and application of resulting technologies. For example, knowledge of exact carbohydrate–lectin binding modes creates the potential for the design of

anti-infectives⁴⁹ that block host–pathogen interactions or may identify an unexpectedly potent antigen that can be used as a carbohydrate vaccine immunogen.⁵⁰ [Au: OK?]

We describe examples that demonstrate strategic and specific potential advances that could be enabled or developed by single-molecule-based technologies in glycobiology. In so doing, we have highlighted both techniques currently used to study carbohydrates in ensembles for which single-molecule counterparts could be valuably developed or techniques that are being used at a single-molecule level in other realms that could be repurposed for sugar-specific interrogation (see **Table 1**). In this way we hope to chart a possible course for development of the field.

[H2] Force spectroscopy. Study of single molecule force spectroscopy (SMFS) by atomic force microscopy (AFM) is a widely and actively used technique for single molecule studies in Biology. By estimating the strength of certain carbohydrate-based interactions, SMFS has allowed researchers to gain insight into the functional roles [Au:OK?] of these interactions^{51,52}. For example, AFM was used to assess differences in carbohydrate binding on the surface of ‘cancerous’ cell lines compared with ‘normal’ cell lines.⁵³ Glycoconjugates on the cell surface of HeLa cells (cancerous) [Au:OK?] and Madin-Darby canine kidney (MDCK) cells (normal) were approached with lectin-coated AFM tips, which were then retracted according to typical SMFS protocols (**Fig. 3a**) to record lectin–glycoconjugate interactions via force-distance cycles. The lectins wheat germ agglutinin (WGA, which is presumed to be selective for glucose (Glc) and *N*-acetylglucosamine (GlcNAc)) and *Polygonatum multiflorum* agglutinin (PMA, which is presumed to be selective for mannose (Man)) were used to probe cell-surface interaction strengths in a glycan-dependent manner. Although the specificity of these lectins for glycans is likely to be more complex than presumed in this study, the data obtained illustrate what AFM can tell us about carbohydrate-based interactions. Specifically, these studies suggested that the energy of activation for sugar–lectin dissociation is lower in cancer cells than in normal cells, which correlates with some established data showing that cancer cells are less adherent than normal cells.⁵³ AFM-based molecular recognition has also been used to study the binding kinetics of the lectin domain of the toxin ricin to galactose moieties on HeLa cells. Using approach–withdrawal cycles [Au: Does this mean that all ‘force-distance cycles’ in the text should be changed to approach-withdrawal?] with a ricin-coated tip gave force-distance measurements that

allowed extrapolation to an *in situ* dissociation constant of 2.2×10^{-3} M.⁵⁴ The use of single molecule techniques to interrogate the binding of glycans to lectins on live cells, as demonstrated by these studies, illustrates the potential contextual advantages of these techniques over bulk studies in obtaining physiologically-relevant information.

Certain glycoconjugates on the surface of pathogens can act as immunogens that trigger host responses;⁵⁵ thus the construction of synthetic glycan mimics to develop carbohydrate-based vaccines is an active area of research.^{56,57} One prerequisite for their systematic, iterative development is the identification of the antigenic moieties that are bound by antibodies raised in this way and SFMS (via AFM) might identify correlations between immunity and the molecular properties of glycoconjugate immunogens. For example, data from bulk, enzyme-linked immunosorbent assays (ELISA) suggest that multi-antennary oligosaccharides, such as $\text{Man}_9(\text{GlcNAc})_2$, are central to the immune responses associated with the natural, neutralizing anti-HIV-1 antibody 2G12.⁵⁸ However, synthetic conjugates containing such or related sugars have failed to stimulate the required level of immune response, prompting further investigation.⁵⁹ When the carbohydrate specificity of 2G12 was probed by SMFS⁶⁰ using a panel of oligosaccharide fragments motifs (**Fig. 3b**), data revealed that 2G12 had lower lifetimes of interaction with segments of $\text{Man}_9(\text{GlcNAc})_2$ at the single molecule level but potentially useful lifetimes with **[Au:OK?]** gp120 glycoprotein displaying multiple glycans. These data highlight the potential value of analyzing the kinetics of interactions, an aspect that is less studied in vaccinology.^{61,62} They also emphasize the need to determine multiple structural and chemical properties in a system as well as the utility of single molecule techniques in highlighting which parameters in a system have more functional influence.

Carbohydrate-containing antigens in glycopeptidic fragments may also undergo self-assembly into complex, ordered arrangements, a concept that has also been studied by single-molecule AFM. For example, force-distance measurements of antigens T ($\text{Gal}\beta(1-3)\text{GalNAc}\alpha\text{Thr/Ser}$) and Tn ($\text{GalNAc}\alpha\text{Thr/Ser}$) derived from porcine submaxillary mucin revealed stable antigen self-interactions with lifetimes ranging from 0.5–4.5 s; these interactions had not previously been observed in such detail.⁶³ These antigens are implicated in cancerous tissues and the authors of the study proposed that the variation of interaction lifetimes observed for these truncated antigens might contribute to aberrant cell-surface signalling.

Protein- or peptide-mediated sugar interactions dominate conventional analyses in glycobiology; direct carbohydrate-to-carbohydrate interactions have been less studied.

However, the sensitivity of single molecule methods, and in particular of AFM, has enabled quantitative analyses of carbohydrate-to-carbohydrate interactions. As an example, the mechanism of carbohydrate-mediated cell adhesion has been studied in marine sponges, providing evolutionary insights into early metazoans that may have broader implications for the development of higher, multicellular organisms.^{64,65} Specifically, ‘self’-interactions amongst sulfated glycosaminoglycans (GAGs) that are evolutionarily preserved in metazoans were probed by force cycles using SFMS, revealing that the presence of chelated calcium and sulfation on these GAGs is vital for their self-interaction and, thus, presumably for cell adhesion.

As well as providing useful information on the binding forces of cell-surface glycans in different contexts, AFM has been used to ‘map’ the localization of cell-surface glycans at the single molecule level (**Fig. 4**). For example, AFM tips functionalized with anti-capsular polysaccharide (CPS) antibodies were used to probe both the localization and density of the outer glycans of CPS and group B carbohydrate (GBC) in wild type and mutant *Streptococcus agalactiae*⁶⁶. AFM tips functionalized with WGA were then used to probe peptidoglycan underlying these outer glycans, revealing that the ‘adhesive probabilities’ of wild-type *S. agalactiae* and mutant *S. agalactiae* that do not display GBC were weak (1–3%), whereas those of mutant *S. agalactiae* that do not display CPS were ~30%. These data highlight and quantitate the vital role that the CPS plays in protecting the peptidoglycan layer of pathogenic bacteria from environmental exposure. A similar study on yeast cells⁶⁷ used anti- β -mannan antibodies on an AFM tip [Au:OK?] to map the localization of α -mannans versus β -mannans. 1,2- β -mannans were markedly detected on pathogenic yeast strains *Candida albicans* and *Candida glabrata* but barely detectable on a non-pathogenic strain, *Saccharomyces cerevisiae*. These data are in line with the fact that 1,2- β -mannans are known to interact with toll-like receptors to induce an inflammatory immune response against pathogenic yeast.

Although it is rare for AFM to provide mechanistic insight, one example of where it did involves family 3 carbohydrate-binding modules (CBM3) and their ‘helper’ role in the enzymatic-processing of cellulose. Combined use of AFM imaging of CBM3-functionalized gold nanoparticles and SFMS using a CBM3-functionalized tip allowed the time-dependent monitoring of the alignment of CBM3 along a single cellulose fibril as the [Au:OK?] assessment of corresponding affinities.⁶⁸⁻⁷⁰ Using different concentrations of CBM3 and a fixed concentration of cellulose a minimum initial concentration of 5.1×10^{-7} M CBM3 was

required for CBM3 to bind to the cellulose after ~5h.⁶⁹ This elegant study enabled the determination of binding kinetics in this semi-heterogeneous system, which is in contrast to bulk studies in which the heterogeneous distribution of ‘free’ molecules limited the understanding of real-time processes.⁷¹ This example highlights the unique insights into kinetics that certain single molecule methods may provide under atypical (for example, interfacial) conditions for substrates that have been difficult to assess, [Au:OK?] such as insoluble or partially-soluble biopolymers (for example, polysaccharides).

Intriguingly, it has been proposed that SMFS can be used as an analytical tool to identify different sugar isomers⁷². A comparison of β -galactan (a polysaccharide of β -1,4-linked D-galactosyl units) and amylose (made up of α -1,4-linked D-glucosyl) moieties showed that this can be achieved by assessing conformation-associated changes in these sugars. Force-extension curves obtained from measurements using, as substrate, [Au:OK?] a single polysaccharide molecule showed a unique ‘fingerprint’ plateau at 600 pN for β -galactans. A comparison of the force spectrograms of β -galactan and amylose revealed that the energy required to stretch a single β -galactosyl residue in the context of β -galactan was 1.1 kcal/mol higher than the energy required to stretch a single α -glucosyl residue in amylose. This suggests that force spectroscopy could potentially discriminate between sugar isomers when cumulative differences in configuration influence chain conformation; viewed in this way, the differences found between β -galactan and amylose occurs through the ‘interchange’ of axial and equatorial bonds at C1 and C4. It will be interesting to see if, in addition to being successfully applied to this example that involves homogeneous repeating units, force spectroscopy could be applied to more heterogeneous polysaccharides (for example, those with alternating subunits) or whether the resulting pulling energy spectra would be insufficiently distinct.

In this subsection we have focused on the use of AFM as a single-molecule method. Given the important influence that trap-based tweezing methods (using optical and magnetic traps, for example) have had on force measurement in other biological fields⁷³ we foresee a wider use of these methods in glycobiology, including through the use of glycan-coated beads.⁷⁴

[H2] Single molecule field-effect transistors. The topology of existing constructs can be altered to provide powerful new systems for use in single molecule glycobiology. Lysozyme-tethered single-wall carbon nanotubes can be used as intimate field-effect transistors to study

the T4 lysozyme-mediated processing of cell wall-derived linear and cross-linked peptidoglycans⁷⁵. By monitoring dynamic electronic signals that correspond to enzyme activity, hydrolytic degradation events were observed. For both linear and cross-linked peptidoglycans, switching rates in signals of 20–50 s⁻¹ were observed that were correlated with substrate processing. Notably, these observations suggested that the enzyme could degrade cross-linked peptidoglycan, albeit at a rate ~15% slower than the rate at which it degraded linear peptidoglycan. Thus, despite the highly heterogeneous structures found in these substrates, single-molecule studies have allowed a more precise understanding of the dynamics and function in what might be considered to be a near-native model of degradation.

[H2] Single molecule fluorescence. Some 20 years ago, fluorescence correlation spectroscopy (FCS) was used to study binding of the lectin concanavalin A (ConA) to a fluorescently labelled β -maltopyranoside. Dilute solutions of this ligand (1 μ M) allowed unbound lectins to be distinguished from carbohydrate-bound lectins⁷⁶, providing valuable proof-of-concept that FCS can be used to study glycobiology. Since then, several examples have given a sense of where single molecule fluorescence is most valuable in glycobiology. For example, CBM–crystalline cellulose interactions have been examined with a method termed defocused orientation and position imaging (DOPI), a technique based on anisotropy of dipole radiation. This was used to study the orientation of CBMs fused to GFP with respect to cellulose fibrils during binding interactions. The images obtained (under TIRF illumination with slight defocusing) exploit the clear transition dipole of GFP allowing a definitive orientation of CBMs along cellulose fibrils to be determined; these proved consistent with current models of CBM molecular recognition processes⁷⁷. Again, in this way, single molecule methods can explore important glycan–protein interfaces (for example, crystalline cellulose) that are less tractable to bulk methods.

As well as the direct monitoring of sugar–protein ligand interactions, the broader (for example, cellular) regulation and function of sugar-processing proteins may be monitored through using fusion fluorescent proteins. For example, single molecule fluorescence spectroscopy allows real-time monitoring of the expression⁷⁸ of the *E. coli lacY* gene that encodes lactose permease (LacY) — a symporter that transports β -galactosides such as lactose into the cell. The *lacY* gene is part of the *lac* operon. Protein production in bacterial cells under non-induced conditions (that is, when a repressor is bound to mRNA) was shown to be a stochastic event that occurs in ‘bursts’⁷⁹. The *lac* operon in *E. coli* is repressed by LacI in the absence of the inducer lactose (**Fig. 5A**). The presence of inducer, however, dislodges

the repressor and triggers protein production. In this way, using LacY fused to a fluorescent protein, gene regulation can be studied at a single molecule and [Au: OK?] single cell level⁸⁰; in the presence of low levels of inducer, cell phenotype can be switched due to the partial dissociation, slow rebinding, or complete dissociation of the repressor from mRNA. [Au:OK?] Specifically, slow rebinding and partial dissociation of the repressor resulted in only basal levels of LacY production, whereas the complete dissociation of the repressor resulted in a large burst of LacY production and a switch in phenotype. As well as monitoring its production, the use of other single molecule fluorescent methods allow the structural dynamics of LacY to be determined; indeed, single-molecule FRET provided a real-time method for monitoring sugar symport in *E. coli*.⁸¹ X-ray crystallography had provided structures that form the basis of an ‘alternating access’ model for LacY-mediated sugar transport^{82,83}. This model proposes that sugar binds at the centre of LacY from either cytoplasmic or periplasmic faces, followed by the sequential opening and closing of the faces to allow symport. FRET-based single molecule fluorescence enabled monitoring of precise distance changes between donor–acceptor fluorophore pairs upon binding of the sugar — specifically a [Au:OK?] galactopyranoside ligand — and hence validated this model.⁸¹

As well as *in cellulo* examination of their protein partners, fluorescence also allows single molecule tracking of glycans themselves. For example, the interaction of ‘red-labelled’ gangliosides with ‘green-labelled’ CD59 domains (used as analogues of glycosylphosphatidylinositol (GPI) anchors) has been studied. Fluorescent single-molecule tracking, using total internal reflection fluorescence (TIRF) spectroscopy that allowed precise localization of fluorophores, revealed that gangliosides and CD59 domains dynamically co-localize with interaction lifetimes of up to 10 ms. Interestingly, cholesterol supplementation enhanced the formation of CD59–ganglioside raft domains, and, although interactions remained extremely dynamic under these conditions, this nonetheless increased the association lifetime of gangliosides and CD59 domains to 40 ms (**Fig. 5b**)⁸⁴.

Cell-surface sugars can also be manipulated to engineer cell-surface morphology (**Fig. 6**). [Au:OK? Otherwise Fig. 6 not cited given your deletion of it below.] For example, techniques can be used to artificially induce network formation between conjugated glycans at cell surfaces^{85 86 87}. [Au: As we do not have software for renumbering references, please could we ask you to insert this reference into your list [Spiegel, S. *et al.* Direct visualization of redistribution and capping of fluorescent gangliosides on lymphocytes. *J Cell Biol.* 99 1575–1581 (1984)]? It will need to become reference 85 and the current reference 85 will become reference 86, and so on.] In one example,⁸⁷ single-molecule

tracking via epifluorescence monitored the response of one such network to changes in tuneable parameters, namely valency of the biotin-binding protein streptavidin (through pre-blocking) and concentrations of streptavidin (Figure 6), revealing that the mobility of the cell membrane could be tuned from one with physiological mobility to one that was stiff and almost 'frozen'. As might be expected, this enhanced artificial network formation decreased membrane mobility and permeability in membrane-dependent processes, such as endocytosis. Controlling the mobility of cell surface glycans, monitored by single molecule fluorescence, might prove valuable in applications that could exploit the use of such artificial cytoskeletal structures.

[H2] Super-resolution microscopy. Several methods in single molecule fluorescence require some level of cellular 'fixing', generating samples that may not represent those in nature. However, the use of stochastic optical reconstruction microscopy (STORM)⁸⁸ can avoid this need by exploiting photoswitchable fluorophores which, following several cycles of activation, enable the localization of biomolecules on cell-surfaces with 'super-resolution'. For example, STORM imaging using Cy5-conjugated lectin allowed the identification of corresponding glycan ligands, which were presumably D-Man-containing based on lectin selectivity, on cell surfaces. Glycan clusters 105–850 nm in diameter were identified, suggesting that these may sit on glycoproteins within membranes that form microdomains. Expansion of this application of STORM through indirect visualization allowed the semi-systematic determination of other carbohydrate types found in clustered states on cell membranes and corresponding membrane components⁸⁹; for example, epidermal growth factor (EGFR) co-localized with GalNAc clusters. The authors propose that these clusters have a more global role in enhancing protein–glycoprotein interactions through the cluster multivalency that is created. It should be noted that the existence of such glycan clusters or microdomains has also been challenged as the direct labelling of unnatural cell-surface glycans with suitable fluorophores (using triazole-forming chemistry) followed by STORM, revealed that they have a homogeneous, non-clustered distribution.⁹⁰ Such method-dependent differences in findings highlight the need to analyse underlying assumptions and possible attributable artefacts. Amongst these assumptions it should be considered that, whereas super-resolution microscopy has a resolution of ~20 nm, which may be suitable for visualizing single protein molecules, this resolution cannot visualize single glycan residues.

The simultaneous fluorescent labelling in live cells of *N*-linked sialic acid and *O*-linked glycans, using alkyne-containing mannose and azido-containing galactosamine, respectively, as metabolic precursors, has allowed their diffusion rates to be estimated by STORM.⁹¹ Glycans on the surface of the mouse mammary cancer cell line PyMT diffused at rates 2–3 times faster than those on the surface of its highly metastatic variant cell line, Met-1. The diffusion rate was hypothesized to reflect the metastatic potential of these cell lines, suggesting that the imaging of single molecule glycans might be useful for studying the morphology, and even potential phenotypes, of cells.

The related super-resolution method photo-activated localization microscopy (PALM) has been used to determine the influence of surface substrate morphologies upon enzyme activity.⁹² Cellulose targeting was used as a model system and PALM was used since AFM-based techniques cannot resolve individual proteins from within a dense, heterogeneous surface. PALM revealed that crystalline-specific carbohydrate-binding modules (CBM) were added to cellulose, they tended to cluster on cellulose ‘ridges’; by contrast amorphous-specific CBMs were found to be uniformly present across the surface of cellulose.⁹² These observations suggest that a combination of CBM types will be more effective when used in aiding cellulose degradation.

[H2] Nanopores. A preliminary system based on heptameric α -haemolysin pores showed the potential of ‘nanopores’ in single-molecule glycobiology (**Fig. 7**). Specifically, the binding kinetics of the *Bauhinia purpurea* (BP)-lectin were examined in a modified variant of α -haemolysin displaying seven glycans.⁹³ By monitoring changes in ionic current flow using single-channel recording, pore events with two lifetimes (suggestive of two binding modes) were detected. Although the presence of multiple glycans on the pore prevented the identification of distinct binding modes from many possible permutations, such methods illustrate that strategies where the ‘sensor’ is also a biomolecular substrate could prove useful for elucidating the mechanism of glycoprotein–lectin interactions.

In this example, the α -haemolysin pore was used as a sensitive platform on which to mount glycans, but it had no inherent role in glycan binding or processing. By contrast, the sugar-transporting pore protein Wza from *E. coli* is involved in capsular polysaccharide (CPS) export. Its ability to record the passage of sugars through single channels has been explored (**Fig. 7**). Using synthetic fragments of CPS and single molecule methods employing droplet bilayers, sugar export was partially recapitulated and an early sense of the inherent

selectivity of this natural sugar pore — even when Wza was not associated with its usual transport helper proteins — was achieved.⁹⁴ The results suggest that, even for small oligomers, coordinated aid may be needed from underlying proteins for sugar transport in such systems. This Wza system allowed a library of putative Wza blockers to be tested and identified an inhibitor of CPS transport through screening at a single molecule level; some inhibitors of CPS transport had dissociation constant (K_d) values < 50 nM.^{94,95} As inhibiting CPS transport prevents the formation of the bacterial capsule, these inhibitors may synergise with antibiotic and anti-bacterial vaccination strategies.^{95,96} Chitoporins, which are involved in sugar transport in marine bacteria, have also been tested in single channel recording. The studies suggest chitoporin monomers have multiple ‘traps’ or binding sites for sugars; stochastic binding of one sugar may cooperatively assist the sequestration of multiple sugar units, perhaps enabling the marine bacteria to achieve high sugar translocation even in extreme environments⁹⁷. It is notable that such nanopore-based ‘traps’ have not yet, to our knowledge, been complemented in glycobiology by other trap types.^{73,98}

[H2] Recognition tunnelling. An approach enabling the identification of sugar anomers and isomers at the single molecule level by ‘recognition tunnelling’ has been identified⁹⁹ In this process, fluctuations in current observed when a single-molecule is trapped in the space between two electrodes that have been functionalized with appropriate recognition molecules can distinguish between anomers or epimers. 4(5)-(2-mercaptoethyl)-1*H*-imidazole-2-carboxamide (ICA), which can form multiple hydrogen bonds, has been proposed to be a suitable recognition molecule for analysing carbohydrates by recognition tunnelling. By analysing changes in the spikes in current produced by different epimers, the authors proposed that α -D-glucopyranoside could be discriminated from β -D-glucopyranoside (these are **[Au:OK?]** C-1 epimers / anomers). They also showed that various C-4 epimers could be distinguished — Glc, GlcNH₂, and GlcNAc from Gal, GalNH₂ and GalNAc, respectively, — and that the C-5 epimers D-glucuronic acid and L-iduronic acid could be discriminated, all with an accuracy > 98 %. Like SMFS in polysaccharides (see above), the applicability of recognition tunnelling to more complex sugar units remains to be explored.

[H1] Challenges and future directions

Although the widespread application of some single molecule techniques in studying glycobiology has been partly impeded by the need for bespoke methods, for example, nanopore technologies or certain fluorescence techniques, the development of commercially available equipment and reagents is increasingly reducing this barrier. That said, for techniques where methods have been more accessible, such as AFM, somewhat time-consuming data recording and result analyses still necessitate high levels of technical expertise (and/or dedication) and prescient experimental design. Moreover, artefacts arising from defective samples, non-specific interactions or the use of indirect probes (for example, labelled lectins) can lead to observations for which caveats need to be carefully considered and declared. Above all, even as mechanisms start to be explored *in vitro* with such some single-molecule methods, *in vivo* (and even *in cellulose*) studies remain limited or barely accessible – these will prove an interesting mid-term challenge.¹⁰⁰

Despite the ongoing challenges, from observing sugar transport through a single pore to testing the mode in which glycans are displayed on cell-surfaces or assessing their multivalency, single molecule interrogation is leading us towards an improved understanding of the more complex aspects of glycobiology. Where bulk studies provided useful possible models but insufficient detail, single molecule interrogation has, in certain cases, already allowed the validation of proposed mechanistic models. This progress may open new avenues in the development of inhibitors of sugar–protein interactions or more potent antigens for use in vaccination, as well as mapping and correlating distribution patterns of glycoconjugates as potentially important functional biomarkers. They also, in turn, raise questions and highlight the current limitations of using single molecule techniques to study glycobiology. For example: Can (or should) all of glycobiology be studied at the single molecule level and which single molecule methods can be developed and utilised to study a given problem without losing contextual or functional relevance? The use of an accessible technique will need to be balanced by its functional relevance.

Although single molecule analysis of sugar biology is still in its relative infancy, a wealth of opportunities exists. Perhaps most importantly (and more so than in other areas of biology) single-molecule methods could prove essential in enabling the proper mechanistic understanding of glycobiology. [Au: OK?] For example, these techniques might be essential for our full understanding of the functional roles of heterogeneity and multivalency in glycans. Thus, with careful experimental design, some molecular causes and drivers of sugar function in the biology of complex organisms may emerge. This, in turn, could enable precise and coherent kinetic analyses of carbohydrate-processing in context and hence allow

sufficient selectivity for more effective medical diagnostics and therapies. By providing specific information and unearthing hitherto unknown details in areas where bulk analyses of mixtures have limits, single molecule technology is a valuable tool in glycoscience.

References

- 1 Schnaar, R. L. Glycobiology simplified: diverse roles of glycan recognition in inflammation. *Journal of Leukocyte Biology* **99**, 825-838, doi:10.1189/jlb.3RI0116-021R (2016).
- 2 Dwek, R. A. Glycobiology: Toward understanding the function of sugars. *Chem Rev* **96**, 683-720, doi:10.1021/cr940283b (1996).
- 3 Varki, A. Biological roles of glycans. *Glycobiology* **27**, 3-49, doi:10.1093/glycob/cww086 (2017).
- 4 Seeberger, P. H. Chemical glycobiology: why now? *Nat Chem Biol* **5**, 368-372 (2009).
- 5 Imperiali, B. The chemistry–glycobiology frontier. *J Am Chem Soc* **134**, 17835-17839, doi:10.1021/ja309542q (2012).
- 6 Wang, L.-X. & Davis, B. G. Realizing the promise of chemical glycobiology. *Chem Sci* **4**, 3381-3394, doi:10.1039/C3SC50877C (2013).
- 7 Bertozzi, C. R., Kiessling & Laura, L. Chemical glycobiology. *Science* **291**, 2357 (2001).
- 8 Krishnamoorthy, L. & Mahal, L. K. Glycomic analysis: An array of technologies. *ACS Chem Biol* **4**, 715-732, doi:10.1021/cb900103n (2009).
- 9 Dube, D. H., Champasa, K. & Wang, B. Chemical tools to discover and target bacterial glycoproteins. *Chem Comm* **47**, 87-101, doi:10.1039/C0CC01557A (2011).
- 10 Grice, I. D. & Wilson, J. C. in *Microbial Glycobiology* (eds Patrick J. Brennan, Mark von Itzstein, & Anthony P. Moran) 233-252 (Academic Press, 2010).
- 11 Mulloy, B., Hart, G. W. & Stanley, P. in *Essentials of Glycobiology* (eds A. Varki *et al.*) 661–678 (Cold Spring Harbor Laboratory Press, 2009).
- 12 Gray, C. J. *et al.* Applications of ion mobility mass spectrometry for high throughput, high resolution glycan analysis. *Biochim Biophys Acta* **1860**, 1688-1709, doi:10.1016/j.bbagen.2016.02.003 (2016).
- 13 Hofmann, J., Hahm, H. S., Seeberger, P. H. & Pagel, K. Identification of carbohydrate anomers using ion mobility-mass spectrometry. *Nature* **526**, 241-244, doi:10.1038/nature15388 (2015).
- 14 Goldberg, D. *et al.* Automated N-Glycopeptide Identification Using a Combination of Single- and Tandem-MS. *Journal of Proteome Research* **6**, 3995-4005, doi:10.1021/pr070239f (2007).
- 15 Novotny, M. V., Alley, W. R. & Mann, B. F. Analytical glycobiology at high sensitivity: current approaches and directions. *Glycoconjugate Journal* **30**, 89-117, doi:10.1007/s10719-012-9444-8 (2013).
- 16 Gaunitz, S., Nagy, G., Pohl, N. L. B. & Novotny, M. V. Recent Advances in the Analysis of Complex Glycoproteins. *Analytical Chemistry* **89**, 389-413, doi:10.1021/acs.analchem.6b04343 (2017).
- 17 Both, P. *et al.* Discrimination of epimeric glycans and glycopeptides using IM-MS and its potential for carbohydrate sequencing. *Nat Chem* **6**, 65-74, doi:10.1038/nchem.1817 (2014).
- 18 Nagy, G. & Pohl, N. L. B. Monosaccharide Identification as a First Step toward de Novo Carbohydrate Sequencing: Mass Spectrometry Strategy for the Identification

- and Differentiation of Diastereomeric and Enantiomeric Pentose Isomers. *Analytical Chemistry* **87**, 4566-4571, doi:10.1021/acs.analchem.5b00760 (2015).
- 19 Nagy, G. & Pohl, N. L. B. Complete Hexose Isomer Identification with Mass Spectrometry. *Journal of The American Society for Mass Spectrometry* **26**, 677-685, doi:10.1007/s13361-014-1072-z (2015).
- 20 Schindler, B. *et al.* Anomeric memory of the glycosidic bond upon fragmentation and its consequences for carbohydrate sequencing. *Nature Communications* **8**, 973, doi:10.1038/s41467-017-01179-y (2017).
- 21 Rees, D. A. & Welsh, E. J. Secondary and Tertiary Structure of Polysaccharides in Solutions and Gels. *Angewandte Chemie International Edition in English* **16**, 214-224, doi:10.1002/anie.197702141 (1977).
- 22 Yuriev, E., Farrugia, W., Scott, A. M. & Ramsland, P. A. Three-dimensional structures of carbohydrate determinants of Lewis system antigens: Implications for effective antibody targeting of cancer. *Immunology And Cell Biology* **83**, 709, doi:10.1111/j.1440-1711.2005.01374.x (2005).
- 23 Yaffe, N. R., Almond, A. & Blanch, E. W. A New Route to Carbohydrate Secondary and Tertiary Structure Using Raman Spectroscopy and Raman Optical Activity. *Journal of the American Chemical Society* **132**, 10654-10655, doi:10.1021/ja104077n (2010).
- 24 Johannessen, C., Pendrill, R., Widmalm, G., Hecht, L. & Barron, L. D. Glycan Structure of a High-Mannose Glycoprotein from Raman Optical Activity. *Angewandte Chemie International Edition* **50**, 5349-5351, doi:10.1002/anie.201008258 (2011).
- 25 Barry, C. S. *et al.* 'Naked' and Hydrated Conformers of the Conserved Core Pentasaccharide of N-linked Glycoproteins and Its Building Blocks. *Journal of the American Chemical Society* **135**, 16895-16903, doi:10.1021/ja4056678 (2013).
- 26 Seeberger, P. H. The Logic of Automated Glycan Assembly. *Accounts of Chemical Research* **48**, 1450-1463, doi:10.1021/ar5004362 (2015).
- 27 Sletten, E. M. & Bertozzi, C. R. Bioorthogonal chemistry: Fishing for selectivity in a sea of functionality. *Angew Chem Int Ed* **48**, 6974-6998, doi:10.1002/anie.200900942 (2009).
- 28 Griffin, M. E. & Hsieh-Wilson, L. C. Synthetic Probes of Glycosaminoglycan Function. *Curr Opin Chem Biol* **17**, 1014-1022, doi:10.1016/j.cbpa.2013.09.015 (2013).
- 29 Griffin, Matthew E. & Hsieh-Wilson, Linda C. Glycan Engineering for Cell and Developmental Biology. *Cell Chemical Biology* **23**, 108-121, doi:10.1016/j.chembiol.2015.12.007.
- 30 Krasnova, L. & Wong, C.-H. Understanding the chemistry and biology of glycosylation with glycan synthesis. *Ann Rev Biochem* **85**, 599-630, doi:10.1146/annurev-biochem-060614-034420 (2016).
- 31 Berti, F. & Adamo, R. Recent mechanistic insights on glycoconjugate vaccines and future perspectives. *ACS Chem Biol* **8**, 1653-1663, doi:10.1021/cb400423g (2013).
- 32 Bernardi, A. *et al.* Multivalent glycoconjugates as anti-pathogenic agents. *Chem Soc Rev* **42**, 4709-4727, doi:10.1039/C2CS35408J (2013).
- 33 Astronomo, R. D. & Burton, D. R. Carbohydrate vaccines: developing sweet solutions to sticky situations? *Nat Rev Drug Discov* **9**, 308-324 (2010).
- 34 Lin, W., Du, Y., Zhu, Y. & Chen, X. A cis-membrane FRET-based method for protein-specific imaging of cell-surface glycans. *J Am Chem Soc* **136**, 679-687, doi:10.1021/ja410086d (2014).

- 35 Belardi, B., O'Donoghue, G. P., Smith, A. W., Groves, J. T. & Bertozzi, C. R. Investigating cell surface galectin-mediated cross-linking on glycoengineered cells. *J Am Chem Soc* **134**, 9549-9552, doi:10.1021/ja301694s (2012).
- 36 Adibekian, A. *et al.* Comparative bioinformatics analysis of the mammalian and bacterial glycomes. *Chemical Science* **2**, 337-344, doi:10.1039/C0SC00322K (2011).
- 37 Laine, R. A. Invited Commentary: A calculation of all possible oligosaccharide isomers both branched and linear yields 1.05×10^{12} structures for a reducing hexasaccharide: the Isomer Barrier to development of single-method saccharide sequencing or synthesis systems. *Glycobiology* **4**, 759-767, doi:10.1093/glycob/4.6.759 (1994).
- 38 Mammen, M., Choi, S.-K. & Whitesides, G. M. Polyvalent Interactions in Biological Systems: Implications for Design and Use of Multivalent Ligands and Inhibitors. *Angewandte Chemie International Edition* **37**, 2754-2794, doi:10.1002/(SICI)1521-3773(19981102)37:20<2754::AID-ANIE2754>3.0.CO;2-3 (1998).
- 39 Lundquist, J. J. & Toone, E. J. The Cluster Glycoside Effect. *Chemical Reviews* **102**, 555-578, doi:10.1021/cr000418f (2002).
- 40 Hunter, C. A. & Anderson, H. L. What is Cooperativity? *Angewandte Chemie International Edition* **48**, 7488-7499, doi:10.1002/anie.200902490 (2009).
- 41 Geissner, A. & Seeberger, P. H. Glycan arrays: From basic biochemical research to bioanalytical and biomedical applications. *Ann Rev Anal Chem* **9**, 223-247, doi:10.1146/annurev-anchem-071015-041641 (2016).
- 42 Laurent, N., Voglmeir, J. & Flitsch, S. L. Glycoarrays-tools for determining protein-carbohydrate interactions and glycoenzyme specificity. *Chemical Communications*, 4400-4412, doi:10.1039/B806983M (2008).
- 43 Fais, M. *et al.* Surface plasmon resonance imaging of glycoarrays identifies novel and unnatural carbohydrate-based ligands for potential ricin sensor development. *Chem Sci* **2**, 1952-1959, doi:10.1039/C1SC00120E (2011).
- 44 Dam, T. K. & Brewer, C. F. Thermodynamic studies of lectin-carbohydrate interactions by isothermal titration calorimetry. *Chem Rev* **102**, 387-430, doi:10.1021/cr000401x (2002).
- 45 Gestwicki, J. E., Cairo, C. W., Strong, L. E., Oetjen, K. A. & Kiessling, L. L. Influencing Receptor-Ligand Binding Mechanisms with Multivalent Ligand Architecture. *Journal of the American Chemical Society* **124**, 14922-14933, doi:10.1021/ja027184x (2002).
- 46 Kitov, P. I. & Bundle, D. R. On the nature of the multivalency effect: A thermodynamic model. *J Am Chem Soc* **125**, 16271-16284, doi:10.1021/ja038223n (2003).
- 47 Mulder, A., Huskens, J. & Reinhoudt, D. N. Multivalency in supramolecular chemistry and nanofabrication. *Org Biomol Chem* **2**, 3409-3424, doi:10.1039/B413971B (2004).
- 48 Tinoco, I. & Gonzalez, R. L. Biological mechanisms, one molecule at a time. *Genes Develop* **25**, 1205-1231, doi:10.1101/gad.2050011 (2011).
- 49 Sauer, M. M. *et al.* Catch-bond mechanism of the bacterial adhesin FimH. *Nature Communications* **7**, 10738, doi:10.1038/ncomms10738 (2016).
- 50 Pejchal, R. *et al.* A Potent and Broad Neutralizing Antibody Recognizes and Penetrates the HIV Glycan Shield. *Science* **334**, 1097-1103, doi:10.1126/science.1213256 (2011).
- 51 Scholl, Z. N., Li, Q. & Marszalek, P. E. Single molecule mechanical manipulation for studying biological properties of proteins, DNA, and sugars. *Wiley Interdisciplinary*

- Reviews: Nanomedicine and Nanobiotechnology* **6**, 211-229, doi:10.1002/wnan.1253 (2014).
- 52 Marszalek, P. E. & Dufrene, Y. F. Stretching single polysaccharides and proteins using atomic force microscopy. *Chemical Society Reviews* **41**, 3523-3534, doi:10.1039/C2CS15329G (2012).
- 53 Zhao, W., Cai, M., Xu, H., Jiang, J. & Wang, H. A single-molecule force spectroscopy study of the interactions between lectins and carbohydrates on cancer and normal cells. *Nanoscale* **5**, 3226-3229, doi:10.1039/C3NR00553D (2013).
- 54 Li, Y. *et al.* Molecular recognition force spectroscopy study of the specific lectin and carbohydrate interaction in a living cell. *Chem Phys Chem* **12**, 909-912, doi:10.1002/cphc.201001008 (2011).
- 55 Mazmanian, S. K. & Kasper, D. L. The love-hate relationship between bacterial polysaccharides and the host immune system. **6**, 849, doi:10.1038/nri1956 (2006).
- 56 Hecht, M.-L., Stallforth, P., Silva, D. V., Adibekian, A. & Seeberger, P. H. Recent advances in carbohydrate-based vaccines. *Current Opinion in Chemical Biology* **13**, 354-359, doi:10.1016/j.cbpa.2009.05.127 (2009).
- 57 Anish, C., Schumann, B., Pereira, Claney L. & Seeberger, Peter H. Chemical Biology Approaches to Designing Defined Carbohydrate Vaccines. *Chemistry & Biology* **21**, 38-50, doi:10.1016/j.chembiol.2014.01.002 (2014).
- 58 Joyce, J. G. *et al.* An oligosaccharide-based HIV-1 2G12 mimotope vaccine induces carbohydrate-specific antibodies that fail to neutralize HIV-1 virions. *Proc Nat Acad Sci USA* **105**, 15684-15689, doi:10.1073/pnas.0807837105 (2008).
- 59 Pashov, A., Garimalla, S., Monzavi-Karbassi, B. & Kieber-Emmons, T. Carbohydrate targets in HIV vaccine research: lessons from failures. *Immunotherapy* **1**, 777-794, doi:10.2217/imt.09.44 (2009).
- 60 Martinez, E., García, I., Marradi, M., Padro, D. & Penadés, S. Dissecting the carbohydrate specificity of the Anti-HIV-1 2G12 antibody by single-molecule force spectroscopy. *Langmuir* **28**, 17726-17732, doi:10.1021/la303484e (2012).
- 61 Hearty, S., Conroy, P. J., Ayyar, B. V., Byrne, B. & O’Kennedy, R. Surface plasmon resonance for vaccine design and efficacy studies: recent applications and future trends. *Expert Review of Vaccines* **9**, 645-664, doi:10.1586/erv.10.52 (2010).
- 62 Lynch, H. E., Stewart, S. M., Kepler, T. B., Sempowski, G. D. & Alam, S. M. Surface plasmon resonance measurements of plasma antibody avidity during primary and secondary responses to anthrax protective antigen. *Journal of Immunological Methods* **404**, 1-12, doi:10.1016/j.jim.2013.11.026 (2014).
- 63 Haugstad, K. E. *et al.* Enhanced self-association of mucins possessing the T and Tn carbohydrate cancer antigens at the single-molecule level. *Biomacromolecules* **13**, 1400-1409, doi:10.1021/bm300135h (2012).
- 64 Garcia-Manyes, S. *et al.* Proteoglycan Mechanics Studied by Single-molecule Force Spectroscopy of Allotypic Cell Adhesion Glycans. *Journal of Biological Chemistry* **281**, 5992-5999, doi:10.1074/jbc.M507878200 (2006).
- 65 Vilanova, E. *et al.* Carbohydrate-Carbohydrate Interactions Mediated by Sulfate Esters and Calcium Provide the Cell Adhesion Required for the Emergence of Early Metazoans. *Journal of Biological Chemistry* **291**, 9425-9437, doi:10.1074/jbc.M115.708958 (2016).
- 66 Beaussart, A. *et al.* Molecular mapping of the cell wall polysaccharides of the human pathogen *Streptococcus agalactiae*. *Nanoscale* **6**, 14820-14827, doi:10.1039/C4NR05280C (2014).

- 67 El-Kirat-Chatel, S. *et al.* Single-molecule analysis of the major glycopolymers of pathogenic and non-pathogenic yeast cells. *Nanoscale* **5**, 4855-4863, doi:10.1039/C3NR00813D (2013).
- 68 Zhang, M., Wang, B. & Xu, B. Measurements of single molecular affinity interactions between carbohydrate-binding modules and crystalline cellulose fibrils. *Physical Chemistry Chemical Physics* **15**, 6508-6515, doi:10.1039/C3CP51072G (2013).
- 69 Zhang, M., Wang, B. & Xu, B. Mapping single molecular binding kinetics of carbohydrate-binding module with crystalline cellulose by atomic force microscopy recognition imaging. *J Phys Chem B* **118**, 6714-6720, doi:10.1021/jp503185n (2014).
- 70 Zhang, M., Wu, S.-C., Zhou, W. & Xu, B. Imaging and Measuring Single-Molecule Interaction between a Carbohydrate-Binding Module and Natural Plant Cell Wall Cellulose. *The Journal of Physical Chemistry B* **116**, 9949-9956, doi:10.1021/jp304686q (2012).
- 71 Lee, S., Mandic, J. & Van Vliet, K. J. Chemomechanical mapping of ligand–receptor binding kinetics on cells. *Proc Nat Acad Sci USA* **104**, 9609-9614, doi:10.1073/pnas.0702668104 (2007).
- 72 Zhang, Q. & Marszalek, P. E. Identification of sugar isomers by single-molecule Force spectroscopy. *J Am Chem Soc* **128**, 5596-5597, doi:10.1021/ja058828e (2006).
- 73 Neuman, K. C. & Nagy, A. Single-molecule force spectroscopy: optical tweezers, magnetic tweezers and atomic force microscopy. *Nature Methods* **5**, 491, doi:10.1038/nmeth.1218 (2008).
- 74 Björnham, O., Bugaytsova, J., Borén, T. & Schedin, S. Dynamic force spectroscopy of the *Helicobacter pylori* BabA–Lewis b Binding. *Biophysical Chemistry* **143**, 102-105, doi:10.1016/j.bpc.2009.03.007 (2009).
- 75 Choi, Y. *et al.* Single-molecule dynamics of lysozyme processing distinguishes linear and cross-linked peptidoglycan substrates. *J Am Chem Soc* **134**, 2032-2035, doi:10.1021/ja211540z (2012).
- 76 La Clair, J. J. Selective Detection of the Carbohydrate-Bound State of Concanavalin A at the Single Molecule Level. *Journal of the American Chemical Society* **119**, 7676-7684, doi:10.1021/ja964366g (1997).
- 77 Dagel, D. J. *et al.* In situ imaging of single carbohydrate-binding modules on cellulose microfibrils. *J Phys Chem B* **115**, 635-641, doi:10.1021/jp109798p (2011).
- 78 Li, G.-W. & Xie, X. S. Central dogma at the single-molecule level in living cells. *Nature* **475**, 308-315 (2011).
- 79 Yu, J., Xiao, J., Ren, X., Lao, K. & Xie, X. S. Probing Gene Expression in Live Cells, One Protein Molecule at a Time. *Science* **311**, 1600-1603, doi:10.1126/science.1119623 (2006).
- 80 Choi, P. J., Cai, L., Frieda, K. & Xie, X. S. A stochastic single-molecule event triggers phenotype switching of a bacterial cell. *Science* **322**, 442-446, doi:10.1126/science.1161427 (2008).
- 81 Majumdar, D. S. *et al.* Single-molecule FRET reveals sugar-induced conformational dynamics in LacY. *Proc Nat Acad Sci USA* **104**, 12640-12645, doi:10.1073/pnas.0700969104 (2007).
- 82 Abramson, J. *et al.* The lactose permease of *Escherichia coli*: overall structure, the sugar-binding site and the alternating access model for transport. *FEBS Lett* **555**, 96-101, doi:10.1016/S0014-5793(03)01087-1 (2003).
- 83 Abramson, J. *et al.* Structure and Mechanism of the Lactose Permease of *Escherichia coli*. *Science* **301**, 610-615, doi:10.1126/science.1088196 (2003).

- 84 Komura, N. *et al.* Raft-based interactions of gangliosides with a GPI-anchored receptor. *Nat Chem Biol* **12**, 402-410, doi:10.1038/nchembio.2059 (2016).
- 85 Spiegel, S., Kassis, S., Wilchek, M. & Fishman, P. H. Direct visualization of redistribution and capping of fluorescent gangliosides on lymphocytes. *The Journal of Cell Biology* **99**, 1575-1581, doi:10.1083/jcb.99.5.1575 (1984).
- 86 Spiegel, S., Kassis, S., Wilchek, M. & Fishman, P. H. Direct visualization of redistribution and capping of fluorescent gangliosides on lymphocytes. *The Journal of Cell Biology* **99**, 1575 (1984).
- 87 Möckl, L., Lindhorst, T. K. & Bräuchle, C. Artificial Formation and Tuning of Glycoprotein Networks on Live Cell Membranes: A Single-Molecule Tracking Study. *ChemPhysChem* **17**, 829-835, doi:10.1002/cphc.201500809 (2016).
- 88 Zhao, W. *et al.* Studying the Nucleated Mammalian Cell Membrane by Single Molecule Approaches. *PLoS ONE* **9**, e91595, doi:10.1371/journal.pone.0091595 (2014).
- 89 Chen, J. *et al.* Systemic localization of seven major types of carbohydrates on cell membranes by dSTORM imaging. *Scientific Reports* **6**, 30247, doi:10.1038/srep30247 (2016).
- 90 Letschert, S. *et al.* Super-Resolution Imaging of Plasma Membrane Glycans. *Angewandte Chemie International Edition* **53**, 10921-10924, doi:10.1002/anie.201406045 (2014).
- 91 Jiang, H., English, B. P., Hazan, R. B., Wu, P. & Ovrn, B. Tracking Surface Glycans on Live Cancer Cells with Single-Molecule Sensitivity. *Angewandte Chemie International Edition* **54**, 1765-1769, doi:10.1002/anie.201407976 (2015).
- 92 Fox, J. M. *et al.* A single-molecule analysis reveals morphological targets for cellulase synergy. *Nat Chem Biol* **9**, 356-361, doi:10.1038/nchembio.1227 (2013).
- 93 Howorka, S., Nam, J., Bayley, H. & Kahne, D. Stochastic detection of monovalent and bivalent protein–ligand interactions. *Angew Chem Int Ed* **43**, 842-846, doi:10.1002/anie.200352614 (2004).
- 94 Kong, L., Almond, A., Bayley, H. & Davis, B. G. Chemical polyglycosylation and nanolitre detection enables single-molecule recapitulation of bacterial sugar export. *Nat Chem* **8**, 461-469, doi:10.1038/nchem.2487 (2016).
- 95 Kong, L. *et al.* Single-molecule interrogation of a bacterial sugar transporter allows the discovery of an extracellular inhibitor. *Nat Chem* **5**, 651-659, doi:10.1038/nchem.1695 (2013).
- 96 Kong, L. *et al.* An antibacterial vaccination strategy based on a glycoconjugate containing the core lipopolysaccharide tetrasaccharide Hep2Kdo2. *Nat Chem* **8**, 242-249, doi:10.1038/nchem.2432 (2016).
- 97 Suginta, W. & Smith, M. F. Single-Molecule Trapping Dynamics of Sugar-Uptake Channels in Marine Bacteria. *Physical Review Letters* **110**, 238102 (2013).
- 98 Wang, Q., Goldsmith, R. H., Jiang, Y., Bockenhauer, S. D. & Moerner, W. E. Probing Single Biomolecules in Solution Using the Anti-Brownian Electrokinetic (ABEL) Trap. *Accounts of Chemical Research* **45**, 1955-1964, doi:10.1021/ar200304t (2012).
- 99 Im, J. *et al.* Electronic single-molecule identification of carbohydrate isomers by recognition tunnelling. *Nature Communications* **7**, 13868, doi:10.1038/ncomms13868 (2016).
- 100 Dufrene, Y. F. *et al.* Five challenges to bringing single-molecule force spectroscopy into living cells. *Nat Meth* **8**, 123-127 (2011).
- 101 Varki, A. *et al.* Symbol Nomenclature for Graphical Representations of Glycans. *Glycobiology* **25**, 1323-1324, doi:10.1093/glycob/cwv091 (2015).

- 102 Li, Y. *et al.* Molecular recognition force spectroscopy of a specific lectin–carbohydrate interaction at single-molecule level. *Journal of Structural Biology* **176**, 46–51, doi:10.1016/j.jsb.2011.05.018 (2011).

Acknowledgements

We thank S. Faulkner, M. Bilyard, K. Wals for proof reading and useful comments.

Author contributions

A.L., M.R. and B.G.D. researched data for the article, made substantial contributions to discussions of the content, wrote the article and reviewed and/or edited the manuscript before submission.

Competing interests

BGD is named as an inventor on patents that describe new antibiotics based on some of the Wza science described in this Review. If licensed, these would afford BGD royalties in line with normal university practice.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Display items

Fig. 1| A snapshot of the glycobiology spectrum. Carbohydrate-containing structures are central to many biological processes. For example, carbohydrate-binding proteins such as galectins and C-type lectins have been implicated in cellular recognition, which may trigger downstream cell signalling pathways that are dependent on ATP. Glycan interactions may also mediate the interaction of cells with carbohydrate-binding toxins, as well as cell–cell interactions and cell–pathogen interactions. Finally, lipopolysaccharides are just one example of a major structural component of bacterial cell walls that may elicit immune responses via glycan-mediated recognition. The key of symbols here and in most other figures in this review follow the symbol nomenclature for graphical representations of glycans (SNFG) convention.¹⁰¹

Fig. 2| Comparison of ensemble measurements and single molecule measurements in a simulated non-sugar system. a| The calculated ensemble fluorescence resonance

energy transfer (FRET) versus time for a reaction of ribosome (R)-tRNA interaction that proceeds via three intermediates with different FRET values. The ensemble measurement fails to show such intermediates. **b|** Using simulated single-molecule FRET to measure the R-tRNA interaction reveals the three different intermediates that are generated in this reaction – namely R-tRNA₁, R-tRNA₂ and R-tRNA₃ – that have FRET values of 0.9, 0.3 and 0.6, respectively. [Au: values OK?] These data suggest that there is a step-by-step ‘movement’ from reactant to product⁴⁸. Such observations, within complex systems that are sampling multiple states from within a potential pool of multiple interactions, highlight how incisive such methods could be in glycoscience. Ra is free ribosome prior to tRNA binding and Rb is free ribosome after release of tRNA. This Figure was reproduced, with permission, from reference ⁴⁸.

Fig. 3| Single molecule force spectroscopy to study sugar–lectin interactions.

a| Schematic representation of a lectin-coated atomic force microscopy (AFM) tip interacting with cell surface glycans to study lectin–glycan binding and dissociation. Representative force-distance curves are also presented. **b|** AFM was used to study the antigenicity of branched and linear oligosaccharides from HIV-1 gp120 to the HIV-1 monoclonal antibody 2G12. The force-distance curves in part a are reproduced, with permission, from reference 53. The structures in part b are adapted, with permission, from reference 60.

Fig. 4| Single molecule cell surface glycan ‘mapping’. **a|** Schematic representation of the use of single molecule atomic force microscopy (AFM) to determine the composition of the cell wall of pathogenic microbes in situ. AFM tips functionalized with anti-capsular polysaccharide (CPS) antibodies were used to probe both the localization and density of the outer glycans of *Streptococcus agalactiae* CPSs. [Au: OK?] [Au: Add sentence on anti-β-glucan antibody?] In addition, anti-β-mannan antibodies on AFM tips were used to map the localization of β-mannans on the surface of yeast strains, revealing that their expression was much greater on pathogenic yeast strains than on non-pathogenic yeast strains. [Au: OK?] [Au: Add sentence on ConA?] **b|** AFM force-volume images of the cell wall of wild type *Streptococcus agalactiae* and of *Streptococcus agalactiae* lacking CPS, probed using anti-CPS antibody. [Au:OK?] The region in the dashed box was probed to generate the frequency-adhesion and force-distance profiles shown in panel (c). [Au:OK?] **(c)** The frequency at which the glycan being probed by AFM was present was plotted with respect to adhesive force and the corresponding force-distance. [Au: OK?] [Au: What does green/red

colouring in graphs represent? What do the percentages in graphs represent?]

Images in part b and c were reproduced, with permission, from reference 66.

Fig. 5| The study of glycan processing and presentation using single molecule

fluorescence. **a|** Stochastic phenotype switching in *E. coli* observed for low or intermediate concentrations of intracellular inducer studied using fluorescent lactose permease (using a LacY-YFP fusion) protein. The observation of precise molecular events allows two distinct states of *E. coli*, based on the expression (and hence function) of the sugar-transporter

LacY, to be identified. **b|** Clustering and distribution of gangliosides and

glycosylphosphoinositol on cell surfaces shown by single molecule fluorescence imaging via total internal reflection fluorescence (TIRF). The observed localization of CD59 and

gangliosides in events with varying lifetimes, in both stimulated and unstimulated cells, implicated ganglioside in the formation of longer lasting CD59-cluster signalling rafts as part of stimulation.

[Au:OK?] Part a was adapted, with permission, from reference 80. Part b was reproduced, with permission, from reference 84.

Fig. 6| Creation of artificial cytoskeletal networks on cell surfaces via glycans.

Artificial cytoskeletal networks can be generated on cell surfaces by cross-linking. In one mode this may be achieved by adding azido sugars to cells (step 1), which are then metabolically incorporated into cell-surface glycans. Copper-free azide+alkyne triazole-forming reaction enables the labelling of glycans with biotin (step 2). Finally, the addition of fluorescent streptavidin (that is, streptavidin labelled with an Alexafluor dye), which can bind to biotin, causes connections to form between glycans (step 3). As the streptavidin is fluorescently labelled, this allows this 'tuneable' artificial process to be monitored at the single molecule level.⁸⁷

Fig. 7| Nanopore single channel recording in glycobiology.

Monitoring changes in current transmitted by ion flow through nanopores has allowed the study of various carbohydrate-protein interactions. **a|** Multivalent carbohydrate–lectin interactions can be monitored by functionalization of the pore with glycans and seeing the modulation of current upon binding.⁹³ **b|** Parts of polysaccharide export in bacteria can be recapitulated by using the relevant export pore itself as the nanopore.⁹⁴ **c|** Screening for inhibitors⁹⁵ of polysaccharide export in bacteria also allows the discovery of compounds that may represent a new class of antibacterial. Note that in this figure the colouring of the spheres is arbitrary and does not follow the symbol nomenclature for graphical representations of glycans.¹⁰¹ The trace in part a was reproduced, with permission, from reference 91. Parts of part b and c were reproduced, with permission, from reference^{93,95}. [Au:OK?]

Table 1| Single molecule techniques with potential for use in glycobiology.

Single molecule technique	Information currently obtained using technique	Chemical tools driving technique	Use of technique in glycobiology	Ref
Force spectrometry	Quantitative analysis of binding interactions and binding kinetics	Atomic force microscopy tip coated with lectin and sugar immobilised on mica or a silicon chip	Analysis of the molecular interactions of sugars, including interaction strength	51-54,60,63-65,67-69,102
Single molecule field effect transistors	Dynamic signals during degradation of glycans	Enzyme-tethered carbon nanotubes	Monitoring dynamic glycan processing	75
Single molecule fluorescence microscopy and spectroscopy	Elucidating the mechanisms of biomolecular interactions; in biological processes within living cells	Fluorescent labels	Probing single molecule events in cells	77,81,84,87
Stochastic optical reconstruction microscopy (STORM)	Distribution and diffusion rate of glycans on the cell surface	Photoswitchable fluorophores	Localization and distribution of glycans and partners on the cell surface	88,89,91

Single channel recording using nanopores	Lifetimes of interaction states between glycans, proteins and small molecule inhibitors	A single protein pore (modified or native), that can monitor interactions	Recapitulation of glycan–protein binding processes and screening for molecules that inhibit them	93-95
Recognition tunnelling	Electrode fluctuations according to identity of sugar	Electrodes functionalized with host molecules	Analysis of carbohydrate epimers	99

Glossary terms

Force-distance cycles: The repeated measurement of force as a function of distance in AFM.

Total internal reflection fluorescence (TIRF): Microscopy using a thin region of a specimen (~100 nm), which uses internal reflection at a surface–sample interface for the selective excitation of surface-associated fluorophores, thereby removing contributions to background.

Stochastic optical reconstruction microscopy (STORM): Dilute fluorophores are switched on-and-off through switching or bleaching and the ‘centres’ of their location ascertained using point-spread of photon emission based on a widefield image. After several rounds a ‘super-resolution’ image is assembled by plotting the measured positions of these centres of the fluorescent probes. [\[Au:Shortened - OK?\]](#)

Photo-activated localization microscopy (PALM): Similar to STORM, but has typically used photoactivation and then bleaching to switch fluorophores on-and-off. It is more often associated with the use of protein fluorophores such as photo-activatable fluorescent proteins.

Subject categories

Biological sciences / Chemical biology / Glycobiology

[URI /631/92/221]

Biological sciences / Biophysics / Single-molecule biophysics

[URI /631/57/2265]

Physical sciences / Chemistry / Analytical chemistry

[URI /639/638/11]

Physical sciences / Chemistry / Chemical biology / Carbohydrates / Glycoconjugates

[URI /639/638/92/72/1202]

Studying glycobiology at the single molecule level

Abirami Lakshminarayanan, Mylène Richard & Benjamin G. Davis

Traditional approaches for studying glycans can lack the precision required to uncover their roles in biomolecular processes. Progress has been made in the use of single molecule techniques for examining individual events in glycobiology — such as the molecular interactions of sugars and their roles in biological processes — within challenging mixtures.