

# Consumption of fruits, vegetables, and fruit juices and differentiated thyroid carcinoma risk in the European Prospective Investigation into Cancer and Nutrition (EPIC) study

Raul Zamora-Ros<sup>1,\*</sup>, Virginie Béraud<sup>1,\*</sup>, Silvia Franceschi<sup>2</sup>, Valerie Cayssials<sup>1</sup>, Konstantinos K. Tsilidis<sup>3,4</sup>, Marie-Christine Boutron-Ruault<sup>5,6</sup>, Elisabete Weiderpass<sup>7,8,9,10</sup>, Kim Overvad<sup>11</sup>, Anne Tjønneland<sup>12</sup>, Anne Kirstine Eriksen<sup>12</sup>, Fabrice Bonnet<sup>5,6,13</sup>, Aurélie Affret<sup>5,6</sup>, Verena Katzke<sup>14</sup>, Tilman Kühn<sup>14</sup>, Heiner Boeing<sup>15</sup>, Antonia Trichopoulou<sup>16,17</sup>, Elisavet Valanou<sup>16</sup>, Anna Karakatsani<sup>16,18</sup>, Giovanna Masala<sup>19</sup>, Sara Grioni<sup>20</sup>, Maria Santucci de Magistris<sup>21</sup>, Rosario Tumino<sup>22</sup>, Fulvio Ricceri<sup>23,24</sup>, Guri Skeie<sup>7</sup>, Christine L Parr<sup>25</sup>, Susana Merino<sup>26</sup>, Elena Salamanca-Fernández<sup>27,28</sup>, Maria-Dolores Chirlaque<sup>28,29,30</sup>, Eva Ardanaz<sup>28,31,32</sup>, Pilar Amiano<sup>28,33</sup>, Martin Almquist<sup>34,35</sup>, Isabel Drake<sup>36</sup>, Joakim Hennings<sup>37</sup>, Maria Sandström<sup>38</sup>, H. Bas Bueno-de-Mesquita<sup>4,39</sup>, Petra H. Peeters<sup>4,40</sup>, Kay-Thee Khaw<sup>41</sup>, Nicholas J. Wareham<sup>42</sup>, Julie A. Schmidt<sup>43</sup>, Aurora Perez-Cornago<sup>43</sup>, Dagfinn Aune<sup>4,44,45</sup>, Elio Riboli<sup>4</sup>, Nadia Slimani<sup>2</sup>, Augustin Scalbert<sup>2</sup>, Isabelle Romieu<sup>2</sup>, Antonio Agudo<sup>1</sup>, Sabina Rinaldi<sup>2</sup>

## *Author affiliations*

<sup>1</sup>Unit of Nutrition and Cancer, Cancer Epidemiology Research Programme, Catalan Institute of Oncology, Bellvitge Biomedical Research Institute (IDIBELL), Barcelona, Spain.

<sup>2</sup>International Agency for Research on Cancer (IARC), Lyon, France.

<sup>3</sup>Department of Hygiene and Epidemiology, University of Ioannina School of Medicine, Ioannina, Greece

<sup>4</sup>School of Public Health, Imperial College London, London, UK

<sup>5</sup>CESP, Université Paris-Sud, UVSQ, INSERM, Université Paris-Saclay, Villejuif, France.

<sup>6</sup>Institut Gustave Roussy, F-94805, Villejuif, France.

<sup>7</sup>Department of Community Medicine, Faculty of Health Sciences, UiT, The Arctic University of Tromsø, Tromsø, Norway.

<sup>8</sup>Department of Research, Cancer Registry of Norway, Institute of Population-Based Cancer Research, Oslo, Norway.

<sup>9</sup>Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden.

<sup>10</sup>Genetic Epidemiology Group, Folkhälsan Research Center, Helsinki, Finland.

<sup>11</sup>Department of Public Health, Section for Epidemiology, Aarhus University, Aarhus, Denmark.

<sup>12</sup>Danish Cancer Society Research Center, Copenhagen, Denmark.

<sup>13</sup>CHU Rennes, Rennes 1 University, Rennes, France

<sup>14</sup>Division of Cancer Epidemiology, German Cancer Research Center, Heidelberg, Germany.

<sup>15</sup>Department of Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany.

<sup>16</sup>Hellenic Health Foundation, Athens, Greece

<sup>17</sup>WHO Collaborating Center for Nutrition and Health, Unit of Nutritional Epidemiology and Nutrition in Public Health, Dept. of Hygiene, Epidemiology and Medical Statistics, School of Medicine, National and Kapodistrian University of Athens, Greece

<sup>18</sup>2nd Pulmonary Medicine Department, School of Medicine, National and Kapodistrian University of Athens, “ATTIKON” University Hospital, Haidari, Greece

<sup>19</sup>Cancer Risk Factors and Life-Style Epidemiology Unit, Cancer Research and Prevention Institute – ISPO, Florence, Italy

<sup>20</sup>Nutritional Epidemiology Unit, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy.

<sup>21</sup>Department of Clinical and Experimental Medicine, Federico II University, Naples, Italy.

<sup>22</sup>Cancer Registry and Histopathology Unit, "Civic M.P. Arezzo" Hospital, ASP Ragusa, Italy.

<sup>23</sup>Department of Clinical and Biological Sciences, University of Turin, Turin, Italy

<sup>24</sup>Unit of Epidemiology, Regional Health Service ASL TO3, Grugliasco (TO), Italy

<sup>25</sup>Domain of Mental and Physical Health, Norwegian Institute of Public Health, Oslo, Norway

<sup>26</sup>Public Health Directorate, Asturias, Spain.

<sup>27</sup>Escuela Andaluza de Salud Pública. Instituto de Investigación Biosanitaria ibs.Granada. Hospitales Universitarios de Granada/Universidad de Granada, Granada, Spain.

<sup>28</sup>CIBER de Epidemiología y Salud Pública (CIBERESP), Spain.

<sup>29</sup>Department of Epidemiology, Regional Health Council, IMIB-Arrixaca, Murcia, Spain

<sup>30</sup>Department of Health and Social Sciences, Universidad de Murcia, Murcia, Spain

<sup>31</sup>Navarra Public Health Institute, Pamplona, Spain

<sup>32</sup>Navarra Institute for Health Research (IdiSNA) Pamplona, Spain

<sup>33</sup>Public Health Division of Gipuzkoa, Regional Government of the Basque Country, Spain

<sup>34</sup>Department of Surgery, University Hospital Lund, Lund, Sweden

<sup>35</sup>Malmö Diet and Cancer Study, University Hospital Malmö, Malmö, Sweden

<sup>36</sup>Department of Clinical Sciences Malmö, Lund University, Malmö, Sweden

<sup>37</sup>Department of Surgical and Perioperative Sciences, Umeå University, Umeå, Sweden

<sup>38</sup>Department for Radiation Sciences, Umeå University, Umeå, Sweden

<sup>39</sup>Department for Determinants of Chronic Diseases (DCD), National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands

<sup>40</sup>Department of Epidemiology, Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, The Netherlands

<sup>41</sup>Department of Public Health and Primary Care, University of Cambridge, UK

<sup>42</sup>MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Cambridge, UK

<sup>43</sup>Cancer Epidemiology Unit, University of Oxford, UK

<sup>44</sup>Department of Public Health and General Practice, Norwegian University of Science and Technology, Trondheim, Norway.

<sup>45</sup>Björknes University College, Oslo, Norway.

\*Both authors equally contributed to this research.

The entire manuscript is 4,181 words (including abstract (229 words), text (3,952 words), 53 references, 3 tables, 3 supplementary table).

**Running Title:** Fruits, vegetables, and fruit juices and thyroid cancer in EPIC

**Corresponding author:** Raul Zamora-Ros, Unit of Nutrition and Cancer, Catalan Institute of Oncology (ICO), Bellvitge Biomedical Research Institute (IDIBELL), Av Gran Via 199-203, 08908 L'Hospitalet de Llobregat, Spain. Phone: +34 932067401. Fax: +34 932607787.

E-mail: rzamora@iconcologia.net

**Key words:** Thyroid cancer, Fruits, Vegetables, Fruit juices, Intake, EPIC

**Abbreviations:** 24-HDR: 24-h dietary recall; BMI: Body Mass Index; CI: Confidence Interval; EPIC: European Prospective Investigation into Cancer and Nutrition; F&V: Fruits and Vegetables; HR: Hazard Ratio; IARC: International Agency for Research on Cancer; NOS: Not Otherwise Specified; TC: Thyroid cancer

**Article category:** Epidemiology

**NOVELTY AND IMPACT OF THE WORK:**

This large prospective cohort does not support any association between the consumption of vegetable, fruit and any individual type of vegetable or fruit and differentiated thyroid carcinoma risk, overall and by histologic type (papillary and follicular tumours). However, a positive trend with fruit juice intake was observed, possibly related to its high sugar content.

## 1    **ABSTRACT**

2    Fruit and vegetable (F&V) intake is considered as probably protective against overall cancer  
3    risk, but results in previous studies are not consistent for thyroid cancer (TC). The purpose of  
4    this study is to examine the association between the consumption of fruits, vegetables, fruit  
5    juices and differentiated thyroid cancer risk within the European Prospective Investigation  
6    into Cancer and Nutrition (EPIC) study. The EPIC study is a cohort including over half a  
7    million participants, recruited between 1991 and 2000. During a mean follow-up of 14 years,  
8    748 incident first primary differentiated TC cases were identified. F&V and fruit juice intakes  
9    were assessed through validated country-specific dietary questionnaires. Hazard ratios (HRs)  
10    and 95% confidence intervals (CIs) were estimated using Cox regression models adjusted for  
11    potential confounding factors. Comparing the highest vs. lowest quartile of intake,  
12    differentiated TC risk was not associated with intakes of total F&V (HR: 0.89; 95% CI: 0.68-  
13    1.15; p-trend=0.44), vegetables (HR: 0.89; 95% CI: 0.69-1.14; p-trend=0.56), or fruit (HR:  
14    1.00; 95% CI: 0.79-1.26; p-trend=0.64). No significant association was observed with any  
15    individual type of vegetable or fruit. However, there was a positive borderline trend with fruit  
16    juice intake (HR: 1.23; 95% CI: 0.98-1.53; p-trend=0.06). This study did not find any  
17    significant association between F&V intakes and differentiated TC risk; however a positive  
18    trend with fruit juice intake was observed, possibly related to its high sugar content.

## INTRODUCTION

The consumption of fruits and vegetables (F&V) has been consistently associated with a reduced risk of many cancers in case-control studies, but these associations become weak or even null in cohorts.<sup>1-3</sup> In 2003 the World Health Organization (WHO)<sup>4</sup> and in 2007 the International Agency for Research on Cancer (IARC)<sup>5</sup> concluded that there was convincing evidence that F&V lower the risk of cardiovascular diseases and probable evidence for lower respiratory and digestive cancer risks. Therefore, they recommended an intake of 400–500 g/day, the well-known campaign of 5 a day (including vegetables, fruits and also juices, although juices are usually recommended  $\leq 1$  portion per day by several national nutrition health programmes, e.g. in the UK, France, Norway, etc.). However, little is known about its association with thyroid cancer (TC) risk, particularly in prospective studies. TC is the most common endocrine cancer (1-1.5% of all new cancers diagnosed each year) worldwide<sup>6</sup> and the incidence has been constantly increasing over the last three decades,<sup>7-10</sup> mainly explained by over-diagnosis.<sup>11</sup>

In 2014, a meta-analysis of 19 case-control studies on dietary factors and TC risk found a weak inverse association with the intake of vegetables and no association with the intake of fruits.<sup>12</sup> Cruciferous vegetables have been studied more closely due to their content in goitrogenic substances,<sup>13</sup> which have tumour promoting effects upon rats' thyroid.<sup>14,15</sup> Two recent meta-analyses of retrospective studies found either no association or a positive association between the intake of cruciferous vegetables and differentiated TC risk.<sup>12,16</sup> For juices, a positive association was suggested between citrus fruit juice intake and TC risk.<sup>17</sup>

The aim of the present study was to prospectively evaluate the association between the consumption of vegetables, fruits, fruit juices and their subtypes and differentiated TC risk in the European Prospective Investigation into Cancer and Nutrition (EPIC) study, a large cohort with a high heterogeneity in dietary intake across the participating countries.<sup>18</sup>



45

## 46 **MATERIAL AND METHODS**

### 47 *Study population*

48 EPIC is a multicentre prospective cohort study principally designed to investigate the  
49 relationships between diet and other environmental factors and cancer risk. The study design  
50 has been explained in detail previously.<sup>19,20</sup> In brief, EPIC enrolled more than half a million  
51 participants, 70% women, mostly aged between 35 and 70 years old from 23 centres in 10  
52 western European countries (Denmark, France, Germany, Greece, Italy, Norway, Sweden,  
53 Spain, the Netherlands, and the United Kingdom) during the period 1991-2000. All  
54 participants gave their written informed consent, and the Internal Review Boards of the IARC  
55 and all ethical committees from participating centres approved the project.

### 56 *Dietary assessment*

57 Validated country-specific dietary questionnaires were used to evaluate participants' diets  
58 over the 12 months preceding recruitment.<sup>19</sup> They were designed to collect the specificity of  
59 local dietary habits and provide optimal compliance. According to the centres, quantitative or  
60 semi-quantitative methods were applied. In Malmö-Sweden, a combination of a 7-day record  
61 and a semi-quantitative food frequency questionnaire was administered. Questionnaires were  
62 mainly self-reported except for Greece, Spain and two centres in Italy (Naples and Ragusa)  
63 where they were administered during a face-to-face interview. Lifestyle questionnaires about  
64 educational and socio-economic status, lifetime history of tobacco smoking and alcohol  
65 consumption, physical activity, menstrual and reproductive history and medical history were  
66 also collected at baseline.<sup>19</sup>

67 The analysed food groups were vegetables (potatoes and legumes were not included), fresh  
68 fruits (excluding nuts, olives, and seeds), and total juices or fruit juices (including citrus fruit

juices, non-citrus fruit juices, vegetable juices and fruit nectars). Vegetable juices were rarely consumed (<2% of total juices) and this item was only recorded in the dietary questionnaires of 3 countries; for this reason we have simplified and used the term fruit juices instead of total juices. Total intake of F&V was calculated as the sum of fruits and vegetables, excluding fruit juices because of their large differences in nutritional composition compared to F&V.<sup>18</sup> Subgroups of vegetables and fruits were also considered: leafy vegetables, fruiting vegetables, root vegetables, cabbages, mushrooms, grain and pod vegetables, onion/garlic, sprouts/stalk vegetables, citrus fruits, hard fruits (apples and pears), berries (strawberries, grapes, blueberries), stone fruits (peach, nectarine, apricot, plum, cherry), banana, and other fruits (fig, kiwi, melon, watermelon, pineapple). Details about the composition of the subgroups are given elsewhere.<sup>18</sup>

#### ***Follow-up and case ascertainment***

Except for Germany and Greece, all information on vital status came from national or regional mortality registries. In Germany and Greece, those data were continuously collected through an active follow-up. Cancer incident cases were identified through regional/national cancer registries, except for German, Greek, and French centres. For those, different methods were used, including health insurance records, contacts with cancer pathology registries, and active follow-up through study participants or relatives. Complete follow-up censoring dates ranged from December 2010 to December 2014, depending on the study centre.<sup>19</sup> Cancer cases were defined as subjects with a first primary TC (code C73 according to the International Classification of Diseases, 10th Revision) during follow-up. Differentiated TC comprises papillary and follicular tumours.

Of the 857 TC cases, anaplastic (n=9), medullary (n=37), lymphoma (n=1) forms and “other morphologies” (n=5) were excluded. Moreover, 29,332 participants (including 45 differentiated TC cases) were excluded from the initial database because of missing or null

follow-time or having a prevalent cancer other than non-melanoma skin cancer. Further 1,277 and 14,555 participants (including 12 differentiated TC cases) were also excluded because their lifestyle and dietary data were not complete or coherent (they were in the top or the bottom 1% of the distribution of the total energy intake to energy requirement ratio), respectively. Finally, 748 first primary incident differentiated TC cases were considered: 601 papillary TCs, 109 follicular TCs, and 38 not otherwise specified (NOS) TCs, most likely papillary TCs. A total of 142,232 men and 333,876 women were included in the present study.

### ***Statistical analysis***

Cox proportional hazard models were used to estimate hazard ratios (HR) and 95% confidence intervals (CI). Intakes of vegetables, fruits, and fruit juices were stated in grams per day. They were analysed as continuous variables (increments of 100 g/d for groups and 50 g/d or 10 g/d for subgroups), and as categorical variables using EPIC-wide quartiles (quartile 1 as reference) or EPIC-specific quartiles by sex, BMI categories, and physical activity status for subgroup analyses. Tests for linear trend were performed by attributing the median of each quartile as a score. Age was the primary time variable in all models. Entry time was age at recruitment and exit time was age at diagnosis of differentiated TC, death, or censoring date (loss of follow-up, end of follow-up), whichever occurred first. Two sets of models were used without (model A) and with (model B) adjustment for total energy intake. Model A1 was stratified by sex, age at recruitment (1-year interval) and centre. Model A2 was model A1 additionally adjusted for non-dietary variables: BMI, smoking status (never, former, current smoker, and unknown), education (primary, secondary, and unknown), physical activity (inactive, active, and unknown according to the Cambridge Physical Activity Index),<sup>21</sup> type of menopause (pre-, peri-, post-, and surgical menopause), use of oral contraceptives (yes, no, and unknown), and fertility problems (yes, no, and unknown), since these reproductive factors

were TC risk factors in our previous study.<sup>22</sup> Model B1 was stratified by sex, age recruitment (1-year interval) and centre, and adjusted for total energy intake (kcal/day). Model B2 was model B1 additionally adjusted for non-dietary variables (as model A2) and dietary variables: total alcohol (g/day) intake. We also created an additional model (model B3) for the main exposure groups, which was model B2 mutually adjusted, when appropriate, for vegetables, fruits, and fruit juices, accordingly. The interpretation of models without total energy intake is focused on the impact of the effect of the absolute amount of compounds of F&V and fruit juices (e.g. vitamins, polyphenols, and contaminants). Whereas in the case of the models with total energy intake adjustment, the relevance is the relative amount compare to other compounds, like in a substitution model. Tests and graphs based on Schoenfeld residuals were used to assess proportional hazards assumptions, which were satisfied. Separate analyses were performed for TC subtypes: follicular and papillary tumours. Heterogeneity of risk between TC subtypes was assessed with the Wald test. Possible interactions with sex, smoking status, BMI (BMI<25 vs. BMI≥25), alcohol intake at recruitment (≤24 g/d vs. >24 g/d), and physical activity (inactive, active) were tested by including an interaction term in the multi-adjusted models. The Wald test was used to evaluate the heterogeneity of risk between BMI and physical activity categories. Sensitivity tests were performed i) excluding female participants from France, since they contributed to 37.2% of all differentiated TC cases among all female participants, and over-diagnosis in France could be relatively high;<sup>11</sup> (ii) excluding participants who had diabetes or unknown diabetes status at baseline, because diabetes is a potential risk factor of TC and iii) excluding cases with a follow-up period below two years, since they could have changed their diets in the pre-diagnostic period. For all analyses, p-values <0.05 were considered as statistically significant and p-values >0.05 and <0.1 as borderline statistically significant. To account for multiple testing for the subgroups of F&V, Bonferroni correction was used and then results were considered statistically significant if

P<0.05/23 (number of tests for the intakes of total F&V, vegetables, fruits, fruit juices and all subgroups)=0.002. All statistical analyses were performed with the R 3.3.1 software (R Foundation for Statistical Computing, Vienna, Austria).

### *Calibration of dietary data*

A second dietary measurement was taken from an 8% random sample of the cohort (36,994 participants) using a detailed computerized 24-h dietary recall (24-HDR) method<sup>23</sup> to calibrate dietary measurements of vegetable, fruit, and fruit juice intake across countries and to correct for systematic overestimation or underestimation of dietary intakes.<sup>24</sup> The 24-HDR values of these participants were regressed on the main dietary questionnaire values for vegetables, fruits, and fruit juices with adjustment for age at recruitment, centre, and total energy intake, and weighting by day of the week and season of the year during which the 24-HDR was collected. Zero consumption values in the main dietary questionnaires were excluded in the regression calibration models, and a zero was directly imputed as a corrected value. Country and sex-specific calibration models were used to obtain individual predicted values of dietary exposure for all participants. Cox regression models were then run using the predicted (calibrated) values for each individual on a continuous scale. The standard error of the calibrated coefficient was estimated with bootstrap sampling in the calibration and disease models and repeated 300 times.<sup>24</sup>

## **RESULTS**

The median (percentile 25<sup>th</sup> and 75<sup>th</sup>) intakes of total F&V, vegetables, fruits, and fruit juices were 391 g/d (250-576 g/d), 175 g/d (110-276 g/d), 194 g/d (106-314 g/d), and 20 g/d (1-94 g/d), respectively. A great heterogeneity of F&V consumption in both men and women among EPIC countries was observed with higher F&V intakes in Southern than in Northern EPIC countries(**Table 1**). Differentiated TC incidence was higher among women (89% of cases) and the most common subtype was papillary (80.3%), followed by follicular (14.6%),

and NOS tumours (5.1%) (Table 1). The analysis of baseline characteristics according to quartiles of total intake of F&V showed that participants in the highest quartile were more likely to be older women, less educated, and less physically active, and to have higher total energy intake and lower tobacco and alcohol consumptions. Women in the highest quartile of F&V tended to be postmenopausal or to have undergone surgical menopause, to have more infertility problems, and to take less oral contraceptives at the baseline (**Supplementary Table 1**).

Four multivariable Cox models with different levels of adjustment (with and without total energy intake) were performed, showing similar results (difference below 10%) (**Table 2 and Supplementary table 2**). In the fully adjusted model (model B2), no association was observed between total intake of F&V and total differentiated TC risk for the highest vs. lowest quartile (HR: 0.89; 95% CI: 0.68-1.15; p-trend=0.44). No association was found with the intake of vegetables (HR: 0.89; 95% CI: 0.69-1.14; p-trend=0.56), or fruits (HR: 1.00; 95% CI: 0.79-1.26; p-trend=0.64). A borderline positive trend between fruit juice intake and total differentiated TC risk was detected (HR: 1.23; 95% CI: 0.98-1.53; p-trend=0.06). After calibration, results were unchanged. In separate analyses by TC type, null results were observed between all exposures and either papillary or follicular TC risk (Table 2). A further model (model B3) was also estimated mutually adjusting for the intake of fruits, vegetables or fruit juices when appropriate and results were practically identical to those in model B2 (Supplementary table 2). Sensitivity tests, excluding French participants, subject with diabetes at baseline, and cases with a follow-up below two years, showed results almost identical to those in the entire cohort (data not shown).

No association between any subgroups of F&V intake and total differentiated TC risk was found (**Table 3**). Associations with papillary and follicular TC risk were also evaluated. For every 10g/day of sprouts/stalk vegetables, the risk of follicular TC was significantly lowered

for the observed data (HR: 0.72; 95% CI: 0.56-0.94; p-value=0.015) and for the calibrated data (HR: 0.56; 95% CI: 0.34-0.91%; p-value=0.020), but not for papillary TC risk. After applying the Bonferroni correction ( $p < 0.002$ ), the inverse association was no longer statistically significant. No associations were observed for any of the other subgroups of F&V with either papillary or follicular TC risk.

No interactions were observed between total intake of F&V and any of the tested potential confounding factors: sex (P for interaction=0.84), smoking status (P for interaction=0.58), BMI (P for interaction=0.37), alcohol (P for interaction=0.54), and physical activity (P for interaction=0.41). Although no heterogeneity was observed between sexes, separate analyses by sex were performed because of the large proportion of women with TC. The study of the association between total intake of F&V and TC risk showed similar results from the previous analyses for both women (HR: 0.88; 95% CI: 0.68-1.15; p-trend=0.34) and men (HR: 0.95; 95% CI: 0.42-2.12; p-trend=0.85) (data not tabulated). A statistically significant interaction between fruit juice intake and BMI (P for interaction=0.03) and borderline significant with physical activity (P for interaction=0.08) was detected. The highest quartile of fruit juice consumption was positively associated with TC risk among inactive participants (HR: 1.31; 95% CI: 0.99-1.72; p-trend=0.05) but not among active subjects (HR: 1.00; 95% CI: 0.70-1.42; p-trend=0.99); and in participants with BMI<25 (HR: 1.43; 95% CI: 1.08-1.89; p-trend=0.002), but not in those with BMI $\geq$ 25 (HR: 1.06; 95% CI: 0.75-1.48; p-trend=0.99). Similar results were found for the analyses by TC subtypes, especially by BMI categories (Supplementary table 3).

## DISCUSSION

In the present study, we prospectively evaluated the association between total F&V and fruit juice intakes and differentiated TC risk. We did not observe any association between total F&V intake and either all differentiated, papillary, or follicular TC risk. Similar null results

were also found in a Korean case-control study on mainly papillary TC cases (90%) among women.<sup>25</sup> Furthermore in our study, no associations with the consumption of either fruits or vegetables were observed; whereas a borderline statistical positive trend with fruit juice intake was detected.

A meta-analysis of 19 case-control studies<sup>12</sup> and a pooled-analysis of 11 case-control studies<sup>26</sup> summarized the associations between vegetable intake and TC risk, showing a weak inverse association with the intake of total vegetables and vegetables excluding cruciferous vegetables. An American cohort study of 300,000 participants (the NIH-AARP diet and health study) showed an impact of adolescent (12-13 years old) and midlife diet (10 years before recruitment when respondents were 41–62 years of age) on TC risk, with a significant inverse association of vegetable intake among women.<sup>27</sup> Although intakes of vegetables consumed by both Americans at midlife and our study population were comparable, we did not find any association between the intake of vegetables and the risk of differentiated TC either overall or by subtype.

In the present cohort study, we observed no association between differentiated TC risk and the consumption of cruciferous vegetables. Similar results were observed in two meta-analyses and one pooled-analysis.<sup>12,26,28</sup> However, in the most recent meta-analysis, a positive association was pointed out, after excluding studies evaluating only one type of cruciferous vegetables and hospital-based case-control studies.<sup>16</sup> The NIH-AARP diet and health study also showed a positive association with the intake of broccoli during midlife in men, however no other cruciferous vegetables were considered in the dietary questionnaire.<sup>27</sup> Overall, no association between the intake of vegetables and cruciferous vegetables and TC was observed in the last 5-10 years before diagnosis (i.e. middle and late adulthood), although potential relationships in earlier life stages, such as adolescence and young adulthood, could not be ruled out according to the results from the NHI-AARP study. Those results suggest that a diet



rich in cruciferous vegetables during early adulthood could have an impact in the early development of TC, but not during the 5-10 years prior to TC diagnosis, which is the mean time of follow-up of TC cases in our study.

Overall, no associations were found with vegetable subclasses with overall, papillary or follicular TC risk. Some previous studies suggested inverse associations with raw vegetables,<sup>25</sup> green vegetables,<sup>29</sup> carrots,<sup>29,30</sup> green salad,<sup>30</sup> raw tomatoes,<sup>31</sup> turnips,<sup>32</sup> rutabagas,<sup>32</sup> and cassava.<sup>33</sup> However, these studies were relatively small (<400 TC cases) and had a retrospective design. Moreover these inverse associations have not been consistently observed in other similar studies and were not even suggested in our large cohort. In the analysis by TC type, there was an inverse association with sprouts/stalk vegetables but only for follicular TC risk (n=109 cases), although no longer after Bonferroni correction. In addition, sprouts/stalk vegetables (including asparagus, celery, fennel and leek) are a very heterogeneous group and are generally consumed in low amounts in Europe, and therefore the results could be due to chance.

In the current study, total fruit intake, as well as intake of fruit subtypes was not associated with TC risk. Although less information is available on this field, our findings are in concordance with the results of previous studies, including the meta-analysis of 19 case-control studies<sup>12</sup> and the American cohort study considering adolescent and midlife periods.<sup>27</sup> Some studies showed a special interest in citrus fruits because of their high content in antioxidants (such as vitamin C and flavonoids). A pooled-analysis of four Italian-based case-control studies showed an inverse association with TC risk.<sup>30</sup> Both Greek and South Korean case-control studies reported inverse associations with the consumption of lemons<sup>31</sup> and tangerines,<sup>25</sup> respectively. However, in the present study, there was no association with citrus fruits intake, similarly to the NIH-AARP cohort where consumption of orange and tangelos was not associated with TC risk in any period.<sup>17</sup>

269 In the present study, a positive trend between the intake of fruit juices and TC risk was  
270 observed, especially in subjects with a BMI<25 or physically inactive. In the NIH-AARP  
271 cohort study, the consumption of orange and grapefruit juices was also associated with a  
272 higher risk of TC.<sup>17</sup> In our study, fruit juices were mainly citrus juices (>61%), but most of  
273 them were commercial fruit juices and fruit nectars (62.3%) with a high sugar content. The  
274 contribution of fruit juices to total sugars varied from <2% in Mediterranean countries to 15%  
275 in Germany.<sup>34</sup> In our study, we cannot differentiate between commercial fruit juice and fruit  
276 nectar (fruit juice with added sugars), and therefore we cannot evaluate the impact of the  
277 consumption of fruit juices with different amounts of sugar in the TC risk. Moreover, we  
278 cannot distinguish between fruit juices and vegetable juices in all participating countries,  
279 although vegetable juices are rarely consumed (only contributing to <2% of total juices in our  
280 study). Our results by BMI categories were in concordance with a recent study within EPIC,  
281 showing that the intake of sugar was positively associated with TC risk among people with a  
282 BMI <25.<sup>35</sup> This finding could be partially due to a higher consumption of fruit juices in  
283 subjects with a BMI<25 (66.9 g/d) than with a BMI≥25 (60.4 g/d). However, the opposite  
284 results would be expected, since overweight is a major determinant of insulin resistance and  
285 hyperinsulinemia,<sup>36</sup> which were associated with a higher prevalence of differentiated TC.<sup>37</sup>  
286 Furthermore, after calibration, no differences in the association between fruit juice intake and  
287 TC by BMI were observed. This could be due to a higher under-reporting of fruit juice intake  
288 among overweight people. Therefore, we need to be cautious in the interpretation of these  
289 sub-analyses. In the analyses by physical activity status, the intake of fruit juices was  
290 positively associated with TC risk among inactive participants. It has been reported that  
291 physical activity significantly improves glycaemia and sensitivity to insulin, which have been  
292 associated with a higher risk of TC.<sup>37,38</sup> Moreover, physical activity modifies female hormone  
293 levels,<sup>39</sup> which may have a weak influence on TC risk.<sup>22,40</sup>

On one hand, various components of F&V and fruit juices could exert a protective effect against TC. The intake of vitamins (in particular vitamin C and E, carotenoids), polyphenols and fibre has been related to a reduced risk of TC in some studies.<sup>41,42</sup> Moreover, a deficiency in magnesium may be involved in the aetiology of some thyroid diseases.<sup>43</sup> However the protective effects of F&V and fruit juices components are not consistent and several studies showed null results,<sup>17,35,44,45</sup> or even positive associations [e.g., with beta-carotene,<sup>46</sup> vitamin C<sup>47</sup> and flavanones].<sup>17</sup>

On the other hand, F&V and fruit juices can also contain traces of chemical pesticides. A high exposure to organophosphate insecticides was associated with a higher TC incidence among spouses of pesticide applicators.<sup>48</sup> Vegetables are also good sources of nitrates, accounting for 80% of their intake in Westernized diets.<sup>49</sup> They are involved in the formation of N-nitroso compounds, which are known carcinogens.<sup>50</sup> Indeed, nitrate intake has been positively associated with TC risk in two cohort studies.<sup>51,52</sup> Cruciferous vegetables contain large amounts of glucosinolates, which are a nutritional source of thiocyanates and isothiocyanates. Those molecules can block the action of carcinogenic substances and suppress the expression of neoplasia in initiated cancer cells,<sup>53</sup> but they also act as goitrogens,<sup>13</sup> promoting thyroid tumour growth in rats.<sup>14,15</sup>

Strengths of the EPIC study are its prospective design, its large number of cancer cases, and the wide range of F&V and fruit juice intake across participants from 10 European countries with standardized information on diet and lifestyle exposures. Several limitations have to be taken into account. Although EPIC is a large study, participants are not representative of the whole population in most of the countries because of the criteria of recruitment, and therefore the results are not totally generalizable to general population. The measurement error in the estimation of F&V and fruit juice intake by dietary questionnaires may have attenuated our findings, although the use of the calibrated F&V and fruit juice data, obtained from the

standardized 24-HDR, did not alter the results. As discussed above, we do not have data on fruit juices and fruit nectars separately. Dietary and lifestyle habits during the follow-up period are not available in EPIC, although diet is relatively stable in adult populations through the years.

To conclude, in the present study, no association with differentiated TC risk was observed for the consumption of total F&V, vegetables, fruits, and their subgroups. This result might be explained by a counterbalance between pro- and anti-carcinogenic effects of their components. However, fruit juice intake was associated with an increased risk of differentiated TC, possibly due to the high sugar content. Our study supports having a diverse intake of F&V, but not consuming too much fruit juice, especially those rich in sugars. Further prospective studies are warranted to confirm these relationships between fruit juice intake and TC risk.

**Conflict of interest:** The authors declare no conflict of interest.

**Acknowledgments:** We thank Mr Bertrand Hémon and Miss Leila Luján-Barroso for their valuable help with the EPIC database.

**Financial support:** This study was supported by the Institute of Health Carlos III, Spain (CP15/00100), and cofounded by the European Regional Development Fund (ERDF) “A way to build Europe”. The coordination of EPIC is financially supported by the European Commission (DG-SANCO) and the International Agency for Research on Cancer. The national cohorts are supported by Danish Cancer Society (Denmark); Ligue Contre le Cancer, Institut Gustave Roussy, Mutuelle Générale de l’Education Nationale, Institut National de la Santé et de la Recherche Médicale (INSERM) (France); German Cancer Aid, German Cancer Research Center (DKFZ), Federal Ministry of Education and Research (BMBF) (Germany); the Hellenic Health Foundation (Greece); Associazione Italiana per la Ricerca sul Cancro-

344 AIRC-Italy and National Research Council (Italy); Dutch Ministry of Public Health, Welfare  
345 and Sports (VWS), Netherlands Cancer Registry (NKR), LK Research Funds, Dutch  
346 Prevention Funds, Dutch ZON (Zorg Onderzoek Nederland), World Cancer Research Fund  
347 (WCRF), Statistics Netherlands (The Netherlands); Health Research Fund (FIS): PI13/00061  
348 to Granada; PI13/01162 to EPIC-Murcia, Regional Governments of Andalucía, Asturias,  
349 Basque Country, Murcia and Navarra, AGAUR, Generalitat de Catalunya (exp. 2014 SGR  
350 726), The Health Research Funds (RD12/0036/0018) (Spain); Swedish Cancer Society,  
351 Swedish Research Council and County Councils of Skåne and Västerbotten (Sweden); Cancer  
352 Research UK (14136 to EPIC-Norfolk; C570/A16491 and C8221/A19170 to EPIC-Oxford),  
353 Medical Research Council (1000143 to EPIC-Norfolk, MR/M012190/1 to EPIC-Oxford)  
354 (United Kingdom). RZ-R would like to thank the “Miguel Servet” program (CP15/00100)  
355 from the Institute of Health Carlos III (Spain) and the European Social Fund (ESF).

356

## References

1. Aune D, Giovannucci E, Boffetta P, Fadnes LT, Keum N, Norat T, Greenwood DC, Riboli E, Vatten LJ, Tonstad S. Fruit and vegetable intake and the risk of cardiovascular disease, total cancer and all-cause mortality—a systematic review and dose-response meta-analysis of prospective studies. *Int J Epidemiol* 2017;In press. doi.org/10.1093/ije/dyw319
2. Boffetta P, Couto E, Wichmann J, Ferrari P, Trichopoulos D, Bueno-de-Mesquita HB, van Duijnhoven FJ, Büchner FL, Key T, Boeing H, Nöthlings U, Linseisen J, Gonzalez CA, Overvad K, Nielsen MR, Tjønneland A, Olsen A, Clavel-Chapelon F, Boutron-Ruault MC, Morois S, Lagiou P, Naska A, Benetou V, Kaaks R, Rohrmann S, Panico S, Sieri S, Vineis P, Palli D, van Gils CH, Peeters PH, Lund E, Brustad M, Engeset D, Huerta JM, Rodríguez L, Sánchez MJ, Dorronsoro M, Barricarte A, Hallmans G, Johansson I, Manjer J, Sonestedt E, Allen NE, Bingham S, Khaw KT, Slimani N, Jenab M, Mouw T, Norat T, Riboli E, Trichopoulou A. Fruit and vegetable intake and overall cancer risk in the European Prospective Investigation into Cancer and Nutrition (EPIC). *J Natl Cancer Inst* 2010;102:529–37.
3. Takachi R, Inoue M, Ishihara J, Kurahashi N, Iwasaki M, Sasazuki S, Iso H, Tsubono Y, Tsugane S; JPHC Study Group. Fruit and vegetable intake and risk of total cancer and cardiovascular disease. *Am J Epidemiol* 2008;167:59–70.
4. World Health Organization. Diet, nutrition and the prevention of chronic diseases. Technical Report Series, No. 916; 2003. 159 p.
5. World Cancer Research Fund, American Institute for Cancer Research. Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective. Cancer Research. 2007. 517 p.
6. Milas ZL, Brainard J, Milas M. Thyroid cancer: classification and cytological diagnosis. In *Clinical Management of Thyroid Cancer*. Future Medicine Ltd. 2013. p18–32.

- 382 7. Wiltshire JJ, Drake TM, Uttley L, Balasubramanian SP. Systematic Review of Trends  
383 in the Incidence Rates of Thyroid Cancer. *Thyroid* 2016;26(11):1541-1552.
- 384 8. Albores-Saavedra J, Henson DE, Glazer E, Schwartz AM. Changing patterns in the  
385 incidence and survival of thyroid cancer with follicular phenotype - Papillary, follicular, and  
386 anaplastic: A morphological and epidemiological study. *Endocr Pathol* 2007;18:1-7.
- 387 9. Hodgson NC, Button J, Solorzano CC. Thyroid cancer: is the incidence still  
388 increasing? *Ann Surg Oncol* 2004;11:1093-7.
- 389 10. Burgess JR. Temporal trends for thyroid carcinoma in Australia: an increasing  
390 incidence of papillary thyroid carcinoma (1982-1997). *Thyroid* 2002;12:141-9.
- 391 11. Vaccarella S, Franceschi S, Bray F, Wild CP, Plummer M, Dal Maso L. Worldwide  
392 Thyroid-Cancer Epidemic? The Increasing Impact of Overdiagnosis. *N Engl J Med*  
393 2016;375:614-7.
- 394 12. Liu ZT, Lin AH. Dietary factors and thyroid cancer risk: a meta-Analysis of  
395 observational studies. *Nutr Cancer* 2014;66: 1165-78.
- 396 13. Eisenbrand G, Gelbke HP. Assessing the potential impact on the thyroid axis of  
397 environmentally relevant food constituents/contaminants in humans. *Arch Toxicol*  
398 2016;90:1841-57.
- 399 14. Philip J., Crooks J, Macgregor AG, McIntosh JA. The growth curve of the rat thyroid  
400 under a goitrogenic stimulus. *Br J Cancer* 1969;23:515-23.
- 401 15. Kanno J, Matsuoka C, Furuta K, Onodera H, Miyajima H, Maekawa A, Hayashi Y.  
402 Tumor promoting effect of goitrogens on the rat thyroid. *Toxicol Pathol* 1990;18:239-46.
- 403 16. Cho YA, Kim J. Dietary Factors Affecting Thyroid Cancer Risk: A Meta-Analysis.  
404 *Nutr Cancer* 2015;67:811-7.

- 405 17. Xiao Q, Park Y, Hollenbeck AR, Kitahara CM. Dietary flavonoid intake and thyroid  
406 cancer risk in the NIH-AARP diet and health study. *Cancer Epidemiol Biomarkers Prev*  
407 2014;23:1102–8.
- 408 18. Agudo A, Slimani N, Ocké MC, Naska A, Miller AB, Kroke A, Bamia C, Karalis D,  
409 Vineis P, Palli D, Bueno-de-Mesquita HB, Peeters PH, Engeset D, Hjartåker A, Navarro C,  
410 Martínez Garcia C, Wallström P, Zhang JX, Welch AA, Spencer E, Stripp C, Overvad K,  
411 Clavel-Chapelon F, Casagrande C, Riboli E. Consumption of vegetables, fruit and other plant  
412 foods in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohorts  
413 from 10 European countries. *Public Health Nutr* 2002;5:1179–96.
- 414 19. Riboli E, Hunt K, Slimani N, Ferrari P, Norat T, Fahey M, Charrondière UR, Hémon  
415 B, Casagrande C, Vignat J, Overvad K, Tjønneland A, Clavel-Chapelon F, Thiébaud A,  
416 Wahrendorf J, Boeing H, Trichopoulos D, Trichopoulou A, Vineis P, Palli D, Bueno-De-  
417 Mesquita HB, Peeters PH, Lund E, Engeset D, González CA, Barricarte A, Berglund G,  
418 Hallmans G, Day NE, Key TJ, Kaaks R, Saracci R. European Prospective Investigation into  
419 Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr*  
420 2002;5:1113-24.
- 421 20. Riboli E, Kaaks R. The EPIC Project: rationale and study design. European  
422 Prospective Investigation into Cancer and Nutrition. *Int J Epidemiol* 1997;26 Suppl 1:S6–14.
- 423 21. Wareham NJ, Jakes RW, Rennie KL, Schuit J, Mitchell J, Hennings S, Day NE.  
424 Validity and repeatability of a simple index derived from the short physical activity  
425 questionnaire used in the European Prospective Investigation into Cancer and Nutrition  
426 (EPIC) study. *Public Health Nutr* 2003;6:407–13.
- 427 22. Zamora-Ros R, Rinaldi S, Biessy C, Tjønneland A, Halkjaer J, Fournier A, Boutron-  
428 Ruault MC, Mesrine S, Tikk K, Fortner RT, Boeing H, Förster J, Trichopoulou A,  
429 Trichopoulos D, Papatesta EM, Masala G, Tagliabue G, Panico S, Tumino R, Polidoro S,



430 Peeters PH, Bueno-de-Mesquita HB, Weiderpass E, Lund E, Argüelles M, Agudo A, Molina-  
 431 Montes E, Navarro C, Barricarte A, Larrañaga N, Manjer J, Almquist M, Sandström M,  
 432 Hennings J, Tsilidis KK, Schmidt JA, Khaw KT, Wareham NJ, Romieu I, Byrnes G, Gunter  
 433 MJ, Riboli E, Franceschi S. Reproductive and menstrual factors and risk of differentiated  
 434 thyroid carcinoma: The EPIC study. *Int J Cancer* 2015;136:1218–27.

435 23. Slimani N, Kaaks R, Ferrari P, Casagrande C, Clavel-Chapelon F, Lotze G, Kroke A,  
 436 Trichopoulos D, Trichopoulou A, Lauria C, Bellegotti M, Ocké MC, Peeters PH, Engeset D,  
 437 Lund E, Agudo A, Larrañaga N, Mattisson I, Andren C, Johansson I, Davey G, Welch AA,  
 438 Overvad K, Tjønneland A, Van Staveren WA, Saracci R, Riboli E. European prospective  
 439 investigation into cancer and nutrition (EPIC) calibration study: rationale, design and  
 440 population characteristics. *Public Health Nutr* 2002;5:1125–45.

441 24. Ferrari P, Kaaks R, Fahey MT, Slimani N, Day NE, Pera G, Boshuizen HC, Roddam  
 442 A, Boeing H, Nagel G, Thiebaut A, Orfanos P, Krogh V, Braaten T, Riboli E; European  
 443 Prospective Investigation into Cancer and Nutrition study. Within- and between-cohort  
 444 variation in measured macronutrient intakes, taking account of measurement errors, in the  
 445 European prospective investigation into cancer and nutrition study. *Am J Epidemiol*  
 446 2004;160:814–22.

447 25. Jung SK, Kim K, Tae K, Kong G, Kim MK. The effect of raw vegetable and fruit  
 448 intake on thyroid cancer risk among women: a case-control study in South Korea. *Br J Nutr*  
 449 2013;109:118–28.

450 26. Bosetti C, Negri E, Kolonel L, Ron E, Franceschi S, Preston-Martin S, McTiernan A,  
 451 Dal Maso L, Mark SD, Mabuchi K, Land C, Jin F, Wingren G, Galanti MR, Hallquist A,  
 452 Glatte E, Lund E, Levi F, Linos D, La Vecchia C. A pooled analysis of case-control studies  
 453 of thyroid cancer. VII. Cruciferous and other vegetables (International). *Cancer Causes*  
 454 *Control* 2002;13:765–75.

- 455 27. Braganza MZ, Potischman N, Park Y, Thompson FE, Hollenbeck AR, Kitahara CM.  
456 Adolescent and mid-life diet and subsequent risk of thyroid cancer in the NIH-AARP diet and  
457 health study. *Int J Cancer* 2015;137:2413–23.
- 458 28. Choi W, Kim J. Dietary Factors and the Risk of Thyroid Cancer: A Review. *Clin Nutr*  
459 *Res* 2014;3:75–88.
- 460 29. Franceschi S, Fassina A, Talamini R, Mazzolini A, Vianello S, Bidoli E, Serraino D,  
461 La Vecchia C. Risk factors for thyroid cancer in northern Italy. *Int J Epidemiol* 1989;18:578–  
462 84.
- 463 30. Franceschi S, Levi F, Negri E, Fassina A, La Vecchia C. Diet and thyroid cancer: a  
464 pooled analysis of four European case-control studies. *Int J Cancer* 1991;48:395–8.
- 465 31. Markaki I, Linos D, Linos A. The influence of dietary patterns on the development of  
466 thyroid cancer. *Eur J Cancer* 2003;39:1912–9.
- 467 32. Mack WJ, Preston-Martin S, Bernstein L, Qian D. Lifestyle and other risk factors for  
468 thyroid cancer in Los Angeles County females. *Ann Epidemiol* 2002;12:395–401.
- 469 33. Cléro É, Doyon F, Chungue V, Rachédi F, Boissin JL, Sebbag J, Shan L, Rubino C, de  
470 Vathaire F. Dietary patterns, goitrogenic food, and thyroid cancer: a case-control study in  
471 French Polynesia. *Nutr Cancer* 2012;64:929–36.
- 472 34. Cust AE, Skilton MR, van Bakel MM, Halkjaer J, Olsen A, Agnoli C, Psaltopoulou T,  
473 Buurma E, Sonestedt E, Chirlaque MD, Rinaldi S, Tjønneland A, Jensen MK, Clavel-  
474 Chapelon F, Boutron-Ruault MC, Kaaks R, Nöthlings U, Chloptsios Y, Zylis D, Mattiello A,  
475 Caini S, Ocké MC, van der Schouw YT, Skeie G, Parr CL, Molina-Montes E, Manjer J,  
476 Johansson I, McTaggart A, Key TJ, Bingham S, Riboli E, Slimani N. Total dietary  
477 carbohydrate, sugar, starch and fibre intakes in the European Prospective Investigation into  
478 Cancer and Nutrition. *Eur J Clin Nutr* 2009;63 Suppl 4:S37-60.

- 479 35. Zamora-Ros R, Rinaldi S, Tsilidis KK, Weiderpass E, Boutron-Ruault MC,  
480 Rostgaard-Hansen AL, Tjønneland A, Clavel-Chapelon F, Mesrine S, Katzke VA, Kühn T,  
481 Förster J, Boeing H, Trichopoulou A, Lagiou P, Klinaki E, Masala G, Sieri S, Ricceri F,  
482 Tumino R, Mattiello A, Peeters PH, Bueno-de-Mesquita HB, Engeset D, Skeie G, Argüelles  
483 M, Agudo A, Sánchez MJ, Chirlaque MD, Barricarte A, Chamosa S, Almquist M, Tosovic A,  
484 Hennings J, Sandström M, Schmidt JA, Khaw KT, Wareham NJ, Cross AJ, Slimani N,  
485 Byrnes G, Romieu I, Riboli E, Franceschi S. Energy and macronutrient intake and risk of  
486 differentiated thyroid carcinoma in the European Prospective Investigation into Cancer and  
487 Nutrition study. *Int J Cancer* 2016;138:65–73.
- 488 36. Kaaks R, Lukanova A, Kurzer MS. Obesity, endogenous hormones, and endometrial  
489 cancer risk: a synthetic review. *Cancer Epidemiol Biomarkers Prev* 2002; 11:1531–43.
- 490 37. Bae MJ, Kim SS, Kim WJ, Yi YS, Jeon YK, Kim BH, Lee BJ, Lee JC, Kim IJ, Wang  
491 SG, Kim YK. High prevalence of papillary thyroid cancer in Korean women with insulin  
492 resistance. *Head Neck* 2016;38:66-71.
- 493 38. Rapp K, Schroeder J, Klenk J, Ulmer H, Concin H, Diem G, Oberaigner W, Weiland  
494 SK. Fasting blood glucose and cancer risk in a cohort of more than 140,000 adults in Austria.  
495 *Diabetologia* 2006;49:945–52.
- 496 39. Rinaldi S, Kaaks R, Friedenreich CM, Key TJ, Travis R, Biessy C, Slimani N,  
497 Overvad K, Østergaard JN, Tjønneland A, Olsen A, Mesrine S, Fournier A, Dossus L,  
498 Lukanova A, Johnson T, Boeing H, Vigl M, Trichopoulou A, Benetou V, Trichopoulos D,  
499 Masala G, Krogh V, Tumino R, Ricceri F, Panico S, Bueno-de-Mesquita HB, Monninkhof  
500 EM, May AM, Weiderpass E, Quirós JR, Travier N, Molina-Montes E, Amiano P, Huerta JM,  
501 Ardanaz E, Sund M, Johansson M, Khaw KT, Wareham N, Scalbert A, Gunter MJ, Riboli E,  
502 Romieu I. Physical activity, sex steroid, and growth factor concentrations in pre- and post-

503 menopausal women: A cross-sectional study within the EPIC cohort. *Cancer Causes Control*  
504 2014;25:111–24.

505 40. Horn-Ross PL, Canchola AJ, Ma H, Reynolds P, Bernstein L. Hormonal factors and  
506 the risk of papillary thyroid cancer in the California Teachers Study cohort. *Cancer Epidemiol*  
507 *Biomarkers Prev* 2011;20:1751–9.

508 41. D’Avanzo B, Ron E, La Vecchia C, Francaschi S, Negri E, Ziegler R. Selected  
509 micronutrient intake and thyroid carcinoma risk. *Cancer* 1997;79:2186–92.

510 42. Haselkorn T, Stewart SL, Horn-Ross PL. Why are thyroid cancer rates so high in  
511 Southeast Asian women living in the United States? The Bay Area thyroid cancer study.  
512 *Cancer Epidemiol Biomarkers Prev* 2003;12:144–50.

513 43. Blaszczyk U, Duda-Chodak A. Magnesium: its role in nutrition and carcinogenesis.  
514 *Rocz Panstw Zakl Hig* 2013;64:165–71.

515 44. Zhang LR, Sawka AM, Adams L, Hatfield N, Hung RJ. Vitamin and mineral  
516 supplements and thyroid cancer : a systematic review. *Eur J Cancer Prev* 2013;22:158–68.

517 45. Lee IM, Cook NR, Manson JE, Buring JE, Hennekens CH. Beta-carotene  
518 supplementation and incidence of cancer and cardiovascular disease: the Women’s Health  
519 Study. *J Natl Cancer Inst* 1999;91:2102–6.

520 46. Cook NR, Le IM, Manson JE, Buring JE, Hennekens CH. Effects of beta-carotene  
521 supplementation on cancer incidence by baseline characteristics in the Physicians’ Health  
522 Study (United States). *Cancer Causes Control* 2000;11:617–26.

523 47. O’Grady TJ, Kitahara CM, DiRienzo AG, Gates MA. The association between  
524 selenium and other micronutrients and thyroid cancer incidence in the NIHAARP diet and  
525 health study. *PLoS One* 2014;9:e110886.

526 48. Lerro CC, Koutros S, Andreotti G, Friesen MC, Alavanja MC, Blair A, Hoppin JA,  
527 Sandler DP, Lubin JH, Ma X, Zhang Y, Beane Freeman LE. Organophosphate insecticide use

528 and cancer incidence among spouses of pesticide applicators in the Agricultural Health Study.  
529 *Occup Environ Med* 2015;72:736–44.

530 49. Hord NG, Tang Y, Bryan NS. Food sources of nitrates and nitrites: the physiologic  
531 contact for potential health benefits. *Am J Clin Nutr* 2009;90:1–10.

532 50. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Ingested  
533 Nitrate and Nitrite, and Cyanobacterial Peptide Toxins IARC *Monogr Eval Carcinog Risks*  
534 *Hum* 2010;94:1-412.

535 51. Kilfoy BA, Zhang Y, Park Y, Holford TR, Schatzkin A, Hollenbeck A, Ward MH.  
536 Dietary nitrate and nitrite and the risk of thyroid cancer in the NIH-AARP Diet and Health  
537 Study. *Int J Cancer* 2011;129:160–72.

538 52. Warda MH, Kilfoy BA, Weyer PJ, Anderson KE, Folsom AR, Cerhan JR. Nitrate  
539 Intake and the Risk of Thyroid Cancer and Thyroid Disease. *Epidemiology* 2010;21:389–95.

540 53. Steinmetz KA, Potter JD. Vegetables, fruit, and cancer. II. Mechanisms. *Cancer*  
541 *Causes Control* 1991;2:427–42.

542