

Sperm and sex peptide stimulate aggression in female *Drosophila*

Eleanor Bath^{a,*}, Samuel Bowden^a, Carla Peters^a, Anjali Reddy^a, Joseph A. Tobias^{a,b}, Evan Easton-Calabria^c, Nathalie Seddon^a, Stephen F. Goodwin^d, Stuart Wigby^a

^a Edward Grey Institute
Department of Zoology
University of Oxford
Oxford, UK OX1 3PS

^b Faculty of Natural Sciences
Department of Life Sciences
Imperial College
London, UK SW7 2AZ

^c Department of International Development,
University of Oxford
Oxford, UK OX1 3TB

^d Centre for Neural Circuits and Behaviour
Department of Physiology, Anatomy and Genetics
University of Oxford
Oxford, UK OX1 3SR

*To whom correspondence should be addressed:

Eleanor Bath: eleanor.bath@zoo.ox.ac.uk.

Phone number: +44 1865 271165

Samuel Bowden: sambbowden@gmail.com

Carla Peters: cmariapeters@gmail.com

Anjali Reddy: anjali.reddy@wadh.ox.ac.uk

Joseph Tobias: j.tobias@imperial.ac.uk

Evan Easton-Calabria: evan.easton-calabria@geh.ox.ac.uk

Nathalie Seddon: nathalie.seddon@zoo.ox.ac.uk

Stephen Goodwin: stephen.goodwin@cncb.ox.ac.uk

Stuart Wigby: stuart.wigby@zoo.ox.ac.uk

Classification: Biological Sciences – Evolution

Keywords: female-female aggression, sex peptide, mating costs, trade-off, behaviour

Introductory paragraph

Female aggression towards other females is associated with reproduction in many taxa, and traditionally thought to be related to the protection or provisioning of offspring, such as through increased resource acquisition. However, the underlying reproductive factors causing aggressive behaviour in females remain unknown. Here we show that female aggression in the fruit fly *Drosophila melanogaster* is strongly stimulated by the receipt of sperm at mating, and in part by an associated seminal fluid protein, the sex peptide. We further show that the post-mating increase in female aggression is decoupled from the costs of egg production and from post-mating decreases in sexual receptivity. Our results suggest that male ejaculates can have a surprisingly direct influence on aggression in recipient females. Male ejaculate traits thus influence the female social competitive environment with potentially far-reaching ecological and evolutionary consequences.

Introduction

Female aggression directed towards other females (hereafter “female aggression”) can have important fitness consequences for females and their offspring^{1–4}. In both vertebrate and invertebrate taxa, females act aggressively to acquire resources or mates, defend territories, and protect and provision eggs or offspring^{5–10}. Females are expected to be more flexible than males in their use of aggression, with aggression heightened at particular stages of reproduction or life-history¹¹, and aggression is generally associated with maximising the production or survival of their offspring¹⁰. One observed pattern is an elevation of female aggression after mating, whether it be during offspring development⁶ or after egg production^{12–14}. These increases in aggression typically occur in parallel with other dramatic changes in female behaviour and physiology, including alterations in sexual receptivity, feeding rates, and hormone levels^{4,15–17}.

In many species, male ejaculate components can induce striking changes in female post-mating behaviour and physiology. For example, in insects such as the fruit fly, *Drosophila melanogaster*, male ejaculate components increase ovulation and egg laying, alter immunity, and decrease female receptivity to remating^{15,18}. Thus, ejaculate components are potential candidates for stimulating post-mating increases in female aggression. Male ejaculates could act either as a cue for females to optimally modify their behaviour, indicating upcoming physiological changes that require females to increase their levels of aggression, or could alter female behaviour for the male’s benefit, which could either be in the interests of females, or represent a source of conflict between the sexes¹⁹.

There are two main pathways through which male ejaculates could stimulate post-mating aggression in females. First, by stimulating increased egg production²⁰, ejaculates increase

the demand for resources which could in turn lead to aggressive behaviour (Figure 1.a). Females lay eggs at high rates after mating and require more, or different, nutrients relative to virgin females^{16,17,21–23}, potentially increasing the motivation for females to compete aggressively over food. Another possibility is a more direct stimulation of female aggression by males during mating, with egg production as a separate, but simultaneous, pathway (Figure 1.b). Disentangling these alternatives is required to understand the physiological and molecular pathways that coordinate female aggression with reproduction, and thus unravelling the processes that drive the evolution of female aggressive behaviour.

In this study, we tested whether mating induces changes in female aggression in *D. melanogaster*. In the wild, this species aggregates on rotting fruit^{24,25}, and lab studies have shown that females will engage in fights when food and oviposition sites are limited^{26,27}. Differential reproductive success may therefore result from variation in female ability to aggressively acquire and defend food and oviposition sites. Males and females both display aggressive behaviours in *D. melanogaster* in the laboratory, although there are significant sex differences in the duration of encounters and the types of behaviour observed^{27,28}. There are also differences in aggressive behaviour between mated and virgin females, with mated females fighting for longer and potentially more likely to escalate to higher levels of intensity²⁷. In addition, female *D. melanogaster* display other dramatic post-mating changes in reproductive behaviour, such as reductions in receptivity to remating, increases in feeding, and reductions in sleep¹⁵. These identified post-mating responses are mediated by male seminal fluid proteins, some of which associate with sperm, such as the ‘sex peptide’^{29,30}, suggesting that female aggression could also be influenced by male ejaculate components.

We hypothesize that mating may stimulate aggression in females via two potential pathways, either indirectly through egg production or directly through ejaculate contents (Figure 1). We tested for these pathways by assessing the levels of aggression in females with genetically reduced egg-laying or blocked vitellogenesis, and in females mated to males lacking specific ejaculate components, such as sperm and sex peptide.

Methods

We conducted five experiments to test our main hypotheses about the causes of female aggression:

1. **Effects of mating:** we measured aggression in pairs of mated females, virgin females, and mated vs. virgin females to test for effects of mating on aggression.
2. **Effects of egg production:** to examine whether any increase in female aggression after mating is mediated by the demands and costs of egg-production^{20,31}, we used the *ovo^{D1}* mutation, which blocks oogenesis prior to vitellogenesis³². The presence of the *ovo^{D1}* mutation in females abolishes the mortality costs associated with egg production: if these same costs drive female aggression, any differences in post-mating aggression between mated and virgin females should be abolished in *ovo^{D1}* flies.
3. **Effects of sperm:** to test whether sperm are necessary to stimulate female post-mating aggression, we measured aggression in females mated to spermless *son-of-Tudor* and control males³³.
4. **Effects of the male seminal fluid protein, 'sex peptide', and the female 'sex-peptide-receptor':** to determine whether the seminal fluid protein 'sex peptide' (SP)

is involved in stimulating increased aggression after mating, we compared aggression in females mated to SP-null males with that of control-mated and virgin females. We also used females that lacked the receptor to SP ('sex peptide receptor', SPR) to examine the downstream pathway through which SP could potentially stimulate female aggression.

Fly stocks and culture

All flies were kept and experiments conducted at 25°C on a 12:12 light: dark cycle. We used the Dahomey genetic background as our stock population, and genetic mutations were backcrossed into this background for >5 generations where appropriate.

Production of wild-type females: Wild-type Dahomey females³⁴ were reared in bottles at standardized larval density³⁵. Virgins were collected within 8hrs of eclosion, and sexes were housed separately (females individually) in vials containing standard fly food media³⁶, with no live yeast. For the experiments using mated vs. virgin flies, *ovo*^{D1}, and *SPR* knockout females, Dahomey males were used as mates.

Production of sterile (ovo^{D1}*) females:* To produce sterile *ovo*^{D1} females, we crossed males carrying the dominant *ovo*^{D1} mutation to Dahomey virgin females³⁷. Sterile *ovo*^{D1} females have ovaries that degenerate prior to S5, meaning that vitellogenesis is blocked and females cannot produce eggs³². Dahomey females, from the stock into which the *ovo*^{D1} flies were crossed, were used as controls. We tested the efficacy of the mutation by counting offspring collected from overnight vials following mating. No *ovo*^{D1} females produced offspring, confirming their complete sterility. Previous research has found mixed evidence for the effects of the *ovo*^{D1} mutation on feeding behaviour³⁷, so we tested the effect of the mutation on feeding behaviour in this study (see Results).

146 *Production of spermless males:* To produce males that did not produce sperm, we used
147 *tudor*³³. *tudor* is a maternal effect mutation which prevents germ plasm assembly. Sons of
148 homozygous *tudor* females have no germline and so do not produce sperm^{33,38}. We
149 collected the male offspring of homozygous *tudor* females mated to wild-type Dahomey
150 males as our spermless males. Males of the same genetic background, but with mothers
151 that did not possess the *tudor* mutation, were used as controls.

152 *Production of sex peptide-less males:* To produce males that did not produce sex peptide (SP
153 null) and their controls (SP+), we used stocks created by Liu & Kubli²⁹. SP null males carry
154 one non-functional SP gene, and a deletion ($\Delta 130$) which removes SP²⁹. The SP+ control
155 males are genetically matched with one non-functional SP gene and one functional SP gene,
156 and show wild-type SP expression. To verify the phenotype of SP null and SP+, we counted
157 offspring from a subset of female overnight vials. Females mated to SP+ males produced
158 more eggs and offspring than females mated to SP null males, confirming that our mutants
159 were acting in the expected way (i.e. SP null males were not transferring SP; GLM with
160 quasipoisson distribution: Block 1 – females mated to SP+ males produced more offspring
161 than those mated to SP null males: $X^2_{1, 109} = 82.917$, $P = 0.009$; Block 2 – females mated to
162 SP+ males produced more eggs than those mated to SP null males or virgins: SP+ vs. SP null:
163 $q_{2, 46} = 11.089$, $P < 0.001$; SP+ vs. V: $q_{3, 46} = 9.711$, $P < 0.001$; Supplementary Figure 1). The SP
164 experiment was carried out over two blocks.

165 *Production of sex peptide receptor-less females:* To produce females that lack the sex peptide
166 receptor (hereon, *SPR* deficiency females), we used the genetic deficiency *Df(1)Exel6234*,
167 which deletes the *SPR* gene, and four other genes of unknown function^{39,40}. *SPR* deficiency
168 females do not produce the SPR. As expected, *SPR* deficiency females did not display

reduced receptivity after remating, nor did they show elevated levels of offspring production relative to virgin females (GLM on offspring production: $X^2_{1, 101} = 111.22$, $P < 0.001$; Supplementary Figure 2)³⁹. *SPR* deficiency females have also previously been shown to have slightly shorter copulation duration than wild-type females⁴¹.

Experimental Design

3d post-eclosion virgin females were marked with acrylic paint (red or yellow) on the thorax to allow individual identification²⁷, and returned to individual vials. 24hrs later, females in the “mated” treatments were placed individually with one male, and a single copulation was observed. After each female mated exactly once, males were discarded and all females (i.e. both mated and virgins) were individually transferred to fresh vials, again containing standard media and no live yeast.

The following morning (5d post-eclosion), females were individually placed in vials containing damp cotton wool and no food for 2hrs after which pairs of females were simultaneously aspirated from these vials into the contest arena. The arena was a single well of a twelve-well plate, containing an Eppendorf tube cap filled with standard fly food media and a ~2 μ L drop of yeast paste, providing a limited resource to compete over²⁷. Females were either both mated, both virgin, or – to test the relative competitive ability of mated vs virgin females – one of each. Females were allowed 5 minutes to acclimatise, and then behaviour was recorded for 30 mins using Toshiba Camileo X400 HD video cameras (short sample videos are available in Supplementary Information).

Behavioural analysis

Videos were scored blind with respect to treatment. Fighting behaviours (head-butt, leg “fencing” (later excluded due to difficulties accurately quantifying the behaviour), shove,

192 retreat²⁷) and feeding behaviours were recorded using the program JWatcher + Video⁴².
193 Specifically, we quantified the number and duration of aggressive encounters, the identity
194 of females initiating each encounter, and the outcome of each encounter (i.e. win, lose,
195 draw). An encounter began when one female head-butted or shoved the other female and
196 continued until the females separated or stopped interacting (e.g. were still within touching
197 distance but both resumed eating). We used total duration of aggressive encounters as our
198 primary response variable, which was measured as the sum of time spent in all encounters.
199 We used total duration as it offered the best indicator of overall aggression as it took into
200 account both the number of encounters and the time spent in encounters, providing a more
201 accurate indicator of aggression than number of encounters alone. We also quantified the
202 time females spent in a feeding posture where females were standing on the food cap with
203 their heads tilted down towards the food in a position consistent with feeding.

204 Statistical analysis

205 The response variables (i.e. duration of aggressive encounters and time spent in feeding
206 posture) were continuous and most closely fitted a Gamma distribution. Thus, we used
207 generalized linear models (GLMs) with a Gamma error distribution to test the effects of
208 mating status, egg production status, and various ejaculate components on total aggressive
209 contest duration and time spent in feeding posture. The models in R followed the following
210 format, with the model from the *ovo*^{D1} experiment given as an example:

```
211      glm(Fight duration + 1 ~ Mating status * Egg production, family=Gamma(link="log"))
```

212 Fight duration + 1 was the response variable measured in seconds, while mating status and
213 egg production were the explanatory variables. The explanatory variables differed
214 depending on which experiment was being analysed.

215 For feeding posture analyses, we additionally included proportion of encounters won as an
216 explanatory variable. The models then followed this format:

```
217     glm(Feeding duration + 1 ~ Treatment * Proportion of encounters won,  
218         family=Gamma(link="log"))
```

219 Because Gamma error distributions use a logistic function, we added 1 to all scores of total
220 contest duration prior to transformation to include replicates with scores of 0. We tested for
221 outliers using the Grubbs outlier test and excluded outliers below a threshold of $P = 0.001$.
222 We excluded one outlier in the egg production experiment in the control mated treatment
223 ($G = 6.15$, $P < 0.0001$). Winsorizing the data did not qualitatively alter the results (in the
224 winsorized analysis, mating status had a significant effect on fighting duration ($\text{Dev}_{1, 127} =$
225 13.413 , $P < 0.001$), egg production capability was marginally non-significant ($\text{Dev}_{1, 126} =$
226 1.364 , $P = 0.069$), and there was no interaction between egg production capability and
227 mating status ($\text{Dev}_{1, 125} = 0.124$, $P = 0.583$). We excluded the outlier in our final analysis even
228 though the results did not qualitatively change as it seemed that even when winsorized, the
229 one outlier was still exerting undue leverage on the results. To allow for non-independence
230 of individuals, we used each dyad as the unit of replication in the GLMs, rather than
231 individual flies. For treatments with one mated and one virgin female, we randomly chose
232 one focal individual from each dyad and analysed the proportion of encounters won using a
233 GLM with a quasibinomial distribution. For multiple comparisons within experiments, we
234 used Tukey tests. All models were run in R version 3.0.1⁴³.

235 **Results**

236 **Mating elevates female aggression**

237 We found that aggression was significantly elevated by mating: pairs with two mated
 238 females spent more time fighting over food (GLM: $\text{Dev}_{1, 240} = 26.01$, $P < 0.001$), than pairs
 239 with two virgins (compare MM and VV in Figure 2.a). Moreover, there was no significant
 240 difference in total contest duration between pairs with two mated females and pairs with
 241 one mated and one virgin female (Tukey test: $q_{2, 20} = 0.81$, $P = 0.58$), whereas pairs with at
 242 least one mated female spent more time fighting than pairs with two virgin females (MM vs
 243 VV: $q_{3, 24} = 4.81$, $P = 0.006$; MV vs VV: $q_{2, 24} = 3.69$, $P = 0.015$; overall model: $\text{Dev}_{2, 74} = 11.36$,
 244 $P = 0.004$; Fig. 2.a). We next paired mated females with virgin females to test relative
 245 fighting ability. We found no evidence that mating status influenced who won or lost
 246 aggressive encounters ($\chi^2_{1, 26} = 2.59$, $P = 0.41$; Fig. 2.b). Thus, the presence of one mated
 247 female is sufficient to elevate the overall aggression level of a dyad, but mated females do
 248 not then win significantly more aggressive encounters.

249 **Effects of egg production and male ejaculate components on female aggression**

250 We found that mating-induced increases in aggression were not significantly
 251 different between sterile (*ovo^{D1}*) females and controls (mating status, $\text{Dev}_{1, 126} = 17.32$, $P <$
 252 0.001 ; sterile vs control, $\text{Dev}_{1, 125} = 1.27$, $P = 0.12$; interaction, $\text{Dev}_{1, 124} = 0.25$, $P = 0.49$; Fig.
 253 3.a). Thus, post-mating increases in female aggression are not restricted to females capable
 254 of vitellogenesis and egg laying, and are thus not driven by the energetic demands of egg
 255 production post-vitellogenesis.

256 Females mated to spermless males, which transfer seminal fluid proteins but no sperm to
 257 females³³, did not differ from virgin females in levels of post-mating aggression, and were
 258 significantly less aggressive than control mated females (overall model: $\text{Dev}_{2, 68} = 14.32$, $P <$
 259 0.001 ; control vs. virgin: $q_{3, 24} = 6.8$, $P < 0.001$; control vs. mates of spermless males: $q_{2, 24} =$

5.3, $P = 0.001$; virgin vs. mates of spermless males: $q_{2, 24} = 1.45$, $P = 0.32$; Fig. 3.b). Sperm is therefore required for the mating-induced increase in female aggression.

Aggression in females mated to males that did not transfer the seminal fluid protein 'sex peptide' (SP null males) was significantly higher than in virgin females ($q_{2, 30} = 3.4$, $P = 0.023$) but significantly lower than in mates of males that transferred SP ($q_{2, 30} = 3.05$, $P = 0.04$; overall model $Dev_{2, 127} = 11.26$, $P < 0.001$; Fig. 3.c). Replicate block had no significant effect on total contest duration ($Dev_{1, 126} = 1.225$, $P = 0.131$). These results suggest that SP contributes partially to increased post-mating female aggression, but that SP cannot fully explain the sperm effect on aggression. *SPR* deficiency females did not significantly differ from wild-type females in their aggression response to mating (mating status, $Dev_{1, 101} = 11.53$, $P < 0.001$; interaction mating status x *SPR*, $Dev_{1, 99} = 0.05$, $P = 0.78$; Fig. 3.d). *SPR* deficiency females also spent more time fighting than wild-type females overall, irrespective of their mating status ($Dev_{1, 100} = 12.71$, $P < 0.001$).

Feeding behaviour responses to mating, egg production

In the majority of our experiments, the time females spent in feeding posture was qualitatively similar to patterns of aggressive behaviour: for example, mated females generally spent more time in feeding posture than virgins (Supplementary Figure 3). In the egg production experiment, there was a marginally non-significant trend for more time spent in feeding position in mated females ($Dev_{1, 102} = 1.128$, $P = 0.056$; Supp. Fig. 3.a) but no significant effect of the *ovo^{D1}* mutation ($Dev_{1, 101} = 0.15$, $P = 0.49$) or interaction ($Dev_{1, 99} = 0.31$, $P = 0.97$). Therefore, *ovo^{D1}* females did not differ from control females in time spent in feeding postures, despite spending less time participating in aggressive encounters.

Virgin females and mates of spermless males did not differ in time spent in feeding posture ($q_{2, 24} = 0.05$, $P = 0.97$), but both fed less than controls (control vs spermless: $q_{3, 25} = 4.67$, $P = 0.008$; control vs virgin: $q_{2, 24} = 4.62$, $P = 0.003$; Supp. Fig. 3.b). There was a non-significant trend for reduced feeding in mates of SP null males compared to controls ($q_{2, 35} = 2.502$, $P = 0.086$) and mates of both SP+ and SP null males fed for longer than virgins (SP+ vs virgin, $q_{3, 35} = 6.05$, $P = 0.0004$; SP null vs virgin; $q_{2, 37} = 3.53$, $P = 0.02$; Supp. Fig. 3.c). However, time spent in feeding posture was not elevated in *SPR* deficiency females ($Dev_{1, 100} = 0.14$, $P = 0.55$), and there was no significant effect of mating status ($Dev_{1, 101} = 0.88$, $P = 0.13$; Supp. Fig. 3.d), or interaction ($Dev_{1, 99} = 0.16$, $P = 0.52$), suggesting that the association between time spent engaging in feeding and aggressive behaviours is not obligatory. The proportion of encounters an individual won was not associated with time spent in feeding posture in any experiment (mated vs. virgin experiment: $Dev_{1, 152} = 0.08$, $P = 0.6$ – results in Supp. Fig. 4; *ovo*^{D1} experiment: $Dev_{1, 100} = 0.0006$, $P = 0.97$; *tudor* experiment: $Dev_{1, 69} = 0.61$, $P = 0.13$; SP experiment: $Dev_{1, 112} = 0.09$, $P = 0.53$; SPR experiment: $Dev_{1, 99} = 0.16$, $P = 0.52$; Supplementary Figure 4).

Discussion

Our results show that mating strongly stimulates aggressive behaviour in female fruit flies. Mated females fought for more than twice as long as virgins, and the full increase in aggression requires the receipt of sperm and sex peptide (SP) in the male ejaculate, but not the ability of females to complete vitellogenesis or begin egg laying, nor the presence of the female sex peptide receptor (SPR)³⁹. *ovo*^{D1} females displayed the same increase in aggression after mating as wild-type females, despite lacking the costs of egg production²⁰,

while females that lacked the SPR gene (*SPR* deficiency) spent more time fighting than wild-type females despite producing fewer eggs (Supplementary Results and Figure 2). These results suggest that the costs of egg production and the levels of aggression can be fully decoupled and are thus modulated by divergent pathways (Figure 1.b).

The receipt of SP was necessary for the full increase in post-mating aggression. It should be noted, however, that as we conducted tests of aggression 24 hours after mating, it is likely that only seminal fluid proteins that bind to sperm remained in the female, as other seminal fluid proteins will no longer be present in females at this time^{44,45}. SP is known to bind to sperm, though it is unclear which other seminal fluid proteins also bind to sperm⁴⁶. It is therefore possible that other seminal fluid proteins bound to sperm may also influence female aggression, or proteins not bound to sperm could influence female aggression on a shorter time scale than the 24 hours we measured.

Seminal fluid proteins (including SP) can, under certain conditions, lower female lifetime survival and reproductive output^{47–49}, raising the possibility that ejaculate-stimulated female aggression could contribute to these costs and thus represent an arena of sexual conflict rather than cooperation. However, the net fitness costs and benefits of female aggression, and the role of males in determining this, remains an area ripe for exploration. Surprisingly, females lacking the SPR displayed full increases in post-mating aggression suggesting that SPR or one of the other deleted genes may affect aggressive behaviour, SP may act through alternative pathways to stimulate female aggression, or that SP may act indirectly through association with other seminal proteins⁵⁰. In addition to acting through the SPR, SP stimulates juvenile hormone production in the *corpora allata*, although the mechanism of this stimulation is unknown^{51,52}. Juvenile hormone facilitates

vitellogenesis in *D. melanogaster*, playing a crucial role in reproduction⁵³. The amount of juvenile hormone present in the haemolymph has also been linked to aggression in both sexes in other insect species, such as burying beetles, paper wasps, and cockroaches^{54–56}. It is therefore possible that SP acts to increase female aggression by stimulating juvenile hormone production, though this has yet to be tested.

Male ejaculates could potentially stimulate post-mating aggression through inducing increased feeding, though our data are not fully consistent with this idea. Results from the sperm and SP experiments suggest an association between feeding and aggression, but results from the *ovo^{D1}* and *SPR* deficiency experiments show weaker or no associations (Supplementary Results 4). Future work directly manipulating food consumption would help to establish to what extent the two behaviours are linked. In addition, although using *ovo^{D1}* females allows us to conclude that aggression is not tied to the costs of reproduction (as *ovo^{D1}* females do not suffer the lifespan costs of egg production)^{20,37}, it is possible that eggs do still play a role in inducing female aggression. Further investigation using females that do not possess a germline may help to further clarify the role of egg production in stimulating female aggression.

It is possible that mating results in a reduction of general social tolerance in females towards other individuals, as females are both less receptive to remating with males and more aggressive towards other females. However, the behaviours involved in rejection of males are very different to those involved in incidents of female aggression^{27,57}. In addition, rejection behaviours and aggressive behaviours appear to be activated through different pathways, as rejection behaviours are primarily stimulated through the SPR but our results show that female aggressive behaviours are not.

Our results fit into a broader trend, across taxa, of mating leading to increased female aggression. For example, gestating and lactating mammals and fish display higher levels of aggression towards conspecifics^{6,12,58,59}. However, our results suggest that it is possible to decouple aggression from increased energetic demands of reproduction after mating in *D. melanogaster*. Instead, we have shown that female aggression in *D. melanogaster* is stimulated more directly by male ejaculates, with downstream indirect effects on other females, whereby post-mating aggression affects the wider female competitive environment. For example, the levels of aggression experienced by females may depend not only on the abundance of resources and rivals in their environment, but also upon the seminal characteristics of their mates and mates of their rivals.

These findings have potential implications for a wide range of sexually reproducing animals. In many insect species, females compete aggressively^{54,60} and male accessory gland products induce striking changes in female behaviour and physiology other than female aggression^{18,61,62}. Females of some mammal species also display increased aggression after mating⁶³, and in some mammal species ejaculate components have been shown to influence female physiology^{64,65}. Thus, it is intriguing to speculate that ejaculate-induced female aggression may occur in mammals, analogous to what we have shown here for *D. melanogaster*. However, further studies are required to verify whether ejaculate-mediated effects on female aggression occur in other species. Further key areas of focus for future research include identifying the neuronal mechanisms producing increased post-mating aggression, and understanding the fitness implications for individual females, their competitors and their mates.

Acknowledgements

This study was funded by NERC ([NE/J018937/1](#)) and BBSRC ([BB/K014544/1](#)) fellowships to SW and scholarships from the Rhodes Trust and St John's College, Oxford to EB.

Author contributions

E.B, S.W and N.S conceived of the project. E.B and S.W designed the experiments, with S.F.G providing additional advice on experimental design for later experiments. E.B, S.B, C.P, A.R performed the behavioural experiments and scored the behavioural data. E.B analysed the data and wrote the manuscript. S.W, S.F.G, N.S, E.E-C, and J.A.T discussed the results and contributed to the manuscript.

Data availability statement

Data will be made publicly available on the Oxford University Research Archive (ORA: <https://ora.ox.ac.uk>).

Competing financial interests

The authors declare no competing financial interests.

References

1. Stockley, P. & Bro-Jørgensen, J. Female competition and its evolutionary consequences in mammals. *Biol. Rev. Camb. Philos. Soc.* **86**, 341–366 (2011).
2. Clutton-Brock, T. & Huchard, E. Social competition and its consequences in female mammals. *J. Zool.* **289**, 151–171 (2013).
3. Cain, K. E. & Ketterson, E. D. Competitive females are successful females; phenotype, mechanism and selection in a common songbird. *Behav. Ecol. Sociobiol.* **66**, 241–252 (2012).
4. Dloniak, S. M., French, J. A. & Holekamp, K. E. Rank-related maternal effects of androgens on behaviour in wild spotted hyaenas. *Nature* **440**, 1190–1193 (2006).
5. Huchard, E. & Cowlshaw, G. Female-female aggression around mating: an extra cost of sociality in a multimale primate society. *Behav. Ecol.* **22**, 1003–1011 (2011).
6. Seebacher, F., Ward, A. J. W. & Wilson, R. S. Increased aggression during pregnancy comes at a higher metabolic cost. *J. Exp. Biol.* **216**, 771–776 (2013).
7. Müller, J. K. & Eggert, A.-K. Time-dependent shifts between infanticidal and parental behavior in female burying beetles: a mechanism of indirect mother-offspring recognition. *Behav. Ecol. Sociobiol.* **27**, 11–16 (1990).
8. Elias, D. O., Botero, C. A., Andrade, M. C. B., Mason, A. C. & Kasumovic, M. M. High resource valuation fuels ‘desperado’ fighting tactics in female jumping spiders. *Behav. Ecol.* **21**, 868–875 (2010).
9. Papadopoulos, N. T., Carey, J. R., Liedo, P., Müller, H.-G. & Sentürk, D. Virgin females compete for mates in the male lekking species *Ceratitis capitata*. *Physiol. Entomol.* **34**, 238–245 (2009).
10. Tobias, J. A., Montgomerie, R. & Lyon, B. E. The evolution of female ornaments and weaponry: social selection, sexual selection and ecological competition. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **367**, 2274–93 (2012).
11. Stockley, P. & Campbell, A. Female competition and aggression: interdisciplinary perspectives. *Philos. Trans. R. Soc. B Biol. Sci.* **368**, 20130073–20130073 (2013).
12. Robinson, M. R. & Kruuk, L. E. B. Function of weaponry in females: the use of horns in intrasexual competition for resources in female Soay sheep. *Biol. Lett.* **3**, 651–654 (2007).
13. Eggert, A.-K. & Müller, J. K. Timing of oviposition enables dominant female burying beetles to destroy brood-parasitic young. *Anim. Behav.* **82**, 1227–1233 (2011).
14. Haney, M., DeBold, J. F. & Miczek, K. A. Maternal aggression in mice and rats towards male and female conspecifics. *Agg. Behav.* **15**, 443–453 (1989).
15. Kubli, E. & Bopp, D. Sexual behavior: How sex peptide flips the postmating switch of

- 426 female flies. *Curr. Biol.* **22**, R520–R522 (2012).
- 427 16. Gittleman, J. L. & Thompson, D. Energy Allocation in Mammalian Reproduction.
428 *Integr. Comp. Biol.* **28**, 863–875 (1988).
- 429 17. Wade, G. N. & Schneider, J. E. Metabolic Fuels and Reproduction in Female Mammals.
430 *Neurosci. Biobehav. Rev.* **16**, 235–272 (1992).
- 431 18. Avila, F. W., Sirot, L. K., LaFlamme, B. A., Rubinstein, C. D. & Wolfner, M. F. Insect
432 seminal fluid proteins: identification and function. *Annu. Rev. Entomol.* **56**, 21–40
433 (2011).
- 434 19. Arnqvist, G. & Rowe, L. *Sexual conflict*. (Princeton University Press, 2005).
- 435 20. Sgrò, C. M. & Partridge, L. A delayed wave of death from reproduction in *Drosophila*.
436 *Science* **286**, 2521–2524 (1999).
- 437 21. Wheeler, D. The role of nourishment in oogenesis. *Annu. Rev. Entomol.* **41**, 407–4731
438 (1996).
- 439 22. Walker, S. J., Corrales-Carvajal, V. M. & Ribeiro, C. Postmating Circuitry Modulates
440 Salt Taste Processing to Increase Reproductive Output in *Drosophila*. *Curr. Biol.* **25**,
441 2621–2630 (2015).
- 442 23. Faas, M. M., Melgert, B. N. & De Vos, P. A brief review on how pregnancy and sex
443 hormones interfere with taste and food intake. *Chemosens. Percept.* **3**, 51–56 (2010).
- 444 24. Wertheim, B., Allemand, R., Vet, L. E. M. & Dicke, M. Effects of aggregation
445 pheromone on individual behaviour and food web interactions: a field study on
446 *Drosophila*. *Ecol. Entomol.* **31**, 216–226 (2006).
- 447 25. Markow, T. A. Reproductive behavior of *Drosophila melanogaster* and *Drosophila*
448 *nigrospiracula* in the field and in the laboratory. *J. Comp. Psychol.* **102**, 169–173
449 (1988).
- 450 26. Ueda, A. & Kidokoro, Y. Aggressive behaviours of female *Drosophila melanogaster* are
451 influenced by their social experience and food resources. *Physiol. Entomol.* (2002).
- 452 27. Nilsen, S. P., Chan, Y.-B., Huber, R. & Kravitz, E. A. Gender-selective patterns of
453 aggressive behavior in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. U. S. A.* **101**,
454 12342–12347 (2004).
- 455 28. Vrontou, E., Nilsen, S. P., Demir, E., Kravitz, E. A. & Dickson, B. J. fruitless regulates
456 aggression and dominance in *Drosophila*. *Nat. Neurosci.* **9**, 1469–1471 (2006).
- 457 29. Liu, H. & Kubli, E. Sex-peptide is the molecular basis of the sperm effect in *Drosophila*
458 *melanogaster*. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 9929–9933 (2003).
- 459 30. Peng, J. *et al.* Gradual Release of Sperm Bound Sex-Peptide Controls Female
460 Postmating Behavior in *Drosophila*. *Curr. Biol.* **15**, 207–213 (2005).
- 461 31. Barton Browne, L. in *Regulatory Mechanisms in Insect Feeding* (eds. Chapman, R. F. &

- 462 de Boer, G.) 307–342 (Chapman & Hall, 1995).
- 463 32. Oliver, B., Perrimon, N. & Mahowald, A. P. The *ovo* locus is required for sex-specific
464 germ line maintenance in *Drosophila*. *Genes Dev.* **1**, 913–923 (1987).
- 465 33. Boswell, R. E. & Mahowald, A. P. *tudor*, a gene required for assembly of the germ
466 plasm in *Drosophila melanogaster*. *Cell* **43**, 97–104 (1985).
- 467 34. Wigby, S. & Chapman, T. Female resistance to male harm evolves in response to
468 manipulation of sexual conflict. *Evolution* **58**, 1028–1037 (2004).
- 469 35. Clancy, D. J. & Kennington, W. J. A simple method to achieve consistent larval density
470 in bottle cultures. *Drosoph. Inf. Serv.* **84**, 168–169 (2001).
- 471 36. Lewis, E. A new standard food medium. *Drosoph. Inf. Serv.* **34**, 117–118 (1960).
- 472 37. Barnes, A. I., Wigby, S., Boone, J. M., Partridge, L. & Chapman, T. Feeding, fecundity
473 and lifespan in female *Drosophila melanogaster*. *Proc. R. Soc. B Biol. Sci.* **275**, 1675–83
474 (2008).
- 475 38. Xue, L. & Noll, M. *Drosophila* female sexual behavior induced by sterile males showing
476 copulation complementation. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 3272–3275 (2000).
- 477 39. Yapici, N., Kim, Y.-J., Ribeiro, C. & Dickson, B. J. A receptor that mediates the post-
478 mating switch in *Drosophila* reproductive behaviour. *Nature* **451**, 33–37 (2008).
- 479 40. Dean, R., Perry, J. C., Pizzari, T., Mank, J. E. & Wigby, S. Experimental Evolution of a
480 Novel Sexually Antagonistic Allele. *PLoS Genet.* **8**, e1002917 (2012).
- 481 41. Perry, J. C. *et al.* Experimental evolution under hyper-promiscuity in *Drosophila*
482 *melanogaster*. *BMC Evol. Biol.* **16**, 131 (2016).
- 483 42. Blumenstein, D., Evans, C. & Daniels, J. C. JWatcher. (2006).
- 484 43. R Core Team. R: A Language and Environment for Statistical Computing. (2012).
- 485 44. Sirot, L. K., Buehner, N. A., Fiumera, A. C. & Wolfner, M. F. Seminal fluid protein
486 depletion and replenishment in the fruit fly, *Drosophila melanogaster*: An ELISA-
487 based method for tracking individual ejaculates. *Behav. Ecol. Sociobiol.* **63**, 1505–
488 1513 (2009).
- 489 45. Pilpel, N., Nezer, I., Applebaum, S. W. & Heifetz, Y. Mating-increases trypsin in female
490 *Drosophila* hemolymph. *Insect Biochem. Mol. Biol.* **38**, 320–330 (2008).
- 491 46. Ram, K. R. & Wolfner, M. F. A network of interactions among seminal proteins
492 underlies the long-term postmating response in *Drosophila*. *Proc. Natl. Acad. Sci. U. S.*
493 *A.* **106**, 15384–15389 (2009).
- 494 47. Wigby, S. & Chapman, T. Sex Peptide Causes Mating Costs in Female *Drosophila*
495 *melanogaster*. *Curr. Biol.* **15**, 316–321 (2005).
- 496 48. Chapman, T., Liddle, L. & Kalb, J. Cost of mating in *Drosophila melanogaster* females is
497 mediated by male accessory gland products. *Nature* **373**, 241–244 (1995).

- 498 49. Fricke, C., Bretman, A. & Chapman, T. Female nutritional status determines the
499 magnitude and sign of responses to a male ejaculate signal in *Drosophila*
500 *melanogaster*. *J. Evol. Biol.* **23**, 157–165 (2010).
- 501 50. Haussmann, I., Hemani, Y., Wijesekera, T., Dauwalder, B. & Soller, M. Multiple
502 pathways mediate the sex-peptide-regulated switch in female *Drosophila*
503 reproductive behaviours. *Proc. R. Soc. B Biol. Sci.* **280**, 20131938 (2013).
- 504 51. Fan, Y. *et al.* Common functional elements of *Drosophila melanogaster* seminal
505 peptides involved in reproduction of *Drosophila melanogaster* and *Helicoverpa*
506 *armigera* females. *Insect Biochem. Mol. Biol.* **30**, 805–812 (2000).
- 507 52. Moshitzky, P. *et al.* Sex-peptide activates juvenile hormone biosynthesis in the
508 *Drosophila melanogaster* corpus allatum. *Arch. Insect Biochem. Physiol.* **32**, 363–374
509 (1996).
- 510 53. Soller, M., Bownes, M. & Kubli, E. Control of oocyte maturation in sexually mature
511 *Drosophila* females. *Dev. Biol.* **208**, 337–351 (1999).
- 512 54. Scott, M. P. Resource defense and juvenile hormone: the ‘challenge hypothesis’
513 extended to insects. *Horm. Behav.* **49**, 276–281 (2006).
- 514 55. Kou, R., Chou, S. Y., Chen, S. C. & Huang, Z. Y. Juvenile hormone and the ontogeny of
515 cockroach aggression. *Horm. Behav.* **56**, 332–338 (2009).
- 516 56. Tibbetts, E. A., Vernier, C. & Jinn, J. Juvenile hormone influences precontest
517 assessment behaviour in *Polistes dominulus* paper wasps. *Anim. Behav.* **85**, 1177–
518 1181 (2013).
- 519 57. Connolly, K. & Cook, R. Rejection Responses By Female *Drosophila Melanogaster* :
520 Their Ontogeny, Causality and Effects Upon the Behaviour of the Courting Male.
521 *Behaviour* **44**, 142–165 (1973).
- 522 58. Clutton-Brock, T. *et al.* Infanticide and expulsion of females in a cooperative mammal.
523 *Proc. R. Soc. B Biol. Sci.* **265**, 2291–2295 (1998).
- 524 59. Bowler, C. M., Cushing, B. S. & Carter, C. S. Social factors regulate female-female
525 aggression and affiliation in prairie voles. *Physiol. Behav.* **76**, 559–566 (2002).
- 526 60. Tibbetts, E. A. Resource value and the context dependence of receiver behaviour.
527 *Proc. Biol. Sci.* **275**, 2201–2206 (2008).
- 528 61. Gillott, C. Male accessory gland secretions: Modulators of Female Reproductive
529 Physiology and Behavior. *Annu. Rev. Entomol.* **48**, 163–184 (2003).
- 530 62. McGraw, L. A., Suarez, S. S. & Wolfner, M. F. On a matter of seminal importance.
531 *BioEssays* **37**, 142–147 (2015).
- 532 63. Kapusta, J. & Marchlewska-Koj, A. Interfemale aggression in adult bank voles
533 (*Clethrionomys glareolus*). *Aggress. Behav.* **24**, 53–61 (1998).
- 534 64. Ratto, M. H. *et al.* The nerve of ovulation-inducing factor in semen. *Proc. Natl. Acad.*

535 *Sci.* **109**, 15042–15047 (2012).

536 65. Bogle, O. A., Ratto, M. H. & Adams, G. P. Evidence for the conservation of biological
537 activity of ovulation-inducing factor in seminal plasma. *Reproduction* **142**, 277–283
538 (2011).

539

540

Figure legends

Figure 1: Two proposed pathways for mating-induced female aggression

- a) The transfer of male ejaculates stimulates increased egg production, which could in turn stimulate increased female aggression
- b) Alternatively, the transfer of male ejaculates could stimulate increased female aggression directly, without requiring elevated egg production.

Figure 2:

Mated females spend more time fighting than virgins but do not win more fights. (a) Contest duration in mated, virgin and mixed mating-status female dyads. MM = two mated females (N = 29), MV = one mated female and one virgin female (N = 22), VV = two virgin females (N = 26). (b) Proportion of encounters won by mated females = black bar (n = 45) and virgin females = grey bar (n = 45) in the mixed treatment (MV in 2.a). Model estimate means and standard errors are shown.

Figure 3:

Effects of male ejaculate components, and female egg-production and sex peptide receptor on female total contest duration. a) Total contest duration of mated sterile females (*ovo^{D1}*) and control females. N pairs: Mated sterile (*ovo^{D1}*) = 33, Virgin sterile (*ovo^{D1}*) = 31, Mated control = 31, Virgin control = 33. b) Effect of sperm transfer on female contest duration. N pairs: Control = 26, Spermless = 25, Virgin = 25. The experiment was carried out in two blocks, but results are pooled in this figure. c) Effect of sex peptide transfer on female contest duration. N pairs: SP+ = 39, SP null = 42, Virgin = 49. d) Effect of SPR on female contest duration. N pairs: Mated control = 25, Virgin control = 24, Mated *SPR* deficiency = 29, Virgin *SPR* deficiency = 25. Bars represent means, while error bars indicate standard errors.