

Calibration of a Hybrid Local-Stochastic Volatility Stochastic Rates Model with a Control Variate Particle Method

Andrei Cozma^{*}, Matthieu Mariapragassam^{*†}, and Christoph Reisinger^{*†}

Abstract. We propose a novel and generic calibration technique for four-factor foreign-exchange hybrid local-stochastic volatility models (LSV) with stochastic short rates. We build upon the particle method introduced by Guyon and Henry-Labordère [Nonlinear Option Pricing, Chapter 11, Chapman and Hall, 2013] and combine it with new variance reduction techniques in order to accelerate convergence. We use control variates derived from: a calibrated pure local volatility model, a two-factor Heston-type LSV model (both with deterministic rates), and the stochastic (CIR) short rates. The method can be applied to a large class of hybrid LSV models and is not restricted to our particular choice of the diffusion. However, we address in the paper some specific difficulties arising from the Heston model, notably by a new PDE formulation and finite element solution to bypass the singularities of the density when zero is attainable by the variance. The calibration procedure is performed on market data for the EUR-USD currency pair and has a comparable run-time to the PDE calibration of a two-factor LSV model alone.

Key words. Heston-type local-stochastic volatility models, calibration, particle method, control variate, Fokker–Planck equation

AMS subject classifications. 60H35, 65C05, 65C30, 65N21

1. Introduction. Efficient pricing and hedging of exotic derivatives requires a model which is rich enough to re-price accurately a range of liquidly traded market products. Calibration to vanilla options has been widely documented in the literature since the work of Dupire [17] in the context of local volatility (LV). Nowadays, the exact re-pricing of call options is a must-have standard, and *Local-Stochastic Volatility* (LSV) models are the *state-of-the-art* in many financial institutions. As discussed in Ren et al. [33], Tian et al. [38], Van der Stoep et al. [16] and Guyon and Henry-Labordère [23], LSV models improve the pricing and risk-management performance when compared to pure local volatility or pure stochastic volatility models. The local volatility component allows a perfect calibration to the market prices of vanilla options. At the same time, the stochastic volatility component already provides built-in smiles and skews which give a rough fit, so that a local volatility component – the so-called leverage function – relatively close to one suffices for a perfect calibration. Moreover, they exhibit superior dynamic properties over pure local volatility models.

We focus on a Heston-type LSV model because of the desirable properties of the Cox–Ingersoll–Ross (CIR) process for the variance, such as mean-reversion and non-negativity, and since semi-analytic formulae are available for calls and puts under Heston’s model (see [25]) and can help calibrate the Heston parameters easily. Various sophisticated calibration techniques for the local

^{*}Mathematical Institute, University of Oxford, OX2 6GG, United Kingdom

[†]Oxford-Man Institute of Quantitative Finance, University of Oxford, OX2 6ED, United Kingdom

matthieu.mariapragassam@gmail.com, andrei.s.cozma@gmail.com, christoph.reisinger@maths.ox.ac.uk

The first author gratefully acknowledges financial support from the EPSRC. The second author gratefully acknowledges financial support from the Oxford–Man Institute and BNP Paribas London.

The authors thank an anonymous review editor and two anonymous referees for their very insightful comments.

volatility component are in use in the financial industry, e.g., based on the Monte Carlo particle method in [23] or the PDE-based approach in [33].

In order to improve the pricing and hedging of foreign exchange (FX) options, we furthermore introduce stochastic domestic and foreign short interest rates into the model. Empirical results (see e.g. [41]) have confirmed that for long-dated FX products the effect of interest rate volatility can be as relevant as that of the FX rate volatility. Extensive research has been carried out in the area of option pricing with stochastic volatility and interest rates in the past few years. Van Haastrecht et al. [41] extended the model of Schöbel and Zhu [35] to currency derivatives by including stochastic interest rates, a model that benefits from analytical tractability even in a full correlation setting due to the processes being Gaussian. On the other hand, Ahlip and Rutkowski [3], Grzelak and Oosterlee [22] and Van Haastrecht and Pelsser [42] examined Heston–CIR/Vasicek hybrid models and concluded that they give rise to non-affine models even under a partial correlation structure of the driving Brownian motions and are not analytically tractable.

The resulting 4-factor model complicates the calibration routine due to the higher dimensionality, especially when PDEs are used to find the joint distribution of all factors. A few papers discuss this problem in simpler settings. Deelstra [15] and Clark [10] mainly consider 3-factor hybrid local volatility models and focus on the theoretical rather than the practical aspects of the calibration, whereas Stoep et al. [40] consider an application to a 2-factor hybrid local volatility. In [23], Guyon and Henry-Labordère discuss an application of Monte Carlo-based calibration methods to a 3-factor LSV equity model with stochastic domestic rate and discrete dividends.

The model of Cox et al. [11] is popular when modeling short rates because the (square-root) CIR process admits a unique strong solution, is mean-reverting and analytically tractable. As of late, the non-negativity of the CIR process is considered to be less desirable when modeling short rates. On one hand, central banks have significantly reduced the interest rates since the 2008 financial crisis and it is now commonly accepted that interest rates need not be positive. On the other hand, if interest rates dropped too far below zero, then large amounts of money would be withdrawn from banks and government bonds, putting a severe squeeze on deposits. Hence, we model the domestic and foreign short rates using the shifted CIR (CIR++) process of Brigo and Mercurio [7]. The CIR++ model allows the short rates to become negative and can fit any observed term structure exactly while preserving the analytical tractability of the original model for bonds, caps, swaptions and other basic interest rate products.

We note that the CIR process is sometimes considered difficult to simulate in practice. Moreover, as factor in the Heston model, it leads to singular probability densities for parameter settings where the variance process can hit zero (i.e., if the so-called Feller condition is violated), which cannot be handled easily in the forward Kolmogorov equation by standard numerical methods. In this paper, we address both these issues by tailored schemes, but note that the variance and interest rate processes can be exchanged without significant changes to the main framework and its benefits, e.g. by exponential Ornstein-Uhlenbeck processes for the volatility and Hull-White processes for the rates, both of which are also popular in the industry.

Based on the above considerations, we study the 4-factor hybrid LSV model defined in (2.1) below, which is a Heston-type LSV model with two shifted CIR short-rate processes. We give a rigorous proof of the calibration condition for the leverage function given in [24, Proposition 12.8] for our model specification; see also the condition given in [15] for a 4-factor LSV–2CIR++ model.

We propose a calibration approach which builds on the particle method of [23], and combines it with a novel and efficient variance reduction technique. The main control variate is the two-factor LSV model obtained by assuming that the domestic and foreign rates are deterministic in the original model. In this case, the leverage function is computed by using a deterministic PDE solver. This allows us to take advantage of the efficiency and accuracy of PDE calibration for a low-dimensional model while keeping the complexity for the high-dimensional model under control by Monte Carlo sampling with drastically reduced variance. We find that around 1000 particles are sufficient in practice. Our numerical experiments suggest that this method recovers the calibration speed from the corresponding 2-factor LSV model with deterministic rates defined in (2.2).

As a result of independent interest, we explain how to effectively deal with violation of the Feller condition for the Heston-type LSV Kolmogorov forward equation and numerically solve the PDE using a finite element method with a *Backward Differentiation Formula* (BDF) time-stepping scheme and an appropriate non-Dirichlet boundary condition. To the best of our knowledge, this represents a new approach which complements the literature on the use of ADI schemes [10, 33, 45] to handle the PDE calibration of an LSV model with deterministic rates.

For Heston type models, the CIR variance process can reach zero if the Feller condition is violated, as is often the case in FX markets (we refer to Table 6.5 in [10] for examples on a large range of currency pairs and maturities). As a consequence, the density is singular at the boundary $V = 0$. In [38], the authors propose to reduce the problem by considering $\log(V_t/v_0)$, whereas [10] suggests to refine the mesh near $V = 0$. While these methods alleviate the problem to some extent, we propose to use a different boundary condition as well as a change of variables which results in a bounded solution in a neighbourhood of $V = 0$.

Moreover, a main advantage of the finite element method compared to ADI schemes, besides the greater flexibility in the mesh construction, is that the Dirac delta initial condition can be handled naturally in the weak formulation. This methodology yields an accurate calibration of the Heston-type LSV model with deterministic rates for a broad set of market data.

Finally, we provide empirical evidence that the inclusion of stochastic rates is important for the pricing of some specific exotic derivatives. In particular, in Section 5.4 we consider the pricing problem for a *Target Accrual Redemption Note* (TARN) and a no-touch option. We demonstrate that the impact of stochastic rates is comparable to the difference between pricing a 5-year no-touch option under a LV or LSV model. Other exotics with similar features, not considered here, are Accumulators and *Power Reverse Dual-Currency notes* (PRDC). Moreover, stochastic rates become necessary for any hybrid product which embeds the rates explicitly. Examples are spread options between an FX rate and the Libor rate.

The remainder of this paper is organised as follows. In Section 2, we specify the model and calibration framework and provide a necessary and sufficient condition for a perfect calibration to vanilla quotes. A rigorous proof emphasising the use of local times and possible moment explosions is given in Appendix A. In Section 3, we introduce the particle method used and detail how the control variates for both conditional expectations and standard expectations are constructed. In Section 4, we describe the calibration of the LSV model with deterministic rates using a carefully constructed finite element method. In Section 5, we present numerical results and show that a low number of particles suffices to provide a very good fit to market quotes, which demonstrates the computational efficiency of the method. The impact of stochastic rates for the pricing of a TARN

and no-touch option is presented. Section 6 concludes with a brief discussion.

2. Model definition and calibration. We consider a domestic and a foreign market with stochastic short rates r^d and r^f , and exchange rate S . The spot S_T is associated with the currency pair ccy1ccy2 (following the notations in [10]) and denotes the amount of units of ccy2 (domestic currency) needed to buy one unit of ccy1 (foreign currency) at time T . We denote by D^d and D^f the domestic and foreign discount factors associated with their respective money market accounts,

$$D_t^d = e^{-\int_0^t r_u^d du}, \quad D_t^f = e^{-\int_0^t r_u^f du}.$$

2.1. Models. We assume the existence of a filtered probability space $(\chi, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{Q}^d)$ with a domestic risk-neutral measure \mathbb{Q}^d . For future reference we also define a foreign risk-neutral measure \mathbb{Q}^f . Under \mathbb{Q}^d , S , r^d and r^f follow a system of SDEs

$$(2.1) \quad \begin{cases} \frac{dS_t}{S_t} = (r_t^d - r_t^f) dt + \alpha(S_t, t) \sqrt{V_t} dW_t \\ r_t^d = g_t^d + h^d(t) \\ r_t^f = g_t^f + h^f(t) \\ dg_t^d = \kappa_d (\theta_d - g_t^d) dt + \xi_d \sqrt{g_t^d} dW_t^d \\ dg_t^f = \left(\kappa_f (\theta_f - g_t^f) - \rho_{Sf} \xi_f \sqrt{g_t^f} \alpha(S_t, t) \sqrt{V_t} \right) dt + \xi_f \sqrt{g_t^f} dW_t^f \\ dV_t = \kappa (\theta - V_t) dt + \xi \sqrt{V_t} dW_t^V, \end{cases}$$

where V is the stochastic variance process and the four-dimensional standard Brownian motion (W, W^V, W^d, W^f) has the correlation structure

$$d\langle W_t, W_t^V \rangle = \rho dt, \quad d\langle W_t, W_t^d \rangle = \rho_{Sd} dt, \quad d\langle W_t, W_t^f \rangle = \rho_{Sf} dt, \quad d\langle W_t^d, W_t^f \rangle = \rho_{df} dt,$$

with $\rho, \rho_{Sd}, \rho_{Sf}, \rho_{df} \in (-1, 1)$, the other correlations being zero (and such that the correlation matrix is positive definite), and for given functions $\alpha : \mathbb{R}^+ \times [0, T] \rightarrow \mathbb{R}^+$, $h^{d/f} : [0, T] \rightarrow \mathbb{R}$, and non-negative numbers $\kappa, \theta, \xi, \kappa_d, \theta_d, \xi_d, \kappa_f, \theta_f, \xi_f$, as well as initial values S_0, g_0^d, g_0^f, V_0 .

Let the call option price under model (2.1) for a notional of one unit of ccy1, with strike $K > 0$ and maturity $T > 0$, be

$$C(K, T) = \mathbb{E}^{\mathbb{Q}^d} \left[D_T^d (S_T - K)^+ \right].$$

If the leverage function $\alpha \equiv 1$ in (2.1), we recover a Heston model with shifted CIR domestic and foreign short rates. We will refer to this model as Heston-2CIR++ model. As this model will only be used for intermediate calibration steps, we will make the additional simplification that the interest rate dynamics are independent of the dynamics of the spot FX rate and the variance process, for analytical tractability (see [3]).

We also define two simpler models which we will refer to in the remainder of the article. In both these models, rates are deterministic, $\bar{r}^d(t) = -\partial \ln P^d(0, t) / \partial t$ and $\bar{r}^f(t) = -\partial \ln P^f(0, t) / \partial t$, with $P^{d/f}(0, T)$ the market zero coupon bond prices for the domestic and foreign money market accounts, respectively.

We can then write the related 2-factor Heston-type LSV model with deterministic rates as

$$(2.2) \quad \begin{cases} \frac{dS_t^{2D}}{S_t^{2D}} = (\bar{r}^d(t) - \bar{r}^f(t)) dt + \alpha^{2D}(S_t^{2D}, t) \sqrt{V_t^{2D}} dW_t, & S_0^{2D} = S_0, \\ dV_t^{2D} = \kappa(\theta - V_t^{2D}) dt + \xi \sqrt{V_t^{2D}} dW_t^V, & V_0^{2D} = V_0, \end{cases}$$

for a given function $\alpha^{2D} : \mathbb{R}^+ \times [0, T] \rightarrow \mathbb{R}^+$, and the pure *Local Volatility* (LV) model as

$$(2.3) \quad \frac{dS_t^{LV}}{S_t^{LV}} = (\bar{r}^d(t) - \bar{r}^f(t)) dt + \sigma_{LV}(S_t^{LV}, t) dW_t, \quad S_0^{LV} = S_0,$$

with a given function $\sigma_{LV} : \mathbb{R}^+ \times [0, T] \rightarrow \mathbb{R}^+$.

Note that while the volatility is “local”, i.e., a function of spot FX and time, the short rates are assumed to be a function of time only. We also note for future reference that under the pure LV model (2.3), call prices C_{LV} satisfy the forward Dupire PDE (see [17])

$$(2.4) \quad \frac{\partial C_{LV}}{\partial T} - \frac{1}{2} \sigma_{LV}(K, T)^2 K^2 \frac{\partial^2 C_{LV}}{\partial K^2} + K (\bar{r}^d(T) - \bar{r}^f(T)) \frac{\partial C_{LV}}{\partial K} + \bar{r}^f(T) C_{LV} = 0.$$

2.2. Calibration outline. The purpose of this paper is to calibrate $h^d, h^f, \kappa_d, \kappa_f, \theta_d, \theta_f, \xi_d, \xi_f, \kappa, \theta, \xi, \rho$ and especially α in (2.1). We will use calibration of (2.2) and (2.3) as “stepping stones”. More precisely, the full calibration process consists of the following steps, illustrated in Figure 2.1.

1. Calibration of Heston-2CIR++ parameters:

- (a) calibrate shifted CIR++ model for domestic and foreign short rates separately (Appendix D in [12]);
- (b) calibrate Heston-2CIR++ SV model assuming volatility, domestic and foreign short rates are all independent processes (Appendix F in [12]);

2. Calibrate local volatility assuming time-dependent domestic and foreign short rates (Appendix E in [12]);

3. Calibration of LSV-2CIR++ LSV model:

- (a) calibrate the leverage function of 2-factor Heston LSV model (2.2) assuming time-dependent domestic and foreign rates using local volatility from Step 2 (Section 4);
- (b) calibrate the leverage function of 4-factor LSV-2CIR++ model (2.1) using Heston-2CIR++ parameters from Step 1, the local volatility from Step 2 and leverage function of 2-factor Heston LSV model as a control variate (Sections 3 and 5).

2.3. A necessary and sufficient condition for exact calibration. In the following, we give the main formula that links market call prices, via the Dupire local volatility, to prices under (2.1).

In [23], the following calibration condition is given¹:

$$(2.5) \quad \alpha^2(K, T) = \frac{\mathbb{E}^{\mathbb{Q}^d}[D_T^d | S_T = K]}{\mathbb{E}^{\mathbb{Q}^d}[D_T^d V_T | S_T = K]} \left(\sigma_{LV}(K, T)^2 + \frac{\mathbb{E}^{\mathbb{Q}^d}[Q_T]}{\frac{1}{2} K^2 \frac{\partial^2 C_{LV}}{\partial K^2}} \right),$$

where σ_{LV} is a local volatility as in (2.3), and

$$(2.6) \quad Q_T = D_T^d \left(r_T^f - \bar{r}^f(T) \right) (S_T - K)^+ - K D_T^d \mathbf{1}_{S_T \geq K} \left[\left(r_T^d - \bar{r}^d(T) \right) - \left(r_T^f - \bar{r}^f(T) \right) \right].$$

¹The (equivalent) context there is an equity with stochastic short rate and dividends.




Figure 2.1: Full calibration routine flowchart.

182 *Assumption 1.* α is Lipschitz and uniformly bounded by α_{\max} , $h^{d,f}$ are uniformly bounded and
 183 both the marginal density $\phi(\cdot, T)$ of S_T in (2.1) and $\mathbb{E}^{\mathbb{Q}^d} [D_T^d V_T | S_T = \cdot]$ are continuous.

184 We note that the continuity and positivity in \mathbb{R}_+^3 of the joint density $\psi(\cdot, \cdot, \cdot, T)$ of (S_T, V_T, D_T^d) in
 185 (2.1) is sufficient for $\mathbb{E}^{\mathbb{Q}^d} [D_T^d V_T | S_T = \cdot]$ to be continuous.

186 We also define $\varphi = 2 + \sqrt{2}$, $\zeta = \xi \alpha_{\max}$, and

$$187 \quad (2.7) \quad \begin{cases} T^* = \frac{2}{\sqrt{\varphi^2 \zeta^2 - \kappa^2}} \left[\frac{\pi}{2} + \arctan \left(\frac{\kappa}{\sqrt{\varphi^2 \zeta^2 - \kappa^2}} \right) \right], & \text{if } \kappa < \varphi \zeta, \\ T^* = \infty, & \text{if } \kappa \geq \varphi \zeta, \end{cases}$$

188 which is a lower bound for the explosion time of S_t^2 (see [13]).

189 We prove the following theoretical results in Appendix A:

190 *Proposition 2.* Under Assumption 1, the call price $C(K, T)$ under model (2.1) satisfies

$$191 \quad (2.8) \quad \frac{\partial C(K, T)}{\partial T} - \frac{1}{2} \alpha^2(K, T) K^2 \frac{\mathbb{E}^{\mathbb{Q}^d} [D_T^d V_T | S_T = K]}{\mathbb{E}^{\mathbb{Q}^d} [D_T^d | S_T = K]} \frac{\partial^2 C(K, T)}{\partial K^2} \\$$

$$192 \quad + \mathbb{E}^{\mathbb{Q}^d} [D_T^d r_T^f (S_T - K)^+] - \mathbb{E}^{\mathbb{Q}^d} [D_T^d \mathbf{1}_{S_T \geq K} K (r_T^d - r_T^f)] = 0$$

for any strike $K > 0$ and maturity $T < T^*$, with T^* given by (2.7).

Theorem 3. Under Assumption 1, the call price $C(K, T)$ under model (2.1) matches the price C^{LV} under the local volatility model (2.3) for any strike $K > 0$ and maturity $T < T^*$ only if (2.5) holds for all $K, T > 0$.

If (2.8) has a unique solution, then the condition (2.5) is also sufficient.

Uniqueness of the solution C to the heat equation (2.8) is normally expected under sufficient regularity of the diffusion coefficient and under a growth condition.

The condition (2.5) expresses when a model of the form (2.1) with *exogenously* given α is consistent with market prices, which are expressed through the local volatility function σ_{LV} . We make no claim about the existence of such a model (see also Remark 4 below), and note that α enters (2.5) not only explicitly but also through the \mathbb{Q}^d -expectations. Existence of a calibrated model is linked to the existence of a solution to the McKean-Vlasov equation which results when inserting α *defined endogenously* by (2.5) in terms of σ_{LV} and the model itself into (2.1). In [1], the existence of a short-time solution of the associated Fokker-Planck equation for the density of LSV processes of this type is shown under certain regularity assumptions. The upper bound on the time in [1] is needed to guarantee that the density stays strictly positive from an assumed strictly positive initial condition, and has no direct link to T^* in this paper.

The ratio on the right-hand side of (2.5) accounts for the stochastic volatility; if there is no stochastic volatility (i.e. $V_T = 1$), we recover the formula in [10]. The term Q_T accounts for the stochastic rates and, if rates are deterministic, $Q_T = 0$ and we recover the formula derived in [17],

$$(2.9) \quad \alpha^{2D}(K, T) = \frac{\sigma_{LV}(K, T)}{\sqrt{\mathbb{E}^{\mathbb{Q}^d}[V_T^{2D} | S_T^{2D} = K]}}.$$

At time $T = 0$, from $r_0^d - \bar{r}^d(0) = r_0^f - \bar{r}^f(0) = 0$ we get $\alpha(K, 0) = \sigma_{LV}(K, 0) / \sqrt{v_0}$.

Theorem 3 provides technical conditions for the formula presented in [23], where a formal proof is given without specification of the rates processes. Here, we consider specifically an LSV-2CIR++ model and derive the result rigorously. In Lemma 9 we provide a sufficient condition for the process

$$\int_0^t \mathbf{1}_{S_u \geq K} D_u^d S_u \alpha(S_u, u) \sqrt{V_u} dW_u$$

to be a true martingale up to T^* , which is an important step in the proof of Theorem 3. On the one hand, T^* is a lower bound for the explosion time of the second moment of the discounted spot process $D_t^d S_t$. On the other hand, [5] show that the moment explodes in finite time for the Heston model, a property that is inherited by our Heston-type LSV-2CIR++ model (2.1) as well as the Heston-type LSV-2Hull-White model in [15]. Therefore, the formula may not hold for certain values of the model parameters and for large maturities T . However, in practice, T^* is very large. For instance, from our calibration given in Section 5 we obtain $\kappa = 1.4124$, $\xi = 0.2988$, $\alpha_{\max} = 1.40$, such that $T^* = 28.6$.

Remark 4. A numerical experiment in [23] raises the question of the existence of a calibrated 2-factor LSV model for large ξ (there, $\xi \approx 350\%$ is used to match forward smiles). In this particular case and with the other model parameters kept the same, we find $T^* = 0.20$, which indicates that moment explosions may occur sooner.

3. Fast calibration with a new control variate particle method. In this and the next section, we describe two of the main components of the calibration routine. We recall the calibration condition (2.5), which involves conditional expectations as well as standard expectations, which have to be estimated under model (2.1).

First, we describe the basic particle method for the estimation of these expectations. Then, we present the various control variates, building on intermediary calibration steps, which we use in order to reduce the computational cost of the calibration of α in the 4-factor model (2.1).

Therefore, we require the prior calibration of the interest rate models in (2.1), the LV model (2.3), the Heston-2CIR++ model, and the LSV model (2.2). The calibration of the latter via a PDE is detailed in Section 4, while we refer to Appendices D, E, and F of [12] for the former three.

Equation (2.5) contains the local volatility, which can be obtained from derivatives of market prices from (2.4) by re-arranging it (into Dupire's formula), and explicitly the second derivative of market prices with respect to strike. Different approximation approaches are used in practice, e.g., one writes the formulae in terms of the implied volatility, and uses a smooth parametrisation for the differentiation. Here, we first calibrate a parametrisation of the local volatility model with a fixed-point iteration as in [32, 39] and then use $\frac{\partial^2 C_{LV}}{\partial K^2}$ obtained from the solution of the forward PDE (2.4) with a smoothing scheme (see Appendix E of [12]).

3.1. Calibration by particle method. A calibrated α is implicitly defined by (2.5), where the right-hand side depends on α in a non-linear way through the (conditional) expectations. Formal insertion of the calibration formula into the SDE (2.1) leads to a process where the diffusion coefficient depends on the distribution of the joint process $X_t = (S_t, V_t, r_t^d, r_t^f, D_t^d)$. The process thus falls in the class of McKean-Vlasov processes [28].

The existence and uniqueness of the solution for this McKean-Vlasov SDE are not established theoretically, to the best of our knowledge. From an empirical perspective, in [23] and in Section 11.8 of [24] the authors encountered problems for very high values of ξ ; see Remark 4. In our case, for ξ calibrated to market smiles ($\approx 30\%$) we are able to reach a high accuracy.

The particle method for processes of this type was introduced in [28] and is discussed in Chapter 2, Section 3 of [37]; it was applied to LSV model calibration in [23] and in Section 11.6 of [24].

We define N -sample path approximations of X_t as $(X_t^{i,N})_{i \in \llbracket 1, N \rrbracket} = (S_t^i, V_t^i, r_t^{d,i}, r_t^{f,i}, D_t^{d,i})_{i \in \llbracket 1, N \rrbracket}$ by the $(5 \times N)$ -dimensional SDE

$$(3.1) \quad \begin{cases} \frac{dS_t^i}{S_t^i} = (r_t^{d,i} - r_t^{f,i}) dt + \hat{\alpha}_N \left(S_t^i, t, (X_t^{j,N})_{j \leq N} \right) \sqrt{V_t^i} dW_t^i \\ r_t^{d,i} = g_t^{d,i} + h^d(t) \\ r_t^{f,i} = g_t^{f,i} + h^f(t) \\ dg_t^{d,i} = \kappa_d (\theta_d - g_t^{d,i}) dt + \xi_d \sqrt{g_t^{d,i}} dW_t^{d,i} \\ dg_t^{f,i} = \left(\kappa_f (\theta_f - g_t^{f,i}) - \rho_{Sf} \xi_f \sqrt{g_t^{f,i}} \hat{\alpha}_N \left(S_t^i, t, (X_t^{j,N})_{j \leq N} \right) \sqrt{V_t^i} \right) dt + \xi_f \sqrt{g_t^{f,i}} dW_t^{f,i} \\ dV_t^i = \kappa (\theta - V_t^i) dt + \xi \sqrt{V_t^i} dW_t^{V,i} \\ dD_t^{d,i} = -r_t^{d,i} D_t^{d,i} dt, \end{cases}$$

where $(W_t^i, W_t^{d,i}, W_t^{f,i}, W_t^{V,i})$, $i \in \llbracket 1, N \rrbracket$ are N i.i.d. copies of the four correlated Brownian motions,

263 and $\hat{\alpha}_N$ is an estimator for α based on $(X_t^{i,N})_{i \leq N}$,

$$264 \quad (3.2) \quad \hat{\alpha}_N \left(K, T, (X_t^{i,N})_{i \leq N} \right) = \sqrt{\frac{\sigma_{LV}(K, T)^2}{\hat{p}_N(K, T)} + \frac{\hat{Q}(K, T)}{\frac{1}{2}\hat{p}_N(K, T) K^2 \frac{\partial^2 C_{LV}}{\partial K^2}}},$$

265 with

$$266 \quad (3.3) \quad \hat{Q} = \frac{1}{N} \sum_{i=1}^N Q_T^i \quad \text{and} \quad \hat{p}_N(K, T) = \frac{\sum_{i=1}^N D_T^{d,i} V_T^i \delta_N(S_T^i - K)}{\sum_{i=1}^N D_T^{d,i} \delta_N(S_T^i - K)},$$

267 where \hat{p}_N is an estimator for $\mathbb{E}^{\mathbb{Q}^d}[D_T^d V_T | S_T = K] / \mathbb{E}^{\mathbb{Q}^d}[D_T^d | S_T = K]$, with δ_N a kernel function,
268 and Q_T^i is the i -th sample of Q_T from (2.6) based on $X_T^{i,N}$.

269 The paths of the $5 \times N$ -dimensional process $(X_t^{i,N})_{i \leq N}$ are now entangled due to the dependence
270 on $\hat{\alpha}_N$ in $(X_t^{i,N})_{i \leq N}$. The process can be seen as a system of N interacting particles evolving in a
271 5-dimensional space, where particle i is defined by its position $X_t^{i,N}$. As in [23], we will therefore
272 use the term “particle” instead of “path”. Because of the four driving factors, we will keep referring
273 to this as a 4-factor model in spite of the extra state variable D^d .

274 A central ingredient for proving convergence of the particle method is the chaos propagation
275 property (see Chapter 2, Section 3 of [37]), which is not proven for the present case.

276 **3.2. Variance reduction for the Markovian projection.** Our goal here is to reduce the variance
277 of the estimator \hat{p}_N from (3.3) to be able to use a minimal number of particles.

278 We assume that the 2-factor LSV model (2.2) is perfectly calibrated to market call prices, i.e.
279 that (2.9) is satisfied. Then we will use

$$280 \quad (3.4) \quad \hat{p}_N^{2D}(K, T) = \frac{\sum_{i=1}^N V_T^{2D,i} \delta_N(S_T^{2D,i} - K)}{\sum_{i=1}^N \delta_N(S_T^{2D,i} - K)},$$

281 which is an estimator for

$$282 \quad p^{2D}(K, T) = \mathbb{E}^{\mathbb{Q}^d}[V_T^{2D} | S_T^{2D} = K],$$

283 as a control variate for \hat{p}_N , and p^{2D} will be computed using a PDE solver. The Kolmogorov forward
284 equation for p^{2D} is commonly used for the calibration of LSV models (see [10, 33, 45]), and we
285 propose in Section 4 a new method which is tailored to the specific difficulties associated with
286 density functions in Heston-style models.

287 We thus define a new estimator p_N^* by

$$288 \quad (3.5) \quad p_N^*(K, T) = \hat{p}_N(K, T) + \lambda (\hat{p}_N^{2D}(K, T) - p^{2D}(K, T)).$$

289 The latter has an asymptotically diminishing bias if we assume the particle method to converge in
290 distribution (and neglect the time stepping bias).

291 In order to get a good estimate for the optimal λ , we can rewrite the above estimator as

$$292 \quad p_N^* = \frac{1}{N} \sum_{i=1}^N m_i + \lambda \left(\frac{1}{N} \sum_{i=1}^N m_i^{2D} - p^{2D} \right)$$

293 with

$$294 \quad m_i = \frac{D_T^{d,i} V_T^i \delta_N(S_T^i - K)}{\frac{1}{N} \sum_{i=1}^N D_T^{d,i} \delta_N(S_T^i - K)}, \quad m_i^{2D} = \frac{V_T^{2D,i} \delta_N(S_T^{2D,i} - K)}{\frac{1}{N} \sum_{i=1}^N \delta_N(S_T^{2D,i} - K)},$$

295 which mimics the standard Monte Carlo control variate form. We can think of m_i and m_i^{2D} roughly
 296 as samples of two random variables m and m^{2D} respectively (but note they are not independent,
 297 although for large N the correlation is very low), and for the best variance reduction (see Section
 298 4.1 in [20]), we take

$$299 \quad \lambda = -\frac{\text{Cov}(m, m^{2D})}{\text{Var}(m^{2D})},$$

300 which we can estimate by

$$301 \quad (3.6) \quad \hat{\lambda}_N = -\frac{\sum_{i=1}^N (m_i - \hat{p}_N) (m_i^{2D} - \hat{p}_N^{2D})}{\sum_{i=1}^N (m_i^{2D} - \hat{p}_N^{2D})^2}.$$

302 We recall that the expected variance reduction factor is

$$303 \quad (3.7) \quad \frac{1}{1 - \text{Corr}(m, m^{2D})^2}.$$

304 Hence, if the stochastic rates are not highly volatile, i.e. if ξ_f and ξ_d are small enough, we expect
 305 a very good variance reduction as the correlation between the particles generated by the 4-factor
 306 hybrid LSV model (2.1) and by the 2-factor LSV model (2.2) will be high. Our numerical tests
 307 performed on a model calibrated to recent EUR and USD market data exhibit a correlation between
 308 model (2.1) and (2.2) of 95% up to 1.5 years and 50% around 5 years. Additionally, our stress test
 309 in Subsection 5.3 suggests that even under high volatility regimes for the rate processes, i.e. when
 310 ξ_f and ξ_d are large, the variance reduction brought by this control variate is significant.

311 **3.3. Variance reduction for standard expectations.** Here, we discuss the variance reduction
 312 for the estimator \hat{Q} in (3.3) for $\mathbb{E}^{\mathbb{Q}^d}[Q_T]$, where we repeat

$$313 \quad (3.8) \quad Q_T = \underbrace{D_T^d \left(r_T^f - \bar{r}^f(T) \right) (S_T - K)^+}_{=: X_{1,T}} - K \underbrace{D_T^d \mathbf{1}_{S_T \geq K} \left[\left(r_T^d - \bar{r}^d(T) \right) - \left(r_T^f - \bar{r}^f(T) \right) \right]}_{=: X_{2,T}},$$

314 from (2.6) for the convenience of the reader. This is an estimator for a standard expectation (in
 315 contrast to conditional expectations). We introduce control variates

$$316 \quad Y_{1,T} = D_T^d (S_T - K)^+, \quad Z_{1,T} = \hat{r}_T^f - \bar{r}^f(T)$$

317 for $X_{1,T}$ defined in (3.8), and

$$318 \quad Y_{2,T} = D_T^d \mathbf{1}_{S_T \geq K}, \quad Z_{2,T} = \left(r_T^d - \bar{r}^d(T) \right) - \left(\hat{r}_T^f - \bar{r}^f(T) \right)$$

319 for $X_{2,T}$, and where \hat{r}^f is the foreign rate process without the quanto adjustment.² We know that
 320 if the model (2.1) is perfectly calibrated to call option prices,

$$321 \quad \mathbb{E}^{\mathbb{Q}^d}[Y_{1,T}] = C_{LV}(K, T),$$

$$322 \quad \mathbb{E}^{\mathbb{Q}^d}[Y_{2,T}] = -\frac{\partial C_{LV}}{\partial K}(K, T),$$

²The last quantity is introduced because the expectation of r^f is not analytically available.

323 estimated from market data via a calibrated LV model. The following are also analytically available:

$$\begin{aligned}
324 \quad \zeta_1 &\equiv \mathbb{E}^{\mathbb{Q}^d} [Z_{1,T}] = g_0^f e^{-\kappa_f T} + \theta_f (1 - e^{-\kappa_f T}) + h^f(T) - \bar{r}^f(T), \\
325 \quad \zeta_2 &\equiv \mathbb{E}^{\mathbb{Q}^d} [Z_{2,T}] = g_0^d e^{-\kappa_d T} + \theta_d (1 - e^{-\kappa_d T}) + h^d(T) - \bar{r}^d(T) - \zeta_1.
\end{aligned}$$

326 We denote the Monte Carlo estimators of the corresponding \mathbb{Q}^d -expectations as $\hat{X}_1, \hat{X}_2, \hat{Y}_1, \hat{Y}_2,$
327 \hat{Z}_1, \hat{Z}_2 , respectively, using the same Brownian paths for W, W^V, W^d, W^f in all estimators.
328 We can define a new Monte Carlo estimator Q^* for $\mathbb{E}^{\mathbb{Q}^d} [Q_T]$ as

$$329 \quad (3.9) \quad Q^* = X_1^* - K X_2^*,$$

330 with

$$\begin{aligned}
331 \quad X_1^* &= \hat{X}_1 + \lambda_1 \left(\hat{Y}_1 - C_{LV}(K, T) \right) + \eta_1 \left(\hat{Z}_1 - \zeta_1 \right), \\
332 \quad X_2^* &= \hat{X}_2 + \lambda_2 \left(\hat{Y}_2 + \frac{\partial C_{LV}(K, T)}{\partial K} \right) + \eta_2 \left(\hat{Z}_2 - \zeta_2 \right).
\end{aligned}$$

333 The weights $\lambda_1, \lambda_2, \eta_1, \eta_2$ above are chosen to minimize the variance of Q^* (see [20]).

334 This approach is particularly useful for out-of-the-money options and digital options as the
335 Monte Carlo estimator will exhibit higher variance in these settings.

336 A computational problem arises if there is no particle with $S_T > K$, which happens if K is large
337 and the total number of particles is relatively small (as it will be the case with control variates),
338 since the estimators of $\mathbb{V} [D_T^d \mathbf{1}_{S_T \geq K}]$ and $\mathbb{V} [D_T^d (S_T - K)^+]$ are then zero. In that case, we pick

$$\begin{aligned}
339 \quad \lambda_1 &= -\zeta_1, \quad \eta_1 = -C_{LV}(K, T), \\
340 \quad \lambda_2 &= -\zeta_2, \quad \eta_2 = \frac{\partial C_{LV}(K, T)}{\partial K},
\end{aligned}$$

341 such that both control variates are of the same order of magnitude.

342 Additionally, the estimators of $\lambda, \lambda_1, \lambda_2, \eta_1, \eta_2$ should ideally be pre-calculated with independent
343 Monte Carlo samples. This to ensure that p_N^* and \hat{p}_N have the same expectation.

344 **3.4. Implementation details.** The leverage function α can in principle be computed for any
345 K and T by the estimator (3.2). However, for computational purposes, we defined it in this way
346 on a grid of points and interpolate it from there with cubic splines in spot and piecewise constant
347 in time. We denote by N_T the number of maturities. Then there are $N_T + 1$ volatility “slices” in
348 total such that we denote the m -th time slice $\alpha(\cdot, T_m)$, by α_m , represented numerically as splines
349 with N_S nodes. While having N_S too small will lead to accuracy problems, choosing it too large
350 will make the surface rougher due to over-fitting. We find 25-30 points to provide a good trade-off
351 between accuracy and smoothness. For a given T_m , the leverage function is thus defined on some
352 interval $[S_{\min}^m, S_{\max}^m]$ and is extrapolated constant outside these bounds. Because we need more grid
353 points around the forward value and less around S_{\min}^m and S_{\max}^m , we use a hyperbolic grid (with
354 $\eta = 0.05$, see Appendix C for more details) refined around the forward value

$$355 \quad F_m = S_0 e^{\int_0^{T_m} (\bar{r}^d(t) - \bar{r}^f(t)) dt}, \quad \text{with} \quad S_{\min}^m = F_m e^{-3\sigma_F(T_m)\sqrt{T_m}}, \quad S_{\max}^m = F_m e^{3\sigma_F(T_m)\sqrt{T_m}},$$

where $\sigma_F(T_m)$ is the at-the-money forward market volatility for maturity T_m (interpolated linearly in variance). Each of the grid values can be seen as a parameter and we denote them by $(\alpha_{m,j})_{m \leq N_T, j \leq N_S}$ with the associated spot grid values $(s_{m,j})_{m \leq N_T, j \leq N_S}$.

We now give the calibration algorithm. As previously, we denote the particle system at time T for the model (2.1) by $(S_T^i, V_T^i, r_T^{d,i}, r_T^{f,i}, D_T^{d,i})_{i \leq N}$. Similarly, we denote the 2-factor particle system at time T for the model (2.2) by $(S_T^{2D,i}, V_T^{2D,i})_{i \leq N}$.

We work with a Gaussian kernel

$$\delta_N(x, T) = \frac{e^{-\frac{1}{2} \left(\frac{x}{h_N(T)} \right)^2}}{h_N(T) \sqrt{2\pi}},$$

with a bandwidth given by a Silverman-type rule (see [36])

$$h_N(T) = \eta S_0 \sigma_{LV}(S_0, T) \sqrt{\max(T, T_{\min})} N^{-\frac{1}{5}},$$

where $\eta = 1.5$ and $T_{\min} = 0.25$ in our tests.

The step-by-step calibration is detailed in Algorithm 1.

Our empirical findings suggest that Quasi-Monte Carlo sampling of the random numbers does not provide significant accuracy gains. In order to speed up the computation of the sums involving kernel functions such as

$$\sum_{i=1}^N V_T^{2D,i} \delta_N(S_T^{2D,i} - K),$$

it is advised (see [23]) to sort the particle state vector by spot value and select only the relevant particles that fall inside an interval $[K - \Delta K, K + \Delta K]$, where we choose

$$\Delta K = \sqrt{-2h_N^2(T) \ln(\epsilon \sqrt{2\pi} h_N)},$$

with $\epsilon = 10^{-5}$.

4. Two-factor Heston-type LSV model calibration by PDE . Here, we describe the calibration of the 2-factor sub-model of (2.1) defined in (2.2) by solution of the forward PDE.³

4.1. Transformation and weak formulation. The following is a small variation of the main result in [27], and the proof is therefore omitted. Note that a new non-Dirichlet boundary condition appears at $z = 0$.

Theorem 5. *Define the region $\Omega = \mathbb{R}_+^2$ and assume that the density ϕ (under \mathbb{Q}^d) of the Markovian process (S_t, V_t) started at (S_0, v_0) at time 0 exists and is $C^{2,2,1}(\Omega \times \mathbb{R}_+)$. Then ϕ is the*

³In this section only, we write S, V and α in lieu of S^{2D}, V^{2D} and α^{2D} , respectively, to ease notation.

Algorithm 1 $\alpha(s, T)$ Calibration with control variate particle method

$$\alpha(s, T_1 = 0) = \frac{\sigma_{LV}(s, 0)}{\sqrt{v_0}}$$

for ($m = 1; m \leq N_T; m++$) **do**

generate $(Z, Z_v, Z_d, Z_f)_{i \leq N}$ and $(U)_{i \leq N}$, i.e. $4 \times N$ independent draws from $\mathcal{N}(0, 1)$ and N draws from $\mathcal{U}([0, 1])$, respectively

evolve the 4-factor particle system from T_m to T_{m+1} with QE -Scheme (B.1) where $\alpha(s, [T_m, T_{m+1}[) = \alpha(s, T_m)$

evolve the 2-factor particle system from T_m to T_{m+1} with QE -Scheme (B.1) with pre-computed α^{2D} and using $(Z, Z_v, U)_{i \leq N}$

solve the Dupire forward PDE (2.4) from T_m to T_{m+1} for C_{LV} , $\frac{\partial C_{LV}}{\partial K}$ and $\frac{\partial^2 C_{LV}}{\partial K^2}$

set $T = T_{m+1}$

for ($j = 1; j \leq N_S; j++$) **do**

set $K = s_{m+1, j}$

compute as in (2.9)

$$p^{2D} = \mathbb{E}^{\mathbb{Q}^d} [V_T^{2D} | S_T^{2D} = K] = \left(\frac{\sigma_{LV}(K, T)}{\alpha^{2D}(K, T)} \right)^2$$

compute as in (3.4)

$$\hat{p}_N^{2D}(K, T) = \frac{\sum_{i=1}^N V_T^{2D, i} \delta_N(S_T^{2D, i} - K)}{\sum_{i=1}^N \delta_N(S_T^{2D, i} - K)}$$

compute $\hat{\lambda}_N$ as in (3.6) and

$$p_N^*(K, T) = \frac{\sum_{i=1}^N D_T^{d, i} V_T^i \delta_N(S_T^i - K)}{\sum_{i=1}^N D_T^{d, i} \delta_N(S_T^i - K)} + \hat{\lambda}_N (\hat{p}_N^{2D}(K, T) - p^{2D}(K, T))$$

as in (3.5)

compute as in (3.9)

$$Q^* = (X_1^* - K X_2^*)$$

with

$$X_1^* = \hat{X}_1 + \lambda_1 \left(\hat{Y}_1 - C_{LV}(K, T) \right) + \eta_1 \left(\hat{Z}_1 - \mathbb{E}^{\mathbb{Q}^d} \left[\left(r_T^f - \bar{r}^f(T) \right) \right] \right)$$

$$X_2^* = \hat{X}_2 + \lambda_2 \left(\hat{Y}_2 + \frac{\partial C_{LV}(K, T)}{\partial K} \right) + \eta_2 \left(\hat{Z}_2 - \mathbb{E}^{\mathbb{Q}^d} \left[\left(r_T^d - \bar{r}^d(T) \right) - \left(r_T^f - \bar{r}^f(T) \right) \right] \right)$$

compute

$$\alpha_{m+1, j} = \sqrt{\frac{1}{p_N^*(K, T)} \left(\sigma_{LV}(K, T)^2 + \frac{Q^*}{\frac{1}{2} K^2 \frac{\partial^2 C_{LV}(K, T)}{\partial K^2}} \right)}$$

end for

end for

383 solution to the Kolmogorov forward equation

$$384 \quad (4.1) \quad \begin{cases} \frac{\partial \phi}{\partial t} + (\bar{r}^d(t) - \bar{r}^f(t)) \frac{\partial x \phi}{\partial x} + \frac{\partial \kappa(\theta-z)\phi}{\partial z} \\ \quad - \frac{1}{2} \left(\frac{\partial^2 x^2 \alpha^2(x,t) z \phi}{\partial x^2} + \frac{\partial^2 \xi^2 z \phi}{\partial z^2} + 2 \frac{\partial^2 \rho \xi x \alpha(x,t) z \phi}{\partial x \partial z} \right) = 0, & (x, z) \in \Omega, t > 0, \\ \left(\frac{\xi^2}{2} \frac{\partial z \phi}{\partial z} - \kappa(\theta-z)\phi + \rho \xi z \frac{\partial x \alpha(x,t) \phi}{\partial x} \right) \Big|_{z=0} = 0, & z = 0, x \geq 0, t > 0, \\ \lim_{z \rightarrow \infty} \phi(x, z, t) = \lim_{x \rightarrow \infty} \phi(x, z, t) = \phi(0, z, t) = 0, & (x, z) \in \Omega, t > 0, \\ \lim_{t \rightarrow 0} \phi(x, z, t) = \delta(x - S_0, z - v_0), & (x, z) \in \Omega. \end{cases}$$

385 *Proof.* Similar to the proof of Lemma 4.1 and Theorem 4.1 in [27]. ■

386 The marginal density function of the CIR process at t is (see [11])

$$387 \quad \phi_v(z, t) = c e^{-u-cz} \left(\frac{cz}{u} \right)^{\beta/2} I_\beta(2\sqrt{cu}z),$$

388 with $c = \frac{2\kappa}{(1-e^{-\kappa t})\xi^2}$, $u = cv_0 e^{-\kappa t}$ and $\beta = \frac{2\kappa\theta}{\xi^2} - 1$, where I_β is the modified Bessel function of
389 the first kind of order β . We can write an asymptotic expression for small z using the asymptotic
390 formula for the modified Bessel function found in [2],

$$391 \quad (4.2) \quad \phi_v(z, t) \sim \frac{c^{(\beta+1)} e^{-u-cz}}{\Gamma(\beta+1)} z^\beta,$$

392 such that $\phi_v(z, t)$ diverges for $z = 0^+$ when $2\kappa\theta < \xi^2$. This agrees with the well-known density ϕ_v^∞
393 of the stationary distribution (see [11]) given by

$$394 \quad (4.3) \quad \phi_v^\infty(z) = \lim_{t \rightarrow \infty} \phi_v(z, t) = \frac{\omega^{(\beta+1)} z^\beta e^{-\omega z}}{\Gamma(\beta+1)},$$

395 with $\omega = 2\kappa/\xi^2$. This motivates a scaling of the density ϕ of the form $p = \phi z^{-\beta}$, and to solve a new
396 PDE for p which we define hereafter. By insertion in (4.1) we get the following.

397 **Corollary 6.** For any $\beta \in \mathbb{R}$, $p = \phi z^{-\beta}$ satisfies the initial boundary value problem

$$398 \quad (4.4) \quad \begin{cases} \frac{\partial p}{\partial t} + (\bar{r}^d(t) - \bar{r}^f(t)) \frac{\partial x p}{\partial x} - \beta \kappa p \\ \quad + \frac{\partial \kappa(\theta-z)p}{\partial z} - (\beta+1) \left(\frac{\partial \xi^2 p}{\partial z} + \frac{\partial \rho \xi x \alpha(x,t) p}{\partial x} \right) \\ \quad - \frac{1}{2} z \left[\frac{\partial^2 x^2 \alpha^2(x,t) p}{\partial x^2} + \frac{\partial^2 \xi^2 p}{\partial z^2} + 2 \frac{\partial^2 \rho \xi x \alpha(x,t) p}{\partial x \partial z} \right] = 0, & (x, z) \in \Omega, t > 0, \\ \left[\frac{\xi^2 z}{2} \frac{\partial p}{\partial z} \right]_{z=0} + \kappa z p \Big|_{z=0} + \rho \xi z \frac{\partial x \alpha(x,t) p}{\partial x} \Big|_{z=0} = 0, & z = 0, t > 0, \\ \lim_{z \rightarrow \infty} p(x, z, t) = \lim_{x \rightarrow \infty} p(x, z, t) = p(0, z, t) = 0, & (x, z) \in \Omega, z \neq 0, t > 0, \\ \lim_{t \rightarrow 0} p(x, z, t) = z^{-\beta} \delta(x - S_0, z - v_0), & (x, z) \in \Omega. \end{cases}$$

399 While this PDE is easier to handle numerically, one wants to work with the original density function
400 ϕ for most of the applications. There are two main calculations one would like to achieve: the
401 expected payoff $f(S_T)$ for a given function f ; the Markovian projection $\mathbb{E}[V_T | S_T = K]$.

As ϕ is still intractable for small z and computing $z^\beta p(x, z, t)$ is not numerically feasible, we perform an integration by parts (noticing $\lim_{z \rightarrow 0} z^{\beta+1} p = \lim_{z \rightarrow \infty} z^{\beta+1} p = 0$ since $\beta + 1 > 0$) to obtain, for deterministic rates,

$$(4.5) \quad \mathbb{E}[D_T f(S_T)] = -D_T \int_0^\infty f(x) \int_0^\infty \frac{z^{\beta+1}}{\beta+1} \frac{\partial p(x, z, T)}{\partial z} dz dx$$

and

$$(4.6) \quad \mathbb{E}[V_T | S_T = K] = -(\beta + 1) \frac{\int_0^\infty z^{\beta+1} p(K, z, T) dz}{\int_0^\infty z^{\beta+1} \frac{\partial p(K, z, T)}{\partial z} dz}.$$

4.2. Finite element method with a two-step BDF time scheme. We combine a finite element approximation in space with a *Backward Differentiation Formula* (BDF) scheme in time, since Crank-Nicholson time-stepping or ADI schemes can give rise to instabilities for Dirac initial data (see [30, 44]; we refer to [6] for a stability analysis and to [18] for some financial applications of BDF schemes).

Equation (4.4) can be written as

$$(4.7) \quad \frac{\partial p}{\partial t} - z \nabla \cdot \mathbf{u} + \nabla \cdot (\mathbf{b} - (\beta + 1) \mathbf{w}) + cp = 0,$$

$$\mathbf{u} = \frac{1}{2} \begin{bmatrix} \frac{\partial x^2 \alpha^2(x, t) p}{\partial x} + \frac{\partial \rho \xi x \alpha(x, t) p}{\partial z} \\ \frac{\partial \xi^2 p}{\partial z} + \frac{\partial \rho \xi x \alpha(x, t) p}{\partial x} \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} (\bar{r}^d(t) - \bar{r}^f(t))x \\ \kappa(\theta - z) \end{bmatrix}, \quad \mathbf{w} = \begin{bmatrix} \rho \xi x \alpha(x, t) p \\ \xi^2 p \end{bmatrix}, \quad c = -\beta \kappa.$$

Denote by $\Gamma_R = \{(x, z) \in \partial\Omega : z = 0\}$ the subset of the boundary of Ω with Robin boundary condition. We derive a weak formulation in the usual way (see, e.g., [31]), i.e., we multiply the PDE by a test function $v \in H^1(\Omega)$, integrate over Ω , using the divergence theorem and boundary conditions, to obtain the weak form of (4.7),

$$\int_\Omega \frac{\partial p}{\partial t} v d\Omega + a(p, v) = 0,$$

with the bi-linear form

$$a(p, v) = \int_\Omega \mathbf{u} \cdot \nabla(zv) + (\nabla \cdot (\mathbf{b} - (\beta + 1) \mathbf{w}) + cp) v d\Omega - \int_{\Gamma_R} \left(\kappa zp + \frac{1}{2} z \frac{\partial \rho \xi x \alpha(x, t) p}{\partial x} \right) v d\Gamma_R,$$

where the last term contains the new boundary condition.

Let us define a uniform time mesh with $t_m = m\Delta_t$, $m \in \llbracket 0, M \rrbracket$. We denote $p_m = p(\cdot, \cdot, t_m)$, in which case the BDF scheme can be written as

$$(4.26) \quad \int_\Omega \left(p_{m+2} - \frac{4}{3} p_{m+1} + \frac{1}{3} p_m \right) v d\Omega + \frac{2}{3} \Delta_t a(p_{m+2}, v) = 0, \quad m = 0, \dots, M-2,$$

where the first time step is divided into two standard fully implicit time steps (see [19]). Then, for the first time step, using the Dirac delta initial condition,

$$(4.29) \quad \int_\Omega p_1 v d\Omega + \Delta_t a(p_1, v) = \int_\Omega p_0 v d\Omega = v_0^{-\beta} v(S_0, v_0).$$





Figure 4.1: \mathbb{P}_2 element with 6 degrees of freedom **Figure 4.2:** Finite element triangular mesh refined around $x = S_0$ and $z = 0$

Note that we initially only assumed $v \in H^1 \not\subset C$, and therefore the operation above with the Dirac delta is not defined for all such v . However, we will next use continuous basis functions. If (S_0, v_0) coincides with a mesh point, this is equivalent to solving a linear system where the right-hand side vector is $v_0^{-\beta}$ for the source point node and zero otherwise.

The PDE solution is approximated by a conforming finite element method with \mathbb{P}_2 elements, i.e., a polynomial of order two on a triangle cell. Each triangle is characterised by 6 local degrees of freedom (nodes) as displayed in Figure 4.1 (see [29] for details). We describe the mesh construction in detail in Appendix C. An example of a thus generated mesh with 30 spot steps and 30 variance steps is illustrated in Figure 4.2.

In order to see the improvement due to the transformed PDE (4.4) for p , we plot both $z^{-\beta}\phi$ and p for a pure Heston model with

$$r = 3\%, \quad q = 1\%, \quad \kappa = 1, \quad \theta = v_0 = 0.04, \quad \rho = -0.3, \quad \xi = 0.5, \quad t = 1,$$

which corresponds to a Feller ratio of 0.32. We use 100 time steps as well as 30 spot steps and 30 variance steps. The solution for $p = z^{-\beta}\phi$ computed with no change of variables is presented in Figure 4.3, where we notice significant numerical instabilities. The tranformed PDE is solved for the same problem and p is also plotted in Figure 4.3.

4.3. Calibration algorithm. The calibration of the 2-factor LSV model (2.2) is performed by finding the leverage function α defined in (2.9). We compute $\mathbb{E}^\mathbb{Q}[V_T \mid S_T = K]$ from (4.6) with the solution p of the dampened PDE (4.4). Both integrals can be computed by double adaptive Clenshaw-Curtis quadrature rules to handle singularities properly when T is small (see [21]).

Furthermore, for very small or very large values of K , both the numerator and denominator will be very small. So we define a smooth extrapolation rule by

$$\mathbb{E}^\mathbb{Q}[V_T \mid S_T = K] \approx \frac{(\beta + 1) \left(\int_0^\infty z^{\beta+1} p(K, z, T) dz + \epsilon \left(v_0 e^{-\kappa T} + (\theta - v_0) (1 - e^{-\kappa T}) \right) \right)}{\epsilon (\beta + 1) - \int_0^\infty z^{\beta+1} \frac{\partial p(K, z, T)}{\partial z} dz},$$

where we pick $\epsilon = 10^{-14}$ in our numerical tests.

The calibration will be done forward in time. Denote by $(\Delta T(i))_{i \leq N_T}$ the interval lengths between maturities and by N_T the number of maturities.




Figure 4.3: $p = z^{-\beta} \phi$ using ϕ computed with the standard (left) and change of variables PDE (right)

Algorithm 2 $\alpha(s, T)$ calibration with the dampened Kolmogorov forward PDE

$$\alpha(s, 0) = \frac{\sigma_{LV}(s, 0)}{\sqrt{v_0}}$$

$T = 0$

for ($i = 1; i \leq N_T; i++$) **do**

solve (4.4) for p on $[T, T + \Delta T]$ with $\alpha(s, [T, T + \Delta T(i)]) = \alpha(s, T)$

for ($j = 1; j \leq N_S; j++$) **do**

$$E_{V_T} = -(\beta + 1) \frac{\int_0^\infty z^{\beta+1} p(s_{i,j}, z, T + \Delta T(i)) dz}{\int_0^\infty z^{\beta+1} \frac{\partial p(s_{i,j}, z, T + \Delta T(i))}{\partial z} dz}$$

$$\alpha_{i,j} = \frac{\sigma_{LV}(s_{i,j}, T + \Delta T(i))}{\sqrt{E_{V_T}}}$$

end for

$T = T + \Delta T(i)$

end for

457 The leverage function α is again defined by splines as detailed in Section 3.1. This approach allows
458 us to compute $\mathbb{E}^\mathbb{Q}[V_T | S_T = K]$ only on the nodes, reducing the computational time considerably.
459 In the calibration routine, we use forward constant interpolation of the leverage function between
460 maturities to handle the non-linearity of the problem. We can then write the calibration procedure
461 as in Algorithm 2.

462 In the test, we use the Heston parameters calibrated in Appendix F of [12] for the Heston–
463 2CIR++ model, i.e.,

464 $v_0 = 0.0094, \quad \theta = 0.0137, \quad \kappa = 1.4124, \quad \rho = -0.1194, \quad \xi = 0.2988,$

465 where the Feller ratio is

466
$$\frac{2\kappa\theta}{\xi^2} \approx 0.4335 < 1,$$

467 which violates the Feller condition.

468 The calibrated leverage function α is plotted in Figure 4.4. For the solution of the forward PDE
 469 between maturities, we use a BDF scheme with constant stepsize and find that 50 time steps per
 470 year and a 80×80 spot-variance finite element mesh give very accurate results.




Figure 4.4: Calibrated leverage function α for (left) the 2-factor LSV model (2.2) and (right) the 4-factor LSV model (2.1).

471 **5. Four-factor Heston-type LSV-2CIR++ model calibration.** In this section, we give the
 472 results for the calibration of the main model (2.1) and test the efficiency of the algorithm.

473 We use vanilla options implied volatility data from Bloomberg from the 18/03/2016 for the
 474 currency pair EURUSD, namely, 10D-Put, 25D-Put, 50D, 25D-Call, 10D-Call, for the maturities⁴
 475 3W, 1M, 2M, 3M, 6M, 1Y, 1Y6M, 2Y, 3Y, 5Y.

476 We use historical correlations estimated in [14] from weekly time series data from 2012–2014,

477
$$\rho_{Sd} = -0.3024, \quad \rho_{Sf} = 0.1226, \quad \rho_{df} = 0.6293.$$

478 We assume that both CIR++ processes are calibrated under their own risk-neutral measure as in
 479 Appendix D in [12], and that the Heston-2CIR++ model is calibrated as in Appendix F of [12] for
 480 the Heston-2CIR++ model. The parameters are as follows

481
$$\begin{cases} v_0 = 0.0094, & \theta = 0.0137, & \kappa = 1.4124, & \rho = -0.1194, & \xi = 0.2988, \\ g_0^d = 0.0001, & \theta_d = 0.5469, & \kappa_d = 0.0837, & \rho_{Sd} = 0, & \xi_d = 0.0274, \\ g_0^f = 0.0001, & \theta_f = 1.1656, & \kappa_f = 0.0110, & \rho_{Sf} = 0, & \xi_f = 0.0370. \end{cases}$$

⁴We skip the 7Y and 10Y quotes as they were not liquid enough.

In order to approximate the particle system (3.1), we use an extension of the QE-scheme from [4] to model (2.1). A full description of the time marching scheme is provided in Appendix B.

5.1. Calibration results and efficiency. For a first illustration of the model fit and the improvement through the control variates, we calibrate the 4-factor model with 800 particles with and without control variates. The associated leverage function is plotted in Figure 4.4 (right).

We then plot, in Figure 5.1, the model implied volatility slices for 3M, 1Y, 2Y, and 5Y. The figure shows a significantly improved fit due to the control variates. We will analyse the accuracy and convergence in detail in Subsection 5.2.




Figure 5.1: Calibration fit with 800 particles for the 4-factor LSV model with and without control variates

As regards computational time, the control variate particle method with 20000 particles, which are enough for a reasonably converged solution (see Subsection 5.2), took approximately 10% of the overall time spent in the PDE calibration of the LSV(2D) model. Most practitioners are familiar with the computational time required to calibrate a 2D LSV model by a forward PDE, a rough estimate being below one second for short- to mid-term expiries (5Y). The extra cost to calibrate the full 4D LSV-2CIR++ model is almost negligible with control variates, while the same accuracy without control variates requires more than 60 times the cost of the 2D PDE solver.

5.2. Variance and error reduction. In this subsection, we compare the results with control variates (CV) to those with the plain particle method (No-CV) as a function of the number of particles. The error measure we use is the absolute error in volatility (in % units). For instance, a maximum error (taken over all quoted deltas and maturities) of 0.03% for a 20% market volatility

signifies that the calibrated volatility can be $20.00\% \pm 0.03\%$ in the worst case scenario.

We use the calibration routine described in Subsection 3.1 with and without control variates for 160, 800, 4 000, 20 000, 100 000, 500 000 and 2 500 000 particles.

The results are presented in Figure 5.2. On the basis that higher correlations between spot-rates and rate-rate will make our control variates more efficient, we also display in Figure 5.2 the calibration errors with no correlations between spot-rates and rate-rate as a presumed worst-case.




Figure 5.2: Error convergence with and without control variates, either with historical $\rho_{Sd}, \rho_{Sf}, \rho_{df}$ correlations (left) or $\rho_{Sd} = 0, \rho_{Sf} = 0, \rho_{df} = 0$ correlations (right). Error is computed on all quoted deltas and maturities (log-scale)

We infer from the data in Figure 5.2 that the use of control variates greatly improves the general calibration routine. Convergence in both the maximum error (0.032%) and in the average error (0.012%) is reached (i.e., the error from here on is dominated by other sources, such as the time discretisation error) for 4 000 particles when using control variates, while the plain particle method (without variance reduction) only reaches the same accuracy with 2 500 000 particles. We infer that the control variates give a 625-fold speed-up.

Moreover, from careful data analysis of Figure 5.2, we find a convergence rate of 0.3 in the number of particles for the plain particle method, for both the average and maximum error. The addition of the control variates preserves the convergence rate but reduces the absolute size of the error significantly.

We note that the error can be further reduced by increasing the number of time steps per year. We used 250 time steps (such that, on average, there is one time step per open day) as it already yields a very accurate calibration at a reasonable computational cost.

Part of the error reduction is due to the conditional control variate described in Subsection 3.2 which can provide very good results for short-term horizons (the other part being due to the control variates for standard expectations). In order to analyse this further, we plot the variance reduction factor from (3.7), estimated with 500 000 particles, as a function of time in Figure 5.3. We estimate a trend line $1 + \frac{C}{T^{1.6}}$, for a given constant C , and thus a very good variance reduction for short-term options. For longer terms, this still yields a good variance reduction factor of 5.2 for the 3Y maturity and 2.34 for 5Y.

In addition, the other two control variates presented in Subsection 3.3 help control the short- to long-term behaviour as well. We plot both variance reduction factors for X_1^* and X_2^* in Figure 5.3. They seem to reach a steady state for maturities around 1.6 and 6.5, respectively. We note that we displayed here average variance reduction values over all strikes, while X_1^* was found to provide significant variance reduction for small strikes.




Figure 5.3: Variance reduction factor as a function of time for the conditional expectation estimator p_N^* (left, using a log-log scale) and the standard expectations estimators X_1^* and X_2^* (right, using a log scale)

5.3. Stress scenario. The last calibration result we present is for an extremely stressed set of rates parameters. This will serve as a robustness test of the control-variate particle method for very volatile short rate processes. In order to perform the test, we multiply the calibrated ξ_d and ξ_f by 20 as well as divide θ_d and θ_f by 20. This leads to strongly violated Feller conditions and very high volatilities for the two CIR++ rate processes. We also set $\rho_{Sd} = 0, \rho_{Sf} = 0, \rho_{df} = 0$ as this has shown to be more challenging for our method (as it makes the control variates less effective). The stress scenario parameter values are

$$\begin{cases} v_0 = 0.0094, & \theta = 0.0137, & \kappa = 1.4124, & \rho = -0.1194, & \xi = 0.2988, \\ g_0^d = 0.0001, & \theta_d = \mathbf{0.0273}, & \kappa_d = 0.0837, & \rho_{Sd} = 0, & \xi_d = \mathbf{0.5480}, \\ g_0^f = 0.0001, & \theta_f = \mathbf{0.0582}, & \kappa_f = 0.0110, & \rho_{Sf} = 0, & \xi_f = \mathbf{0.7400}. \end{cases}$$

We emphasise that in practice, the volatility parameters $\xi_{d/f}$ are rarely above 0.06. We refer the reader to [8] for more details. A calibration summary for the average error in absolute volatility is displayed in Table 1. In this stress scenario, the calibration via control-variate particle method reaches an error of 0.0165% for 20 000 particles, whereas 2 500 000 particles are required with the plain particle method (without control variates) to reach the same accuracy. Hence, the control-variate particle method shows a consistent improvement over the plain particle method even under stress scenarios. The conditional control variate for p_N^* yields a variance reduction factor of almost 2 for the last maturity (5Y), similar to the correlated case.

5.4. Impact of stochastic rates. In this subsection, we discuss the pricing of more exotic products, namely a no-touch option and a target accrual redemption note (TARN); see Chapter 8

N	4 000	20 000	100 000	500 000	2 500 000
Plain particle method	0.0855%	0.0440%	0.0320%	0.0255%	0.0160%
Control variate particle method	0.0312%	0.0165%	0.0133%	0.0133%	0.0133%

Table 1: Average error in absolute volatility (% unit) for a high-volatility stress scenario on the rate processes. N is the number of particles.

in [10] and Section 2.2 in [46] respectively for a discussion of these products. Specifically, we assess the impact of stochastic rates on products embedding knock-out features with mid- to long-term expiries.

No-touches. The foreign no-touch up option pays one EUR at maturity if the exchange rate has not breached an upper barrier during the product lifespan. The payout under an arbitrage-free model with foreign risk-neutral measure \mathbb{Q}^f , for a premium expressed in foreign units (EUR), a notional N_{EUR} in foreign units (EUR) and for a maturity T is

$$N_{\text{EUR}} \mathbb{E}^{\mathbb{Q}^f} \left[D_T^f \mathbf{1}_{M_T < B_1} \right],$$

where $M_t = \sup_{0 \leq u \leq t} S_u$ is the running-maximum of the spot S and B_1 is the upper barrier. For the tests, we pick

$$T = 5.0, \quad B_1 = 1.4 \times S_0.$$

TARNs. The specification of the TARN is here as follows: the buyer receives the forward value $K - S_{t_i}$ at fixing date t_i if $K > S_{t_i}$; the buyer has to pay $S_{t_i} - K$ if $K < S_{t_i}$. At each fixing date, if the amount received is positive, the accrued value is increased by the paid amount. If at some point in the deal life-cycle, the accrued amount breaches a target H_{target} , the deal is terminated early. To protect the buyer, an additional knock-out barrier redeems the deal early if the spot fixes at or above an upper barrier B_2 . The payout under an arbitrage-free model with domestic risk-neutral measure \mathbb{Q}^d , for a premium expressed in domestic units (USD) and a notional N_{EUR} in foreign units (EUR), for a maturity T is

$$N_{\text{EUR}} \mathbb{E}^{\mathbb{Q}^d} \left[D_T^d \sum_{i=0}^{n_f} \mathbf{1}_{\tau > t_i} (K - S_{t_i}) \right],$$

where τ is the (early) redemption date that is either triggered by the accrual breaching the target H_{target} or the spot S breaching the upper barrier B_2 . In this test, we pick monthly fixings and

$$T = 5.0, \quad n_f = 12 \times 5, \quad K = 1.09 \times S_0, \quad B_2 = 1.45 \times S_0, \quad H_{\text{target}} = 6 \times (K - S_0).$$

We display in Table 2 the prices computed with 1 048 575 quasi-Monte Carlo paths (with Sobol sequences and Brownian bridge construction from [9]) and 365 time steps per year for the foreign no-touch and TARN contracts under the LV, LSV and LSV-2CIR++ models. The running maximum is sampled with the Brownian bridge technique, as described in Chapter 6 of [20].

Model	NPV - No-touch	NPV - TARN
LV	72.42%	29.16%
LSV	74.59%	14.69%
LSV-2CIR++	73.60%	17.50%

Table 2: Monte Carlo NPVs, in % of N_{EUR} and N_{USD} respectively, for the 5Y foreign no-touch and target accrual redemption note

We infer from the data in Table 2 a relative difference of 1.35% in price for the foreign no-touch and -16.07% for the TARN between the LSV and LSV-2CIR++ models, which confirms that adding stochastic rates has a significant impact even for a 5Y contract.

Interestingly, in this setup, the price of the foreign no-touch option under the LSV-2CIR++ lies in between the LV and LSV models. It is common to observe a market price that lies between the LV model price and a Heston-type LSV model price, when the Heston parameters have been calibrated to the same vanilla options. Practitioners therefore introduce a mixing factor to control the amount of stochastic volatility of the LSV model and to manually match the no-touch options quotes (see [10] for details). The introduction of stochastic rates seems to achieve a similar behaviour, at least in this example.

6. Conclusion. In this paper, we have provided a new and numerically effective method to calibrate a 4-factor LSV model to vanilla options. In our numerical tests with market data, we managed to achieve an approximate 625-fold speed-up for the calibration using control variates, as compared to the plain particle method. We have shown that a high accuracy can be obtained with as few as 4000 particles (with a maximum error in absolute volatility of 0.03%), and we were able to get a good fit with only 800 particles (with a maximum error in absolute volatility of 0.05%).

Using the calibrated leverage function from this paper, we showed that the addition of stochastic rates has a significant impact on structured products, even more so when barrier features and coupon detachments are combined for longer-dated contracts. Stochastic rates become necessary in the modelling if one wishes to price hybrid products where the rates appear explicitly (for instance, a spread option on the FX performance and the Libor rate). One could use a second factor in the CIR++ processes to improve the fit to caps and use the method presented in this article to calibrate the leverage function.

REFERENCES

- [1] F. ABERGEL AND R. TACHET, *A nonlinear partial integro-differential equation from mathematical finance*, Discrete and Continuous Dynamical Systems – Series A, 27 (2010), pp. 907–917.
- [2] M. ABRAMOWITZ, *Handbook of Mathematical Functions, with Formulas, Graphs, and Mathematical Tables*, Dover Publications, 1974.
- [3] R. AHLIP AND M. RUTKOWSKI, *Pricing of foreign exchange options under the Heston stochastic volatility model and CIR interest rates*, Quantitative Finance, 13 (2013), pp. 955–966.
- [4] L. ANDERSEN, *Simple and efficient simulation of the Heston stochastic volatility model*, The Journal of Computational Finance, 11 (2008), pp. 1–42.
- [5] L. ANDERSEN AND V. PITERBARG, *Moment explosions in stochastic volatility models*, Finance and Stochastics,

- 11 (2007), pp. 29–50.
- [6] O. BOKANOWSKI, A. PICARELLI, AND C. REISINGER, *Stability results for second order backward differentiation schemes for parabolic Hamilton–Jacobi–Bellman equations*, 2017. HAL preprint, hal-01628040.
- [7] D. BRIGO AND F. MERCURIO, *A deterministic-shift extension of analytically-tractable and time-homogeneous short-rate models*, Finance and Stochastics, 5 (2001), pp. 369–387.
- [8] D. BRIGO AND F. MERCURIO, *Interest Rate Models: Theory and Practice*, Springer, 2006.
- [9] R. E. CAFLISCH, W. MOROKOFF, AND A. B. OWEN, *Valuation of mortgage-backed securities using Brownian bridges to reduce effective dimension*, Journal of Computational Finance, 1 (1997), pp. 27–46.
- [10] I. J. CLARK, *Foreign Exchange Option Pricing: A Practitioner’s Guide*, John Wiley & Sons, 2010.
- [11] J. COX, J. INGERSOLL, AND S. ROSS, *A theory of the term structure of interest rates*, Econometrica, 53 (1985), pp. 385–407.
- [12] A. COZMA, M. MARIAPRAGASSAM, AND C. REISINGER, *Calibration of a hybrid local-stochastic volatility stochastic rates model with a control variate particle method (an extended version)*, 2018. arXiv preprint arXiv:1701.06001.
- [13] ———, *Convergence of an Euler scheme for a hybrid stochastic-local volatility model with stochastic rates in foreign exchange markets*, SIAM Journal of Financial Mathematics, (2018). Forthcoming.
- [14] C. S. DE GRAAF, D. KANDHAI, AND C. REISINGER, *Efficient exposure computation by risk factor decomposition*, arXiv preprint arXiv:1608.01197, (2016).
- [15] G. DEELSTRA AND G. RAYEE, *Local volatility pricing models for long-dated FX derivatives*, Applied Mathematical Finance, 20 (2013), pp. 380–402.
- [16] A. W. V. DER STOEP, L. A. GRZELAK, AND C. W. OOSTERLEE, *The Heston stochastic-local volatility model: efficient Monte Carlo simulation*, International Journal of Theoretical and Applied Finance, 17 (2014), pp. 1–30.
- [17] B. DUPIRE, *A unified theory of volatility*, in Derivatives Pricing: The Classic Collection, P. Carr, ed., Risk Books, 2004.
- [18] F. L. FLOCH, *TR-BDF2 for stable American option pricing*, Journal of Computational Finance, 17 (2014).
- [19] M. GILES AND R. CARTER, *Convergence analysis of Crank-Nicolson and Rannacher time-marching*, Journal of Computational Finance, 9 (2006), pp. 89–112.
- [20] P. GLASSERMAN, *Monte Carlo Methods in Financial Engineering*, vol. 53 of Stochastic Modelling and Applied Probability, Springer, 2003.
- [21] B. GOUGH, *GNU Scientific Library Reference Manual - Third Edition*, Network Theory Ltd., 3rd ed., 2009.
- [22] L. A. GRZELAK AND C. W. OOSTERLEE, *On the Heston model with stochastic interest rates*, SIAM Journal on Financial Mathematics, 2 (2011), pp. 255–286.
- [23] J. GUYON AND P. HENRY-LABORDÈRE, *Being particular about calibration*, Risk Magazine, (2012).
- [24] J. GUYON AND P. HENRY-LABORDÈRE, *Nonlinear Option Pricing*, Chapman and Hall/CRC Financial Mathematics, Chapman and Hall/CRC, 2013.
- [25] S. HESTON, *A closed-form solution for options with stochastic volatility with applications to bond and currency options*, Review of Financial Studies, 6 (1993), pp. 327–343.
- [26] T. R. HURD AND A. KUZNETSOV, *Explicit formulas for Laplace transforms of stochastic integrals*, Markov Processes and Related Fields, 14 (2008), pp. 277–290.
- [27] V. LUCIC, *Boundary conditions for computing densities in hybrid models via PDE methods*, Stochastics: An International Journal of Probability and Stochastic Processes, 84 (2012), pp. 705–718.
- [28] H. P. MCKEAN, *A class of Markov processes associated with nonlinear parabolic equations*, Proceedings of the National Academy of Sciences of the United States of America, 56 (1966), pp. 1907–1911.
- [29] R. T. O.C. ZIENKIEWICZ AND J. ZHU, *The Finite Element Method Set. Its Basis and Fundamentals*, Butterworth-Heinemann, 6 ed., 2005.
- [30] D. M. POOLEY, K. R. VETZAL, AND P. A. FORSYTH, *Convergence remedies for non-smooth payoffs in option pricing*, Journal of Computational Finance, 6 (2003), pp. 25–40.
- [31] A. QUARTERONI AND A. VALLI, *Numerical approximation of partial differential equations*, vol. 23, Springer Science & Business Media, 2008.
- [32] A. REGHAI, *The hybrid most likely path*, Risk Magazine, April (2006).
- [33] Y. REN, D. MADAN, AND M. QIAN QIAN, *Calibrating and pricing with embedded local volatility models*, Risk Magazine, September (2007).

- [34] C. L. C. G. ROGERS AND D. WILLIAMS, *Diffusions, Markov processes, and Martingales. Vol. 2*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2000.
- [35] R. SCHÖBEL AND J. ZHU, *Stochastic volatility with an Ornstein–Uhlenbeck process: an extension*, European Finance Review, 3 (1999), pp. 23–46.
- [36] B. W. SILVERMAN, *Density estimation for statistics and data analysis*, Biometrical Journal, 30 (1988), pp. 876–877.
- [37] A.-S. SZNITMAN, *Ecole d’Été de Probabilités de Saint-Flour XIX*, Springer, 1991, ch. Topics in propagation of chaos, pp. 165–251.
- [38] Y. TIAN, Z. ZHU, G. LEE, F. KLEBANER, AND K. HAMZA, *Calibrating and pricing with a stochastic-local volatility model*, Journal of Derivatives, 22 (2015), pp. 21–39.
- [39] L. TUR, *Local volatility calibration with fixed-point algorithm*, GDF Suez Trading, (2014). Internal.
- [40] A. W. VAN DER STOEP, L. A. GRZELAK, AND C. W. OOSTERLEE, *A novel Monte Carlo approach to hybrid local volatility models*, Quantitative Finance, forthcoming, (2017).
- [41] A. VAN HAASTRECHT, R. LORD, A. PELSSER, AND D. SCHRAGER, *Pricing long-maturity equity and FX derivatives with stochastic interest rates and stochastic volatility*, Insurance: Mathematics and Economics, 45 (2009), pp. 436–448.
- [42] A. VAN HAASTRECHT AND A. PELSSER, *Generic pricing of FX, inflation and stock options under stochastic interest rates and stochastic volatility*, Quantitative Finance, 11 (2011), pp. 665–691.
- [43] R. WHITE, *Numerical solution to PDEs with financial applications*, OpenGamma Quantitative Research, (2013). <https://developers.opengamma.com/quantitative-research/numerical-solutions-to-pdes-with-financial-applications-opengamma.pdf>.
- [44] M. WYNS, *Convergence analysis of the Modified Craig-Sneyd scheme for two-dimensional convection-diffusion equations with nonsmooth initial data*, arXiv preprint arXiv:1508.04296, (2015).
- [45] M. WYNS AND J. D. TOIT, *A finite volume - alternating direction implicit approach for the calibration of stochastic local volatility models*, arXiv preprint arXiv:1611.02961, (2016).
- [46] U. WYSTUP, *FX Options and Structured Products (The Wiley Finance Series)*, The Wiley Finance Series, Wiley, 2007.

Appendix A. Proof of Proposition 2 and Theorem 3.

We first state a necessary auxiliary result, which is an adaptation of Tanaka's formula in Chapter 4 of [34], where the integrand D in the local time integral is 1.

Proposition 7. *On a filtered probability space $(\chi, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{Q}^d)$, let D and X be two \mathcal{F}_t -adapted continuous semi-martingales, with D positive and integrable, $(l_t^a)_{t \geq 0}$ the local time of X at level a and, for all $n > 0$,*

$$(A.1) \quad \delta_n^a(x) = \begin{cases} 0, & |x - a| > \frac{1}{n}, \\ \frac{n}{2}, & |x - a| \leq \frac{1}{n}. \end{cases}$$

Then, for any $T > 0$ and $0 < t \leq T$, $\int_0^t D_s \delta_n^a(X_s) d\langle X \rangle_s$ converges almost surely, and uniformly in time, to $\int_0^t D_s dl_s^a$.

Proof. Follows as in Section 45 of [34] with a few additional steps. A complete proof is given in Appendix A of [12].

We now state and prove two lemmas necessary for the derivation of Theorem 3. Lemma 8 provides a link between the local time of a process and its density function.

Lemma 8. *Given a filtered probability space $(\chi, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{Q}^d)$, let W be a standard Brownian motion and μ, Y two \mathcal{F}_t -adapted processes with Y continuous, with finite second moment and*

$$\int_0^t (|\mu_u| + Y_u^2) du < \infty.$$

Consider a continuous Itô process X given by

$$X_t = X_0 + \int_0^t \mu_u du + \int_0^t Y_u dW_u$$

whose marginal density function $\phi(\cdot, t)$ and $\mathbb{E}^{\mathbb{Q}^d} [D_t Y_t^2 | X_t = \cdot]$ are assumed to be continuous. Further denote by $(l_t^a)_{t \geq 0}$ the local time of X at level a . Then, for any continuous, integrable and positive \mathcal{F}_t -adapted semi-martingale D and any $a \in \mathbb{R}$,

$$\mathbb{E}^{\mathbb{Q}^d} \left[\int_0^t D_u dl_u^a \right] = \int_0^t \mathbb{E}^{\mathbb{Q}^d} [D_u Y_u^2 | X_u = a] \phi(a, u) du.$$

Proof. From Proposition 7, we know that the following holds almost surely:

$$\int_0^t D_u dl_u^a = \lim_{n \rightarrow \infty} \int_0^t D_u \delta_n^a(X_u) Y_u^2 du.$$

This implies convergence in distribution, so that we can write

$$\mathbb{E}^{\mathbb{Q}^d} \left[\int_0^t D_u dl_u^a \right] = \lim_{n \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^d} \left[\int_0^t D_u \delta_n^a(X_u) Y_u^2 du \right],$$

and, by the stochastic Fubini theorem, we get

$$\begin{aligned}\mathbb{E}^{\mathbb{Q}^d} \left[\int_0^t D_u dl_u^a \right] &= \lim_{n \rightarrow \infty} \int_0^t \mathbb{E}^{\mathbb{Q}^d} [D_u \delta_n^a(X_u) Y_u^2] du \\ &= \lim_{n \rightarrow \infty} \int_0^t \mathbb{E}^{\mathbb{Q}^d} [\delta_n^a(X_u) \mathbb{E}^{\mathbb{Q}^d} [D_u Y_u^2 | X_u]] du.\end{aligned}$$

We denote $\gamma(x, u) = \mathbb{E}^{\mathbb{Q}^d} [D_u Y_u^2 | X_u = x]$, such that

$$\begin{aligned}\mathbb{E}^{\mathbb{Q}^d} \left[\int_0^t D_u dl_u^a \right] &= \lim_{n \rightarrow \infty} \int_0^t \int_0^\infty \delta_n^a(x) \gamma(x, u) \phi(x, u) dx du \\ &= \lim_{n \rightarrow \infty} \int_0^\infty \delta_n^a(x) \left(\int_0^t \gamma(x, u) \phi(x, u) du \right) dx,\end{aligned}$$

where we have used Fubini's Theorem in the second line. By the continuity assumptions on ϕ and γ , we deduce that

$$\mathbb{E}^{\mathbb{Q}^d} \left[\int_0^t D_u dl_u^a \right] = \int_0^t \left(\mathbb{E}^{\mathbb{Q}^d} [D_u Y_u^2 | X_u = a] \phi(a, u) \right) du.$$

Lemma 9. *Given the set-up of Theorem 3,*

$$M_t = \int_0^t \mathbf{1}_{S_u \geq K} D_u^d \alpha(S_u, u) S_u \sqrt{V_u} dW_u$$

is a true martingale up to T^ given by (2.7).*

Proof. Since α and $\mathbf{1}_{S_u \geq K}$ are bounded, the process

$$M_t = \int_0^t \mathbf{1}_{S_u \geq K} D_u^d \alpha(S_u, u) S_u \sqrt{V_u} dW_u,$$

is a true martingale if

$$\mathbb{E}^{\mathbb{Q}^d} \left[\int_0^t \left(D_u^d S_u \right)^2 V_u du \right] < \infty.$$

On the one hand, since $t < T^*$, from Proposition 3.13 in [13], we can find $\omega > 2$ such that

$$\sup_{u \in [0, t]} \mathbb{E}^{\mathbb{Q}^d} \left[\left(D_u^d S_u \right)^\omega \right] < \infty.$$

On the other hand, from Theorem 3.1 in [26],

$$\sup_{u \in [0, t]} \mathbb{E}^{\mathbb{Q}^d} \left[V_u^{\frac{\omega}{\omega-2}} \right] < \infty.$$

Using Hölder's inequality with the pair $\left(\frac{\omega}{2}, \frac{\omega}{\omega-2} \right)$,

$$\mathbb{E}^{\mathbb{Q}^d} \left[\left(D_u^d S_u \right)^2 V_u \right] \leq \mathbb{E}^{\mathbb{Q}^d} \left[\left(D_u^d S_u \right)^\omega \right]^{\frac{2}{\omega}} \mathbb{E}^{\mathbb{Q}^d} \left[V_u^{\frac{\omega}{\omega-2}} \right]^{\frac{\omega-2}{\omega}} < \infty.$$

Finally, using the Fubini theorem, $\mathbb{E}^{\mathbb{Q}^d} \left[\int_0^t (D_u^d S_u)^2 V_u du \right] < \infty$ and hence M is a true martingale of zero expectation. ■

Combining Lemmas 8 and 9, we can derive Proposition 2.

Proof of Proposition 2. Let $K \in \mathbb{R}^+$, $0 < t < T^*$ and $H_t = (S_t - K)^+$. The Trotter-Meyer theorem [34] gives

$$(S_t - K)^+ - (S_0 - K)^+ = \int_0^t \mathbf{1}_{S_u \geq K} dS_u + \frac{1}{2} l_t^K,$$

which we can write in differential form as

$$dH_t = \mathbf{1}_{S_t \geq K} S_t \left(r_t^d - r_t^f \right) dt + \frac{1}{2} dl_t^K + \mathbf{1}_{S_t \geq K} \alpha(S_t, t) S_t \sqrt{V_t} dW_t.$$

Also,

$$\begin{aligned} d \left(D_t^d H_t \right) &= D_t^d \left[-r_t^d H_t + \mathbf{1}_{S_t \geq K} S_t \left(r_t^d - r_t^f \right) \right] dt + \frac{1}{2} D_t^d dl_t^K \\ &\quad + \mathbf{1}_{S_t \geq K} D_t^d \alpha(S_t, t) S_t \sqrt{V_t} dW_t. \end{aligned} \tag{A.2}$$

Hence, by applying Lemma 8 with $X_t = S_t$, $D_t = D_t^d$ and $Y_t = \alpha(S_t, t) S_t \sqrt{V_t}$, we can write

$$\mathbb{E}^{\mathbb{Q}^d} \left[\int_0^t D_u^d dl_u^K \right] = \int_0^t \left(\alpha^2(K, u) K^2 \mathbb{E}^{\mathbb{Q}^d} \left[D_u^d V_u | S_u = K \right] \phi(K, u) \right) du,$$

where ϕ is the marginal density function of S at time t . Furthermore, one can define $\bar{\phi}_n$ as

$$\bar{\phi}_n(K, u) = \mathbb{E}^{\mathbb{Q}^d} \left[D_u^d \delta_n^K(S_u) \right] = \int_0^\infty \delta_n^K(x) \mathbb{E}^{\mathbb{Q}^d} \left[D_u^d | S_u = x \right] \phi(x, u) dx,$$

with δ_n^K defined as in (A.1) by

$$\delta_n^K(x) = \begin{cases} 0, & |x - K| > \frac{1}{n}, \\ \frac{n}{2}, & |x - K| \leq \frac{1}{n}, \end{cases}$$

and by a similar reasoning to that of Lemma 8 we get

$$\lim_{n \rightarrow \infty} \bar{\phi}_n = \mathbb{E}^{\mathbb{Q}^d} \left[D_u^d | S_u = K \right] \phi(K, u).$$

Since

$$\frac{\partial^2 C(K, u)}{\partial K^2} = \lim_{n \rightarrow \infty} \mathbb{E}^{\mathbb{Q}^d} \left[D_u^d \delta_n^K(S_u) \right],$$

we write

$$\frac{\partial^2 C(K, u)}{\partial K^2} = \mathbb{E}^{\mathbb{Q}^d} \left[D_u^d | S_u = K \right] \phi(K, u).$$

Combining (A.3) and (A.4) allows to write

$$\mathbb{E}^{\mathbb{Q}^d} \left[\int_0^t D_u^d dL_u^K \right] = \int_0^t \alpha^2(K, u) K^2 \frac{\mathbb{E}^{\mathbb{Q}^d} [D_u^d V_u | S_u = K]}{\mathbb{E}^{\mathbb{Q}^d} [D_u^d | S_u = K]} \frac{\partial^2 C(K, u)}{\partial K^2} du.$$

Hence, integrating (A.2),

$$\begin{aligned} C(K, t) &= \mathbb{E}^{\mathbb{Q}^d} [D_t^d H_t] = \int_0^t \left(-\mathbb{E}^{\mathbb{Q}^d} [D_u^d r_u^d (S_u - K)^+] + \mathbb{E}^{\mathbb{Q}^d} [D_u^d \mathbf{1}_{S_u \geq K} S_u (r_u^d - r_u^f)] \right) du \\ &+ \frac{1}{2} \int_0^t \alpha^2(K, u) K^2 \frac{\mathbb{E}^{\mathbb{Q}^d} [D_u^d V_u | S_u = K]}{\mathbb{E}^{\mathbb{Q}^d} [D_u^d | S_u = K]} \frac{\partial^2 C(K, u)}{\partial K^2} du \\ &+ \mathbb{E}^{\mathbb{Q}^d} \left[\int_0^t \mathbf{1}_{S_u \geq K} D_u^d \alpha(S_u, u) S_u \sqrt{V_u} dW_u \right]. \end{aligned} \quad (\text{A.5})$$

Furthermore, on a fixed time interval $[0, T^*]$, D_t^d is uniformly bounded by $\exp(T^* \max_{u \in [0, T^*]} |h^d(u)|)$. Then, from Lemma 9 we know that $\int_0^t \mathbf{1}_{S_u \geq K} D_u^d \alpha(S_u, u) S_u \sqrt{V_u} dW_u$ is a true martingale of zero expectation.

We write (A.5) at time T , differentiate with respect to T and, upon noticing that $\mathbf{1}_{S_T \geq K} S_T = (S_T - K)^+ + \mathbf{1}_{S_T < K} K$, we get (2.8). ■

We are now ready to give the proof of Theorem 3.

Proof of Theorem 3. First, we want to ensure that (2.5) is a necessary condition for

$$C(K, T) = C_{LV}(K, T).$$

Hence, by subtracting the Dupire PDE (2.4) from (2.8), we obtain

$$\begin{aligned} \frac{1}{2} K^2 \left(\alpha^2(K, T) \frac{\mathbb{E}^{\mathbb{Q}^d} [D_T^d V_T | S_T = K]}{\mathbb{E}^{\mathbb{Q}^d} [D_T^d | S_T = K]} - \sigma_{LV}^2(K, T) \right) \frac{\partial^2 C_{LV}}{\partial K^2} &= \mathbb{E}^{\mathbb{Q}^d} [D_T^d r_T^f (S_T - K)^+] \\ &- \bar{r}^f(T) C_{LV} - \mathbb{E}^{\mathbb{Q}^d} [D_T^d \mathbf{1}_{S_T \geq K} K (r_T^d - r_T^f)] - K (\bar{r}^d(T) - \bar{r}^f(T)) \frac{\partial C_{LV}}{\partial K}, \end{aligned}$$

so

$$\alpha^2(K, T) = \frac{\mathbb{E}^{\mathbb{Q}^d} [D_T^d | S_T = K]}{\mathbb{E}^{\mathbb{Q}^d} [D_T^d V_T | S_T = K]} (\sigma_{LV}^2(K, T) + \bar{q}(K, T)),$$

where

$$\bar{q}(K, T) = \frac{\mathbb{E}^{\mathbb{Q}^d} [\bar{Q}_T]}{\frac{1}{2} K^2 \frac{\partial^2 C_{LV}}{\partial K^2}},$$

$$\bar{Q}_T = D_T^d r_T^f (S_T - K)^+ - \bar{r}^f(T) C_{LV} - K \left(D_T^d \mathbf{1}_{S_T \geq K} (r_T^d - r_T^f) + (\bar{r}^d(T) - \bar{r}^f(T)) \frac{\partial C_{LV}}{\partial K} \right).$$

It remains to show that we can replace \bar{Q}_T by Q_T in \bar{q} . First, $D_T^d (S_T - K)^+$ is weakly differentiable with respect to K with $\frac{\partial D_T^d (S_T - K)^+}{\partial K} = D_T^d \mathbf{1}_{S_T \geq K}$, which is bounded by the integrable process D_T^d . We can interchange differentiation and expectation to get $\mathbb{E}^{\mathbb{Q}^d} [D_T^d \mathbf{1}_{S_T \geq K}] = -\frac{\partial C}{\partial K}$. Since the models agree, $\mathbb{E}^{\mathbb{Q}^d} [D_T^d \mathbf{1}_{S_T \geq K}] = -\frac{\partial C_{LV}}{\partial K}$ and $\mathbb{E}^{\mathbb{Q}^d} [D_T^d (S_T - K)^+] = C_{LV}$, and (2.5) holds. By re-tracing the steps in reverse order, one sees that (2.5) is also a sufficient condition for (A.6) provided the solution to (2.8) is unique. \blacksquare

Appendix B. Monte Carlo QE-scheme. The *Quadratic-Exponential (QE)* scheme [4], that is used to discretise the square-root process, employs moment-matching techniques and can significantly reduce the Monte Carlo discretisation error. While the full truncation Euler scheme and the QE scheme have shown to perform well in our tests, we experienced a faster convergence in time for the QE scheme when the Feller condition is broken. Hence, we choose the QE scheme for the variance process and the full truncation Euler for both stochastic rates, as the computational cost will be smaller. We briefly write a generalised QE scheme based on the original scheme from [4] to incorporate a leverage function and stochastic rates in the discretisation.

We follow our time interpolation rule for the calibration of α and interpolate forward-flat in time. We assume for simplicity that each Monte Carlo time step belongs to the α time grid. We can write

$$V_{t+\Delta t} = V_t + \int_t^{t+\Delta t} \kappa (\theta - V_u) du + \xi \int_t^{t+\Delta t} \sqrt{V_u} dW_u^V,$$

and hence

$$\int_t^{t+\Delta t} \sqrt{V_u} dW_u^V = \frac{V_{t+\Delta t} - V_t - \int_t^{t+\Delta t} \kappa (\theta - V_u) du}{\xi},$$

and

$$d \ln S_t = \left(r_t^d - r_t^f - \frac{1}{2} \alpha^2 (S_t, t) V_t \right) dt + \alpha (S_t, t) \rho \sqrt{V_t} dW_t^V + \alpha (S_t, t) \sqrt{1 - \rho^2} \sqrt{V_t} dW_t^S,$$

where W_t^S is a Brownian motion independent of W_t^V . Therefore,

$$\begin{aligned} \ln S_{t+\Delta t} &= \ln S_t + \int_t^{t+\Delta t} \left(r_u^d - r_u^f \right) du - \frac{1}{2} \alpha^2 (S_t, t) \int_t^{t+\Delta t} V_u du \\ &\quad + \frac{\alpha (S_t, t) \rho \left(V_{t+\Delta t} - V_t - \kappa \theta \Delta t + \kappa \int_t^{t+\Delta t} V_u du \right)}{\xi} \\ &\quad + \alpha (S_t, t) \sqrt{1 - \rho^2} \int_t^{t+\Delta t} \sqrt{V_u} dW_u^S. \end{aligned}$$

We approximate $\int_t^{t+\Delta t} V_u du$ by $\left(\frac{V_{t+\Delta t} + V_t}{2} \right) \Delta t$, and note that conditional on V_t and $\int_t^{t+\Delta t} V_u du$, since W_u^V and W_u^S are independent, the Itô integral $\int_t^{t+\Delta t} \sqrt{V_u} dW_u^S$ is normally distributed with mean zero and variance $\int_t^{t+\Delta t} V_u du$. We write the full scheme below

$$\begin{aligned}
812 \quad (B.1) \quad & \left\{ \begin{aligned}
g_{t+\Delta t}^d &= g_t^d + \kappa_d \left(\theta_d - (g_t^d)^+ \right) \Delta t + \xi_d \sqrt{(g_t^d)^+} \sqrt{\Delta t} Y_d \\
g_{t+\Delta t}^f &= g_t^f + \left(\kappa_f \left(\theta_f - (g_t^f)^+ \right) - \left(\rho_{Sf} \xi_f \sqrt{(g_t^f)^+} \alpha(S_t, t) \sqrt{V_t} \right) \right) \Delta t \\
&\quad + \xi_f \sqrt{(g_t^f)^+} \sqrt{\Delta t} Y_f \\
\ln S_{t+\Delta t} &= \ln S_t + \left(\frac{(r_{t+\Delta t}^d - r_{t+\Delta t}^f) + (r_t^d - r_t^f)}{2} - \frac{1}{4} \alpha^2(S_t, t) (V_{t+\Delta t} + V_t) \right) \Delta t \\
&\quad + \frac{\alpha(S_t, t) \rho (V_{t+\Delta t} - V_t + \kappa \left(\frac{V_{t+\Delta t} + V_t}{2} - \theta \right) \Delta t)}{\xi} \\
&\quad + \alpha(S_t, t) \sqrt{1 - \rho^2} \sqrt{\frac{V_{t+\Delta t} + V_t}{2}} \sqrt{\Delta t} Z \\
V_{t+\Delta t} : &\begin{cases} \text{if } \psi \leq \psi_c & : V_{t+\Delta t} = a (b + Z_v)^2 \\ \text{else} & \begin{cases} \text{if } U \leq p & : V_{t+\Delta t} = 0 \\ \text{else} & : V_{t+\Delta t} = \ln \left(\frac{1-p}{1-U} \right) \frac{m}{1-p}, \end{cases} \end{cases}
\end{aligned} \right.
\end{aligned}$$

813 with

$$\begin{aligned}
814 \quad & \begin{cases} m &= \theta + (V_t - \theta) e^{-\kappa \Delta t}, \\ \gamma^2 &= \frac{V_t \xi^2 e^{-\kappa \Delta t}}{\kappa} (1 - e^{-\kappa \Delta t}) + \frac{\theta \xi^2}{2\kappa} (1 - e^{-\kappa \Delta t})^2, \\ \psi &= \frac{\gamma^2}{m^2}, \quad p = \frac{\psi-1}{\psi+1}, \quad \beta = \frac{1-p}{m}, \\ b^2 &= \frac{2}{\psi} - 1 + \sqrt{\frac{2}{\psi}} \sqrt{\frac{2}{\psi} - 1}, \quad a = \frac{m}{1+b^2}, \\ \psi_c &= 1.5, \end{cases}
\end{aligned}$$

815 Let the Cholesky decomposition of the correlation matrix ⁵

$$816 \quad \begin{bmatrix} 1 & \rho & 0 & 0 \\ \rho & 1 & \rho_{Sd} & \rho_{Sf} \\ 0 & \rho_{Sd} & 1 & \rho_{df} \\ 0 & \rho_{Sf} & \rho_{df} & 1 \end{bmatrix},$$

817 be $\mathbf{L}\mathbf{L}^T$. Y_d and Y_f are defined as

$$818 \quad \begin{bmatrix} Y_v \\ Y_s \\ Y_d \\ Y_f \end{bmatrix} = \mathbf{L} \begin{bmatrix} Z_v \\ Z \\ Z_d \\ Z_f \end{bmatrix},$$

819 where Z, Z_v, Z_d, Z_f are independent draws from a standard normal distribution and U is a draw
820 from a uniform distribution.

⁵If the correlation matrix is not positive definite, one can rely on spectral decomposition instead, see 2.3 in [20] for details.

Appendix C. Finite element mesh construction.

In order to refine the mesh in the most relevant area, we use an exponential mesh on the variance axis and a hyperbolic mesh (see [43]) in the spot direction. This makes the mesh finer around $z = 0$ and $x = S_0$. In order to build our mesh, we first define the grids in spot $(x_i)_{i \in \llbracket 0, N_S \rrbracket}$ and variance $(z_j)_{j \in \llbracket 0, N_V \rrbracket}$ separately. Additionally, to solve the PDE numerically, we need to truncate at the boundary and use $\bar{\Omega} = \{(x, z) \in [0, S_{\max}] \times [0, V_{\max}]\}$ on a time interval $[0, T]$. We choose

$$S_{\max} = S_0 e^{5\alpha(S_0, T)} \sqrt{(v_0 e^{-\kappa T} + \theta(1 - e^{-\kappa T}))T}$$

and recall that the stationary distribution of the CIR process is a gamma distribution of density ϕ_v^∞ as defined in (4.3). We compute V_{\max} with the inverse cumulative density function such that

$$\mathbb{P}(z > V_{\max}) = 0.01\%.$$

We write

$$\begin{aligned} x_i &= f_h(g_h(\bar{x}_i)), \\ z_j &= f_e(g_e(\bar{z}_j)), \end{aligned}$$

with

$$\begin{aligned} f_h(x) &= S_0 + b \sinh(\nu x + d), & f_e(z) &= c + c \exp(\lambda z), \\ b &= \eta(S_{\max} - S_{\min}), & c &= \frac{V_{\max}}{e^\lambda - 1}, \\ d &= \operatorname{arcsinh}\left(\frac{S_{\min} - S_0}{b}\right), & \lambda &= \max\left(1, 4 - \frac{3\kappa\theta}{\xi^2}\right), \\ \nu &= \operatorname{arcsinh}\left(\frac{S_{\max} - S_0}{b}\right) - d, & \bar{z}_j &= \frac{j}{(N_V + 1)}, \\ \bar{x}_i &= \frac{i}{(N_S + 1)}, & & \\ \eta &= 0.02, & & \end{aligned}$$

where η is defined according to our numerical experiments and g_h is the quadratic polynomial that passes through the points $(0, 0)$, $(1, 1)$,

where λ is defined according to our numerical experiments and g_e is the quadratic polynomial that passes through the points $(0, 0)$, $(1, 1)$,

$$\left(\frac{\lfloor f_h^{-1}(S_0)(N_S + 1) + 0.5 \rfloor}{N_S + 1}, f_h^{-1}(S_0)\right) \quad \left(\frac{\lfloor f_e^{-1}(v_0)(N_V + 1) + 0.5 \rfloor}{N_V + 1}, f_e^{-1}(v_0)\right).$$

The latter intermediate step makes sure that both S_0 and v_0 are vertices of their respective grids. The construction of the finite element triangular mesh can be achieved by creating a vertex at each point (x_i, z_j) and defining two triangular cells (upper left and lower right) in each rectangle.