

Cellular census of human fibrosis defines functionally distinct stromal cell types and states

Authors

Thomas B Layton¹, Lynn Williams¹, Fiona McCann¹, Mingjun Zhang², Marco Fritzsche¹, Huw Colin-York³, Marisa Cabrita¹, Michael T H Ng³, Marc Feldmann¹, Stephen Samson¹, Dominic Furniss⁴, Weilin Xie^{2, 5}, Jagdeep Nanchahal^{1†}

Affiliations

¹The Kennedy Institute of Rheumatology, Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, UK

²Department of Inflammation Research, Celgene Corporation, San Diego, California, United States of America.

³MRC Human Immunology Unit, Weatherall Institute of Molecular Medicine, University of Oxford, Headley Way, Oxford, UK.

⁴Nuffield Department of Orthopaedics, Rheumatology, and Musculoskeletal Sciences, University of Oxford, Botnar Research Centre, UK.

⁵Present address: Institute of Materia Medica, Shandong First Medical University & Shandong, Academy of Medical Sciences, Jinan, Shandong Province, P R China

†Corresponding author. Email: jagdeep.nanchahal@kennedy.ox.ac.uk

Abstract

Fibrotic disorders are some of the most devastating and poorly treated conditions in developed nations, yet effective therapeutics are not identified for many of them. A major barrier for the identification of targets and successful clinical translation is a limited understanding of the human fibrotic microenvironment. Here, we construct a stromal cell atlas of human fibrosis at single cell resolution from patients with Dupuytren's disease, a localized fibrotic condition of the hand. A molecular taxonomy of the fibrotic milieu characterizes functionally distinct stromal cell types and states, including a subset of immune regulatory ICAM1⁺ fibroblasts. In developing fibrosis, myofibroblasts exist along an activation continuum of phenotypically distinct populations. We also show that the tetraspanin CD82 regulates cell cycle progression and can be used as a cell surface marker of myofibroblasts. These findings have important implications for targeting core pathogenic drivers of human fibrosis.

Introduction

Fibrosis is defined as the accumulation of excess matrix proteins and contributes to a significant proportion of the mortality in the Western World^{1,2}. The key mediators of fibrosis are collagen-producing stromal cells, whose activation drives a coordinated pathogenic system that ultimately compromises organ function^{1,2}. To develop effective novel therapeutics, we must first gain a detailed understanding of the states, subtypes and functional properties of fibrotic stromal cells³⁻⁷. A major challenge in studying fibrosis is the availability of well-characterized patient samples to dissect the molecular landscape of these disorders.

Localized fibrotic diseases represent a rich source of readily accessible early stage tissue⁸. Dupuytren's disease (DD) is a common and progressive fibroproliferative disorder of the palmar and digital fascia of the hand and, in Western populations, affects 12% of those aged 55 years, increasing to 29% of those 75 years and older⁹. The initial clinical presentation is the appearance of a firm nodule in the palm, which expands into fibrous collagenous cords that extend into the digits. Dupuytren's nodules, which represent the early stage of the disease, are a highly cellular fibrotic ecosystem and are an important model to examine developing fibrosis in humans.

In this study, using clinical samples from DD patients, we construct a molecular taxonomy of human fibrosis. Our single cell atlas of the fibrotic milieu elucidates functionally distinct stromal cell types and states, including ICAM1⁺ fibroblasts and CD82^{high} myofibroblasts that contribute discrete pro-fibrotic functions. In addition, we demonstrate that the tetraspanin CD82 is expressed on human myofibroblasts and functions to regulate cell cycle progression in this population.

Results

Single cell profiling of developing human fibrosis

To profile human fibrosis, we examined Dupuytren's nodules using mass cytometry (CyTOF) and single cell RNA-seq, constructing a single-cell atlas of an active and cellular fibrotic microenvironment (Fig. 1a-e, Supplementary Figure 1a-f). We generated a dataset yielding high quality profiles of over 300,000 cells from 18 patients (12 single cell RNA-seq & 6 CyTOF) to study cellular heterogeneity in this pathological ecosystem (Fig. 1a-e, Supplementary Figure 1a-f). Initially, we employed graph-based clustering¹⁰ that resolved a complex disease landscape comprised of fibroblasts, myofibroblasts, immune, pericytes, cycling and endothelial cells (Fig. 1b-e, Supplementary Figure 1c,d). Next, we performed a GWAS meta-analysis of DD from UK Biobank and the BSSH-GODD study¹¹, and mapped the associated regions to candidate genes and individual cellular subtypes using SNPsea¹². GWAS associated SNPs were strongly associated with particular stromal subtypes (Supplementary Figure 2a,b), supporting our classification. Marker genes for pericytes included *JAG1* and *MCAM*, *SFPR4* and *PLA2G2A* for fibroblasts, and myofibroblasts were marked by *MMP14* and *MAFB*. The single cell RNA-seq and CyTOF aligned with significant markers for stromal cell types, where CyTOF further elucidated cell surface expression of these markers for fibroblasts, myofibroblasts and pericytes.

A dynamic and immunoregulatory ICAM1⁺ fibroblast in fibrosis

We then undertook detailed characterization of fibroblasts and myofibroblasts. Paired single cell profiling using scRNA-seq and CyTOF uncovered distinct molecular profiles of ACTA2⁻ fibroblasts (CD9, PLA2G2A, C1R, CXCL14) and ACTA2⁺ myofibroblasts (α -SMA, ITG- β 1, TGF- β 1, β -catenin, MMP14). (Supplementary Figure 2c-g). Myofibroblasts were the predominant cell type representing ~60% of the total stromal cell population (Supplementary Figure 2d). This observation is consistent with previous histological descriptions of DD nodules that showed these structures are composed of densely packed myofibroblasts¹³. Functional annotation¹⁴ of cell marker genes using gene ontology

(GO) revealed a diversification of function, with inflammation, proliferation and ECM remodelling pathways localizing to distinct cell types (Fig. 2a, Supplementary Figure 2e). Myofibroblasts showed strongest enrichment for pathways involved with cell contraction and ECM remodelling (Fig. 2a, Supplementary Figure 2e) and core enrichment genes including *ACTA2*, *TPM2* and *POSTN*. In contrast, fibroblasts showed an enrichment for genes involved with ‘chemokine activity’ and ‘immune response’ pathways (Fig. 2a, Supplementary Figure 2e) including *CXCL14*, *CXCL8*, *IL6*, *C1R* and *PLA2G2A*.

Next, we performed a focused analysis of *ACTA2*⁺ fibroblasts, subsampling this population from the entire dataset and repeating the computational workflow used to delineate major cell types. Graph-based clustering of transcriptomic profiles defined three major subsets, *CD34*⁺, *PDPN*⁺ and *ICAM1*⁺ fibroblasts (*ICAM1*⁺*IL6*^{high}, *ICAM1*⁺*IL6*^{low}) (Fig. 2b-d, Supplementary Figure 3a,b). Moreover, these populations were conserved at the protein level in the CyTOF dataset. GO analysis of fibroblast marker genes demonstrated *ICAM1*⁺*IL6*^{high} fibroblasts were enriched for immune responsive pathways driven by the expression of several chemokines including *IL6* and *IL8* (Fig. 2d,e). We noted *ICAM1*⁺*IL6*^{high} fibroblasts were conserved across multiple patient samples and confirmed this subset showed the highest protein expression of IL-6 and IL-8 (Supplementary Figure. 3b,c) using flow cytometry of freshly isolated DD nodular cells. Subsequently, to explore potential relationships between subsets we applied diffusion maps to the fibroblasts. This uncovered a complex topography with discrete trajectories linking *CD34*⁺ and *ICAM1*⁺ subsets with *PDPN*⁺ fibroblasts, suggesting a putative underlying developmental path (Supplementary Figure. 3d).

Next, we sought to define the dynamics of fibroblast subsets in fibrosis pathogenesis. To assess this, we used flow cytometry to determine their proportions within two distinct Dupuytren’s structures, the early disease state myofibroblast and immune cell-rich nodule^{15,16} and later disease stage matrix-rich cord^{17,18} (Fig. 2f,h, Supplementary Figure 3e). We observed a higher proportion of *ICAM1*⁺

fibroblasts in nodules, which have been shown to harbour the majority of inflammatory cells in DD and are present at the early stages of the disease (Fig. 2h). Subsequently, we tested whether ICAM1⁺ fibroblasts could induce immune cell chemotaxis as predicted by their gene expression profiles. For this, we sorted freshly isolated ICAM1⁺ and ICAM1⁻ fibroblasts (CD45⁻CD31⁻CD146⁻ITGβ1^{low}) from Dupuytren's nodules and incubated each with THP-1 mononuclear immune cells (Fig. 2g). This confirmed that ICAM1⁺IL6^{high} fibroblasts produced significantly higher immune cell chemotaxis (Fig. 2g). Together, this identifies a dynamic ICAM1⁺IL6^{high} fibroblast in human fibrosis which act to promote immune-cell recruitment.

Distinct myofibroblast states along an activation continuum

Myofibroblasts are central mediators of the dysregulated wound-healing program that defines fibrosis^{3,4} therefore we studied this population in detail. Graph based clustering of the single cell RNA-seq data defined four major subsets (Fig. 3a) that included a cycling population (Cycling MFB) (Fig. 3a, b). After this, we sought to confirm that proliferating stromal cells (Ki67⁺) were myofibroblasts (Fig. 3c-e, Supplementary Figure 4a-d). Using flow cytometry, we stained freshly disaggregated nodular cells and gated on myofibroblasts (CD45⁻CD31⁻CD146⁻ITGβ1^{high}) and fibroblasts (CD45⁻CD31⁻CD146⁻ITGβ1^{low}) and demonstrated Ki67^{high} cells were a subset of myofibroblasts. In the single cell RNA-seq, a second subset was characterized by lower expression of *ACTA2* and intermediate expression of fibroblast marker genes (*CXCL14*, *PLA2G2A*, *C1R*), we termed the ACTA2^{low} myofibroblast (Fig. 3a and b, Supplementary Figure 5a). A third comprised a discrete CD82^{high}OX40L⁺ population (CD82, OX40L, MMP11, MMP14, CD82) (Fig. 3a,g, Supplementary Figure 5b-e). Finally, a fourth less distinct community we termed the intermediate myofibroblast (Fig 3a and b, 2 Supplementary Figure 5a). GO analysis demonstrated that CD82^{high}OX40L⁺ myofibroblast marker genes showed the strongest enrichment for cell contraction pathways (Fig. 3f). In contrast, differential gene expression and pathway analysis revealed ACTA2^{low} myofibroblasts resembled a general fibroblast phenotype, without clear characteristics of any one subtype. In addition to *ACTA2*, significant ACTA2^{low}

myofibroblasts markers were general fibroblasts markers including PLA2G2A and CXCL14. ACTA2^{low} myofibroblasts also shared pathways enriched in fibroblasts such as ‘interleukin signalling’ and ‘chemotaxis (Fig. 3f).

As we discovered the membrane tetraspanin CD82 marked the myofibroblast population showing high expression of ACTA2 and ITGB1 (CD82^{high}OX40L⁺ myofibroblast), we confirmed its co-expression with α -SMA and ITG β -1 proteins (Fig. 3g,h, Supplementary Figure 5f,g) using flow cytometry and multiplex immunofluorescence. This validated a cell surface marker of human myofibroblasts and showed tight co-expression of CD82 with established myofibroblast markers α -SMA and ITG- β 1 (Supplementary Figure 5f). Finally, using immunohistochemistry we confirmed CD82⁺ myofibroblasts were enriched in Dupuytren’s nodules as compared to cord (Supplementary Figure 5g).

The overall topography of the myofibroblast clusters suggested an overarching trajectory structure in which the two distinct cell populations represent diverging ends of a continuum separated by an intermediate cellular state (Fig. 3a). This concept reflects a central principle of myofibroblast biology, which describes an ‘activation’ paradigm along a differentiation path towards a fully pathogenic phenotype^{19,20}. Supporting this concept, principal component analysis (PCA) of the myofibroblasts demonstrated the first principal component to be strongly associated with ACTA2 expression and inversely correlated with a gene signature of ACTA2^{low} myofibroblasts (Supplementary Figure 6a,b).

To better understand this activation signature, we used diffusion maps to reconstruct a potential developmental path. In accordance with the PCA, the first two diffusion map components^{21,22} captured a trajectory associated with increasing ACTA2 and CD82 expression (Supplementary Figure 6c-f). Fitting a principal curve²³ through the first two diffusion components enabled us to define a dynamic gene module along this continuum (Supplementary Figure 6c). We observed a coordinated decrease in the ACTA2^{low} myofibroblast markers and progressive increase in CD82^{high}OX40L⁺

myofibroblast markers along this path (Supplementary Figure. 6f). This observation is compatible with the CD82^{high}OX40L⁺ myofibroblast representing a fully ‘activated’ phenotype, at the terminal stage of a differentiation path. In light of this finding, we performed RNA velocity analysis on human myofibroblasts to explore potential directionality in trajectories using the proportion of spliced and unspliced reads. This demonstrated a complex vector field showing a transition from cycling myofibroblasts into the major cell populations and a putative dynamic equilibrium moving between CD82^{high} and ACTA2^{low} myofibroblasts.

After this, we looked for the spatial expression of CD82^{high} myofibroblast markers in DD nodules using immunohistochemistry. Within nodules, CD82^{high}OX40L⁺ myofibroblast markers (e.g. MMP11, CD82, MMP14) had the highest expression in densely packed, α -SMA⁺ myofibroblast-rich foci (Supplementary Figure 7).

Functional heterogeneity of myofibroblasts in human fibrosis

We next investigated whether myofibroblasts along the activation trajectory were phenotypically distinct. We sorted CD82^{high} and CD82^{low} myofibroblasts and performed traction force microscopy (TFM) to measure cell contractility at single-cell resolution (Fig. 4a-d). As force generation is a key *in vivo* characteristic of myofibroblasts it is a highly relevant functional readout. In addition, pathway analysis of CD82^{high} myofibroblasts associated this population with cell contraction pathways and thus TFM enabled validation of a functional characteristic predicted in the single cell RNA-seq (Fig 4A). This demonstrated human myofibroblasts are functionally heterogeneous with CD82^{high} myofibroblasts marking a highly contractile population (Fig. 4b-d). In addition, CD82^{high} myofibroblasts had distinct biophysical profiles with localized areas of high force magnitude that enables these cells to bind and remodel matrix proteins (Fig. 4b).

To further understand the functional role of *CD82*, we performed bulk RNA-sequencing following siRNA mediated knock-down in freshly-isolated myofibroblasts (Fig. 4e, Supplementary Figure. 8a-d). This revealed that *CD82* regulated a wide array of genes and GO analysis of *CD82* regulated genes demonstrated an enrichment for diverse processes, including cell cycle and p53 signalling (Supplementary Figure 8e). Given this, we sought to validate a functional role of *CD82* in myofibroblasts. Using a flow cytometry-based cell cycle assay and siRNA mediated knockdown of *CD82*, we confirmed *CD82* directly regulated the myofibroblast cell cycle, promoting entrance into the G2-M phase (Fig. 4e). Together, these results support a role of *CD82* in maintaining the proliferative potential of myofibroblasts and may explain how these cells accumulate in this condition.

To explore the relevance of our findings in other fibrotic diseases, next we examined *CD82*⁺ stromal cells in a murine model of pulmonary fibrosis (Fig. 4f-h, Supplementary Figure 9a-c). Remarkably, we discovered that *Cd82* marked a distinct *Pdgfra*⁺ fibroblast which was the most upregulated population during fibrosis progression (Fig 4f-h, Supplementary Figure 9b-c). Based on these findings, we looked for *CD82* protein expression in human pulmonary fibrosis (IPF), co-staining for pertinent markers of *Pdgfra*⁺ fibroblasts (*PDGFR-α* & *COL13A1*) (Fig. 4i, Supplementary Figure 9d). This confirmed the presence of *CD82*⁺ fibroblasts in human IPF adjacent to fibrotic foci. Collectively, these data show a *Cd82*⁺*Prgfra*⁺ fibroblast in murine and human pulmonary fibrosis and demonstrate that *CD82* marks distinct myofibroblast and fibroblast populations across visceral and localised human fibrosis. It is therefore possible that at different anatomical sites undergoing fibrosis *CD82* may drive diverse cellular responses to propagate this disease process.

Discussion

The fibrotic microenvironment houses multiple cell subpopulations with diverse genetic and phenotypic characteristics²⁴⁻²⁷. How this heterogeneity emerges in developing fibrosis remains unclear

and a major barrier to this understanding is the limited availability of high-quality human samples^{8,28}. Here, we build a single cell atlas of a human fibrotic microenvironment and describe functionally distinct fibroblast and myofibroblast types and states. A number of recent studies have utilized single cell RNA-seq to profile human fibrotic disease, including lung and liver fibrosis, but these reports highlight the challenge in studying small biopsy samples from these disorders²⁹. Crucially, our results from Dupuytren's nodules enabled description of stromal cells from a complete fibrotic ecosystem, excised in its entirety, and thus forms a powerful resource for future studies in fibrosis.

Fibroblasts are increasingly recognized as central mediators of diverse pathological systems and here we uncovered three major subsets (ICAM1⁺, CD34⁺ and PDPN⁺) in human fibrosis. Crucially, we discovered a dynamic immune regulatory ICAM1⁺IL6^{high} fibroblast expressing high levels of chemokines and exhibiting a direct chemotactic activity. In addition, we show this population is expanded in developing and immune-cell rich stages of fibrosis in Dupuytren's nodules. We have previously shown in DD immune cell factors, including TNF, are crucial for myofibroblast development and activation¹⁶. Thus, together these findings support ICAM1⁺IL6^{high} fibroblasts as key drivers of inflammation that feedbacks to sustain stromal activation during fibrosis progression.

Tetraspanins are a family of highly conserved proteins containing four transmembrane domains with diverse functions in human health and disease. CD82 represents a canonical membrane tetraspanin, most studied in the context T cell activation and tumour suppression³⁰. Molecular functions of CD82 are broad and include directing intracellular signalling³¹, binding to integrins^{32,33} and modulating p53³⁴. To our knowledge CD82 has not been described in relation to myofibroblasts. Here, we show CD82 is a specific protein and gene marker of human myofibroblasts whose expression tightly correlates with α -SMA and ITG- β 1. In addition, we demonstrate it acts to regulate myofibroblast cell cycle progression and provides a mechanism by which these cells may accumulate in Dupuytren's disease.

Our stromal cell census delineates a unified molecular program of myofibroblasts in fibrosis and demonstrates these cells exist along an activation trajectory housing ACTA2^{low}, intermediate and CD82^{high}OX40L⁺ subsets, the latter being highly contractile and housing a cycling population. RNA velocity analysis suggests a complex and dynamic topography of this activation signature containing a proliferative subset that supports a precursors pool. Additionally, we show CD82 expression follows the activation trajectory that transitions between ACTA2^{low} and CD82^{high}OX40L⁺ myofibroblasts.

A central function of myofibroblasts is the generation of traction force which plays a key role in remodelling the matrix and also modulates the activities of the embedded stromal cells in wound healing and fibrosis⁷⁻⁹. Using traction force microscopy, we validate functional heterogeneity along the myofibroblast activation trajectory marked by CD82 and show CD82^{high} myofibroblasts have distinct biophysical profiles exerting high traction force on ECM coated hydrogels. It is reasonable that force generation in this population contributes to their proliferative and ECM remodelling capacity. However, the biophysical profile of CD82^{high} myofibroblasts shows distinct force foci of corresponding size and morphology to fibrillary matrix proteins³⁶ and supports a predominant role of binding to and remodelling of the matrix.

Given the lack of an animal model for DD we are unable to directly validate our findings in an *in vivo* model, including potential functional and lineage tracing studies of stromal cell populations. In addition, limitations of murine models of IPF must be acknowledged with regards to their relevance to human disease. Nonetheless, the differential expression of CD82 on fibroblasts in visceral fibrosis and myofibroblasts in localised human fibrosis suggests discrete functions during fibrosis at different anatomical sites. We uncovered a functional role of CD82 in human myofibroblasts in localised fibrosis, but the exact molecular function of CD82⁺ stromal cells in pulmonary fibrosis remains unknown and represents an exciting avenue for future research.

In summary, our findings provide important insights into mesenchymal subpopulations in human fibrosis, classify a division of labour between fibrotic stromal cells and form a powerful translational resource to help inform development of future treatments.

Methods

Patient samples

After approval by the local ethical review committee (REC 07/H0706/81, University of Oxford), tissue samples were obtained with informed consent from patients with Dupuytren's disease. Dupuytren's nodular tissue were obtained from individuals with Dupuytren's disease undergoing dermofasciectomy. Given the lack of well-defined controls in Dupuytren's disease we instead focused on dissection heterogeneity within the fibrotic microenvironment.

Cell culture

Cells from Dupuytren's disease were isolated from α -SMA-rich nodules as described previously^{17,37}. Tissue samples were dissected into small pieces and digested in DMEM (Lonza) with Type I collagenase (Worthington Biochemical Corporation) + DNase I (Roche Diagnostics) for up to 2 h at 37 °C. Cells were cultured in DMEM with 5% (vol/vol) FBS and 1% penicillin–streptomycin at 37 °C in a humidified incubator with 5% (vol/vol) CO₂. Cells before passage 2 were used for experiments and only freshly isolated cells were used for single cell RNA-seq, CyTOF and flow cytometry.

Antibodies

Antibodies used in flow cytometry: PE-ICAM1 (Biolegend HA58), PE/Cy7-CD34 (Biolegend 581-BL), APC-PDPN (Biolegend NC-08), APC-CD31 (Biolegend WM59), APC-CD146 (Biolegend SHM-57), FITC-CD146 (Abcam PIH12), BV421-CD45 (Biolegend HI30), BV605-ki67 (Biolegend Ki67), APC-IL8 (eBioscience BCH), APC-IL6 (Biolegend MQ2-13A5), AF700- α -SMA (R&D Systems ICI420N), APC-CD29

(Biolegend 152/16) and PE-CD82 (Biolegend ASL-24). Antibodies used for immunohistochemistry and Immunofluorescence: α -SMA (Abcam A5228), PLA2G2A (Novus Bio. 620501), CXCL14 (Abcam ab36622), C1R (Sigma-Aldrich P00736), ITG- β 1 (Abcam EP1041Y), β -Catenin (Abcam ab16051), MMP14 (Abcam EP1264Y), FAPB5 (Abcam ab84028), IL-11RA (Abcam ab109697), MMP11 (Abcam ab53143) and CD82 (Abcam TS82b)

Single cell Isolation

Following surgical resection Dupuytren's disease samples were rapidly transported to the research facility. On arrival, samples were then rinsed in PBS and the nodule isolated using a scalpel. Each nodule was minced using a scalpel and transferred to 5ml of digestion medium described above. Dissociated cells were then washed in 5% FBS DMEM (Gibco) and passed through a 100 μ m cell strainer. This single cell solution was slowly frozen using a Mr Frosty Container (ThermoFisher). Single cell suspensions were thawed on the day of sequencing, diluted at a concentration of 1000 cells/ μ L in 0.04% BSA/PBS for loading into 10X Chromium Single Cell A Chips.

Droplet-Based scRNA-seq

Single cell libraries were prepared using the Chromium 3' v2 platform (10X Genomics, Pleasanton, CA) following the manufacturer's protocol. In brief, single cells were encapsulated into Gel beads in Emulsions (GEMs) in the GemCode instrument followed by cell lysis and barcoded reverse transcription of RNA, amplification, shearing and 3' adaptor and sample index attachment. Approximately 10,000 single cells were loaded on each channel and approximately 1,500-6,000 cells were recovered. Libraries were sequenced on the Illumina HiSeq 4000 (Paired end reads: Read 1, 26bp, Read 2, 98bp).

Computational analysis

Sample de-multiplexing, alignment to the GRCh38 human transcriptome and UMI-collapsing were performed using the Cellranger toolkit (v2, 10X Genomics). Aspects of the downstream analysis were done in Seurat R package (Sajita Lab). We first performed quality control on each patient sample separately. We excluded poor quality cells that expressed fewer than 200 genes and with over 10% of UMIs mapping to mitochondrial genes as well as cells that expressed over 4500 genes. After this, the donors were aligned from the first and second batches. Batch correction was performed using Combat as implemented in the 'sva' R package using the default parametric adjustment mode³⁸. Batch correction was assessed before downstream analysis and we observed minimal batch effects following the Combat method. We employed a global-scaling normalization procedure as per³⁹ calling Seurat's LogNormalize() function. Briefly, the cell ranger UMI count value for each gene was divided by the sum of the total UMI counts per cell to normalize for differences in library size and then multiplied by a scaling factor that represented the median library complexity (10,000) producing TPM-like values. We then took the log transform of this procedure for downstream analysis. Downstream analysis was performed using all patient donors, and first and second patient batches were displayed separately for visualization only.

Feature selection was first undertaken by defining highly variable genes using the FindVariableGenes() function (1,856 genes). Values were then centred and scaled before input to PCA, which was implemented using the R function 'prcomp' from the "stats" R package. After PCA, significant PCs were identified using the permutation test implemented using the 'permutationPA' function from the 'jackstraw' R package⁴⁰. This test identified 20 significant PCs and these were used as the input to further analysis.

To partition the data into clusters of transcriptionally related cells, we used unsupervised clustering based on the Louvain algorithm with the Jaccard correction implemented in the FindClusters()

function. Initially, we used clustering to delineate the major cell types. Cell clusters were then annotated to known biological cell types using canonical marker genes. In many of the patient samples and pooled dataset, we noted a small (<2%) population of proliferative stromal cells. Regression of the cell cycle effect did not significantly influence clustering and therefore we opted not to undertake correction for cell cycle effects. After defining the major cell types we subsampled each (Fibroblasts, myofibroblast, Pericytes and endothelial cells) to identify potentially meaningful sub-clusters. We applied the same steps as described above to delineate sub-clusters including selection of highly variable genes, PCA and selection of significant PCs. As the average UMI count varied between distinct cell types when analysing individual cell types, we performed a second round of quality control to remove doubles and contaminants. We used a set of canonical markers for each cell type and excluded those cells that shared markers of multiple cell types and those cells with a UMI count 3 standard deviations above the median cell-type UMI count. For visualization we used the Barnes-Hut approximate version of t-distributed stochastic neighbour embedding (tSNE) using the 'Rtsne' R package.

On inspection of the the PCA projection of the myofibroblast dataset we noted a strong association between *ACTA2* and CD82⁺ marker expression to the first principal component. This was apparent in both pooled datasets and in individual patient samples. To investigate this further, we extracted those genes with the top PC1 loading and again saw a strong negative correlation to ACTA2^{low} myofibroblast marker genes (eg. *PLA2G2A* & *C1R*) and strong positive correlation to *ACTA2* expression and markers of the CD82⁺ population. Given this, we reasoned that this may represent the transcriptional profile of a putative differentiation or activation pathway. To visualize this, we defined a gene module of the ACTA2^{low} myofibroblasts and CD82⁺ myofibroblast taking top 100 genes correlated (Spearman) to *ACTA2* and *CXCL14*. Then, we plotted PC1 score of each cell against the average expression of these gene modules minus a control gene set and obtained the Pearson correlation coefficient. To account for differences in library complexity between cells we calculated a control gene set from these

scores⁴¹. This was selected by binning all genes in the dataset and randomly selected 100 genes from each bin that contained every gene of the test gene set. Thus, the control gene set has a comparable distribution of expression levels to that of the ACTA2 and CXCL14 gene-sets and the control gene set is 100-fold larger, such that its average expression is analogous to averaging over 100 randomly-selected gene-sets of the same size as the ACTA2 and CXCL14 gene-sets. This revealed a strong positive correlation to the ACTA2 gene module ($R=0.46$) and a strong negative correlation to the fibroblast-like gene module ($R=-0.71$) along the first principal component, suggesting a strong transcriptome signature of this putative trajectory in the data.

To assign each myofibroblast along the putative activation trajectory we first performed dimensional reduction using diffusion maps, implemented in the R package 'destiny'⁴². The distance metric was the Euclidean distance between pairs of cells in the reduced dimension space of the significant PCs ($n=20$), as defined above. In concordance with the PCA, the first two diffusion components were highly associated with ACTA2 expression. We then fitted a principal curve (R package princurve, smoother = 'lowess', $f = 1/3$) through the first two diffusion co-ordinates⁴³. As the λ value of the curve reflects the arc-length from the beginning of the curve for each point we utilised this to assign each cell to a "Pseudotime" trajectory. Importantly, this modelled trajectory was focused on capturing myofibroblast differentiation associated with ACTA2 expression, but did not ascribed any distinct cell type as the myofibroblast precursor. In addition to this approach, we also the 'slingshot' R package to infer trajectories and align cells along developmental pseudo-time. Slingshot revealed a very similar trajectory structure with a path between ACTA2^{low} and CD82^{high} myofibroblasts. Expression smoothing was performed using generalized additive models implement in the "gam" function in the 'mgcv' R package. RNA velocity analysis was performed using Velocity using default parameters.

GO enrichment of cluster markers and differentially expressed genes was performed using the R package 'clusterProfiler'⁴⁴ with a Benjamini-Hochberg multiple testing adjustment and a false-

discovery rate cut-off of 0.1, using all expressed genes expressed in >3 cells as background. Visualization was performed using the R packages 'ggplot2' and 'igraph'.

The collection of DNA samples, genotype calling, quality control and imputation of the BSSH-GODD cohort has previously been described¹¹. Similarly, information pertaining to the UKBiobank genotype and phenotype data has been previously described⁴⁵. Additional QC thresholds were implemented to the UKBiobank genotyping data to remove potential spurious variants and samples, namely 1) call rate <98%, 2) heterozygosity > 3 standard deviation from the mean, 3) individuals with aneuploidy or demonstrating a discrepancy between self-reported sex and genetically inferred sex and 4) non white British, 5) SNP doesn't conform the Hardy-Weinberg equilibrium ($p < 0.004$), and 6) MAF < 1%. The UKBiobank imputed data is a combination of the Haplotype Reference consortium and the UK 10K/1K Genomes Phase 3 panel comprising 92,693,895 variants and 487,442 individuals, which was reduced to 401,667 after QC. A total of 3736 Dupuytren disease patients were identified within the UKBiobank using a combination of diagnostic codes; ICD-10 (M720, M7204), OPCS (T521, T522, T525, T526, T561, T562), non-cancer self-reported illness code 20002 (1544), and self-reported operation code 20004 (1535), and were age, sex and genotyping platform matched at a ratio of 1 to 5 to select 18680 controls. The summary association statistics of both datasets were calculated using BOLT-LMM⁴⁶, and variants with an info score < 0.9, a standard error > 5 or an estimated minor allele count in cases < 3 were removed prior meta-analysis, which was performed using the inverse variance weighted method provided in GWAMA⁴⁷.

To generate a normalised count from the single cell RNA expression data, raw counts of each gene of the same cell type were amalgamated and subsequently normalised using DESeq2⁴⁸. Associations between the normalised RNA expression and DD associative variants were identified using SNPsea⁴⁹.

Flow cytometry

After tissue disaggregation into a single cell suspension, cells were first stained with a panel of fluorescently labelled antibodies to surface antigens, washed with FACS wash buffer (1% BSA, 0.01% NaN₃ in PBS) then fixed using CytoFix (eBiosciences Foxp3 staining buffer set # 00-5523-00) for a minimum of 30 min at 4°C. After 3 washes in perm wash (eBioscience Foxp3 staining buffer), intracellular antigens were stained with another panel of fluorescently labelled antibodies, and cells were then washed with perm wash and analysed by flow cytometry (BD LSR Fortessa X20) and FlowJo software. BD comp beads (anti-mouse Ig k #552843) were used to establish compensation settings. Dead cells were deselected using live/dead stain, added to surface staining panel prior to cell fixation (Life Technologies Live/Dead™ Near IR fixable dead cell stain kit #L10119). Isotype controls were used during antibody optimization.

FACS Sorting

Dupuytren's myofibroblasts were stained without prior magnetic bead enrichment into CD82^{high} and CD82^{low} based on the top and bottom 30% expressing cells, respectively. Samples were filtered through a 70 µm strainer before sorting commenced. Single cell sorting was performed using a FACS Aria (BD Biosciences). After doublet exclusion, isolated single cells were sorted into 5ml FACS tubes containing 1ml of complete medium (5% DMEM + 1% pen/strep). Cell purity was assessed for each sorted population and was approximately 95%. Sorted stromal cell populations were seeded onto hydrogels for traction force microscopy.

Mass cytometry

Where possible antibodies were purchased from Fluidigm, (San Francisco , CA) or were labelled using metal tags using Maxpar antibody labelling kits (Fluidigm). All antibodies were titrated and used at a concentration between (0.25 and 0.5 ug/ml). For each sample, 1-3 million cells were first stained with a solution containing rhodium DNA intercalator (Fluidigm) to distinguish live/dead, prior to Fc receptor

blocking (Miltenyi Biotec). Samples were then stained with a mixture of metal conjugated antibodies recognising cell surface antigens (see staining panel). After washing in Maxpar cell staining buffer (Fluidigm), samples were fixed and permeabilised using the FoxP3 transcription factor staining buffer kit (ThermoFisher) prior to washing and incubation with metal conjugated antibodies recognising intracellular antigens. Samples were washed twice in cell staining buffer, fixed by incubation with 1.6% PFA (Pierce) for 10 minutes and finally incubated overnight with iridium DNA intercalator in Maxpar fix and perm buffer (Fluidigm). Prior to acquisition samples were washed twice in Maxpar cell staining buffer and twice in Maxpar water and filtered through a 40 um cell strainer before being acquired on a Helios mass cytometer (Fluidigm).

After acquisition, all .fcs files in the experiment were normalised using tools within the Helios software and then uploaded to Cytobank (www.cytobank.org) for all gating and further analysis including using the clustering and dimensionality reduction algorithm viSNE.

CyTOF Antibodies

CD252 169-Tm (3166007B, ML5, Fluidigm), CD34 166-Er (Catalog:3166012B, Clone:581, Fluidigm), CD45 141-Pr (Catalog: 3141009B, Clone: HI30, Fluidigm), HLA-D A, B C 142-Nd (Catalog: 3142007B, Clone: HCD57, Fluidigm), CD19 143-Nd (Catalog: 3144007A, Clone:NP6G4, Fluidigm), HLA-DR 89-Y (Catalog: 3173005B, L243, Fluidigm), CD3 189-Y (Catalog:3158021A, Clone: 24E10, Fluidigm), B-catenin 176-Lu (Catalog:3147005A, Clone:D10A8, Fluidigm), CD55 174-Yb (Catalogue; 3148015B, Clone: JS11, Fluidigm), CD146 155-Nd (3155006B, P1H12, Fluidigm), CD29 156-Gd (3156007B, TS2/16, Fluidigm), CD82 158Gd (3158025B, ASL-24, Fluidigm), CD90 172-Yb (3173011B, 5E10, Fluidigm), TGFb 150-Nd (3163010B, CloneTW46H10, Fluidigm) CD163 145-Nd (3145010B, GHI/61, Fluidigm), Ki-67 168-Er (3168007B, B56, Fluidigm), CD54 170-Er (3170014B, HA58, Fluidigm), CD68 171-Yb (3171011B, Y1/82A, Fluidigm), CD9 172-Yb (Cat. No. 312102, Biolegend), CD252 150-Nd (326302, Biolegend), PGLA2 172-Yb (MAB5374, R&D Systems), a-SMA 156-Gr (MAB1420, 1A4, R&D Systems), gp38 155-Gd

(AF3670-SP, R&D Systems), PDGFRB 154-Sm (SAB4700458, 18A2, Sigma Aldrich), MMP14 149-Sm (MAB9181-SP, 128527 R&D Systems), Cad-11 148-Nd (AF1790-SP, R&D Systems) and FAP 147-Sm (MAB3715-SP, 427819 RD Systems).

Immunohistochemistry

Dupuytren's tissue samples were fixed with 4% paraformaldehyde in PBS for 20 min, longitudinally bisected, and embedded in paraffin wax, and 7- μ m sections from the cut surface were processed for immunohistochemistry¹⁷. Sequential sections were stained with mouse monoclonal antibodies. Antibodies were detected using a two-staged polymer enhancer system (Sigma). Murine or Rabbit IgG isotypes at the same protein concentration as the monoclonal antibody solution were used as a control. Images were acquired with an Olympus BX51 Microscope (Olympus).

Immunofluorescence and confocal microscopy

Dupuytren's tissue samples were fixed with 4% paraformaldehyde in PBS for 20 min, longitudinally bisected, and embedded in paraffin wax, and 7- μ m sections from the cut surface were processed for immunofluorescence¹⁷. Then, the tissue sections were stained with antibodies listed above followed by incubation with fluorescent-dye-conjugated secondary antibodies (Life Technologies). Nuclei were counterstained with DAPI (4, 6-diamidino-2-phenylindole; Sigma-Aldrich) and mounted using Prolong[™] Gold anti-fade (Life Technologies). Fluorescent images were obtained with a confocal system (Zeiss LSM 710).

Traction force microscopy

Polyacrylamide (PAA) hydrogels for traction force microscopy were prepared as previously described⁵⁰. Briefly, acid washed glass coverslips (18mm) were incubated with poly-L-lysine (10 μ g/ml in H₂O) for 30 mins at 4 °C, then coated with 0.04 μ m carboxylate-modified red FluoSpheres (1:5000 in ddH₂O). PAA gel formation was initiated with APS (10% solution in ddH₂O, Sigma) and *TEMED*

(Sigma). Polymerized PAA gels were functionalized with sulfo-SANPAH (Invitrogen) and coated with Type 1 Collagen (200ug/ml Rat Tail, Thermo Fisher). The Young's modulus of the PAA gels were 2.55 ± 0.5 KPa. Cells were allowed to adhere to the gel for 4-6 hours before image acquisition. Images were captured using a x40 Plan Apo objective on a Zeiss LSM 710 Confocal microscope and the experiments were performed at 37 °C and 5% CO₂ in Phenol red free DMEM (Thermo Fisher) in a microscope stage incubation chamber. Bead positions were acquired before and after the addition of trypsin which removed cells from the gel surface. Bead displacement was tracked with an ImageJ PIV plugin⁵¹ and cellular forces reconstructed using a FTTC algorithm also implemented in ImageJ^{51,52}.

Downstream analysis and visualization was performed using MATLAB (Mathworks) and R (R Version 3.5). Single cell force maps of CD82^{high} and CD82^{low} sorted cells were first normalized per cell area and scaled before PCA implemented in the "prcomp" function. Significant components were determined as described above for the single cell expression data using visualization of a scree plot and/or the Jackstraw procedure. Significant PCs were used as input for tSNE clustering using the Barnes-Hut implementation in Rtsne() function.

Transfections

For transfection experiments, cells were seeded at 2×10^5 /6 well plate and the following day transfected with siRNA against CD82 (AM16708) or non-targeting control oligo at 1-20nM (Life Technologies). All transfections were performed with Dharmafect 1 (GE Healthcare) and Optimem™ (Life Technologies) according to the manufacturer's instructions. Media was replaced 2 hrs later with 5% FBS phenol red free DMEM (Gibco). Experiments were performed 6 days post transfection.

Chemotaxis Assay

5 x 10⁵ THP-1 cells (ATCC) were allowed to adhere for 2 hrs to a Corning Transwell® plate with 8µM pore in 1% FBS DMEM, the reservoir plate was then replaced with conditioned media generated from freshly sorted DD fibroblasts. For the last 3 days of the assay, the media was replaced with 1% FBS, DMEM. Supernatants harvested, filtered and used neat, alongside CCL2 (2 ng/ml) (Peprotech, London UK) as a positive control. Cells were allowed to migrate for 6 hrs, the number of migrated cells were counted in the reservoir plate following the addition of Hoechst 3342 Fluorescent Stain (Life Technologies) at 0.1µg/ml for 1 hr using the Celigo Imaging Cytometer, Nexcelom Bioscience (Lawrence, MA).

Bulk RNA-seq library preparation

Passage 2 DD myofibroblasts from 7 donors were transfected with siRNA against *CD82* (AM16708) or non-targeting control oligo at 1-20nM (Life Technologies). RNA was extracted using Direct-Zol™ RNA kit. The library preparation was started with the NEBNext® Poly(A) mRNA Magnetic Isolation Module followed by the NEBNext® Ultra™ Directional RNA Library Prep Kit for Illumina with the NEBNext® Multiplex Oligos for Illumina® (Index Primers Set 1) and (Index Primers Set 2). The concentration of each library was determined using the NEBNext® Library Quant Kit for Illumina® and High Sensitivity D1000 Screentape Bioanalyzer (Agilent).

Bulk RNA-seq data analysis

FASTQ files were assessed using FASTQC followed by the generation of TPM values with kallisto v0.42.4⁵³. TPM values were summed to obtain gene-level expression values using tximport and differential expression analysis was undertaken with DESeq2⁵⁴. GO enrichment of differentially expressed genes was performed using the R package 'clusterProfiler'⁴⁴ with a Benjamini-Hochberg multiple testing adjustment and a false-discovery rate cut-off of 0.1. Visualization was performed using the R packages 'ggplot2' and 'igraph'.

Data availability

Data supporting the findings of this study are available within the paper and its Supplementary Information files. The source data underlying all figures are provided as a Source Data file, and are further available from the authors upon reasonable request. RNA-Seq are deposited in SRA under accession number PRJNA607098 (<https://www.ncbi.nlm.nih.gov/sra/PRJNA607098>)

References

1. Wynn, T.A. & Ramalingam, T.R. Mechanisms of fibrosis: therapeutic translation for fibrotic disease. *Nature medicine* **18**, 1028-1040 (2012).
2. Wynn, T.A. Cellular and molecular mechanisms of fibrosis. *The Journal of pathology* **214**, 199-210 (2008).
3. Mora, A.L., Rojas, M., Pardo, A., *et al.* Emerging therapies for idiopathic pulmonary fibrosis, a progressive age-related disease. *Nature Reviews Drug Discovery* **16**, 755 (2017).
4. Shook, B.A., Wasko, R.R., Rivera-Gonzalez, G.C., *et al.* Myofibroblast proliferation and heterogeneity are supported by macrophages during skin repair. *Science* **362**, eaar2971 (2018).
5. Mizoguchi, F., Slowikowski, K., Wei, K., *et al.* Functionally distinct disease-associated fibroblast subsets in rheumatoid arthritis. *Nature Communications* **9**, 789 (2018).
6. Schneider, R.K., Mullally, A., Dugourd, A., *et al.* Gli1(+) Mesenchymal Stromal Cells Are a Key Driver of Bone Marrow Fibrosis and an Important Cellular Therapeutic Target. *Cell stem cell* **20**, 785-800.e788 (2017).
7. Xie, T., Wang, Y., Deng, N., *et al.* Single-Cell Deconvolution of Fibroblast Heterogeneity in Mouse Pulmonary Fibrosis. *Cell reports* **22**, 3625-3640 (2018).
8. Nanchahal, J. & Hinz, B. Strategies to overcome the hurdles to treat fibrosis, a major unmet clinical need. *Proceedings of the National Academy of Sciences of the United States of America* **113**, 7291-7293 (2016).
9. Lanting, R., Broekstra, D.C., Werker, P.M.N., *et al.* A Systematic Review and Meta-Analysis on the Prevalence of Dupuytren Disease in the General Population of Western Countries. *Plastic and Reconstructive Surgery* **133**, 593-603 (2014).
10. Blondel, V.D., Guillaume, J.-L., Lambiotte, R., *et al.* Fast unfolding of communities in large networks. *Journal of Statistical Mechanics: Theory and Experiment* **2008**, P10008 (2008).
11. Ng, M., Thakkar, D., Southam, L., *et al.* A Genome-wide Association Study of Dupuytren Disease Reveals 17 Additional Variants Implicated in Fibrosis. *Am J Hum Genet* **101**, 417-427 (2017).
12. Slowikowski, K., Hu, X. & Raychaudhuri, S. SNPsea: an algorithm to identify cell types, tissues and pathways affected by risk loci. *Bioinformatics* **30**, 2496-2497 (2014).
13. Verjee, L.S., Midwood, K., Davidson, D., *et al.* Myofibroblast Distribution in Dupuytren's Cords: Correlation With Digital Contracture. *Journal of Hand Surgery* **34**, 1785-1794.
14. Subramanian, A., Tamayo, P., Mootha, V.K., *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 15545-15550 (2005).
15. van Beuge, M.M., Ten Dam, E.-J.P.M., Werker, P.M.N., *et al.* Matrix and cell phenotype differences in Dupuytren's disease. *Fibrogenesis & tissue repair* **9**, 9-9 (2016).
16. Verjee, L.S., Verhoekx, J.S.N., Chan, J.K.K., *et al.* Unraveling the signaling pathways promoting fibrosis in Dupuytren's disease reveals TNF as a therapeutic target. *Proceedings of the National Academy of Sciences* **110**, E928-E937 (2013).
17. Verjee, L.S., Midwood, K., Davidson, D., *et al.* Myofibroblast distribution in Dupuytren's cords: correlation with digital contracture. *The Journal of hand surgery* **34**, 1785-1794 (2009).
18. Verhoekx, J.S.N., Beckett, K.S., Bisson, M.A., *et al.* The mechanical environment in dupuytren's contracture determines cell contractility and associated MMP-mediated matrix remodeling. *Journal of Orthopaedic Research* **31**, 328-334 (2013).
19. Lagares, D., Ghassemi-Kakroodi, P., Tremblay, C., *et al.* ADAM10-mediated ephrin-B2 shedding promotes myofibroblast activation and organ fibrosis. *Nature medicine* **23**, 1405 (2017).
20. Liu, F., Lagares, D., Choi, K.M., *et al.* Mechanosignaling through YAP and TAZ drives fibroblast activation and fibrosis. *Am J Physiol Lung Cell Mol Physiol* **308**, L344-357 (2015).

21. Coifman, R.R., Lafon, S., Lee, A.B., *et al.* Geometric diffusions as a tool for harmonic analysis and structure definition of data: Diffusion maps. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 7426 (2005).
22. Angerer, P., Haghverdi, L., Büttner, M., *et al.* destiny: diffusion maps for large-scale single-cell data in R. *Bioinformatics* **32**, 1241-1243 (2015).
23. Petropoulos, S., Edsgard, D., Reinius, B., *et al.* Single-Cell RNA-Seq Reveals Lineage and X Chromosome Dynamics in Human Preimplantation Embryos. *Cell* **165**, 1012-1026 (2016).
24. Lagares, D., Santos, A., Grasberger, P.E., *et al.* Targeted apoptosis of myofibroblasts with the BH3 mimetic ABT-263 reverses established fibrosis. *Science translational medicine* **9**(2017).
25. Gieseck, R.L., 3rd, Ramalingam, T.R., Hart, K.M., *et al.* Interleukin-13 Activates Distinct Cellular Pathways Leading to Ductular Reaction, Steatosis, and Fibrosis. *Immunity* **45**, 145-158 (2016).
26. Lodyga, M., Cambridge, E., Karvonen, H.M., *et al.* Cadherin-11-mediated adhesion of macrophages to myofibroblasts establishes a profibrotic niche of active TGF- β . *Science Signaling* **12**, eaao3469 (2019).
27. Rinkevich, Y., Walmsley, G.G., Hu, M.S., *et al.* Identification and isolation of a dermal lineage with intrinsic fibrogenic potential. *Science* **348**, aaa2151 (2015).
28. Martinez, F.J., Collard, H.R., Pardo, A., *et al.* Idiopathic pulmonary fibrosis. *Nature Reviews Disease Primers* **3**, 17074 (2017).
29. Reyfman, P.A., Walter, J.M., Joshi, N., *et al.* Single-Cell Transcriptomic Analysis of Human Lung Provides Insights into the Pathobiology of Pulmonary Fibrosis. *American journal of respiratory and critical care medicine* **199**, 1517-1536 (2019).
30. Shibagaki, N., Hanada, K., Yamashita, H., *et al.* Overexpression of CD82 on human T cells enhances LFA-1 / ICAM-1-mediated cell-cell adhesion: functional association between CD82 and LFA-1 in T cell activation. *European journal of immunology* **29**, 4081-4091 (1999).
31. Lebel-Binay, S., Lagaudriere, C., Fradelizi, D., *et al.* CD82, tetra-span-transmembrane protein, is a regulated transducing molecule on U937 monocytic cell line. *Journal of leukocyte biology* **57**, 956-963 (1995).
32. Spring, F.A., Griffiths, R.E., Mankelow, T.J., *et al.* Tetraspanins CD81 and CD82 Facilitate $\alpha 4\beta 1$ -Mediated Adhesion of Human Erythroblasts to Vascular Cell Adhesion Molecule-1. *PLOS ONE* **8**, e62654 (2013).
33. Termini, C.M., Cotter, M.L., Marjon, K.D., *et al.* The membrane scaffold CD82 regulates cell adhesion by altering $\alpha 4$ integrin stability and molecular density. *Mol Biol Cell* **25**, 1560-1573 (2014).
34. Wu, J., Liang, S., Bergholz, J., *et al.* Δ Np63 α activates CD82 metastasis suppressor to inhibit cancer cell invasion. *Cell Death & Disease* **5**, e1280-e1280 (2014).
35. Vianay, B., Senger, F., Alamos, S., *et al.* Variation in traction forces during cell cycle progression. *Biol Cell* **110**, 91-96 (2018).
36. Goffin, J.M., Pittet, P., Csucs, G., *et al.* Focal adhesion size controls tension-dependent recruitment of alpha-smooth muscle actin to stress fibers. *J Cell Biol* **172**, 259-268 (2006).
37. Verjee, L.S., Verhoekx, J.S., Chan, J.K., *et al.* Unraveling the signaling pathways promoting fibrosis in Dupuytren's disease reveals TNF as a therapeutic target. *Proceedings of the National Academy of Sciences of the United States of America* **110**, E928-937 (2013).
38. Leek, J.T., Johnson, W.E., Parker, H.S., *et al.* The sva package for removing batch effects and other unwanted variation in high-throughput experiments. *Bioinformatics (Oxford, England)* **28**, 882-883 (2012).
39. Macosko, Evan Z., Basu, A., Satija, R., *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015).
40. Chung, N.C. & Storey, J.D. Statistical significance of variables driving systematic variation in high-dimensional data. *Bioinformatics (Oxford, England)* **31**, 545-554 (2015).
41. Tirosh, I., Izar, B., Prakadan, S.M., *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science* **352**, 189 (2016).

42. Angerer, P., Haghverdi, L., Buttner, M., *et al.* destiny: diffusion maps for large-scale single-cell data in R. *Bioinformatics* **32**, 1241-1243 (2016).
43. Hastie, T. & Stuetzle, W. Principal Curves. *Journal of the American Statistical Association* **84**, 502-516 (1989).
44. Yu, G., Wang, L.-G., Han, Y., *et al.* clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics : a journal of integrative biology* **16**, 284-287 (2012).
45. Bycroft, C., Freeman, C., Petkova, D., *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203-209 (2018).
46. Loh, P.R., Tucker, G., Bulik-Sullivan, B.K., *et al.* Efficient Bayesian mixed-model analysis increases association power in large cohorts. *Nat Genet* **47**, 284-290 (2015).
47. Magi, R. & Morris, A.P. GWAMA: software for genome-wide association meta-analysis. *BMC Bioinformatics* **11**, 288 (2010).
48. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
49. Slowikowski, K., Hu, X. & Raychaudhuri, S. SNPsea: an algorithm to identify cell types, tissues and pathways affected by risk loci. *Bioinformatics* **30**, 2496-2497 (2014).
50. Colin-York, H., Eggeling, C. & Fritzsche, M. Dissection of mechanical force in living cells by super-resolved traction force microscopy. *Nature protocols* **12**, 783-796 (2017).
51. Martiel, J.-L., Leal, A., Kurzawa, L., *et al.* Chapter 15 - Measurement of cell traction forces with ImageJ. in *Methods in Cell Biology*, Vol. 125 (ed. Paluch, E.K.) 269-287 (Academic Press, 2015).
52. Butler, J.P., Tolic-Norrelykke, I.M., Fabry, B., *et al.* Traction fields, moments, and strain energy that cells exert on their surroundings. *American journal of physiology. Cell physiology* **282**, C595-605 (2002).
53. Bray, N.L., Pimentel, H., Melsted, P., *et al.* Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology* **34**, 525 (2016).
54. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

Acknowledgments

We are grateful to our clinical collaborators Christopher Bainbridge, David Warwick, Lorraine Harry and Dominique Davidson for provision of tissue samples from their patients. We thank the Wellcome Trust Centre for Human Genetics for preparation of the single cell libraries. In addition, we are grateful to the Imaging Centre at the Kennedy Institute of Rheumatology for their support in microscopy.

Author contributions

JN, SS, WX, M Feldmann and TL conceived and designed the study. TL, LW, MN, DF, FM, MC, MZ, CY, M Fritsche and HCY performed and contributed to the analysis of the experiments. JN and TL wrote the manuscript with contribution from all co-authors.

Competing interests

JN and M Feldmann are co-founders and have equity in 180 Therapeutics, which has exclusively licensed IP from the University of Oxford for the treatment of Dupuytren's disease. None of the other authors declare any competing interests.

Figure Legends

Fig. 1. Single cell profiling of developing human fibrosis

a, Schematic illustrating experimental protocol for single cell profiling of Dupuytren's nodules with representative immunohistochemistry image showing α -SMA protein expression in nodules. **b-c**, tSNE projections of single cell RNA-seq for 36,864 cells from the first batch ($n = 6$ DD patients) showing major cell types (**a**) and fibroblasts (*CXCL14* expression) and myofibroblasts (*ACTA2* expression) (**b**) in Dupuytren's nodules. Scale bar in scaled $\log(\text{UMI}+1)$. Cycling MFB represents cycling myofibroblasts. **d**, Heatmap of single cell RNA-seq showing z-score normalized mean expression of cell type marker genes in nodular cells. ($n = 12$ DD patients). **e**, tSNE projections of CyTOF analysis from representative Dupuytren's nodule, coloured by normalized protein expression of cell type markers ($n = 6$ DD patients).

Fig. 2. A dynamic and immunoregulatory ICAM1⁺ fibroblast in fibrosis

a, Heatmap of single cell RNA-seq showing z-score mean expression of genes driving GO signatures in cell types with exemplar GO terms annotated. GO analysis performed on significant cell type markers (2-sided Wilcoxon Rank sum Test, FDR correction, $p\text{-adjust} < 0.01$). **b**, tSNE projection of fibroblasts coloured by major subsets ($n = 12$ DD patients) in single cell RNA-seq. **c**, tSNE projections of CyTOF analysis for representative DD nodule showing three major fibroblasts subsets ($n = 6$ DD patients). Scale bar represents normalized protein expression. **d**, Heatmap of single cell RNA-seq showing mean z-score expression of top 10 marker genes for each fibroblast subset ($n = 12$ DD patients). **e**, tSNE projections of single cell RNA-seq showing fibroblasts coloured by *ICAM1* and *IL6* expression in scaled ($\log(\text{UMI}+1)$) ($n = 12$ DD patients). **f**, Representative density plots of flow cytometry analysis showing ICAM1⁺ fibroblast in freshly isolated cells from matched DD nodule and cord ($n = 8$ DD patients). **g**, Dot plot of chemotaxis assay showing migration of THP-1 cells after incubation with ICAM1^{-/+} sorted populations ($n=11$ DD patients). 2-sided paired t -test, $p\text{-value} = 0.0051$. **h**, Box and whisker plots of

flow cytometry analysis showing the percentage of cells (proportion) for fibroblast subsets (CD34⁺, ICAM1⁺ & PDPN⁺) in Dupuytren's nodules and cords as a proportion of total fibroblasts. 2-sided unpaired *t*-test, mean \pm SEM (*n* = 8 DD patients). ICAM1 range 3-32%, median 8%, box bounds 5-17%. CD34 range 13-50%, median 25%, box bounds 22-30%. PDPN range 42-63%, median 48%, box bounds 45-55%. Box bounds are first and third quantiles.

Fig. 3. Distinct myofibroblast states along an activation continuum

a, Force directed graph (Fruchterman-Reingold layout) of single cell RNA-seq showing myofibroblasts coloured by major subsets (*n* = 12 DD patients). MFB represents myofibroblast. **b**, Heatmap of single cell RNA-seq showing the z-score expression of top 10 myofibroblast subset markers. 2-sided Wilcoxon Rank Sum Test with FDR correction (BH correction (*n* = 12 DD patients)). **c**, tSNE projections of single cell RNA-seq showing myofibroblasts coloured by the expression of *ACTA2* and *MKI67* in scaled $\log(\text{UMI}+1)$ (*n* = 12 DD patients). **d**, Violin plots showing gene expression of *MKI67* and *ITGB1* in fibroblasts and myofibroblasts in scaled($\log(\text{UMI}+1)$) from single cell RNA-seq. **e**, Box and whisker plot of flow cytometry analysis showing Ki67 protein expression in myofibroblasts ITG β 1^{high} myofibroblasts (range 21-46%, mean 28%, box bounds 24-27% representing first to third quantiles) and ITG β 1^{low} fibroblasts (range 0-0%, mean 0.0%, percentiles 0%). 2-sided unpaired *t*-test, *p*-value = 0.00014, mean \pm SEM. (*n* = 8 DD patients). **f**, Dot plot of single cell RNA-seq showing pathways enriched in myofibroblast subsets. Gene ratio is number of marker genes associated with pathway and *p.adjust* is adjusted *p*-value (2-sided Wilcoxon Rank Sum test, BH FDR-correction). **g**, tSNE projections of CyTOF analysis for representative DD patient showing distinct CD82^{high}OX40L⁺ myofibroblast. Scale bar is normalized protein expression. (*n* = 6 DD patients). **h**, Confocal images of immunofluorescence showing co-expression of CD82, α -SMA and ITG- β 1 in DD nodules (*n* = 3 DD patients). Scale bar 20 μ m.

Fig. 4. Functional heterogeneity of myofibroblasts in human fibrosis

a, Line plot demonstrating *CD82* expression in myofibroblasts (*loess* smoothed normalized counts \pm SE) along pseudotime (myofibroblast activation trajectory) in single cell RNA-seq ($n=12$ DD patients).

b, Above, Heatmaps of traction force microscopy analysis showing magnitude of traction force (Pascals) in freshly isolated FACS sorted myofibroblasts ($CD82^{\text{high}}$ & $CD82^{\text{low}}$). Below, Surface plots of traction force microscopy showing representative force foci from sorted myofibroblasts (Scaled to 800 Pa). Color scale represents force range per cell.

c, Bar plots of traction force microscopy analysis showing mean and maximum traction force per cell in $CD82^{\text{high}}$ and $CD82^{\text{low}}$ sorted myofibroblast populations ($n = 3$ DD patients, >15 cells per condition). 2-sided unpaired *t*-test, mean \pm SEM. *p*-values = 0.0056 (max traction), and 0.0098 (mean traction).

d, tSNE projection of traction force microscopy analysis showing clustering of single cell force signatures in sorted populations ($CD82^{\text{high}}$ & $CD82^{\text{low}}$) ($n = 3$ DD patients, >15 cells per condition).

e, Box and whisker plot of flow cytometry analysis showing cell cycle phases in siControl and siCD82 transfected DD myofibroblasts ($n = 3$ DD patients). 2-sided paired *t*-test, *p*-value = 0.023, mean \pm SEM). (siRNA range 2.6–85.5%, mean 32.8%, box bounds 9.75-45.2% represent first and third quantiles. siCD82 range 1.1-87.1%, mean 32.7%, box bounds 3.1-59.8% represent first and third quantiles.)

f, tSNE projections of single cell RNA-seq showing fibroblasts and myofibroblasts in bleomycin murine model marked by the expression of selected genes in scaled log(UMI+1) ($n = 2$ mice).

g, tSNE projection of single cell RNA-seq showing murine fibroblasts and myofibroblasts in bleomycin model coloured by Louvain clusters ($n = 2$ mice).

h, Bar plot of single cell RNA-seq showing the percentage of stromal cells before (Day 0) and after (Day 21) the installation of bleomycin in murine lung fibrosis model.

i, Confocal images of immunofluorescence showing co-expression of *CD82*, α -SMA and PDGFR- α in human IPF ($n= 3$ independent IPF patients). Scale bar = 20 μm .

Fig 1. Single cell profiling of developing human fibrosis




Fig. 2. A dynamic and immunoregulatory ICAM1⁺ fibroblast in fibrosis




Fig. 3. Distinct myofibroblast states along an activation continuum




Fig. 4. Functional heterogeneity of myofibroblasts in human fibrosis

