



Senescence-related myocardial dysfunction: keeping a young heart

Ramzi A. Ajjan ^{1,†}, Robert T.R. Huckstepp ^{2,*,†}, Naveed Akbar ³, Johann Bauersachs ⁴, Jeshier Ching Wai Lok⁵, Juel Choppy-Madeleine⁶, Gary Christopher ⁷, Mayank Dalakoti ^{8,9}, Emily Dawson-Plincke⁵, John Dodds⁶, Georgina M. Ellison-Hughes ¹⁰, Costanza Emanuelli ¹¹, Christopher H. George ¹², Anushka Goyal¹³, Victoria Higginbotham¹², Lorenz Holzner¹⁴, Venkateswarlu Kanamarlapudi ¹², Paolo Madeddu ¹⁵, Claudio Mauro ¹⁶, Fiona Lewis-McDougall ¹⁷, Andrew J. Murray ¹⁸, Renita Peter Damien Genetus¹⁹, Emma Short ^{20,21}, Mark A. Sussman ²², Adriana A.S. Tavares ²³, Samuel Thompson²⁴, Morya Wadodkar¹³, Stuart R.G. Calimport ^{20,25}, and Barry L. Bentley ^{20,26,27}

¹Leeds Institute of Cardiovascular and Metabolic Medicine, University of Leeds, Leeds, UK; ²School of Life Sciences, University of Warwick, Coventry CV4 7AL, UK; ³Division of Cardiovascular Medicine, Radcliffe Department of Medicine, University of Oxford, Oxford, UK; ⁴Department of Cardiology and Angiology, Hannover Medical School, Hannover, Germany; ⁵Medical School, University of East Anglia, Norwich UK; ⁶Medical School, University of Glasgow, Glasgow, UK; ⁷Centre for Ageing and Dementia Research, Swansea University, Swansea, UK; ⁸Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK; ⁹Department of Cardiology, National University Heart Centre, Singapore; ¹⁰Centre for Human and Applied Physiological Sciences, School of Basic & Medical Biosciences, Faculty of Life Sciences & Medicine, King's College London, London, UK; ¹¹National Heart and Lung Institute, Imperial College London, London, UK; ¹²Medical School, Swansea University, Swansea, Wales, UK; ¹³Medical School, University of Bristol, Bristol, UK; ¹⁴Epigenetics, Babraham Institute, Babraham, UK; ¹⁵Bristol Heart Institute, School of Translational Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK; ¹⁶College of Medicine and Health, University of Birmingham, Birmingham, UK; ¹⁷Centre for Microvascular Research, William Harvey Research Institute, Faculty of Medicine and Dentistry, Queen Mary University, London, UK; ¹⁸Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge, UK; ¹⁹Medical School, University of Birmingham, Birmingham, UK; ²⁰Cardiff School of Technologies, Cardiff Metropolitan University, Cardiff, UK; ²¹Department of Cellular Pathology, Swansea Bay University Health Board, Swansea, UK; ²²Biology Department and Integrated Regenerative Research Institute, San Diego State University, San Diego, CA, USA; ²³University-BHF Centre for Cardiovascular Science, Queen's Medical Research Institute, University of Edinburgh, Edinburgh, UK; ²⁴New College, Oxford University, Oxford, UK; ²⁵Collaboration for the Advancement of Sustainable Medical Innovation (CASMI), University College London, London, UK; ²⁶Harvard Medical School, Harvard University, Boston, MA, USA; and ²⁷Department of Surgery, Massachusetts General Hospital, Boston, MA, USA

Received 13 September 2025; revised 4 December 2025; accepted 26 January 2026

* Corresponding author. Email: R.Huckstepp@warwick.ac.uk


† Joint first authors.

© The Author(s) 2026. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Graphical Abstract

Senescence-related myocardial dysfunction



microRNA, micro ribonucleic acid

Ajjan RA, et al. *European Heart Journal*.

The ageing heart undergoes progressive and substantial changes. Myocardial cells enter a senescence-associated secretory phenotype (SASP), associated with alterations in extracellular vesicles (EVs) containing microRNAs, creating an inflammatory environment. These molecular changes, combined with mitochondrial dysfunction, induce myocardial cell hypertrophy, multinucleation, and also result in altered responses to autonomic regulation. Alongside the stress of the continuous workload on the heart over a lifetime, there is a reduction in repair mechanisms, which coupled with fibrosis, steatosis, valve stenosis and compromised blood supply, due to atherosclerosis, ultimately lead to organ dysfunction, and myocardial ageing. Created in Biorender.

Abstract

The heart, a vital organ, works without interruption and constantly adjusts to the ever-changing demands on our body. It adapts to physiological and pathological changes, including exercise and emotional state, as well as metabolic, respiratory, and vascular abnormalities. The pumping action of the heart is determined by the health of the myocardium, which undergoes changes with ageing that are both under-investigated and incompletely understood, potentially impacting our approach to pathological conditions. Here, the alterations in cellular, tissue, and gross physiological function of the heart with age are discussed. At the molecular level, non-coding RNAs influence cellular senescence, and extracellular vesicles induce fibrosis through matrix remodelling. Mitochondrial dysfunction and altered fatty acid oxidation reduce cellular energetics, whilst accumulation of reactive oxygen species and steatosis, as well as telomere shortening coupled with reduced autophagy, limit the myocardium's regenerative capability. Loss of cardiomyocytes, combined with senescence, requires compensatory hypertrophy, inducing myocardial stiffness and altered muscle function. In addition to these direct alterations in myocardial characteristics with ageing, other factors that can affect the myocardium indirectly are addressed, including valve calcification, resulting in regurgitation and/or stenosis; vascular abnormalities, reducing compliance and exacerbating hypertension; fibrosis leading to cardiac arrhythmias; and autonomic dysregulation, reducing cardiac adaptability. Finally, potential modulation of cardiac ageing is discussed whilst also addressing which senescent modifications should be considered as

ageing-related physiological changes of the myocardium. A better understanding of myocardial ageing will differentiate physiological changes from early, preventable, and reversible pathological changes, consequently helping to optimize management of individuals with or at risk of myocardial disease by taking into account diverse trajectories of myocardial ageing.

Introduction

The adult human heart comprises 11 major cell types and contains 3–4 billion cells.¹ Over our lifetime, we generate ~3 billion heartbeats at a substantial energy requirement; turning over ~6 kg of ATP daily, 15–20-fold the heart's weight, supporting myofibrillar contraction, ion pumping, and other functional demands.²

The heart exhibits senescence as it ages. Cellular senescence is the irreversible arrest of the cell cycle and release of a senescence-associated secretory phenotype (SASP), including cytokines, proteases, and microRNAs (miRNAs), often associated with extracellular vesicles (EVs). Tissue senescence, defined as an ageing-related decline in the functional capacity or structure of a tissue,³ follows cell loss and consequently results in morphological alterations, reduced regeneration, and fibrosis of the heart (*Graphical Abstract*); however, differentiating the effects of normal physiological ageing from co-existent pathology can be challenging. Ultimately, these processes may culminate in organ senescence, defined as an ageing-related decline in the functional capacity or structure of an organ.³

Health systems are encountering new challenges and opportunities in addressing the increasing prevalence of ageing-related cardiovascular conditions. Myocardial dysfunction is a critical contributor to morbidity and mortality in older populations, including both diastolic and systolic changes in cardiac function, valvular abnormalities, and alterations in the conduction system increasing the risk of arrhythmias.³ Heart failure in individuals below 55 years is ~1%, increasing to >10% in those above 70 years,⁴ with numbers rising as diagnostics improve. Ageing may lead to progressive cardiac dysfunction, but it may also imply increased susceptibility to pathological conditions as risk factors for cardiac dysfunction are more common with increasing age,⁵ adding further complexity to understanding the effects of physiological age-related changes.

To improve clinical management of the older populations, there is a need for the systematic and comprehensive classification of ageing-related pathologies at metabolic, tissue, organ, and systemic levels.⁶ To address this, the International Consortium to Classify Ageing-Related Pathologies (ICCARP) was established⁷ and has recently defined the criteria for an ageing-related pathology⁷ and reached an international consensus on senescence definitions.⁸ A cardiovascular working group formed from ICCARP members, with broad expertise but a common cardiovascular interest, was formed to write this comprehensive review on age-related functional changes of the myocardium.

Ageing-related changes of cardiomyocytes, the cellular 'machines of contraction'

Morphological changes in cardiomyocytes. Cardiomyocytes constitute the majority of the atrial and ventricular volume but only comprise 30% and 50% of the cells in these tissues, respectively.¹

Catastrophic cardiomyocyte loss is associated with acute and chronic cardiac disease. However, the extent to which cardiomyocyte loss is a feature of normative ageing is relatively underexplored, partly related to the difficulties in separating ageing-related myocyte loss from the aggravated loss secondary to cardiac disease.⁹ Left ventricular and septal mass decreases by ~0.5% per year, and given the exceptionally low cardiomyocyte loss with age, other changes must underpin the observed reduction in heart mass.^{10,11} Even with extreme longevity, the number of human cardiomyocytes renewed over the life course is likely fewer than half of the perinatal population. Although this renewal rate decreases with age, cardiomyocyte numbers are largely constant across a CVD-free lifetime.¹² Whilst the mode of cardiomyocyte exchange in humans is not yet known, in mice new cardiomyocytes are generated predominantly by division of pre-existing cardiomyocytes¹³ and not through transdifferentiation processes or *de novo* cardiomyocyte generation.¹⁴ Following myocardial ischaemia, however, myocyte proliferation increases within and adjacent to the injury site in human and mouse hearts, and therefore the presence of pathology affects cardiomyocyte proliferation

A feature of human ageing is cardiomyocyte hypertrophy.¹¹ Senescent cardiomyocytes, with impaired contractile output, can accumulate in the ageing myocardium, and loss of 'functional cells' redistributes workload placing a greater contractile demand upon remaining cardiomyocytes, leading to hypertrophy.¹⁵ Moreover, the long-term persistence of senescent cells disrupts tissue structure and function with deleterious paracrine, autocrine, and systemic effects in humans and animals,¹⁶ remaining part of the tissue architecture but are functionally defunct.¹⁷

Human cardiomyocytes are normally polyploid, and increased polyploidization correlates with cardiomyocyte enlargement and variable cellular DNA content.¹⁸ However, the relative proportions of mono and multinucleated cardiomyocytes can remain stable.¹⁹ Whilst human ageing promotes cardiomyocyte hypertrophy, some pathological states associated with ageing have similar effects,²⁰ such as hypertension, making the distinction between physiological and pathological age-related changes challenging.

Importantly, time- and age-dependent changes in cardiomyocytes are often represented in linear relationship models. However, many studies lack adequate power to reveal complex non-linear behaviour, and age-linked dynamic processes are likely characterized by non-linear behaviour exhibiting thresholds or 'tipping points', which remains an area for active research.²¹

Molecular changes in cardiac ageing

DNA methylation is associated with cardiovascular ageing.²² Cardiovascular risk factors, such as smoking and hyperhomocysteinaemia, interact with ageing, inducing dysregulated DNA methylation. The methylation status of CpG sites forms an epigenetic clock to measure human ageing rates. DNA methylation-based estimation correlates well with cardiovascular disease (CVD) incidence,²³ and dysregulation of epigenetic regulators in




Figure 1 Ageing-related myocardial cellular and molecular alterations. Cellular changes occur with myocardial ageing, encompassing cardiomyocyte hypertrophy, cell senescence, mitochondrial alterations, impaired autophagy and mitophagy, increased fibrosis, and vascular alterations, including arterial stiffening and endothelial dysfunction. These cellular alterations are governed by molecular mechanisms involving increased senescence-associated secretory phenotype (SASP) and pro-inflammatory factors, up-regulation and down-regulation of different microRNAs, increased oxidative stress and reactive oxygen species (ROS), telomere attrition, polyploidy of cardiomyocytes, increased tissue inhibitors of metalloproteinases (TIMPs), and decreased matrix metalloproteinases (MMPs). Extracellular vesicles of myocardial cells release SASP and pro-inflammatory factors, and miRNAs. These molecular alterations directly and indirectly influence the cellular changes contributing to myocardial ageing. The up red and down black arrows indicate an increase or decrease, respectively. Created in Biorender.

histone modifications predisposes to cardiac ageing. Down-regulation of genes clustered in oxidative stress and gene transcription regulation is associated with epigenetic changes, including decreased H3K9ac, increased H3K9me3, and H3K27me3,²⁴ which are implicated in myocardial hypertrophy.

MicroRNAs are post-transcriptional regulators of gene expression that target specific mRNAs for promoting degradation or translational arrest, thus influencing inflammatory, fibrotic, and senescence pathways, with the first miRNA inhibitor currently trialled for heart failure.²⁵ Arterial stiffening, endothelial dysfunction, and increased oxidative stress are all pathological processes and part of normative ageing, controlled by miRNAs. For example, miR-21-5p, miRNA-22-3p, and miRNA-22-5p promote fibrosis and cardiovascular remodelling,^{26,27} whilst miR-92a-3p, miR-146a-5p, and miR-126 are essential for maintaining endothelial integrity and function^{28–30} and miRNA-34a regulates cardiac ageing and vascular senescence.^{31,32} Cardiovascular-enriched miRNAs may act as minimally invasive biomarkers of age; e.g. circulating miR-146a-5p, miR-126-3p, miR-21-5p, and miR-181a-5p progressively increase with age, except in ultra-centenarians in whom levels are comparable to young individuals.³³

The interaction of miRNAs with long non-coding RNAs (lncRNAs) and circular RNAs (circRNAs) adds further complexity to the regulation of cardiovascular ageing. CircRNAs sequester

miRNAs, modulating their availability to target mRNAs. circRNA–miRNA–mRNA networks associated with ageing-related macular degeneration,³⁴ and lncRNA–miRNA–mRNA networks are also involved in age-associated CVD.³⁵ A summary of cardiomyocyte changes with ageing is provided in *Figure 1*.


Ageing-related alterations in myocardial energetics and metabolism

Myocardial energetics

In humans, myocardial energetic reserve declines with age,^{36,37} through either a proportional depletion of both ATP and phosphocreatine (PCr) contents³⁶ or a fall in the phosphocreatine-to-ATP ratio (PCr/ATP).³⁷ The ageing-related decline in PCr/ATP occurs alongside reduced left ventricular mass and ejection fraction³⁷ and correlates with diastolic dysfunction. Lower PCr in the myocardium may result from impaired creatine kinase activity.³⁸

Substrate metabolism

ATP requirements are met through oxidation of glucose, ketone bodies, lactate, and fatty acids.³⁹ Myocardial fatty acid oxidation



declines with age, suggesting increased reliance on glucose metabolism,⁴⁰ coupled with mitochondrial impairment.⁴¹ Altered fatty acid oxidation may relate to ageing-related down-regulation of the fatty acid-activated transcription factor PPAR α , and its downstream targets *Cpt1* and *HADH*, as well as lower *HADH* activity.⁴² Mechanistically, cardiac-specific deletion of *GCN5L1* (a mitochondrial acetyltransferase) restores fatty acid (and glucose) oxidation, consequently rescuing diastolic function.⁴³ Impaired fatty acid oxidation, increased circulating lipids, and myocardial fatty acid uptake can contribute to lipid accumulation in the aged myocardium.⁴⁴ This can cause metabolic and functional changes, including myocardial triacylglycerol accumulation, which correlates with ageing-related loss of diastolic function⁴⁵ and, in rodents, mitochondrial alterations.⁴⁶

The plasma membrane fatty acid transporter CD36 is implicated in ageing-related steatosis⁴⁶ and myocardial lipotoxicity.⁴⁷ The accumulation of toxic lipid species may also occur with ageing⁴⁴ and is linked to altered nerve signalling⁴⁴ and increased myocardial insulin resistance.⁴⁸ In ageing populations, hearts exhibit impaired anaplerotic response to insulin stimulation,⁴¹ and depletion of protein and mRNA of the insulin-responsive glucose transporter, GLUT-4⁴⁹ leads to impaired insulin-stimulated glucose uptake.⁵⁰ Insulin resistance and impaired myocardial glucose uptake provide additional challenges to cardiac substrate metabolism in later life, occurring secondary to and alongside suppressed fatty acid oxidation.⁵¹

Mitochondria

Ageing-associated mitochondrial impairments include lower tricarboxylate cycle enzyme activities,⁵² diminished oxidative

capacity,⁵³ and increased reactive oxygen species (ROS).⁵⁴ A population of cytochrome c oxidase [electron transport chain (ETC) complex IV]-deficient cardiomyocytes increases with advanced age.⁵⁵ Lower cellular respiration rates supported by substrates for Complexes III⁵⁶ and IV⁵⁷ are seen in the intermyofibrillar mitochondria (IFM) subpopulation, with the subsarcolemmal mitochondria (SSM) relatively unaffected.⁵⁷ Unexpectedly, enzyme activities of Complexes I, II, III, and IV are preserved in older human hearts,⁵⁸ highlighting that not all aspects of mitochondrial function decline with age. The discrepancy between these findings and reports of lower mitochondrial oxidative capacities can be explained by decreased assembly of mitochondrial supercomplexes in the aged heart,⁵⁹ compromising electron transfer, consequently resulting in functional impairment.

Role of oxidative stress

Markers of oxidative stress show age-dependent accumulation in the myocardium, including increased lipid peroxidation,^{54,58} mitochondrial DNA (mtDNA) oxidation, and mutational load,⁶⁰ with increased mitochondrial biogenesis as a potential compensatory mechanism.⁶¹ Whilst mitochondria are a source of ROS in the aged myocardium, they might also be susceptible to ROS-mediated dysfunction. Peroxidation of mitochondrial lipids can result in loss of cardiolipin,⁶² a phospholipid important for mitochondrial dynamics.⁶³ This impairs the activity of cytochrome c oxidase,⁶² whilst molecules that target cardiolipin enhance ETC respiratory function.⁶⁴ mtROS production might also increase the opening of the mitochondrial permeability transition pore (mtPTP) during ischaemia,⁵³ leading to mitochondrial swelling, functional impairment, and potentially apoptosis.

Increased mtPTP opening in response to Ca^{2+} overload can be mitigated by sirtuin-3 (SIRT3)-mediated deacetylation, preventing ageing-related cardiac hypertrophy,⁶⁵ suggesting that some ageing-related changes are reversible. A summary of myocardial energetics and metabolism changes is provided in *Figures 1 and 2*.

Loss of protection, inflammation, and cardiac ageing

Telomeres and cardiac ageing

Telomeres are repeated DNA sequences at the end of the chromosome, tasked with protecting DNA from degradation. Both premature and chronological ageing are associated with telomere shortening, which contributes to reduced regenerative capacity and heightened inflammation. Telomere shortening is associated with CVD pathology, making the independent contribution of ageing difficult to decouple. Importantly, interaction between different ageing-related changes can accelerate myocardial dysfunction; e.g. telomeric DNA sequences are more susceptible to DNA damage from oxidative stress, and thus ROS production augments telomere attrition.^{66,67} In parallel, telomeric shortening induces mitochondrial reprogramming, resulting in functional alterations,⁶⁶ processes exacerbated by oxidative stress.⁶⁸

Reduction in autophagy

The minimal replicative capacity of cardiomyocytes, coupled with a metabolically demanding redox-enriched intracellular environment, makes cardiomyocytes vulnerable to sustained damage. Autophagy, a cardioprotective mechanism, targets damaged proteins and organelles for lysosomal-mediated degradation, which is critical for maintaining myocardial baseline homeostasis and stress responses.⁶⁹ There is good evidence linking myocardial ageing with impaired autophagy and cardiomyocyte senescence,⁶⁹ including in progeria, a premature ageing syndrome.⁷⁰ In highly aerobic tissues like the heart, with its extensive mitochondrial networks, mitophagy (a specialized autophagy) preserves mitochondrial function and 'fitness',⁷¹ although the exact role of mitophagy in ageing-related changes remains to be defined.

The role of immune cells

The immune system impacts cardiac development, composition, and function.⁷² After myocardial injury, immune cells are recruited to remove dying heart tissue, scavenge pathogens, and promote healing.⁷² However, immune cells can also cause irreversible damage, contributing to heart failure.⁷² Ageing alters the relative proportions of cardiac leukocytes with increased monocyte-derived cardiac macrophages and CD8+ T cells.⁷³ Increased cardiac fibrosis with age can also be due to the modulation of macrophage fibrotic function (discussed below).

Haematopoietic stem cells can acquire mutations causing expansion of mutated clones (clonal haematopoiesis of indeterminate potential, CHIP), inducing inflammatory changes.⁷⁴ Such mutations predominantly occur in epigenetic regulators (such as DNMT3a or TET2) and adversely affect heart function.⁷⁴ Clonal haematopoiesis is enriched in patients with heart failure, increases with age, and predicts mortality.^{75,76} This is partly due to the CHIP-induced inflammaging processes including up-

regulation of pro-inflammatory genes, increasing cardiac leukocyte infiltration,⁷⁵ which contribute to cardiac dysfunction.^{75,76}

Mice with heart failure and CHIP display enlarged hearts with increased fibrosis and reduced output.⁷⁵ In addition, CHIP contributes to atherosclerosis⁷⁵ and a thrombotic environment,⁷⁷ thus increasing coronary artery disease risk.⁷⁶ Ageing increases senescent cardiac progenitor cells producing pro-inflammatory SASP factors, which promote cellular senescence via paracrine mechanisms, reducing cardiac repair capability.⁷⁸ SASPs also recruit monocytes to the heart, which promotes further senescence in a vicious feed-forward loop.⁷⁹ Whilst the immune system changes with age, further studies are required to directly link inflammaging to cardiac ageing.

Extracellular vesicles and myocardial senescence

Exosomes are small EVs exocytosed from cells, surrounded by a lipid bilayer that facilitates intercellular communication by transferring bioactive molecules including RNA, DNA, and proteins. EVs are both mediators and markers of cellular dysfunction.⁸⁰ In mice and humans, senescent cardiomyocytes and endothelial cells release EVs enriched with pro-inflammatory and profibrotic factors, which contribute to myocardial remodelling, extracellular matrix (ECM) degradation, and propagation of senescence to neighbouring cells, exacerbating tissue dysfunction.⁸¹

EV-associated plasma miRNAs are linked to senescence and myocardial dysfunction and may be useful diagnostic markers. For example, patients with heart failure and preserved ejection fraction have EVs with specific miRNA signatures.⁸² EVs from senescent endothelial cells, isolated from older human coronary arteries, carry calcification-promoting proteins and inflammatory mRNAs, contributing to vascular stiffness and ageing.^{80,83}


EVs also hold promise as therapeutic agents given that progenitor cell-derived EVs demonstrate anti-inflammatory and anti-fibrotic effects *in vitro* and *ex vivo* on senescent human myocardial tissue.⁸⁴ Additionally, EV-based therapies to deliver anti-senescent molecules, such as siRNAs targeting SASP components, are under investigation. These strategies aim to mitigate the spread of senescence and restore cardiac function in ageing myocardium.

Cells within the myocardium and vasculature are intimately linked; in human and animal tissues, the release of EV by activated endothelial cells can affect myocardial function.⁸⁵ This opens a new avenue for managing cardiac dysfunction, an area under active investigation. *Figure 3* summarizes the potential role of EVs in cardiac ageing.

Additional factors contributing to cardiac ageing

Valvular and vascular abnormalities

Ageing is associated with valvular abnormalities, including stenosis and regurgitation.⁸⁶ For example, aortic stenosis affects 2%–7% of individuals >65 years of age⁸⁷ and predisposes to heart failure. Vascular abnormalities also increase with age due to reduced elastin, increased collagen and calcification,



thus narrowing the vessels and increasing stiffness,⁸⁸ consequently leading to hypertension and compromising cardiac function.⁸⁹

Cardiac perfusion

Vascular changes with age reduce coronary blood flow,⁹⁰ increasing the risk of myocardial ischaemia.⁹¹ Moreover, microvascular rarefaction contributes to reduced coronary flow through reduced angiogenesis, and increased apoptosis resulting in diminished nitric oxide bioavailability and increased inflammation.⁹²⁻⁹⁵ Additionally, age-related decreases in vascular and platelet-derived growth factors reduce vascular repair and maturation of endothelial cells.^{92,93,95} Senescent endothelial cells and pericytes contribute to apoptosis whilst inhibiting angiogenesis.^{92,94} In addition, pericyte coverage is reduced, and the crosstalk with neighbouring cells is weakened in aged hearts, resulting in depressed pro-angiogenic activity, profibrotic commitment, and loss of ribosome biogenesis.⁹⁶ These defects contribute to vascular fragility, loss of microvascular barrier integrity, and increased severity of ischaemic injury. Several pathways central to vessel formation, such as hypoxia-inducible factor (HIF)-1 α , peroxisome proliferator-activated receptor- γ coactivator (PGC)-1 α , and endothelial nitric oxide synthase (eNOS), interact with ageing-related mechanisms.⁹⁷ Nucleolar proteins play critical roles in lifespan extension and DNA repair, regulate cellular plasticity in cardiomyocytes, and protect against ischaemia-induced damage, explaining why rescuing nuclear proteins reactivates angiogenesis in the ageing heart.⁹⁶

Cardiac fibrosis

In preclinical models of ageing, soluble collagen rises, whilst insoluble collagen falls,⁹⁸ increasing net intracardiac collagen.⁹⁹ In humans, endomysial and perimysial collagen types I and III increase in aged hearts.¹⁰⁰ Conversely, tissue inhibitors of metalloproteinases (TIMPs) are reduced,⁹⁹ whilst matrix metalloproteinases (MMPs), collagen type I and III, and Lysyl oxidase are all up-regulated in cardiac remodelling.^{101,102} SMAD2/3/4 [proteins involved in cardiac remodelling via extracellular matrix (ECM) gene expression] are altered with age.¹⁰² Importantly, miRNA profiling shows miR-1468-3p is elevated in aged cardiac fibrosis patients,¹⁰³ which can result in arrhythmias,^{104,105} whilst also predisposing to heart failure.¹⁰⁶ Late gadolinium-enhanced cardiac magnetic resonance imaging can be used to assess myocardial scarring and is a predictor of mortality, particularly in individuals with non-ischaemic dilated cardiomyopathy.¹⁰⁷ Cardiac fibroblast proliferation is dependent on extracellular matrix stiffness, which modulates NF- κ B protein levels, altering the mechanosensitive transcription factors pathways,¹⁰⁸ whilst calcium handling plays an important role in fibroblast proliferation and myocardial fibrosis.¹⁰⁹ Finally, β -adrenergic receptors regulate fibroblast proliferation, suggesting a role for the autonomic nervous system in myocardial fibrosis.¹¹⁰

The conducting system and autonomic innervation

Cardiac innervation changes with age, with maximal heart rate the most significantly affected,¹¹¹ in contrast to mean heart rate which shows minor changes,¹¹² and resting pulse rate

that remains consistent.¹¹³ Changes in maximal and mean heart rate were thought to be due to fibrosis and subsequent atrophy of the sinoatrial node (SAN).¹¹⁴ Lipids begin infiltrating the SAN around age 30, collagen and elastic tissue equal muscle tissue by age 40, and the node is delineated by fat and elastic fibres at age 50.¹¹⁵ By age 75, the remodelling of the node is so pronounced that the muscular distribution in the sinoatrial node shows almost no overlap with younger hearts.¹¹⁶

Similar changes are found throughout the conducting system, atrioventricular (AV) node, the bundle of His, and the beginning of the right and left branches,¹¹⁷ without any ageing-related calcification or inflammation.¹¹⁵ Diseases of conduction, such as AV block, are not usually associated with fibrosis of the conduction pathway unless myocardial ischaemia impacts the nodes. Normal sinus rhythm occurs even with most fibrotic sinoatrial nodes¹¹⁸ and heart block is only seen in the most severe fibrotic lesions of the AV node.¹¹⁸ It is now evident that fibrosis provides structural integrity by forming a scaffold, with fatty depositions and blood vessels, providing electrical insulation.¹¹⁹ Therefore, fibrosis of the conducting system is not, as previously thought, a disease of ageing but instead is essential for normative ageing.

Autonomic innervation contributes to inotropic, dromotropic, and chronotropic properties of the heart. Parasympathetic (acetylcholinergic) and sympathetic (noradrenergic) neurons decrease with age,¹²⁰ accompanied by loss of dopamine- β -hydroxylase neurons, which support sympathetic activity.¹²⁰ Whilst sympathetic nerves are decreased, sympathetic nerve activity reciprocally increases with age.¹²¹ The increased neurotransmitter release is met by ageing-related β -adrenergic desensitization.^{122,123}

Coupling increased noradrenaline release with age-associated reductions in its transporter density and activity,¹²⁴ causing a higher rate of noradrenaline spillover from the heart to the plasma,¹²⁵ although whether this is global or occurs in specific areas of the heart remains controversial.¹²⁶⁻¹²⁸ Conversely, parasympathetic drive decreases with age,¹²⁹ reducing heart rate variability.¹³⁰ In animal models, ageing-associated cardiac denervation can be diminished by senolytic therapy, suggesting this is a reversible process.¹³¹

The aforementioned loss of parasympathetic tone, accompanied by increased sympathetic tone, contributes to age-related hypertension.¹³²⁻¹³⁴ In conjunction, fibrosis of heart chamber walls and hypertrophy increase stiffness,¹³⁵ predisposing to ventricular and atrial arrhythmias.^{136,137} The combination of the above processes predisposes to heart failure and cardiac arrhythmias, frequently encountered in the older population.^{134,138} The effects of non-myocardial cell changes on cardiac ageing are summarized in [Figure 4](#).

Biomarkers or cardiovascular ageing

N-terminal pro-B-type natriuretic peptide is released by the heart under excessive strain and is an established biomarker of myocardial health.¹³⁹ However, due to its large overall inter-individual variability with older age, it would not appear to be a useful marker of normative ageing.¹⁴⁰ As such, NT-proBNP appears to better signal the presence of cardiovascular disease, rather than changes seen in ageing *per se*.¹⁴¹ This is a common feature of biomarkers of cardiac ageing, such as troponin, which




Figure 4 Physical changes within the myocardium and associated structures. The conducting tissue undergoes lipid infiltration and fibroblast activation, leading to fibrosis and steatosis. The alterations in the conducting system are accompanied by autonomic dysregulation, with a decrease in sympathetic and parasympathetic innervation accompanied by sympathetic overdrive that is compensated for by adrenergic desensitization. In contrast, parasympathetic drive decreases with age, which reduces heart rate variability and may contribute to cellular senescence. Altered collagen and elastin in coronary arteries, coupled with calcification, alter function and induce microvascular rarefaction. Similar mechanisms in the heart valves result in stenosis. Cellular loss and senescence are accompanied by compensatory hypertrophy and remodelling of the myocardium, including increased fibrosis. Created in Biorender.




Figure 5 Prevention of ageing-related cardiac senescence. Regular moderate physical activity and a balanced diet, rich in antioxidants and low in calories, can mitigate oxidative stress and inflammation whilst improving mitochondrial capacity, myocardial efficiency, and muscle mass, thus promoting myocardial health. Pharmacological intervention strategies, such as rapamycin, metformin, and senolytics, as well as gene therapies [sirtuins, extracellular vesicles (EVs), and growth differentiation factor 11 (GDF11)], have demonstrated early effectiveness in reducing cell senescence and oxidative damage. These therapies are also associated with improved endothelial function, restoring myocardial tissue, and aiding the reversal of cardiac hypertrophy. The up and down arrows indicate an increase or decrease, respectively. Created in Biorender.

also increase with age but are mainly related to the association with myocardial pathology.¹⁴¹

A gene expression-based age monitoring clock (GamC) has been created for cardiovascular health, comprising three key genes (*ABLIM1*, *CCR7*, and *LEF1*) involved in muscle contraction, immune function, cardiac repair, and response to injury.¹⁴² GamC shows a positive correlation with chronological age, with variation decreasing as age increases, but fails to correlate with markers of autonomic function, casting doubts over its relevance as a reliable measure of biological ageing.¹⁴²

Prevention of ageing-related changes

Regular physical activity and a balanced diet mitigate ageing-related cardiac decline and support cardiovascular health by improving myocardial metabolism⁴² and reducing oxidative stress.¹⁴³ A diet rich in antioxidants and anti-inflammatory components may further support cardiac function by combating

oxidative damage and inflammation.¹⁴⁴ Calorie restriction (CR), defined as a sustained reduction in caloric intake without malnutrition, may attenuate age-associated structural and functional changes in the myocardium by enhancing mitochondrial function, reducing oxidative stress, and modulating metabolic pathways.^{145,146} However, hard evidence that such a dietary approach improves myocardial health or increases longevity is lacking.

Gene therapy offers innovative approaches to counteract myocardial ageing. For instance, overexpression of the gene encoding growth differentiation factor 11 reverses ageing-related cardiac hypertrophy and improves cardiac function in animal models.¹⁴⁷ Intravenous injection of small EVs from young mice mitigates cardiac ageing by reducing cellular senescence.¹⁴⁸ Furthermore, therapies that enhance the expression of sirtuins, a family of proteins involved in cellular regulation, are under exploration for their potential to mimic the effects of calorie restriction.¹⁴⁹

Myosin heavy chain (MHC) proteins in cardiomyocytes are differentially modulated during ageing.^{150,151} Although the adult heart primarily contains the β -MHC isoform, a significant amount of α -MHC, which is more efficient, is present in neonatal hearts and can be reactivated by antiaging gene therapy.^{96,152} Administration of a polymorphic gene variant associated with extreme longevity or its encoded protein improves capillary density and perfusion, reduces fibrosis, and reverses diastolic dysfunction associated with ageing.^{153,154} Potential strategies to prevent cardiac ageing are summarized in [Figure 5](#).

Effects of senolytics on cardiac disorders

Navitoclax

In aged mice, navitoclax ameliorates left ventricular mass loss by decreasing myocyte hypertrophy and senescence whilst reducing cardiac fibrosis, consequently improving cardiac function.¹⁵⁵⁻¹⁵⁷ Moreover, this agent rejuvenates the vasculature by reducing stiffness and improving endothelial dysfunction.^{157,158} Navitoclax reduces mortality by improving ejection fraction following myocardial infarction in aged mice but not in ageing alone.^{156,158,159} Whilst navitoclax reduces senescence markers in primates and is well tolerated,¹⁶⁰ clinical uses may be limited as it increases endothelial-to-mesenchymal transition in advanced atherosclerotic lesions and raises mortality in some animal studies.¹⁵⁵

Dasatinib and quercetin (D + Q)

Quercetin monotherapy is unable to improve cardiac function or longevity in aged mice but can reduce mortality following hypoxic cardiac damage.^{161,162} In rodents, this agent is useful as a preventative measure where cardiovascular risk is high, such as sepsis and diabetes.^{161,162} Quercetin is often used intermittently in conjunction with dasatinib; monotherapy with the latter agent is avoided due to adverse effects in heart failure patients.^{160,162} D + Q induces apoptosis in senescent cells, decreasing SASP, whilst promoting cardiomyocyte production from progenitor cells, enhancing cardiac regenerative capacity and reversing ageing.^{78,160,162} Concurrently, D + Q reduces fibrosis, recovering cardiac function and preserving electrical stability, consequently improving survival in aged rodents.^{155,157,160,162} In high cardiovascular risk animals, such as obese mice, D + Q reduces senescent cells, improves insulin sensitivity, and enhances cardiac function.¹⁵⁸ Even a single dose of D + Q improves cardiovascular function, boosts health, and increases rodent survival.^{162,163} Their positive effects in cell line models of human age-related cardiomyopathy, including organoid models, promise future clinical potential.^{158,162,164}

Other senolytics and senomorphics

Hesperidin and hesperetin display senomorphic effects diminishing SASP levels and enhancing antioxidant enzymes to protect aged rat hearts.¹⁶¹ In models of ischaemia-reperfusion injury, kaempferol, apigenin, baicalin, fisetin, naringenin, and luteolin all protected cardiomyocytes by ameliorating mitochondrial dysfunction and ROS production, attenuating myocardial apoptosis and fibrosis, thus restoring cardiac function.¹⁶¹ Spermidine induces autophagy in hearts and vessels, delaying cardiac and

vascular ageing and extending lifespan in mice, whilst also improving function in animal models with different cardiovascular pathologies.¹⁵⁷ In elderly people, spermidine may lower the risk of cardiovascular disease and mortality,¹⁵⁷ although definitive trials are needed. Canagliflozin and dapagliflozin counteract myocyte senescence in a model of diabetic cardiomyopathy,¹⁶⁰ which may underlie their beneficial effects in heart failure.

In aged mice, senolytic and senomorphic agents acting on endothelial cells (anthocyanins), vascular smooth muscle cells (rutin)¹⁶¹ or both (fisetin, naringenin, apigen) have been tested.^{155,157,160,161} Fisetin ameliorates atherosclerosis in murine models of the disease, as does rutin in diabetic mice.¹⁶¹ Luteolin decreases vascular inflammation in mice and in human cell lines¹⁶¹ but human *in vivo* studies are lacking.

mTOR inhibitors

Rapamycin and its analogues demonstrate cardiac benefits in aged rodents with and without cardiomyopathies.^{155,160} However, these benefits are not necessarily translatable to humans,¹⁶¹ and therefore results from animal studies should be interpreted with caution.¹⁶⁰ Metformin, a widely used agent clinically, suppresses SASP expression, vascular inflammation, and mitochondrial ROS generation in preclinical models of ageing¹⁵⁵ and ameliorates endothelial dysfunction, arterial stiffness, and left ventricular hypertrophy.¹⁵⁵ Whilst observational studies suggest a cardiovascular benefit for metformin,^{155,160,165} robust data from randomized controlled trials are lacking.

Cardiovascular inflammaging

Inflammaging is age-related persistent low-grade inflammation that adversely impacts ageing, morbidity, and mortality. Immune ageing impedes the resolution of inflammatory processes, perpetuating inflammation.¹⁶⁶ At the organ level, inflammaging is more complex, consisting of three interrelated processes: chronic inflammation, cellular senescence, and maladaptive tissue remodelling.¹⁶⁷

Chronic inflammation, including various pro-inflammatory mediators, drives cellular senescence, tissue damage, and irreversible remodelling.^{166,167} In the heart, macrophage-mediated signalling and altered ECM degradation promote fibrosis through up-regulation of MMPs, NF- κ B, and activation of the NLRP3 inflammasome.¹⁶⁷ Oxidative stress and subsequent mitochondrial dysfunction promote the release of mtDNA, stimulating damage-associated molecular pattern secretion, further activating the NLRP3 inflammasome.^{166,167} Inflammation drives oxidative stress and mitochondrial dysfunction,¹⁶⁶ promoting cellular senescence. Interestingly, deficiency in mitochondrial transcription factor A in T cells alone replicates inflammaging and accelerates senescence.¹³⁹ SASP cells modify the local tissue environment by secreting pro-inflammatory cytokines, impairing stem cells through activin A secretion, and exacerbating fibrosis through fibroblast activation.¹⁶⁷ Maladaptive tissue remodelling increases cellular senescence by suppressing p53, Rb1, and Meis2 and initiating inflammatory processes through matrikines, further driving fibrosis and reducing contractility. In the coronary vasculature, these processes induce atherosclerosis and in heart valves can result in fibrosis and calcification.^{166,167} Age-related decline in 5' AMP-activated protein kinase and sirtuins (which

ameliorate inflammatory gene expression, DNA damage, and mitochondrial dysfunction) further potentiates inflammation.¹⁶⁷

The myocardium, where pathology meets physiology

Whilst one would predict that any form of ageing-related senescence in an organ would be classified as an ageing-related pathology, in the heart, the process is more complicated and nuanced. The loss or cell arrest of cardiomyocytes and hypertrophy of valvular cells are classical examples of ageing-related pathologies that fulfil the three criteria defined by the ICCARP,⁷ as they (i) develop and progress with increasing chronological age; (ii) are associated with, and contribute to, functional decline and an increased susceptibility to functional decline; and (iii) are evidenced by studies in humans.

Whilst some ageing-related alterations may cause concerns, some of these changes are not necessarily detrimental. For example, cardiomyocyte hypertrophy may be regarded as problematic, but cardiomyocyte loss, or senescence, without reciprocal hypertrophy would lead to a significant reduction in heart volume and loss of structural integrity, compromising pump function. Therefore, hypertrophy can be a compensatory protective mechanism that does not comply with the second criterion above. Cardiac fibrosis and steatosis must also be considered carefully, as they are not always inherently detrimental. We recognize that myocardial fibrosis and lipid infiltration may contribute to a decline in cardiac function and risk of arrhythmias. However, in the absence of overt disease or injury, fibrosis and steatosis of the conduction pathway are required to ensure appropriate pacing of the adult heart by providing increased insulation and structural stability to this tissue.

The understanding that certain changes within the myocardium, which have always been seen as pathological (e.g. hypertrophy and fibrosis), may constitute, under specific circumstances, protective mechanisms or be required for normal function in the heart has only just begun to emerge. This new discovery has significant implications for future myocardial disease treatment, where apparently similar processes are both pathological and pseudophysiological.

Conclusion: clinical implications of cardiac ageing

Disentangling physiological myocardial ageing processes from pathology, encountered in older populations, is key to understanding the impact of age on heart health. It can be argued that changes associated with cardiac ageing, which negatively impact function, do not practically differ from pathological dysfunction as both impact a patient's capacity to perform daily tasks. Furthermore, cardiac ageing may compromise the ability of the organ to adapt to pathological conditions, worsening symptom severity, and adding another dimension to the adverse role of cardiac ageing.

To simplify a rather complicated area, one option is to classify myocardial changes with ageing into those that are adaptive and have neutral/protective effects on cardiac function and those that negatively impact the function of the myocardium,

regardless of whether there is an associated pathological condition. Examples of adaptive changes include compensatory hypertrophy of myocytes, which can be regarded as protective as it helps to maintain cardiac volume in response to myocyte loss and gradual atrophy. However, pathological conditions commonly occurring in the elderly, such as hypertension, can cause myocyte hypertrophy, which is usually associated with cardiac enlargement. Another example is the age-related fibrotic changes around the conducting system, which preserve function and prevent arrhythmias. In contrast, fibrosis in the cardiac chambers can result in impaired myocardial relaxation/contraction and can predispose to arrhythmias. Therefore, it is not only the type of change but also the site and context that decide whether age-related alterations result in unwanted functional symptoms. These compensatory changes represent potentially important clinical challenges: e.g. could treating the valvular hypertrophy that underlies stenosis worsen cardiac function if it also removes the myocardial hypertrophy that acts to lessen the effect of myocardial cell loss? Likewise, treating myocardial fibrosis to prevent arrhythmias from re-entry loops could lead to even more inappropriate pacing of the heart if it disrupts the essential fibrosis of the conducting system.

A better understanding of myocardial changes with ageing has the potential to help with the clinical management of older people at a number of levels. First, elucidating the exact relationships between age-related changes and cardiac function will ensure appropriate management strategies for older people by avoiding the start of therapies when structural changes are detected that are protective or have no functional effects. Second, understanding the mechanisms for cardiac ageing, particularly those that cause functional myocardial decline, can help devise early and simple preventative strategies, thus improving longer-term outcomes. Third, elucidating the additive, or even synergistic, effects of myocardial ageing on an existing cardiac pathology will ensure that treatment modalities are both personalized and optimized. Finally, developing a new generation of therapies that specifically target cardiac ageing, such as senolytic therapies, will help to normalize myocardial function with ageing, thus improving the general well-being of older people and increasing productivity of the society.

Supplementary data

Supplementary data are not available at [European Heart Journal](#) online.

Declarations

Disclosure of Interest

Nothing to declare.

Although they have nothing to declare with regard to this review, for transparency, A.A.S.T. declares submission of two patent applications (PCT/EP2019/066546 and PG450503GB/YR), one of which is currently being explored with Life Molecular Imaging. A.A.S.T. has an active collaboration with Unilever. A.J.M. declares a paid consultancy role with Rivos Pharmaceuticals. J.B. received honoraria for lectures/consulting from Novartis, Abbott, Bayer, Pfizer, Boehringer Ingelheim, AstraZeneca, Cardior, CVRx, BMS, Amgen, Edwards, Roche,

and Zoll and research support for the department from Zoll, CVRx, Abiomed, Norgine, and Roche. All other authors declare no conflict of interest related to this work.

Data Availability

No data were generated or analysed for or in support of this paper.

Funding

The project was supported by a Longevity Impetus Grant from Norn Group, awarded to Drs Bentley and Calimport. A.A.S.T is supported by the Wellcome Trust (221295/Z/20/Z); Chan Zuckerberg Initiative (DAF2021-225273); and an advised fund of the Silicon Valley Community Foundation (<https://doi.org/10.13039/100014989>). A.J.M. is supported by Research Councils UK (Grant Number: EP/E500552/1), Wellcome Trust (Grant Number: 220033/Z/19/Z), British Heart Foundation (Grant Numbers: FS/17/61/33473, FS/4yPhD/F/20/34124, and FS/19/54/34889C). L.H. is supported by the Wellcome Trust (Grant Number: 220033/Z/19/Z). C.H.G. is supported by the British Heart Foundation (RG/15/6/31436) and the Health and Care Research Wales Infrastructure Award. R.T.R.H. is supported by the BBSRC (Grant Number: BB/X008290/1). R.A.A. is supported by grants from the BBSRC, British Heart Foundation Accelerator Award (RE/23/130040), Diabetes UK, National Institute for Health and Care Research (NIHR), and Abbott Diabetes Care. C.M. is supported by the British Heart Foundation (Grant Number: FS/SBSRF/22/31031) and by the HORIZON programme UNION-HORIZON-MSCA-DN-2024-111167421. N.A. is supported by the British Heart Foundation (BHF) Centre of Research Excellence, Oxford (RE/13/1/30181 and RE/18/3/34214), and a British Heart Foundation Intermediate Fellowship (FS/IBSRF/22/25110). M.A.S. is supported by the University of California Office of the President through the Tobacco-Related Disease Research Program Pilot Award (T31IP1790) and Impact Award (T31IR1585). C.E. is supported by the British Heart Foundation (BHF) Imperial Centre of Research Excellence Award (RE/24/130023), BHF Chair (CH/15/1/31199), and the National Institute for Health and Care Research Imperial Biomedical Research Centre (BRC).

References

- Litviňuková M, Talavera-López C, Maatz H, Reichart D, Worth Catherine L, Lindberg EL, et al. Cells of the adult human heart. *Nature* 2020;**588**: 466–72. <https://doi.org/10.1038/s41586-020-2797-4>
- Neubauer S. The failing heart—an engine out of fuel. *N Engl J Med* 2007;**356**: 1140–51. <https://doi.org/10.1056/NEJMr063052>
- Ribeiro ASF, Zerolo BE, López-Espuela F, Sánchez R, Fernandes VS. Cardiac system during the aging process. *Aging Dis* 2023;**14**:1105–22. <https://doi.org/10.14336/ad.2023.0115>
- van Riet EE, Hoes AW, Wagenaar KP, Limburg A, Landman MA, Rutten FH. Epidemiology of heart failure: the prevalence of heart failure and ventricular dysfunction in older adults over time. A systematic review. *Eur J Heart Fail* 2016;**18**:242–52. <https://doi.org/10.1002/ejhf.483>
- Rodgers JL, Jones J, Bolleddu SI, Vanthenapalli S, Rodgers LE, Shah K, et al. Cardiovascular risks associated with gender and aging. *J Cardiovasc Dev Dis* 2019;**6**:19. <https://doi.org/10.3390/jcdd6020019>
- Calimport SRG, Bentley BL, Stewart CE, Pawelec G, Scuteri A, Vinciguerra M, et al. To help aging populations, classify organismal senescence. *Science* 2019;**366**:576–8. <https://doi.org/10.1126/science.aay7319>
- Short E, Adcock IM, Al-Sarireh B, Ager A, Ajjan R, Akbar N, et al. Defining an ageing-related pathology, disease or syndrome: international consensus statement. *GeroScience* 2024;**47**:1713–20. <https://doi.org/10.1007/s11357-024-01315-9>
- Short E, Huckstepp Robert TR, Alavian K, Amoaku WMK, Barber TM, van Beek EJ, et al. International Consortium to Classify Ageing-related Pathologies (ICCARP) senescence definitions: achieving international consensus. *GeroScience* 2025;**47**:6789–96. <https://doi.org/10.1007/s11357-025-01509-9>
- Hastings MH, et al. Cardiac ageing: from hallmarks to therapeutic opportunities. *Cardiovasc Res* 2024;**121**:1474–88. <https://doi.org/10.1093/cvr/cvae124>
- Mallat Z, Fornes P, Costagliola R, Esposito B, Belmin J, Lecomte D, et al. Age and gender effects on cardiomyocyte apoptosis in the normal human heart. *J Gerontol A Biol Sci Med Sci* 2001;**56**:M719–23. <https://doi.org/10.1093/gerona/56.11.m719>
- Olivetti G, Melissari M, Capasso JM, Anversa P. Cardiomyopathy of the aging human heart. Myocyte loss and reactive cellular hypertrophy. *Circ Res* 1991;**68**:1560–8. <https://doi.org/10.1161/01.res.68.6.1560>
- Bergmann O, Zdunek S, Felker A, Salehpour M, Alkass K, Bernard S, et al. Dynamics of cell generation and turnover in the human heart. *Cell* 2015;**161**:1566–75. <https://doi.org/10.1016/j.cell.2015.05.026>
- Senyo SE, Steinhauser ML, Pizzimenti CL, Yang VK, Cai L, Wang M, et al. Mammalian heart renewal by pre-existing cardiomyocytes. *Nature* 2013;**493**:433–6. <https://doi.org/10.1038/nature11682>
- Eschenhagen T, Bolli R, Braun T, Field LJ, Fleischmann BK, Frisén J, et al. Cardiomyocyte regeneration: a consensus statement. *Circulation* 2017;**136**: 680–6. <https://doi.org/10.1161/circulationaha.117.029343>
- Suda M, Paul KH, Minamino T, Miller JD, Lerman A, Ellison-Hughes GM, et al. Senescent cells: a therapeutic target in cardiovascular diseases. *Cells* 2023;**12**:1296. <https://doi.org/10.3390/cells12091296>
- Tchkonina T, Kirkland JL. Aging, cell senescence, and chronic disease: emerging therapeutic strategies. *JAMA* 2018;**320**:1319–20. <https://doi.org/10.1001/jama.2018.12440>
- Evangelou K, Vasileiou PVS, Papaspyropoulos A, Hazapis O, Petty R, Demaria M, et al. Cellular senescence and cardiovascular diseases: moving to the “heart” of the problem. *Physiol Rev* 2023;**103**:609–47. <https://doi.org/10.1152/physrev.00007.2022>
- Rosenberg B, Pfitzer P. Ploidy in the hearts of elderly patients. *Virchows Arch B Cell Pathol Incl Mol Pathol* 1983;**42**:19–24. <https://doi.org/10.1007/bf02890366>
- Olivetti G, Cigola E, Maestri R, Corradi D, Lagrasta C, Gambert SR, et al. Aging, cardiac hypertrophy and ischemic cardiomyopathy do not affect the proportion of mononucleated and multinucleated myocytes in the human heart. *J Mol Cell Cardiol* 1996;**28**:1463–77. <https://doi.org/10.1006/jmcc.1996.0137>
- Tripodiadis F, Xanthopoulos A, Butler J. Cardiovascular aging and heart failure: JACC review topic of the week. *J Am Coll Cardiol* 2019;**74**:804–13. <https://doi.org/10.1016/j.jacc.2019.06.053>
- Pitrez PR, Monteiro LM, Borgogno O, Nissan X, Mertens J, Ferreira L. Cellular reprogramming as a tool to model human aging in a dish. *Nat Commun* 2024;**15**:1816. <https://doi.org/10.1038/s41467-024-46004-5>
- Rando TA, Chang HY. Aging, rejuvenation, and epigenetic reprogramming: resetting the aging clock. *Cell* 2012;**148**:46–57. <https://doi.org/10.1016/j.cell.2012.01.003>
- Lind L, Ingelsson E, Sundström J, Siegbahn A, Lampa E. Methylation-based estimated biological age and cardiovascular disease. *Eur J Clin Invest* 2018;**48**: e12872. <https://doi.org/10.1111/eci.12872>
- Papait R, Cattaneo P, Kunderfranco P, Greco C, Carullo P, Guffanti A, et al. Genome-wide analysis of histone marks identifying an epigenetic signature of promoters and enhancers underlying cardiac hypertrophy. *Proc Natl Acad Sci U S A* 2013;**110**:20164–9. <https://doi.org/10.1073/pnas.1315155110>
- Bauersachs J, Solomon SD, Anker SD, Antorrena-Miranda I, Batkai S, Viereck J, et al. Efficacy and safety of CDR132L in patients with reduced left ventricular ejection fraction after myocardial infarction: rationale and design of the HF-REVERT trial. *Eur J Heart Fail* 2024;**26**:674–82. doi:<https://doi.org/10.1002/ejhf.3139>
- Thum T, Gross C, Fiedler J, Fischer T, Kissler S, Bussen M, et al. MicroRNA-21 contributes to myocardial disease by stimulating MAP kinase signalling in fibroblasts. *Nature* 2008;**456**:980–4. <https://doi.org/10.1038/nature07511>
- Jazbutyte V, Fiedler J, Kneitz S, Galuppo P, Just A, Holzmann A, et al. MicroRNA-22 increases senescence and activates cardiac fibroblasts in the aging heart. *Age (Dordr)* 2013;**35**:747–62. <https://doi.org/10.1007/s11357-012-9407-9>
- Nikolajevic J, Ariaee N, Liew A, Abbasnia S, Fazeli B, Sabovic M. The role of MicroRNAs in endothelial cell senescence. *Cells* 2022;**11**:1185. <https://doi.org/10.3390/cells11071185>

29. Rogg E-M, Abplanalp WT, Bischof C, John D, Schulz MH, Krishnan J, et al. Analysis of cell type-specific effects of MicroRNA-92a provides novel insights into target regulation and mechanism of action. *Circulation* 2018; **138**:2545–58. <https://doi.org/10.1161/CIRCULATIONAHA.118.034598>
30. Cheng HS, Sivachandran N, Lau A, Boudreau E, Zhao JL, Baltimore D, et al. MicroRNA-146 represses endothelial activation by inhibiting pro-inflammatory pathways. *EMBO Mol Med* 2013; **5**:1017–34. <https://doi.org/10.1002/emmm.201202318>
31. Ito T, Yagi S, Yamakuchi M. MicroRNA-34a regulation of endothelial senescence. *Biochem Biophys Res Commun* 2010; **398**:735–40. doi:<https://doi.org/10.1016/j.bbrc.2010.07.012>
32. Boon RA, Iekushi K, Lechner S, Seeger T, Fischer A, Heydt S, et al. MicroRNA-34a regulates cardiac ageing and function. *Nature* 2013; **495**:107–10. <https://doi.org/10.1038/nature11919>
33. Accardi G, Bono F, Cammarata G, Aiello A, Herrero MT, Alessandro R, et al. miR-126-3p and miR-21-5p as hallmarks of bio-positive ageing; correlation analysis and machine learning prediction in young to ultra-centenarian Sicilian population. *Cells* 2022; **11**:1505. <https://doi.org/10.3390/cells11091505>
34. Wu J, Jiang Y, Sun J, Sun X. Identification and validation of an aging-associated circRNA-miRNA-mRNA network in neovascular age-related macular degeneration. *Gerontology* 2023; **69**:1218–31. <https://doi.org/10.1159/000531287>
35. Greco S, Gaetano C, Martelli F. Long noncoding competing endogenous RNA networks in age-associated cardiovascular diseases. *Int J Mol Sci* 2019; **20**:3079. <https://doi.org/10.3390/ijms20123079>
36. Okada M, Mitsunami K, Inubushi T, Kinoshita M. Influence of aging or left ventricular hypertrophy on the human heart: contents of phosphorus metabolites measured by ³¹P MRS. *Magn Reson Med* 1998; **39**:772–82. <https://doi.org/10.1002/mrm.1910390515>
37. Schocke MF, Metzler B, Wolf C, Steinboeck P, Kremser C, Pachinger O, et al. Impact of aging on cardiac high-energy phosphate metabolism determined by phosphorus-31 2-dimensional chemical shift imaging (³¹P 2D CSI). *Magn Reson Imaging* 2003; **21**:553–9. [https://doi.org/10.1016/s0730-725x\(03\)00079-1](https://doi.org/10.1016/s0730-725x(03)00079-1)
38. Chesky JA, Rockstein M, Lopez T. Changes with age of myocardial creatine phosphokinase in the male Fischer rat. *Mech Ageing Dev* 1980; **12**:237–43. [https://doi.org/10.1016/0047-6374\(80\)90046-9](https://doi.org/10.1016/0047-6374(80)90046-9)
39. Stanley WC, Recchia FA, Lopaschuk GD. Myocardial substrate metabolism in the normal and failing heart. *Physiol Rev* 2005; **85**:1093–129. <https://doi.org/10.1152/physrev.00006.2004>
40. Kates AM, Herrero P, Dence C, Soto P, Srinivasan M, Delano DG, et al. Impact of aging on substrate metabolism by the human heart. *J Am Coll Cardiol* 2003; **41**:293–9. [https://doi.org/10.1016/s0735-1097\(02\)02714-6](https://doi.org/10.1016/s0735-1097(02)02714-6)
41. Hyyti OM, Ledee D, Ning X-H, Ge M, Portman MA. Aging impairs myocardial fatty acid and ketone oxidation and modifies cardiac functional and metabolic responses to insulin in mice. *Am J Physiol Heart Circ Physiol* 2010; **299**:H868–75. <https://doi.org/10.1152/ajpheart.00931.2009>
42. Iemitsu M, Miyauchi T, Maeda S, Tanabe T, Takanashi M, Irukayama-Tomobe Y, et al. Aging-induced decrease in the PPAR- α level in hearts is improved by exercise training. *Am J Physiol Heart Circ Physiol* 2002; **283**:H1750–60. <https://doi.org/10.1152/ajpheart.01051.2001>
43. Stewart JE, Crawford JM, Mullen WE, Jacques A, Stoner MW, Scott I, et al. Cardiomyocyte-specific deletion of GCN5L1 reduces lysine acetylation and attenuates diastolic dysfunction in aged mice by improving cardiac fatty acid oxidation. *Biochem J* 2024; **481**:423–36. <https://doi.org/10.1042/bcj20230421>
44. Drosatos K. Fatty old hearts: role of cardiac lipotoxicity in age-related cardiomyopathy. *Pathobiol Aging Age Relat Dis* 2016; **6**:32221. <https://doi.org/10.3402/pba.v6.32221>
45. van der Meer RW, Rijzewijk LJ, Diamant M, Hammer S, Schar M, Bax JJ, et al. The ageing male heart: myocardial triglyceride content as independent predictor of diastolic function. *Eur Heart J* 2008; **29**:1516–22. <https://doi.org/10.1093/eurheartj/ehn207>
46. Koonen DP, Febbraio M, Bonnet S, Nagendran J, Young ME, Michelakis ED, et al. CD36 expression contributes to age-induced cardiomyopathy in mice. *Circulation* 2007; **116**:2139–47. <https://doi.org/10.1161/circulationaha.107.712901>
47. Glatz JFC, Angin Y, Steinbusch LKM, Schwenk RW, Luiken JJFP. CD36 as a target to prevent cardiac lipotoxicity and insulin resistance. *Prostaglandins Leukot Essent Fatty Acids* 2013; **88**:71–7. <https://doi.org/10.1016/j.plefa.2012.04.009>
48. Zhang L, Ussher JR, Oka T, Cadete VJJ, Wagg C, Lopaschuk GD. Cardiac diacylglycerol accumulation in high fat-fed mice is associated with impaired insulin-stimulated glucose oxidation. *Cardiovasc Res* 2011; **89**:148–56. <https://doi.org/10.1093/cvr/cvq266>
49. Hall JL, Mazzeo RS, Podolin DA, Cartee GD, Stanley WC. Exercise training does not compensate for age-related decrease in myocardial GLUT-4 content. *J Appl Physiol* (1985) 1994; **76**:328–32. <https://doi.org/10.1152/jappl.1994.76.1.328>
50. Bhashyam S, Parikh P, Bolukoglu H, Shannon AH, Porter JH, Shen Y-T, et al. Aging is associated with myocardial insulin resistance and mitochondrial dysfunction. *Am J Physiol Heart Circ Physiol* 2007; **293**:H3063–71. <https://doi.org/10.1152/ajpheart.00163.2007>
51. Abel ED. Free fatty acid oxidation in insulin resistance and obesity. *Heart Metab* 2010; **48**:5–10.
52. Hansford RG, Castro F. Age-linked changes in the activity of enzymes of the tricarboxylate cycle and lipid oxidation, and of carnitine content, in muscles of the rat. *Mech Ageing Dev* 1982; **19**:191–200. [https://doi.org/10.1016/0047-6374\(82\)90010-0](https://doi.org/10.1016/0047-6374(82)90010-0)
53. Lesnefsky EJ, Chen Q, Hoppel CL. Mitochondrial metabolism in aging heart. *Circ Res* 2016; **118**:1593–611. <https://doi.org/10.1161/circresaha.116.307505>
54. Rizvi F, Preston CC, Emelyanova L, Yousufuddin M, Viqar M, Dakwar O, et al. Effects of aging on cardiac oxidative stress and transcriptional changes in pathways of reactive oxygen species generation and clearance. *J Am Heart Assoc* 2021; **10**:e019948. <https://doi.org/10.1161/jaha.120.019948>
55. Müller-Höcker J. Cytochrome-c-oxidase deficient cardiomyocytes in the human heart—an age-related phenomenon. A histochemical ultracytochemical study. *Am J Pathol* 1989; **134**:1167–73.
56. Lesnefsky EJ, Guduz TI, Moghaddas S, Migita CT, Ikeda-Saito M, Turkaly PJ, et al. Aging decreases electron transport complex III activity in heart inter-fibrillar mitochondria by alteration of the cytochrome c binding site. *J Mol Cell Cardiol* 2001; **33**:37–47. <https://doi.org/10.1006/jmcc.2000.1273>
57. Fannin SW, Lesnefsky EJ, Slabe TJ, Hassan MO, Hoppel CL. Aging selectively decreases oxidative capacity in rat heart inter-fibrillar mitochondria. *Arch Biochem Biophys* 1999; **372**:399–407. <https://doi.org/10.1006/abbi.1999.1508>
58. Miró Ó, Casademont J, Casals E, Perea M, Urbano-Márquez A, Rustin P, et al. Aging is associated with increased lipid peroxidation in human hearts, but not with mitochondrial respiratory chain enzyme defects. *Cardiovasc Res* 2000; **47**:624–31. [https://doi.org/10.1016/S0008-6363\(00\)00122-X](https://doi.org/10.1016/S0008-6363(00)00122-X)
59. Gómez LA, Monette JS, Chavez JD, Maier CS, Hagen TM. Supercomplexes of the mitochondrial electron transport chain decline in the aging rat heart. *Arch Biochem Biophys* 2009; **490**:30–5. <https://doi.org/10.1016/j.abi.2009.08.002>
60. Hayakawa M, Hattori K, Sugiyama S, Ozawa T. Age-associated oxygen damage and mutations in mitochondrial DNA in human hearts. *Biochem Biophys Res Commun* 1992; **189**:979–85. [https://doi.org/10.1016/0006-291x\(92\)92300-m](https://doi.org/10.1016/0006-291x(92)92300-m)
61. Dai DF, Rabinovitch PS. Cardiac aging in mice and humans: the role of mitochondrial oxidative stress. *Trends Cardiovasc Med* 2009; **19**:213–20. <https://doi.org/10.1016/j.tcm.2009.12.004>
62. Paradies G, Ruggiero FM, Petrosillo G, Quagliariello E. Peroxidative damage to cardiac mitochondria: cytochrome oxidase and cardiolipin alterations. *FEBS Lett* 1998; **424**:155–8. [https://doi.org/10.1016/s0014-5793\(98\)00161-6](https://doi.org/10.1016/s0014-5793(98)00161-6)
63. Paradies G, Paradies V, Ruggiero FM, Petrosillo G. Role of cardiolipin in mitochondrial function and dynamics in health and disease: molecular and pharmacological aspects. *Cells* 2019; **8**:728. <https://doi.org/10.3390/cells8070728>
64. Szeto HH, Birk AV. Serendipity and the discovery of novel compounds that restore mitochondrial plasticity. *Clin Pharmacol Ther* 2014; **96**:672–83. doi: <https://doi.org/10.1038/clpt.2014.174>
65. Hafner AV, Dai J, Gomes AP, Xiao C-Y, Palmeira CM, Rosenzweig A, et al. Regulation of the mPTP by SIRT3-mediated deacetylation of CypD at lysine 166 suppresses age-related cardiac hypertrophy. *Aging (Albany NY)* 2010; **2**:914–23. <https://doi.org/10.18632/aging.100252>
66. Zheng Q, Huang J, Wang G. Mitochondria, telomeres and telomerase subunits. *Front Cell Dev Biol* 2019; **7**:274. <https://doi.org/10.3389/fcell.2019.00274>
67. von Zglinicki T, Pilger R, Sitte N. Accumulation of single-strand breaks is the major cause of telomere shortening in human fibroblasts. *Free Radic Biol Med* 2000; **28**:64–74. [https://doi.org/10.1016/s0891-5849\(99\)00207-5](https://doi.org/10.1016/s0891-5849(99)00207-5)
68. Borghini A, Ndreu R, Canale P, Campolo J, Marinaro I, Mercuri A, et al. Telomere length, mitochondrial DNA, and micronucleus yield in response to oxidative stress in peripheral blood mononuclear cells. *Int J Mol Sci* 2024; **25**:1428. <https://doi.org/10.3390/ijms25031428>
69. Ren J, Zhang Y. Targeting autophagy in aging and aging-related cardiovascular diseases. *Trends Pharmacol Sci* 2018; **39**:1064–76. <https://doi.org/10.1016/j.tips.2018.10.005>
70. Aman Y, Schmauck-Medina T, Hansen M, Morimoto RI, Simon AK, Bjedov I, et al. Autophagy in healthy aging and disease. *Nat Aging* 2021; **1**:634–50. <https://doi.org/10.1038/s43587-021-00098-4>

71. Picca A, Faitg J, Auwerx J, Ferrucci L, D'Amico D. Mitophagy in human health, ageing and disease. *Nat Metab* 2023;5:2047–61. <https://doi.org/10.1038/s42255-023-00930-8>
72. Swirski FK, Nahrendorf M. Cardioimmunology: the immune system in cardiac homeostasis and disease. *Nat Rev Immunol* 2018;18:733–44. <https://doi.org/10.1038/s41577-018-0065-8>
73. Molawi K, Wolf Y, Kandalla PK, Favret J, Hagemeyer N, Frenzel K, et al. Progressive replacement of embryo-derived cardiac macrophages with age. *J Exp Med* 2014;211:2151–8. <https://doi.org/10.1084/jem.20140639>
74. Sano S, Oshima K, Wang Y, Katanasaka Y, Sano M, Walsh K. CRISPR-mediated gene editing to assess the roles of Tet2 and Dnmt3a in clonal hematopoiesis and cardiovascular disease. *Circ Res* 2018;123:335–41. <https://doi.org/10.1161/circresaha.118.313225>
75. Chavkin NW, Min KD, Walsh K. Importance of clonal hematopoiesis in heart failure. *Trends Cardiovasc Med* 2022;32:198–203. <https://doi.org/10.1016/j.tcm.2021.04.005>
76. Sano S, Thel MC, Walsh K. Clonal hematopoiesis: the nonhereditary genetics of age-associated cardiovascular disease. *Curr Opin Cardiol* 2023;38:201–6. <https://doi.org/10.1097/hco.0000000000001032>
77. Karakasis P, Theofilis P, Lefkou E, Antoniadis AP, Patoulias D, Korantzopoulos P, et al. Clonal hematopoiesis of indeterminate potential and atrial fibrillation: insights into pathophysiology and clinical implications. *Int J Mol Sci* 2025;26:2739. <https://doi.org/10.3390/ijms26062739>
78. Lewis-McDougall FC, Ruchaya PJ, Domenjo-Vila E, Shin Teoh T, Prata L, Cottle BJ, et al. Aged-senescent cells contribute to impaired heart regeneration. *Aging Cell* 2019;18:e12931. <https://doi.org/10.1111/acer.12931>
79. Martini H, Iacovoni JS, Maggiorani D, Dutaur M, Marsal DJ, Roncalli J, et al. Aging induces cardiac mesenchymal stromal cell senescence and promotes endothelial cell fate of the CD90+ subset. *Aging Cell* 2019;18:e13015. doi: <https://doi.org/10.1111/acer.13015>
80. Mas-Bargues C, Borrás C, Alique M. The contribution of extracellular vesicles from senescent endothelial and vascular smooth muscle cells to vascular calcification. *Front Cardiovasc Med* 2022;9:854726. <https://doi.org/10.3389/fcvm.2022.854726>
81. Caño-Carrillo S, Castillo-Casas JM, Franco D, Lozano-Velasco E. Unraveling the signaling dynamics of small extracellular vesicles in cardiac diseases. *Cells* 2024;13:265. <https://doi.org/10.3390/cells13030265>
82. Watson CJ, Gupta SK, O'Connell E, Thum S, Glezeva N, Fendrich J et al. MicroRNA signatures differentiate prenatally from reduced ejection fraction heart failure. *Eur J Heart Fail*. 2015;17:405–15. <https://doi.org/10.1002/ehfj.244>
83. Abbas M, Jesel L, Auger C, Amoura L, Messas N, Manin G, et al. Endothelial microparticles from acute coronary syndrome patients induce premature coronary artery endothelial cell aging and thrombogenicity: role of the ang II/AT1 receptor/NADPH oxidase-mediated activation of MAPKs and PI3-kinase pathways. *Circulation* 2017;135:280–96. <https://doi.org/10.1161/circulationaha.116.017513>
84. Grigorian Shamagian L, Rogers RG, Luther K, Angert D, Echavez A, Liu W, et al. Rejuvenating effects of young extracellular vesicles in aged rats and in cellular models of human senescence. *Sci Rep* 2023;13:12240. <https://doi.org/10.1038/s41598-023-39370-5>
85. Akbar N, Braithwaite AT, Corr EM, Koelwyn GJ, van Solingen C, Cochain C, et al. Rapid neutrophil mobilization by VCAM-1+ endothelial cell-derived extracellular vesicles. *Cardiovasc Res* 2023;119:236–51. <https://doi.org/10.1093/cvr/cvac012>
86. Ohukainen P, Ruskoaho H, Rysa J. Cellular mechanisms of valvular thickening in early and intermediate calcific aortic valve disease. *Curr Cardiol Rev* 2018;14:264–71. <https://doi.org/10.2174/1573403X14666180820151325>
87. Martinsson A, Li X, Andersson C, Nilsson J, Smith JG, Sundquist K. Temporal trends in the incidence and prognosis of aortic stenosis: a nationwide study of the Swedish population. *Circulation* 2015;131:988–94. <https://doi.org/10.1161/circulationaha.114.012906>
88. Kohn JC, Lampi MC, Reinhart-King CA. Age-related vascular stiffening: causes and consequences. *Front Genet* 2015;6:112. <https://doi.org/10.3389/fgene.2015.00112>
89. Huynh J, Nishimura N, Rana K, Peloquin JM, Califano JP, Montague CR, et al. Age-related intimal stiffening enhances endothelial permeability and leukocyte transmigration. *Sci Transl Med* 2011;3:112ra122. <https://doi.org/10.1126/scitranslmed.3002761>
90. Thijssen DH, Carter SE, Green DJ. Arterial structure and function in vascular ageing: are you as old as your arteries? *J Physiol* 2016;594:2275–84. <https://doi.org/10.1113/jp270597>
91. Kelshiker MA, Seligman H, Howard JP, Rahman H, Foley M, Nowbar AN, et al. Coronary flow reserve and cardiovascular outcomes: a systematic review and meta-analysis. *Eur Heart J* 2022;43:1582–93. <https://doi.org/10.1093/eurheartj/ehab775>
92. Ahmed B, Rahman AA, Lee S, Malhotra R. The implications of aging on vascular health. *Int J Mol Sci* 2024;25:11188. <https://doi.org/10.3390/ijms252011188>
93. Rivard A, Fabre J-E, Silver M, Chen D, Murohara T, Kearney M, et al. Age-dependent impairment of angiogenesis. *Circulation* 1999;99:111–20. <https://doi.org/10.1161/01.cir.99.1.111>
94. Ungvari Z, Tarantini S, Kiss T, Wren JD, Giles CB, Griffin CT, et al. Endothelial dysfunction and angiogenesis impairment in the ageing vasculature. *Nat Rev Cardiol* 2018;15:555–65. <https://doi.org/10.1038/s41569-018-0030-z>
95. Goligorsky MS. Microvascular rarefaction. *Organogenesis* 2010;6:1–10. <https://doi.org/10.4161/org.6.1.10427>
96. Cattaneo M, Beltrami AP, Thomas AC, Spinetti G, Alvino VV, Avolio E, et al. The longevity-associated BPIFB4 gene supports cardiac function and vascularization in ageing cardiomyopathy. *Cardiovasc Res* 2023;119:1583–95. <https://doi.org/10.1093/cvr/cvad008>
97. Lähteenvuori J, Rosenzweig A, Sinclair D, North B. Effects of aging on angiogenesis. *Circ Res* 2012;110:1252–64. <https://doi.org/10.1161/CIRCRESAHA.111.246116>
98. Lindsey ML, Goshorn D, Squires C, Escobar G, Hendrick J, Mingoia J, et al. Age-dependent changes in myocardial matrix metalloproteinase/tissue inhibitor of metalloproteinase profiles and fibroblast function. *Cardiovasc Res* 2005;66:410–9. <https://doi.org/10.1016/j.cardiores.2004.11.029>
99. Horn MA, Graham HK, Richards MA, Clarke JD, Greensmith DJ, Briston SJ, et al. Age-related divergent remodeling of the cardiac extracellular matrix in heart failure: collagen accumulation in the young and loss in the aged. *J Mol Cell Cardiol* 2012;53:82–90. <https://doi.org/10.1016/j.yjmcc.2012.03.011>
100. Gazoti Debessa CR, Mesiano Maifrino LB, Rodrigues de Souza R. Age related changes of the collagen network of the human heart. *Mech Ageing Dev* 2001;122:1049–58. [https://doi.org/10.1016/s0047-6374\(01\)00238-x](https://doi.org/10.1016/s0047-6374(01)00238-x)
101. Biernacka A, Frangogiannis NG. Aging and cardiac fibrosis. *Aging Dis* 2011;2:158–73.
102. Dworatzek E, Baczko I, Kararigas G. Effects of aging on cardiac extracellular matrix in men and women. *Proteomics Clin Appl* 2016;10:84–91. <https://doi.org/10.1002/prca.201500031>
103. Lin R, Rahtu-Korpela L, Magga J, Ulvila J, Swan J, Kemppi A, et al. miR-1468-3p promotes aging-related cardiac fibrosis. *Mol Ther Nucleic Acids* 2020;20:589–605. <https://doi.org/10.1016/j.omtn.2020.04.001>
104. Li CY, Zhang JR, Hu WN, Li SN. Atrial fibrosis underlying atrial fibrillation (review). *Int J Mol Med* 2021;47:9. <https://doi.org/10.3892/ijmm.2020.4842>
105. Morita N, Mandel WJ, Kobayashi Y, Karagueuzian HS. Cardiac fibrosis as a determinant of ventricular tachyarrhythmias. *J Arrhythm* 2014;30:389–94. <https://doi.org/10.1016/j.joa.2013.12.008>
106. Sweeney M, Corden B, Cook SA. Targeting cardiac fibrosis in heart failure with preserved ejection fraction: mirage or miracle? *EMBO Mol Med* 2020;12:e10865. <https://doi.org/10.15252/emmm.201910865>
107. Theerasuwipakorn N, Chokesuwattanaskul R, Phannajit J, Marsukjai A, Thapanasuta M, Klem I, et al. Impact of late gadolinium-enhanced cardiac MRI on arrhythmic and mortality outcomes in nonischemic dilated cardiomyopathy: updated systematic review and meta-analysis. *Sci Rep* 2023;13:13775. <https://doi.org/10.1038/s41598-023-41087-4>
108. Ebrahimighaei R, Tarassova N, Bond SC, McNeill MC, Hathway T, Vohra H, et al. Extracellular matrix stiffness controls cardiac fibroblast proliferation via the nuclear factor-Y (NF-Y) transcription factor. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 2024;1871:119640. doi: <https://doi.org/10.1016/j.bbamcr.2023.119640>
109. Feng J, Armillei MK, Yu AS, Liang BT, Runnels LW, Yue L. Ca(2+) signaling in cardiac fibroblasts and fibrosis-associated heart diseases. *J Cardiovasc Dev Dis* 2019;6:34. <https://doi.org/10.3390/jcdd6040034>
110. Tanner MA, Thomas TP, Maitz CA, Grisanti LA. β_2 -adrenergic receptors increase cardiac fibroblast proliferation through the G α s/ERK1/2-dependent secretion of interleukin-6. *Int J Mol Sci* 2020;21:8507. <https://doi.org/10.3390/ijms21228507>
111. Tanaka H, Monahan KD, Seals DR. Age-predicted maximal heart rate revisited. *J Am Coll Cardiol* 2001;37:153–6. [https://doi.org/10.1016/s0735-1097\(00\)01054-8](https://doi.org/10.1016/s0735-1097(00)01054-8)
112. Santos MA, Sousa AC, Reis FP, Santos TR, Lima SO, Barreto-Filho JA. Does the aging process significantly modify the mean heart rate? *Arq Bras Cardiol* 2013;101:388–98. <https://doi.org/10.5935/abc.20130188>
113. Ostchega Y, Porter KS, Hughes J, Dillon CF, Nwankwo T. Resting pulse rate reference data for children, adolescents, and adults: United States, 1999–2008. *Natl Health Stat Report* 2011:1–16.
114. Shiraiishi I, Takamatsu T, Minamikawa T, Onouchi Z, Fujita S. Quantitative histological analysis of the human sinoatrial node during growth and aging. *Circulation* 1992;85:2176–84. <https://doi.org/10.1161/01.cir.85.6.2176>

115. Comunoglu C, Comunoglu N, Eren B, Tanrıloer O, Türkmen N, Gündoğmuş UN, et al. Age-related histopathological changes in the cardiac conducting system in the Turkish population: an evaluation of 202 autopsy cases. *Folia Morphol (Warsz)* 2012;**71**:178–82.
116. Davies MJ, Pomerance A. Quantitative study of ageing changes in the human sinoatrial node and internodal tracts. *Br Heart J* 1972;**34**:150–2. <https://doi.org/10.1136/hrt.34.2.150>
117. Erickson EE, Lev M. Aging changes in the human atrioventricular node, bundle, and bundle branches. *J Gerontol* 1952;**7**:1–12. <https://doi.org/10.1093/geronj/7.1.1>
118. Thery C, Gosselin B, Lekieffre J, Warembourg H. Pathology of sinoatrial node. Correlations with electrocardiographic findings in 111 patients. *Am Heart J* 1977;**93**:735–40. [https://doi.org/10.1016/s0002-8703\(77\)80070-7](https://doi.org/10.1016/s0002-8703(77)80070-7)
119. Fedorov VV, Glukhov AV, Chang R, Kostecki G, Aferol H, Hucker WJ, et al. Optical mapping of the isolated coronary-perfused human sinus node. *J Am Coll Cardiol* 2010;**56**:1386–94. <https://doi.org/10.1016/j.jacc.2010.03.098>
120. Chow LT, Chow SS, Anderson RH, Gosling JA. Autonomic innervation of the human cardiac conduction system: changes from infancy to senility—an immunohistochemical and histochemical analysis. *Anat Rec* 2001;**264**:169–82. <https://doi.org/10.1002/ar.1158>
121. de Lucia C, Piedepalumbo M, Paolisso G, Koch WJ. Sympathetic nervous system in age-related cardiovascular dysfunction: pathophysiology and therapeutic perspective. *Int J Biochem Cell Biol* 2019;**108**:29–33. doi:<https://doi.org/10.1016/j.biocel.2019.01.004>
122. Fleg JL, Schulman S, O'Connor F, Becker LC, Gerstenblith G, Clulow JF, et al. Effects of acute beta-adrenergic receptor blockade on age-associated changes in cardiovascular performance during dynamic exercise. *Circulation* 1994;**90**:2333–41. <https://doi.org/10.1161/01.cir.90.5.2333>
123. Brodde OE, Zerkowski HR, Schranz D, Broede-Sitz A, Michel-Reher M, Schäfer-Beisenbusch E, et al. Age-dependent changes in the beta-adrenoceptor-G-protein(s)-adenylyl cyclase system in human right atrium. *J Cardiovasc Pharmacol* 1995;**26**:20–6. <https://doi.org/10.1097/00005344-199507000-00004>
124. Leineweber K, Wangemann T, Giessler C, Bruck H, Dhein S, Kostelka M, et al. Age-dependent changes of cardiac neuronal noradrenaline reuptake transporter (uptake1) in the human heart. *J Am Coll Cardiol* 2002;**40**:1459–65. [https://doi.org/10.1016/s0735-1097\(02\)02168-x](https://doi.org/10.1016/s0735-1097(02)02168-x)
125. Esler MD, Thompson JM, Kaye DM, Turner AG, Jennings GL, Cox HS, et al. Effects of aging on the responsiveness of the human cardiac sympathetic nerves to stressors. *Circulation* 1995;**91**:351–8. <https://doi.org/10.1161/01.cir.91.2.351>
126. Tsuchimochi S, et al. Age and gender differences in normal myocardial adrenergic neuronal function evaluated by iodine-123-MIBG imaging. *J Nucl Med* 1995;**36**:969–74.
127. Nakajima K, Okuda K, Matsuo S, Wakabayashi H, Kinuya S. Is (123)I-metaiodobenzylguanidine heart-to-mediastinum ratio dependent on age? From Japanese Society of Nuclear Medicine normal database. *Ann Nucl Med* 2018;**32**:175–81. <https://doi.org/10.1007/s12149-018-1231-6>
128. Sakata K, Iida K, Mochizuki N, Ito M, Nakaya Y. Physiological changes in human cardiac sympathetic innervation and activity assessed by (123)I-metaiodobenzylguanidine (MIGB) imaging. *Circ J* 2009;**73**:310–5. <https://doi.org/10.1253/circj.cj-08-0614>
129. Kuga K, Yamaguchi I, Sugishita Y. Age-related changes of sinus node function and autonomic regulation in subjects without sinus node disease—assessment by pharmacologic autonomic blockade. *Jpn Circ J* 1993;**57**:760–8. <https://doi.org/10.1253/cj.57.760>
130. Jandackova VK, Scholes S, Britton A, Steptoe A. Are changes in heart rate variability in middle-aged and older people normative or caused by pathological conditions? Findings from a large population-based longitudinal cohort study. *J Am Heart Assoc* 2016;**5**:e002365. <https://doi.org/10.1161/jaha.115.002365>
131. Wagner JUG, Tombor LS, Malacarne PF, Kettenhausen L-M, Panthel J, Kujundzic H, et al. Aging impairs the neurovascular interface in the heart. *Science* 2023;**381**:897–906. <https://doi.org/10.1126/science.ade4961>
132. Baker SE, Limberg JK, Dillon GA, Curry TB, Joyner MJ, Nicholson WT. Aging alters the relative contributions of the sympathetic and parasympathetic nervous system to blood pressure control in women. *Hypertension* 2018;**72**:1236–42. <https://doi.org/10.1161/hypertensionaha.118.11550>
133. Franklin SS. Ageing and hypertension: the assessment of blood pressure indices in predicting coronary heart disease. *J Hypertens Suppl* 1999;**17**:S29–36.
134. Wallace SM, Yasmin, McEniery CM, Mäki-Petäjä KM, Booth AD, Cockcroft JR, et al. Isolated systolic hypertension is characterized by increased aortic stiffness and endothelial dysfunction. *Hypertension* 2007;**50**:228–33. <https://doi.org/10.1161/hypertensionaha.107.089391>
135. Chaturvedi RR, et al. Passive stiffness of myocardium from congenital heart disease and implications for diastole. *Circulation* 2010;**121**:979–88. <https://doi.org/10.1161/circulationaha.109.850677>
136. Nguyen TP, Qu Z, Weiss JN. Cardiac fibrosis and arrhythmogenesis: the road to repair is paved with perils. *J Mol Cell Cardiol* 2014;**70**:83–91. <https://doi.org/10.1016/j.yjmcc.2013.10.018>
137. Dun W, Boyden PA. Aged atria: electrical remodeling conducive to atrial fibrillation. *J Interv Card Electrophysiol* 2009;**25**:9–18. <https://doi.org/10.1007/s10840-008-9358-3>
138. Feinberg WM, Blackshear JL, Laupacis A, Kronmal R, Hart RG. Prevalence, age distribution, and gender of patients with atrial fibrillation. Analysis and implications. *Arch Intern Med* 1995;**155**:469–73. <https://doi.org/10.1001/archinte.1995.00430050045005>
139. Collerton J, Kingston A, Yousaf F, Davies K, Kenny A, Neely D, et al. Utility of NT-proBNP as a rule-out test for left ventricular dysfunction in very old people with limiting dyspnoea: the Newcastle 85+ Study. *BMC Cardiovasc Disord* 2014;**14**:128. <https://doi.org/10.1186/1471-2261-14-128>
140. Fradley MG, Larson MG, Cheng S, McCabe E, Coglianese E, Shah RV, et al. Reference limits for N-terminal-pro-B-type natriuretic peptide in healthy individuals (from the Framingham Heart Study). *Am J Cardiol* 2011;**108**:1341–5. <https://doi.org/10.1016/j.amjcard.2011.06.057>
141. Forman DE, de Lemos JA, Shaw LJ, Reuben DB, Lyubarova R, Peterson ED, et al. Cardiovascular biomarkers and imaging in older adults: JACC council perspectives. *J Am Coll Cardiol* 2020;**76**:1577–94. <https://doi.org/10.1016/j.jacc.2020.07.055>
142. Hernández-Bellido N, Hernández-Vicente A, García-Mendivil L, Ramos-Marqués E, Hermando D, Cebollada A, et al. A simple blood biomarker based on gene expression describes cardiovascular health-related biological age. *GeroScience* 2025;**47**:6613–29. <https://doi.org/10.1007/s11357-025-01784-6>
143. Hao Y, Liu W. Metabolic changes in cardiac aging. *Rev Cardiovasc Med* 2023;**24**:82. <https://doi.org/10.31083/j.rcm.2403082>
144. Pulakat L, Chen HH. Pro-senescence and anti-senescence mechanisms of cardiovascular aging: cardiac MicroRNA regulation of longevity drug-induced autophagy. *Front Pharmacol* 2020;**11**:774. <https://doi.org/10.3389/fphar.2020.00774>
145. Most J, Tosti V, Redman LM, Fontana L. Calorie restriction in humans: an update. *Ageing Res Rev* 2017;**39**:36–45. <https://doi.org/10.1016/j.arr.2016.08.005>
146. Minor RK, Allard JS, Younts CM, Ward TM, de Cabo R. Dietary interventions to extend life span and health span based on calorie restriction. *J Gerontol A Biol Sci Med Sci* 2010;**65**:695–703. <https://doi.org/10.1093/gerona/gdq042>
147. Loffredo FS, Steinhauser ML, Jay SM, Gannon J, Pancoast JR, Yalamanchi P, et al. Growth differentiation factor 11 is a circulating factor that reverses age-related cardiac hypertrophy. *Cell* 2013;**153**:828–39. <https://doi.org/10.1016/j.cell.2013.04.015>
148. Chen X, Luo Y, Zhu Q, Zhang J, Huang H, Kan Y, et al. Small extracellular vesicles from young plasma reverse age-related functional declines by improving mitochondrial energy metabolism. *Nat Aging* 2024;**4**:814–38. <https://doi.org/10.1038/s43587-024-00612-4>
149. Wu QJ, Zhang TN, Chen HH, Yu XF, Lv JL, Liu YY, et al. The sirtuin family in health and disease. *Signal Transduct Target Ther* 2022;**7**:402. <https://doi.org/10.1038/s41392-022-01257-8>
150. Long X, Boluyt MO, O'Neill L, Zheng JS, Wu G, Nitta YK, et al. Myocardial retinoid X receptor, thyroid hormone receptor, and myosin heavy chain gene expression in the rat during adult aging. *J Gerontol A Biol Sci Med Sci* 1999;**54**:B23–7. <https://doi.org/10.1093/gerona/54.1.b23>
151. Mercadier JJ, de la Bastie D, Ménasché P, Cao ANV, Bouveret P, Lorente P, et al. Alpha-myosin heavy chain isoform and atrial size in patients with various types of mitral valve dysfunction: a quantitative study. *J Am Coll Cardiol* 1987;**9**:1024–30. [https://doi.org/10.1016/s0735-1097\(87\)80303-0](https://doi.org/10.1016/s0735-1097(87)80303-0)
152. Dang Z, Avolio E, Thomas AC, Faulkner A, Beltrami AP, Cervellini C, et al. Transfer of a human gene variant associated with exceptional longevity improves cardiac function in obese type 2 diabetic mice through induction of the SDF-1/CXCR4 signalling pathway. *Eur J Heart Fail* 2020;**22**:1568–81. <https://doi.org/10.1002/ehfj.1840>
153. Faulkner A, Dang Z, Avolio E, Thomas AC, Batstone T, Lloyd GR, et al. Multi-omics analysis of diabetic heart disease in the db/db model reveals potential targets for treatment by a longevity-associated gene. *Cells* 2020;**9**:1283. <https://doi.org/10.3390/cells9051283>
154. Alvino VV, Slater S, Qiu Y, Cattaneo M, Mohammed KAK, Gate S, et al. Healthy longevity-associated protein improves cardiac function in murine models of cardiomyopathy with preserved ejection fraction. *Cardiovasc Diabetol* 2024;**23**:397. <https://doi.org/10.1186/s12933-024-02487-6>
155. Sutanto H, Fetarayani D, Narendra MR, Nasution SA. The role of the senescence-associated secretory phenotype in cardiovascular disease

- among the elderly. *Eur J Intern Med* 2025;**141**:106488. <https://doi.org/10.1016/j.ejim.2025.106488>
156. Walaszczyk A, Dookun E, Redgrave R, Tual-Chalot S, Victorelli S, Spyridopoulos I, et al. Pharmacological clearance of senescent cells improves survival and recovery in aged mice following acute myocardial infarction. *Aging Cell* 2019;**18**:e12945. <https://doi.org/10.1111/acer.12945>
157. Abdellatif M, Schmid ST, Fuerlinger A, Kroemer G. Anti-ageing interventions for the treatment of cardiovascular disease. *Cardiovasc Res* 2025;**121**:1524–36. <https://doi.org/10.1093/cvr/cvae177>
158. Atlante S, Gottardi Zamperla M, Cis L, Farsetti A, Gaetano C. Senolytic therapies for cardiovascular aging: tackling fibrosis and metabolic dysfunction. *Eur J Intern Med* 2025;**140**:106413. <https://doi.org/10.1016/j.ejim.2025.07.009>
159. Dookun E, Walaszczyk A, Redgrave R, Palmowski P, Tual-Chalot S, Suwana A, et al. Clearance of senescent cells during cardiac ischemia-reperfusion injury improves recovery. *Aging Cell* 2020;**19**:e13249. <https://doi.org/10.1111/acer.13249>
160. Stojanović SD, Thum T, Bauersachs J. Anti-senescence therapies: a new concept to address cardiovascular disease. *Cardiovasc Res* 2025;**121**:730–47. <https://doi.org/10.1093/cvr/cvaf030>
161. Zheng H, Li T, Hu Z, Zheng Q, Wang J. The potential of flavonoids to mitigate cellular senescence in cardiovascular disease. *Biogerontology* 2024;**25**:985–1010. <https://doi.org/10.1007/s10522-024-10141-7>
162. Nieto M, Konigsberg M, Silva-Palacios A. Quercetin and dasatinib, two powerful senolytics in age-related cardiovascular disease. *Biogerontology* 2024;**25**:71–82. <https://doi.org/10.1007/s10522-023-10068-5>
163. Zhu Y, Tchkonja T, Pirtskhalava T, Gower AC, Ding H, Giorgadze N, et al. The Achilles' heel of senescent cells: from transcriptome to senolytic drugs. *Aging Cell* 2015;**14**:644–58. <https://doi.org/10.1111/acer.12344>
164. Scalise M, Cianflone E, Quercia C, Pagano L, Chieffalo A, Stincelli A, et al. Senolytics rejuvenate aging cardiomyopathy in human cardiac organoids. *Mech Ageing Dev* 2025;**223**:112007. <https://doi.org/10.1016/j.mad.2024.112007>
165. Scherthaner G, Brand K, Bailey CJ. Metformin and the heart: update on mechanisms of cardiovascular protection with special reference to comorbid type 2 diabetes and heart failure. *Metabolism* 2022;**130**:155160. <https://doi.org/10.1016/j.metabol.2022.155160>
166. Müller L, Di Benedetto S. Inflammaging, immunosenescence, and cardiovascular aging: insights into long COVID implications. *Front Cardiovasc Med* 2024;**11**:1384996. <https://doi.org/10.3389/fcvm.2024.1384996>
167. Młynarska E, Kowalik A, Krajewska A, Krupińska N, Marcinkowska W, Motor J, et al. Inflammaging and senescence-driven extracellular matrix remodeling in age-associated cardiovascular disease. *Biomolecules* 2025;**15**:1452. <https://doi.org/10.3390/biom15101452>