

Masked Alkyne Equivalents for the Synthesis of Mechanically Interlocked Polyynes

Przemyslaw Gawel, Steffen L. Woltering, Yaoyao Xiong, Kirsten E. Christensen, and Harry L. Anderson*

Dedicated in memory of François Diederich (1952–2020)

[*] Dr. P. Gawel,[‡] Dr. S. L. Woltering, Dr. Y. Xiong, Dr. K. E. Christensen, Prof. H. L. Anderson

Department of Chemistry, Oxford University

Chemistry Research Laboratory, Oxford, OX1 3TA (UK)

E-mail: harry.anderson@chem.ox.ac.uk

[†] Current address: Institute of Organic Chemistry, Polish Academy of Sciences, Kasprzaka 44/52, Warsaw (Poland)

Supporting information for this article is given via a link at the end of the document.

Abstract: Polyynes, polyrotaxanes, encapsulated cyclocarbon catenanes and other fascinating mechanically interlocked carbon-rich architectures should become accessible if masked alkyne equivalents (MAEs) can be developed that are large enough to prevent unthreading of a macrocycle, and that can be cleanly unmasked under mild conditions. Here we report the synthesis of a new bulky MAE based on *t*-butylbicyclo[4.3.1]decatriene. This MAE was used to synthesize a polyynes [2]rotaxane and a masked-polyynes [3]rotaxane by Cadiot-Chodkiewicz coupling. Glaser cyclo-oligomerization of the [2]rotaxane gave masked cyclocarbon catenanes. The unmasking behavior of the catenanes and rotaxanes was tested by photolysis at a range of UV wavelengths. Photochemical unmasking did not proceed cleanly enough to prepare extended encapsulated polyynes polyrotaxanes. We highlight the scope and challenges involved with this approach to interlocked carbon-rich architectures.

Introduction


The synthesis of polyynes has attracted attention ever since the invention of the Glaser and Cadiot-Chodkiewicz coupling reactions.^[1] Polyynes are models for the elusive carbon allotrope carbyne,^[2] which consists of infinite linear chains of *sp*-hybridized carbon atoms. The electronic and mechanical properties predicted for this material have inspired generations of experimentalists and theorists,^[3] but its synthesis is precluded by the tendency of long polyynes chains to undergo exothermic crosslinking reactions.^[4] Bulky end groups have been shown to stabilize carbon chains of up to 48 atoms.^[5,6] Other strategies for stabilizing carbon chains include rotaxane formation^[7–9] and helical wrapping.^[10] Stabilization has been achieved by encapsulation inside carbon nanotubes, but this strongly modifies the properties of the 1D carbon chain.^[11]

Polyrotaxane formation is an appealing approach to stabilizing polyynes because the macrocyclic components could cover the whole length of the carbon chain, shielding regions that are unaffected by bulky end groups; in principle, this strategy could stabilize polyynes of any length. Synthetic approaches to polyynes [2]rotaxanes are well developed,^[7,8]


particularly using active metal templates,^[12,13] but there is a need for methods for threading many macrocycles onto a long polyynes thread to form a polyrotaxane.

Masked alkyne equivalents (MAEs) have been used since the 1980s to release polyynes chains from suitable precursors. Unmasking is achieved using light,^[14–16] heat,^[17] chemical reagents^[18–21] and on-surface using voltage pulses.^[22,23] MAEs have opened synthetic avenues to cyclocarbons, rings consisting entirely of *sp*-hybridized carbon atoms.^[17–19,23] While cyclocarbons were successfully studied on surfaces using AFM,^[23] they are only stable at low temperatures. One strategy for making them stable under ambient conditions may be to encapsulate them as catenanes,^[24] by analogy with the stabilization of cyclobutadiene inside a hemicarcerand, by Cram and co-workers.^[25] The synthesis of cyclocarbon catenanes could be achieved using MAEs that act as stoppers in a rotaxane intermediate. Such a rotaxane would be a versatile building block for encapsulated cyclocarbons and carbyne polyrotaxanes, as a form of insulated molecular wire.^[26]

Most reported MAEs are too small to act as stoppers in rotaxanes. Here we explore a MAE based on a bicyclo[4.3.1]decatriene pioneered by Tobe,^[15] but with a *t*-butyl group on the indane ring to increase the steric bulk. When irradiated with UV light, bicyclo[4.3.1]decatrienes **1** generate a carbene, which rearranges rapidly to form an alkyne **2** (Scheme 1).^[15] Here we demonstrate that this masking group is large enough to prevent slipping of a small phenanthroline macrocycle **3**^[27] by synthesizing rotaxane **4** (Scheme 2). We demonstrate that this MAE can be used to synthesize masked polyynes polyrotaxanes and masked cyclocarbon catenanes, and we test photochemical unmasking of these compounds.



Scheme 1. Photochemical unmasking of bicyclo[4.3.1]decatriene **1** and subsequent rearrangement of the carbene to form TMS-triyne **2**.^[15]



Scheme 2. Proposed synthetic route to cyclocarbon catenanes and carbyne polyrotaxanes. (PG = protecting group.)

Results and Discussion


1. [2]Rotaxane Synthesis

Designing a bulky photocleavable MAE was the first challenge in this project. The most promising MAE was identified as Tobe's bicyclo[4.3.1]decatriene. In contrast to the other photocleavable MAE developed by Tobe's group ([4.3.2]propellatriene),^[14] it is not fused as an annulated ring but attached to a double bond that is cross-conjugated with the acetylene network; as a result, it protrudes further from the polyyne chain, increasing the steric bulk. However, model studies using polyynes masked with unsubstituted bicyclo[4.3.1]decatriene and macrocycle **3** did not yield rotaxanes, indicating that the macrocycle can slip over this MAE. This problem was solved by adding a *t*-butyl group.

Masked triyne **13** is synthesized as shown in Scheme 3. Friedel-Crafts alkylation of indane gives **5**.^[28] Subsequent Birch reduction yields cyclohexadiene **6**. Dibromocarbene attacks **6** predominantly on the central double bond to yield **7**, whereas in absence of the *t*-butyl group both double bonds react.^[29] Reaction conditions for addition of bromine to **7** to give tetrabromide **8** had to be optimized, because Tobe's procedure (using pyridinium tribromide)^[15] gave an overbrominated product

S1 (which was identified by single-crystal X-ray analysis; Supporting Information S7). Use of elemental bromine in carbon tetrachloride at 0 °C gave the desired tetrabromide **8**, which was immediately submitted for elimination using DBU to furnish **9** in 40% yield over two steps. Intermediate **9** can be prepared in large quantities and stored without special precautions, and its structure was confirmed by X-ray crystallography (see Supporting Information S7).


When **9** is treated with *t*-BuLi, one bromine is exchanged for lithium and the carbanion can be captured with ketone **10**^[30] to install alkyne units in the periphery of the MAE, to give **11**, after quenching with TMS-chloride. Another lithium-bromine exchange with *t*-BuLi eliminates the OTMS group to give masked triyne **12**. This elimination is accompanied by a skeletal rearrangement: the [4.3.1]propellane undergoes an electrocyclic reaction to form the bicyclo[4.3.1]decane, as revealed by the fact that the bridgehead carbon signal is shifted from 47 ppm in **11** to 140 ppm in **12**. This is in accord with previous reports^[15,31] and the rearrangement is probably driven by the strain in the cyclopropane subunit with an exocyclic double bond.



Scheme 3. Synthesis of bulky masked alkyne equivalent building blocks **16** and **17**.

Deprotection of the TMS-alkynes with aqueous LiOH yields masked triyne **13**, which is desymmetrized by statistically protecting one of the terminal acetylenes with acetone, to give **14**. The two alkynes in **13** are nonequivalent, due to the presence of the *t*-Bu substituent, and this reaction gives an inseparable mixture of the *E* and *Z* stereoisomers of **14** in a ratio of 2:3. It was not possible to assign the stereoisomers, as the distance between *t*-butyl group and terminal acetylene or acetone group is too far to observe NOEs by ^1H NMR spectroscopy and attempts at crystallizing these compounds were unsuccessful. Glaser coupling with TMS-acetylene yields the masked tetrayne **15** and separation of the *E* and *Z* stereoisomers was achieved at this stage, although it was still not possible to assign the isomers. The TMS group of **15** is removed by treatment with aqueous potassium carbonate to give the terminal alkyne **16**, which is converted to the corresponding bromide **17** with NBS in the presence of AgNO_3 . Both **16** and **17** are unstable and quickly decompose (even at low temperature) in pure form. Therefore, these compounds were prepared immediately before use.

With compounds **16** and **17** in hands, we screened conditions for [2]rotaxane formation. Homocoupling of alkyne **16** in the presence of macrocycle **3**, mediated by $\text{Cu(I)}^{[12]}$ gave traces of interlocked product that could be detected by mass spectrometry, but the rotaxane could not be isolated. In contrast, Cadiot-Chodkiewicz coupling between the terminal alkyne **16** and bromide **17** gives [2]rotaxane **18** in 22% yield (Scheme 4). Deprotection of the acetone groups with $\text{K}_2\text{CO}_3/18\text{-crown-6}^{[32]}$ gives [2]rotaxane **4** in 60% yield. This [2]rotaxane is the first example of an interlocked compound in which a masked alkyne acts as a stopper. The interlocked character is unambiguously confirmed by ESI-MS and ^1H NMR spectroscopy. Tightly fitting macrocycles can sometimes enhance the yield of rotaxanes, $^{[33]}$ but the synthesis of **18** is less efficient than similar reactions using larger macrocycles. $^{[7,8]}$



Scheme 4. Cadiot-Chodkiewicz coupling of **16** and **17** in the cavity of **3** yields [2]rotaxane **18**, which is deprotected with K_2CO_3 to give **4**.

The ^1H NMR spectrum of [2]rotaxane **4** is compared with the spectra of free macrocycle **3** and free thread **19** in Figure 1. As the axle is formed as a mixture of diastereomers, additional splittings for the macrocycle peaks are observed. The macrocycle also affects the peaks of the axle and increases the splitting between the *E* and *Z* isomers; for example, the signal for the proton of the terminal acetylene (d in Figure 1) appears as a singlet in free thread **19**, but it is split in [2]rotaxane **4**. The effect of interlocking is stronger for proton a (shielded by 0.2 ppm in the [2]rotaxane) of the cycloheptatriene ring compared to signals b and c, which are almost unaffected by the threaded macrocycle. [2]Rotaxanes such as **4** have potential as building blocks for numerous systems involving encapsulated alkynes. To demonstrate this, we show how interlocked systems of higher order can be derived from this subunit.





Figure 1. ^1H NMR spectra (CDCl_3 , 298 K, 400 MHz) of macrocycle **3**, rotaxane **4** and free thread **19**. The signals of the macrocycle are strongly split because the thread is a mixture of diastereomers.

2. [3]Rotaxane Synthesis

Synthesis of a [3]rotaxane was investigated as a step towards the preparation of carbyne polyrotaxanes. An asymmetric [2]rotaxane **21**, with the MAE stopper on one side and a bulky supertrityl stopper on the other side, was synthesized via a Cadiot-Chodkiewicz coupling between bromo alkyne **17** and supertrityl triyne **20**^[5] in the presence of macrocycle **3** (Scheme 5). In an attempt to keep the NMR spectra simple, pure samples of diastereomer A of **16** (unknown *E* or *Z* stereochemistry) were used in this line of experiments. The bulky supertrityl stoppers were chosen to ensure that the interlocking of components is maintained after unmasking of the MAE. Removal of the acetonide protecting group with K_2CO_3 /18-crown-6 yields **22**. Oxidative Glaser homo-coupling of **22** gives [3]rotaxane **23** in 51% yield. The structure of **23** was unambiguously confirmed by NMR spectroscopy and mass spectrometry. At this stage, the [3]rotaxane is compartmentalized: the macrocycles are trapped between the supertrityl stopper unit and the MAE and cannot shuttle freely along the polyyne thread. These findings show that it is possible to synthesize masked encapsulated polyyne chains using this approach, which opens avenues to the synthesis of encapsulated carbyne (Scheme 1).




Scheme 5. Synthesis of [3]rotaxane **23**.

3. Synthesis of Masked Cyclocarbon Catenanes

Catenanes were prepared from [2]rotaxane **4** (consisting exclusively of diastereomer B) by Glaser cyclo-oligomerization under dilute conditions. A range of conditions was screened, and

only coupling with copper(I) chloride in pyridine under oxygen yielded catenanes (Scheme 6). [3]Catenane **24** and [4]catenane **25** were separated by recycling gel permeation chromatography and isolated in 5% and 4% yield, respectively. The constitution of **24** and **25** was demonstrated by MALDI mass spectrometry, but we were unable to record informative NMR spectra of these catenanes; they gave broad and poorly resolved spectra, which probably reflects the fact that they are complex mixtures of diastereomers. (Even using a single diastereomer of **4** is expected to give 7 stereoisomers of **24** and 24 stereoisomers for **25**).



Scheme 6. Synthesis of masked cyclocarbon catenanes **24** and **25** via copper-mediated cyclo-oligomerization of rotaxane **4**.

3. Photochemical Unmasking

Tobe reported photochemical unmasking of compound **1** to give triyne **2** in 43% yield (Scheme 1),^[15] so we decided to investigate whether the additional *t*-butyl group in the present system influences its photochemical reactivity. TMS-protected MAE **12** was chosen as a model to study the photochemistry of this system.

The ¹H NMR spectra of compound **12** before and after irradiation at 250 nm in CDCl₃ are compared in Figure 2. Under the conditions of our experiment, unmasking was complete after 4 h of irradiation. The unmasking reaction appears to proceed cleanly to the corresponding TMS-triyne **2** and *t*-butyl indane **5**, but decomposition via an unknown pathway takes place and the

signal-to-noise-ratio decreases over time. This contrasts with the photochemistry of unsubstituted **1**, in which an isomerized product is formed in 30% yield via a 1,5-sigmatropic shift.^[34]





Figure 2. ¹H NMR spectra (CDCl₃, 298 K, 400 MHz) of UV unmasking (250 nm) of **12** in CDCl₃.

Using adamantane as an internal ¹H NMR standard, the yield for **5** and **2** from irradiation of **12** was determined to be 37% and 47%, respectively (with an uncertainty of about 3%), which is consistent with the isolated yields (23% **5**, 36% **2**). The unmasking performance could not be improved by switching to longer wavelengths (300 nm, 350 nm) or by varying concentration, or changing the solvent to cyclohexane. Unfortunately, the photochemical performance of this MAE is not suitable for the unmasking of extended systems with multiple MAE units, because the yield is expected to drop exponentially with the number of masking units present in the molecule. In spite of this, we tested the unmasking of [3]rotaxane **23**, [3]catenane **24** and [4]catenane **25**. In the case of [3]rotaxane **23**, evidence for unmasking was observed on irradiation at 250 nm (Scheme 7). Mass spectrometric analysis of the reaction mixture showed strong peaks for the partially unmasked [3]rotaxane **26** and weak signals for fully unmasked [3]rotaxane **27**. UV-vis analysis of the reaction mixture confirmed that unmasking was achieved, and the characteristic peaks of a tetradecayne stoppered by supertrityl groups were observed at 428, 394, and 378 nm (see Supporting Information S6),^[5] but the unmasked rotaxanes **26** and **27** were not isolated.



Scheme 7. Photochemical unmasking of [3]rotaxane **23**. When irradiated with UV light (250 nm) partially unmasked rotaxane **26** can be identified in the reaction mixture by ESI-MS. The UV-vis of the reaction mixture shows signature peaks for a supertrityl-stoppered tetradecayne, suggesting the formation of **27**. Unmasking was not efficient enough to isolate **26** or **27**.

These findings suggest that unmasking of the catenanes **24** and **25** is unlikely to be viable using this MAE. Irradiation with UV light (254 nm or 350 nm) in either CDCl_3 (at 20 °C) or *n*-hexane (at –80 °C) was monitored by mass spectrometry, but no peaks for fully or partially unmasked cyclocarbon catenanes were observed. Based on the performance of the model compound, the theoretical yield for complete unmasking would be 4–6% for the [3]catenane and up to 1.5% for the [4]catenane.

Conclusion

We have demonstrated the first synthesis of a [2]rotaxane in which both stoppers are MAEs. Rotaxanes of this type are potential precursors of polyyne polyrotaxanes and cyclocarbon catenanes. The reactivity of this MAE (as tested in compounds **12**, **23**, **24** and **25**) is unsuitable for the preparation of interlocked polynes, and highlights the need for better understanding of the structure-property relationships controlling photochemical unmasking. Recently, we showed that a different MAE based on indane is capable of unmasking within rotaxanes but is too small to act as a stopper.^[8] When modified appropriately to increase the bulk, such a system might unmask more efficiently than the MAE presented here. An alternative to photochemically activated systems might be the use of metal complexes as stoppering groups as they are inherently sterically demanding.^[21]

Acknowledgements


We thank Daniel R. Kohn and Alexandru Vlasceanu for helpful discussions, and F. Wieland Goetzke and Maria A. Gafitescu for help with photoreactors. This project was funded by Leverhulme Trust project grant RPG-2017-032. P.G. was supported by Swiss National Science Foundation Postdoc.Mobility fellowship P300P2-177829.

Keywords: polynes • rotaxanes • masked alkyne • supramolecular encapsulation • photochemistry

- [1] (a) C. Glaser, *Ber. Dtsch. Chem. Ges.* **1869**, 2, 422–424; b) W. Chodkiewicz, P. Cadiot, *Hebdomadae Acad. Sci.* **1955**, 241, 1055–1057; c) P. Siemen, R. C. Livingston, F. Diederich, *Angew. Chem. Int. Ed.* **2000**, 39, 2632–2637; *Angew. Chem.* **2000**, 112, 2740–2767.
- [2] a) F. Diederich *Nature* **1994**, 369, 199–207; b) W. A. Chalifoux, R. R. Tykwinski, *C. R. Chim.* **2009**, 12, 341–358; c) R. R. Tykwinski, *Chem. Rec.* **2015**, 15, 1060–1074; d) C. S. Casari, M. Tommasini, R. R. Tykwinski, *Nanoscale* **2016**, 8, 4414–4435; e) C. S. Casari, A. Milani, *MRS Commun.* **2018**, 8, 207–219.
- [3] a) M. Kertesz, J. Koller, A. Azman, *J. Chem. Phys.* **1978**, 68, 2779–2782; b) S. Tongay, R. T. Senger, S. Dag, S. Ciraci, *Phys. Rev. Lett.* **2004**, 93, 136404; c) S. Yang, M. Miklos Kertesz, *J. Phys. Chem. A* **2006**, 110, 9771–9774; d) Z. Zanolli, G. Onida, J.-C. Charlier, *ACS Nano* **2010**, 4, 5174–5180; e) M. Liu, V. I. Artyukhov, H. Lee, F. Xu, B. I. Yakobson, *ACS Nano* **2013**, 7, 10075–10082.
- [4] a) P. Tarakeshwar, P. R. Buseck, H. W. Kroto, *J. Phys. Chem. Lett.* **2016**, 7, 1675–1681; b) T. R. Hoye, B. Baire, D. Niu, P. H. Willoughby, B. P. Woods, *Nature* **2012**, 490, 208–211; c) S. Schrettl, C. Stefaniu, C. Schwiager, G. Pasche, E. Oveisi, Y. Fontana, A. Fontcuberta i Morral, J. Reguera, R. Petraglia, C. Corminboeuf, G. Brezesinski, H. Frauenrath, *Nat. Chem.* **2014**, 6, 468–467; d) E. T. Chernick, R. R. Tykwinski, *J. Phys. Org. Chem.* **2013**, 26, 742–749.
- [5] a) Y. Gao, Y. Hou, F. Gordillo Gámez, M. J. Ferguson, J. Casado, R. R. Tykwinski, *Nat. Chem.* **2020**, 12, 1143–1149; b) W. Chalifoux, R. R. Tykwinski, *Nat. Chem.* **2010**, 2, 967–9714.
- [6] a) B. Pigulski, N. Gulia, S. Szafert, *Eur. J. Org. Chem.* **2019**, 1420–1445; b) M. Jevic, M. B. Nielsen, *Asian J. Org. Chem.* **2015**, 4, 286–295.
- [7] a) L. D. Movsisyan, M. Franz, F. Hampel, A. L. Thompson, R. R. Tykwinski, H. L. Anderson, *J. Am. Chem. Soc.* **2016**, 138, 1366–1376; b) L. D. Movsisyan, D. V. Kondratuk, M. Franz, A. L. Thompson, R. R. Tykwinski, H. L. Anderson, *Org. Lett.* **2012**, 14, 3424–3426; c) N. Weisbach, Z. Baranova, S. Gauthier, J. H. Reibenspies, J. A. Gladysz, *Chem. Commun.* **2012**, 48, 7562–7564.
- [8] S. L. Woltering, P. Gawel, K. E. Christensen, A. L. Thompson, H. L. Anderson, *J. Am. Chem. Soc.* **2020**, 142, 13523–13532.
- [9] S. Schrettl, E. Contal, T. Hoheisel, M. Fritzsche, S. Balog, R. Szilluweit, H. Frauenrath, *Chem. Sci.* **2015**, 6, 564–574.

- [10] J. Stahl, J. C. Bohling, E. B. Bauer, T. B. Peters, W. Mohr, J. M. Martín-Alvarez, F. Hampel, J. A. Gladysz, *Angew. Chem. Int. Ed.* **2002**, *41*, 1871–1876; *Angew. Chem.* **2002**, *114*, 1951–1957.
- [11] a) C. Zhao, R. Kitaura, H. Hara, S. Irle, H. Shinohara, *J. Phys. Chem. C* **2011**, *115*, 13166–13170; b) L. Shi, P. Rohringer, K. Suenaga, Y. Niimi, J. Kotakoski, J. C. Meyer, H. Peterlik, M. Wanko, S. Cahangirov, A. Rubio, Z. J. Lapin, L. Novotny, P. Ayala, T. Pichler, *Nat. Mater.* **2016**, *15*, 634–640.
- [12] S. Saito, E. Takahashi, K. Nakazono, *Org. Lett.* **2006**, *8*, 5133–5136.
- [13] J. Berná, J. D. Crowley, S. M. Goldup, K. D. Hänni, A.-L. Lee, D. A. Leigh, *Angew. Chem. Int. Ed.* **2007**, *46*, 5709–5713; *Angew. Chem.* **2007**, *119*, 5811–5815.
- [14] a) Y. Tobe, T. Fujii, K. Naemura, *J. Org. Chem.* **1994**, *59*, 1236–1237; b) Y. Tobe, T. Fujii, H. Matsumoto, K. Naemura, Y. Achiba, T. Wakabayashi, *J. Am. Chem. Soc.* **1996**, *118*, 2758–2759.
- [15] Y. Tobe, R. Umeda, N. Iwasa, M. Sonoda, *Chem. Eur. J.* **2003**, *9*, 5549–5559.
- [16] D. C. Zecher, R. West, *J. Am. Chem. Soc.* **1967**, *89*, 153–155.
- [17] a) Y. Rubin, F. Diederich, *J. Am. Chem. Soc.* **1989**, *111*, 6870–6871; b) Y. Rubin, C. B. Knobler, F. Diederich, *J. Am. Chem. Soc.* **1990**, *112*, 1607–1617; c) Y. Rubin, M. Kahr, C. B. Knobler, F. Diederich, C. L. Wilkins, *J. Am. Chem. Soc.* **1991**, *113*, 495–500; d) F. Diederich, Y. Rubin, C. B. Knobler, R. L. Whetten, K. E. Schriver, K. N. Houk, Y. Li, *Science* **1989**, *245*, 1088–1090.
- [18] a) Y. Rubin, C. B. Knobler, F. Diederich, *J. Am. Chem. Soc.* **1990**, *112*, 4966–4968; b) J. Lewis, B. Lin, M. S. Khan, M. R. A. Al-Mandhary, P. R. Raithby, *J. Organomet. Chem.* **1994**, *484*, 161–167; c) M. M. Haley, B. L. Langsdorf, *Chem. Commun.* **1997**, 1121–1122.
- [19] G. A. Adamson, C. W. Rees, *J. Chem. Soc., Perkin Trans. 1* **1996**, 1535–1543.
- [20] a) S. Eisler, R. R. Tykwinski, *J. Am. Chem. Soc.* **2000**, *122*, 10736–10737; b) T. Luu, Y. Morisaki, N. Cunningham, R. R. Tykwinski, *J. Org. Chem.* **2007**, *72*, 9622–9629.
- [21] D. R. Kohn, P. Gawel, Y. Xiong, K. E. Christensen, H. L. Anderson, *J. Org. Chem.* **2018**, *83*, 2077–2086.
- [22] N. Pavliček, P. Gawel, D. R. Kohn, Z. Majzik, Y. Xiong, G. Meyer, H. L. Anderson, L. Gross, *Nat. Chem.* **2018**, *10*, 853–858.
- [23] a) K. Kaiser, L. M. Scriven, F. Schulz, P. Gawel, L. Gross, H. L. Anderson, *Science* **2019**, *365*, 1299–1301; b) L. M. Scriven, K. Kaiser, F. Schulz, A. J. Sterling, S. L. Woltering, P. Gawel, K. E. Christensen, H. L. Anderson, L. Gross, *J. Am. Chem. Soc.* **2020**, *142*, 12921–12924.
- [24] a) G. Gil-Ramírez, D. A. Leigh, A. J. Stephens, *Angew. Chem. Int. Ed.* **2015**, *54*, 6110–6150; *Angew. Chem.* **2015**, *127*, 6208–6249; b) S. Mena-Hernando, E. M. Pérez, *Chem. Soc. Rev.* **2019**, *48*, 5016–5032.
- [25] D. J. Cram, M. E. Tanner, R. Thomas, *Angew. Chem. Int. Ed.* **1991**, *30*, 1024–1027; *Angew. Chem.* **1991**, *103*, 1048–1054.
- [26] M. J. Frampton, H. L. Anderson, *Angew. Chem. Int. Ed.* **2007**, *46*, 1028–1064; *Angew. Chem.* **2007**, *119*, 1046–1083.
- [27] A. M. Blanco-Rodríguez, M. Towrie, J.-P. Collin, S. Záliš, A. Vlček, Jr., *Dalton Trans.* **2009**, 3941–3949.
- [28] M. Melmer, H. Neudeck, *Monatsh. Chem.* **1996**, *127*, 275–290.
- [29] E. Vogel, W. Wiedemann, H. D. Roth, J. Eimer, H. Günther, *Justus Liebigs Ann. Chem.* **1972**, *759*, 1–36.
- [30] A. H. Alberts, H. Wynberg, *Chem. Commun.* **1988**, 748–749.
- [31] P. Bönzli, M. Neuenschwander, *Helv. Chim. Acta* **1991**, *74*, 255–274.
- [32] D. Boyall, F. López, H. Sasaki, D. Frantz, E. M. Carreira, *Org. Lett.* **2000**, *2*, 4233–4236.
- [33] H. Lahlali, K. Jobe, M. Watkinson, S. M. Goldup, *Angew. Chem. Int. Ed.* **2011**, *50*, 4151–4155; *Angew. Chem.* **2011**, *123*, 4237–4241.
- [34] Y. Tobe, N. Iwasa, R. Umeda, M. Sonoda, *Tetrahedron Lett.* **2001**, *42*, 5485–5488.

Entry for the Table of Contents



Text for Table of Contents: Active metal template synthesis has been used to prepare a [2]rotaxane in which the whole dumbbell component is a masked $C_{16}H_2$ polyynediyl. *t*-Butyl substituents make the photo-labile terminal groups bulky enough to act as stoppers and prevent unthreading of the macrocycle. Rotaxanes of this type are promising intermediates in the synthesis of mechanically interlocked carbon-rich architectures, such as cyclocarbons catenanes and carbyne polyrotaxanes.

Institute and/or researcher Twitter usernames: @HLAGroupOx