Social Organization, Home Ranges, and Extraterritorial Forays of Black-backed Jackals on a Game Reserve in South Africa

Journal:	<i>Journal of Wildlife Management and Wildlife Monographs</i>
Manuscript ID	JWM-19-0021.R1
Wiley - Manuscript type:	Note
Date Submitted by the Author:	n/a
Complete List of Authors:	Kamler, Jan; Oxford University, Wildlife Conservation Research Unit Stenkewitz, Ute; University of Iceland's Research Centre at Snaefellsnes Gharajehdaghipour, Tazarve; Mount Revelstokeadn Glacier National Parks Macdonald, David; Oxford University, Wildlife Conservation Research Unit
Keywords:	black-backed jackal, <i>Canis mesomelas</i> , home range overlap, excursion, extraterritorial foray, group formation, social organization, South Africa
Abstract:	<p>We monitored 15 black-backed jackals (<i>Canis mesomelas</i>) from 8 adjacent family groups on Benfontein Game Farm (hereafter Benfontein) in South Africa to investigate their movement patterns and social organization. Jackal family groups consisted of mated pairs (alphas), 0-3 nonbreeding adults (betas), and pups, depending on the season. Mean ($\pm$SE) home-range size of alphas (11.0 ± 0.8 km², $n = 4$) did not significantly differ ($P = 0.321$) from betas (9.8 ± 0.7 km², $n = 8$). Most beta jackals (8 of 10) remained philopatric on Benfontein, apparently due to the high density of springbok (<i>Antidorcas marsupialis</i>), their preferred prey. Three of 5 alphas (60%) and all 8 betas (100%) went on extraterritorial forays (hereafter forays). Generally, betas spent more of their active time on forays (2-20% of time) than alphas (0-3%; $P = 0.048$), and betas went farther on forays (2-8 km) than alphas (2-3 km; $P = 0.003$). The number of forays differed ($P < 0.001$) among seasons, as most forays occurred during summer (64%) when jackals visited neighboring livestock farms, apparently to predate on sheep (<i>Ovis aries</i>). Overall, our results indicate forays by jackals are affected by social status, seasonal availability of preferred prey, and the reproductive cycle of jackals. We recommend several preventative measures to reduce jackal predation on livestock farms near reserves, based on the preferred prey of jackals and the periods when jackals are most likely to travel outside of reserves.</p>

23 June 2019
Jan F. Kamler
Wildlife Conservation Research Unit
Dept. of Zoology, Oxford University
The Recanati-Kaplan Centre, Tubney House
Abingdon Road, Tubney
Abingdon OX13 5QL
United Kingdom
Phone: (+44) 0-1865-611100; Fax: (+44) 0-1865-611101
E-mail: jan.f.kamler@gmail.com

RH: Kamler et al. · Jackal Ecology in South Africa

**Social Organization, Home Ranges, and Extraterritorial Forays of Black-backed Jackals
on a Game Reserve in South Africa**

JAN F. KAMLER¹, Wildlife Conservation Research Unit, The Recanati-Kaplan Centre,
Department of Zoology, Oxford University, Tubney House, Abingdon Road, Tubney,
Abingdon, OX13 5QL, United Kingdom

UTE STENKEWITZ², Wildlife Conservation Research Unit, The Recanati-Kaplan Centre,
Department of Zoology, Oxford University, Tubney House, Abingdon Road, Tubney,
Abingdon, OX13 5QL, United Kingdom

TAZARVE GHARAJEHDAGHIPOUR, Mount Revelstoke and Glacier National Parks, Parks
Canada, P.O. Box 350, Revelstoke, British Columbia, V0E 2S0, Canada

DAVID W. MACDONALD, Wildlife Conservation Research Unit, The Recanati-Kaplan Centre,
Department of Zoology, Oxford University, Tubney House, Abingdon Road, Tubney,
Abingdon, OX13 5QL, United Kingdom

¹ Email: jan.f.kamler@gmail.com

² Present address: University of Iceland's Research Centre at Snæfellsnes, Hafnargata 3, 340
Stykkishólmur, Iceland

2 | Kamler et al.

ABSTRACT We monitored 15 black-backed jackals (*Canis mesomelas*) from 8 adjacent family groups on Benfontein Game Farm (hereafter Benfontein) in South Africa to investigate their movement patterns and social organization. Jackal family groups consisted of mated pairs (alphas), 0-3 nonbreeding adults (betas), and pups, depending on the season. Mean ($\pm$ SE) home-range size of alphas (11.0 ± 0.8 km², $n = 4$) did not significantly differ ($P = 0.321$) from betas (9.8 ± 0.7 km², $n = 8$). Most beta jackals (8 of 10) remained philopatric on Benfontein, apparently due to the high density of springbok (*Antidorcas marsupialis*), their preferred prey. Three of 5 alphas (60%) and all 8 betas (100%) went on extraterritorial forays (hereafter forays). Generally, betas spent more of their active time on forays (2-20% of time) than alphas (0-3%; $P = 0.048$), and betas went farther on forays (2-8 km) than alphas (2-3 km; $P = 0.003$). The number of forays differed ($P < 0.001$) among seasons, as most forays occurred during summer (64%) when jackals visited neighboring livestock farms, apparently to predate on sheep (*Ovis aries*). Overall, our results indicate forays by jackals are affected by social status, seasonal availability of preferred prey, and the reproductive cycle of jackals. We recommend several preventative measures to reduce jackal predation on livestock farms near reserves, based on the preferred prey of jackals and the periods when jackals are most likely to travel outside of reserves.

KEY WORDS black-backed jackal, *Canis mesomelas*, home range overlap, excursion, extraterritorial foray, group formation, social organization, South Africa

Black-backed jackals (*Canis mesomelas*) are common mesocarnivores that have a wide distribution in eastern and southern Africa. Home ranges of black-backed jackals reportedly are 1-34 km² for territorial adults, with larger home ranges for subadults and smaller home ranges

3 | Kamler et al.

for juveniles, in Botswana (McKenzie 1990, Kaunda 2001), Namibia (Hiscocks and Perrin 1988, Jenner et al. 2011), Zimbabwe (Loveridge and Macdonald 2002), and South Africa (Rowe-Rowe 1982, Ferguson et al. 1983, Humphries et al. 2016). Black-backed jackals appear to have a relatively complex social organization, which is based on the degree of territoriality and breeding status within groups. For example, black-backed jackals in South Africa were classified as resident and transient (Humphries et al. 2016), similar to that reported for coyotes (*C. latrans*; Andelt 1985, Kamler and Gipson 2000, Kamler et al. 2005, Morin and Kelly 2017). Residents typically are family groups comprised of a mated pair (hereafter alphas) along with 0-6 other non-breeding adults (McKenzie 1990; Loveridge and Macdonald 2001, Walton and Joly 2003, Jenner et al. 2011). The non-breeding adults in the group, often called helpers (Moehlman 1979, 1983, Ferguson et al. 1983, Walton and Joly 2003), typically are grown offspring from the previous litter that ‘help’ their parents raise the next litter (hereafter betas). However, it is not known if alphas and betas have the same home range sizes, and little information exists about how often they travel together.

For beta jackals, there are several benefits and costs for remaining philopatric, instead of dispersing. The benefits of philopatry include increasing their survivorship, gaining experience raising pups, potential of inheriting the territory from their parents, and increasing their inclusive fitness via kin selection (Moehlman 1983, 1986). Regarding the last point, betas are typically full siblings to the pups in their parents’ next litter, thus betas are on average as closely related to the pups as they would be to their own offspring (Moehlman 1979, 1983). Because the presence of betas was shown to increase pup survival in black-backed jackals, then betas increase their inclusive fitness via kin selection, in lieu of having their own pups (Moehlman 1979, 1983). The costs of betas remaining philopatric include delaying reproduction, and spending time feeding

4 | Kamler et al.

and guarding pups (Moehlman 1983, 1986). Despite the fact that the benefits of philopatry for both betas and alphas appear to outweigh the costs, only 24% of surviving young black-backed jackals remained philopatric as helpers in East Africa (Moehlman 1986). However, the costs and benefits of philopatry are likely affected by availability of food resources throughout the year and the availability of vacant territories (Moehlman 1986, Gese et al. 1996), thus betas in other populations might remain philopatric at higher or lower rates. When betas do disperse, they presumably find mates and establish territories in vacant territories, although little information exists concerning group formation in black-backed jackals.

Territorial animals, from rodents to birds, sometimes travel outside of their territories for short periods, often called extra-territorial forays (hereafter, forays; Soulsbury et al. 2011, Mayer et al. 2017). There appear to be three main purposes for forays, which includes: 1) food acquisition, 2) breeding adults seeking extra-pair copulations, and 3) pre-dispersal exploratory movements by non-breeding subordinates that are seeking vacant territories and possible mates (Soulsbury et al. 2011, Mayer et al. 2017). Among canid species, extraterritorial forays have been reported in wolves (*C. lupus*; van Ballenberghe 1983, Messier 1985, Ballard et al. 1997, Mancinelli and Ciucci 2018), coyotes (Andelt 1985, Kamler and Gipson 2000, Kamler et al. 2004b, Morin and Kelly 2017), bat-eared foxes (*Otocyon megalotis*; Kamler et al. 2017), gray foxes (*Urocyon cinereoargenteus*; Deuel et al. 2017), and *Vulpes* spp. (Kamler et al. 2004a, Soulsbury et al. 2011, Iossa et al. 2008, Kamler and Macdonald 2014), and appear to be common in black-backed jackals as well (Kaunda et al. 2001, McKenzie 1990, Jenner et al. 2011). Similar to that reported for other animal species, one of the main reasons for forays in black-backed jackals appear to be for food acquisition. For example, black-backed jackals went on forays to access marine food resources on the coast of Namibia (Jenner et al. 2011), to scavenge ungulate

5 | Kamler et al.

carcasses in the Kalahari (Nel 1984), and to form larger groups to hunt large ungulates in Botswana (McKenzie 1990). However, the seasonality of forays and degree to which alphas and betas use forays differently (e.g., extra-pair copulations vs. food acquisition) has not been studied in black-backed jackals.

Due to extirpations of large carnivores, black-backed jackals are the dominant predator over most of South Africa, including on small livestock farms where sheep (*Ovis aries*) are often their main prey (Klare et al. 2010, Minnie et al. 2016, Drouilly et al. 2018, Minnie et al. 2018). Consequently, jackals are heavily persecuted on small livestock farms due to their real or perceived predation on livestock (Drouilly et al. 2018, Du Plessis et al. 2015, Minnie et al. 2016). In contrast, on many nature reserves, jackals are not hunted, or are hunted at a much lower intensity compared to livestock farms (Kamler et al. 2013b, Drouilly et al. 2018, Minnie et al. 2018). This has created a source-sink system of jackal populations in South Africa based on anthropogenic mortality, whereby reserves act as source populations of jackals for surrounding livestock farms which act as sinks (Minnie et al. 2018). Furthermore, jackals living on some reserves may go on forays to surrounding livestock farms, apparently due to the abundance of food there (Kaunda 2001), as sheep and other livestock has been found in scats of jackals living in several different reserves (Bothma 1971, Rowe-Rowe 1976, Rowe-Rowe 1983, Kaunda and Skinner 2003). In contrast, in other reserves there was no evidence that jackals left to predate on sheep in surrounding farms (Drouilly et al. 2018), although reasons for these differences among reserves are unknown. Detailed data is needed on movements of jackals between reserves and surrounding livestock farms, as such information would not only help elucidate reasons for forays in this species, but it would also help assess the potential management implication of jackal movements, especially within the mosaic of reserves and livestock farms in South Africa.

We monitored 15 black-backed jackals on Benfontein Game Farm and two small adjoining game reserves (hereafter Benfontein), South Africa, to investigate their movement patterns, social organization, and group formation, with an emphasis on comparing differences between alphas and betas. We compared home-range and core-area overlap within and between groups, and determined how often they foraged together. We predicted that: 1) alphas would have smaller home ranges than betas, 2) home ranges and core areas would overlap more within compared to between groups, and 3) most betas would disperse before their parents' next litter, and would attempt to establish territories with other betas in vacant territories. Benfontein is surrounded by a mosaic of differently managed lands, primarily small livestock farms that raise sheep. Thus, we also compared the frequency, distance, and timing of extraterritorial forays between alphas and betas to help elucidate the purpose of the forays and the extent to which jackals used neighboring livestock farms. We predicted that: 4) betas would go on forays more frequently, and for longer distances, than alphas, 5) alphas would mostly go on forays during the breeding season (winter) to search for extra-pair copulations within Benfontein, 6) betas would mostly go on forays during the breeding season to search for potential mates and vacant territories outside of Benfontein, and 7) both alphas and betas would go on forays to surrounding livestock farms to consume sheep during winter, which is the main lambing season (i.e., food pulse outside Benfontein).

STUDY AREA

We conducted research on Benfontein Game Farm (hereafter, Benfontein; 110 km²), near Kimberley, South Africa (28°53' S, 24°49' E). During our study, Benfontein was managed primarily for wild ungulate species, including springbok (*Antidorcas marsupialis*), blesbok

7 | Kamler et al.

(*Damaliscus dorcas*), and black wildebeest (*Connochaetes gnou*), along with some domestic cattle (*Bos taurus*). All large (>15 kg) carnivore species were extirpated from this area prior to 1900 (Skinner and Chimimba 2005). In addition to jackals, other carnivore species present on Benfontein included cape foxes (*Vulpes chama*), bat-eared foxes (*Otocyon megalotis*), aardwolves (*Proteles cristatus*), caracals (*Caracal caracal*), African wild cats (*Felis silvestris*), black-footed cats (*Felis nigripes*), small-spotted genets (*Genetta genetta*), striped polecats (*Ictonyx striatus*), and at least 5 species of Herpestidae. Aside from one or two annual culls for ungulates on Benfontein, there is relatively little human activity. No carnivore species was heavily persecuted on Benfontein during the study, although jackals were occasionally shot (<3 jackal/yr) during culling operations for ungulates. On other game farms in South Africa, jackals often are heavily persecuted due to their perceived predation on game species (Minnie et al. 2018). Because jackals were not systematically persecuted in Benfontein, this site functioned as a reserve rather than a typical game farm, thus we considered Benfontein a game reserve for the purposes of this paper. In contrast, jackals were heavily persecuted on surrounding livestock farms, to the extent to that territorial jackals were almost completely excluded on the livestock farms (Kamler et al. 2013b). Surrounding livestock farms contained mostly sheep, in addition to small herds of goats (*Capra aegagrus*) and cattle (Kamler et al. 2012a). Many owners of livestock farms also managed small herds of wild ungulates, such as springbok, impala (*Aepyceros melampus*), or gemsbok (*Oryx gazella*), to supplement their incomes with safari hunting (Kamler et al. 2012a). Vegetation on Benfontein and surrounding lands contained elements of 3 major biomes, Savanna, Nama Karoo, and Grassland, although the most dominant was Nama Karoo vegetation (66% of study area; Kamler et al. 2012b). The area had a semi-arid continental climate, with a distinct cold and dry period during winter (Jun to Aug) and a hot and

8 | Kamler et al.

rainy period during summer (Dec to Feb), with intermediate rainfall and temperatures during spring (Sep to Nov) and autumn (Mar to May; Kamler et al. 2012b). See Kamler et al. (2012b, 2013b) for more detailed information about the study site.

One male and one female jackal captured and collared on Benfontein had home ranges which occurred partially and mostly outside of Benfontein, respectively. The male jackal had a home range that extended onto Banksfontein Farm, which was an adjacent small game reserve along the southeastern border of Benfontein (Fig. 1). A female jackal had a home range that was almost entirely within the Mauritzfontein Farm, which was an adjacent game reserve along the western border of Benfontein (Fig. 1). The fences between both reserves and Benfontein were porous, thus jackals could easily move between the properties. Both of these adjacent reserves had similar game species as Benfontein, they did not contain sheep, and jackals were not persecuted, thus we considered them as part of the study site because they were managed similarly to Benfontein.

METHODS

Capture and monitoring

We captured black-backed jackals using padded foothold traps (Woodstream Corp., Lititz, Pennsylvania, USA) set along dirt roads and intersections throughout Benfontein (Kamler et al. 2008). Padded foothold traps were set with a high pan tension to exclude species smaller than jackals (Kamler et al. 2008). Trapping only occurred during the winter (May-August) to avoid capturing pups younger than 9 months old. We set traps just prior to sunset, checked at sunrise, then left closed during the day. We fitted captured jackals with radio collars (Advanced Telemetry Systems, Inc., Isanti, Minnesota, USA) weighing 190 g (i.e., ~2% of body mass). All

jackals were sexed, weighed to the nearest 0.1 kg, and aged according to tooth wear and reproductive condition, then released immediately at the capture site. If jackals showed heavy wear on incisors (Gier 1968) and had large testes ($n = 4$ males) or dark elongated teats (indicating previous nursing; $n = 1$ female) then they were considered adults (>12 months old) when captured and classified as alphas. All other jackals (5 male, 5 female) had minor wear on incisors and small testes or small pink teats, therefore they were considered 9-12 months old when captured and classified as betas. For aging purposes, we assumed a birth date of 1 September based on the approximate date of birth for the pups we observed at den sites during the study. One lower canine tooth was pulled from the skulls of 4 jackals (2 alphas, 2 betas) that died during the study, and sent to Matson's Laboratory (Milltown, Montana, USA) to count cementum annuli and determine ages. The cementum annuli counts indicated that the age at capture was 5 years for an alpha male, 4 years for an alpha female, and 1 year for the betas (1 male, 1 female), indicating our age classifications based on tooth wear were accurate. We followed the animal care and use guidelines of the American Society of Mammalogists (Sikes et al. 2016) and our research protocol (#0401/05) was approved by the Department of Tourism, Environment and Conservation (now Department of Environment and Nature Conservation), Northern Cape Province, Kimberley, South Africa.

Individual jackals were radio-tracked 2-3 times per week throughout the year from a vehicle using a null-peak system consisting of dual four-element yagi antennas. We also radio tracked on foot using 3-element hand-held antennas to locate den sites. When locating jackals, observers took ≥ 2 readings from known telemetry stations < 5 minutes apart. We calculated location estimates using the maximum likelihood estimation option in the program Locate II (Pacer, Inc., Truro, Nova Scotia, Canada). Mean ($\pm$ SE) error of estimated locations was 57.3 ($\pm$

6.3) m when using reference collars ($n = 29$) placed at known locations 0.8-1.5 km from observers (i.e., typical distance when tracking animals). Locations for jackals were assumed to be independent because only 1 location per animal was obtained during 6 to 8-hr tracking sessions. During each tracking session, jackals were located in a random order at random times. However, when a jackal was located, we checked if other jackals were traveling with it. Group size was determined by observing natal dens of collared jackals several times per month during spring (Sep-Nov) to count the number of adult jackals that visited or associated with the pups. Dens were observed using binoculars from >100 m away to avoid disturbing the jackals. For collared jackals within the same group, we determined the percent time they travelled together by counting the total number of times they were located within 100 m of each other when foraging, and dividing by the total number of locations for each jackal.

Home ranges

We determined the home range size of collared jackals using 95% fixed-kernel density estimates (KDE; Worton 1989). We also calculated 50% KDEs for each jackal to represent their core areas, which are areas of concentrated use within home ranges (Laundre and Keller 1981, Springer 1982). To allow for comparisons with previous studies, we also calculated home ranges and core areas using 95% minimum convex polygon (MCP) and 50% MCP, respectively. Home ranges and core areas were calculated for jackals with 40-100 locations and 6-12 months of tracking. If a jackal was monitored in multiple years, then we only included in the analysis the home range from the first year, unless the jackal changed social status and shifted ranges between years. We quantified KDEs, MCPs, isopleths and areas (km²) in the Geospatial Modelling Environment (GME; Beyer 2012). Gaussian KDEs were calculated using the kernel

density command in GME (Beyer 2012) and plug-in bandwidth estimator algorithm in the “ks” package (Duong 2018) of R software (R Version 3.3.2, www.r-project.org, accessed 17 Dec 2016). We compared mean home range sizes between alphas and betas using independent *t*-tests. We excluded from the statistical comparison the home ranges of the two betas that formed a mated pair and became alphas, because their newly established home ranges, which was situated between their two natal territories, were atypically small compared to the other alpha and beta home ranges.

The utilization distribution (UD) provides a useful summary of an animal’s use of space within its home range, by transforming the spatial data into a probability distribution (Fieberg and Kochanny 2005). The UD also provides a more informative measure of overlap between individuals through indices that are a function of the UD. To measure home-range overlap, we used the utilization distribution overlap index (UDOI) which compares the 3-dimensional UD between the intersecting and exclusive areas of use (Fieberg and Kochanny 2005). The UDOI equals 0 when 2 UD’s do not overlap, and 1 if both animals’ UD have 100% overlap with uniformly distributed locations. For 2 non-uniformly distributed UD’s, values >1 indicate a high degree of overlap, whereas values <1 indicate less overlap than expected relative to uniform space use (Fieberg and Kochanny 2005). The biological interpretation of UDOI is similar to that of Hurlbert’s index of niche overlap (Hurlbert 1978) that assumes 2 animals use space independently of one another. We investigated the spatial overlap between all jackals within the same group, and between jackals of neighboring groups. The UDOI was calculated in R software at both 95% and 50% UD isopleth levels, to represent home ranges and core areas, respectively, using KernSmith (Wand 2015) and adehabitatHR (Calenge 2015) packages, and code from Fieberg (2014). The UDOI was calculated between neighboring jackals only if they were

monitored during the same period, and results were compared between seasons using Mann-Whitney *U* tests. To allow comparisons with previous studies, we also calculated percent area overlap using 95% and 50% MCPs. Percent area overlap was determined for 2 jackals by multiplying the area overlap by 2, and then dividing by the total area of both polygons (Cole 1949).

Extraterritorial forays

Extraterritorial forays, hereafter forays, of collared jackals were defined as solitary movements of short duration (1-3 days) ≥ 2 km from their home range boundary (Kamler and Gipson 2000). All known forays were either to surrounding livestock farms outside of Benfontein, or to the core areas of neighboring jackal groups within Benfontein. Foray locations were not included in home-range calculations (Andelt 1985, Kamler and Gipson 2000). We determined the length of known forays by measuring the distance between the foray location and the jackal's home-range boundary, as calculated by 95% KDE. Of 36 total forays recorded during the study, we could not calculate distance for 10 forays because the beta jackals traveled too far away to obtain a location (we did not have permission to radio track on all surrounding properties). Thus, our estimates of foray distances should be considered conservative, at least for the beta jackals. We calculated the percent time that each jackal went on forays by dividing the total number of forays by the total number of active locations. We compared the percent time alphas went on forays with that of betas using a Mann-Whitney *U* test.

To assess seasonal differences in number of forays, we divided the year into spring (Sep-Nov), summer (Dec-Feb), autumn (Mar-May), and winter (Jun-Aug), to parallel major changes in climate and vegetation, and different biological periods for jackals and their main prey (Klare

et al. 2010). For example, the two birthing periods of springbok, the main prey of jackals on Benfontein, occurred during spring and autumn (Klare et al. 2010). Winter was the main lambing season for sheep on surrounding livestock farms (Kamler et al. 2012a). Jackals gave birth during early spring, thus their breeding season occurred in winter given their gestation lasts 60-65 days (Walton and Joly 2003). Number of forays from all jackals were grouped and compared among seasons using a chi-square goodness-of-fit test.

RESULTS

We captured, radio-collared, and monitored 15 black-backed jackals from 8 adjacent family groups from July 2005 to March 2008 (Fig. 1). At the time of capture, jackals were classified as 5 alphas and 10 betas. Two betas later formed a mated pair on Benfontein and had pups, thus they were classified as alphas after they left their respective natal territories and began associating with each other. Jackal family group consisted of 2-5 adults, which included the mated pair (alphas) and 0-3 nonbreeding adults (betas). The only jackal group that we observed which did not have betas was the new group formed by the two former betas mentioned above. Within the same group, betas travelled with other betas or alphas only 4.0% (range = 2.1-5.4%, $n = 5$) of the time when foraging, although all monitored betas and alphas often associated with each other at natal dens when pups were present. Two alphas travelled with each other 23.9% of the time when foraging.

There was sufficient data to calculate home ranges for 11 of 15 jackals, because 2 jackals were killed by humans 1-3 months after being collared, and another 2 jackals dispersed from the study site about 2 months after being collared. The mean ($\pm$ SE) home-range size for all jackals was $10.24 \pm 0.55 \text{ km}^2$ using KDE, and $9.12 \pm 0.77 \text{ km}^2$ using MCP. Mean home-range sizes

(KDE) were not significantly different ($t_{10} = 1.045$, $P = 0.321$) between alphas and betas (Table 1). Similarly, mean core-area sizes (KDE) were not significantly different ($t_{10} = 0.018$, $P = 0.986$) between alphas and betas (Table 1). Home-range overlap (UDOI 95) between individual jackals was higher ($Z = -3.450$, $n = 30$, $P < 0.001$) within groups ($\bar{X} \pm SE = 1.177 \pm 0.113$) than between groups (0.010 ± 0.003 , Fig. 1). The mean ($\pm SE$) core-area overlap (UDOI 50) for jackals within the same group was 0.538 ± 0.139 , whereas core areas never overlapped between jackals in adjacent groups. When using MCP, individual home ranges overlapped 79.2% (range = 64.0-90.0%, $n = 4$) within groups and 1.6% (0.0-16.1%, $n = 25$) between groups, whereas core areas overlapped 68.8% (53.2-90.9%) within groups, but never overlapped between groups.

A male (13-M) and female (16-F) beta from neighboring groups formed a pair when they were both about 18 months old, and thereafter used a home range in the area between their respective core areas (Fig. 2). The subsequent home range of 13-M (6.1 km²) over the next 12 months was the smallest we recorded for any jackal, and the home range of the 16-F (6.4 km²) was only slightly larger (Table 1). Their new home ranges overlapped extensively with jackals from both his natal home range (MCP = 14.2-19.4%) and her natal home range (26.9-31.0%, Fig. 2). This pair produced at least one litter of pups, but then 16-F died due to a non-traumatic cause, possibly disease. Thereafter, 13-M joined the neighboring group, which was the natal group of 16-F, and his home range expanded to 11.0 km², which was the typical size of a jackal home range on Benfontein (Table 1).

We recorded 36 forays during the study. Most jackals went on forays, including 3 of 5 alphas (60%) and all 8 betas (100%) that were monitored more than 2 months. Overall, betas spent more of their active time on forays (2-20% of time) than alphas (0-3%; $P = 0.048$), and betas went farther on forays (2-8 km) than alphas (2-3 km; $P = 0.003$). The number of forays

differed ($P < 0.001$) among seasons, as most forays occurred during summer (63.9%) followed by winter (19.4%, Fig. 3). In summer, 22 of 23 forays (95.7%) occurred when alpha and beta jackals visited neighboring livestock farms. In winter, 5 of 7 forays (71.4%) occurred when beta jackals visited core areas of neighboring jackal groups. There was only 1 foray recorded by an alpha during winter, and that was to feed on the carcass of a springbok that had been shot in a culling operation on Benfontein. During spring and autumn combined, there were 6 total forays: 3 to livestock farms and 3 to neighboring jackal groups. Within the same family group, some betas rarely went on forays (2-4%), similar to alphas, whereas other betas went on forays more frequently (12-20%). Only 2 betas dispersed during the study, and both occurred during winter when the jackals were 11-12 months old.

DISCUSSION

The home range sizes of alphas and betas were similar, which did not support our first prediction. However, because sample size and statistical power were relatively low, a difference might have been detected with a larger sample size. Nonetheless, betas went on forays more frequently and farther than alphas, consequently most betas did have more wide-ranging movements than alphas. Thus, if we had not removed forays from home-range calculations, then some betas would have had much larger home range sizes. When foraging, the alphas spent more time with each other than with betas, although all jackals spent a majority of their time foraging alone. Home range and core area overlap was higher within groups than between groups (Fig. 1), which supported our second prediction. Our results were similar to that previously reported for black-backed jackals, in which overlap between neighboring groups was $<10\%$ whereas overlap within groups was $>50\%$ (Rowe-Rowe 1982, Ferguson et al. 1983, Loveridge and Macdonald

2001). Although betas seldom traveled with alphas, betas often visited natal dens and associated with pups, whether or not alphas were present, similar to that reported in other jackal studies (Moehlman 1983). We could not confirm if betas brought food to pups, although we assumed they did because food provisioning by betas was previously reported in black-backed jackals (Moehlman 1979, 1983). Additionally, the presence of betas at natal dens likely provided extra protection for the pups, especially when the alphas were absent (Moehlman 1979, 1983, Jenner et al. 2011). In contrast to other studies, we did not record transient jackals during our study, probably due to the high rates of philopatry by the beta jackals.

Most beta jackals (8 of 10) stayed on Benfontein within their natal groups beyond 12 months of age, and after their presumed parents had the next litter, which did not support our third prediction. Additionally, 2 betas (both females) stayed within their natal territory beyond 24 months of age without producing pups. This was surprising given that betas made numerous forays to surrounding livestock farms, yet none attempted to establish territories there. Although the rate of philopatry by betas was higher than that observed in East Africa (24%; Moehlman 1986), our results were consistent with that reported by Minnie et al. (2016) in South Africa, who showed that on reserves with low human-caused mortality, only about 50% of female black-backed jackals that were 3-4 year old had become pregnant, suggesting a relatively high proportion of older jackals were betas. Beta females staying in their natal ranges for two subsequent litters to help their parents was reported previously for black-backed jackals (Moehlman 1979, Ferguson et al. 1983) and coyotes (Andelt 1985, Gese et al. 1996, Kamler and Gipson 2000), so it may not be unusual behavior for medium-sized canids. Because black-backed jackals can reproduce at 12 months of age (Minnie et al. 2016), there must have been

substantial benefits for beta jackals on Benfontein to forego their own reproduction and remain within their natal range beyond 12 months old.

There are several possible explanations for the high rates of philopatry and lack of group formation by betas outside of Benfontein. Firstly, because all philopatric betas associated with the next litter of their presumed parents, the betas likely benefited from gaining experience with raising pups while increasing their inclusive fitness via kin selection (Moehlman 1979, 1986). Secondly, although betas made numerous forays to livestock farms, they probably did not meet potential mates during their short trips there because many of livestock farms lacked territorial groups due to high rates of human persecution (Kamler et al. 2013b). Thirdly, the preferred prey and most highly consumed species of jackals on Benfontein was springbok (*Antidorcas marsupialis*), which had a relatively high density (22 springbok/km²; Klare et al. 2010). In contrast, surrounding livestock farms either did not contain springbok, or had a relatively low density (5 springbok/km²; Kamler et al. 2012a). Additionally, although sheep were consumed by jackals on surrounding livestock farms, wild ungulates were selectively consumed over sheep in most seasons (Kamler et al. 2012a), and densities of sheep (16 sheep/km²) were lower than that of springbok on Benfontein (Kamler et al. 2012a). Therefore, the higher density of preferred prey on Benfontein may have caused most betas to remain philopatric, instead of dispersing into relatively unknown areas with much lower preferred prey. Lack of dispersal by subordinate group members, despite the existence of vacant neighboring territories, has been observed in other canid species (Macdonald et al. 2004) although the reasons for this behavior are not always clear. In a review of helping behavior, Emlen (1982) indicated that grown offspring can delay dispersal even if breeding opportunities or vacant territories exist, if the cost of rearing their own young is perceived to be prohibitive. Thus, beta jackals might have perceived the livestock farms

18 | Kamler et al.

as too risky to establish territories, either due to the lower abundance of preferred prey or to lack of other jackals. Nonetheless, perhaps with a larger sample size of betas, some betas might have established territories on the livestock farms.

Betas spent more of their active time on forays, and went farther on forays, than alphas, which supported our fourth prediction. More extensive forays by betas has been reported in other mammal species (Mayer et al. 2017), including coyotes (Kamler and Gipson 2000, Kamler et al. 2004b) and gray wolves (Messier 1985), as forays allow betas to become familiar with surrounding areas and search for potential mates and vacant territories. Overall, the number of forays differed among seasons, as most forays for alphas and betas occurred during summer, which did not support our fifth, sixth, and seventh predictions. We predicted that both alphas and betas would go on forays mostly in winter, which is the jackal breeding season, because we assumed alphas would search for extra-pair matings in neighboring groups (prediction 5) and betas would search for new mates in vacant areas outside of Benfontein (prediction 6). Additionally, most sheep lambing on surrounding farms occurred in winter, and we assumed jackals on Benfontein would take advantage of this food pulse and travel to the livestock farms to consume lambs (prediction 7). Instead, most forays occurred in summer when jackals visited surrounding livestock farms. On Benfontein, summer coincided with two important aspects of the jackal diet: the lowest amount of springbok consumed (26% of biomass consumed), and the highest amount of sheep consumed by jackals (8%; Fig. 3; Klare et al. 2010). Sheep did not occur on Benfontein, so the consumption of sheep by jackals in summer had to occur during their forays to surrounding livestock farms, similar to that reported on other reserves (Bothma 1971, Rowe-Rowe 1976, Rowe-Rowe 1983, Kaunda and Skinner 2003). Similar movements of jackals

were reported in Namibia, where resident jackals often went on forays of up to 20 km to feed on seal pups along the coast (Jenner et al. 2011).

On Benfontein, consumption of springbok by jackals peaked during the two springbok birthing seasons in spring and autumn, indicating jackals consumed mostly springbok fawns (Fig. 3, Klare et al. 2010). During summer, only subadult and adult springbok would have been available to jackals on Benfontein, decreasing the accessibility of this prey species to jackals. Although jackals can kill adult springbok (Krofel 2008, Klare et al. 2010), they are clearly more difficult and dangerous to bring down compared to small fawns. Thus, it appears jackals chose to consume more subadult and adult sheep on surrounding livestock farms in summer when springbok fawns became unavailable, probably because adult sheep are easier and less dangerous to bring down than adult springbok. Our results indicate that food consumption was the main purpose of forays for both alphas and betas, and that the number of forays peaked during low availability or accessibility of preferred prey within Benfontein (summer), but not during a major food pulse outside of Benfontein (winter). Our results are consistent with previous studies that showed forays in jackals, coyotes, and wolves occurred more during periods of food stress (Bekoff and Wells 1980, Messier 1985, McKenzie 1990, Ballard et al. 1997).

During winter, the jackal breeding season, there was a secondary peak in number of forays (Fig. 3). Most forays during winter were by beta jackals visiting neighboring groups within Benfontein, indicating the primary purpose for forays during this season was to seek mates or inter-group breeding opportunities. In other canid species, alphas sometimes went on forays to neighboring groups during the breeding season in search for extra-pair copulations (Gese et al. 1996, White et al. 2000, Iossa et al. 2008, Deuel et al. 2017), although this was not recorded on Benfontein, which could have been due to the low sample size of alphas. Betas

sought breeding opportunities within Benfontein instead of the livestock farms, and consequently this resulted in pair bonding and group formation by two neighboring betas within Benfontein (Fig. 2). Group formation by betas on the edge of or in between their natal territories was not reported in previous studies of jackals, although it was reported in other canid species, including gray wolves (Mech and Boitani 2003), red foxes (*Vulpes vulpes*; Baker et al. 2000), and crab-eating foxes (*Cerdocyon thous*; Macdonald and Courtenay 1996). Such territorial division within canid family groups often leads to clusters of related individuals (see Kamler et al. 2013a and references within) and may serve to reduce conflict and infanticide between neighboring groups (Macdonald and Courtenay 1996). Our results support the conclusion that territorial inheritance or sharing in canids is one of the benefits for betas remaining philopatric (Bekoff and Wells 1980, Moehlman 1986, Gese et al. 1996, Macdonald et al. 2004).

MANAGEMENT IMPLICATIONS

If predation by black-backed jackals on sheep is too high on livestock farms near reserves, our results can help managers identify several preventative measures to reduce livestock losses, because forays in black-backed jackals were affected by social status, seasonal availability of preferred prey, and the reproductive cycle of jackals. However, caution should be used when extrapolating our results to other areas, because forays onto livestock farms by jackals are likely dependent on several factors (e.g., abundance of jackals, their preferred prey, and sheep) and probably vary spatially and temporally. Nonetheless, we recommend several preventative measures to reduce livestock losses based on the results of our study. Firstly, because there was a strong seasonal variation of forays onto livestock farms, managers of livestock farms could implement seasonal measures to reduce livestock predation during summer, the peak season

when forays occurred. For example, most livestock farms in the area used rotational grazing, and thus sheep could intentionally be moved to fields further away from reserves during summer. Additionally, herders could be seasonally employed during summer, thereby reducing the cost of annually employing them. Furthermore, hunting of jackals on livestock farms could increase during summer. Secondly, several of the livestock farms in the area maintained small herds of springbok, a small hider species (Klare et al. 2010), to supplement their income with safari hunting. However, this species is the most preferred prey of black-backed jackals (Klare et al. 2010), thus maintaining herds of this species on their property is likely to attract jackals, potentially more so than the sheep. Instead, livestock managers could switch to ungulate species that are less preferred by jackals, such as follower species (e.g., blesbok) or hidiers with larger body sizes (e.g., gemsbok; Klare et al. 2010). Finally, other alternative methods such as using livestock guarding dogs might be particularly effective (Potgieter et al. 2013), especially if this reduces forays onto livestock farms and discourages beta jackals from establishing territories there. Realistically, it may be impossible to completely prevent jackal predation on sheep, especially on farms that are near to reserves, but using one or more of the above preventative measures could help reduce livestock losses to acceptable levels.

ACKNOWLEDGMENTS

We thank De Beers Consolidated Mines for allowing us access to their property, and providing support for this project. A special thanks goes to S. Cade, P. Gibbs, M. Hauptfleisch, and J. Kruger of De Beers; C. Anderson, T. Anderson, A. Sliwa, and B. Wilson from the McGregor Museum, Kimberley, and; M. Anderson and E. Herrmann from the Department of Tourism, Environment and Conservation, Kimberley. We also thank L. Mayer and S. Visagie for making

traps, Z. Davidson and A. Loveridge for equipment loans, B. Cypher for donation of radio collars, and N. F. Jacobsen, U. Klare, B. Dean, D. Nkosi, T. Keswick, and I. Midoko for help with data collection. Funding for J. F. Kamler was provided by a Research Fellowship from the Wildlife Conservation Society, New York, and a Marie Curie Fellowship from the European Commission, Brussels, Belgium. Additional support was provided by British Airways, and grants to DWM from the Peoples' Trust for Endangered Species.

LITERATURE CITED

- Andelt, W. F. 1985. Behavioral ecology of coyotes in South Texas. *Wildlife Monographs* 94:1-45.
- Baker, P. J., S. M. Funk, S. Harris, and P. C. L. White. 2000. Flexible spatial organization of urban foxes, *Vulpes vulpes*, before and during and outbreak of sarcoptic mange. *Animal Behaviour* 59:127-146.
- Ballard, W. B., L. A. Ayres, P. R. Krausman, D. J. Reed, and S. G. Fancy. 1997. Ecology of wolves in relation to a migratory caribou herd in northwest Alaska. *Wildlife Monographs* 135:1-47.
- Bekoff, M., and W. C. Wells. 1980. The social ecology of coyotes. *Scientific American* 242:130-148.
- Beyer. H. L. 2012. Geospatial Modelling Environment (Version 0.7.2.1).
www.spataleecology.com/gme
- Bothma, J. D. P. 1971. Food of *Canis mesomelas* in South Africa *Zooligca Africana* 6:195-203.

- 506 Calenge, C. 2015. Home range estimation in R: the adehabitatHR package. The Comprehensive
507 R Archive Network. [https://cran.r-](https://cran.r-project.org/web/packages/adehabitatHR/adehabitatHR.pdf)
508 [project.org/web/packages/adehabitatHR/adehabitatHR.pdf](https://cran.r-project.org/web/packages/adehabitatHR/adehabitatHR.pdf)
- 509 Cole, L. C. 1949. The measurement of interspecific association. Ecology 30:411-424.
- 510 Deuel, N. R., L. M. Conner, K. V. Miller, M. J. Chamberlain, M. J. Cherry, and L. V.
511 Tannenbaum. 2017. Gray fox home range, spatial overlap, mated pair interactions and
512 extra-territorial forays in southwestern Georgia, USA. Wildlife Biology 2017:wlb.00326.
- 513 Duong, T. (2018). ks: Kernel Smoothing. R package version 1.11.0. [https://CRAN.R-](https://CRAN.R-project.org/package=ks)
514 [project.org/package=ks](https://CRAN.R-project.org/package=ks)
- 515 Drouilly, M, N. Nattrass, and M. J. O’Riain. 2018. Dietary niche relationships among predators
516 on farmland and a protected area. Journal of Wildlife Management 82:507-518.
- 517 Du Plessis, J. J., N. L. Avenant, and H. O. De Wall. 2015. Quality and quantity of the scientific
518 information available on black-backed jackals and caracals: contributing to human-
519 predator conflict management? African Journal of Wildlife Research 45:138-157.
- 520 Emlen, S. T. 1982. The evolution of helping. I. An ecological constraints model. American
521 Naturalist 119:29-39.
- 522 Ferguson, J. W. H., J. A. J. Nel, and M. J. De Wet. 1983. Social organization and movement
523 patterns of black-backed jackals *Canis mesomelas* in South Africa. Journal of Zoology
524 (London) 199:487-502.
- 525 Fieberg, J. 2014. Home range overlap indices implemented using kernel density estimators with
526 plug-in smoothing parameters and Program R. Retrieved from the Data Repository for
527 the University of Minnesota. <http://dx.doi.org/10.13020/D6G59W>

- 528 Fieberg, J., and C. O. Kochanny. 2005. Quantifying home-range overlap: The importance of the
529 utilization distribution. *Journal of Wildlife Management* 69:1346-1359.
- 530 Gese, E. M., R. L. Ruff, and R. L. Crabtree. 1996. Social and nutritional factors influencing the
531 dispersal of resident coyotes. *Animal Behaviour* 52:1025-1043.
- 532 Gier, H. T. 1968. Coyotes in Kansas. Revised. Kansas State College Agricultural Experiment
533 Station Bulletin 393.
- 534 Hiscocks, K., and M. R. Perrin. 1988. Home range and movements fo black-backed jackals at
535 Cape Cross Seal Reserve, Namibia. *South African Journal of Wildlife Research* 18:97-
536 100.
- 537 Humphries, B. D., T. Ramesh, T. R. Hill, and C. T. Downs. 2016. Habitat use and home range of
538 black-backed jackals (*Canis mesomelas*) on farmlands in the Midlands of KwaZulu-
539 Natal, South Africa. 51:37-45.
- 540 Hurlbert, S. H. 1978. The measurement of niche overlap and some relatives. *Ecology* 59:67-77.
- 541 Iossa, G., C. D. Soulsbury, P. J. Baker, and S. Harris. 2008. Body mass, territory size, and life
542 history tactics in a socially monogamous canid, the red fox *Vulpes vulpes*. *Journal of*
543 *Mammalogy* 89:1481-1490.
- 544 Jenner, N., J. Groombridge, and S. M. Funk. 2011. Commuting, territoriality and variation in
545 group and territory size in a black-backed jackal population reliant on a clumped,
546 abundant food resource in Namibia. *Journal of Zoology (London)* 284:231-238.
- 547 Kamler, J. F., W. B. Ballard, E. M. Gese, R. L. Harrison, and S. M. Karki. 2004a. Dispersal
548 characteristics of swift foxes. *Canadian Journal of Zoology* 82:1837-1842.
- 549 Kamler, J. F., W. B. Ballard, R. L. Gilliland, P. R. Lemons, and K. Mote. 2005. Home range and

- 550 habitat use of coyotes in an area of native prairie, farmland, and CRP fields. American
551 Midland Naturalist 153:203-211.
- 552 Kamler, J. F., W. B. Ballard, R. L. Gilliland, and K. Mote. 2004b. Coyote (*Canis latrans*)
553 movements relative to cattle (*Bos taurus*) carcass areas. Western North American
554 Naturalist 64:53-58.
- 555 Kamler, J. F., and P. S. Gipson. 2000. Space and habitat use by resident and transient coyotes.
556 Canadian Journal of Zoology 78:2106-2111.
- 557 Kamler, J. F., M. M. Gray, A. Oh, and D. W. Macdonald. 2013a. Genetic structure, spatial
558 organization, and dispersal in two populations of bat-eared foxes. Ecology and Evolution
559 3:2892-2902.
- 560 Kamler, J. F., N. F. Jacobsen, and D. W. Macdonald. 2008. Efficiency and safety of Soft Catch
561 traps for capturing black-backed jackals and excluding non-target species. South African
562 Journal of Wildlife Research 38:113-116.
- 563 Kamler, J. F., U. Klare, and D. W. Macdonald. 2012a. Seasonal diet and prey selection of black-
564 backed jackals on a small-livestock farm in South Africa. African Journal of Ecology
565 50:299-307.
- 566 Kamler, J. F., and D. W. Macdonald. 2014. Social organization, survival, and dispersal of cape
567 foxes (*Vulpes chama*) in South Africa. Mammalian Biology 79:64-70.
- 568 Kamler, J. F., S. Rostro-García, and D. W. Macdonald. 2017. Seasonal changes in social
569 behavior and movements of bat-eared foxes in South Africa: disease implications. Journal
570 of Mammalogy 98:1426-1433.
- 571 Kamler, J. F., U. Stenkewitz, U. Klare, N. F. Jacobsen, and D. W. Macdonald. 2012b. Resource

- 572 partitioning among cape foxes, bat-eared foxes, and black-backed jackals in South Africa.
573 Journal of Wildlife Management 76:1241-1253.
- 574 Kamler, J. F., U. Stenkewitz, and D. W. Macdonald. 2013b. Lethal and sublethal effects of
575 black-backed jackals on cape foxes and bat-eared foxes. Journal of Mammalogy 94:295-
576 306.
- 577 Kaunda, S. K. K. 2001. Spatial utilization by black-backed jackals in southeastern Botswana.
578 African Zoology 36:143-152.
- 579 Kaunda, S. K. K., and J. D. Skinner. 2003. Black-backed jackal diet at Mokolodi Nature
580 Reserve, Botswana. African Journal of Ecology 41:39-46.
- 581 Klare, U., J. F. Kamler, U. Stenkewitz, and D. W. Macdonald. 2010. Diet, prey selection, and
582 predation impact of black-backed jackals in South Africa. Journal of Wildlife
583 Management 74:1030-1042.
- 584 Krofel, M. 2008. Opportunistic hunting behavior of black-backed jackals in Namibia. African
585 Journal of Ecology 46:220-222.
- 586 Laundre, J. W., and B. L. Keller. 1981. Home-range use by coyotes in Idaho. Animal Behaviour
587 29:449-461.
- 588 Loveridge, A. J., and D. W. Macdonald. 2001. Seasonality in spatial organization and dispersal
589 of sympatric jackals (*Canis mesomelas* and *C. adustus*): implications for rabies
590 management. Journal of Zoology (London) 253:101-111.
- 591 Macdonald, D. W., and O. Courtenay. 1996. Enduring social relationships in a population
592 of crab-eating zorros, *Cerdocyon thous*, in Amazonian Brazil (Carnivore, Canidae).
593 Journal of Zoology (London) 239:329-355.
- 594 Macdonald, D. W., S. Creel, and M. G. L. Mills. 2004. Canid society. Pages 85-106 in D. W.

- 595 Macdonald and C. Sillero-Zubiri, editors. The biology and conservation of wild canids.
596 Oxford University Press, Oxford, United Kingdom.
- 597 Mancinelli, S, and P. Ciucci. 2018. Beyond home: Preliminary data on wolf extraterritorial
598 forays and dispersal in Central Italy. *Mammalian Biology* 93:51-55.
- 599 Mayer, M., A. Zedrosser, and F. Rossell. 2017. Extra-territorial movements differ between
600 territory holders and subordinates in a large, monogamous rodent. *Scientific Reports*
601 7:15261.
- 602 McKenzie, A. A. 1990. Cooperative hunting in the black-backed jackal *Canis mesomelas*,
603 Schreber. Dissertation, University of Pretoria, Pretoria, South Africa.
- 604 Mech, L. D., and L. Boitani. 2003. Wolf social ecology. Pages 1-34 in L. D. Mech and L.
605 Boitani, editors. *Wolves: behavior, ecology, and conservation*. University of Chicago
606 Press, Chicago, Illinois, USA.
- 607 Messier, F. 1985. Solitary living and extraterritorial movements of wolves in relation to social
608 status and prey abundance. *Canadian Journal of Zoology* 63:239-245.
- 609 Minnie, L., A. Gaylard, and G. I. H. Kerley. 2016. Compensatory life-history responses of a
610 mesopredator may undermine carnivore management efforts. *Journal of Applied Ecology*
611 53:379-387.
- 612 Minnie, L., A. Zalewski, H. Zalewska, and G. I. H. Kerley. 2018. Spatial variation in
613 anthropogenic mortality induces a source-sink system in a hunted mesopredator.
614 *Oecologia* 186:939-951.
- 615 Moehlman, P. D. 1979. Jackal helpers and pup survival. *Nature* 277:382-383.
- 616 Moehlman, P. D. 1983. Socioecology of silverbacked and golden jackals (*Canis mesomelas* and
617 *Canis aureus*). Pages 423-453 in J. F. Eisenberg, and D. G. Kleiman editors. *Advances*

- 618 in the study of mammalian behavior. American Society of Mammalogists, Special
619 Publication Number 7, Pittsburgh, Pennsylvania, USA.
- 620 Moehlman, P. D. 1986. Ecology and cooperation in canids. Pages 64-86 *in* D. I. Rubenstein and
621 R. W. Wrangham, editors. Ecological aspects of social evolution: birds and mammals.
622 Princeton University Press, Princeton, New Jersey, USA.
- 623 Morin, D. J., and M. J. Kelly. 2017. The dynamic nature of territoriality, transience and biding in
624 an exploited coyote population. *Wildlife Biology* 2017:wlb.00335.
- 625 Nel, J. A. J. 1984. Behavioural ecology of canids in the south-western Kalahari. *Koedoe*
626 S:229-235.
- 627 Potgieter, G. C., L. L. Marker, N. L. Avenant, and G. I. H. Kerley. 2013. Why Namibian farmers
628 are satisfied with the performance of their livestock guarding dogs. *Human Dimensions*
629 *of Wildlife* 18:403-415.
- 630 Rowe-Rowe, D. T. 1976. Food of the black-backed jackal in nature conservation and farming
631 areas in Natal. *East African Wildlife Journal* 14:345-348.
- 632 Rowe-Rowe, D. T. 1982. Home range and movements of black-backed jackals in an African
633 montane region. *South African Journal of Wildlife Research* 12:79-84.
- 634 Rowe-Rowe, D. T. 1983. Black-backed jackal diet in relation to food availability in the Natal
635 Drakensberg. *South African Journal of Wildlife Research* 13:17-23.
- 636 Sikes, R. S., and The Animal Care and Use Committee of the American Society of
637 Mammalogists. 2016. 2016 Guidelines of the American Society of Mammalogists for the
638 use of wild mammals in research and education. *Journal of Mammalogy* 97:663-688.
- 639 Skinner, J. D., and C. T. Chimimba. 2005. The mammals of the southern African subregion.
640 Cambridge University Press, Cambridge, United Kingdom.

- 641 Soulsbury, C. D., G. Iossa, P. J. Baker, P. C. L. White, and S. Harris. 2011. Behavioral and
642 spatial analysis of extraterritorial movements in red foxes (*Vulpes vulpes*). Journal of
643 Mammalogy 92:190-199.
- 644 Springer, J. T. 1982. Movement patterns of coyotes in south central Washington. Journal of
645 Wildlife Management 46:191-200.
- 646 van Ballenberghe, V. 1983. Extraterritorial movements and dispersal of wolves in southcentral
647 Alaska. Journal of Mammalogy 64:168-171.
- 648 Walton, L. R., and D. O. Joly. 2003. *Canis mesomelas*. Mammalian Species 715:1-9.
- 649 Wand, M. 2015. KernSmooth: Functions for kernel smoothing Supporting Wand & Jones (1995).
650 R package version 2.23-15. <https://CRAN.R-project.org/package=KernSmooth>
- 651 White, P. J., K. Ralls, and D. B. Sniff. 2000. Nocturnal encounters between kit foxes. Journal of
652 Mammalogy 81:456-461.
- 653 Worton, B. J. 1989. Kernel methods for estimating the utilization distribution in home-range
654 studies. Ecology 70:164-168.
- 655 *Associate Editor: Ryan Long*
- 656
- 657 **Summary for online Table of Contents**
- 658 On a reserve in South Africa, forays by black-backed jackals were shown to be affected by social
659 status, seasonal availability of preferred prey, and the reproductive cycle of jackals. We
660 recommend several preventative measures to reduce jackal predation on livestock farms near
661 reserves, based on the preferred prey of jackals and the periods when jackals are most likely to
662 travel outside of reserves.
- 663

Figure 1. Home ranges (95% fixed kernel density estimates [KDE]; thin black outlines) and core areas (50% KDE; solid gray polygons) of 9 black-backed jackals, comprising 7 family groups, that were monitored simultaneously on Benfontein Game Farm and two smaller adjoining game reserves (thick black outlines), South Africa, during 2006-2007. Note that the smaller ovals represent disjunct parts of jackal home ranges and core areas. The 3 dots and connected dashed lines are examples of extraterritorial forays taken by 3 different jackals to neighboring sheep farms.

Figure 2. Group formation by black-backed jackals on Benfontein Game Farm (thick black outline), South Africa. During the initial year of monitoring (2005-2006), a beta male and alpha male occupied similar home ranges (95% fixed kernel density estimate [KDE]) that represented a family group (A; thin black outlines), whereas a beta female occupied a home range that represented an adjacent family group (B; thin black outline). During the second year of monitoring (2006-2007), the beta male and beta female formed a mated pair and produced pups within the overlap area of their respective family groups (C; light gray represents their home ranges [95% KDE], whereas dark gray represents their core areas [50% KDE]). Note that the smaller ovals represent disjunct parts of jackal home ranges.

Figure 3. Number of extraterritorial forays by season for black-backed jackals on Benfontein Game Farm, South Africa, during 2005-2008. Also shown is the seasonal amount of springbok and domestic sheep consumed by black-backed jackals on Benfontein during the same period, taken from Klare et al. (2010).

Table 1. Home-range sizes (95%) and core-area sizes (50%) of 11 black-backed jackals (*Canis mesomelas*) that were monitored on Benfontein Game Farm, South Africa, 2005-2008 (KDE = fixed kernel density estimate, MCP = minimum convex polygon).

ID-sex	KDE (km ²)		MCP (km ²)	
	95%	50%	95%	50%
Alphas				
13-M II ^a	6.11	1.54	5.22	1.33
13-M III ^b	10.95	2.37	9.81	2.74
16-F II ^a	6.38	1.47	5.39	1.49
17-M	11.87	2.84	8.84	3.90
53-M	8.80	1.45	9.50	1.16
54-M	12.55	2.60	11.25	2.51
Mean ± SE	11.04 ± 0.82	2.32 ± 0.30	9.85 ± 0.51	2.58 ± 0.56
Betas				
13-M I	10.60	2.49	7.28	1.64
16-F I	10.41	3.07	7.53	2.05
35-F	7.85	1.51	5.11	1.53
45-F	6.23	1.40	5.17	0.58
52-F	11.96	3.50	9.37	3.00
48-M	8.95	2.18	9.29	1.37
49-M	10.92	2.19	14.24	1.74

710	51-M	11.73	2.11	12.01	1.73
711	Mean ± SE	9.83 ± 0.71	2.31 ± 0.25	8.75 ± 1.13	1.70 ± 0.23

712 _____

713 ^a Home ranges of 13-M II and 16-F II were not included in the mean because they were former
714 betas who had atypically small home range sizes for alphas after they had mated and established
715 their home ranges in between their respective natal territories.

716 ^b The home range of 13-M III was included in the mean because after his mate (16-F) died, he
717 joined the adjacent jackal group, which was the natal group of his former mate.

718

Fig. 1

Fig 2