

Isotopic evidence for a change in mobility and landscape use patterns between the Neolithic and Early Bronze Age in western Ireland

Christophe Snoeck^{1,2,3,4,*}, Carleton Jones⁵, John Pouncett¹, Steven Goderis², Philippe Claeys², Nadine Mattielli³, Antoine Zazzo⁶, Paula J. Reimer⁷, Julia A. Lee-Thorp¹, Rick J. Schulting¹

¹Research Laboratory for Archaeology and the History of Art, School of Archaeology, University of Oxford, Dyson Perrins Building, South Parks Rd, Oxford, OX1 3TG, UK

²Research Unit: Analytical, Environmental & Geo-Chemistry, Dept. of Chemistry, Vrije Universiteit Brussel, AMGC-WE-VUB, Pleinlaan 2, 1050 Brussels, Belgium

³G-Time Laboratory, Université Libre de Bruxelles, CP 160/02, 50, Avenue F.D. Roosevelt, B-1050 Brussels, Belgium

⁴Maritime Cultures Research Institute, Dept. of Art Sciences & Archaeology, Vrije Universiteit Brussel, MARI-LW-VUB, Pleinlaan 2, 1050 Brussels, Belgium

⁵School of Geography and Archaeology, Arts/Science Building, National University of Ireland Galway, University Road, Galway, Ireland

⁶Archéozoologie, Archéobotanique : sociétés, pratiques et environnements (AASPE), Muséum national d'histoire naturelle, CNRS, Sorbonne université, CP 56, 55 rue Buffon 75005 Paris, France

⁷¹⁴CHRONO Centre for Climate, the Environment, and Chronology, Queen's University of Belfast, Belfast BT7 1NN, Northern Ireland, UK

*Corresponding author – christophe.snoeck@vub.be

ABSTRACT – This paper presents the results of a study using strontium, oxygen and carbon isotopes, strontium concentrations, infrared analyses and radiocarbon dating to investigate human mobility as seen in individuals from the Neolithic court tomb of Parknabinnia, Co. Clare, Ireland. Taking advantage of the recent demonstration that it is possible to obtain reliable *in vivo* strontium isotope signals from calcined bone, we compare measurements on cremated bone (n = 4) and uncremated tooth enamel (n = 4). The results suggest that two out of four uncremated enamel samples can be considered ‘local’ while the other two, and all four cremated bone samples, represent ‘non-local’ individuals. New radiocarbon dates obtained on two of the cremated bone fragments place them in the Chalcolithic/Early Bronze Age, rather than the Neolithic ages for the uncremated remains, demonstrating re-use of the monument. Assuming that our small sample is representative, it seems that the court tomb was used for burial by both ‘locals’ and ‘non-locals’ during the Neolithic and predominantly by ‘non-locals’ in the Chalcolithic/Early Bronze Age. This stands in contrast to the nearby Early Neolithic portal tomb of Poulmabrone where only one individual (of 17 analysed) appears to be an ‘outsider’. This study suggests that, even within a small region, mobility and landscape use may have differed significantly within the Neolithic and also between the Neolithic and the Chalcolithic/Early Bronze Age.

INTRODUCTION

Funerary practices in the Irish Neolithic included the placement of both uncremated and cremated human remains in mortuary monuments, including court tombs. These monuments were constructed in the earlier Neolithic (from c. 3750 cal. BC), but use could have continued for centuries during the Neolithic and also sometimes re-used in the Chalcolithic (c. 2450–2150 BC) and the Early Bronze Age (c. 2150–1500 BC). Use and re-use involved both cremation and inhumation (Jones et al. 2015; Schulting et al. 2012). The reasons behind the choice of one funerary rite over the other are poorly understood. One possibility is that the cremated human bone represents only token deposits brought to the site from cremations that took place elsewhere, while non-cremated human remains represent the local dead. Given the long use-histories of some court tombs, might the occurrence of both practices within a single tomb reflect a diachronic shift in funerary rites?

In this paper, carbon, oxygen and strontium isotope ratios, strontium concentrations, radiocarbon dating, and infrared analyses are applied to address these questions of human mobility, funerary practice and tomb re-use at the earlier Neolithic court tomb at Parknabinnia, Co. Clare. We analysed both cremated bone and uncremated tooth enamel to investigate the possibility whether the former were part of token deposits brought to the monument from other locations. If so, we might expect their strontium isotope results to reflect a wider variety of lithologies than seen in the uncremated enamel, which could represent the deceased of the local community. Our results are compared to those previously published on uncremated enamel from the nearby Early Neolithic portal tomb at Poul nabrone, Co. Clare (Ditchfield 2014; Kador 2010; Kador et al. 2014).

The skeletal tissue of choice in C, O and Sr isotope studies is dental enamel, since this has been shown to be far more resistant to diagenesis than either dentine or bone (Tuross et al. 1989; Budd et al. 2000; Hoppe et al. 2003; Lee-Thorp & Sponheimer 2003). Recent experimental work, however, has demonstrated that calcined bone, because of its substantially greater crystallinity and thus resistance to external influences, also provides a reliable substrate for strontium isotope analyses (Harvig et al. 2014; Snoeck et al. 2015). Carbon and oxygen isotopes, however, have been heavily altered by the very high temperature reached during cremation but can still be used to look into the cremation ritual. These observations open up new possibilities not only for Irish prehistoric archaeology

(Snoeck et al. 2016a), but for that of other periods and places as well (e.g. Snoeck et al. 2018).

Strontium isotope analyses

Two isotopes of strontium, ^{86}Sr and ^{87}Sr , are widely used in mobility studies of humans and fauna. Their ratio ($^{87}\text{Sr}/^{86}\text{Sr}$) varies between different types of bedrock because of the radioactive decay of ^{87}Rb , with values ranging from about 0.7 to more than 4.0 due to different geological age, original Rb/Sr values and initial $^{87}\text{Sr}/^{86}\text{Sr}$ (Faure 1986). After being incorporated into the body through drinking water and the consumption of plants and animals, $^{87}\text{Sr}/^{86}\text{Sr}$ can be measured in bone and teeth to reflect the consumer's location at the time the tissue in question formed. Measurements on tooth enamel relate to the time during which the tooth crown formed, and so present an indication of conditions during infancy through to early adolescence, depending on the tooth measured, from c. 4 months to 2-3 years in the case of the first molar (M1) and from c. 3 to 7 years in the case of the second molar (M2), etc. Bone on the other hand continues to remodel, and so provides information relating to the later years of adult life (Hedges et al. 2007; Robin & New 1997).

To study the strontium isotope variations observed in human remains, it is necessary to construct a baseline for the biologically available strontium (BASr) around the site of interest (Bentley 2006; Evans et al. 2009; 2010). The map of the BASr for Ireland (Snoeck et al. accepted) is based on many samples coming from the Burren and surroundings and can thus be used here to contextualize the strontium isotope results obtained on the human remains. With this map, it is possible to compare the local BASr values to those measured on uncremated enamel and cremated bone and identify individuals who consumed the majority of their food close to or far from the site, following the method described in Snoeck et al. (2016a). In short, individuals having strontium isotope ratios compatible with the BASr values of catchments with a radius up to 5 km will be considered to be 'locals'. Farmers are likely to grow most of their crops and keep their animals within this range most of the time (Chisholm 1968; Jones et al. 1999). Individuals are defined as 'regional' if they exhibit a strontium isotope ratio consistent with catchments with a radius of 5–20 km and as 'outsiders' if their strontium isotope ratio is more than two standard deviations from the average BASr value measured for the 20 km radius catchment.

Carbon and oxygen isotope analyses and infrared measurements

In tooth enamel, carbon isotopes reflect overall sources of dietary carbon (Ambrose and Norr 1993), while oxygen isotopes primarily reflect drinking water. Following the expected rainfall gradient, oxygen isotopes in groundwaters are more depleted in ^{18}O in the east than in the west of Ireland (Darling et al. 2003). However, because of the high pyre temperatures reached during cremation, carbon and oxygen in bone exchange with the surrounding combustion atmosphere and in particular with carbon dioxide from the fuel. The final isotope composition represents a complex mixture of carbon and oxygen from the endogenous bone mineral and collagen fractions and that contributed by the fuel used for the cremation (e.g. Zazzo et al. 2011; 2013; Snoeck et al. 2014a; 2016b). It is therefore not possible to directly compare carbon and oxygen isotopes from cremated bone and tooth enamel. Nevertheless, in the case of uncremated enamel, it is possible to observe variations in diet and place of origin, and in the case of cremated bone, together with FTIR (Fourier Transform Infrared) spectroscopy, to assess pyre characteristics such as temperature and ventilation (Lebon et al. 2010; Snoeck et al. 2014b; 2016b). Infrared analyses allow, among other things, for the detection of cyanamide, which has been suggested to appear in bone burned under reducing conditions (Zazzo et al. 2013; Snoeck et al. 2014b; 2016b)

Radiocarbon dating of calcined bone

Twenty years ago, radiocarbon dating of calcined bone (completely white bone burned at temperatures above 650°C) provided reliable results (Lanting and Brindley 1998; Lanting et al. 2001). This has been put to good use in a wide range of archaeological contexts, including Neolithic Ireland (Schulting et al. 2012). While it is important to keep in mind that the old wood effect might impact on the radiocarbon dates of calcined bone (see Hüls et al. 2010; Zazzo et al. 2011; 2012; Snoeck et al. 2014a), reliable results can still be obtained if one can assume that the wood used to burned the deceased, and the deceased are of contemporary age.

Parknabinnia court tomb and its context

The Parknabinnia court tomb (Cl 153 – Figure 1) is located on Roughan Hill within an upland limestone region known as the Burren in north-west County Clare on the west coast of Ireland (Figure 2), an area that seems to have been a particular focus for Neolithic farmers (Jones 2003). The tomb is atypical in that it has a narrow, straight-sided ‘court’ rather than the more usual open court with curving sides, and a short heel-shaped cairn rather than a

long, trapezoidal cairn (Jones & Walsh 1996). These features, however, are shared with other court tombs in the immediate and wider region and they seem to form a morphologically distinct north Munster type (Jones 2019). Radiocarbon dates from an initial sample of twelve uncremated human bones suggest that, like many other court tombs across the country, Parknabinnia was probably initially used c. 3700–3570 BC (the earliest date is 3690–3375 cal. BC at 2 σ ; GU–10578; 4785 \pm 60 BP), but unusually, it continued to be used, if intermittently, up into the first half of the third millennium (GU–10575; 4195 \pm 55 BP; 2905–2620 cal. BC at 2 σ) (Schulting et al. 2012). As part of the current project, another two radiocarbon dates were obtained for two of the four cremated bone fragments on which isotope analyses were carried out (see below).

The skeletal remains from Parknabinnia were recovered from the tomb's two chambers, both of which were partially filled with a matrix of loose stones interspersed with disarticulated human bone (both inhumed and cremated), animal bone, potsherds, and lithic and bone artefacts. The potsherds probably represent 5–10 vessels including two Carinated Bowls, at least two simple bowls, and a decorated bowl. These vessels all probably date to the second quarter of the 4th millennium BC (Brindley 2010). The lithics recovered include leaf-shaped arrowheads, a plano-convex knife and a bifacial knife, and two flat stone beads. The bone artefacts include what appear to be two halves of the same toggle with expanded 'golf-tee' shaped heads, and a portion of what may be a long, tubular bead with an expanded head. The skeletal remains were highly fragmented, but at least twenty individuals are represented, comprising fifteen adults and five sub-adults. The stratigraphy of the site, the radiocarbon dates, and a taphonomic analysis of the skeletal collection indicate that the Neolithic burial rite at Parknabinnia consisted of successive primary inhumations that were rearranged and disturbed by subsequent inhumations (Beckett 2005a; b; 2011; Beckett & Robb 2006).




Figure 1 – Plan for Parknabinnia court tomb (CI 153); re-drawn by Libby Mulqueeny (Schulting et al. 2012)

Geology, palaeoenvironment, and archaeology of the Burren and surrounding region

In order to interpret the strontium isotope results from Parknabinnia, it is important to understand the region's geological, landscape, and human histories. The site is situated on Carboniferous limestone, while Carboniferous shale and sandstone occurs less than 5 km to the west, with older Silurian and Upper-Devonian formations to the east and south-east in the Slieve Aughty Mountains (Geological Survey of Ireland). The soils on Roughan Hill today are thin rendzinas (Finch 1971) with bare bedrock exposed in places, but they are nevertheless more developed than in many parts of the Burren. Micromorphological analysis of the sediments underlying the Parknabinnia court tomb indicated the pre-tomb presence of rendzinas and a clay-rich soil that was probably developed, in part, from glacial drift (Lewis 2003).

There is of course the possibility of soil change since the Neolithic on the Burren, which is now a karstic landscape with limited and fragmentary pockets of soil development (Drew 1982, 1983, 2001; Drew & Treacy 2014). In the past these may have been more extensive, and would have included shale and granite-bearing soils deriving from Co. Galway to the north, via glacial movement (Moles & Moles 2002). These materials would be expected to have higher $^{87}\text{Sr}/^{86}\text{Sr}$ signatures than the Carboniferous limestone. Analyses of pollen cores from the Burren and its immediate surrounds indicate changes in the vegetation that are probably related to changes in soil cover, beginning with initial clearances of the original pine and hazel forest by Neolithic farmers in the earlier 4th millennium BC (Crabtree 1982; Jelacic & O'Connell 1992; Lamb & Thompson 2005; O'Connell & Molloy 2001; Watts 1984).

190

191 The palaeoenvironmental evidence, together with the distribution of known Neolithic
192 monuments, suggests a marked preference for well-drained limestone substrates. A core area
193 of Neolithic settlement is present in the south-east Burren with other, possibly less dense,
194 occupation to the west on the Atlantic coast, to the north around the inner shores of Galway
195 Bay, and to the east across the lower-lying limestone of east Clare and west Tipperary (De
196 Valera & Ó Nualláin 1961; 1972; 1982). In contrast, only limited archaeological evidence of
197 Neolithic settlement exists on the poorly drained sandstone and shale south-west of the
198 Burren, although there is a possible court tomb (Cl 50) located in this area, where the small
199 Moy River meets the Atlantic (De Valera and Ó Nualláin 1961). While the preference for
200 siting Neolithic monuments on limestone is clear, there also appears to be a preference for
201 geological/environmental boundaries. This may reflect economies that utilized ecotonal
202 environments and can be seen in the concentration of Neolithic monuments near the southern
203 edge of the Burren where it borders sandstone and shale-derived soils and also in the siting of
204 the east Clare monuments which are within 3–6 km of the contrasting geological formations
205 of the Slieve Bernagh and Slieve Aughty mountains (De Valera and Ó Nualláin 1961).

206

207 Although not particularly abundant, evidence for the spatial distribution of Neolithic
208 settlement other than megalithic tombs does seem to be consistent with the monument
209 distributions, which show a particular focus on the southern Burren and lower levels of
210 activity elsewhere (Hull & Comber 2011; Jones 2003). This impression is reinforced by pre-
211 development work in advance of pipelines and roads passing through the region but avoiding
212 the Burren. Although there are likely sampling biases in these studies, the low number of
213 sites and features pre-dating the Chalcolithic encountered on these projects is striking
214 (Delaney et al. 2012; Grogan et al. 2007; Hull & Taylor 2010).

215

216 In the subsequent Chalcolithic and Early Bronze Age periods, areas which had previously
217 witnessed Neolithic activity seem to become particularly important activity foci (Jones et al.
218 2015), and many formerly more ‘quiet’ parts of the landscape show evidence for settlement
219 and ritual activity as well (Delaney et al. 2012; Hull & Taylor 2010). The Neolithic
220 Parknabinnia court tomb is, in fact, set in the midst of a ‘busy’ Chalcolithic/Early Bronze
221 Age landscape consisting of the densest concentration of Chalcolithic megalithic wedge
222 tombs in the country along with Chalcolithic/Early Bronze habitation enclosures and field
223 walls (Jones 2016).

MATERIALS AND METHODS

Samples

Only a small number of calcined bone fragments (c. 50 identifiable as human, with a larger number of small unidentified fragments; Beckett 2005a: 227) were encountered in the excavations, and are insufficient in weight to represent even a single individual. The assumption is, then, that these are token deposits from an unknown number of cremations carried out elsewhere. Uncremated teeth, all isolated specimens, were also limited in number. Here, four enamel (T) and four cremated bone samples (B) were selected for infrared, strontium, carbon and oxygen isotope analyses (Table 1). Two cremated bone samples (B1 and B2) were selected for radiocarbon dating.

Table 1 – Human remains from Parknabinnia

Find	Chamber	Context	Element	Notes
B1 986.01	1	443	skull frag	/
B2 1018.01		443	skull frag	/
B3 1959.02		582	skull frags	/
T1 1728.04		565	lower M1	minimal wear
T2 1908.11		582	upper M2	minimal wear; root fully formed
B4 1433.01	2	533	skull frag	/
T3 1952.01		584	incisor	lightly worn, dentine exposed
T4 2099.01		586	lower M1	cusps lightly worn, exposing dentine

Three cremated bone and two enamel samples came from Chamber 1, while the other two enamel samples and one cremated bone sample came from Chamber 2 (Figure 1). One of the cremated bone samples was analysed in triplicate (B3a–c). Based on the stratigraphic evidence and their state of attrition, T3 and T4 (slightly worn, exposing dentine) could be from the same individual, as could T1 and T2 (minimally worn, with no dentine exposure). The calcined bone fragments are assumed to belong to different individuals than those represented by the teeth, since cremation would not leave these unaffected. B1 and B2 were found in the same context and could, therefore, derive from the same individual. As discussed below, however, radiocarbon dating of these two samples shows that they represent distinct individuals, both post-dating the Neolithic. The demonstrable problems with stratigraphic integrity make it difficult to determine the number of individuals represented. At minimum, there must be four individuals: two uncremated and two cremated. We return to this below.

Analytical procedures and definition of 'local'

Details on the various analytical procedures and the geographic assignment of each individual can be found in Supplementary Information.

RESULTS

Biologically available strontium (BASr) in the Burren and surroundings

The map of the biologically available strontium for Ireland (Snoeck et al. accepted) shows that only one geological area within 75 km of Parknabinnia presents lower strontium isotope values relative to the local BASr range (0.7086 ± 0.0004 – Table 2): Carboniferous volcanics and minor intrusions to the south-east (Figure 2). However, several geological formations present higher $^{87}\text{Sr}/^{86}\text{Sr}$ values: Upper Devonian to Lower Carboniferous Old Red Sandstone of the Slieve Bernagh mountains east of Parknabinnia; the Ordovician granite formation in Connemara to the north; and the Carboniferous sandstone and shales of west Clare, etc. To evaluate their possible contribution to the strontium isotope values of the studied individuals, average BASr values for 1, 5, 10 and 20 km catchments were calculated for both Parknabinnia and Poul nabrone (Table 2).




Figure 2 – BASr map of the region around Parknabinnia and Poul nabrone (left = median and right = median absolute deviation) based on Snoeck et al. (accepted). A thin coastal strip (50m) is expected to exhibit values near 0.7092 due to the sea-spray effect.

Table 2 – BASr for the outcrop on which each site lies ('local BASr') and the average BASr values calculated for 1, 5, 10 and 20 km catchments ($\pm 2\text{SD}$); the values between brackets represent the number of different geological formations included in the calculation of the average BASr

	Site BASr	1km BASr	5km BASr	10km BASr	20km BASr
Poul nabrone	0.7086 ± 0.0004 (1)		0.7089 ± 0.0004 (2)	0.7090 ± 0.0005 (2)	0.7091 ± 0.0006 (2)
Parknabinnia			0.7089 ± 0.0004 (2)	0.7091 ± 0.0006 (2)	0.7093 ± 0.0006 (3)

Radiocarbon dating results

Two cremated bone fragments were radiocarbon dated (B1 and B2). Stratigraphically, these were incorporated into Neolithic deposits, but the results show that they are much later intrusions. B1 returned an early Chalcolithic date (UBA-31468: 2456 – 2415 cal. BC) while B2 is significantly more recent, falling within the Early Bronze Age (UBA-32467: cal. 1971 – 1687 BC) (Table 3). Both dates are several centuries younger than the latest dates on uncremated bone from the court tomb and indicate a previously unknown re-use of the tomb (Figure 3).

Table 3 – Radiocarbon dating results for the cremated bone

Find no.	Lab code	Element	¹⁴ C yr BP	cal BC (95%)
B1_986.01	UBA-31468	skull frag	3824 ± 34	2456 – 2415 BC
B2_1018.01	UBA-32467	skull frag	3502 ± 55	1971 – 1687 BC

OxCal v4.3.2 Bronk Ramsey (2017); r:5 IntCal13 atmospheric curve (Reimer et al 2013)




Figure 3 – Radiocarbon dates obtained on uncremated (Schulting et al. 2012) and cremated bone (this study) from Parknabinnia.

Infrared, elemental and isotope results

The infrared and carbon isotope results (Table 4) show that cremated bone samples B1 and B4 differ substantially from B2 and B3, having lower carbonate content (BPI – Type B Carbonate to Phosphate Index) and higher cyanamide content (CN/P). Furthermore, B1 and B4 are more enriched in ¹³C than B2 and B3. B2 also has higher $\delta^{18}\text{O}$ values compared to the other samples (Figure 4a). The variability observed in the $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values of the three

B3 samples is not altogether surprising, as this depends mainly on the composition of the combustion atmosphere during cremation, which will be highly variable (Snoeck et al. 2014a; Snoeck et al. 2016b). Even taking into account the large variability observed in B3, the $\delta^{13}\text{C}$ values of B1 and B4 are still distinguished from B2 and B3.

The tooth enamel samples have similar carbon and oxygen isotope ratios (Figure 4b). The $\delta^{13}\text{C}$ values are typical of a terrestrial C_3 -plant based diet (Zazzo et al. 2010). As would be expected, the oxygen isotope values are all enriched in ^{18}O compared to those observed for human and faunal remains from Dublin on Ireland's east coast (Knudson et al. 2012).




Figure 4 – $\delta^{13}\text{C}_{\text{VPDB}}$ vs $\delta^{18}\text{O}_{\text{VPDB}}$ of (a) the cremated bone and (b) the tooth enamel samples; note the difference in the $\delta^{13}\text{C}$ scales

Table 4 – Infrared, elemental and isotopic results for tooth enamel (T) and cremated bone (B)

	BPI	IRSF	CN/P	$^{87}\text{Sr}/^{86}\text{Sr} (\pm 2\sigma)$	[Sr] (ppm)	$\delta^{13}\text{C}_{\text{VPDB}} (\text{‰})$	$\delta^{18}\text{O}_{\text{VPDB}} (\text{‰})$
B1	0.21	4.17	0.52	0.711166 ± 10	47	-20.8	-15.3
B2	0.49	3.76	0.12	0.709881 ± 08	39	-28.4	-13.6
B3a	0.40	4.62	0.12	0.710134 ± 07	37	-25.3	-16.3
B3b	0.45	4.69	0.13	0.709818 ± 07	*	-26.1	-15.2
B3c	0.48	4.36	0.12	0.710212 ± 10	36	-28.0	-15.0
B4	0.18	4.29	1.34	0.711371 ± 08	44	-18.0	-14.9
T1	0.53	3.52	/	0.709612 ± 11	81	-16.2	-2.9
T2	0.50	3.47	/	0.710142 ± 07	98	-16.7	-3.2
T3	0.60	3.40	/	0.708883 ± 07	50	-16.7	-2.8
T4	0.56	3.37	/	0.708728 ± 08	48	-16.9	-2.2

*Not enough material remaining from B3b for strontium concentration measurement

The $^{87}\text{Sr}/^{86}\text{Sr}$ results clearly fall into three groups: one (B1 and B4 – Chamber 1 and 2 respectively) with ratios higher than 0.7112; a second (B2, B3, T1 and T2 – all Chamber 1) with intermediate ratios from 0.7096 to 0.7102; and a final group (T3 and T4 – both Chamber 2) with lower ratios of 0.7086/7 (Figure 5). As would be expected, the results obtained for the three sub-samples (B3a, B3b and B3c) of the same calcined bone fragment exhibit limited variation in $^{87}\text{Sr}/^{86}\text{Sr}$ values (0.0004). Once again, B1 and B4 differ from B2 and B3.




Figure 5 – Strontium isotope results for the uncremated enamel and calcined bone samples from Parknabinnia; the grey shaded areas correspond to the average BASr values calculated for the different catchments ($\pm 2SD$) (see Table 2)

Combining the strontium isotope ratios with strontium concentrations further highlights the differences between B1/B4 and B2/B3 as well as between T1/T2 and T3/T4. Two subsamples of B3 (a and c) returned identical concentrations. The similar isotope ratios and concentrations measured for B1/B4, B2/B3 and T3/T4 suggest that each group originated from a similar place and had a similar dietary plant/meat ratio. T1 and T2, however, present different strontium isotope ratios and concentrations. Strong positive linear correlations are observed for both the tooth enamel ($r^2 = 0.99$, $p = 0.003$) and calcined bone samples ($r^2 = 0.78$, $p = 0.116$). While the correlation for the enamel is statistically significant, both results should be taken as provisional given the small sample sizes, especially as it is difficult to be certain of the number of individuals represented by these samples.




Figure 6 – Strontium isotope ratios versus strontium concentration results for the uncremated enamel and calcined bone samples from Parknabinnia

DISCUSSION

The present project sought to compare $^{87}\text{Sr}/^{86}\text{Sr}$ values for uncremated dental enamel and cremated bone from Parknabinnia, testing the hypothesis that the former would be more likely to represent individuals from the immediate locality, while the latter might represent the token remains of individuals from further afield, since cremation would facilitate their transportation. The results suggest that the situation is more complex than this, particularly since radiocarbon dates reveal that at least some of the cremated bone post-dates the Neolithic use of the tomb. This highlights the need for caution when material from potentially comingled deposits is being analysed. Without the dates, the interpretation of the $^{87}\text{Sr}/^{86}\text{Sr}$ results would have been very different, offering some support for the hypothesised difference between the subsistence catchments represented by uncremated and cremated human remains.




Figure 7 – Geographic assignments of ‘non-local’ samples B1 (left) and B4 (right), based on the absolute values of the residuals between the sample strontium isotope ratio and the 5km focal mean for the BASr.

Two enamel samples (T3 and T4) from Chamber 2 have $^{87}\text{Sr}/^{86}\text{Sr}$ values consistent with the biologically available strontium isotope range for the immediate vicinity of the site, as well as its 1 and 5 km catchments (together defined as ‘local’). The two other enamel and two cremated bone samples (defined as ‘regional individuals’ and all from Chamber 1) have strontium isotope ratios intermediate between the local BASr value and the adjacent Carboniferous sandstone and shale BASr value located 2–3 km to the west. This falls outside

of the ‘local’ range because of the small proportion of the more radiogenic lithological formations contained within the 0-5 km catchment. The last two samples (B1 and B4) can be clearly characterised as ‘outsiders’ (see SI), having values above 0.7110, consistent with the Slieve Aughty uplands to the east and south-east (Figure 7). The strontium isotope ratios of T3 and T4 (0.7087–0.7088) suggest that the limestone $^{87}\text{Sr}/^{86}\text{Sr}$ signal of the Burren was only minimally affected by surficial glacial soils with significantly different values, and hence, at least some of the individuals represented by the calcined bone are likely to be ‘outsiders’. While it is relatively straightforward to evaluate the possible place of origin of T3 and T4 (‘locals’), and the later lives of B1 and B4 (‘outsiders’ probably from the Slieve Aughty uplands), the situation with the remaining four samples (T1, T2, B2 and B3) requires further discussion. Indeed, their $^{87}\text{Sr}/^{86}\text{Sr}$ values could be the result of consuming a combination of local resources and resources originating from the Slieve Aughty mountains (mudstone/sandstone), and/or those of the adjacent shale and sandstone region of west Clare.

The provisional linear correlation observed in the strontium concentration versus strontium isotope ratios of the tooth enamel fragments (Figure 6) suggests two main sources of strontium, with different strontium-to-calcium ratios. The individual or individuals represented by T3 and T4 clearly consumed foods locally, while T1 and T2 moved between a region with higher Sr/Ca ratios and $^{87}\text{Sr}/^{86}\text{Sr}$ values reflecting the local limestone. The difference in strontium isotope ratio and concentration for T1 and T2 suggests that these represent two distinct individuals, and that T2 either consumed more food with a higher Sr/Ca ratio than T1, or, those resources were obtained from a geological context with higher Sr/Ca ratio.

The fact that B1 and B4 differ in infrared and in $\delta^{13}\text{C}$ and $^{87}\text{Sr}/^{86}\text{Sr}$ values compared to B2 and B3 suggests that the former two derive from a different region. Incidentally, this, together with non-overlapping radiocarbon dates, also indicates that B1 and B2 are distinct individuals as both are cranial fragments in which we would expect to see similar turnover. Furthermore, the low $\delta^{13}\text{C}$ values and the presence of cyanamide in B1 and B4 could indicate that these bones were cremated in different conditions compared to B2 and B3 (Zazzo et al. 2013; Snoeck et al. 2014b), possibly on smaller and/or less well ventilated pyres, where low O_2 facilitates the incorporation of cyanamide (Snoeck et al. 2016b). Considering all of the above, at least five individuals are represented: T1, T2, T3/T4, B1/B4 and B2/B3.

Parknabinnia and Poul nabrone

While there are limited strontium isotope data available to compare with Parknabinnia, the nearby portal tomb of Poul nabrone, Co. Clare, situated on the same geological formations, provides a $^{87}\text{Sr}/^{86}\text{Sr}$ dataset on uncremated enamel (Ditchfield 2014; Kador 2010; Kador et al. 2014). The onset of burial activity begins earlier here than at Parknabinnia by one to two centuries, but the two sites overlap in use (Schulting 2014). All but one individual at Poul nabrone exhibit values between 0.7082 and 0.7092, consistent with the local BASr value (0.7086 ± 0.0004 ; $n = 17$) as well as with those calculated for 1, 5, 10 and 20 km catchments, suggesting that most individuals interred at the site originated in this part of north-west Clare. The single ‘non-local’ (‘regional’ in the present framework) at Poul nabrone has a value of 0.7102, similar to those from chamber 1 at Parknabinnia (T1, T2, B2, B3), and could have consumed food from the limestone/shale geological divide to the west or from the Slieve Aughty mountains. This individual has been directly dated to the earlier Neolithic (Ditchfield 2014: Table 4.38). The evidence for only one ‘non-local’ (6%) at Poul nabrone (Figure 8), contrasts with Parknabinnia where only T3/T4 is consistent with the local BASr.




Figure 8 – Strontium isotope results for the uncremated enamel samples from Poul nabrone (Kador 2010; Kador 2014; Ditchfield 2014); the grey shaded areas correspond to the average BASr values calculated for the different catchments ($\pm 2\text{SD}$)

The calcined remains at Parknabinnia could have been brought to the site as token deposits after cremation at a more distant location during the Chalcolithic and Early Bronze Age re-use of the tomb, which is in line with the observation that the cyanamide content and carbon isotope ratios are highly variable, and also with the small amount of material recovered. In contrast, it appears that inhumation of local and regional individuals characterized the Neolithic deposits. It is also possible that the differing local and regional $^{87}\text{Sr}/^{86}\text{Sr}$ values

obtained from the tooth samples indicate that different subsets of the local population at Parknabinnia and Poul nabrone used different areas of the landscape as their fields and pastures in the Neolithic (e.g., Bogaard et al. 2011).

Given that the strontium isotope analysis of cremated bone is relatively new, few data are available for comparison; nevertheless, cremated bone from northern Ireland dating to the Neolithic and Bronze Age also showed a majority of ‘non-local’ individuals with nine out of fifteen samples showing a ‘regional’ or ‘outsider’ signal (Snoeck et al. 2016a) which is consistent with the results obtained for Parknabinnia where all four calcined samples can be classified as ‘non-local’ (‘regional’ or ‘outsider’).

CONCLUSION

Taking advantage of the recent demonstration that cremated bone provides a reliable substrate for strontium isotope mobility studies, we investigated human mobility represented by calcined and uncremated human remains at the earlier Neolithic court tomb of Parknabinnia on the Burren of Co. Clare, western Ireland. At both Parknabinnia and Poul nabrone, uncremated tooth samples belong to local or regional individuals, while calcined bone from Parknabinnia belongs to regional individuals or individuals from other parts of Ireland or further afield. The CN/P and the $\delta^{13}\text{C}$ values obtained on the cremated bone also highlight the possibility that different pyre settings were used, suggesting that these remains may have been brought to Parknabinnia after cremation elsewhere. Both calcined bone samples that were radiocarbon dated in this study (two out of the four samples subjected to strontium isotope analysis) dated to the Chalcolithic and the Early Bronze Age, and so it is possible that all the cremated material post-dates the Neolithic use of the tomb. However, leaving the calcined bones aside, there is still a contrast with Poul nabrone in that two out of three (or four) individuals represented by the uncremated tooth samples from Parknabinnia belong to regional individuals.

Many excavated court and portal tombs have been found to contain both cremated and uncremated human remains; some, such as Creggandevosky, Co. Tyrone, held only cremated remains, though this may be because of differential survival. Further studies incorporating this material will no doubt provide new data for the investigation of mobility not only in the Irish Neolithic, but also in the Chalcolithic and Early Bronze Age (periods that saw the re-use

of both monument types and the continued use of both funerary rites), as well as new insights into the reasons underlying the two funerary rites. In this regard, the finding that the cremated remains at Parknabinnia appear to represent Chalcolithic and Early Bronze Age individuals spending most of their final years away from the site is of particular interest. The extent to which it can be extended to other sites remains to be determined.

The results of this study clearly show that, by combining a wide range of analytical methods, it is possible to better evaluate the minimum number of individuals, ascertain the re-use of a Neolithic monument in later periods, and assess landscape use and possible origins of both uncremated and cremated individuals in prehistory.

ACKNOWLEDGMENTS

This research was made possible by the financial support of the British Academy (British Academy grant SG130690 ‘Coming to Knowth’, the Quaternary Research Association, and the generous support of the Philippe Wiener-Maurice Anspach Foundation (<http://fwa.ulb.ac.be>) towards the DPhil of C.S. The VUB and VUB Strategic Research fund are thanked for their financial support for analyses; the Research Foundation – Flanders (FWO) is thanked for C.S.’s and S.G.’s postdoctoral fellowships. We thank D. Fiorillo and J. Ughetto (SSMIM) for their help with the isotope measurements at the Muséum national d’Histoire naturelle, Paris. W. Debouge and J. de Jong from the G-Time Laboratory (Géochimie: Traçage isotopique, minéral et élémentaire) of the Université Libre de Bruxelles (Belgium) are acknowledged for their help with the strontium isotope analyses by MC-ICP-MS. We are grateful to the National Museum of Ireland for permitting the analysis of the Parknabinnia material. We would like to thank the two anonymous reviewers for their time and useful comments.

BIBLIOGRAPHY

Ambrose, S.H. and Norr, L. 1993. Experimental evidence for the relationship of the carbon isotope ratios of whole diet and dietary protein to those of bone collagen and carbonate. In: J.B. Lambert and G. Grupe (eds.), *Prehistoric Human Bone: Archaeology at the Molecular Level*: pp. 1-37. New York: Springer-Verlag.

Beckett J.F. & Robb J. 2006. Neolithic burial taphonomy, ritual, and interpretation in Britain and Ireland: a review. *Social archaeology of funerary remains*. 57–80

Beckett J.F. 2005a. Death and Burial on the Burren; A Taphonomic Study of Three Megalithic Monuments in County Clare, Ireland. Unpublished PhD thesis, Department of

- 494 Archaeology, University of Cambridge.
- 495 Beckett J.F. 2005b. Selective burial in Irish megalithic tombs; burial practice, age, sex, and
 496 representation in the Neolithic. In: *Proceedings of the Fifth Annual Conference of the British*
 497 *Association for Biological Anthropology and Osteoarchaeology*. 31–39
- 498 Beckett J.F. 2011. Interactions with the dead: a taphonomic analysis of burial practices in
 499 three megalithic tombs in County Clare, Ireland. *European Journal of Archaeology*. 14(3):
 500 394–418
- 501 Bentley R.A. 2006. Kinship and mobility during the prehistoric spread of farming: isotope
 502 evidence from the skeletons. *General Anthropology*. 13(1):1–10
- 503 Bentley R.A. 2013. Mobility and the diversity of early Neolithic lives: Isotopic evidence from
 504 skeletons. *Journal of Anthropological Archaeology*. 32(3):303–312
- 505 Bentley R.A. & Knipper C. 2005. Transhumance at the early Neolithic settlement at
 506 Vaihingen (Germany). *Antiquity*. 79 (<http://antiquity.ac.uk>)
- 507 Bentley R.A., Price T.D. & Stephanc E. 2004. Determining the 'local' $^{87}\text{Sr}/^{86}\text{Sr}$ range for
 508 archaeological skeletons: a case study from Neolithic Europe. *Journal of Archaeological*
 509 *Science*. 31(4):365–375
- 510 Bentley R.A., Price T.D., Lüning J., Gronenborn D., Wahl J. & Fullagar P.D. 2002.
 511 Prehistoric migration in Europe: strontium isotope analysis of Early Neolithic skeletons.
 512 *Current Anthropology*. 43(5):799–804
- 513 Bogaard A., Krause R. & Strien H.C. 2011. Towards a social geography of cultivation and
 514 plant use in an early farming community: Vaihingen an der Enz, south-west Germany.
 515 *Antiquity*. 85(328):395–416
- 516 Brindley A. 2010. Report on the prehistoric pottery from the court tomb C153 at
 517 Parknabinnia, Co. Clare (98E0230). Unpublished report.
- 518 Budd P., Montgomery J., Barreiro B. & Thomas G.C. 2000. Differential diagenesis of
 519 strontium in archaeological human dental tissues. *Applied Geochemistry*. 15(5):687–694
- 520 Chisholm M. 1968. *Rural Settlement and Land Use*, Hutchinson & Co., London.
- 521 Crabtree K. 1982 Evidence for the Burren's Forest Cover. In M. Bell and S. Limbrey (eds.)
 522 *Archaeological Aspects of Woodland Ecology*. British Archaeological Reports International
 523 Series 146:105–113. Oxford: BAR.
- 524 Darling W.G. & Talbot J.C. 2003. The O & H stable isotopic composition of fresh waters in
 525 the British Isles: 1, Rainfall. *Hydrology and earth System Sciences*, 7(2):163–181
- 526 Delaney S., Bayley D., Lyne E., McNamara S., Nunan J. & Molloy K. (eds.) 2012.
 527 *Borderlands. Archaeological investigations along the route of the M18 Gort to Crusheen*
 528 *road scheme*. NRA Scheme Monographs 9. Dublin: The National Roads Authority.
- 529 De Valera R. & Ó Nualláin S. 1961. *Survey of the Megalithic Tombs of Ireland. Volume I*
 530 *County Clare*. Dublin: The Stationery Office.

- 531 De Valera R. & Ó Nualláin S. 1972. *Survey of the Megalithic Tombs of Ireland. Volume III*
 532 *Galway, Roscommon, Leitrim, Longford, Westmeath, Laoighis, Offaly, Kildare, Cavan.*
 533 Dublin: The Stationery Office.
- 534 De Valera R. & Ó Nualláin S. 1982. *Survey of the Megalithic Tombs of Ireland. Volume IV*
 535 *Cork, Kerry, Limerick and Tipperary.* Dublin: The Stationery Office.
- 536 Ditchfield P. 2014. Stable isotope analysis. In: Lynch A. (ed.), *Poulnabrone, Co. Clare.*
 537 *Excavation of an Early Neolithic Portal Tomb:*86–92. Dublin: Department of Arts, Heritage
 538 and the Gaeltacht, Archaeological Monograph Series.
- 539 Drew D.P. 1982. Environmental Archaeology and Karstic Terrains: The Example of the
 540 Burren, Co. Clare, Ireland. In: Bell M. & Limbrey S. (eds.), *Archaeological Aspects of*
 541 *Woodland Ecology:* 115-127. BAR International Series 146. Oxford: British Archaeological
 542 Reports.
- 543 Drew D.P. 1983. Accelerated Soil Erosion In A Karst Area: The Burren, Western Ireland.
 544 *Journal of Hydrology*, 61: 113-124.
- 545 Drew D.P. 2001. *Classic Landforms of the Burren Karst.* Sheffield: Geographical
 546 Association and the British Geomorphological Research Group.
- 547 Drew, D.P. & Treacy A. 2014. An examination of micro-solutional landforms (karren)
 548 developed on Poulnabrone portal tomb. In Lynch A., *Poulnabrone: An Early Neolithic Portal*
 549 *Tomb in Ireland:* 31-37. Archaeological Monograph Series 9. Dublin: The Stationery Office.
- 550 Evans J.A., Chenery C.A. & Fitzpatrick A.P. 2006. Bronze Age childhood migration of
 551 individuals near Stonehenge revealed by strontium and oxygen isotope tooth enamel analysis.
 552 *Archaeometry*. 48(2):309–321
- 553 Faure G. 1986. The Rb-Sr method of dating. In: *Principles of Isotope Geology, Second*
 554 *Edition:* 117–140. John Wiley and Sons, New York.
- 555 Finch T. 1971. *Soils of County Clare.* Soil Survey Bulletin 23. Dublin: An Foras Tuluntais.
- 556 Frei K.M., Frei R., Mannering U., Gleba M., Nosch M. & Lyngstrøm H. 2009. Provenance of
 557 ancient textiles – a pilot study evaluating the strontium isotope system in wool.
 558 *Archaeometry*, 51(2):252–276
- 559 Giblin J.I., Knudson K.J., Bereczki Z., Pálfi G. & Pap I. 2013. Strontium isotope analysis and
 560 human mobility during the Neolithic and Copper Age: a case study from the Great Hungarian
 561 Plain. *Journal of Archaeological Science*. 40:227–239
- 562 Grogan E., O'Donnell L., & Johnston P. 2007 *The Bronze Age landscapes of the Pipeline to*
 563 *the West.* Bray: Wordwell.
- 564 GSI – Geological Survey of Ireland, accessed 08.09.2014, <http://www.gsi.ie/>
- 565 Hess J., Bender M.L. & Schilling J.G. 1986. Evolution of the ratio of strontium-87 to
 566 strontium-86 in seawater fromm Cretaceous to present. *Science*, 231:979–985

- 567 Hoppe K.A., Koch P.L. & Furutani T.T. 2003. Assessing the preservation of biogenic
568 strontium in fossil bones and tooth enamel. *International Journal of Osteoarchaeology*.
569 13:20–28
- 570 Hull G. & Comber M. 2011. Caherconnell, Carron, Co. Clare, Final Archaeological
571 Excavation Report, Licence Nos: 08E0535 and 08R0171. Unpublished report submitted to
572 National Monuments Service.
- 573 Hull G. & Taylor K. (eds.) 2012. *Beneath the Banner - Archaeology of the M18 Ennis Bypass*
574 *and N85 Western Relief Road, Co Clare. NRA Scheme Monographs 10*. Dublin: The National
575 Roads Authority.
- 576 Hüls C.M., Erlenkeuser H., Nadeau M.-J., Grootes P.M. & Andersen N. 2010. Experimental
577 study on the origin of cremated bone apatite carbon. *Radiocarbon*. 52(2–3):587–599
- 578 Jelacic L. & O’Connell M. 1992. History of vegetation and land use from 3200 BP to the
579 present in the north-west Burren, a karstic region of western Ireland. *Vegetation History and*
580 *Archaeobotany*. 1(3):119–140
- 581 Jones C. 2003. Neolithic beginnings on Roughan Hill and the Burren. In Armit I., Murphy
582 E., Nelis E. & Simpson D. (eds.) *Neolithic Settlement in Ireland and Western Britain: pp.*
583 *188-194*. Oxford: Oxbow.
- 584 Jones C. 2007. *Temples of Stone: Exploring the Megalithic Tombs of Ireland*. Collins
585 Press. Jones, C. 2016. Dating ancient field walls in karst landscapes using differential bedrock
586 lowering. *Geoarchaeology – An International Journal* 31, 77-100.
- 587 Jones, C. 2019. The north Munster atypical court tombs of western Ireland – social dynamics,
588 regional trajectories and responses to distant events over the course of the Neolithic. In
589 Müller, J. Hinz, M., & Wunderlich, M. (eds) *Megaliths - Societies - Landscapes. Early*
590 *Monumentality and Social Differentiation in Neolithic Europe*. Proceedings of the
591 international conference in Kiel 16-20 June 2015, 983 – 1004. Bonn: Habelt Verlag.
- 592 Jones C., McVeigh T., Ó Maoldúin R. 2015. Monuments, Landscape and Identity in
593 Chalcolithic Ireland. In Springs K. (ed.) *Landscape and Identity – Archaeology and Human*
594 *Geography*. British Archaeology Reports International Series 2709, 3-25. Oxford:
595 Archaeopress.
- 596 Jones G., Bogaard A., Halstead P., Charles M. & Smith H. 1999. Identifying the intensity of
597 crop husbandry practices on the basis of weed floras. *The Annual of the British School at*
598 *Athens* 94:167–89.
- 599 Jones C. & Walsh P. 1996. Recent Discoveries on Roughan Hill, County Clare. *Journal of*
600 *the Royal Society of Antiquaries of Ireland*. 126:86–107
- 601 Kador T. 2010. Searching for Neolithic migrants at the introduction of agriculture to Ireland.
602 Unpublished report prepared for the Heritage Council, Republic of Ireland.
- 603 Kador T., Fibiger L., Cooney G., & Fullagar P. 2014. Movement and diet in early Irish
604 prehistory: first evidence from multi-isotope analysis. *Journal of Irish Archaeology* 23, 83 –
605 96.

- 606 Kador T., Cassidy L.M., Geber J., Hensey R., Meehan P. & Moore S. 2018. Rites of Passage:
607 Mortuary Practice, Population Dynamics, and Chronology at the Carrowkeel Passage Tomb
608 Complex, Co. Sligo, Ireland. *Proceedings of the Prehistoric Society*, 84:225–255
- 609 Lamb H. & Thompson A. 2005. Unusual mid-Holocene abundance of *Ulmus* in western
610 Ireland – human impact or the absence of a pathogen? *The Holocene*. 15(3):447–452
- 611 Lanting J.N, Aerts-Bijma A.T & van der Plicht J. 2001. Dating of cremated bones.
612 *Radiocarbon*. 43:249–254
- 613 Lanting J.N, Brindley A.L. 1998. Dating cremated bone: the dawn of a new era. *Journal of*
614 *Irish Archaeology*. 9:1–7.
- 615 Lebon M., Reiche I., Bahain J.-J., Chadeaux C., Moigne A.-M., Frohlich F., Sémah F.,
616 Schwarcz H.P. & Falguères C. 2010. New parameters for the characterization of diagenetic
617 alterations and heat-induced changes of fossil bone mineral using Fourier transform infrared
618 spectrometry. *Journal of Archaeological Science*, 37:2265–2276
- 619 Lee-Thorp J.A. & Sponheimer M. 2003. Three case studies used to reassess the reliability of
620 fossil bone and enamel isotope signals for paleodietary studies. *Journal of Anthropological*
621 *Archaeology*, 22(3):208–216
- 622 LeGeros R.Z. & LeGeros J.P. 1983. Carbonate analyses of synthetic, mineral and biological
623 apatites. *Journal of Dental Research*, 82:259
- 624 Lewis H. 2003 *Soil micromorphology and geoarchaeology at Parknabinnia court tomb*
625 *(Clare Megalith 153), Co. Clare, Ireland*. Unpublished report.
- 626 Madgwick R., Lamb A.L., Sloane H., Nederbragt A.J., Albarella U., Pearson M.P. & Evans J.
627 A. 2019. Multi-isotope analysis reveals that feasts in the Stonehenge environs and across
628 Wessex drew people and animals from throughout Britain. *Science advances*, 5(3):eaau6078
- 629 Moles N.R. & Moles R.T. 2002. Influence of geology, glacial processes and land use on soil
630 composition and Quaternary landscape evolution in The Burren National Park, Ireland.
631 *Catena*. 47(4):291–321
- 632 Nehlich O., Montgomery J., Evans J., Schade-Lindig S., Pichler S.L., Richards M.P. & Alt
633 K.W. 2009. Mobility or migration: a case study from the Neolithic settlement of Nieder-
634 Mörlen (Hessen, Germany). *Journal of Archaeological Science*. 36(8):1791–1799
- 635 O’Connell M. & Molloy K. 2001. Farming and Woodland Dynamics in Ireland during the
636 Neolithic. *Proceedings of the Royal Irish Academy*. 101B (1–2):99–128
- 637 Price T.D., Grupe G. & Schröter P. 1998. Migration in the Bell Beaker period of Central
638 Europe. *Antiquity*. 72:405–411
- 639 Price T.D., Wahl J. & Bentley R.A. 2006. Isotopic evidence for mobility and group
640 organization among Neolithic farmers at Talheim, Germany, 5000 BC. *European Journal of*
641 *Archaeology*. 9(2–3):259–284
- 642 Ryan S.E., Reynard L.M., Crowley C.G., Snoeck C. & Tuross N. 2018a. Early Medieval
643 Reliance on the Land and the Local: An Integrated Multi-Isotope Study ($^{87}\text{Sr}/^{86}\text{Sr}$, $\delta^{18}\text{O}$,

- 644 $\delta^{13}\text{C}$, $\delta^{15}\text{N}$) of Diet and Migration in Co. Meath, Ireland. *Journal of Archaeological Science*,
645 98:59–71
- 646 Ryan S.E., Snoeck C., Crowley Q.G. & Babechuk M.G. 2018b. High Spatial Resolution
647 Geochemical Mapping of geosphere-hydrosphere-biosphere interactions: A Case Study in
648 Ireland. *Applied Geochemistry*, 92:209–224
- 649 Schulting, R.J. 2014. The dating of Poul nabrone, Co. Clare. In: A. Lynch (ed.), *Poul nabrone:*
650 *An Early Neolithic Portal Tomb in Ireland*: pp. 93–113. Dublin: Department of Arts, Heritage
651 and the Gaeltacht, Archaeological Monograph Series no. 9.
- 652 Schulting R.J., Murphy E., Jones C. & Warren G. 2012. New dates from the north, and a proposed
653 chronology for Irish court tombs. *Proceedings of the Royal Irish Academy*. 112C:1–60
- 654 Sheridan A., Jay M., Montgomery J., Pellegrini M. and Wilson J.C. 2013. ‘Tara Boy’: local
655 hero or international man of mystery? In: M. O’Sullivan, C. Scarre and M. Doyle (eds.), *Tara*
656 *- From the Past to the Future. Towards a New Research Agenda*: pp. 207–232. Dublin:
657 Wordwell.
- 658 Sjögren K.-G. & Price T.D. 2013. A complex Neolithic economy: isotope evidence for the
659 circulation of cattle and sheep in the TRB of western Sweden. *Journal of Archaeological*
660 *Science*. 40:690–704
- 661 Sjögren K.-G., Price T.D. & Ahlström T. 2009. Megaliths and mobility in south-western
662 Sweden. Investigating relationships between a local society and its neighbours using
663 strontium isotopes. *Journal of Anthropological Archaeology*. 28(1):85–101
- 664 Snoeck C. 2014. Impact of strontium sea spray effect on the isotopic ratio ($^{87}\text{Sr}/^{86}\text{Sr}$) of
665 plants in coastal Ireland. *Quaternary Newsletter*, 134:37–39
- 666 Snoeck C., Brock F. & Schulting R.J. 2014a. Carbon exchanges between bone apatite and
667 fuels during cremation: Impact on radiocarbon dates. *Radiocarbon*, 56(2):591–602
- 668 Snoeck C., Lee-Thorp J.A. & Schulting R.J. 2014b. From bone to ash: Compositional and
669 structural studies of cremated bone. *Palaeogeography, Palaeoclimatology, Palaeoecology*,
670 416:55–68
- 671 Snoeck C., Lee-Thorp J.A., Schulting R.J., de Jong J., Debouge W. & Mattielli N. 2015.
672 Calcined bone provides a reliable substrate for strontium isotope ratios as shown by an
673 enrichment experiment. *Rapid Communication in Mass Spectrometry*, 29:107–114
- 674 Snoeck C., Pouncett J., Ramsey G., Meighan I., Mattielli N., Goderis S., Lee-Thorp J.A. &
675 Schulting R.J. 2016a. Mobility during the Neolithic and Bronze Age in Northern Ireland
676 explored using strontium isotope analysis of cremated human bone. *American Journal of*
677 *Physical Anthropology*, 160(3):397–413
- 678 Snoeck C., Ryan S., Pouncett J., Pellegrini M., Claeys Ph., Wainwright A.N., Mattielli N.,
679 Lee-Thorp J.A. & Schulting R.J. *Accepted*. Towards a biologically available strontium
680 isotope baseline for Ireland. *Science of The Total Environment*.

- 681 Snoeck C., Schulting R.J., Lee-Thorp J.A., Zazzo A. & Lebon M. 2016b. Impact of heating
682 conditions on the carbon and oxygen isotope composition of calcined bone. *Journal of*
683 *Archaeological Science*, 65:32–43
- 684 Snoeck C., Pouncett J., Claeys Ph., Goderis S., Mattielli N., Parker Pearson M., Willis C.,
685 Zazzo A., Lee-Thorp J.A. & Schulting R.J. 2018. Strontium isotope analysis on cremated
686 human remains from Stonehenge support links with west Wales. *Scientific Reports*, 8:10790
- 687 Sponheimer M. & Lee-Thorp J.A. 1999. Alteration of enamel carbonate environments during
688 fossilization. *Journal of Archaeological Science*, 26:143–150
- 689 Tuross N., Behrensmeyer A.K. & Eanes E.D. 1989. Strontium increases and crystallinity
690 changes in taphonomic and archaeological bone. *Journal of Archaeological Science*,
691 16(6):661–672
- 692 Viner S., Evans J., Albarella U. & Parker Pearson M. 2010. Cattle mobility in prehistoric
693 Britain: strontium isotope analysis of cattle teeth from Durrington Walls (Wiltshire, Britain).
694 *Journal of Archaeological Science*. 37(11):2812–2820
- 695 Watts W.A. 1984. The Holocene vegetation of the Burren, western Ireland. In: E. Haworth
696 and J. Lund (eds.) *Lake Sediments and Environmental History, Studies in palaeolimnology*
697 *and paleoecology in honor of Winfred Tutin*, 359–376. Minneapolis: University of Minnesota
698 press.
- 699 Weiner S. & Bar-Yosef O. 1990. States of Preservation of Bones from Prehistoric Sites in the
700 Near East: A Survey. *Journal of Archaeological Science*, 17:187–196
- 701 Whipkey C.E., Capo R.C., Chadwick O.A. & Stewart B.W. 2000. The importance of sea
702 spray to the cation budget of a coastal Hawaiian soil: a strontium isotope approach. *Chemical*
703 *Geology*, 168(1):37–48
- 704 Wilson J.C., Osborne H., Taylor C., Ditchfield P. & Pike A.W.G. 2012. Strontium and
705 oxygen isotope analysis on iron age and early historic burials around the Great Mound at
706 Knowth, Co. Meath. In: G. Eogan (ed.) *Excavations at Knowth 5: The Archaeology of*
707 *Knowth in the First and Second Millennia AD*: pp. 775–788. Dublin: Royal Irish Academy.
- 708 Zazzo A., Balasse M., Passey B.H., Moloney A.P., Monahan F.J. & Schmidt O. 2010. The
709 isotope record of short-and long-term dietary changes in sheep tooth enamel: implications for
710 quantitative reconstruction of paleodiets. *Geochimica et Cosmochimica Acta*, 74(12):3571–
711 3586
- 712 Zazzo A., Lebon M., Chiotti L., Comby C., Delqué-Količ E., Nespoulet R. & Reiche I. 2013.
713 Can we use calcined bone for ^{14}C dating the Paleolithic? *Radiocarbon*, 55(2–3):1409–1421
- 714 Zazzo A. & Saliège J.-F. 2011. Radiocarbon dating of biological apatites: a review.
715 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 310:52–61
- 716 Zazzo A., Saliège J.-F., Lebon M., Lepetz S., Moreau C. 2012. Radiocarbon dating of calcined bones:
717 insights from combustion experiments under natural conditions. *Radiocarbon*. 54(3–4):1–12.