

Structure-based development of new RAS-effector inhibitors from a combination of active and inactive RAS-binding compounds

Abimael Cruz-Migoni^{a,b,1}, Peter Canning^{a,1,2}, Camilo E. Quevedo^{a,1}, Carole J. R. Bataille^c, Nicolas Bery^a, Ami Miller^a, Angela J. Russell^c, Simon E. V. Phillips^{b,d}, Stephen B. Carr^{b,d}, and Terence H. Rabbitts^{a,3}

^aWeatherall Institute of Molecular Medicine, MRC Molecular Haematology Unit, John Radcliffe Hospital, University of Oxford, OX3 9DS Oxford, United Kingdom; ^bResearch Complex at Harwell, Rutherford Appleton Laboratory, OX11 0FA Didcot, United Kingdom; ^cChemistry Research Laboratory, University of Oxford, OX1 3TA Oxford, United Kingdom; and ^dDepartment of Biochemistry, University of Oxford, OX1 3QU Oxford, United Kingdom

Edited by James A. Wells, University of California, San Francisco, CA, and approved December 17, 2018 (received for review July 6, 2018)

The *RAS* gene family is frequently mutated in human cancers, and the quest for compounds that bind to mutant *RAS* remains a major goal, as it also does for inhibitors of protein–protein interactions. We have refined crystallization conditions for KRAS₁₆₉^{Q61H}-yielding crystals suitable for soaking with compounds and exploited this to assess new *RAS*-binding compounds selected by screening a protein–protein interaction-focused compound library using surface plasmon resonance. Two compounds, referred to as PPIN-1 and PPIN-2, with related structures from 30 initial *RAS* binders showed binding to a pocket where compounds had been previously developed, including *RAS* effector protein–protein interaction inhibitors selected using an intracellular antibody fragment (called Abd compounds). Unlike the Abd series of *RAS* binders, PPIN-1 and PPIN-2 compounds were not competed by the inhibitory anti-*RAS* intracellular antibody fragment and did not show any *RAS*-effector inhibition properties. By fusing the common, anchoring part from the two new compounds with the inhibitory substituents of the Abd series, we have created a set of compounds that inhibit *RAS*-effector interactions with increased potency. These fused compounds add to the growing catalog of *RAS* protein–protein inhibitors and show that building a chemical series by crossing over two chemical series is a strategy to create *RAS*-binding small molecules.

cancer | *RAS* | drugs | antibody | intracellular antibody

The oncogenic family of *RAS* genes is of significant interest in the fight against cancer because of the frequency of activating mutations (1). Their presence in almost all major cancers makes them a highly valued therapeutic target, in particular the *KRAS* gene, since it has been identified as one of the most frequently mutated oncogenes (2, 3). *RAS* proteins are linked to the plasma membrane by COOH-terminal prenylation mediated by farnesyl transferases (4). All family members function by signal transduction to the nucleus of cells via interaction with effectors (such as RAF, RALGDS, and PI3K) that catalyze phosphorylation of downstream proteins (5). When *KRAS* is bound to GDP, the protein is in the inactive state and becomes activated by nucleotide exchange from GDP to GTP. Normally, the activation/deactivation cycle is catalyzed by guanine nucleotide exchange factors and GTPase-activating proteins (GAPs) (6, 7). Mutant *RAS* proteins remain in the active state and hydrolyze GTP at a much slower rate than wild-type (WT) *RAS* (8). Mutations reduce GAP activity leading to constitutive activation of *RAS* effector pathways (2), constantly generating a signaling cascade that activates cell functions such as division, survival, and invasion (9).

Despite its great potential as a cancer target, *KRAS* has proved to be very difficult to inhibit in a therapeutic setting. *KRAS* signaling works via protein–protein interactions (PPI) that can be very difficult to disrupt (10). In addition, the nucleotides that regulate *KRAS* function (GTP and GDP) bind to the protein with picomolar affinity, making them problematic to displace (11). Attempts at targeting *RAS* function using farnesyl

transferase inhibitors also proved to be ineffective, failing to demonstrate antitumor activity in *KRAS*-driven cancers (12). As an alternative to compounds, various macromolecules [called macrodrugs (13)] have been developed that can bind to *RAS* and prevent PPI with the *RAS* effectors, such as has been shown with intracellular antibody fragments (14, 15). The possible clinical use of these macrodrugs has not been implemented thus far due to difficulties in their delivery into cells, although methods are becoming available that may solve this problem (16).

Although there are a large number of mutant *RAS* protein isoforms, their structural conformation is highly conserved (17) because of the invariant N-terminal domain up to amino acid 166. The interest in inhibition of *RAS* proteins by small molecules has increased again recently (18), and several compounds have been described that bind to *RAS* (19–27). Recently, we have defined a chemical series based on an intracellular antibody-binding domain (28) that interact with a hydrophobic pocket (designated pocket I, *SI Appendix*, Fig. S14), previously identified in silico (29) and confirmed as the binding site for 4,6-dichloro-2-methyl-3-aminoethyl-indole (DCAI) near the switch I region of *KRAS* (23).

Significance

The *RAS* family of oncogenic proteins is important as therapy targets because of the frequency of activating mutations in almost all major cancers. An important approach is development of small molecules with drug-like properties that can inhibit *RAS*-effector protein interactions inside cells. We present a strategy for identification of such compounds, and their development as *RAS*-effector interaction inhibitors, utilizing a structure-based design approach and cell-based assays. By combining moieties from two distinct sets of *RAS*-binding molecules, we generated cross-over compounds that showed improved efficacy in vitro and in cell-based assays.

Author contributions: C.E.Q. and T.H.R. designed research; A.C.-M., P.C., C.J.R.B., N.B., and A.M. performed research; A.C.-M., P.C., C.E.Q., C.J.R.B., N.B., A.M., A.J.R., S.E.V.P., S.B.C., and T.H.R. analyzed data; and A.C.-M., P.C., C.E.Q., C.J.R.B., N.B., A.M., S.E.V.P., S.B.C., and T.H.R. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under [Creative Commons Attribution License 4.0 \(CC BY\)](#).

Data deposition: The atomic coordinates have been deposited in the Protein Data Bank, www.pdb.org (PDB ID codes 6GOD, 6GOE, 6GOF, 6GOG, 6GOM, 6GQT, 6GQW, 6GQX, and 6GQY).

¹A.C.-M., P.C., and C.E.Q. contributed equally to this work.

²Present address: Assay Development, LifeArc, Open Innovation Campus, SG1 2FX Stevenage, United Kingdom.

³To whom correspondence should be addressed. Email: terence.rabbitts@imm.ox.ac.uk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1811360116/-DCSupplemental.

Published online January 25, 2019.




Fig. 2. Crystal structure of PPIN-1 and PPIN-2 bound to KRAS₁₆₉^{Q61H}-GppNHp. The crystal structure of KRAS₁₆₉^{Q61H} with PPIN-1 and PPIN-2 was derived by crystal soaking with the compounds (their structures are shown in A and B, respectively). (C and D) Surface representations of the binding of PPIN-1 and PPIN-2 into pocket I close to the switch regions I (red) and II (blue). Good 2mF_o-DFc electron density (green mesh) was found for the whole of PPIN-1 and for the biphenyl head group of PPIN-2 (green mesh) but less contiguous for the rest of the molecule. (E and F) Expanded views of the interactions of PPIN-1 and PPIN-2 with KRAS with the following residues in contact: K5, L6, V7, S39, Y40, R41, D54, I55, L56, Y71, and T74.

to assess intracellular RAS-protein interactions. As predicted from the waterLOGSY data using the intracellular antibody fragment, the PPIN compounds did not disrupt the interaction of KRAS₁₆₆^{G12D} with the anti-RAS iDAb (VHY6) with a dematured version of the iDAb (VHY6dm) or with full-length CRAF (CRAF^{FL}) (*SI Appendix, Fig. S4D*). This discrepancy between *in vitro* affinity and *in cell* potency could be attributed to targets with high conformational variability (like RAS switch regions) interacting with allosteric binders (32, 33).

Crystal Soaking of KRAS₁₆₉Q61H with PPIN-1 and -2. We determined the RAS-binding sites of PPIN-1 and PPIN-2 by soaking KRAS₁₆₉^{Q61H} crystals. Crystals of KRAS soaked with PPIN-1 diffracted to 1.63 Å (Fig. 2 C and E and *SI Appendix, Table S3*). Good electron density was observed for the ligand in one chain of the asymmetric unit (chain A, in which the switch regions are stabilized due to additional interactions on the opposite face with residues Arg102 and Lys101 from chain B in a neighboring asymmetric unit) and located adjacent to the C-terminal end of the switch I and switch II regions in pocket I. PPIN-1 primarily contacts KRAS₁₆₉^{Q61H} with the biphenyl head group via van der Waals interactions. No hydrogen bonds are formed with the protein. Crystals soaked with PPIN-2 diffracted to 1.7 Å. For this compound, clear electron density corresponding to the methoxy-biphenyl anchor group was observed in four of the six chains (Fig. 2 D and F). Weaker electron density was observed around the linker and tail groups, suggesting that the rest of the compound remains flexible when bound to the protein. It was noteworthy that both PPIN compounds have the same biphenyl anchoring group

but different tail functional groups, suggesting that the biphenyl-type groups of PPIN-1 and -2 are key in targeting these compounds to the pocket I-binding site.

Design and Characterization of RAS-Binding Cross-Over Compounds.

Our data show that PPIN-1 and PPIN-2 bind to KRAS at the same pocket I as several previously identified compounds (23), but they do not disrupt RAS function. To understand the lack of RAS inhibition of the PPIN compounds, we selected one of our RAS-binding intracellular antibody-derived compounds (Abd-7) able to interfere with RAS PPI in cells (28). We used the computational chemistry suite FORGE (<https://www.cresset-group.com/forge/>; ref. 34) that employs a ligand comparison method to align and score molecules independently using their shape and electrostatic properties, aiding the understanding of structure-activity relationships. FORGE was used to compare the structures of PPIN-1 and -2 with Abd-7 by performing an alignment based on the surfaces of the compounds only and detected similarities in the lower half of the molecules (Fig. 3 A and B). The aromatic ring of the benzodioxane moiety in Abd-7 aligned with the terminal biphenyl aryl ring in both PPIN compounds, and the pyridine ring of Abd-7 aligned with the middle aromatic ring of both PPIN compounds. This suggested that the PPIN biphenyl system might act as the anchor to the RAS protein. The alignment with the functional groups on the upper half of the molecules was poor. We used these analyses to design of cross-over compounds linking the common diphenyl-anchoring moiety of PPIN-1 and PPIN-2 to the Abd-7 aniline fragment.




Fig. 3. Abd and PPI-net compound alignment and cross-over compound crystallography. Alignments were carried out with the computational chemistry suite FORGE. (A) Abd-7 and PPIN-1 with alignments. (B) Abd-7 and PPIN-2. Three cross-over compounds were synthesized after the alignments, which are shown in C (Left, Ch-1; Middle, Ch-2; Right Ch-3). These compounds were soaked into KRAS₁₆₉^{Q61H}-GppNHp crystals. (D) A surface representation of the binding of Ch-1 (Left), Ch-2 (Middle), and Ch-3 (Right) in KRAS pocket I, close to the switch regions I (red) and II (blue). Full electron density (2Fo-Fc) was found for the three compounds, all depicted as a green mesh. (E) An expanded view of the interaction of the compounds with KRAS with the following residues in contact: K5, L6, V7, S39, R41, D54, I55, L56, Y71, and T74.

Three cross-over compounds (Fig. 3C: Ch-1, Ch-2, and Ch-3) were synthesized and their binding geometries were determined by X-ray crystallography using KRAS₁₆₉^{Q61H} crystal soaking (Fig. 3D and E and *SI Appendix, Table S4*). All three compounds showed very similar binding modes to Abd-7 with van der Waals contacts to K5, L6, V7, S39, Y40, R41, D54, I55, L56, G70, Y71, T74, and G75. Ch-1 was found in four of the six KRAS₁₆₉^{Q61H} chains (A, B, C, and F); Ch-2 was again found in three of the six chains (A, B, and C) and Ch-3 was found in four of the six chains (A, B, C, and D). Thus, by combining the anchor constituent of the PPIN compounds with the aniline fragment of Abd-7, we generated a compound series showing good electron density for the entire molecule when bound to KRAS pocket 1.

The RAS-Binding Cross-Over Compounds Are PPI Inhibitors in Cells. The purpose of the compounds was to generate inhibitors of RAS PPI. The ability of the compounds to interfere with RAS-associated PPI was analyzed with our BRET-based RAS biosensor toolbox (31). The interaction of full-length KRAS^{G12D} and either the anti-RAS iDAb VHY6, a dematured version of this iDAb (VHY6dm), or the natural RAS partner CRAF^{FL} was assessed as a dose-response with the three compounds (respectively, Fig. 4A–C). While

the interaction of the high-affinity WT iDAb with RAS was minimally affected, even at the highest dose of compound (i.e., 20 μ M), both the lower-affinity-dematured iDAb and CRAF binding to RAS were progressively impaired, starting at the lowest dose of 5 μ M. No alteration in the BRET signal for the PPI of a non-relevant protein pair (LMO2-VH576dm) was observed (Fig. 4D), confirming that the dose-response effects of the Ch compounds was not due to loss of cell viability.

We also tested the effect of compound Ch-3 in the BRET assay using five different full-length KRAS^{G12} mutations interacting with full-length CRAF (Fig. 4E: G12A, G12C, G12R, and G12V) or CRAF RAS-binding domain (RBD) (Fig. 4G: G12D). Each of these PPIs was inhibited by the Ch-3 compound in a dose-response assay. Furthermore, the BRET interaction signal between KRAS^{G12D} and either PI3K (α or γ) or RALGDS was inhibited by Ch-3 (Fig. 4G), demonstrating that the BRET data are not restricted to KRAS-CRAF interaction. Finally, we show that Ch-3 interferes with NRAS and HRAS isoforms using the BRET biosensor assay. Interactions between full-length NRAS^{Q61H} or HRAS^{G12V} and full-length CRAF (Fig. 4F), between NRAS^{Q61H} and RBD for PI3K (α or γ), CRAF, and RALGDS (Fig. 4H), or between HRAS^{G12V} and RBD for PI3K (α or γ), CRAF, and RALGDS (Fig. 4I) are inhibited in the BRET assay by Ch-3.

These BRET data show that the three cross-over compounds can enter cells and reach their target protein (RAS) in the cytoplasmic environment. The BRET assay relies on cotransfected donor and acceptor expression plasmids, and we confirmed this inhibitory capability by testing the effect of the compounds on biomarker phosphorylation in DLD-1 colorectal cancer cells. The cells were incubated with our previously described antibody-derived compounds Abd-2 and Abd-7, the original PPIN-2 compound, or the three cross-over compounds Ch-1, -2, and -3. Phosphorylation of AKT (downstream of RAS-PI3K signaling) or phosphorylation of ERK (downstream of RAS-RAF signaling) was determined following EGF stimulation (Fig. 5A–C). The PPIN-2 (and control Abd-2) had no effect on the levels of phospho-AKT or phospho-ERK even at 20 μ M (addressed using Western blotting, Fig. 5A–C), but we found that the three cross-over compounds caused loss of phospho-AKT or phospho-ERK, as did the previously described compound Abd-7. The most potent compound appears to be Ch-3, which invokes an almost complete reduction of phospho-AKT at a concentration of 10 μ M while not affecting AKT protein levels (Fig. 5C).

The biomarker Western blotting assay was carried out 2.5 h after addition of the compounds, at which time no loss of viability was observed. The survival of DLD-1 cells was determined over 48- and 72-h periods using a dose-response (0–20 μ M) (Fig. 5D; data shown at 72 h), allowing a calculation of IC₅₀ for each compound (*SI Appendix, Table S5*). The previously characterized low-affinity Abd-2 compound does not affect DLD-1 viability over the range of concentrations nor does the PPIN-2 compound. Conversely, the intracellular antibody-derived compound Abd-7 causes loss of viability with IC₅₀ of 10.8 μ M at 48 h and 8.2 μ M at 72 h. The potency of the two of the cross-over compounds (Ch-1 and Ch-3) is improved, relative to Abd-7, as these show IC₅₀ at 72 h of 5.3 and 4.5 μ M, respectively (*SI Appendix, Table S5*). This increased efficacy of Ch-3 in the challenge of DLD-1 viability matches the most efficacious compound in the signaling biomarker assay.

Discussion

The RAS family of genes is among the most frequently mutated in human cancer (e.g., up to 96% in pancreatic cancer) and therefore an important target for drug development. Targeting the RAS-effector PPI is one possible route to RAS inhibitors. Screening chemical compound libraries per se does not guarantee selection of compounds that will act as PPI inhibitors unless there is a method to guide the screen to functionally specific locations. Strategies are needed that will allow protein-binding compounds




Fig. 4. Compound Ch-3 disrupts RAS-effector interactions. Assessment of the inhibition of RAS protein–protein interactions in cells by the chemical series compounds Ch-1, Ch-2, and Ch-3 using different BRET-based RAS biosensor expression vectors. (A–C) Data from BRET assays using RLuc8-KRAS^{G12D} with either anti-RAS VHY6-GFP² (A) with dematured anti-RAS VHY6dm-GFP² (B) or with full-length CRAF^{FL}-GFP² (C). (D) Data from BRET assay using a negative control BRET-based biosensor LMO2/VH576dm. The VH576dm is a dematured anti-LMO2 VH. The data are computed relative to cells treated with DMSO vehicle only (open bar) or with Ch-1, Ch-2, or Ch-3 (shaded bars). (E and F) The effect of the Ch-3 compound on mutant KRAS^{G12X} (E) and NRAS^{Q61H} and HRAS^{G12V} (F) interactions with CRAF^{FL}. (G–I) The effect of Ch-3 on the interaction of KRAS^{G12D} (G), NRAS^{Q61H} (H), and HRAS^{G12V} (I) with various RAS effector domains (PI3K α , PI3K γ , CRAF, and RALGDS). The range of concentration of the compounds was 5, 10, and 20 μ M. Each experiment was repeated at least twice (biological replicates). Statistical analyses were performed using a one-way ANOVA followed by Dunnett's posttests (* P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001). Error bars correspond to mean values \pm SD of biological repeats. RLuc8-KRAS, NRAS, and HRAS all comprised full-length RAS components.




Fig. 5. Activity of compounds in a mutant KRAS human cancer cell. The new chemical series compounds Ch-1, -2, and -3 were assessed in two cell-based assays. (A–C) Western blot analysis of EGF-stimulated DLD-1 cells treated with 5, 10, and 20 μM of Abd-2 or Abd-7 (A), PPIN-2 or Ch-1 (B), and Ch-2 or Ch-3 (C). Cell extracts were fractionated by SDS/PAGE and transferred to PVDF membranes that were incubated with antibodies detecting the indicated proteins. These data are quantitated in *SI Appendix, Fig. S5*. (D) DLD-1 cell viability 72 h after treatment with a single application of compound at the indicated concentrations. Viability was determined using the CellTiterGlo method and carried out in triplicate. The data are plotted as normalized cell viability mean with error bars showing SDs.

to be selected and improved in their properties related to the functional mechanism to be disrupted. In particular, X-ray crystallography of RAS proteins is a key method to determine the location and geometry of bound compounds. We optimized production of KRAS₁₆₉^{G61H} crystals for soaking of compounds for this purpose. It should be noted that these conditions, in principle, could be used to produce crystals of other RAS mutants suitable for compound-soaking experiments. Surface analysis of KRAS (using the CASTp server: sts.bioe.uic.edu/castp/) confirmed three pockets with internal volumes greater than 80 Å³ (*SI Appendix, Fig. S14*: pockets I, III, and IV) and a less deep, more like a shallow groove, pocket (*SI Appendix, Fig. S14*: pocket II). Pocket I was previously identified in silico (29) and was also the site where DCAI was first selected (23). In addition, this pocket is close to the switch region and could act as a point of inhibition for PPI. Furthermore, we have focused on the identification and optimization of compounds inhibiting RAS-effector interactions and not other RAS modulation effects, such as DCAI has shown. Structural comparison of the six chains in the asymmetric unit show that compound binding has a minimal effect on the conformation of the switch regions in RAS protein. Any observed differences are no more than twice the coordinate error (35) and therefore cannot be considered significant. Comparison of the temperature factors between RAS molecules with compound bound and 3GFT (where no compound is present) show no evidence that compound

binding results in significant reduction in flexibility of switch II relative to the rest of the molecule, suggesting that any stabilization of the switch regions by compound binding is, at most, slight. Therefore, we conclude that any inhibitory effect observed is due to a disruption of RAS-effector PPI rather than any other modulation of the RAS protein.

We have screened a chemical library that yielded two RAS-binding compounds (the PPINs) and have shown, by crystallography, that they bind to pocket I near to the effector binding sites. However, when these were tested in an orthogonal NMR waterLOGSY binding assay, their binding to KRAS₁₆₆^{G12V} was not impaired by the presence of the anti-RAS intracellular antibody fragment, nor did they interfere with PPI using a cell-based BRET assay (*SI Appendix, Fig. S4*). Crystallography and medicinal chemistry was undertaken to improve the initial PPIN hits using structure-based design combining the crystal information and molecular fragments from two different chemical series, namely PPIN and the Abd series (28). The resulting crossover compounds illustrate that this strategy guided the conversion of the PPIN RAS-binding compounds to RAS PPI inhibitors in a series (compounds designated Ch-1–3). These compounds have a low molecular weight suitable for further medicinal chemistry to improve drug-like properties and with better ligand efficiencies than their progenitors (*SI Appendix, Table S5*). Furthermore, the compounds have also shown better cell viability results than their progenitors. Their low molecular weight makes them a better starting point for the development of RAS inhibitors based on this promising chemical series.

We have previously shown that the Abd chemical series affected RAF, RAL, and PI3K interactions with RAS (28) and, due to the similarities in binding mode and orientation of the Ch series with the Abd series, we expect the Ch compounds to have a similar range of profiles with other KRAS mutants and also with NRAS and HRAS isoforms. This was confirmed using various KRAS^{G12} mutants and the NRAS^{Q61H} and HRAS^{G12V} mutants with four effector molecules in BRET assays.

This approach shows that compounds binding in pocket I are not necessarily able to inhibit RAS PPI, but synthetically linking components of two classes of RAS-binding compounds can generate new active molecules that inhibit PPI. It should also be noted that the presence of pocket I in nonmutated forms of RAS as well as in mutant RAS (*SI Appendix, Fig. S1 B–G*) presents a technical challenge for the development of anti-RAS drugs since these will bind to the pocket in WT-activated RAS as well as in mutant RAS. Development of methodologies for specific drug delivery could avoid drug interference with non-mutated RAS. Approaches such as Antibody-Drug Conjugates (ADC) offer one route to avoid drugs entering normal cells (reviewed in ref. 36) by targeting antigens expressed on tumors. While this, in turn, has difficulties, since few surface antigens are tumor-specific, surfaceome studies of tumors (37, 38) can find possible markers or pairs of markers that may be useful for mono- or bispecific ADCs. An alternative approach, based on structure-activity relationships, could be the development of compounds anchoring at pocket I and moving toward the nucleotide-binding region of RAS or perhaps linking the “unselective” but potent compounds identified in pocket I to those binding in pocket II.

In conclusion, our approach demonstrates the importance in drug development of combining assays for PPI with the identification of compounds that bind at important locations but not necessarily with PPI properties. Thus, compounds binding with good potency to the target but not showing any effect on PPI could be utilized in combination chemistry to create new chemical series. A combination of high-resolution crystallography from different chemical series with biophysical competition assays, such as using high-affinity antibody fragments, is thus a powerful way to identify hit compounds of interest in analogous

settings. It is also useful in the development of new chemical series, even when initial compounds are inactive, and should allow directed medicinal chemistry for drug development.

Methods

Detailed methods on protein expression and purification, SPR screening, NMR analysis, crystallography experiments, cell-based assays, and chemical experiments can be found in *SI Appendix, Methods*. The atomic coordinates have been deposited in the Protein Data Bank (39–47).

- Cox AD, Fesik SW, Kimmelman AC, Luo J, Der CJ (2014) Drugging the undruggable RAS: Mission possible? *Nat Rev Drug Discov* 13:828–851.
- Prior IA, Lewis PD, Mattos C (2012) A comprehensive survey of Ras mutations in cancer. *Cancer Res* 72:2457–2467.
- Simanshu DK, Nissley DV, McCormick F (2017) RAS proteins and their regulators in human disease. *Cell* 170:17–33.
- Gutierrez L, Magee AJ, Marshall CJ, Hancock JF (1989) Post-translational processing of p21ras is two-step and involves carboxyl-methylation and carboxy-terminal proteolysis. *EMBO J* 8:1093–1098.
- Wittinghofer A, Nassar N (1996) How Ras-related proteins talk to their effectors. *Trends Biochem Sci* 21:488–491.
- Bos JL, Rehmann H, Wittinghofer A (2007) GEFs and GAPs: Critical elements in the control of small G proteins. *Cell* 129:865–877.
- Cheffils J, Zeghouf M (2013) Regulation of small GTPases by GEFs, GAPs, and GDIs. *Physiol Rev* 93:269–309.
- Hunter JC, et al. (2015) Biochemical and structural analysis of common cancer-associated KRAS mutations. *Mol Cancer Res* 13:1325–1335.
- Wood KW, Sarnecki C, Roberts TM, Blenis J (1992) Ras mediates nerve growth factor receptor modulation of three signal-transducing protein kinases: MAP kinase, Raf-1, and RSK. *Cell* 68:1041–1050.
- Scott DE, Bayly AR, Abell C, Skidmore J (2016) Small molecules, big targets: Drug discovery faces the protein-protein interaction challenge. *Nat Rev Drug Discov* 15: 533–550.
- Spiegel J, Cromm PM, Zimmermann G, Grossmann TN, Waldmann H (2014) Small-molecule modulation of Ras signaling. *Nat Chem Biol* 10:613–622.
- Whyte DB, et al. (1997) K- and N-Ras are geranylgeranylated in cells treated with farnesyl protein transferase inhibitors. *J Biol Chem* 272:14459–14464.
- Tanaka T, Rabbitts TH (2008) Functional intracellular antibody fragments do not require invariant intra-domain disulfide bonds. *J Mol Biol* 376:749–757.
- Tanaka T, Rabbitts TH (2012) Intracellular antibody capture (IAC) methods for single domain antibodies. *Methods Mol Biol* 911:151–173.
- Cochet O, et al. (1998) Intracellular expression of an antibody fragment-neutralizing p21 ras promotes tumor regression. *Cancer Res* 58:1170–1176.
- Dewhirst MW, Secomb TW (2017) Transport of drugs from blood vessels to tumour tissue. *Nat Rev Cancer* 17:738–750.
- Ostrem JM, Shokat KM (2016) Direct small-molecule inhibitors of KRAS: From structural insights to mechanism-based design. *Nat Rev Drug Discov* 15:771–785.
- Ledford H (2015) Cancer: The Ras renaissance. *Nature* 520:278–280.
- Burns MC, et al. (2014) Approach for targeting Ras with small molecules that activate SOS-mediated nucleotide exchange. *Proc Natl Acad Sci USA* 111:3401–3406.
- Lito P, Solomon M, Li LS, Hansen R, Rosen N (2016) Allele-specific inhibitors inactivate mutant KRAS G12C by a trapping mechanism. *Science* 351:604–608.
- Maurer T, et al. (2012) Small-molecule ligands bind to a distinct pocket in Ras and inhibit SOS-mediated nucleotide exchange activity. *Proc Natl Acad Sci USA* 109: 5299–5304.
- Ostrem JM, Peters U, Sos ML, Wells JA, Shokat KM (2013) K-Ras(G12C) inhibitors allosterically control GTP affinity and effector interactions. *Nature* 503:548–551.
- Sun Q, et al. (2012) Discovery of small molecules that bind to K-Ras and inhibit Sos-mediated activation. *Angew Chem Int Ed Engl* 51:6140–6143.
- Winter JJ, et al. (2015) Small molecule binding sites on the Ras:SOS complex can be exploited for inhibition of Ras activation. *J Med Chem* 58:2265–2274.
- Athuluri-Divakar SK, et al. (2016) A small molecule RAS-mimetic disrupts RAS association with effector proteins to block signaling. *Cell* 165:643–655.
- Patricelli MP, et al. (2016) Selective inhibition of oncogenic KRAS output with small molecules targeting the inactive state. *Cancer Discov* 6:316–329.
- Welsch ME, et al. (2017) Multivalent small-molecule Pan-RAS inhibitors. *Cell* 168: 878–889.e29.
- Quevedo CE, et al. (2018) Small molecule inhibitors of RAS-effector protein interactions derived using an intracellular antibody fragment. *Nat Commun* 9:3169.
- Grant BJ, et al. (2011) Novel allosteric sites on Ras for lead generation. *PLoS One* 6: e25711.
- Sewell H, et al. (2014) Conformational flexibility of the oncogenic protein LMO2 primes the formation of the multi-protein transcription complex. *Sci Rep* 4:3643.
- Bery N, et al. (2018) BRET-based RAS biosensors that show a novel small molecule is an inhibitor of RAS-effector protein-protein interactions. *eLife* 7:e37122.
- Johnstone S, Albert JS (2017) Pharmacological property optimization for allosteric ligands: A medicinal chemistry perspective. *Bioorg Med Chem Lett* 27:2239–2258.
- Lawson ADG, MacCoss M, Heer JP (2018) Importance of rigidity in designing small molecule drugs to tackle protein-protein interactions (PPIs) through stabilization of desired conformers. *J Med Chem* 61:4283–4289.
- Andersson V, et al. (2016) Macrocyclic prodrugs of a selective nonpeptidic direct thrombin inhibitor display high permeability, efficient bioconversion but low bio-availability. *J Med Chem* 59:6658–6670.
- Cruikshank DWJ (1999) Remarks about protein structure precision. *Acta Crystallogr D Biol Crystallogr* 55:583–601.
- Beck A, Goetsch L, Dumontet C, Corvaia N (2017) Strategies and challenges for the next generation of antibody-drug conjugates. *Nat Rev Drug Discov* 16:315–337.
- Town J, et al. (2016) Exploring the surfaceome of Ewing sarcoma identifies a new and unique therapeutic target. *Proc Natl Acad Sci USA* 113:3603–3608.
- Martinko AJ, et al. (2018) Targeting RAS-driven human cancer cells with antibodies to upregulated and essential cell-surface proteins. *eLife* 7:e31098.
- Cruz-Migoni A, et al. (2018) Structure-based development of new RAS-effector inhibitors from a combination of active and inactive RAS-binding compounds. Protein Data Bank. Available at <https://www.rcsb.org/structure/6GOD>. Deposited June 1, 2018.
- Cruz-Migoni A, et al. (2018) Structure-based development of new RAS-effector inhibitors from a combination of active and inactive RAS-binding compounds. Protein Data Bank. Available at <https://www.rcsb.org/structure/6GOE>. Deposited June 1, 2018.
- Cruz-Migoni A, et al. (2018) Structure-based development of new RAS-effector inhibitors from a combination of active and inactive RAS-binding compounds. Protein Data Bank. Available at <https://www.rcsb.org/structure/6GOF>. Deposited June 1, 2018.
- Cruz-Migoni A, et al. (2018) Structure-based development of new RAS-effector inhibitors from a combination of active and inactive RAS-binding compounds. Protein Data Bank. Available at <https://www.rcsb.org/structure/6GOG>. Deposited June 1, 2018.
- Cruz-Migoni A, et al. (2018) Structure-based development of new RAS-effector inhibitors from a combination of active and inactive RAS-binding compounds. Protein Data Bank. Available at <https://www.rcsb.org/structure/6GOM>. Deposited June 1, 2018.
- Cruz-Migoni A, et al. (2018) Structure-based development of new RAS-effector inhibitors from a combination of active and inactive RAS-binding compounds. Protein Data Bank. Available at <https://www.rcsb.org/structure/6GQT>. Deposited June 8, 2018.
- Cruz-Migoni A, et al. (2018) Structure-based development of new RAS-effector inhibitors from a combination of active and inactive RAS-binding compounds. Protein Data Bank. Available at <https://www.rcsb.org/structure/6GQW>. Deposited June 5, 2018.
- Cruz-Migoni A, et al. (2018) Structure-based development of new RAS-effector inhibitors from a combination of active and inactive RAS-binding compounds. Protein Data Bank. Available at <https://www.rcsb.org/structure/6GQX>. Deposited June 5, 2018.
- Cruz-Migoni A, et al. (2018) Structure-based development of new RAS-effector inhibitors from a combination of active and inactive RAS-binding compounds. Protein Data Bank. Available at <https://www.rcsb.org/structure/6GQY>. Deposited June 5, 2018.