

Strong constraints on aerosol-cloud interactions from volcanic eruptions

Authors: Florent F. Malavelle^{1*}, Jim M. Haywood^{1,2}, Andy Jones², Andrew Gettelman³, Lieven Clarisse⁴, Sophie Bauduin⁴, Richard P. Allan^{5,6}, Inger Helene H. Karset⁷, Jón Egill Kristjánsson^{7,S}, Lazaros Oreopoulos⁸, Nayeong Cho^{8,9}, Dongmin Lee^{8,10}, Nicolas Bellouin⁵, Olivier Boucher¹¹, Daniel P. Grosvenor¹², Ken S. Carslaw¹², Sandip Dhomse¹², Graham W. Mann^{12,13}, Anja Schmidt¹², Hugh Coe¹⁴, Margaret E. Hartley¹⁴, Mohit Dalvi², Adrian A. Hill², Ben T. Johnson², Colin E. Johnson², Jeff R. Knight², Fiona M. O'Connor², Daniel G. Partridge^{15,16,17,#}, Philip Stier¹⁷, Gunnar Myhre¹⁸, Steven Platnick⁸, Graeme L. Stephens¹⁹, Hanii Takahashi^{20,19}, Thorvaldur Thordarson²¹.

Affiliations:

¹College of Engineering, Mathematics, and Physical Sciences, University of Exeter, Exeter, UK.

²Met Office Hadley Centre, Exeter, UK.

³National Center for Atmospheric Research, Boulder, Colorado, USA.

⁴Chimie Quantique et Photophysique CP160/09, Université Libre de Bruxelles (ULB), Bruxelles, Belgium.

⁵Department of Meteorology, University of Reading, Reading, UK.

⁶National Centre for Earth Observation, University of Reading, UK.

⁷Department of Geosciences, University of Oslo, Oslo, Norway.

⁸Earth Sciences Division, NASA GSFC, Greenbelt, Maryland, USA.

⁹USRA, Columbia, Maryland, USA.

¹⁰Morgan State University, Baltimore, Maryland, USA.

¹¹Laboratoire de Météorologie Dynamique, IPSL, UPMC/CNRS, Jussieu, France.

¹²School of Earth and Environment, University of Leeds, Leeds, UK.

¹³National Centre for Atmospheric Science, University of Leeds, Leeds, UK.

¹⁴School of Earth and Environmental Sciences, University of Manchester, Manchester, UK.

¹⁵Department of Environmental Science and Analytical Chemistry, University of Stockholm, Stockholm, Sweden

¹⁶Bert Bolin Centre for Climate Research, University of Stockholm, Stockholm, Sweden

¹⁷Atmospheric, Oceanic and Planetary Physics, Department of Physics, University of Oxford, Oxford, UK.

¹⁸Center for International Climate and Environmental Research, Oslo, Norway.

¹⁹Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA.

²⁰Joint Institute for Regional Earth System Science and Engineering, University of California, Los Angeles, California, USA

²¹Faculty of Earth Sciences, University of Iceland, Reykjavik, Iceland.

*Corresponding author: f.malavelle@exeter.ac.uk

^{\$}Deceased 14th August 2016.

[#]Now at College of Engineering, Mathematics, and Physical Sciences, University of Exeter, Exeter, UK.

Summary (149 words of referenced text):

The climate impact of aerosols is highly uncertain owing primarily to their poorly quantified influence on cloud properties. During 2014-15, a fissure eruption in Holuhraun (Iceland) emitted huge quantities of sulphur dioxide, resulting in significant reductions in liquid cloud droplet size. Using satellite observations and detailed modelling, we estimate a global mean radiative forcing from the resulting aerosol-induced cloud brightening for the time of the eruption of around -0.2 W.m^{-2} . Changes in cloud amount or liquid water path are undetectable, indicating that these aerosol-cloud indirect effects are modest. It supports the

idea that cloud systems are well buffered against aerosol changes as only impacts on cloud effective radius appear relevant from a climate perspective, thus providing a strong constraint on aerosol-cloud interactions. This result will reduce uncertainties in future climate projections as we are able to reject the results from climate models with an excessive liquid water path response.

Main Text: (3103 words of referenced text, including concluding paragraph)

1. The 2014-15 eruption at Holuhraun (486 words of referenced text):

Anthropogenic emissions that affect climate are not just confined to greenhouse gases. Sulphur dioxide and other pollutants form atmospheric aerosols that can scatter and absorb sunlight and can influence the properties of clouds, modulating the Earth-atmosphere energy balance. Aerosols act as cloud condensation nuclei (CCN); an increase in CCN translates into a higher number of smaller, more reflective cloud droplets that scatter more sunlight back to space¹ (the ‘first’ indirect effect of aerosols). Smaller cloud droplets decrease the efficiency of collision-coalescence processes that are pivotal in rain initiation, thus aerosol-influenced clouds may retain more liquid water and extend coverage/lifetime^{2,3} (the ‘second’ or ‘cloud lifetime’ indirect effect). Aerosols usually co-vary with key environmental variables making it difficult to disentangle aerosol-cloud impacts from meteorological variability⁴⁻⁶. Additionally, clouds themselves are complex transient systems subject to dynamical feedbacks (e.g. cloud top entrainment/evaporation, invigoration of convection) which influence cloud response⁷⁻¹². These aspects present great challenges in evaluating and constraining aerosol-cloud interactions (ACI) in General Circulation Models (GCM)¹³⁻¹⁷, with particular contentious debate surrounding the relative importance of these feedback mechanisms.

Nonetheless, anthropogenic aerosol emissions are thought to cool the Earth via indirect effects¹⁷, but the uncertainty ranges from -1.2 to -0.0 W.m⁻² (90% confidence interval) due to *i)* a lack of characterization of the pre-industrial aerosol state^{15,18,19}, and *ii)* model parametric

and structural errors in representing cloud responses to aerosol changes^{16,18,20,21}. It is estimated that uncertainty in the pre-industrial state can account for approximately 30% of total ACI uncertainty^{18,21} while representation of chemistry-aerosol-cloud processes in models is responsible for the remaining 70% uncertainty^{16,21}. Recently, a framework to break down uncertainties in the causal chain from emission to radiative forcing showed that the sources of uncertainty within different GCMs differ greatly¹⁶.

Volcanic eruptions provide invaluable natural experiments to investigate the role of large-scale aerosol injection in the Earth system²²⁻²⁶. There have been several Icelandic volcanic eruptions over recent years; Eyjafjallajökull erupted in 2010, Grímsvötn in 2011 and Holuhraun in 2014-15. At its peak, the 2014-15 eruption at Holuhraun emitted ~120 kt of sulphur dioxide (SO₂) per day into the atmosphere, a rate some four times higher than all 28 European Union member states or over a third of global emission rates. Iceland became in effect a continental-scale pollution source of SO₂; SO₂ is readily oxidised via gas- and aqueous-phase reactions, producing a massive aerosol plume in a near-pristine environment where clouds should be most susceptible to aerosol concentrations^{16,18,27}.

We advance upon preliminary observational assessments of the impact of the 2014-15 eruption at Holuhraun^{28,29} through an extensive observational analysis that includes a statistical evaluation of the significance of the observed spatial distribution of the cloud perturbations to untangle the impacts of aerosol/meteorological impacts. We then assess the simulation from a range of different climate models and assess the performance against available observations. Finally, we show that observations of a volcanic plume (Mt. Kilauea, Hawaii) in an entirely different meteorological regime exhibit similar overall impacts.

2. Impact of the eruption on clouds (2140 - 20 = 2120 words of referenced text):

Following the lifecycle of sulphur from emission, our initial analysis concentrates on the coherence of SO₂ detected by the Infrared Atmospheric Sounding Interferometer (IASI) sensor (Supplementary M1) and the HadGEM3 GCM that is constrained by observed

temperatures and winds (i.e. nudged, Supplementary M2). IASI retrievals use the discrete spectral absorption structure of SO₂ to determine concentrations³⁰. Comparisons of IASI SO₂ observations from explosive volcanic eruptions against model simulations have proven valuable in the past^{31,32}. The processing procedure for quantitative comparison between IASI and HadGEM3 data uses only data that are spatially and temporally coherent (Supplementary M3).

There is considerable uncertainty in the quantitative emission of SO₂ from the 2014-15 eruption at Holuhraun. A previous study²⁸ assumed a constant emission rate of 40 kt[SO₂]/day based on initial estimates of degassing. As our standard scenario (STAN) we use an empirical relationship between degassed sulphur and TiO₂/FeO ratios and lava production derived from Icelandic basaltic flood lava eruptions³³ which suggests significantly higher emissions during the early phase of the eruption in September, but we also investigate a simulation where a constant 40 ktSO₂/day is released (40KT scenario). The model simulations and IASI retrievals of column SO₂ are shown in Figure 1 (40KT emission scenario shown in Supplementary S1).

*****Insert Figure 1 here*****

The distribution and the magnitude of the column loading of SO₂ detected by IASI are similar to those derived from HadGEM3, showing that the GCM nudging scheme and the assumed altitude of the emissions in the STAN scenario (surface to 3 km) reproduces the week to week spatial variability and magnitude of observed column SO₂ (SI-SO2_animation.mp4).

While the spatial distribution of sulphate aerosol optical depth (*AOD*) caused by the eruption can be determined easily in the model (Supplementary Fig. S2.1), detection of the aerosol plume over the north Atlantic in the MODIS data is hampered by the mutual exclusivity of aerosol and cloud retrievals. The predominance of cloudy scenes makes accurate detection of the aerosol plume in monthly-mean MODIS data extremely challenging (Supplementary S2).

Nonetheless, despite lacking observations of AOD , we can look for evidence of perturbations caused by aerosols on cloud properties. We examine the perturbation to retrieved cloud top droplet effective radius (r_{eff}) in September and October 2014 using collection 051 monthly mean data from MODIS AQUA (MYD08, Supplementary M4) over the period 2002-2014. MODIS AQUA data are not subject to the degradation in performance of the sensors at visible wavelengths that has recently been documented for the MODIS TERRA³⁴ sensor (Supplementary S3). We present a summary of the change in r_{eff} , Δr_{eff} , for October 2014 compared to the long term 2002-2013 mean in Figure 2a. A full analysis of the year-to-year variability in Δr_{eff} is presented in Supplementary S4.

*****Insert Figure 2 here*****

There is clear evidence of a signal in Δr_{eff} in October (Figures 2a) and September (Supplementary Fig. S5.1a). Pixels that are statistically significantly different from the 2002-2013 climatological mean at 95% confidence occur over the entire breadth of the north Atlantic. The spatial distribution of Δr_{eff} is governed by the prevailing wind conditions that advect the volcanic plume and are quantitatively similar to those noted in Collection 006 MODIS data²⁹.

Figures 3a show the corresponding Δr_{eff} derived from the model in October (for September, Supplementary Fig. S5.2a). The observations and modelling show obvious similarities in spatial distribution. In addition to the spatial coherence in Δr_{eff} , the changes in the model of $-1.21 \mu\text{m}$ (September) and $-0.68 \mu\text{m}$ (October) are within 30% of MODIS Δr_{eff} of $-0.98 \mu\text{m}$ (September) and $-0.97 \mu\text{m}$ (October) for the domain shown in Figure 2.

*****Insert Fig 3 here*****

161 There are similarities between the MODIS and HadGEM3 probability distribution functions
 162 (Figures 2b and 3b) with a shift to smaller r_{eff} for the year of the eruption. Almost all high
 163 values of r_{eff} (i.e. $r_{eff} > \sim 16 \mu\text{m}$ for MODIS and $r_{eff} > \sim 11 \mu\text{m}$ for HadGEM3) are absent in
 164 2014 suggesting that clouds with high r_{eff} are entirely absent from the domain in both the
 165 observations and the model. There are obvious discrepancies in the absolute magnitude of r_{eff}
 166 between MODIS and HadGEM3. MODIS retrievals of r_{eff} from the MYD06 product in liquid
 167 water cloud regimes have been shown to be significantly larger than those derived from other
 168 satellite sensor products, mainly due to the algorithm's use of a different primary spectral
 169 channel relative to other products^{35,36}. Nevertheless, Δr_{eff} is in encouraging agreement as this
 170 quantity, along with changes in cloud liquid water path (LWP), needs to be accurately
 171 represented if aerosol-cloud interactions are to be better quantified. As with r_{eff} , there are
 172 similarities between the MODIS and HadGEM3 for ΔLWP (Figure 2c-d and Figure 3c-d),
 173 however, evidence of a clear signal due to the volcano is neither observed or modelled.
 174 Additionally, we also found that perturbations in the monthly mean cloud fraction from
 175 MODIS are negligible, both in September and October as previously reported²⁹.
 176 It is incumbent on any study attributing Δr_{eff} to volcanic emissions to prove the causality
 177 beyond reasonable doubt, i.e. that the changes are not due to natural meteorological
 178 variability. The meteorological analyses in Supplementary S6 suggest that, while in
 179 September 2014 the southern part of the spatial domain shown in Figure 2 is somewhat
 180 influenced by anomalous easterlies bringing pollution from the European continent over the
 181 easternmost Atlantic Ocean and hence influencing r_{eff} , the perturbations to r_{eff} during October
 182 2014 are entirely of volcanic origin.
 183 MODIS and HadGEM3 show a similar spatial distribution and magnitude for October for the
 184 perturbation in cloud droplet number concentration (ΔN_d), but a smaller ΔN_d in MODIS than
 185 in HadGEM3 for September 2014 (Supplementary S7.2). Once r_{eff} is reduced, the
 186 autoconversion process whereby cloud droplets grow to sufficient size to form precipitation

may be inhibited, leading to clouds with increased liquid water path³. The cloud optical depth, τ_{cloud} , is related to r_{eff} and LWP and the density of water (ρ) by the approximation:

$$\tau_{cloud} \cong \frac{3LWP}{2\rho r_{eff}} \quad (1)$$

We use HadGEM3 to assess the detectability of perturbations against natural variability. Two different methods are pursued using the nudged model; firstly, assessing model simulations with and without the emissions from the eruption for the year 2014 (HOL₂₀₁₄-NO_HOL₂₀₁₄), and secondly assessing model simulations including emissions from Holuhraun for 2014 against simulations for 2002-2013 (HOL₂₀₁₄-NO_HOL₂₀₀₂₋₂₀₁₃). While the former method allows the ‘cleanest’ assessment of the impacts of the eruption (as the meteorology is effectively identical and meteorological variability is removed), the second method allows assessment of the statistical significance against the natural meteorological variability. This provides an assessment that is directly comparable to observations and can be used to effectively isolate signal from noise³⁷ (Supplementary S7).

Insert Figure 4 here

Figure 4 shows that ΔAOD , ΔN_d , and Δr_{eff} are statistically significant at 95% confidence across the majority of latitudes. The fact that the simulations from [HOL₂₀₁₄-NO_HOL₂₀₁₄] and [HOL₂₀₁₄-NO_HOL₂₀₀₂₋₂₀₁₃] are similar for these variables again indicates that the impacts of natural meteorological variability on these variables is small (i.e. NO_HOL₂₀₁₄ \approx NO_HOL₂₀₀₂₋₂₀₁₃). For ΔLWP , no statistically significant changes are evident at either 95% or 67% confidence, suggesting that meteorological variability provides a far stronger control on cloud LWP than aerosol (Supplementary S7.3). With ΔLWP being due to meteorological noise, $\Delta \tau_{cloud}$ is driven by Δr_{eff} and Figure 4e suggests that the perturbations to τ_{cloud} north of around 67°N/57°N, which are significant at the 95%/67% confidence level, are due to the 2014-15 Holuhraun eruption. Our simulations suggest that Top of Atmosphere changes in

short wave radiation (ΔToA_{SW}) are unlikely to be detectable at 95% or even 67% confidence when compared to natural variability. More details supporting this assertion are given in Supplementary S7.5 which uses satellite observations of the Earth's radiation budget.

We have shown that HadGEM3 is capable of representing observations of aerosol-cloud interactions with a reasonable representation of the perturbation to r_{eff} but minimal perturbation to LWP . To demonstrate the practical value of the study, we repeat the simulations with other models. First, we use HadGEM3 but using the older single moment CLASSIC³⁸ aerosol scheme instead of the new two-moment UKCA/GLOMAP-mode scheme³⁹. We also perform calculations with the NCAR Community Atmosphere Model²⁸ (CAM5-NCAR) and the atmospheric component of an intermediate version of the Norwegian Earth System Model⁴⁰ (CAM5-Oslo), driven using nominally the same emissions and plume top height. CAM5-NCAR has been used previously in free-running mode to provide an initial estimate of the radiative forcing of the 2014-15 Holuhraun eruption²⁸, but as in the HadGEM3 simulations we run CAM5-NCAR and CAM5-Oslo in nudged mode to simulate the meteorology during the eruption as closely as possible. Figure 5 shows a comparison of Δr_{eff} and ΔLWP derived from $HOL_{2014}-NO_HOL_{2014}$ simulations from HadGEM3, HadGEM3-CLASSIC, CAM5-NCAR, CAM5-Oslo and MODIS for October. We chose October as the contribution from continental Europe pollution to cloud property anomalies has been shown to be small (Supplementary S4-6-7; Supplementary S8 shows the impacts on cloud properties in September).

****Insert Figure 5 here****

It is immediately apparent from the first column of Figure 5 that HadGEM3 using UKCA, CAM5-NCAR, and CAM5-Oslo are able to accurately model the impact on Δr_{eff} , while HadGEM3-CLASSIC produces an impact that is too strong when compared to the MODIS observations owing to the single moment nature of the aerosol scheme (Supplementary S9).

240 For ΔLWP , as we have seen from the multi-year analysis of MODIS (Supplementary Fig.
241 S7.3), the meteorological variability is the controlling factor. Even with meteorological
242 variability suppressed in these [HOL₂₀₁₄-NO_HOL₂₀₁₄] results, HadGEM3 using UKCA
243 shows only a very limited increase in LWP (Fig. 5f), HadGEM3-CLASSIC and CAM5-Oslo
244 show a progressively more significant response whereas CAM5-NCAR shows a much larger
245 response (Fig. 5h).

246 It is insightful to examine the influence of the eruption on precipitation in both observations
247 and models using a similar analysis (Supplementary S10). We observe that there is little
248 impact on precipitation indicating that the cloud system readjusts to a new equilibrium with
249 little impact on either LWP or precipitation. The larger response in CAM5-NCAR ($\Delta LWP >$
250 16 g.m^{-2}) is not supported by the MODIS observations where the 2002-2013 domain mean
251 standard deviation in ΔLWP is $\sim 4.5 \text{ g.m}^{-2}$. Thus, we are able to use the eruption to evaluate
252 the models: HadGEM3 using UKCA and CAM5-Oslo perform in a manner consistent with
253 the MODIS observations while HadGEM3-CLASSIC and CAM5-NCAR do not. Moreover,
254 the fact that changes in LWP are not detectable above natural variability suggests that
255 aerosol-cloud interactions beyond the impact on r_{eff} are small (i.e. net second indirect effects
256 are small).

257 The effective radiative forcing (ERF) from the event may be estimated from the difference
258 between the top of atmosphere net irradiances from simulations including and excluding the
259 volcanic emissions. The global ERF from HadGEM3 over the September-October 2014
260 period is estimated at -0.21 W.m^{-2} . Tests using an offline version of the radiation code reveal
261 that the presence of overlying ice-cloud weakens the ERF by approximately 20%
262 (Supplementary S11).

263 We also investigate whether a fissure eruption of this magnitude could have a more
264 significant radiative impact if the timing/location of the eruptions were different
265 (Supplementary S12). Our simulations suggest that for contrasting scenarios the global ERF
266 would *i)* strengthen to -0.29 W.m^{-2} (+40%) if the eruption commenced at the beginning of

June, *ii*) strengthen to -0.49 W.m^{-2} (+140%) if the fissure eruption had occurred in an area of South America where it could affect clouds in a stratocumulus-dominated regime, *iii*) strengthen to -0.32 W.m^{-2} (+55%) if the eruption had occurred in pre-industrial times when the background concentrations of aerosols was reduced¹⁸ indicating that climatic impact of fissure eruptions such as Laki⁴¹ in 1783-1784 would not have been as large if it had occurred in the present day.

Many studies^{9,11,42,43} suggest that cloud adjustments may be dependent upon meteorological regime, so we ask whether the cloud *LWP* invariance observed near Holuhraun is simply a special case. We have reproduced the cloud regimes analysis derived from satellite measurements presented in a recent study⁴⁴. We find that, when examining the 2014-15 eruption at Holuhraun, we are far from examining a meteorological ‘special case’, in fact rather the opposite (Supplementary S13); we are examining a region that contains the whole spectrum of liquid-dominated cloud regimes and deducing that, overall, the impact on *LWP* is minimal.

To further support our conclusion, we report results from a different event (Mount Kilauea, Hawaii, Supplementary S14), which degassing rate significantly increased during June-August 2008. The outflow of the plume affected the surrounding trade maritime cumuli^{24,45,46}, increasing the SW reflectance; the causal interpretations of this in the literature have varied^{24,46}. affecting the surrounding trade maritime cumuli^{24,45,46} and increased the SW reflectance in the outflow of the plume, although with different causal interpretations^{24,46}. Again, *LWP* does not vary, either in the AMSR-E data⁴⁶ or in the MODIS monthly retrievals (Supplementary S14) which again suggests *LWP* insensitivity in the trade cumulus regime as well. Thus, for a very different meteorological environment dominated by very different cloud regimes, similar conclusions emerge.

4. Discussion and Conclusion (507 words of referenced text):

293 The 2014-15 eruption at Holuhraun presents a unique opportunity to investigate continental-
294 scale aerosol-cloud climatic effects. Using synergistic observations and models driven by an
295 empirical estimate of SO₂ emissions³³ we simulate spatial distributions of SO₂ that compare
296 favourably with satellite observations. The HadGEM3 model is able to predict an impact
297 from aerosol-cloud interactions of similar magnitude to the signal found in the MODIS data.
298 Our analysis further highlights that cloud properties are largely unaffected by the eruption
299 beyond the impact on r_{eff} .

300 We repeated the experiment with two additional GCMs and show that HadGEM3 using
301 UKCA, CAM5-NCAR and CAM5-Oslo are able to capture the magnitude of the observed
302 impacts on r_{eff} despite the lack of explicit representation of processes such as sub-cloud
303 updraft velocities and entrainment, enhancing our confidence in GCMs' ability in predicting
304 the aerosol first indirect effect. However, in line with recent work¹⁶, modelled responses in
305 the LWP differ significantly. The fact that cloud adjustments via LWP are not identified in the
306 observations of the 2014-15 eruption at Holuhraun indicates that clouds are buffered against
307 LWP changes^{9-10,12}, providing evidence that models with a low LWP response display a more
308 convincing behaviour. These findings have wide scientific relevance in the field of climate
309 modelling as, in terms of climate forcing, they suggest that aerosol second indirect effects
310 appear small and climate models with a significant LWP feedback need reassessment^{15-16,47}.

311 Despite such massive emissions and large anomalies in r_{eff} , we estimate a moderate global-
312 mean radiative forcing of $-0.21 \pm 0.08 \text{ W.m}^{-2}$ (1 standard deviation, Supplementary S15) for
313 September-October which equates to a global annual mean effective radiative forcing of
314 $-0.035 \pm 0.013 \text{ W.m}^{-2}$ (1 standard deviation) assuming that a forcing only occurs in
315 September and October 2014. Global emissions of anthropogenic SO₂ currently total around
316 100 TgSO₂/year and the Intergovernmental Panel on Climate Change^{17,47} suggests a best
317 estimate for the aerosol forcing of -0.9 W.m^{-2} , yielding a forcing efficiency of -0.009
318 $\text{W.m}^{-2}/\text{TgSO}_2$. The emissions for September and October 2014 total approximately 4 TgSO₂,
319 thus the global annual mean radiative forcing efficiency for the 2014-15 eruption at

Holuhraun yields a forcing efficiency of $-0.0088 \pm 0.0024 \text{ W.m}^{-2}/\text{TgSO}_2$ (1 standard deviation). The similarity is remarkable, but may be by chance given the modelled sensitivity to emission location and time (Supplementary S12).

Our study is not without caveats given that the observations themselves are uncertain owing to the limitations of satellite retrievals. The modelling is not completely constrained owing to the lack of detailed in-situ observations of e.g. the background aerosol concentrations and plume height. We cannot rule out that models showing small *LWP* sensitivity to aerosol emission behave as they do because they lack the resolution to represent fine-scale dynamical feedbacks^{9,12}. Further high-resolution modelling of the 2014-15 Holuhraun eruption is necessary to evaluate more thoroughly how processes such as autoconversion or droplet evaporation plays a role in buffering the aerosol effect^{9,12,48,49}. Bringing many of the different global models together and inter-comparing results of Holuhraun simulations is merited to provide a traceable route for reducing the uncertainty in future climate projections.

References:

¹Twomey, S., The influence of pollution on the shortwave albedo of clouds. *J. Atmos. Sci.*, 34:1149–1152 (1977).

²Albrecht, B. A., Aerosols, cloud microphysics, and fractional cloudiness. *Science*, 245(4923):1227–1230 (1989).

³Haywood, J.M., and Boucher, O., Estimates of the direct and indirect radiative forcing due to tropospheric aerosols: a review. *Reviews of Geophysics*, 38, 513-543 (2000).

⁴Lohmann, U., Koren, I. and Kaufman, Y. J., Disentangling the role of microphysical and dynamical effects in determining cloud properties over the Atlantic. *Geophys. Res. Lett.*, 33, L09802, doi:10.1029/2005GL024625 (2006).

⁵Mauger, G. S., and J. R. Norris, Meteorological bias in satellite estimates of aerosol-cloud relationships. *Geophys. Res. Lett.*, 34, L16824, doi:10.1029/2007GL029952 (2007).

⁶Gryspeerd, E., Quaas, J. and Bellouin, N., Constraining the aerosol influence on cloud fraction. *J. Geophys. Res. Atmos.*, 121, 3566–3583, doi:10.1002/2015JD023744 (2016).

348 ⁷Ackerman, A. S. et al., The impact of humidity above stratiform clouds on indirect climate
349 forcing. *Nature*, 432, 1014–1017 (2004).

350 ⁸Sandu, I., J. L. Brenguier, O. Geoffroy, O. Thoueron, and V. Masson, Aerosol impacts on the diurnal
351 cycle of marine stratocumulus. *J. Atmos. Sci.*, 65, 2705–2718, doi:10.1175/2008JAS2451.1 (2008).

352 ⁹Stevens, B. and Feingold, G., Untangling aerosol effects on clouds and precipitation in a buffered
353 system. *Nature*, 461, 607–613 (2009).

354 ¹⁰Seifert, A., Köhler, C., and Beheng, K. D., Aerosol-cloud-precipitation effects over Germany as
355 simulated by a convective-scale numerical weather prediction model. *Atmos. Chem. Phys.*, 12, 709-
356 725, doi:10.5194/acp-12-709-2012 (2012).

357 ¹¹Lebo, Z. J. and Feingold, G., On the relationship between responses in cloud water and precipitation
358 to changes in aerosol. *Atmos. Chem. Phys.*, 14:11817–11831 (2014).

359 ¹²Seifert, A., T. Heus, R. Pincus, and B. Stevens, Large-eddy simulation of the transient and near-
360 equilibrium behaviour of precipitating shallow convection. *J. Adv. Model. Earth Syst.*, 7, 1918–1937,
361 doi:10.1002/2015MS000489 (2015).

362 ¹³Quaas, J. et al., Aerosol indirect effects - general circulation model intercomparison and evaluation
363 with satellite data. *Atmos. Chem. Phys.*, 9, 8697–8717, doi:10.5194/acp-9-8697-2009 (2009).

364 ¹⁴Penner, J. E., Xu, L. & Wang, M. H., Satellite methods underestimate indirect climate forcing by
365 aerosols. *Proc. Natl Acad. Sci.*, USA 108, 13404–13408, doi:10.1073/pnas.1018526108 (2011).

366 ¹⁵Stevens, B., Rethinking the Lower Bound on Aerosol Radiative Forcing. *J. Clim.*, 28, 4794–4819,
367 doi:10.1175/JCLI-D-14-00656.1 (2015).

368 ¹⁶Ghan, S. et al., Challenges in constraining anthropogenic aerosol effects on cloud radiative forcing
369 using present-day spatiotemporal variability. *Proc. Natl. Acad. Sci. USA*, 113, 5804–5811,
370 doi:10.1073/pnas.1514036113 (2016).

371 ¹⁷Boucher, O. et al., Clouds and Aerosols. In: Climate Change 2013: The Physical Science Basis.
372 Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on
373 Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A.
374 Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United
375 Kingdom and New York, NY, USA (2013).

376 ¹⁸Carslaw, K. S. et al., Large contribution of natural aerosols to uncertainty in indirect Forcing.
377 *Nature*, 503(7474):67–71 (2013).

378 ¹⁹Hamilton, D. S. *et al.*, Occurrence of pristine aerosol environments on a polluted planet.
379 *Proceedings of the National Academy of Sciences of the United States of America*,
380 doi:10.1073/pnas.1415440111 (2014).

381 ²⁰Lohmann, U. *et al.*, Total aerosol effect: radiative forcing or radiative flux perturbation?. *Atmos.*
382 *Chem. Phys.*, 10, 3235-3246, doi:10.5194/acp-10-3235-2010 (2010).

383 ²¹Gettelman, A., Putting the clouds back in aerosol-cloud interactions. *Atmos. Chem. Phys.*,
384 15:12397–12411, doi:10.5194/acp-15-12397-2015 (2015).

385 ²²McCormick, M.P., Thomason, L.W., and Trepte, C.R., Atmospheric effects of the Mt. Pinatubo
386 eruption. *Nature*, v. 373, p. 399—404, doi:10.1038/373399a0 (1995).

387 ²³Gassó, S., Satellite observations of the impact of weak volcanic activity on marine clouds. *J.*
388 *Geophys. Res.*, 113, D14S19, doi:10.1029/2007JD009106 (2008).

389 ²⁴Yuan, T., Remer, L. A., and Yu, H., Microphysical, macrophysical and radiative signatures of
390 volcanic aerosols in trade wind cumulus observed by the A-Train. *Atmos. Chem. Phys.*, 11, 7119-
391 7132, doi:10.5194/acp-11-7119-2011 (2011).

392 ²⁵Schmidt, A. *et al.*, Importance of tropospheric volcanic aerosol for indirect radiative forcing of
393 climate. *Atmos. Chem. Phys.*, 12, 7321-7339, doi:10.5194/acp-12-7321-2012 (2012).

394 ²⁶Haywood, J. M., Jones, A. and Jones, G. S., The impact of volcanic eruptions in the period 2000–
395 2013 on global mean temperature trends evaluated in the HadGEM2-ES climate model. *Atmos. Sci.*
396 *Lett.*, 15: 92–96. doi:10.1002/asl2.471 (2014).

397 ²⁷Penner, J. E., C. Zhou, and L. Xu, Consistent estimates from satellites and models for the first
398 aerosol indirect forcing. *Geophys. Res. Lett.*, 39, L13810, doi:10.1029/2012GL051870 (2012).

399 ²⁸Gettelman, A., A. Schmidt, and J.-E. Kristjánsson, Icelandic volcanic emissions and climate. *Nature*
400 *Geoscience*, 8, 243, doi:10.1038/ngeo2376 (2015).

401 ²⁹McCoy, D. T., and D. L. Hartmann, Observations of a substantial cloud-aerosol indirect effect
402 during the 2014–2015 Bárðarbunga-Veiðivötn fissure eruption in Iceland. *Geophys. Res. Lett.*, 42,
403 10,409–10,414, doi:10.1002/2015GL067070 (2015).

404 ³⁰Clarisse, L. *et al.*, Tracking and quantifying volcanic SO₂ with IASI, the September 2007 eruption
405 at Jebel at Tair. *Atmos. Chem. Phys.*, 8, 7723–7734, doi:10.5194/acp-8-7723-2008 (2008).

406 ³¹Haywood, J.M. *et al.*, Observations of the eruption of the Sarychev volcano and simulations using
407 the HadGEM2 climate model. *J. Geophys. Res.*, 115, D21212, doi:10.1029/2010JD014447 (2010).

408 ³²Schmidt, A. *et al.*, Satellite detection, long-range transport, and air quality impacts of volcanic sulfur
409 dioxide from the 2014–2015 flood lava eruption at Bárðarbunga (Iceland). *J. Geophys. Res. Atmos.*,
410 120, doi:10.1002/2015JD023638 (2015).

411 ³³Thordarson, T., Self, S., Miller, D. J., Larsen, G., & Vilmundardóttir, E. G., Sulphur release from
412 flood lava eruptions in the Veidivötn, Grímsvötn and Katla volcanic systems, Iceland. Geological
413 Society, London, Special Publications, 213(1), 103-121 (2003).

414 ³⁴Polashenski, C. M. *et al.*, Neither dust nor black carbon causing apparent albedo decline in
415 Greenland's dry snow zone: Implications for MODIS C5 surface reflectance. *Geophys. Res. Lett.*, 42,
416 doi:10.1002/2015GL065912 (2015).

417 ³⁵Platnick, S. *et al.*, MODIS Atmosphere L2 Cloud Product (06_L2). NASA MODIS Adaptive
418 Processing System, Goddard Space Flight Center, USA:
419 http://dx.doi.org/10.5067/MODIS/MOD06_L2.006 (2015).

420 ³⁶Zhang, Z. and Platnick, S., An assessment of differences between cloud effective particle radius
421 retrievals for marine water clouds from three MODIS spectral bands. *Journal of Geophysical*
422 *Research: Atmospheres* (1984–2012), 116(D20) (2011).

423 ³⁷Stevens, B., and J-L Brenguier, Cloud Controlling Factors – Low Clouds, Heintzenberg, J., and R. J.
424 Charlson, eds. *Clouds in the Perturbed Climate System: Their Relationship to Energy Balance,*
425 *Atmospheric Dynamics, and Precipitation.* Strüngmann Forum Report, vol. 2. Cambridge, MA: MIT
426 Press ISBN 978-0-262-01287-4 (2009).

427 ³⁸Bellouin, N. *et al.*, Aerosol forcing in the CMIP5 simulations by HadGEM2-ES and the role of
428 ammonium nitrate. *J. Geophys. Res.*, doi:10.1029/2011JD016074 (2011).

429 ³⁹Dhomse, S. S. *et al.*, Aerosol microphysics simulations of the Mt. Pinatubo eruption with the UM-
430 UKCA composition-climate model. *Atmos. Chem. Phys.*, 14, 11221-11246, doi: 10.5194/acp-14-
431 11221-2014 (2014).

432 ⁴⁰Kirkevåg, A. *et al.*, Aerosol-climate interactions in the Norwegian Earth System Model - NorESM1-
433 M. *Geosci. Model Dev.*, 6, 207-244, doi:10.5194/gmd-6-207-2013 (2013).

434 ⁴¹Schmidt, A. *et al.*, The impact of the 1783–1784 AD Laki eruption on global aerosol formation
435 processes and cloud condensation nuclei. *Atmos. Chem. Phys.*, 10, 6025-6041, doi:10.5194/acp-10-
436 6025-201 (2010).

- ⁴²Zhang, S. *et al.*, On the characteristics of aerosol indirect effect based on dynamic regimes in global climate models. *Atmos. Chem. Phys.*, 16, 2765–2783, doi:10.5194/acp-16-2765-2016 (2016).
- ⁴³Michibata, T., Suzuki, K., Sato, Y., and Takemura, T., The source of discrepancies in aerosol–cloud–precipitation interactions between GCM and A-Train retrievals. *Atmos. Chem. Phys.*, 16, 15413–15424, doi:10.5194/acp-16-15413-2016 (2016).
- ⁴⁴Oreopoulos, L., N. Cho, D. Lee, and S. Kato, Radiative effects of global MODIS cloud regimes. *J. Geophys. Res. Atmos.*, 121, 2299–2317, doi:10.1002/2015JD024502 (2016).
- ⁴⁵Eguchi, K. *et al.*, Modulation of cloud droplets and radiation over the North Pacific by Sulfate Aerosol Erupted from Mount Kilauea. *SOLA*, 7, 77–80, doi:10.2151/sola.2011-020 (2011).
- ⁴⁶Mace, G. G., and A. C. Abernathy, Observational evidence for aerosol invigoration in shallow cumulus downstream of Mount Kilauea. *Geophys. Res. Lett.*, 43, 2981–2988, doi:10.1002/2016GL067830 (2016).
- ⁴⁷Myhre, G. *et al.*, Anthropogenic and Natural Radiative Forcing. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA (2013).
- ⁴⁸Golaz, J.-C., L. W. Horowitz, and H. Levy, Cloud tuning in a coupled climate model: impact on 20th century warming. *Geophys. Res. Lett.*, 40, 2246–2251, doi:10.1002/grl.50232 (2013).
- ⁴⁹Zhou, C. and Penner, J. E.: Why do general circulation models overestimate the aerosol cloud lifetime effect? A case study comparing CAM5 and a CRM. *Atmos. Chem. Phys.*, 17, 21–29, doi:10.5194/acp-17-21-2017 (2017).

List of Supplementary Materials:

SUPPLEMENTARY_INFORMATION.docx

SI-Cloud-Animation.mp4

SI-SO2_animation.mp4

Acknowledgements: JMH, AJ, MD, BTJ, CEJ, JRK and FMOC were supported by the Joint UK BEIS/Defra Met Office Hadley Centre Climate Programme (GA01101). The National Center for Atmospheric Research is sponsored by the U.S. National Science Foundation. SB and LC are respectively Research Fellow and Research Associate funded by F.R.S.-FNRS. PS acknowledges support from the European Research Council (ERC) project ACCLAIM (Grant Agreement FP7-280025). JMH, FFM, DGP and PS were part funded by the UK Natural Environment Research Council project ACID-PRUF (NE/I020148/1). AS was funded by an Academic Research Fellowship from the University of Leeds and a NERC urgency grant NE/M021130/1 (The source and longevity of sulphur in an Icelandic flood basalt eruption plume). RA was supported by the NERC SMURPHS project NE/N006054/1. GWM was funded by the National Centre for Atmospheric Science, one of the UK Natural Environment Research Council's research centres. DPG is funded by the School of Earth and Environment at the University of Leeds. GWM and SD acknowledge additional EU funding from the ERC under the FP7 consortium project MACC-II (grant agreement 283576) and Horizon 2020 project MACC-III (grant agreement 633080). GWM, KSC and DG were also supported through the financial support via the Leeds-Met Office Academic Partnership (ASCI project). The work done with CAM5-Oslo is supported by the Research Council of Norway through the EVA project (grant 229771), NOTUR project nn2345k and NorStore project ns2345k. The following researchers have contributed substantially to the development version of CAM5-Oslo used in this study: Kari Alterskjær, Alf Grini, Matthias Hummel, Trond Iversen, Alf Kirkevåg, Dirk Olivié, Michael Schulz, Øyvind Seland. The AQUA/MODIS MYD08 L3 Global 1 Deg. dataset was acquired from the Level-1 and Atmosphere Archive & Distribution System (LAADS) Distributed Active Archive Center (DAAC), located in the Goddard Space Flight Center in Greenbelt, Maryland (<https://ladsweb.nascom.nasa.gov/>). This work is dedicated to the memory of Jón Egill Kristjánsson who died in a climbing accident in Norway. Jón Egill was a very active, talented and internationally respected researcher, and will be sadly missed.

Author contributions: FFM (Text, processing and analysis of the satellite data and the model results), JMH (Text, analysis of the satellite data and the model results, radiative transfer calculations), AJ, AG, IHHK and JEK (model runs), RA (processing of the CERES data and

contribution to the text), LC and SB (processing of the IASI data and contribution to the text), LO, NC and DL (MODIS cloud regimes), DPG (estimate of CDNC from MODIS data), TT and MEH (provide emission estimates for the 2014-15 eruption at Holuhraun). AJ, NB, OB, KSC, SD, GWM, AS, HC, MD, AAH, BTJ, CEJ, FMOC, DGP, PS, (contribution to the development of UKCA), GM, SP, GLS, HT, JRK (discussion contributing to text and/or help with the MODIS data).

Author Information: The authors declare no competing financial interests. Correspondence and material requests should be addressed to Florent Malavelle (f.malavelle@exeter.ac.uk)

Figure legends:

Figure 1. The column loading of sulphur dioxide. First column: processed data from HadGEM3 masked using positive detections of SO_2 from IASI and spatially and temporally coherent plume data from HadGEM3. Second column: processed data from IASI re-gridded onto the regular HadGEM3 grid. The column loading are expressed in Dobson Units (DU), with 1 DU equivalents to approximately $0.0285 \text{ g}[\text{SO}_2].\text{m}^{-2}$. In each case ‘avg’ represents the average concentration derived within the plume.

Figure 2. Changes in cloud properties detected by MODIS AQUA for October 2014. The mean changes in (a) cloud droplet effective radius (μm) and (c) liquid water path (g.m^{-2}) with corresponding zonal means. The probability distributions of absolute cloud droplet effective radius (b) and liquid water path (d) for the year 2014 (blue) and the 2002-2013 mean (green). Changes correspond to the deviation from the 2002-2013 mean. Stippling in a) and c) represent areas of 95% confidence level significant perturbation based on a two-tailed Student’s t-test. Grey shading in the zonal means represent the standard deviation over 2002-2013.

Figure 3. Changes in cloud properties modelled by HadGEM3 for October 2014. The mean changes in (a) cloud droplet effective radius (μm) and (c) liquid water path (g.m^{-2}) with corresponding zonal means. The probability distributions of absolute cloud droplet effective radius (b) and liquid water path (d) for 2014 including (blue) or excluding (gold) the Holuhraun emissions, and the 2002-2013 mean (green). Changes correspond to the deviation from the 2002-2013 mean. Stippling in a) and c) represent areas of 95% confidence level significant perturbation based on a

two-tailed Student's *t*-test. Grey shading in the zonal means represent the standard deviation over 2002-2013.

Figure 4. Modelled perturbations from HadGEM3 using UKCA during the Sept-Oct 2014 period.

Showing perturbations for a) AOD, b) N_d , c) τ_{eff} , d) LWP, e) τ_{cloud} , and f) Top of Atmosphere (ToA) net SW radiation. Zonal means are shown for the 44°N-80°N, 60°W-30°E analysis region. The shaded regions represent the natural variability in the simulations from 2002-2013. Values outside of the light grey (respectively dark grey, bottom row) shaded regions represent significant perturbations at the 95% (respectively 67%) confidence level based on a two-tailed Student's *t*-test. Red lines represent HOL_{2014} minus NO_HOL_{2014} and blue lines represent HOL_{2014} minus $NO_HOL_{2002-2013}$.

Figure 5. Multi-model estimates of the changes in cloud properties for October 2014. Left column

shows Δr_{eff} (μm) and right column ΔLWP (g.m^{-2}) determined from HadGEM3 using the 2-moment UKCA/GLOMAP-mode aerosol scheme (first row), HadGEM3 using the single moment CLASSIC aerosol scheme (second row) CAM5-NCAR (third row), CAM5-Oslo (fourth row) and AQUA MODIS (last row). Note that MODIS anomalies show the aerosol impacts plus the meteorological variability while the model simulations show the impact of aerosols only (Supplementary S7).