

# Gaussian-type Isoperimetric Inequalities in $\mathrm{RCD}(K, \infty)$ probability spaces for positive $K$

Luigi Ambrosio \*

Andrea Mondino †

July 11, 2018

## Abstract

In this paper we adapt the well-established  $\Gamma$ -calculus techniques to the context of  $\mathrm{RCD}(K, \infty)$  spaces, proving Bobkov’s local isoperimetric inequality [12],[13] and, when  $K$  is positive, the Gaussian isoperimetric inequality in this class of spaces. The proof relies on the measure-valued  $\Gamma_2$  operator introduced by Savaré in [22].

*Keywords:* metric geometry, metric analysis, Isoperimetric Inequality, Gaussian measures, Ricci curvature.

## 1 Introduction

The goal of the paper is to extend to  $\mathrm{RCD}(K, \infty)$  spaces, a class of non-smooth “Riemannian” and possibly infinite-dimensional metric measure spaces with lower Ricci curvature bounds, the Lévy-Gromov Gaussian isoperimetric inequality. Our proof is strongly inspired by the arguments of Bakry-Ledoux [12] (see also the recent monograph [10]), where the authors gave a proof of such an inequality for infinite dimensional diffusion operators via  $\Gamma$ -calculus. This calculus, also according to the presentation in [10], relies on the assumption of the existence of a dense algebra of functions stable under the  $\Gamma$  operator, an assumption that seems to be problematic in the category of  $\mathrm{RCD}(K, \infty)$  spaces. Savaré realized in [22] that, by looking at the iterated  $\Gamma$  operator  $\Gamma_2$  as a measure-valued operator, the assumption on the algebra can be relaxed, and that most techniques of  $\Gamma$ -calculus still apply (heuristically, because the negative part of the tensor  $\Gamma_2$  is still absolutely continuous w.r.t.  $\mathfrak{m}$ ); this point of view has also been exploited in [19] for the development of Hessian estimates and a full theory of covariant derivatives. In this paper we follow the same line of thought to prove Bobkov’s inequality and the Lévy-Gromov Gaussian isoperimetric inequality in sharp form. On the technical side, while the computations are essentially the same (see in particular Chapter 8.5.2 of [10]), we have tried to emphasize the technical issues arising from the not-so-smooth context of  $\mathrm{RCD}(K, \infty)$  spaces, in particular the lack of smoothness of the function  $\sqrt{\mathcal{I}^2(g) + c_\alpha(t)\Gamma(g)}$  appearing in the proof of Bobkov’s inequality.

One of the motivations for writing this note comes from [6], where the isoperimetric result of this paper is used to prove new compactness results in variable Sobolev spaces

---

\*Scuola Normale Superiore di Pisa (Italy) *Email:* luigi.ambrosio@sns.it

†MSRI-Berkeley&Universität Zürich, Institut für Mathematik. *Email:* andrea.mondino@math.uzh.ch

$H^{1,p}(X, \mathbf{d}_i, \mathbf{m}_i)$  relative to  $\text{RCD}(K, \infty)$  metric measure structures  $(X, \mathbf{d}_i, \mathbf{m}_i)$ , in the whole range of exponents  $p \in [1, \infty)$  ( $p = 1$  corresponds to  $BV(X, \mathbf{d}_i, \mathbf{m}_i)$ ).

Let us mention that the isoperimetric inequality for Gaussian measures is originally due to C. Borell [14] and V. Sudakov - B. Cirel'son [25]. The functional form of the Gaussian isoperimetric inequality presented in Theorem 4.1 was introduced by S. Bobkov [13] who established it first on the two-point space and then in the limit for the Gaussian measure by the central limit Theorem. On the basis of the Bobkov inequality, the local inequalities of Theorem 3.1 were established by D. Bakry and M. Ledoux [12] in the context of smooth infinite dimensional spaces with lower Ricci curvature bounds. Theorem 4.2 may be considered as the *non-smooth, infinite-dimensional* extension of a celebrated result of P. Lévy and M. Gromov [20] comparing the isoperimetric profile of a Riemannian manifold with a strictly positive lower bound on the Ricci curvature to that of the sphere with the same (constant) curvature and dimension. Let us mention that the dimensional Lévy-Gromov inequality has been recently extended to non-smooth spaces with lower Ricci curvature bounds in [15].

**Acknowledgment:** Part of the work has been developed while A. M. was lecturer at the Institut für Mathematik at the Universität Zürich and part when he was in residence at the Mathematical Science Research Institute in Berkeley, California, during the spring semester 2016 and was supported by the National Science Foundation under the Grant No. DMS-1440140. He wishes to express his gratitude to both the institutes for the stimulating atmosphere and the excellent working conditions.

## 2 Preliminaries

Throughout the paper  $(X, \mathbf{d}, \mathbf{m})$  will be a metric measure space, m.m.s. for short, i.e.  $(X, \mathbf{d})$  is a complete and separable metric space and  $\mathbf{m}$  is a nonnegative Borel measure. Even if some of the statements of this paper hold in case  $\mathbf{m}$  is a sigma finite measure, for simplicity we will always assume  $\mathbf{m}(X) = 1$  and  $\text{supp}(\mathbf{m}) = X$ . We shall denote by  $\text{LIP}(X)$  the space of Lipschitz functions, by  $\mathcal{P}(X)$  the space of Borel probability measures on the complete and separable metric space  $(X, \mathbf{d})$ . We also denote by

$$\mathcal{P}_2(X) := \left\{ \mu \in \mathcal{P}(X) : \int_X \mathbf{d}^2(x_0, x) \, \mathrm{d}\mu(x) < \infty \text{ for some (and hence all) } x_0 \in X \right\}.$$

the subspace consisting of all the probability measures with finite second moment. The *relative entropy functional*  $\text{Ent}_{\mathbf{m}} : \mathcal{P}_2(X) \rightarrow (-\infty, \infty]$  with respect to a probability measure  $\mathbf{m}$  is defined by

$$\text{Ent}_{\mathbf{m}}(\mu) := \begin{cases} \int_X f \log f \, \mathrm{d}\mathbf{m} & \text{if } \mu = f\mathbf{m}; \\ \infty & \text{otherwise.} \end{cases} \quad (2.1)$$

Note that, by Jensen's inequality, this functional is nonnegative.

A curve  $\gamma : [0, 1] \rightarrow X$  is a *geodesic* if

$$\mathbf{d}(\gamma_s, \gamma_t) = |t - s| \mathbf{d}(\gamma_0, \gamma_1) \quad \forall s, t \in [0, 1]. \quad (2.2)$$

We will denote by  $\text{Geo}(X)$  the space of all constant speed geodesics  $\gamma : [0, 1] \rightarrow X$ , namely  $\gamma \in \text{Geo}(X)$  if (2.2) holds.

## 2.1 Lower Ricci curvature bounds

In the sequel we briefly recall those basic definitions and properties of spaces with lower Ricci curvature bounds that we will need later on.

For  $\mu_0, \mu_1 \in \mathcal{P}_2(X)$  the quadratic transportation distance  $W_2(\mu_0, \mu_1)$  is defined by

$$W_2^2(\mu_0, \mu_1) = \inf_{\gamma} \int_X d^2(x, y) d\gamma(x, y), \quad (2.3)$$

where the infimum is taken over all  $\gamma \in \mathcal{P}(X \times X)$  with  $\mu_0$  and  $\mu_1$  as the first and the second marginal. It turns out that any geodesic  $(\mu_t) \in \text{Geo}(\mathcal{P}_2(X))$  can be lifted to a measure  $\pi \in \mathcal{P}(\text{Geo}(X))$ , in such a way that  $(e_t)_\# \pi = \mu_t$  for all  $t \in [0, 1]$ .

We turn to the formulation of the  $\text{CD}(K, \infty)$  condition, coming from the seminal works of Lott-Villani [21] and Sturm [23].

**Definition 2.1** ( $\text{CD}(K, \infty)$  condition). *Given  $K \in \mathbb{R}$ , we say that a m.m.s.  $(X, d, \mathbf{m})$  is a  $\text{CD}(K, \infty)$ -space if for any two measures  $\mu_0, \mu_1 \in D(\text{Ent}_{\mathbf{m}})$  there exists a geodesic  $(\mu_t) \in \text{Geo}(\mathcal{P}_2(X))$  which satisfies the convexity inequality*

$$\text{Ent}_{\mathbf{m}}(\mu_t) \leq (1-t)\text{Ent}_{\mathbf{m}}(\mu_0) + t\text{Ent}_{\mathbf{m}}(\mu_1) - \frac{K}{2}t(1-t)W_2^2(\mu_0, \mu_1), \quad \text{for all } t \in [0, 1]. \quad (2.4)$$

Notice that the above-mentioned lifting property shows that the  $\text{CD}(K, \infty)$  condition implies that the class  $\text{Geo}(X)$  is sufficiently rich to provide  $\pi \in \mathcal{P}(\text{Geo}(X))$  connecting any two measures with finite entropy. Using this, one can prove that  $X = \text{supp } \mathbf{m}$  is length, i.e.  $d(x, y)$  is the infimum of the length of the curves connecting  $x$  to  $y$  for all  $x, y \in \text{supp } \mathbf{m}$ .

Two crucial properties of the  $\text{CD}(K, \infty)$ -condition are the consistency with the smooth counterpart and the stability under convergence. More precisely, it was shown both in [21] and [23] that a smooth Riemannian manifold satisfies the  $\text{CD}(K, \infty)$ -condition if and only if it has Ricci curvature bounded below by  $K$ . Regarding the stability, it was proved by Lott-Villani [21] that the  $\text{CD}(K, \infty)$ -condition is stable under pointed measured Gromov-Hausdorff convergence of *proper* pointed metric measure spaces; Sturm [23] instead introduced a new distance, denoted with  $\mathbb{D}$ , between normalized metric measure spaces and proved the stability of the  $\text{CD}(K, \infty)$ -condition under such a convergence. The stability in the general case of non-proper and non-normalized spaces was established in [18] where a new notion of convergence, called pointed measured Gromov convergence, was introduced (let us mention that such a convergence coincides with the measured Gromov Hausdorff one in case of uniformly doubling spaces and with the  $\mathbb{D}$ -convergence in case of normalized spaces) and the stability of the  $\text{CD}(K, \infty)$ -condition under such a convergence was established.

## 2.2 The Cheeger energy, its gradient flow and the $\text{RCD}(K, \infty)$ -condition

First of all recall that the metric slope of a Lipschitz function  $f \in \text{LIP}(X)$  is defined by

$$\text{lip } f(x) := \limsup_{y \rightarrow x} \frac{|f(x) - f(y)|}{d(x, y)} \quad (2.5)$$

(with the convention  $\text{lip } f(x) = 0$  if  $x$  is an isolated point). The Cheeger energy (introduced in [16] and further studied in [3]) is defined as the  $L^2$ -lower semicontinuous envelope of the functional  $f \mapsto \frac{1}{2} \int_X \text{lip}^2 f d\mathbf{m}$ , i.e.:

$$\text{Ch}_{\mathbf{m}}(f) := \inf \left\{ \liminf_{n \rightarrow \infty} \frac{1}{2} \int_X \text{lip}^2 f_n d\mathbf{m} : f_n \in \text{LIP}(X), f_n \rightarrow f \text{ in } L^2(X, \mathbf{m}) \right\}. \quad (2.6)$$

If  $\text{Ch}_{\mathbf{m}}(f) < \infty$  it was proved in [16, 3] that the set

$$G(f) := \{g \in L^2(X, \mathbf{m}) : \exists f_n \in \text{LIP}(X), f_n \rightarrow f, \text{lip } f_n \rightarrow h \geq g \text{ in } L^2(X, \mathbf{m})\}$$

is closed and convex, therefore it admits a unique element of minimal norm called *minimal weak upper gradient* and denoted by  $|Df|_w$ . The Cheeger energy can be then represented by integration as

$$\text{Ch}_{\mathbf{m}}(f) := \frac{1}{2} \int_X |Df|_w^2 \, d\mathbf{m}.$$

It is not difficult to see that  $\text{Ch}_{\mathbf{m}}$  is a 2-homogeneous, lower semi-continuous, convex functional on  $L^2(X, \mathbf{m})$ , whose proper domain  $D(\text{Ch}_{\mathbf{m}}) := \{f \in L^2(X, \mathbf{m}) : \text{Ch}_{\mathbf{m}}(f) < \infty\}$  is a dense linear subspace of  $L^2(X, \mathbf{m})$ . It then admits an  $L^2$ -gradient flow which is a continuous semigroup of contractions  $(H_t)_{t \geq 0}$  in  $L^2(X, \mathbf{m})$ , whose continuous trajectories  $t \mapsto H_t f$ , for  $f \in L^2(X, \mathbf{m})$ , are locally Lipschitz curves from  $(0, \infty)$  with values into  $L^2(X, \mathbf{m})$ . We can now define the  $\text{RCD}(K, \infty)$ -condition introduced in [4] (see also [1] for the present simplified axiomatization and for the generalization to  $\sigma$ -finite reference measures).

**Definition 2.2** ( $\text{RCD}(K, \infty)$ -space). *Let  $(X, d, \mathbf{m})$  be a probability metric measure space. We say that  $(X, d, \mathbf{m})$  is an  $\text{RCD}(K, \infty)$ -space if it satisfies the  $\text{CD}(K, \infty)$ -condition, and moreover Cheeger's energy is quadratic, i.e. it satisfies the parallelogram identity*

$$\text{Ch}_{\mathbf{m}}(f + g) + \text{Ch}_{\mathbf{m}}(f - g) = 2\text{Ch}_{\mathbf{m}}(f) + 2\text{Ch}_{\mathbf{m}}(g), \quad \forall f, g \in D(\text{Ch}_{\mathbf{m}}). \quad (2.7)$$

If  $(X, d, \mathbf{m})$  is an  $\text{RCD}(K, \infty)$ -space then the Cheeger energy induces the Dirichlet form  $\mathcal{E}(f) := 2\text{Ch}_{\mathbf{m}}(f)$  which is strongly local and admits the Carré du Champ

$$\Gamma(f) := |Df|_w^2, \quad \text{for every } f \in D(\text{Ch}_{\mathbf{m}}). \quad (2.8)$$

Moreover the sub-differential  $\partial \text{Ch}_{\mathbf{m}}$  is single-valued and coincides with the linear generator  $\Delta$  of the heat flow semi-group  $(H_t)_{t \geq 0}$  defined above.

The  $\text{RCD}(K, \infty)$ -condition was introduced in order to single out in a more precise way, compared to the  $\text{CD}$  theory, non-smooth “Riemannian” structures; indeed, while on the one hand it is possible to give examples of smooth Finsler manifolds satisfying the  $\text{CD}(K, \infty)$ -condition, on the other hand it is not difficult to see that if a Finsler manifold satisfies (2.7) then it is actually Riemannian. Let us finally mention that the stability of the  $\text{RCD}(K, \infty)$  condition was proved in [4] for normalized metric measure spaces with respect to the  $\mathbb{D}$ -convergence of Sturm, and the general case of pointed-Gromov convergence of spaces endowed with  $\sigma$ -finite reference measures was settled in [18].

### 2.3 The $\text{BE}(K, \infty)$ -condition and its self-improvement

We set by definition  $\mathbb{V} := D(\text{Ch}_{\mathbf{m}})$  and we endow such space with the norm

$$\|f\|_{\mathbb{V}} := \sqrt{\|f\|_{L^2(X, \mathbf{m})}^2 + 2\text{Ch}_{\mathbf{m}}(f)}.$$

If  $(X, d, \mathbf{m})$  is an  $\text{RCD}(K, \infty)$ -space then  $\mathbb{V}$  is a Hilbert space, while it is not difficult to see that in a general m.m.s.  $\mathbb{V}$  is a Banach space (indeed (2.7) is equivalent to require that  $\mathbb{V}$  is Hilbert). Moreover, still under the  $\text{RCD}(K, \infty)$  assumption, the very construction of  $\text{Ch}_{\mathbf{m}}$  and

the fact that  $\text{Ch}_m$  is quadratic imply that  $\text{LIP}(X) \cap L^\infty(X, m)$  is dense in  $\mathbb{V}$  ([4, Proposition 4.10]).

The Laplace operator, introduced above as the infinitesimal generator of the Heat semi-group, can be defined also via integration by parts as the operator  $-\Delta : \mathbb{V} \rightarrow \mathbb{V}'$  given by

$$\int_X (-\Delta f) g \, dm := \int_X \Gamma(f, g) dm, \quad \forall f, g \in \mathbb{V}, \quad (2.9)$$

and it is an unbounded self-adjoint non-negative operator with domain  $D(\Delta)$ . Let us also introduce the functional spaces

$$D_{\mathbb{V}}(\Delta) = \{f \in \mathbb{V} : \Delta f \in \mathbb{V}\} \quad (2.10)$$

and

$$D_{L^\infty}(\Delta) = \{f \in \mathbb{V} \cap L^\infty(X, m) : \Delta f \in L^\infty(X, m)\}. \quad (2.11)$$

If  $(X, d, m)$  is an  $\text{RCD}(K, \infty)$ -space, the multilinear form  $\Gamma_2 : D_{\mathbb{V}}(\Delta) \times D_{\mathbb{V}}(\Delta) \times D_{L^\infty}(\Delta) \rightarrow \mathbb{R}$  is defined by

$$\Gamma_2[f, g; \varphi] := \frac{1}{2} \int_X [\Gamma(f, g) \Delta \varphi - (\Gamma(f, \Delta g) + \Gamma(g, \Delta f)) \varphi] \, dm.$$

When  $f = g$  we set

$$\Gamma_2[f; \varphi] := \Gamma_2[f, f; \varphi] := \int_X \left( \frac{1}{2} \Gamma(f) \Delta \varphi - \Gamma(f, \Delta g) \varphi \right) dm,$$

so that

$$\Gamma_2[f, g; \varphi] = \frac{1}{4} \Gamma_2[f + g; \varphi] - \frac{1}{4} \Gamma_2[f - g; \varphi].$$

The  $\Gamma_2$  operator provides a weak version of the Bakry-Émery condition [11] (see also [9] and the recent monograph [10]). More precisely it was proved in [4] (see also [1] for the  $\sigma$ -finite case) that  $\text{RCD}(K, \infty)$  implies the Bakry-Émery condition  $\text{BE}(K, \infty)$ :

**Theorem 2.3.** *Let  $(X, d, m)$  be an  $\text{RCD}(K, \infty)$ -space, then  $\text{BE}(K, \infty)$ -condition holds, i.e.*

$$\Gamma_2[f; \varphi] \geq K \int_X \Gamma(f) \varphi \, dm, \quad \text{for every } (f, \varphi) \in D_{\mathbb{V}}(\Delta) \times D_{L^\infty}(\Delta), \varphi \geq 0 \text{ } m\text{-a.e.} \quad (2.12)$$

Under natural regularity assumptions on the distance canonically associated to the Dirichlet form it is also to prove the converse implication, i.e. from  $\text{BE}(K, \infty)$  to  $\text{RCD}(K, \infty)$ , see [5] (see also [17] and [8] for the dimensional case).

A crucial property of the  $\text{BE}(K, \infty)$  condition that will be used later in the paper, is the self-improvement established by Savaré [22, § 3] in the present setting.

**Theorem 2.4.** *Let  $(X, d, m)$  be an  $\text{RCD}(K, \infty)$ -space.*

(1) *For every  $f \in D_{\mathbb{V}}(\Delta) \cap \text{LIP}(X)$  we have  $\Gamma(f) \in \mathbb{V}$  with*

$$\text{Ch}_m(\Gamma(f)) \leq - \int_X \left( 2K\Gamma(f)^2 + 2\Gamma(f)\Gamma(f, \Delta f) \right) dm. \quad (2.13)$$

(2)  $D_{\mathbb{V}}(\Delta) \cap \text{LIP}(X)$  is an algebra (i.e. closed w.r.t. pointwise multiplication) and, more generally, if  $\mathbf{f} = (f_i)_{i=1}^n \in (D_{\mathbb{V}}(\Delta) \cap \text{LIP}(X))^n$  then  $\Phi(\mathbf{f}) \in D_{\mathbb{V}}(\Delta) \cap \text{LIP}(X)$  for every smooth function  $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}$  with  $\Phi(0) = 0$ .

(3) If  $f \in D_{\mathbb{V}}(\Delta) \cap \text{LIP}(X)$ , then the linear functional

$$\varphi \in \text{LIP}(X) \cap L^\infty(X, \mathbf{m}) \mapsto - \int_X \Gamma(\varphi, \Gamma(f)) \, d\mathbf{m}$$

can be represented by a signed Borel measure  $\Delta^* \Gamma(f)$  which can be extended to a unique element in  $\mathbb{V}'$  (by integration of the  $\text{Ch}_{\mathbf{m}}$ -quasicontinuous representative) and, by defining

$$\Gamma_{2,K}^*[f] := \frac{1}{2} \Delta^* \Gamma(f) - \left( \Gamma(f, \Delta f) + K \Gamma(f) \right) \mathbf{m}, \quad (2.14)$$

the measure  $\Gamma_{2,K}^*[f]$  is nonnegative and satisfies

$$\Gamma_{2,K}^*[f](X) \leq \int_X \left( (\Delta f)^2 - K \Gamma(f) \right) \, d\mathbf{m}. \quad (2.15)$$

(4) There exists a continuous, symmetric and bilinear map

$$\gamma_{2,K} : (D_{\mathbb{V}}(\Delta) \cap \text{LIP}(X))^2 \rightarrow L^1(X, \mathbf{m})$$

such that for every  $f \in D_{\mathbb{V}}(\Delta) \cap \text{LIP}(X)$  one has

$$\Gamma_{2,K}^*[f] = \gamma_{2,K}[f, f] \mathbf{m} + \Gamma_{2,K}^\perp[f], \quad \text{with} \quad \Gamma_{2,K}^\perp[f] \geq 0, \quad \Gamma_{2,K}^\perp[f] \perp \mathbf{m}. \quad (2.16)$$

Setting  $\gamma_{2,K}[f] := \gamma_{2,K}[f, f] \geq 0$ , one has for every  $f \in D_{\mathbb{V}}(\Delta) \cap \text{LIP}(X)$

$$\Gamma(\Gamma(f)) \leq 4\gamma_{2,K}[f] \Gamma(f) \quad \mathbf{m}\text{-a.e. in } X. \quad (2.17)$$

Notice that the measures  $\Gamma_{2,K}^*[f]$ ,  $K \in \mathbb{R}$ , just differ by a multiple of  $\Gamma(f) \mathbf{m}$ , so the (non-negative) singular part in the Lebesgue decomposition (2.16) is independent of  $K$ . For more results about functional inequalities and regularity in  $\text{BE}(K, \infty)$ -spaces the interested reader is referred to [5] and [7].

## 2.4 Improved regularity of the heat flow in $\text{RCD}(K, \infty)$ -spaces

In  $\text{RCD}(K, \infty)$ -spaces the heat flow  $\mathbf{H}_t$  can be viewed in two conceptually different, but consistent, ways: the first one, as in the theory of Dirichlet form and in the more general theory of gradient flows of lower semicontinuous functionals in Hilbert spaces, is based on  $\mathbf{H}_t$  as the  $L^2(X, \mathbf{m})$  gradient flow of  $\text{Ch}_{\mathbf{m}}$ ; the second way looks at  $\mathbf{H}_t$  as the gradient flow  $\mathcal{H}_t$  of the Shannon entropy functional in the Wasserstein space. It has been proved in full generality in [3] that the two points of view coincide in the common domain of probability densities in  $L^1(X, \mathbf{m})$ . In addition, using this identification, in [4] several regularity properties of  $\mathbf{H}_t$ , besides self-adjointness, Markov, mass-preserving, have been deduced. We recall some of them. When  $f \in L^\infty(X, \mathbf{m})$ , for all  $t > 0$  the function  $\mathbf{H}_t f$  has a continuous representative, denoted by  $\tilde{\mathbf{H}}_t f$ , which is defined as follows (see Theorem 6.1 in [4])

$$\tilde{\mathbf{H}}_t f := \int_X f \, d\mathcal{H}_t(\delta_x). \quad (2.18)$$

Moreover, for each  $f \in L^\infty(X, \mathbf{m})$  the map  $(t, x) \mapsto \tilde{\mathbf{H}}_t f(x)$  belongs to  $C_b((0, \infty) \times X)$ . Finally, for all  $f \in L^\infty(X, \mathbf{m})$  with  $\Gamma(f) \in L^\infty(X, \mathbf{m})$  the classical Bakry-Émery gradient estimate holds even in the pointwise form (see Theorem 6.2 in [4])

$$\text{lip } \tilde{\mathbf{H}}_t f \leq e^{-2Kt} \tilde{\mathbf{H}}_t(\Gamma(f)) \quad \text{in } X = \text{supp } \mathbf{m}, \text{ for all } t > 0, \quad (2.19)$$

while if we drop the boundedness assumption on  $\Gamma(f)$  one has (with  $\text{Lip}$  denoting the global Lipschitz constant)

$$\text{Lip}(\tilde{\mathbf{H}}_t f) \leq \frac{1}{\sqrt{2l_{2K}(t)}} \|f\|_{L^\infty(X, \mathbf{m})} \quad \text{with } l_{2K}(t) = \frac{e^{2Kt} - 1}{2K}. \quad (2.20)$$

For functions  $f \in \mathbb{V}$ , instead, one has the traditional form of  $\Gamma$ -calculus

$$\Gamma(\mathbf{H}_t f) \leq e^{-2Kt} \mathbf{H}_t(\Gamma(f)) \quad \mathbf{m}\text{-a.e.}, \text{ for all } t \geq 0. \quad (2.21)$$

From now on, when  $f \in L^\infty(X, \mathbf{m})$ , we will always choose  $\tilde{\mathbf{H}}_t f$  as representative of  $\mathbf{H}_t f$  and we will simply write  $\mathbf{H}_t f$ .

An immediate consequence of (2.19), (2.21) and of the length property of  $X = \text{supp } \mathbf{m}$  is that the Dirichlet form induced by  $\text{Ch}_\mathbf{m}$  is irreducible, i.e.  $\text{Ch}_\mathbf{m}(f) = 0$  implies that  $f$  is equivalent to a constant (this holds because all functions  $\tilde{\mathbf{H}}_t f$ ,  $t > 0$ , have null slope in  $X$  and therefore are  $\mathbf{m}$ -equivalent to a constant). Then, since irreducibility implies ergodicity (see for instance [10, Section 3.8]) we get

$$\lim_{t \rightarrow +\infty} \mathbf{H}_t f = \int_X f \, d\mathbf{m} \quad \text{in } L^2(X, \mathbf{m}), \text{ for all } f \in L^2(X, \mathbf{m}). \quad (2.22)$$

The following lemma will be useful in the proof of the Bobkov inequality.

**Lemma 2.5.** *Let  $(X, d, \mathbf{m})$  be an  $\text{RCD}(K, \infty)$  probability space. Then, for every fixed  $f \in L^2(X, \mathbf{m})$ , the map  $\mathbf{H}_{(\cdot)} f : (0, T] \rightarrow \mathbb{V}$  is locally Lipschitz, continuous up to  $t = 0$  if  $f \in \mathbb{V}$ , with also  $t \mapsto \Gamma(\mathbf{H}_t f)$  locally Lipschitz in  $(0, T]$  as a  $L^1(X, \mathbf{m})$ -valued map, and*

$$\lim_{s \rightarrow t} \frac{\Gamma(\mathbf{H}_s f) - \Gamma(\mathbf{H}_t f)}{s - t} = \Gamma(\Delta \mathbf{H}_t f, f) \quad \text{in } L^1(X, \mathbf{m}) \text{ for } \mathcal{L}^1\text{-a.e. } t > 0. \quad (2.23)$$

Moreover, if  $f \in L^\infty(X, \mathbf{m})$ , then for every  $t > 0$  we have  $\mathbf{H}_t f \in D_\mathbb{V}(\Delta) \cap \text{LIP}(X)$ .

*Proof.* The proof of the continuity of  $\mathbf{H}_t f$  and the  $\Delta$ -regularity of  $\mathbf{H}_t f$  follow immediately by the regularization estimates provided by the theory of gradient flows in Hilbert spaces (see for instance [2]), namely

$$\begin{aligned} \text{Ch}_\mathbf{m}(\mathbf{H}_t f) &\leq \inf_{g \in \mathbb{V}} \frac{\|g - f\|_{L^2(X, \mathbf{m})}^2}{2t}, \\ \|\Delta \mathbf{H}_t f\|_2^2 &\leq \inf_{g \in D(\Delta)} \frac{\|g - f\|_{L^2(X, \mathbf{m})}^2}{t^2}, \end{aligned}$$

using also the semigroup property and the commutation  $\mathbf{H}_t \circ \Delta = \Delta \circ \mathbf{H}_t$  for  $t > 0$ . Formula (2.23) is a simple consequence of  $\Gamma(\mathbf{H}_s f) - \Gamma(\mathbf{H}_t f) = \Gamma(\mathbf{H}_s f - \mathbf{H}_t f, \mathbf{H}_s f + \mathbf{H}_t f)$  and of the differentiability of  $t \mapsto \mathbf{H}_t f$  as a  $\mathbb{V}$ -valued map.

Finally, if  $f \in L^\infty(X, \mathbf{m})$  the Lipschitz regularity of  $\mathbf{H}_t f$  follows directly from (2.20).  $\square$

Thanks to Lemma 2.5 we will be able to apply Theorem 2.4 to  $\mathbf{H}_t f$  for  $f \in L^\infty(X, \mathbf{m})$ ,  $t > 0$ .

## 2.5 The Gaussian isoperimetric profile

Let  $H(r) := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^r e^{-x^2/2} dx$ ,  $r \in \mathbb{R}$ , be the distribution function of the standard Gaussian measure on the real line and set  $h := H'$  to be its density with respect to the Lebesgue measure  $\mathcal{L}^1$ . Then

$$\mathcal{J} := h \circ H^{-1} : [0, 1] \rightarrow \left[0, \frac{1}{\sqrt{2\pi}}\right] \quad (2.24)$$

defines the *Gaussian isoperimetric profile* (in any dimension). Note that the function  $\mathcal{J}$  is concave, continuous, symmetric with respect to  $1/2$ , it satisfies  $\mathcal{J}(0) = \mathcal{J}(1) = 0$  and the fundamental differential equation  $\mathcal{J}\mathcal{J}'' = -1$ .

## 3 Local Bobkov inequality in $\text{RCD}(K, \infty)$ -spaces, $K \in \mathbb{R}$

The goal of this section is to prove the following result.

**Theorem 3.1** (Local Bobkov Inequality). *Let  $(X, d, \mathbf{m})$  be an  $\text{RCD}(K, \infty)$  probability space for some  $K \in \mathbb{R}$ . Then, for every  $f \in \text{LIP}(X)$  with values in  $[0, 1]$ , every  $\alpha \geq 0$  and every  $t \geq 0$ , it holds*

$$\sqrt{\mathcal{J}^2(\mathbf{H}_t f) + \alpha \Gamma(\mathbf{H}_t f)} \leq \mathbf{H}_t \left( \sqrt{\mathcal{J}^2(f) + c_\alpha(t) \Gamma(f)} \right), \quad \mathbf{m}\text{-a.e.}, \quad (3.1)$$

where, for  $t \geq 0$ , we have set

$$c_\alpha(t) := \frac{1 - e^{-2Kt}}{K} + \alpha e^{-2Kt}, \quad \text{if } K \neq 0; \quad c_\alpha(t) := 2t + \alpha, \quad \text{if } K = 0. \quad (3.2)$$

In the next lemma we isolate a key computation for proving Theorem 3.1.

**Lemma 3.2.** *Let  $(X, d, \mathbf{m})$  be an  $\text{RCD}(K, \infty)$  probability space for some  $K \in \mathbb{R}$ . Let  $\Psi(t, u, v) : \mathbb{R}^3 \rightarrow \mathbb{R}$  be a function of class  $C^4$  with  $\partial_v \Psi \geq 0$ , fix  $T > 0$ ,  $f \in \text{LIP}(X) \cap L^\infty(X, \mathbf{m})$ ,  $\varphi \in L^\infty(X, \mathbf{m})$  with  $\varphi \geq 0$   $\mathbf{m}$ -a.e., and set*

$$\Phi(t) := \int_X \mathbf{H}_t(\Psi(t, \mathbf{H}_{T-t}(f), \Gamma(\mathbf{H}_{T-t}(f)))) \varphi \, d\mathbf{m}, \quad \forall t \in [0, T]. \quad (3.3)$$

Then  $\Phi$  is continuous in  $[0, T]$ , locally Lipschitz in  $(0, T)$ , and for  $\mathcal{L}^1$ -a.e.  $t_0 \in (0, T)$  it holds

$$\frac{d}{dt} \Big|_{t=t_0} \Phi(t) \geq \int_X \zeta \, \mathbf{H}_{t_0} \varphi \, d\mathbf{m}, \quad (3.4)$$

where, denoting  $g := \mathbf{H}_{T-t_0} f$  and writing simply  $\Psi$  in place of  $\Psi(t_0, g, \Gamma(g))$ , we have set

$$\zeta := \partial_t \Psi + \partial_u^2 \Psi \Gamma(g) + 2 \partial_u \partial_v \Psi \Gamma(g, \Gamma(g)) + \partial_v^2 \Psi \Gamma(\Gamma(g)) + 2K \partial_v \Psi \Gamma(g) + 2 \partial_v \Psi \gamma_{2,K}[g]. \quad (3.5)$$

*Proof. Step 1.*  $\Phi : [0, T] \rightarrow \mathbb{R}$  is locally Lipschitz in  $(0, T)$ , continuous up to  $t = 0, T$ , and Leibniz rule.

To this aim recall that, in virtue of Lemma 2.5, for every fixed  $f \in L^2(X, \mathbf{m})$  the heat flow  $\mathbf{H}_{(\cdot)} f : (0, T] \rightarrow \mathbb{V}$  is locally Lipschitz. Moreover, since by assumption  $f \in \text{LIP}(X) \cap L^\infty(X, \mathbf{m})$ , on the one hand by maximum principle it holds  $\|\mathbf{H}_t f\|_{L^\infty(X, \mathbf{m})} \leq \|f\|_{L^\infty(X, \mathbf{m})}$  and on the other hand, in virtue of (2.21), it holds  $\mathbf{H}_t f \in \text{LIP}(X)$  with uniform Lipschitz bounds for



$t \in [0, T]$ . It follows that, for a fixed  $f \in \text{LIP}(X) \cap L^\infty(X, \mathbf{m})$ , the pair  $(\mathbf{H}_{T-t}f, \Gamma(\mathbf{H}_{T-t}f))$  is essentially bounded with values in  $\mathbb{R}^2$  for  $t \in [0, T]$ . Therefore  $\Psi$  is Lipschitz on the range of  $(t, \mathbf{H}_{T-t}f, \Gamma(\mathbf{H}_{T-t}f))$  for  $t \in [0, T]$ , and thus

$$[0, T] \ni t \mapsto G(t) := \Psi(t, \mathbf{H}_{T-t}f, \Gamma(\mathbf{H}_{T-t}f)) \in L^1(X, \mathbf{m}) \text{ is locally Lipschitz in } [0, T],$$

taking the local Lipschitz property of  $t \mapsto \Gamma(\mathbf{H}_t f)$  as a  $L^1(X, \mathbf{m})$ -valued map into account. In addition, (2.23) and a standard chain rule provide the existence for  $\mathcal{L}^1$ -a.e.  $t_0 \in (0, T)$  of the strong  $L^1(X, \mathbf{m})$  derivative of  $G$ , given by

$$G'(t_0) = \partial_t \Psi - \partial_u \Psi \Delta g - 2\partial_v \Psi \Gamma(g, \Delta g), \quad (3.6)$$

with  $g = \mathbf{H}_{T-t_0}f$  and  $\Psi = \Psi(t_0, g, \Gamma g)$ . Notice also that, by gradient contractivity,  $\|G\|_{L^\infty(X, \mathbf{m})}$  is bounded in  $[0, T]$ , hence  $G$  is continuous in  $(0, T]$  as a  $L^2(X, \mathbf{m})$ -valued map. Using that  $\mathbf{H}_t$  is self-adjoint, we can write  $\Phi(t) = \int_X G(t) \mathbf{H}_t \varphi \, d\mathbf{m}$ , and writing

$$\Phi(t) - \Phi(t_0) = \int_X (G(t) - G(t_0)) \mathbf{H}_{t_0} \varphi \, d\mathbf{m} + \int_X (\mathbf{H}_t \varphi - \mathbf{H}_{t_0} \varphi) G(t) \, d\mathbf{m}$$

we can use the  $L^2(X, \mathbf{m})$  continuity of  $G$  to obtain the Leibniz rule

$$\Phi'(t_0) = \int_X (\partial_t \Psi - \partial_u \Psi \Delta g - 2\partial_v \Psi \Gamma(g, \Delta g)) \mathbf{H}_{t_0} \varphi \, d\mathbf{m} + \int_X G(t_0) \Delta \mathbf{H}_{t_0} \varphi \, d\mathbf{m} \quad (3.7)$$

for  $\mathcal{L}^1$ -a.e.  $t_0 \in (0, T)$ . This proves also the local Lipschitz property of  $\Phi$  in  $(0, T)$ . Continuity up to  $t = 0, T$  follows by (2.21) and the assumption  $f \in \text{LIP}(X)$ .

**Step 2.** An intermediate approximation.

In this intermediate step, for  $t \in (0, T)$  fixed, first we compute by chain rule the laplacian of  $\Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g)))$  and then show that  $\Delta \Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g)))$  admits an explicit (weak) limit as  $\varepsilon \downarrow 0$  which formally can be considered as  $\Delta^* \Psi(t, g, \Gamma(g))$ ; the expression of the (weak) limit will be then used in Step 3.

Occasionally we shall use the notation  $\Psi_\varepsilon$  for  $\Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g)))$ ,  $\Psi$  for  $\Psi(t, g, \Gamma(g))$  and analogous ones for  $\Psi_u$ ,  $\Psi_v$  and for second order partial derivatives.

Since we know that  $g := \mathbf{H}_{T-t}(f) \in \text{LIP}(X)$ , thus  $\Gamma(g) \in L^\infty(X, \mathbf{m})$  and  $\mathbf{H}_\varepsilon(\Gamma(g)) \in L^\infty(X, \mathbf{m}) \cap \text{LIP}(X) \cap \mathbf{D}(\Delta)$  for all  $\varepsilon > 0$ . Therefore, by standard chain rule (see for instance [22, (2.4)]) we get that  $\Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g))) \in \text{LIP}(X) \cap \mathbf{D}(\Delta)$  with

$$\begin{aligned} \Delta \Psi_\varepsilon &= \partial_u \Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g))) \Delta g + \partial_u^2 \Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g))) \Gamma(g) \\ &\quad + 2\partial_u \partial_v \Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g))) \Gamma(g, \mathbf{H}_\varepsilon(\Gamma(g))) + \partial_v^2 \Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g))) \Gamma(\mathbf{H}_\varepsilon(\Gamma(g))) \\ &\quad + \partial_v \Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g))) \Delta \mathbf{H}_\varepsilon(\Gamma(g)). \end{aligned} \quad (3.8)$$

We now pass to the limit as  $\varepsilon \downarrow 0$  in the last formula. To this aim observe that by (2.17) we know that  $\Gamma(g) \in \mathbb{V}$ , so that  $\mathbf{H}_\varepsilon(\Gamma(g)) \rightarrow \Gamma(g)$  in  $\mathbb{V}$  and therefore, since  $\Psi, \partial \Psi, \partial^2 \Psi$  are Lipschitz on the range of  $(t, g, \mathbf{H}_\varepsilon(\Gamma(g)))$  uniformly for  $\varepsilon \in [0, 1]$  it follows that

$$\begin{aligned} \Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g))) &\rightarrow \Psi(t, g, \Gamma(g)), \quad \partial \Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g))) \rightarrow \partial \Psi(t, g, \Gamma(g)), \\ \partial^2 \Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g))) &\rightarrow \partial^2 \Psi(t, g, \Gamma(g)) \quad \text{in } \mathbb{V}, \end{aligned} \quad (3.9)$$

where for brevity we wrote  $\partial\Psi, \partial^2\Psi$  in place of  $\partial_u\Psi, \partial_v\Psi, \partial_u^2\Psi, \partial_v^2\Psi, \partial_u\partial_v\Psi$ . Therefore the first two lines of (3.8) pass to the limit in  $L^1(X, \mathbf{m})$  topology as  $\varepsilon \downarrow 0$ :

$$\begin{aligned} & \partial_u\Psi_\varepsilon \Delta g + \partial_u^2\Psi_\varepsilon \Gamma(g) + 2\partial_u\partial_v\Psi_\varepsilon \Gamma(g, \mathbf{H}_\varepsilon(\Gamma(g))) + \partial_v^2\Psi_\varepsilon \Gamma(\mathbf{H}_\varepsilon(\Gamma(g))) \\ \rightarrow & \partial_u\Psi \Delta g + \partial_u^2\Psi \Gamma(g) + 2\partial_u\partial_v\Psi \Gamma(g, \Gamma(g)) + \partial_v^2\Psi \Gamma(\Gamma(g)) \quad \text{in } L^1(X, \mathbf{m}). \end{aligned} \quad (3.10)$$

Regarding the convergence of the last line of (3.8), observe that  $\Delta^*\Gamma(g) \in \mathbb{V}'$ , so that  $\Delta\mathbf{H}_\varepsilon\Gamma(g)\mathbf{m} \rightarrow \Delta^*\Gamma(g)$  weakly in  $\mathbb{V}'$  topology as  $\varepsilon \downarrow 0$ . Therefore, the combination with (3.9) gives

$$\langle \partial_v\Psi_\varepsilon, \Delta\mathbf{H}_\varepsilon(\Gamma(g))\mathbf{m} \rangle_{\mathbb{V}, \mathbb{V}'} \rightarrow \langle \partial_v\Psi, \Delta^*\Gamma(g) \rangle_{\mathbb{V}, \mathbb{V}'}. \quad (3.11)$$

Combining (3.8), (3.10) and (3.11) we conclude that for every  $\varphi \in \text{LIP}(X) \cap L^\infty(X, \mathbf{m})$  it holds

$$\begin{aligned} \int_X \varphi \Delta\Psi_\varepsilon \, d\mathbf{m} & \rightarrow \int_X [\partial_u\Psi \Delta g + \partial_u^2\Psi \Gamma(g) + 2\partial_u\partial_v\Psi \Gamma(g, \Gamma(g)) + \partial_v^2\Psi \Gamma(\Gamma(g))] \varphi \, d\mathbf{m} \\ & + \langle \varphi \partial_v\Psi, \Delta^*\Gamma(g) \rangle_{\mathbb{V}, \mathbb{V}'}, \quad \text{as } \varepsilon \downarrow 0. \end{aligned} \quad (3.12)$$

**Step 3.** Formula (3.4) holds.

Set

$$\Phi_\varepsilon(t) := \int_X \Psi(t, \mathbf{H}_{T-t}(f), \mathbf{H}_\varepsilon(\Gamma(\mathbf{H}_{T-t}f))) \mathbf{H}_t(\varphi) \, d\mathbf{m}.$$

By analogous arguments of Step 1, we get that  $\Phi_\varepsilon$  is locally Lipschitz on  $(0, T)$  with

$$\begin{aligned} \Phi'_\varepsilon(t_0) &= \int_X [\partial_t\Psi(t_0, g, \mathbf{H}_\varepsilon(\Gamma(g))) - \partial_u\Psi(t_0, g, \mathbf{H}_\varepsilon(\Gamma(g))) \Delta g] \mathbf{H}_{t_0}\varphi \, d\mathbf{m} \\ &\quad - 2 \int_X \partial_v\Psi(t_0, g, \mathbf{H}_\varepsilon(\Gamma(g))) \mathbf{H}_\varepsilon(\Gamma(g, \Delta g)) \mathbf{H}_{t_0}\varphi \, d\mathbf{m} \\ &\quad + \int_X \Psi(t_0, g, \mathbf{H}_\varepsilon(\Gamma(g))) \Delta \mathbf{H}_{t_0}\varphi \, d\mathbf{m}, \end{aligned} \quad (3.13)$$

for  $\mathcal{L}^1$ -a.e.  $t_0 \in (0, T)$ . Combining (3.7), (3.9) and (3.13) we infer that, given a sequence  $\varepsilon_n \downarrow 0$  it holds

$$\Phi'_{\varepsilon_n}(t_0) \rightarrow \Phi'(t_0), \quad \text{for } \mathcal{L}^1\text{-a.e. } t_0 \in (0, T). \quad (3.14)$$

By Step 2 we know that  $\Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g))) \in \text{LIP}(X) \cap \mathbf{D}(\Delta)$  with  $\Delta\Psi(t, g, \mathbf{H}_\varepsilon(\Gamma(g)))$  given by (3.8); therefore we can integrate by parts the laplacian in the last integral in (3.13) and use (3.9), (3.12) in order to pass to the limit as  $\varepsilon_n \downarrow 0$ :

$$\begin{aligned} \Phi'(t_0) &= \lim_{n \rightarrow \infty} \Phi'_{\varepsilon_n}(t_0) \\ &= \int_X (\partial_t\Psi - \partial_u\Psi \Delta g - 2\partial_v\Psi \Gamma(g, \Delta g)) \mathbf{H}_{t_0}\varphi \, d\mathbf{m} \\ &\quad + \int_X (\partial_u\Psi \Delta g + \partial_u^2\Psi \Gamma(g) + 2\partial_u\partial_v\Psi \Gamma(g, \Gamma(g)) + \partial_v^2\Psi \Gamma(\Gamma(g))) \mathbf{H}_{t_0}\varphi \, d\mathbf{m} \\ &\quad + \int_X \partial_v\Psi \mathbf{H}_{t_0}\varphi \, d\Delta^*\Gamma(g), \quad \text{for } \mathcal{L}^1\text{-a.e. } t_0 \in (0, T), \end{aligned} \quad (3.15)$$

where in the first line we used (3.14) and we wrote simply  $\Psi$  in place of  $\Psi(t_0, g, \Gamma(g))$ .

Recall that, thanks to Theorem 2.4,

$$\frac{1}{2}\Delta^*\Gamma(g) = \left(\Gamma(g, \Delta g) + K\Gamma(g)\right)\mathbf{m} + \Gamma_{2,K}^\perp[g] + \gamma_{2,K}[g]\mathbf{m},$$

where  $\Gamma_{2,K}^\perp[g]$  is nonnegative and singular with respect to  $\mathbf{m}$  and  $\gamma_{2,K}[g]\mathbf{m}$  is the nonnegative absolutely continuous part. Since  $\mathbf{H}_{t_0}\varphi$  is nonnegative by the minimum principle of the heat flow and  $\partial_v\Psi \geq 0$  by assumption, neglecting the contribution of the singular part the thesis follows.  $\square$

### Proof of Theorem 3.1

**Step 1.** It is enough to show the validity of (3.1) for every  $f \in \text{LIP}(X)$  taking values in  $[\varepsilon, 1 - \varepsilon]$ , for some  $\varepsilon \in (0, 1/2)$ .

Let  $f \in \text{LIP}(X)$  taking values in  $[0, 1]$  and for every  $\varepsilon \in (0, 1/2)$  define  $f_\varepsilon$  with values in  $[\varepsilon, 1 - \varepsilon]$  by truncation as

$$f_\varepsilon := \max(\min(f, 1 - \varepsilon), \varepsilon).$$

In this first step we show that, assuming the validity of (3.1) for the truncated  $f_\varepsilon$ , then we can pass into the limit and get the validity of (3.1) also for  $f$ . To this aim, first observe that since the truncation satisfies  $\Gamma(f_\varepsilon) \leq \Gamma(f)$  which is essentially bounded by the assumption  $f \in \text{LIP}(X)$ , and since  $\mathcal{I}$  takes values into  $[0, \frac{1}{\sqrt{2\pi}}]$ , then there exists  $C > 0$  depending on  $f$  but not on  $\varepsilon$  such that

$$\sqrt{\mathcal{I}^2(f_\varepsilon) + c_\alpha(t)\Gamma(f_\varepsilon)} \leq C \quad \mathbf{m}\text{-a.e.}, \quad \forall \varepsilon \in (0, 1/2). \quad (3.16)$$

Moreover it is readily seen that  $f_\varepsilon \rightarrow f$  in  $\mathbb{V}$ -topology and then, since  $\mathbf{H}_t(\cdot)$  is continuous as map from  $\mathbb{V}$  to  $\mathbb{V}$ , also  $\mathbf{H}_t f_\varepsilon \rightarrow \mathbf{H}_t f$  in  $\mathbb{V}$ -topology, as  $\varepsilon \rightarrow 0$ . In particular, for every fixed  $t \geq 0$ , there exists a sequence  $\varepsilon_n \downarrow 0$  such that

$$f_{\varepsilon_n} \rightarrow f, \quad \mathbf{H}_t f_{\varepsilon_n} \rightarrow \mathbf{H}_t f, \quad \Gamma(f_{\varepsilon_n}) \rightarrow \Gamma(f), \quad \Gamma(\mathbf{H}_t f_{\varepsilon_n}) \rightarrow \Gamma(\mathbf{H}_t f), \quad \mathbf{m}\text{-a.e.} \quad (3.17)$$

as  $n \rightarrow \infty$ . We can then pass into the limit  $\mathbf{m}$ -a.e. as  $n \rightarrow \infty$  in the left hand side of (3.1). In order to pass into the limit  $\mathbf{m}$ -a.e. also in the right hand side of (3.1) observe that thanks to (3.16) and (3.17) we can apply the Dominated Convergence Theorem to infer that

$$\sqrt{\mathcal{I}^2(f_{\varepsilon_n}) + c_\alpha(t)\Gamma(f_{\varepsilon_n})} \rightarrow \sqrt{\mathcal{I}^2(f) + c_\alpha(t)\Gamma(f)} \quad \text{in } L^2(X, \mathbf{m})\text{-strong topology, as } n \rightarrow \infty.$$

Using again the continuity of the heat flow  $\mathbf{H}_t(\cdot)$  as map from  $L^2(X, \mathbf{m})$  to  $L^2(X, \mathbf{m})$  we conclude that, possibly along a subsequence, it holds

$$\mathbf{H}_t \left( \sqrt{\mathcal{I}^2(f_{\varepsilon_n}) + c_\alpha(t)\Gamma(f_{\varepsilon_n})} \right) \rightarrow \mathbf{H}_t \left( \sqrt{\mathcal{I}^2(f) + c_\alpha(t)\Gamma(f)} \right) \quad \mathbf{m}\text{-a.e. as } n \rightarrow \infty.$$

Therefore we can pass into the limit  $\mathbf{m}$ -a.e. also in the right hand side of (3.1) and conclude that (3.1) holds also for  $f$ .

### Step 2. Explicit computation of $\zeta$ .

Thanks to Step 1 we can assume  $f$  to take values into  $[\varepsilon, 1 - \varepsilon]$ , for some  $\varepsilon \in (0, 1/2)$ . Therefore by maximum principle also  $\mathbf{H}_t f$  will take values into  $[\varepsilon, 1 - \varepsilon]$ , and thus  $\mathcal{I}(\mathbf{H}_{T-t} f)$  will take

values into  $[\delta, \frac{1}{\sqrt{2\pi}} - \delta]$  for some  $\delta \in (0, 1/2\sqrt{2\pi})$  depending on  $\varepsilon$ . It follows that, in the range of  $(t, H_{T-t}f, \Gamma(H_{T-t}f))$ , the function

$$\Psi(t, u, v) := \sqrt{\mathcal{J}^2(u) + c_\alpha(t)v}, \quad (t, u, v) \in (0, T) \times \left[\delta, \frac{1}{\sqrt{2\pi}} - \delta\right] \times [0, \infty) \quad (3.18)$$

is of class  $C^4$  and we can then apply Lemma 3.2. Writing  $\mathcal{J}, \mathcal{J}'$  in place of  $\mathcal{J}(u), \mathcal{J}'(u)$  for brevity, it is immediate to check that

$$\Psi \partial_t \Psi = \frac{c'_\alpha}{2} v, \quad \Psi \partial_u \Psi = \mathcal{J} \mathcal{J}', \quad \Psi \partial_v \Psi = \frac{c_\alpha}{2},$$

and, by differentiating once more and using that  $\mathcal{J} \mathcal{J}'' = -1$ ,

$$\Psi^3 \partial_u^2 \Psi = -\mathcal{J}^2 \mathcal{J}'' + \Psi^2 (\mathcal{J}'' - 1), \quad \Psi^3 \partial_u \partial_v \Psi = -\frac{c_\alpha}{2} \mathcal{J} \mathcal{J}', \quad \Psi^3 \partial_v^2 \Psi = -\frac{c_\alpha^2}{4}.$$

Therefore, the corresponding function  $\zeta$  defined in (3.5), satisfies **m-a.e.**

$$\begin{aligned} \Psi^3 \zeta &= \Psi^2 \frac{c'_\alpha(t)}{2} \Gamma(g) - \mathcal{J}^2(g) \mathcal{J}'^2(g) \Gamma(g) + \Psi^2 (\mathcal{J}''(g) - 1) \Gamma(g) \\ &\quad - c_\alpha(t) \mathcal{J}(g) \mathcal{J}'(g) \Gamma(g, \Gamma(g)) - \frac{c_\alpha^2(t)}{4} \Gamma(\Gamma(g)) + K \Psi^2 c_\alpha(t) \Gamma(g) + \Psi^2 c_\alpha(t) \gamma_{2,K}[g], \end{aligned}$$

where we set  $g = H_{T-t}f$  and  $\Psi = \Psi(t, g, \Gamma(g))$  for brevity. Since  $\Psi^2(t, u, v) = \mathcal{J}^2(u) + c_\alpha(t)v$ , it follows that the following equalities hold **m-a.e.**:

$$\begin{aligned} \Psi^3 \zeta &= [\mathcal{J}^2(g) + c_\alpha(t) \Gamma(g)] \frac{c'_\alpha(t)}{2} \Gamma(g) - \mathcal{J}^2(g) \mathcal{J}'^2(g) \Gamma(g) + [\mathcal{J}^2(g) + c_\alpha(t) \Gamma(g)] (\mathcal{J}''(g) - 1) \Gamma(g) \\ &\quad - c_\alpha(t) \mathcal{J}(g) \mathcal{J}'(g) \Gamma(g, \Gamma(g)) - \frac{c_\alpha^2(t)}{4} \Gamma(\Gamma(g)) + K [\mathcal{J}^2(g) + c_\alpha(t) \Gamma(g)] c_\alpha(t) \Gamma(g) \\ &\quad + [\mathcal{J}^2(g) + c_\alpha(t) \Gamma(g)] c_\alpha(t) \gamma_{2,K}[g] \\ &= c_\alpha(t) \Gamma^2(g) \left( \frac{c'_\alpha(t)}{2} - 1 \right) + c_\alpha^2(t) \left[ -\frac{1}{4} \Gamma(\Gamma(g)) + \Gamma(g) \gamma_{2,K}[g] \right] + \mathcal{J}^2(g) \Gamma(g) \left( \frac{c'_\alpha(t)}{2} - 1 \right) \\ &\quad - c_\alpha(t) \mathcal{J}'(g) \mathcal{J}(g) \Gamma(g, \Gamma(g)) + c_\alpha(t) \mathcal{J}''(g) \Gamma^2(g) + K [\mathcal{J}^2(g) + c_\alpha(t) \Gamma(g)] c_\alpha(t) \Gamma(g) \\ &\quad + \mathcal{J}^2(g) c_\alpha(t) \gamma_{2,K}[g]. \end{aligned}$$

By the very definition (3.2) of  $c_\alpha(t)$ , we have that  $\frac{c'_\alpha(t)}{2} - 1 = -K c_\alpha(t)$  so that the above formula for  $\zeta$  simplifies to

$$\begin{aligned} \Psi^3 \zeta &= c_\alpha^2(t) \left[ -\frac{1}{4} \Gamma(\Gamma(g)) + \Gamma(g) \gamma_{2,K}[g] \right] \\ &\quad + c_\alpha(t) [-\mathcal{J}'(g) \mathcal{J}(g) \Gamma(g, \Gamma(g)) + \mathcal{J}''(g) \Gamma^2(g) + \mathcal{J}^2(g) \gamma_{2,K}[g]] \quad \text{m-a.e.} \quad (3.19) \end{aligned}$$

**Step 3.** Conclusion of the proof.

First of all observe that thanks to Lemma 2.5 we can apply Theorem 2.4 (which, let us stress, is implied by the curvature condition  $\text{RCD}(K, \infty)$ ) to  $g = H_{T-t}f$  and get

$$\Gamma(\Gamma(g)) \leq 4\Gamma(g) \gamma_{2,K}[g], \quad \text{m-a.e.} \quad (3.20)$$

Since  $\Psi = \Psi(t, H_{T-t}f, \Gamma(H_{T-t}f)) > 0$  as we are assuming  $f$  to take values into  $[\varepsilon, 1 - \varepsilon]$ , the combination of (3.4), (3.19) and (3.20) gives

$$\frac{d}{dt}|_{t=t_0} \Phi(t) \geq \int_{\{\Gamma(g) > 0\}} \frac{c_\alpha(t)}{\Psi^3} \left[ \mathcal{J}^2(g) \frac{\Gamma(\Gamma(g))}{4\Gamma(g)} - \mathcal{J}'(g) \mathcal{J}(g) \Gamma(g, \Gamma(g)) + \mathcal{J}'^2(g) \Gamma^2(g) \right] H_{t_0} \varphi \, d\mathbf{m}, \quad (3.21)$$

for all  $\varphi \in L^\infty(X, \mathbf{m})$ , with  $\varphi \geq 0$   $\mathbf{m}$ -a.e. in  $X$ . Notice that the restriction of the region of integration to  $\{\Gamma(g) > 0\}$  is possible because the nonnegativity of  $\gamma_{2,K}[g]$  and  $\Gamma(\Gamma(g)) = 0$   $\mathbf{m}$ -a.e. on  $\{\Gamma(g) = 0\}$  imply that  $\zeta$  is nonnegative  $\mathbf{m}$ -a.e. on  $\{\Gamma(g) = 0\}$ .

Now observe that the right hand side of (3.21) is a quadratic form in  $\mathcal{J}(g), \mathcal{J}'(g)$  which is positive definite since

$$\frac{\Gamma(\Gamma(g))}{4\Gamma(g)} \geq 0, \quad \Gamma(g)^2 \geq 0, \quad \text{and} \quad \Gamma(g, \Gamma(g)) \leq \Gamma^{1/2}(g) \Gamma^{1/2}(\Gamma(g)), \quad \mathbf{m}\text{-a.e.}$$

We conclude that for  $\mathcal{L}^1$ -a.e.  $t_0 \in (0, T)$  it holds  $\frac{d}{dt}|_{t=t_0} \Phi(t) \geq 0$ . Since by Lemma 3.2 we know that  $\Phi$  is continuous in  $[0, T]$  and locally Lipschitz in  $(0, T)$ , we can integrate to get

$$\begin{aligned} 0 &\leq \int_0^T \Phi'(t) \, dt = \Phi(T) - \Phi(0) \\ &= \int_X \left[ H_T \left( \sqrt{\mathcal{J}^2(f) + c_\alpha(T)\Gamma(f)} \right) - \sqrt{\mathcal{J}^2(H_T f) + \alpha\Gamma(H_T f)} \right] \varphi \, d\mathbf{m}, \end{aligned} \quad (3.22)$$

for all  $\varphi \in L^\infty(X, \mathbf{m})$ , with  $\varphi \geq 0$   $\mathbf{m}$ -a.e., which completes the proof of Theorem 3.1.  $\square$

## 4 Bobkov and Gaussian isoperimetric inequalities for $\text{RCD}(K, \infty)$ probability spaces, $K > 0$

In this section we specialize the local Bobkov Inequality (3.1) to the case  $K > 0$ , and use it to prove the Gaussian Isoperimetric Inequality.

**Theorem 4.1** (Bobkov Inequality). *Let  $(X, d, \mathbf{m})$  be an  $\text{RCD}(K, \infty)$  probability space for some  $K > 0$ . Then, for every  $f \in \mathbb{V}$  with values in  $[0, 1]$ , it holds*

$$\sqrt{K} \mathcal{J} \left( \int_X f \, d\mathbf{m} \right) \leq \int_X \sqrt{K\mathcal{J}^2(f) + \Gamma(f)} \, d\mathbf{m}. \quad (4.1)$$

*Proof.* Inequality (4.1) for  $f \in \text{LIP}(X)$  taking values in  $[0, 1]$  follows by the combination of the ergodicity of the Heat flow and the local Bobkov inequality. More precisely, set  $\alpha = 1/K$  (so that  $c_\alpha(t) = 1/K$ ) and let  $t \rightarrow +\infty$  in

$$\int_X \mathcal{J}(H_t f) \, d\mathbf{m} \leq \int_X \sqrt{\mathcal{J}^2(f) + \frac{1}{K}\Gamma(f)} \, d\mathbf{m}, \quad \mathbf{m}\text{-a.e.},$$

derived from (3.1) dropping one nonnegative term and integrating both sides; (4.1) follows then by applying the ergodicity property (2.22).

The validity of (4.1) for all  $f \in \mathbb{V}$  taking values in  $[0, 1]$  follows by the density of Lipschitz functions in  $\mathbb{V}$ , see [4, Proposition 4.10].  $\square$

Applying (4.1) to functions approximating the characteristic function of a set  $E \subset X$  we get the desired Gaussian isoperimetric inequality. More precisely, recall the standard definition of  $BV(X, \mathbf{d}, \mathbf{m}) \subset L^1(X, \mathbf{m})$  space and of total variation  $|Df|(X)$

$$|Df|(X) := \inf \left\{ \liminf_{n \rightarrow \infty} \int_X \text{lip } f_n \, d\mathbf{m} : f_n \in \text{LIP}(X), \lim_{n \rightarrow \infty} \int_X |f_n - f| \, d\mathbf{m} = 0 \right\},$$

we can pass to the limit in the inequality (derived from (4.1) from the subadditivity of the square root and the inequality  $\sqrt{\Gamma(f)} \leq \text{lip } f$   $\mathbf{m}$ -a.e.)

$$\sqrt{K} \, \mathcal{J} \left( \int_X f \, d\mathbf{m} \right) \leq \int_X (\sqrt{K} \, \mathcal{J}(f) + \text{lip } f) \, d\mathbf{m}$$

for all  $f \in \text{LIP}(X)$  with values in  $[0, 1]$  to get

$$\sqrt{K} \, \mathcal{J} \left( \int_X f \, d\mathbf{m} \right) \leq \sqrt{K} \int_X \mathcal{J}(f) \, d\mathbf{m} + |Df|(X) \quad \forall f \in BV(X, \mathbf{d}, \mathbf{m}).$$

In particular, by applying this to characteristic functions  $f = \chi_E$ , since  $\mathcal{J}(0) = \mathcal{J}(1) = 0$ , we obtain the Gaussian isoperimetric inequality with the perimeter  $\mathcal{P}(E) := |D\chi_E|(X)$ :

**Theorem 4.2.** *Let  $(X, \mathbf{d}, \mathbf{m})$  be an  $\text{RCD}(K, \infty)$  probability space for some  $K > 0$ . Then, for every Borel subset  $E \subset X$  it holds*

$$\mathcal{P}(E) \geq \sqrt{K} \, \mathcal{J}(\mathbf{m}(E)). \quad (4.2)$$

## References

- [1] L. AMBROSIO, N. GIGLI, A. MONDINO AND T. RAJALA, *Riemannian Ricci curvature lower bounds in metric spaces with  $\sigma$ -finite measure*, Trans. Amer. Math. Soc., **367**, (7), (2015), 4661–4701.
- [2] L. AMBROSIO, N. GIGLI AND G. SAVARÉ, *Gradient flows in metric spaces and in the space of probability measures*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, second ed., 2008.
- [3] —, *Calculus and heat flow in metric measure spaces and applications to spaces with Ricci bounds from below*, Invent. Math., **195**, (2), (2014), 289–391.
- [4] —, *Metric measure spaces with Riemannian Ricci curvature bounded from below*, Duke Math. J., **163**, (2014), 1405–1490.
- [5] —, *Bakry-Émery curvature-dimension condition and Riemannian Ricci curvature bounds*, Ann. Probab., **43**, (1), (2015), 339–404.
- [6] L. AMBROSIO, S. HONDA, *New stability results for sequences of metric measure spaces with uniform Ricci bounds from below*. In preparation.
- [7] L. AMBROSIO, A. MONDINO AND G. SAVARÉ, *On the Bakry-Émery condition, the gradient estimates and the Local-to-Global property of  $\text{RCD}^*(K, N)$  metric measure spaces*, J. Geom. Anal., **26**, (1), (2016), 24–56.

- [8] ———, *Nonlinear diffusion equations and curvature conditions in metric measure spaces*, preprint arXiv:1509.07273.
- [9] D. BAKRY, *L'hypercontractivité et son utilisation en théorie des semigroupes*, in Lectures on probability theory (Saint-Flour, 1992), vol. 1581 of Lecture Notes in Math., Springer, Berlin, (1994), 1–114.
- [10] D. BAKRY, I. GENTIL AND M. LEDOUX, *Analysis and Geometry of Markov Diffusion Operators*, Grundlehren der math. Wiss., Springer, (2014).
- [11] D. BAKRY AND M. ÉMERY, *Diffusions hypercontractives*, Sémin. de probabilités XIX, Univ. Strasbourg, Springer, (1985).
- [12] D. BAKRY AND M. LEDOUX, *Lévy-Gromov's isoperimetric inequality for an infinite dimensional diffusion generator*, Invent. Math., **123**, (1996), 259–281.
- [13] S.G. BOBKOV, *An isoperimetric inequality on the discrete cube, and an elementary proof of the isoperimetric inequality in Gauss space*, Ann. Probab., **25**, (1), (1997), 206–214.
- [14] C. BORELL, *The Brunn-Minkowski inequality in Gauss space*, Invent. Math., **30**, (2), (1975), 207–216.
- [15] F. CAVALLETTI AND A. MONDINO *Sharp and rigid isoperimetric inequalities in metric-measure spaces with lower Ricci curvature bounds*, preprint arXiv:1502.06465, (2015).
- [16] J. CHEEGER, *Differentiability of Lipschitz functions on metric measure spaces*, Geom. Funct. Anal., **9** (1999), 428–517.
- [17] M. ERBAR, K. KUWADA, K.-T. STURM, *On the equivalence of the entropic curvature-dimension condition and Bochner's inequality on metric measure spaces*, Invent. Math., **201** (2015), 993–1071.
- [18] N. GIGLI, A. MONDINO AND G. SAVARÉ, *Convergence of pointed non-compact metric measure spaces and stability of Ricci curvature bounds and heat flows*, Proc. London Math. Soc., **111**, (5), (2015), 1071–1129.
- [19] N. GIGLI, *Nonsmooth differential geometry – An approach tailored for spaces with Ricci curvature bounded from below*. ArXiv preprint 1407.0809. To appear on Mem. Am. Math. Soc.
- [20] M. GROMOV, *Metric structures for Riemannian and non Riemannian spaces*, Modern Birkhäuser Classics, (2007).
- [21] J. LOTT AND C. VILLANI, *Ricci curvature for metric-measure spaces via optimal transport*, Ann. of Math. **169** (2009), 903–991.
- [22] G. SAVARÉ, *Self-improvement of the Bakry-Émery condition and Wasserstein contraction of the heat flow in  $RCD(K, \infty)$  metric measure spaces*, Disc. Cont. Dyn. Syst. A, **34**, (2014), 1641–1661.
- [23] K.T. STURM, *On the geometry of metric measure spaces. I*, Acta Math. **196** (2006), 65–131.

- [24] ———, *On the geometry of metric measure spaces. II*, Acta Math. **196** (2006), 133–177.
- [25] V.N. SUDAKOV, B.S. CIREL'SON, *Extremal properties of half-spaces for spherically invariant measures*, Zap. Nauc. Semin. LOMI 41, 14–24, 165 (1974). Problems in the theory of probability distributions, II.
- [26] C. VILLANI, *Optimal transport. Old and new*, vol. 338 of Grundlehren der Mathematischen Wissenschaften, Springer-Verlag, Berlin, 2009.