

Human responses to climate change in the late prehistoric Western Loess Plateau, northwest China

Tingting Chen ¹, Menghan Qiu ¹, Ruiliang Liu ^{2,*}, Haiming Li ¹, Hongwei Hou ³,

Philly Howarth ², Samantha Bowring ², Aifeng Zhou ^{1,*}

¹ MOE Key Laboratory of Western China's Environmental System, College of Earth & Environmental Sciences, Lanzhou University, Lanzhou, 730000, China

² School of Archaeology, University of Oxford, Oxford OX1 3TG, United Kingdom

³ Gansu Provincial Institute of Cultural Relics and Archaeology, Lanzhou, 730000, China

* Corresponding author

E-mail: * ruiliang.liu@arch.ox.ac.uk; * zhouaf@lzu.edu.cn

Abstract Human responses to the rapid climate change in antiquity has been extensively discussed and debated. This paper reports 91 new archaeobotanical results including 13567 charred plant seeds and 19 radiocarbon dates from 41 late prehistoric sites in the upper Wei River valley of Western Loess Plateau (WLP). Based on the new dating results as well as the culture attributes, the 41 sites can be divided into four chronological phases (Phase 1: Late Yangshao and Majiayao culture; Phase 2: Qijia culture; Phase 4: Siwa culture). A significant gap has been identified around 3600-3000 cal yr BP in this region. A further comparison with the high-resolution

paleoclimate records (Tianchi) suggests that this gap could be attributed to the dramatic drop in the temperature around 3600 cal yr BP. But archaeobotanical evidence with refined chronology shows that adoption of cold-tolerant subsistence such as barley in NETP. Drawing from various lines of information (chronology, palaeoclimate, archaeobotany and archaeology), it is reasonable to conclude that even confronting the similar magnitude of climate change, the local human responses could vary tremendously. It was essentially result of the asynchronous adoption of the new subsistence strategies brought by the prehistoric trans-Eurasia culture exchange.

Keywords: archaeobotanical analysis, radiocarbon dating, subsistence strategy, late Neolithic and Bronze periods, climate change

INTRODUCTION

Climate change matters in the human history (e.g. Weiss et al., 1993; D'Andrea et al., 2011; Wu et al., 2018). Large-scale migration or social expansions, technological innovations, and the rise and fall of ancient civilisations were often triggered by important climate events, such as significant cold periods, droughts (Marshall et al., 2011; Timmermann and Friedrich, 2016; Staubwasser et al., 2018). **Climate events declined the ancient civilizations (Staubwasser et al., 2003; Dong et al., 2017a)** in the Late Pleistocene and the Early to Mid-Holocene. From Late Paleolithic to Neolithic, when the climate became **favorable for human survival**, human activity was markedly increased **and** culture flourished (Hu et al., 2016). However, when climate became deteriorated, we can see a series of cultural collapse and migration to lower elevations or latitudes (Bhattacharya et al., 2015; Bai et al., 2017; Cui et al., 2018). The relationship between climate change and human activity during the transition period from Late Neolithic to Bronze Age, which was characterized by global cooling but marked expansion of human living space. Such as Central Asia (Spengler and Willcox, 2013; Spengler, 2015) and the Tibetan plateau (Chen et al., 2015). Different trajectories of human response to climate change in the prehistoric period in the world is various. During 8450-5650 cal yr BP, as a result of major changes in temperature and humidity, the Apulia region of southeast Italy saw considerable fluctuations of the population density. Local farmers therefore had to adjust the sorts of crops that were available to them and the cultivation time (Fiorentino et al., 2013). Likewise,

although radical changes happened in Greece in 6500-500 cal yr BP with the three continuous droughts (5800-5700, 5450 and 5000-4900 cal yr BP), they appeared to have very little impact on the human activities. Local people became well adapted to the climate changes by adjusting the agricultural and pastoral practice (Lespez et al., 2016). The Canaan region in the western side of Eurasia, controlled by the Egyptian empire in the late Bronze Age had experienced a dry climate around 3200-3050 cal yr BP. The governors chose to develop new technologies such as dry farming and ploughing in order to cope with the increasingly drier conditions reflected in increasing cattle frequencies and animals that were kept to colder age (Finkelstein et al., 2017). In the Indus valley during the post-urban period (3950-3150 cal yr BP), the Late Harappan civilization increased the cultivation of drought-resistant crops (e.g. millet) in the peripheral areas, which was also an echo to the decline of SW monsoon during the late Holocene, to which millets are better suited (Pokharia et al., 2014). Except them, the same gap around 3.5 ka BP between the Lower Xiajiadian Culture and Upper Xiajiadian Culture in the West Liao River Basin (Jia et al., 2016), the birth of civilization in the Central Plains, and the cultural transformation from Baodun Culture to Sanxingdui Culture in Chengdu Plain (Zhang et al., 2019).

In the northeast Tibetan Plateau (NETP), people migrated toward the high-altitude areas from the upper Yellow River valleys and became permanently settled above 3000 m.a.s.l (meter above sea level) after 3600 cal yr BP (the calibrated year before present, before AD 1950) (Chen et al., 2015; Dong et al., 2016) when climate became cooler (Marcott et al., 2013). Archaeobotanical and zooarchaeological evidence suggests that adoption of barley growing, sheep and yak herding, greatly facilitated the year-round activities in the high-cold environment of the Tibetan Plateau in the

Bronze Age. The results of archaeological survey and radiocarbon dating (Bureau of National Cultural Relics, 1996; Dong et al., 2013; Chen et al., 2015) suggest a clear continuation in settlement between Neolithic and Bronze Age in NETP (Fig 1a) and the number of Bronze Age sites dated to 4000-2300 cal yr BP in that area is much more abundant than in the Neolithic (5200-4000 cal yr BP).

However, the trajectory of the culture evolution in the adjacent Longxi Loess Plateau (WLP) appears rather different. The Yangshao and Majiayao Neolithic cultures (~6000-4000 cal yr BP) and the Qijia Chalcolithic culture (~4300-3600 cal yr BP) were prosperous but the Bronze Age cultures, including Siwa (~3600-2600 cal yr BP), show a downward trend (Bureau of National Cultural Relics, 2011). The spatial pattern for the culture evolution in WLP and NETP during the Bronze Age was abnormal. In theory, the lower altitude environment in WLP to be more suitable for human survival and cultural development than NETP, which is located on higher altitudes. Therefore, it is necessary to further understand the different patterns for the evolution of prehistoric human-environment interaction in these two areas and the underlying mechanism of development.

The Tibetan Plateau has been the focus of a large number of research projects (Chen et al., 2015; Dong et al., 2016). Given its remarkable geographical feature such as low-high elevation, changing temperature and humidity or the different vegetation, human inhabitation to the Tibetan Plateau in prehistory must have required a variety of strategies and adaptations that were dramatically different from those in the lower flat plains. This paper focused on the two adjacent areas (NETP and WLP) to attempt to illustrate different human responses to the same climatic change. The Western

Loess Plateau is really important. It is the link between the Central Plains and the northeast Tibetan Plateau and the prehistoric culture in the Western Loess Plateau is also a key node for the spread westward from Central Plains culture to the Northeast Tibetan Plateau. But the history of human settlement and subsistence strategy in WLP during the late prehistoric period remains rather ambiguous, owing to a lack of archaeobotanical and chronological data. While NETP has been comprehensively studied (Chen et al. 2015; Dong et al. 2016). This paper has investigated a substantial number of Neolithic and Bronze Age sites in the upper Wei River valley of WLP (Bureau of National Cultural Relics, 2011). With the aid of 13567 charred plant seeds and 19 AMS ^{14}C dates, we have reconstructed the chronology of local settlements and associated crop subsistence. A further comparative study between the charred plant seeds of WLP/NETP and the high-resolution paleoclimate record illustrates the rather different patterns of social evolution affected by the climate change and cultural exchanges around the northeastern area of Tibetan Plateau during the late prehistoric period.

STUDY AREA

The upper Wei River valley (104°41'-106°18' E, 34°24'-35°8' N) is located in the Gansu Province (Fig 1b). The study area is approximately 200 km long from west to east. The trunk stream flows from 1600-900 m.a.s.l and forms a gentle slope and a broad river valley. The climate of this area was classified as continental winter dry (Dwb) according to Köppen classes classification (Chan et al., 2016). The mean annual temperature is about 8-10°C and the mean annual precipitation is around 500-650 mm. Modern agriculture in this area includes millets, wheat, corn, potato etc.

The **Second** National Archaeological Survey in China have found more than two hundred archaeological sites in this area, ranging from the Late Neolithic to the Bronze Age (Bureau of National Cultural Relics, 2011). A majority of them are distributed in the valley of the upper Wei River and its main tributaries. The late Neolithic and Chalcolithic sites can be further divided into three cultures, among which 125 have been attributed to the Late Yangshao Culture (~5500-5000 cal yr BP), 42 to the Majiayao Culture (~5300-4000 cal yr BP) and 371 to the Qijia Culture (~4300-3600 cal yr BP). Merely 14 belong to the Siwa Culture (~3600-2600 cal yr BP) and one site belongs to the Xindian Culture (~3600-2600 cal yr BP).

MATERIALS AND METHODS

In 2016, 41 Late Neolithic and Bronze Age sites in **WLP** were investigated by the Gansu Institute of Cultural Relics and Archaeology (Fig 1b). At each site, one can identify archaeological deposits such as single or multiple cultural layers and ash pits that have been exposed on the profile. These deposits in different archaeological units were sampled for floatation. The units were avoided in case of any latter natural or anthropogenic disturbances. The average volume of the 91 sampled soils was 19.78 L. Over the process of flotation, lighter remains such as charcoal and charred plant seeds were floated upward and gathered by 80-mesh sifter (with 0.2 mm aperture). These were subsequently wrapped by gauze and hung in shady and cool area for desiccation. The species of the charred plant seeds were identified **in** the Paleoethnobotany Laboratory, Institute of Archaeology, Chinese Academy of Social Sciences and the MOE Key Laboratory of Western China's Environmental Systems, College of Earth & Environmental Sciences, Lanzhou University.

The number of different plant species were recorded to analyze the structure of the past agricultural activities in the targeted area. However, different crops have different weight and behave rather differently in the process of harvesting, utilizing and carbonizing (Yang et al., 2011). As a result, the proportions revealed by the unearthed charred seeds may not precisely reflect the true proportion of crops utilized in past agriculture. Therefore, a modified method of using the Weight Ratio Function for different crops proposed by Zhou (Zhou et al., 2016). The statistical analysis of the Weight Ratio Function of charred plant seeds can better reflect the actual proportion of different sorts of crops.

Where N_1 =number of common millet grains, $F_1=7.5$, N_2 = number of foxtail millet grains, $F_2=2.6$, N_3 =number of wheat grains, $F_3=35$, N_4 =number of barley grains, $F_4=45$, and $P(S)$ =actual yield percentage of that particular crop.

19 of plant remains were selected for radiocarbon dating, including foxtail millet, wheat and barley (Table 1). The samples were pretreated at the MOE Key Laboratory of Western China's Environmental Systems, College of Earth & Environmental Sciences, Lanzhou University using standard pre-treatment (acid-alkali-acid) and were subsequently measured by accelerator mass spectrometry (AMS) at Peking University. All dates were calibrated using Calib 7.0.2 calibration software (Reimer et al., 2009) with the IntCal13 calibration curve (Reimer et al., 2013). The cumulative probability distribution (CPD) curve of radiocarbon dates has been applied to estimate the intensity of human activities in the following discussion (Michczyńska and

Pazdur., 2004).

RESULTS

The 19 AMS ^{14}C dates are listed at Table 1, ranging between 5500 and 2300 cal yr BP. Based on the new dating results as well as the culture attributes (Phase 1: Late Yangshao and Majiayao culture; Phase 2: Qijia culture; Phase 4: Siwa culture), the 41 sites can be divided into four chronological phases. Phases 1, 2, 4 are contemporary with the Late Yangshao-Majiayao cultures, the Qijia culture and the Siwa culture, respectively.

A total number of 13567 charred plant seeds have been recovered from the 91 flotations (Table 2), including 9809 foxtail millet (*Setaria italica*) (Fig 2a), 1649 broomcorn millet (*Panicum miliaceum*) (Fig 2b), 1016 wheat (*Triticum aestivum*) (Fig 2c) and 259 barley (*Hordeum vulgare*). The rest 834 seeds are weed remains (such as *Chenopodium album*, *Setaria viridis*, *Melilotus suaveolens*, *Xanthium sibiricum*, *Sphaerophysa salsula*, *Atriplex patens*, *Galium tricornis*, *Glycyrrhiza uralensis*, *Kochia scoparia*, *Salsola collina* pall, *Polygonum nepalense*, *Rumex acetosa* and et al) (Fig 2d).

Based on the new radiocarbon dates and the results of archaeobotanical analysis, one can quickly draw an agriculture development for WLP during the late prehistoric period (Table 1, 2; Fig 3e). During the Phase 1 (5500-4300 cal yr BP), the 4934 foxtail millets (72.46% of weight) and 650 broomcorn millets (27.54% of weight) identified from 42 samples (855 L soil in total) indicate a millet-based agriculture. Only one charred barley grain has been identified from the Shizhaocun, which is very

likely due to the later disturbance of the site or context. Similar issues were also encountered in previous research (Dodson et al., 2013; Jia et al., 2013). During Phase 2 (4300-3600 cal yr BP), a mixed farming practice gradually emerged: 3783 foxtail millets (75.44% of weight), 239 broomcorn millets (13.75% of weight), 39 wheats (10.47% of weight) and 1 barley were collected from 21 samples of 387 L soil. Phase 3, spanning from 3600 to 3000 cal yr BP, reveals a significant gap in human activities in **WLP**. No traces of agriculture within the study area are found during this phase. During Phase 4 (3000-2300 cal yr BP), A more mixed agricultural tradition is in evidence when compared to Phase 2, as the addition of wheat and barley to the dominant crops occurs. 977 wheat (62.98% of weight), 257 barley (21.30% of weight), 760 broomcorn millet (10.50% of weight) and 1092 foxtail millet (5.22% of weight) were identified from 28 samples (560 L soil).

DISCUSSION

Different temporospatial patterns for the human activities and subsistence strategies in **WLP and NETP during the late prehistoric periods**

Archaeobotanical studies and radiocarbon dating have greatly contributed to the study of the density of human activities and subsistence strategies in the prehistoric societies (Hageman and Goldstein, 2009; Gaudzinski-Windheuser and Kindler, 2012; Dong et al., 2016, 2018). New chronological framework can be established for the occupation sequence in **WLP** by the new results of radiocarbon dating together with the published crop grain dates (Fig 3f). **Relative to Phase 2 (4300-3600 cal yr BP), the probability density of radiocarbon dating appears more gentle of Phase 1 (5500-4300 cal yr BP).** This is probably representative of an early stage of settlement by the farming group in

WLP. Phase 2 (4300-3600 cal yr BP) witnessed a sharp increase in the probability density, which reaches the maximum between 3900-3800 cal yr BP, suggestive of rapid intensification of human activities in the Qijia period in **WLP**. This theory is reinforced by the results of national archaeological survey (Bureau of National Cultural Relics, 2011). Based on the updated radiocarbon data, the lack of radiocarbon dates between ~3600-3000 cal yr BP suggests a gap in human activities in **WLP** (Table 1).

Comparing with **WLP**, the history of human activities during the late prehistoric periods in NETP depicts the same trajectory during Phase 1 and 2, but radically differ in Phase 3 and 4 (Fig 3h). During Phase 1 and Phase 2, the intensity of human activity also fluctuated at a relatively low profile, then increased and reached a peak at ~4100-3800 cal yr BP before dropping again. However, during next two phases of NETP, the intensity of human practice maintained a relatively high level from ~3500-3100 cal yr BP which continued to ~2300 cal yr BP (Fig 3h, Chen et al., 2015).

Archaeobotanical and zooarchaeological evidence can also make vital contributions to the understanding of subsistence strategy in these two areas during the late prehistoric period (Lee et al., 2007; Yuan et al., 2008; Zhao et al., 2010). In Phase 1, people in **WLP** only cultivated foxtail millet and broomcorn millet, which was initially domesticated in north China (Zhao, 2011). As seen in Fig 3e, the importance of foxtail millet appears far greater than the other. These millet farmers also raised domestic animals (pig and dog) and hunted wild animals (**Caprinae** and deer) as auxiliary animal resource (Zhou, 1999). In Phase 2 sheep, wheat and barley that were initially domesticated in west Asia (Zeder, 2008; Riehl et al., 2013) were introduced into

northwest China (Flad et al., 2007; Dodson et al., 2013; Dong et al., 2018). While wheat became the most important cultivated crop in the Hexi Corridor from ~3700 cal yr BP (Zhou et al., 2016) and was utilized in WLP during Phase 2, the most dominant local crop was still millet (Fig 3e). Phase 2 also saw an increasing variety of domestic animals, including sheep, cattle, pig and dog in WLP (Zhou, 1999; Jia et al., 2013). There was no sites to excavating crops in WLP of Phase 3, but wheat was the most important crops in Phase 4 (Fig 3e).

The history for the transition of subsistence strategies in NETP was different in comparison to that of WLP. Local NETP people relied mainly on broomcorn millet instead of foxtail millet in Phase 1 (Chen et al., 2015; Fig 3g). Hunting was another important subsistence strategy in NETP during the same period, Millet continued to be the major local crop in Phase 2 and pigs and dogs domesticated (Ren, 2017; Wang, 2017). And a supplementary proportion of barley being used simultaneously (Fig 3g). This is also verified by the stable isotopic analysis of human bones unearthed from sites dated to the same period (Ma et al., 2014; 2016). Zooarchaeological analysis from the Lajia site of NETP also suggests an interesting change in the meat supply, shifting from pig to sheep (Ye, 2015). Phase 3 in NETP, in contrast to Phase 3 in WLP, witnessed not only the radical expansion of humans' living space, the increasing dependence on barley (Fig 3g), but also different subsistence strategies taken by different groups of people along different altitudes (Zhang and Dong, 2017). This spatial differentiation of the human subsistence was retained in Phase 4 in NETP and the most characteristic pattern was the increasing ratio of barley in NETP (Chen et al., 2015; Zhang and Dong, 2017).

In summary, the temporospatial difference of human subsistence strategy in NETP and WLP was mainly reflected by the variation in cultivated crop types. During Phase 1, the most important crops were foxtail and broomcorn millet in WLP and NETP. Stepping into the next phase, the primary crop in both regions was foxtail millet, broomcorn millet, and wheat began to increase in WLP and barley began to increase in NETP (Figs 3e and 3g). In Phases 3 and 4, barley became the dominant plant subsistence in NETP, while WLP shifted to wheat, followed by barley, broomcorn and foxtail millet (Figs 3e and 3g).

Contrasting different human responses to the same climate change between WLP and NETP

The late Neolithic and Bronze Age were unique periods to studying the evolution of the human-land relation, because this period has the rise and fall of the earliest civilizations, trans-Eurasia culture exchange (Spengler et al., 2014; Dong et al., 2017b; Liu et al., 2019). According to high-resolution paleoclimate records in WLP, including bio-markers from the sediment of the Liupan Tianchi lake (Sun, 2011; Sun et al., 2018) and oxygen isotopes of stalagmite in the Jiuxian cave (Cai et al., 2010), the temperature remained relatively high during 5500-3600 cal yr BP as a whole, but followed by several cold periods between ~4900-4500 cal yr BP and 3600-2000 cal yr BP (Fig 3d). Meanwhile, four droughts can be detected in the periods of ~4900-4700 cal yr BP, ~3800-3600 cal yr BP, ~3000-2800 cal yr BP and ~2700-2500 cal yr BP, respectively (Figs 3a and 3b).

The correlation between paleoclimate records and CPD curves of radiocarbon dates suggests that human settlement in WLP and NETP during the late prehistoric period

was mainly affected by the variation in temperature rather than precipitation. Compared with precipitation, the decline in temperature appears more dramatic. The shortage of water supply caused by decreased precipitation (with fluctuations) can also be eased by the abundant local river system, such as the Wei River and its branches. The most notable decline in the CPD distribution of the two areas occurred around 3600 cal yr BP, when the temperature reached its lowest (Figs 3c and 3d). In NETP settlement was soon re-established but WLP experienced a period of nearly six hundred years with virtually no human activities. The next inhabitation of WLP can be seen during 3000-2300 cal yr BP (Phase 4). It was an interesting period as there were two droughts identified by the record obviously (Figs 3a and 3b). This phenomenon might be related to the physiological characteristics of foxtail and broomcorn millets, both of which are drought-tolerant but sensitive to low temperature (Baltensperger, 1996).

The difference of life-styles in NETP and WLP during the Late Neolithic period was also affected by the climate. The temperature becomes much lower in NETP than in WLP due to the higher elevation. Therefore, human mainly cultivated foxtail millet in WLP, compared to broomcorn millet in NETP during 5500-4300 cal yr BP (Phase 1) as broomcorn millet is much more resistant to a cold environment (Baltensperger, 1996). Temperature between 4300-4000 cal yr BP was higher than Phase 1 (Fig 3d), foxtail millet became the primary crop in NETP during that period (Chen et al., 2015). Wheat and barley were subsequently introduced into northwest China around 4000 cal yr BP (Dodson et al., 2013; Dong, 2018). When the temperature declined again between 4000-3600 cal yr BP (Figs 3c and 3d), human in NETP and WLP might have added barley and wheat into their cultivated crops (Figs 3e and 3g), sheep was also

frequently utilized in NETP during that period (Ye, 2015; Ren, 2017).

The temperature remained persistently low in Phases 3 and 4 (Figs 3c and 3d), which might have significantly affected the yield of millet and therefore resulted in the absence of human activities in **WLP** (Table1; Fig 3f). However, people survived in NETP and even successfully expanded to high-cold areas during this period. This could be explained by the development of new strategies such as the cultivation of barley and adopting sheep and yak (Chen et al., 2015; Dong et al., 2016). In Phase 4, **WLP** decided to have wheat that has the higher production than barley as the primary plant subsistence, along with the emergence of the Siwa culture in the area. Undoubtedly, the choice made by **WLP** was completely different from NETP. Compared to wheat, barley requires a much shorter growth period and appears more adapted to high-cold areas of the Tibetan Plateau (Páldi et al., 2001; D'Alpoim Guedes et al., 2014, 2015).

CONCLUSION

The new archaeobotanical and radiocarbon results in the upper Wei River valley, combined with previously published data, have demonstrated that people in **WLP** were mainly dependent on foxtail millet and to a lesser extent, broomcorn millet during 5500-4300 cal yr BP. Wheat was subsequently added to the local choice during 4300-3600 cal yr BP. The current data shows a serious gap of human activity in **WLP** during 3600-3000 cal yr BP. The area/region was repopulated between 3000-2300 cal yr BP, during which phase wheat became the most important subsistence, followed by barley, broomcorn and foxtail millet. Further comparisons to high-resolution paleoclimate records suggest that human activity in the late prehistoric **WLP** was not

affected by variation in precipitation levels but variation in temperature.

Despite the rapid decline of the local temperature, by adopting new crop species with different physiological characteristics, people successfully adapted to the new climatic conditions in NETP and WLP during 5500-3600 cal yr BP. One of the ultimate reasons that allowed people to survive (and even migrate to higher elevations in NETP) was the quick adoption of new cold-tolerant barley and sheep, which were brought into northwest China by trans-Eurasia culture exchange during early Bronze Age.

ACKNOWLEDGEMENTS

This research was supported by the National Key R&D Program of China (2018YFA0606402), Strategic Priority Research Program of the Chinese Academy of Sciences, Pan-Third Pole Environment Study for a Green Silk Road (Pan-TPE) (Grant No. XDA2004010101), the National Natural Science Foundation of China (Grant Nos. 41671077 and 41620104007), the International Partnership Program of Chinese Academy of Sciences (grant no. 131C11KYSB20160061).

REFERENCE

- Bai YJ, Zhang PZ, Gao T, Yu RZ, Zhou PC, Cheng H. 2017. The 5400 a BP extreme weakening event of the Asian summer monsoon and cultural evolution. *Science China Earth Sciences* 60: 1171-1182.
- Baltensperger DD. 1996. Foxtail and proso millet. In: Janick J (ed) Progress in new crops. *Alexandria: ASHS Press* 182-190.
- Bhattacharya T, Byrne R, Böhnelt H, Wogau K, Kienel U, Ingram BL, Zimmerman S. 2015. Cultural implications of late Holocene climate change in the Cuenca Oriental, Mexico. *Proceedings of the National Academy of Sciences of USA* 112: 1693-1698.
- Bureau of National Cultural Relics. 1996. Atlas of Chinese Cultural Relics-Fascicule of Qinghai Province. *Beijing: China Cartographic Publishing House Press*. In Chinese.
- Bureau of National Cultural Relics. 2011. Atlas of Chinese Cultural Relics-Fascicule of Gansu Province. *Beijing: Surveying and Mapping Press*. In Chinese.
- Cai YJ, Tan LC, Cheng H, An ZS, Edwards RL, Kelly MJ, Kong XG, Wang XF. 2010. The variation of summer monsoon precipitation in central China since the last deglaciation. *Earth and Planetary Science Letters* 291: 21-31.

- Chan D, Wu Q, Jiang GX, Dai XL. 2016. Projected shifts in Köppen climate zones over China and their temporal evolution in CMIP5 multi-model simulations. *Advances in Atmospheric Sciences* 33: 283-293.
- Chen FH, Dong GH, Zhang DJ, Liu XY, Jia X, An CB, Ma MM, Xie YW, Barton L, Ren XY, Zhao ZJ, Wu XH, Jones MK. 2015. Agriculture facilitated permanent human occupation of the Tibetan Plateau after 3600 BP. *Science* 347: 248-250.
- Chen TT, Jia X, Li HM, Dong GH. 2019. The analysis of spatiotemporal transformations of agricultural and its influence factors during Qijia culture period in Gansu-Qinghai region. *Quaternary Sciences* 39: 132-144. In Chinese.
- Cui YF, Liu YJ, Ma MM. 2018. Spatiotemporal evolution of prehistoric Neolithic-Bronze Age settlements and influencing factors in the Guanting Basin, northeast Tibetan Plateau. *Science China Earth Sciences* 61: 149-162.
- D’Alpoim Guedes J, Lu HL, Li YX, Spengler R, Wu XH, Aldenderfer M. 2014. Moving agriculture onto the Tibetan plateau: the archaeobotanical evidence. *Archaeological and Anthropological Sciences* 6: 255-269.
- D’Alpoim Guedes J, Lu HL, Hein AM, Schmidt AH. 2015. Early evidence for the use of wheat and barley as staple crops on the margins of the Tibetan Plateau. *Proceedings of the National Academy of Sciences of USA* 112: 5625-5630.
- D’Andrea WJ, Huang YS, Fritz SC, Anderson NJ. 2011. Abrupt Holocene climate change as an important factor for human migration in West Greenland. *Proceedings of the National Academy of Sciences of USA* 108: 9765-9769.
- Dodson JR, Li XQ, Zhou XY, Zhao KL, Sun N, Atahan P. 2013. Origin and spread of

wheat in China. *Quaternary Science Reviews* 72: 108-111.

Dong GH, Jia X, Robert E, Chen FH, Li SC, Wang L, Cai LH, An CB. 2013. Spatial and temporal variety of prehistoric sites and its influencing factors in the upper Yellow River valley Qinghai Province China. *Journal of Archaeological Science* 40: 2538-2546.

Dong GH, Wang ZL, Ren LL, Matuzeviciute GM, Wang H, Ren XY, Chen FH. 2014. A comparative study of ^{14}C dating on charcoal and charred seeds from Late Neolithic and Bronze Age sites in Gansu and Qinghai Provinces NW China. *Radiocarbon* 56: 157-163.

Dong GH, Ren LL, Jia X, Liu XY, Dong SM, Li HM, Wang ZX, Xiao YM, Chen FH. 2016. Chronology and subsistence strategy of Nuomuhong culture in the Tibetan Plateau. *Quaternary International* 426: 42-49.

Dong GH, Liu FW, Chen FH. 2017a. Environmental and technological effects on ancient social evolution at different spatial scales. *Science China Earth Sciences* 60: 2067-2077.

Dong GH, Yang YS, Han JY, Wang H, Chen FH. 2017b. Exploring the history of cultural exchange in prehistoric Eurasia from the perspectives of crop diffusion and consumption. *Science China Earth Sciences* 60: 1110-1123.

Dong GH. 2018. A new story for wheat into China. *Nature Plants* 4: 243-244.

Dong GH, Yang YS, Liu XY, Li HM, Cui YF, Wang H, Chen GK, Dodson J, Chen FH. 2018. Prehistoric trans-continental cultural exchange in the Hexi Corridor

- northwest China. *The Holocene* 28: 621–628.
- Flad RK, Yuan J, Li SC. 2007. Zooarcheological evidence for animal domestication in northwest China. *Developments in Quaternary Sciences* 9: 167-204.
- Finkelstein I, Langgut D, Meiri M, Sapir-Hen L. 2017. Egyptian imperial economy in Canaan: Reaction to the climate crisis at the end of the Late Bronze Age. *Ägypten und Levante/Egypt and the Levant* 27: 249-259.
- Fiorentino G, Caldara M, De Santism V, D’Oronzom C, Muntoni IM, Simone O, Primavera M, Radina F. 2013. Climate changes and human–environment interactions in the Apulia region of southeastern Italy during the Neolithic period. *The Holocene* 23: 1297-1316.
- Gaudzinski-Windheuser S, Kindler L. 2012. Research perspectives for the study of Neandertal subsistence strategies based on the analysis of archaeozoological assemblages. *Quaternary International* 247: 59-68.
- Hageman JB, Goldstein DJ. 2009. An integrated assessment of archaeobotanical recovery methods in the Neotropical rainforest of northern Belize: flotation and dry screening. *Journal of Archaeological Science* 36: 2841-2852.
- Hu YQ, Cao XY, Zhao ZJ, Li YY, Sun YG, Wang H. 2016. The palaeoenvironmental and palaeoclimatic reconstruction and the relation with the human activities during the early and middle Holocene in the upper western Liao River region. *Quaternary Sciences* 36: 530-541. In Chinese.
- Ji DX. 2007. Environmental archaeological perspective: dispersal of anatomically modern human and origin of agriculture in Northern China-A case study in

Gansu and Ningxia. Doctoral Dissertation Lanzhou: Lanzhou University 72-85.
In Chinese.

Jia X, Dong GH, Li H, Brunson K, Chen FH, Ma MM, Wang H, An CB, Zhang KR.
2013. The development of agriculture and its impact on cultural expansion
during the late Neolithic in the Western Loess Plateau China. *The Holocene* 237:
85-92.

Jia X, Sun YG, Wang L, Sun WF, Zhao ZJ, Lee HF, Huang WB, Wu SY, Lu HY.
2016. The transition of human subsistence strategies in relation to climate change
during the Bronze Age in the West Liao River Basin, Northeast China. *The
Holocene* 26: 781-789.

Lee GA, Bestel S. 2007. Contextual Analysis of Plant Remains at the Erlitou Period
Huizui Site Henan China. *Bulletin of the Indo-Pacific Prehistory Association* 27:
49-60.

Lespez L, Glais A, Lopez-Saez JA, Le Drezen Y, Tsirtsoni Z, Davidson R, Biree L,
Malamidou D. 2016. Middle Holocene rapid environmental changes and human
adaptation in Greece. *Quaternary Research* 85: 227-244.

Li XQ, Zhou XY, Zhou J, Dodson J, Zhang HB, Shang X. 2007. The earliest
archaeobiological evidence of the broadening agriculture in China recorded at
Xishanping site in Gansu Province. *Science in China Series D: Earth Sciences*
50: 1707-1714.

Liu XY, Jones PJ, Matuzeviciute GM, Hunt HV, Lister DL, An T, Przelomska N,
Kneale CJ, Zhao ZJ, Jones MK. 2019. From ecological opportunism to multi-

cropping: Mapping food globalisation in prehistory. *Quaternary Science Reviews* 206: 21-28.

Ma MM, Dong GH, Lightfoot E, Wang H, Liu XY, Jia X, Zhang KR, Chen FH. 2014. Stable Isotope Analysis of Human and Faunal Remains in the Western Loess Plateau Approximately 2000 Cal BC. *Archaeometry* 56: 237-255.

Ma MM, Dong GH, Jia X, Wang H, Cui YF, Chen FH. 2016. Dietary shift after 3600 cal yr BP and its influencing factors in northwestern China: Evidence from stable isotopes. *Quaternary Science Reviews* 145: 57-70.

Marcott SA, Shakun JD, Clark PU, Mix AC. 2013. A reconstruction of regional and global temperature for the past 11300 years. *Science* 339: 1198-1201.

Marshall MH, Lamb HF, Huws D, Davies SJ, Bates CR, Bloemendal J, Boyle J, Leng MJ, Umer M, Bryant C. 2011. Late Pleistocene and Holocene drought events at Lake Tana the source of the Blue Nile. *Global and Planetary Change* 78: 147-161.

Michczyńska DJ, Pazdur A. 2004. Shape analysis of cumulative probability density function of radiocarbon dates set in the study of climate change in the Late Glacial and Holocene. *Radiocarbon* 46: 733-744.

Páldi E, Szalai G, Janda T, Horváth E, Rácz I, Lásztity D. 2001. Determination of frost tolerance in winter wheat and barley at the seedling stage. *Biologia Plantarum* 44: 145-147.

Pokharia AK, Kharakwal JS, Srivastava A. 2014. Archaeobotanical evidence of millets in the Indian subcontinent with some observations on their role in the

Indus civilization. *Journal of Archaeological Science* 42: 442-455.

Reimer PJ, Baillie MGL, Bard E, Bayliss A, Beck JW, Blackwell PG, Bronk Ramsey C, Buck CE, Burr GS, Edwards RL, Friedrich M, Grootes PM, Guilderson TP, Hajdas I, Heaton TJ, Hogg AG, Hughen KA, Kaiser KF, Kromer B, McCormac FG, Manning SW, Reimer RW, Richards DA, Southon JR, Talamo S, Turney CSM, van der Plicht J, Weyhenmeyer CE. 2009., IntCal09 and Marine09 Radiocarbon age calibration curves 0-50000 years cal BP. *Radiocarbon* 51: 1111–1150.

Reimer PJ, Bard E, Bayliss A, Beck JW, Blackwell PG, Bronk Ramsey C, Buck CE, Cheng H, Edwards RL, Friedrich M, Grootes PM, Guilderson TP, Haflidason H, Hajdas I, Hatte C, Heaton TJ, Hoffmann DL, Hogg A G, Hughen KA, Kaiser KF, Kromer B, Manning SW, Niu M, Reimer RW, Richards DA, Scott EM, Southon JR, Staff RA, Turney CSM, van der Plicht J. 2013. IntCal 13 and Marine 13 radiocarbon age calibration curves 0-50000 years cal BP. *Radiocarbon* 55: 1869-1887.

Riehl S, Zeidi M, Conard NJ. 2013. Emergence of agriculture in the foothills of the Zagros Mountains of Iran. *Science* 341: 65-67.

Ren LL. 2017. A study on animal exploitation strategies from the late Neolithic to Bronze Age in northeastern Tibetan Plateau and its surrounding areas China. Doctoral Dissertation Lanzhou: Lanzhou University 70-96. In Chinese.

Spengler RN, Willcox G. 2013. Archaeobotanical results from Sarazm Tajikistan an Early Bronze Age Settlement on the edge: Agriculture and exchange.

Environmental Archaeology 18: 211-221.

Spengler RN, Frachetti M, Doumani P, Rouse L, Cerasetti B, Bullion E, Mar'yashev A. 2014. Early agriculture and crop transmission among Bronze Age mobile pastoralists of Central Eurasia. *Proceedings of the Royal Society B: Biological Sciences* 281: 20133382.

Spengler RN. 2015. Agriculture in the Central Asian Bronze Age. *Journal of World Prehistory* 28: 215-253.

Staubwasser M, Sirocko F, Grootes PM, Segl M. 2003. Climate change at the 42 ka BP termination of the Indus valley civilization and Holocene south Asian monsoon variability. *Geophysical Research Letters* 30: 1-7.

Staubwasser M, Drăgușin V, Onac BP, Assonov S, Ersek V, Hoffmann DL, Veres D. 2018. Impact of climate change on the transition of Neanderthals to modern humans in Europe. *Proceedings of the National Academy of Sciences of USA* 115: 9116-9121.

Sun HL. 2011. Palaeoenvironment Reconstruction of the Middle to Late Holocene in a High-resolution Sediment Core from Tianchi lake on Liupan Mountain. Doctoral Dissertation Lanzhou: Lanzhou University 77-89. In Chinese.

Sun HL, Bendle J, Seki O, Zhou AF. 2018. Mid-to-late Holocene hydroclimatic changes on the Chinese Loess Plateau: evidence from n-alkanes from the sediments of Tianchi Lake. *Journal of paleolimnology* 60: 511-523.

Timmermann A, Friedrich T. 2016. Late Pleistocene climate drivers of early human migration. *Nature* 538: 92.

- Wang YR. 2017. Identifying the beginning of sheep husbandry in western China. Doctoral Dissertation University of Cambridge 134-233.
- Weiss H, Courty MA, Wetterstrom W, Guichard F, Senior L, Meadow R, Curnow A. 1993. The genesis and collapse of third millennium north Mesopotamian civilization. *Science* 261: 995-1004.
- Wu WX, Zheng HB, Hou M, Ge QS. 2018. The 5.5 cal ka BP climate event population growth circumscription and the emergence of the earliest complex societies in China. *Science China Earth Sciences* 61: 134-148.
- Yang Q, Li XQ, Liu WG, Zhou XY, Zhao KL, Sun N. 2011. Carbon isotope fractionation during low temperature carbonization of foxtail and common millets. *Organic geochemistry* 42: 713-719.
- Yang Y. 2014. The Analysis of Charred Plant Seeds at Jinchankou Site and Lijiaping Site during Qijia Culture Period in the Hehuang Region China. Masteral Dissertation Lanzhou: Lanzhou University 28-30. In Chinese.
- Ye ML. 2015. Ecological adaptation of Qijia cultural agricultural development: Preliminary exploration of originagrass-agriculture: Taking the Lajia site Qinghai Province as an example. *Agricultural archaeology* 6: 19-26. In Chinese.
- Yuan J, Flad R, Luo YB. 2008. Meat-acquisition patterns in the Neolithic Yangzi river valley China. *Antiquity* 82: 351-366.
- Zeder MA. 2008. Domestication and early agriculture in the Mediterranean Basin: **Origins, diffusion, and impact**. *Proceedings of the national Academy of Sciences of USA* 105: 11597-11604.

Zhang C, Pollard AM, Rawson J, Huan LM, Liu RL, Tang XJ. 2019. China's major Late Neolithic centres and the rise of Erlitou. *antiquity* 93: 588-603.

Zhang SJ, Dong GH. 2017. Human adaptation strategies to different altitude environment during mid-late bronze age in northeast tibetan plateau. *Quaternary Sciences* 37: 696-708. In Chinese.

Zhao ZJ. 2010. Paleoethnobotany: Theories Methods and Practice. *Beijing: Science Press*. In Chinese.

Zhao ZJ. 2011. New archaeobotanic data for the study of the origins of agriculture in China. *Current Anthropology* 52: S295–S306.

Zhou BX. 1999. Animal remains from the sites of Shizhaocun and Xishanping In: Institute of Archaeology Chinese Academy of Social Sciences eds Shizhaocun and Xishanping. *Beijing: The Encyclopedia of China Publishing House* 335-339. In Chinese.

Zhou XY, Li XQ, Dodson J, Zhao KL. 2016. Rapid agricultural transformation in the prehistoric Hexi corridor China. *Quaternary International* 426: 33-41.
Table 1. Calibrated radiocarbon dates and associated information of samples from the sites in WLP.

Site	Lab number	Dated material	^{14}C age	Calibrated age (cal yr BP)		Culture	References
				1 sigma	2 sigma		
Bupo	LZU17206	Foxtail millet	4495±25	5167±113	5168±122	Majiaya	This study
Xishanping	TKa13889	Foxtail millet	4490±35	5166±117	5140±158	Majiaya	Li et al., 2007

Yujiawa	LZU171 96	Foxtail millet	4465±25	5131±14 3	5129±15 4	Late Yangsh ao	This study
Yangwa	LZU171 95	Foxtail millet	4450±25	5119±14 4	5123±15 7	Majiaya o	This study
Xishanp ing	TKa138 90	Rice	4430±10 0	5075±19 9	5077±23 6	Majiaya o	Li et al., 2007
Huidier	LZU172 08	Foxtail millet	4425±35	5049±16 2	5073±20 2	Late Yangsh ao	This study
Buzipin g	BA1108 82	Foxtail millet	4300±25	4854±10	4894±64	Majiaya o	Jia et al., 2013
Lixinzh en	LZU172 10	Foxtail millet	4185±35	4738±91	4712±12 6	Majiaya o	This study
Shizhao cun	LZU171 91	Foxtail millet	4000±25	4471±43	4470±50	Majiaya o	This study
Laohuz ui	BA1108 66	Charre d millets	3870±30	4323±81	4287±12 6	Qijia	Dong et al., 2014
Shangm iangua	LZU142 30	Foxtail millet	3845±30	4253±95	4279±12 5	Qijia	Chen et al., 2019
Jiangjia zui	BA1108 71	Broomc orn millet	3835±25	4222±65	4277±12 7	Qijia	Chen et al., 2019
Zhengji ashanlia ng	LZU172 12	Foxtail millet	3725±25	4069±75	4067±82	Qijia	This study
Qiaocun	BA1108 70	Foxtail millet	3635±25	3943±35	3972±10 0	Qijia	Chen et al., 2019

Guanzizui	BA120217	Charred millets	3630±30	3939±40	3965±111	Qijia	Dong et al., 2014
Buziping	BA110883	Foxtail millet	3610±20	3927±39	3919±59	Qijia	Jia et al., 2013
Yuanguan	LZU17207	Foxtail millet	3600±30	3914±49	3908±70	Qijia	This study
Sumiaoyuantou	BA110879	Charred millets	3600±25	3916±47	3908±64	Qijia	Dong et al., 2014
Wujiaping	LZU17193	Foxtail millet	3585±25	3877±33	3902±68	Qijia	This study
Shipocun	LZU17201	Foxtail millet	3570±35	3875±44	3850±124	Qijia	This study
Shuiquan	LZU17211	Foxtail millet	3570±25	3866±28	3849±116	Qijia	This study
Yierjidianguan	LZU17203	Foxtail millet	3520±35	3784±62	3792±94	Qijia	This study
Yaoshan	LZU17189	Wheat	2865±25	3001±54	2976±89	Siwa	This study
Longxihetan	LZU17188	Wheat	2765±30	2856±62	2863±80	Siwa	This study
Qinjiaping	LZU17197	Wheat	2545±25	2644±100	2625±122	Siwa	This study
Dongjiaping	LZU17198	Wheat	2510±25	2613±106	2614±123	Siwa	This study
Ruji	LZU17190	Wheat	2445±25	2532±152	2529±170	Siwa	This study

Laoyeshan	LZU17202	Wheat	2420±20	2412±48	2517±161	Siwa	This study
Sanjiaodi	LZU17200	Barley	2320±25	2343±8	2335±22	Siwa	This study

Table 2. Charred plant seeds identified from the 41 archaeological sites during 5500-2300 cal yr BP in **WLP.**

Site	Sample number	Soil floatation quantity (L)	Crop				Weed	Total
			<i>Setaria italica</i>	<i>Panicum miliaceum</i>	<i>Triticum aestivum</i>	<i>Hordeum vulgare</i>		
Nuanquanshan	1	20		29			5	34
Zhoujiawan	2	30	46	50				96
Chaijiaping	2	30	35	16				51
Xishanping	3	53	71	37			1	109
Shanpingli	2	45	139	72			6	217
Caikeding	1	17	5					5
Shizidao	2	38	1634	37				1671
Yangwa	3	60	1591	32			1	1624
Xiewanding	1	28	231	12			2	245
Chunshuwan	2	46	32				3	35

Yujiawa	4	106	497	183			3	683
Wayaodi ng	1	16		1				1
Shangren wan	2	40	21				1	22
Yangjiap ing	1	16	39				2	41
Douwash an	2	42	77	26			13	116
Bupo	2	52	28					28
Huidier	3	56	48	5			7	60
Lixinzhe n	4	80	209	50			3	262
Yingwu a	2	44	180	79			1	260
Shizhaoc un	2	36	51	21		1	2	75
zhengjias hanliang	4	93	270	30			7	307
Wujiapin g	2	40	855	13			7	875
Yierjidia nguandi	1	13	38	9				47
Shuiquan gou	2	50	267	45			5	317
Fanjiach eng	2	36	26	4	1		11	42
Siping	2	24	137	56		1	18	212
Sishilipu	2	33	1350	64	33		40	1487
Shipocun	1	16	257	4	3		5	269
Weishuiy u	2	27	21	4	1			26
Yuangud ui	3	55	562	10	1			573

Laoyeshan	1	18	191	500	31	9	1	732
Yaoshan	2	30	184	21	17	1	56	279
Qinjiaping	4	88	180	27	69	32	6	314
Xihetan	5	96	83	75	38	29	98	323
Sanjiaodi	3	56	88	14	62	26	3	193
Ruji	3	51	57	35	127	41	25	285
Wapenyao	2	47	112	11	163	19	28	333
Diantiandi	3	74	34	1	197	28	15	275
Dongjiaping	3	56	37	71	86	1	2	197
Majiayao	1	24	119	1	127	51	44	747
Beiposi	1	20	7	4	60	20	8	99
Total	91	1802	9809	1649	1016	259	83	1356
							4	7

Figure captions

Fig 1. (a) Distribution of prehistoric sites with radiocarbon dates in NETP (Dong et al., 2014; Yang, 2014; Chen et al., 2015). (b) Location of the 41 sampled sites in the study area in the paper. Sample sites: 1. Nuanquanshan, 2. Longxixihetan, 3. Lixinzhen, 4. Yingwuwa 5. Wujiaping, 6. Laoyeshan, 7. Yujiawa, 8. Bupo, 9. Majiayao, 10. Dongjiaping, 11. Diantiandi, 12. Qinjiaping, 13. Sishilipu, 14. Wapenyao, 15. Chunshuwan, 16. Beiposi 17. Shuiquangou, 18. Yuangudui, 19. Yierjidianguandi, 20. Huidier, 21. Wayaoding, 22. Shangrenwan, 23. Yangwa, 24. Shizidao, 25. Sanjiaodi, 26. Yangjiaping, 27. Siping, 28. Shipocun, 29.

Xiewanding, 30. Zhengjiashanliang, 31. Douwashan, 32. Weishuiyu, 33. Yaoshang, 34. Shanpingli, 35. Fanjiacheng, 36. Shizhaocun, 37. Xishanping, 38. Ruji, 39. Chaijiaping, 40. Caikeding 41. Zhoujiawan.

Fig 2. Charred plant seeds collected from WLP (scale bar: 1mm).

a. *Setaria italica*, b. *Panicum miliaceum*, c. *Triticum aestivum*, d. *Hordeum vulgare*, e. *Chenopodium album*, f. *Setaria viridis*, g. *Melilotus suaveolens*, h. *Galium tricornne*, i. *Sphaerophysa salsula*, j. *Atriplex patens*, k. *Galium tricornutum*, l. *Glycyrrhiza uralensis*, m. *Kochia scoparia*, n. *Salsola pestifer*, o. *Polygonum nepalense*, p. *Rumex acetosa*.

Fig 3. Human-environment interaction in WLP and NETP.

a. The $\delta^{18}\text{O}$ record of the Jiuxian Cave stalagmites (Cai et al., 2010), b. Paq values based-on n-alkanes in Liupan Tianchi Lake (Sun et al., 2018), c. Northern Hemisphere (30° to 90°N) temperature record compared to 1961–1990 instrumental mean temperature (Marcott et al., 2013), d. MBT/CBT-derived MAT in Liupan Tianchi Lake (Sun, 2011), e. The weight percentage of crops in WLP, f. CPD curve formed by the probability density of radiocarbon dates from charred plant seeds in WLP (Li et al., 2007; Jia et al., 2013; Dong et al., 2014; Chen et al., 2019), g. The weight percentage of crops in NETP (Chen et al., 2015), h. CPD curve formed by the probability density of radiocarbon dates from charred plant seeds in NETP (Dong et al., 2014; Chen et al., 2015).