

Bringing a time-depth perspective to collective animal behaviour

5

Dora Biro^{1*}, Takao Sasaki¹ & Steven J. Portugal²

¹Department of Zoology, University of Oxford, UK. ²School of Biological Sciences, Royal Holloway, University of London, UK

10

* Corresponding author: Biro, D. (dora.biro@zoo.ox.ac.uk).

Keywords

Collective behaviour, collective learning, decision-making, collective intelligence,

15 energetics, time-depth

20 **Abstract**

The field of collective animal behaviour examines how relatively simple, local interactions between individuals in groups combine to produce global-level outcomes. Existing mathematical models and empirical work have identified candidate mechanisms for numerous collective phenomena, but have typically focused on one-off or short-term performance. We
25 argue that feedback between collective performance and learning – giving the former the capacity to become an adaptive, and potentially cumulative, process – is a currently poorly explored, but crucial mechanism in understanding collective systems. We synthesise material ranging from swarm intelligence in social insects, through collective movements in vertebrates, to collective decision-making in animal and human groups, to propose avenues
30 for future research to identify the potential for changes in these systems to accumulate over time.

What are Collective Behaviours and How Do They Arise?

35 Some of the most impressive biological phenomena emerge out of interactions among members of animal groups. Bird flocks, fish schools and insect swarms perform highly coordinated collective movements that can encompass thousands of individuals, producing complex group-level patterns that are difficult to predict from the behaviour of only isolated individuals. Animal groups are also able to solve problems that are beyond the capacities of
40 single individuals [1] – ant colonies, for example, tackle certain types of optimisation problems so effectively that they have inspired an entire field of computer science [2]. Despite the appearance of synchronised organisation, it is increasingly well understood that no central control acts on the collective as a whole; instead, the global patterns result from simple, local interactions among the group's neighbouring members: a form of biological
45 self-organisation [3] (see Glossary). Recent years have seen a proliferation of both empirical and theoretical work on the mechanistic underpinnings of collective animal behaviour [4], with self-organisation emerging as a major principle in a variety of contexts including

collective motion [5], decision-making [6] and construction [7], activity synchronisation [8], and the spontaneous emergence of leader-follower relations [9].

50

Nonetheless, a rigorous adaptive framework is yet to be applied to collective animal behaviour: little is known about the nature of the selective forces that act at the level of the individual behavioural rules to shape pattern formation at group-level. Over shorter time-scales, and crucially for this review, no major synthesis has yet examined collective behaviour from a *time-depth perspective*: we do not know (i) what changes group-level organisation might undergo over the course of repeated executions of collective tasks, (ii) to what extent solutions arrived at collectively are retained (learned), either at the individual or at the collective level, with the potential to influence future interactions, or (iii) what effect changes in group composition, due to natural demographic processes, have on whether solutions are “inherited” from previous generations.

60

Why Time-depth?

We use the term “time-depth” as applied primarily in linguistics and archaeology, where it is used to refer to the length of time a trait in question (language, behaviour, technology, etc) has been undergoing change (e.g. [10]). Thus, implicit in the term is an appreciation that any current observations of a phenomenon are only snapshots that represent the outcome of a potentially long history of previous states. Correspondingly, we argue that in the case of collective behaviour, collective performance we observe at any given time has a history upon which its current state is contingent. Such contingencies can be rooted both phylogenetically and ontogenetically. First, natural selection can fine-tune individual interaction rules in ways that modulate global-level phenomena [3,11], even in systems with very low levels of relatedness [12]. Second, individuals can adjust their contributions as a function of, for example, the quality of a previous collective action as they perceive it. In this review we focus on the latter scenario, and examine the changes that collective phenomena can undergo over repeated performances of a collective task. Crucial to our perspective is the idea that

75

individuals can learn from their experiences of acting collectively with others, making collective behaviour a plastic process that can allow groups to adapt their collective problem-solving dynamically. In that sense, time-depth is what distinguishes collective behaviour in biological systems from those in the physical or chemical domains: the component units possess memory and are capable of learning. By considering those changes to collective outcomes that are the products of learning as a result of collective experience, rather than merely of the individual, we can pursue a novel perspective on collective animal behaviour.

The Case for Collective Learning

Although pedagogical research and developmental psychology have long acknowledged that humans interacting in a group context influence each other's learning, this has typically been framed in terms of sophisticated cognitive mechanisms such as joint attention and mental-state attribution [13]. However, the same premise – that knowledge can be *constructed from the interactions* of multiple individuals – applies equally to collective behaviour. For example, previous research has shown that during collective navigation by homing pigeon flocks, birds less well informed about the terrain nonetheless contribute to the route-finding process, and can thus improve the performance of both naïve and knowledgeable flight partners [14] (see Box 1 for more detail). We refer to this phenomenon as collective learning [15]. A theoretical treatment of this topic by Kao *et al.* [16] modelled collective learning to demonstrate that individual experience gained during collective action results in superior group decisions under a range of hypothesised environmental conditions. Empirical data on how such predictions relate to the performance of real animal groups is, however, largely lacking.

We suggest that collective learning not only influences knowledge held by individuals (and hence these individuals' subsequent behaviour whether alone or in a group setting), but also has the potential to affect *how* collective decisions are made on future occasions. For example, following a successful collective action, links between specific individuals might be

reinforced as they recognise the usefulness of the information received, or, conversely, a
105 failed collective decision might weaken bonds between individuals and promote social
reorganisation. Agent-based models suggest many interesting potential outcomes of such
reorganisation, including social stratification and elite formation [17], but the empirical
relevance of such models to real biological systems is unclear. Figure 1 summarises the
interrelationships among the different conceptual elements we have so far highlighted.

Groups as Generators, Rather than Only Repositories of Information

The progressive increase in the breadth, complexity and efficiency of cultural phenomena in
hominins is commonly described as cumulative cultural evolution (CCE) [18]. With
behavioural innovations continually building on previous innovations, CCE gives rise to
115 behaviours that go beyond what individuals are capable of inventing in a single lifetime. Such
“ratcheting” [19] is argued to have been key to the scope that culture has attained in humans
but not in other species [20]. From religion to the Mars rover, much of present-day human
behaviour and technology is the product of information accumulation over thousands of
generations.

120 Models that attempt to explain what factors might have driven CCE in hominins frequently
incorporate demography, focusing on population size or density [21,22]. These parameters
(representing the pool of social learners) determine how likely novel behaviours – generated
with a given probability – are to be retained. In a recent review, Fogarty *et al.* [23] briefly
125 suggest that these models fall short on taking into account interactions between individuals as
potential factors influencing innovation rates. We strongly agree with this suggestion and
propose it deserves much more detailed consideration. What previous approaches lack is a
role for groups as *generators*, rather than simply repositories of information upon which
culture is built and can accumulate. In other words, not only is the final product (knowledge
130 accumulation) dependent on group size, but so is the mechanism: larger groups might (i)
generate more innovations because they have a higher probability of including an innovator,

and/or (ii) generate more innovations because collective intelligence operates more strongly the more individuals contribute to problem solving.

135 As an example, laboratory studies of CCE in humans, pioneered by Caldwell and Millen
[24,25], have shown progressive improvements in solving tasks (such as building increasingly
tall towers out of spaghetti and plasticine) when these are given to successions of “micro-
societies” consisting of a mixture of previous solvers and novices. These transmission chain
designs are notable for their use of groups of participants at each stage, and are highly
140 informative in terms of outcome (the accumulation of improvements) as well as mechanisms
(emphasising features such as prosociality, teaching and collaboration [20,26]). However,
they are not explicit about the potential role that solving the task as a group might itself have
had on the generation of innovations, particularly if each link in the chain had consisted of
more than the study’s maximum of three individuals [25].

145
Issues explored above raise many interesting questions about what is necessary for collective
tasks to benefit from pooling the contributions of multiple individuals (“collective
intelligence”). What are the necessary social, ecological and cognitive prerequisites for
animal groups to generate and retain solutions to problems in ways that allow the
150 accumulation of these over time? In what measurable aspect can collective solutions
improve? In the next section we examine how we can evaluate behavioural solutions, before
returning to address these questions.

Measureable Outcomes

155 How can we measure the quality of collective performance? This question is relevant both to
researchers seeking to identify changes in said quality, and to the individuals themselves
involved in collective action (i.e. how does an individual within a group assess success and
effectiveness in a group task?). The former speaks to our ability to study changes in collective
outcomes longitudinally, and the latter to the mechanisms that would allow such changes to

160 effect learning by individual agents within the collective. As broad categories, the speed, accuracy, cohesion and energetic efficiency of collective performance are all credible candidates – theoretically detectable by individuals in collectives, and subject to adjustment as a function of individual behaviour. We illustrate each briefly below.

165 The capacity of groups to make accurate consensus decisions due to information pooling has entered popular science lore (as the “wisdom of crowds” [27]), and the relationship between group size and decision accuracy has extensive theoretical and empirical support. Shoals of fish become capable of finer-scale discriminations [28] and of better predator avoidance [29], flocks of birds select routes closer to the beeline path to their nests [30,31], and human
170 crowds move more accurately toward a target destination [32], as the number of individuals in these groups increases. Condorcet’s jury theorem, the “many wrongs” principle, and increased information processing power are typically relied on to explain the mechanism [6,33,34]. With the assumption that there is no population-level bias in opinions and that group members contribute information independently and equally, individual errors are
175 averaged out to approach the optimum, and/or the population majority tends towards the correct decision.

Often just as vital as decision accuracy, decision speed provides another measure of collective performance. This is particularly evident when under threat through predation or other forms
180 of ecological pressure, where a group’s capacity to respond rapidly is of fundamental importance. Here too increases in performance quality have been documented with increases in group size. For example, how quickly shoals of fish choose a path that avoids a predator [35] or how quickly honeybee colonies acquire and evaluate information about suitable nest sites [36] are improved by larger numbers of individuals contributing to the processing of
185 available information.

Quorums often contribute to these effects, allowing groups to switch from information gathering to rapid convergence on a decision. Cross-inhibition, one mechanism through which such convergence operates, shows interesting parallels between social insect and neuronal decision-making [37]. Although quorums link decision speed and accuracy, the two can also be involved in a trade-off. For example, much like in individual decision-making [38], decision accuracy can be traded off against decision speed: theoretically, the speed of a collective decision can be increased by decreasing the steepness of the quorum function, but this will also cause a decrease in the accuracy of the decision [33]. Ants in harsh environments, where decisions have to be made rapidly, potentially sacrificing accuracy, respond just so [39].

Since many of the benefits of social living depend on group cohesion [33,40], group fragmentation might be viewed as a sub-optimal outcome during collective action. Anti-predatory effects such as predator confusion and dilution will be compromised [41], while information-processing advantages will be correspondingly scaled back [6,36]. However, differing interests or preferences over the best course of action can generate conflicts, where individuals will typically pay a “consensus cost” for remaining with the group [42]. Under such circumstances, groups can fragment: differing preferences in the direction of travel have been shown to break up homing pigeons [43] and king penguins [44] (although, perhaps due to different balancing of long vs. short-term costs, not meerkats [45] or baboons [46]). Cohesion can also be involved in trade-offs with speed and accuracy [47]. Analyses of baboon group movements suggest that decisions are delayed when opinions within the group diverge widely [46], likely because forces maintaining cohesion compete with individual preferences, reducing decision speed.

Lastly, collective action can generate energetic savings that might be detectable to individuals. These savings can be accrued through, for example, positive aero- or hydro-dynamic interactions: crustaceans [48], fish [49], adult [50] and juvenile [51] marine

215 mammals, and birds during both flight [52] and surface swimming [53] have been shown to
benefit energetically from moving together with conspecifics. Box 3 details a case study for
flying birds.

How can Measureable Outcomes Feed Back into Collective Behaviour?

220 It seems reasonable to assume that individuals in groups are sensitive to some combination of
the measurable outcomes of collective action outlined above. Although absolute evaluation
might not be possible in many circumstances (a bird in a flock might not know if the flock is
flying an efficient route to a destination, or an ant with limited knowledge of the environment
might not be able to judge if the colony was delayed in choosing a new nest site), relative
225 judgements based on comparisons to previous group performance could be available to guide
evaluation. Based on such comparisons, individuals might, for example, (i) choose to adjust
their own contribution on subsequent occasions, (ii) redistribute the relative weighting they
assign their personal vs social information, or (iii) change the way they interact with specific
group-mates. Similarly, increases in an individual's experience or competence as a result of
230 previous collective action might affect what information it contributes and how it interacts
with others in future. We now explore examples of both these processes – adjustments based
on judgement of previous performance quality and on learning as a result of previous
collective action – with reference to theoretical and empirical examples.

235 Changing one's relative contribution to collective decisions might depend on a judgment of
the quality of one's own information. That such adjustments – a function of individual
certainty – are possible has been demonstrated in a variety of species. How well informed
human participants in a collective decision-making task judge themselves to be, influences
how readily and quickly they contribute information to the group [54]. Male bottlenose
240 dolphins perform specific behavioural signals that initiate group travel more frequently the
greater their knowledge about the optimal timing of such activity shifts [55].

Changes in the organisation of decision-making represent perhaps more subtle adjustments. Modelling work examining changes in information flow within groups over repeated
245 iterations of task solving found that links between individuals were reinforced when they judged each other to have contributed high-quality information on previous occasions [17]. In a sense, individuals chose to rely on group-mates that had proved themselves competent. Similar mechanisms might be at work in several of the systems we discuss in previous sections and in Boxes 1-4, although cases could be limited to species that have stable and
250 small enough groups, and the requisite cognitive capacities for individual recognition. Through such recognition and selective targeting of attention, the contributions not only of competent group-mates but, conversely, also of “persistent offenders” might change over time.

255 In systems that use quorums in reaching consensus decisions, quorum size can be adjusted to tune decision speed, and thus to modulate how decision speed is traded off against decision accuracy (see Box 2). Honeybee colonies vary in how they trade off speed for accuracy according to their size [36]: larger colonies invest larger numbers of scouts into discovering nest sites but make decisions at the same speed as smaller swarms, allowing for higher
260 accuracy. Combined with the observation that prior knowledge contributes to collective decisions in social insects [56], flexibility in lowering quorum size when individuals are well informed could lead to progressive improvements in colony performance in terms of speed, without sacrificing accuracy.

265 While many of the examples above deal with collective decisions, improvements in collective performance are also attainable in cases where there is no explicit “decision”. Groups of prey, for example, might streamline their escape responses following successful interactions with predators (much like certain types of collective motion rules are suggested to reduce group fragmentation following predator attacks [43]). Similarly, increases in energy savings derived
270 from moving in formation can be obtained by individuals learning, during group movements,

where best to position themselves for more efficient exploitation of the aero- or hydro-dynamic benefits offered by group travel (Box 3). On the other side of predator-prey interactions, cooperative hunting presents an interesting case study in which we suggest there is potential for collective learning and time-depth (Box 4). Where there is division of labour –

275 not only in group hunting but in more discretised roles within society such as in social insect temporal and physical castes – flexibility in the roles assumed by individuals, coupled with feedback on how well they fulfil their roles [57], can tune collective performance over time.

Finally, it is worth noting that the idea that iterative collective performance in animal groups

280 might be influenced by the group's previous states has been suggested to present intriguing parallels with neuronal processes (e.g., [58]). In both cases, interactions among populations of units, as well as the properties of the units themselves, can be tuneable as a function of prior history; the succession of collective states thus assumed can be regarded as reflecting 'collective memory' [59]. Future work linking processes such as memory formation in

285 organismal and neural collectives is likely to provide cross-disciplinary insights on both sides of this analogy [58].

When might Capacity for Time-depth be Most Useful?

Although in our descriptions above we deal with examples where time-depth is both feasible

290 and potentially operates, we acknowledge there are situations in which it will either be impossible to implement, or of limited use. First, in cases where collective outcomes are not necessarily or directly linked to mechanisms at the individual level, but are instead "emergent" properties, by definition behaviours that improve group performance cannot be learnt. Second, it might be that adjusting the collective outcome only has utility in certain

295 situations, where, for example there is need, room or capacity for improvement (Box 4). In this second case, changes might be generated and implemented flexibly, thus increasing the mechanism's functionality, and fine-tuning its effectiveness to the given scenario. The role, or best use, of time-depth is, therefore, situation dependent.

300 In a task or event, a time-depth element might be utilised to either be (a) *in progress*, to learn, innovate and problem-solve as a collective, for future use, or (b) *static*, to benefit from previous innovation and iterative interactions as a collective, to maximise potential gains to be made through working cooperatively. The propensity of a group to work collectively will require alternate functions, *in progress* or *static*, of a time-depth element, depending on the

305 task in-hand. During collective tasks where solutions are open-ended or shifting, groups composed of knowledgeable and naïve individuals might facilitate finding the best solutions (see e.g. Box 1). During such tasks, innovations (or, more simply, “noise”) from naïve individuals, added to the knowledge of those more experienced, can work together to bring about improvements in the measurable outcome. In this instance, a stable, static, society

310 would perform worse than one with immigration or demographic turnover, with the time-depth element needing to be considered *in progress*. If, however, solution quality can or has reached a plateau where no further innovation will better any measurable outcome, then a *static* state would be more effective, reducing the element of risk. For example, if a group has found a continually productive foraging site, once the best route (e.g., straightest and/or

315 safest) has been located between the foraging site and home, the best solution would be to continue benefiting from route innovations prior to that point, but to remain *static*. The decision, or feedback, to remain static and cease innovation can be spontaneous, or a consequence of a lack of demographic turnover at a given time point.

320 It is likely, therefore, there will be times and events where the potential noise from innovation could have deleterious effects. Such events might be at specific times in the annual cycle where resources are limited, or due to an energetic bottleneck whereby deleterious noise could have a significant impact on survival rates and/or energy expenditure. For example, where collective action results in energetic savings through co-operative group locomotion,

325 the situation could be considered quite different. To maximise energy savings during a long migratory flight in a V formation, an important component of success is learning the correct

positioning ([60], Box 3) and the requisite social rules for positional swapping within the V [61]. In this scenario, a stable group of ‘experts’ would be best: a *static* use of time-depth. During critical events, such as migration, innovations might be too risky.

330

Concluding Remarks

We have highlighted a hitherto largely overlooked aspect of collective animal behaviour: that many collective outcomes we observe and study at a given time might be contingent on the collective’s previous history and memory. There is evidence that collective performance –
335 measured in terms of the speed and accuracy of group decisions, group cohesion and/or energetic efficiency – can change over time, both in groups where the same members solve the same task repeatedly, and in those that experience at least partial turnovers in group membership over the course of such repetition (e.g. [15,31,62–64], Boxes 1-4).

340 Key to our argument is that if collective learning not only influences individual knowledge, but also has the potential to affect how future collective decisions are made, then we must acknowledge collective behaviour as a flexible process and explore its capacity to adapt using feedback from the group’s prior performance. We suggest that in future research on biological self-organisation and collective animal behaviour, crucial insights will be achieved by
345 focusing explicitly on the following four questions. 1. To what extent are solutions arrived at collectively retained by individuals, and to what extent do they to influence future contributions to and interactions during subsequent task solving? 2. What changes does group-level organisation undergo over repeated executions of collective tasks? 3. What effect do changes in group composition due to natural demographic processes have on solutions
350 “inherited” from previous generations and on producing innovations that modify these solutions further? 4. What role does energetic optimization play in the streamlining of collective actions? Through synthesis of these questions with mechanistic and functional studies of collective behaviour, it will be possible to illuminate in hitherto unprecedented detail how animal groups acquire, process and store information.

Acknowledgements

We thank Máté Nagy, Benjamin Pettit and Tim Guilford for useful discussions, Máté Nagy for help with preparing Figure 1, Damien Farine and two anonymous referees for valuable comments on a previous version of the manuscript. DB was supported by a Royal Society

360 University Research Fellowship.

Figure Legends

Figure 1. Schematic representation showing how different processes combine to produce time-depth in collective behaviour. Coloured circles represent individuals and thin arrows between them represent their interactions. Collective behaviour (the appearance of patterns at group-level based on interactions between individuals) combines with individual learning capacities to allow individuals to acquire novel information through their interactions with others ('collective learning'). Through repeated executions of a collective task, collective behaviour becomes iterative and personal information regarding the quality of the collective outcome continues to accumulate from each round of feedback, with the potential to inform subsequent collective action. Both adjustments based on repeated performance of collective tasks and through learning via such experiences gives collective behaviour time-depth: groups can adapt their problem-solving based on feedback detected at the individual level from the group's performance. Procedural (*how* to solve tasks) as well as content-based (*what information* to use to solve tasks) knowledge can effect these changes.

Figure I. (to be included in Box 1) Homing pigeons solving a navigational task collectively.

Photo by Zsuzsa Ákos.

Figure II. (to be included in Box 2) Ants of the species *Temnothorax rugatulus* inside their nest in the laboratory. Photo by Takao Sasaki.

Figure III. (to be included in Box 3) **(A)** Development of V formation flight in juvenile white ibis (redrawn from Fig. 1 in [65]). **(B-D)** Three-dimensional location histogram showing position of individual juvenile ibises (n=14) flying as a flock, with respect to flock centroid, measured by a 5 Hz GPS data logger. The colour scale refers to the duration (in seconds) a bird was present in each 0.25m × 0.25m grid. The sequence of histograms shows the development of organized V formation flight over time, with the birds flying in training

flights in **(B)** late July and **(C)** early August before **(D)** embarking on the first migratory flight in late August (2012). Data adapted from [60] and from online supplementary data in [61].

Figure IV. (to be included in Box 4) **(A)** Hypothesised likelihood and/or degree of usefulness

of time-depth element in collective behaviour as a function of various parameters relating to group composition and the task undertaken. We suggest that a capacity for time-depth will be least relevant in cases where groups are composed of individuals of low relatedness and are transient or unstable in composition, and where tasks are repeated only rarely within

individual lifetimes. At the other end of the scale, time-depth is envisaged as most relevant

where groups are small and stable, members have high relatedness, and the task frequently recurs. **(B)** Killer whales (Antarctic Type B) coordinate to “wave-wash” a Weddell Seal off an ice floe in Antarctica [66]. Such cooperative hunting falls at the “high relatedness, high group stability, high task frequency” end of the spectrum in (A), and hence might be a potential candidate for time-depth. Photo by John Durban, NOAA Southwest Fisheries

Science Center.

Boxes

410

Box 1. Navigational problem-solving in homing pigeon flocks: leadership hierarchies, collective learning and competence

Homing pigeons (*Columba livia*) have long served as model animals in the study of large-scale spatial cognition [67]. One of the most consistent findings emerging in recent research is that, with experience, pigeons establish idiosyncratic routes home (based on memorised chains of landmarks) that they recapitulate faithfully whenever flying solo [68]. When flying as a flock, the collective route emerges as a compromise between individuals' preferred paths via a self-organised process, often, but not always, leading to "better" (closer to the beeline) routes overall ([14,30,69], Fig. 1). Furthermore, pairwise leader-follower relations are spontaneously generated within the group, and condense into robust, fully transitive leadership hierarchies [70] that reflect the flow of information within the flock. Consequently, how these leadership hierarchies are structured will have important implications for the quality of the group's navigational performance [71,72], and changes in rank allocations have the capacity to dynamically modulate group performance.

425

Interestingly, since leaders are by definition responsible for more of the flock's navigational decisions than followers, recent work has shown that they are also the ones that learn most through the experience of moving collectively [31]. This raises the possibility – as yet unexplored – that there exists a feedback loop between leadership, learning and competence, with the potential to effect improvements in collective performance over time. In other words, although leaders might not necessarily be the most competent navigators at the outset, they improve in their roles *through the experience of leading*, which can in turn improve the flock's performance and reinforce their leadership role in future.

435

Pettit *et al.* [14] have shown that while individual birds eventually reach a plateau in the efficiency of their routes, adding a locally naïve individual as a flight partner allows the pair

to improve beyond this individual constraint. This tantalisingly suggests that collective intelligence and social (collective) learning can interact to produce increasingly efficient group solutions over successive “generations”. Input from new individuals, combined with what experienced individuals had previously learnt, effectively acts as the “innovation” upon which novel, better solutions are built. Such improvements – that go beyond the capacities of single individuals – are hallmarks of cumulative culture [18], a process so far argued to be unique to humans [20].

Key questions for future work will be to determine how flocks’ organisational structure changes as a function of individuals’ prior experiences (do leadership hierarchies become progressively more stable, more stratified, or more or less heavily weighted in favour of input by birds at the top?), and whether improvements (reflected in increasingly more efficient homing routes) can accumulate over time through iterative rounds of navigational “innovation” followed by collective learning.

Box 2. Nest emigrations in social insects: do ant colonies get better at house-hunting?

When their nest becomes uninhabitable, ants of the genus *Temnothorax* (Fig. II) make collective house-hunting decisions, which emerge out of differential recruitment efforts for different potential new nest-sites by scouts [73]. These decisions can be superior to those made by individuals, as colonies are less susceptible to error when the discrimination task is difficult [74], involves a larger number of choices [6] or in cases where a “decoy” leads to irrational decisions in single ants [75].

How the organisation and quality of house-hunting collective decisions change over repeated emigrations by the same colony has received surprisingly little attention, despite the fact that such successive events have powerful ecological relevance. Not only are colonies likely to face similar emigration problems repeatedly over their lifespan, but the process might be

undertaken after at least partial turnover in colony membership due to normal demographic processes.

Langridge *et al.* [62] were the first to examine the effect that repeated emigrations have on a colony's collective problem solving. They demonstrated that emigration time decreased with repeated task solving, with the improvement apparently due to learning by individuals. All components of the total emigration time (discovery, assessment and transport of colony-mates) were reduced upon repetition, however, interestingly, there was no change in division of labour (scouts vs non-scouts, transporters vs non-transporters) across the colony. Instead, further work by the same authors identified changes in the behaviour of ants actively involved in previous emigrations: they switched to carrying colony contents (other individuals or brood items) sooner than ants that had not previously acted as transporters [76]. Thus decision speed was accelerated. However, whether colonies could also improve in other ways – for example in the resolution of their discriminative capacity, or in their resistance to decoys – as a result of repeated task solving remained to be established. Interestingly, Sasaki and Pratt [77] showed that colonies are indeed capable of more subtle improvements: they can adapt the weighting they place on different attributes used to distinguish between potential nest sites as a function of which of these attributes had proved the more informative during previous emigrations. Again, learning by ants (specifically, how scouts changed their individual weightings for different nest attributes) is implicated in the observed improvements.

As results in both sets of studies rely on individual rather than collective learning (in other words, ants learn things through their own independent actions rather than through collective action), it seems likely that demographic turnover would limit the extent to which any improvement is able to accumulate over time in these systems. This is in contrast with cases where naïve individuals introduce novel innovations that can build upon previously reached solutions, and where learning is influenced not just by an individual's own actions but by what it experiences as a consequence of group action (e.g. Box 1). Nonetheless, much

remains to be explored with respect to *Temnothorax* collective decision-making, and if individual ants also change their interactions with others as a function of previous experience (suggested, but not confirmed in [62]) and if these interactions in turn shape learning in new recruits, then longer term effects indeed become possible.

Box 3. Energetics of group movement in bald ibis: practise makes perfect?

Aerodynamic theory predicts that when birds fly in V formations, energy savings can be achieved by capturing the upwash produced by the preceding bird – positive aerodynamic interactions occur between members of the V formation [78,79]. As impressively coordinated as such flocks appear, developmental studies reveal they do not spontaneously assemble, but result from learning by individuals in a collective setting.

The critically endangered Northern Bald Ibis (*Geronticus eremita*) is currently being reintroduced back into its central European range, a process involving imprinted birds following a micro-light para-plane containing a human foster parent [60]. Such migratory flights would traditionally be undertaken in groups comprising juveniles and adults in small family groups, implicating kin selection [80]. Training flights pre-migration are critical, particularly for juveniles, since, like in many other species, the first migration is the highest cause of mortality in the lifespan of an individual [81].

The onset of V formation in juvenile birds post-fledging had previously been investigated in American White Ibis (*Eudocimus albus*), and was assumed to develop through repeated interactions and flights with adult birds [65]. During the course of the observations, the tendency of juveniles to fly in formation increased from 17.8% of all juveniles immediately post fledging in late June, to 88.0% of juveniles by late August (Fig. IIIA). Out of 64,000 observations, only once was a juvenile seen flying out of a mixed-age flock, suggesting the presence of adult birds plays a role in the development of formation flight in young birds.

The imprinted Northern Bald Ibis, however, present a different scenario. Unlike in a wild-type setting, the imprinted ibis did not have knowledgeable leaders to follow or learn from – there were no adult birds to demonstrate V formation flight, and no experienced individuals to impart knowledge via social interactions. Using biologging technology [60], it was possible to document the onset of V formation flight in the young birds (Fig. IIIB-D). Successive training flights, followed by actual migratory flight, show a clear and gradual move from apparently uncoordinated flight, akin to that of cluster flight in pigeons [82] to the distinctive V formation. While it is possible the delayed onset of formation flight is linked to flight capabilities, and younger birds do not have the requisite skills to fly in such a controlled manner, the results do suggest that a group of naïve birds are able to self-sort over a period of time, and learn the optimal positions to maximise upwash capture. It is likely that positive feedback fine-tunes positioning within the flock, while the motivation to fly in a V is genetically determined [64]. The group was able to work as a collective to progressively find not only the most energetically profitable flock shape, but also and where, within that flock, each bird should be optimally positioned.

Box 4. Candidates for time-depth?

Animal groups come in many shapes and sizes, and the degree of usefulness of a time-depth component to collective behaviour is likely to vary along with certain key parameters. We suggest that the usefulness and likelihood of time-depth will primarily be related to three important factors: (a) levels of inter-individual conflict within the collective (itself linked to the relatedness of individuals comprising the group), (b) stability of group membership, and (c) regularity of the collective task undertaken (Fig. IVA). A group is unlikely to benefit from a capacity for time-depth if group members are (a) unrelated, (b) fluid in composition, and (c) only perform a given collective task sporadically. For example, collectively migrating passerines that travel in large clusters of unrelated individuals are unlikely to accumulate significant improvements over time: the task is so rarely undertaken and the fission-fusion

nature of groups means time invested in developing individual roles or expertise would not be
550 recouped in the absence of recurring interactions with the same individuals, or with
individuals with closely aligned interests. Similarly, large group sizes can negate advantages
that time-depth can bring, if sheer numbers mean repeated interactions between individuals
will be limited, and feedback between individual and group performance will not be
transparent. In contrast, a group is likely to benefit greatly from time-depth if members are (a)
555 related, (b) static, and (c) regularly perform tasks as a collective.

A good example of the latter scenario is provided by co-operative hunting (Fig. IVB).
Cooperative hunting has been reported in several mammal species [83] and one bird [84]. It is
particularly prevalent in the delphinids (e.g. [85,86]), with a variety of hunting approaches
560 utilized depending on prey type, habitat and group size, with some dolphin species even
hunting cooperatively with humans [87]. Many of the cooperative hunting strategies (e.g.
intentional beach stranding [88,89] and pack ice breaking [66,90]) used by dolphins appear
region- or pod- specific [66,89], suggesting an element of culture in cetacean society [91,92].
Furthermore, delphinids exhibit role specialization, where specific group members repeatedly
565 take the same role over many years in each cooperative hunt. Such division of labour within a
stable social group potentially allows an individual to perfect its role. It remains unclear why
a division of labour with role specialization is so rare in species that hunt cooperatively. One
theory proposes that practice might not improve performance sufficiently to warrant such role
specialization [85]. Why it should prevail in marine mammals is likely to be linked to prey
570 diversity, prey biomass, mobility, and crucially, practice rewards [93]. In cooperative group
hunters, some highly skilled individuals can exert more influence during hunts. The full effect
that these ‘keystone individuals’ [94] have, and most importantly, how long their influence
lingers after their departure, is a topic of current research effort. What remains unknown is
how such cooperative hunting techniques improve over time, both within the lifespan of an
575 individual, and over successive generations. As such, cooperative collective hunting in

cetaceans can potentially offer an intriguing future case study for examining time-depth in collective action.

580 **Glossary Box**

Collective behaviour: behaviour observed at one level of a biological, physical or chemical system that emerges from interactions between lower-level units of the system. When these units comprise whole organisms (animals), collective patterns are those that are observed at
585 the level of the social group.

Collective intelligence: shared or group intelligence that emerges from pooling information from many individuals.

Collective learning: the process of acquiring knowledge through interactive mechanisms where individual knowledge is shared. The content of what is learnt is generated through co-
590 action or interactions between individuals, and are thus unavailable to the same individuals when learning alone.

Cumulative culture: The accumulation of sequential modifications over time, and typically over generations, in culturally transmitted traits (i.e., those passed on through social learning) in a population. Cumulative cultural evolution is often likened to a ratchet-like effect, where
595 successful iterations are maintained until they are improved upon, reflected in incremental increases in the efficiency and/or complexity of the behaviour.

Energetics: the study or exploitation of energy contained in chemical bonds. In respiration, some fraction of this energy is converted into biologically useful forms for biosynthesis, membrane transport, muscle contraction, nerve conduction, movement etc.

600 **Innovation:** a process resulting in new or modified behaviour, which can be learnt by the innovator, by others it acts collectively with, or by neither.

Quorum: the minimum number of individuals that that need to agree on a course of action for others in the group to copy them. Quorums speed up decisions by effectively ending

deliberations when the group is in the process of deciding between multiple options.

605 **Self-organisation:** the emergence of group-level patterns from local interactions between the group's neighbouring component units, resulting in organised behaviour without global or centralized control.

Time-depth: the interpretation of a trait in question (language, behaviour, technology, process, species, etc) as the product of a series of changes in state that it has undergone over
610 time. Changes can be due to selective forces acting on evolutionary timescales, or to learning in the lifetimes of individuals or groups.

References

- 1 Krause, J. *et al.* (2009) Swarm intelligence in animals and humans. *Trends Ecol. Evol.*
615 25, 28–34
- 2 Dorigo, M. *et al.* (2006) Ant colony optimization. *IEEE Comput. Intell. Mag.* 1, 28–39
- 3 Camazine, S. *et al.* (2001) *Self-organisation in biological systems*, Princeton University Press.
- 4 Sumpter, D.J.T. (2010) *Collective Animal Behavior*, Princeton University Press.
- 620 5 Vicsek, T. and Zafeiris, A. (2012) Collective motion. *Phys. Rep.* 517, 71–140
- 6 Sasaki, T. and Pratt, S.C. (2012) Groups have a larger cognitive capacity than individuals. *Curr. Biol.* 22, R827–R829
- 7 Latty, T. *et al.* (2011) Structure and formation of ant transportation networks. *J. R. Soc. Interface* 8, 1298–1306
- 625 8 Néda, Z. *et al.* (2000) The sound of many hands clapping. *Nature* 403, 849–850
- 9 Couzin, I.D. *et al.* (2005) Effective leadership and decision-making in animal groups on the move. *Nature* 433, 513–516
- 10 Renfrew, C. *et al.* (2000) *Time depth in historical linguistics.*, McDonald Institute for Archaeological Research.
- 630 11 Seeley, T.D. (1997) Honey bee colonies are group-level adaptive units. *Am. Nat.* 150, S22–S41

- 12 Farine, D.R. and Montiglio, P. (2015) From Individuals to Groups and Back : The Evolutionary Implications of Group Phenotypic Composition. *Trends Ecol. Evol.* 30, 609–621
- 635 13 Tomasello, M. *et al.* (1993) Cultural learning. *Behav. Brain Sci.* 16, 495–552
- 14 Pettit, B. *et al.* (2013) Not just passengers: pigeons, *Columba livia*, can learn homing routes while flying with a more experienced conspecific. *Proc. R. Soc. B Biol. Sci.* 280, 20122160
- 15 Flack, A. and Biro, D. (2013) Collective learning in route navigation. *Commun. Integr. Biol.* 6, e26521
- 640 16 Kao, A.B. *et al.* (2014) Collective Learning and Optimal Consensus Decisions in Social Animal Groups. *PLoS Comput. Biol.* 10, e1003762
- 17 Dávid-Barrett, T. and Dunbar, R.I.M. (2014) Social elites can emerge naturally when interaction in networks is restricted. *Behav. Ecol.* 25, 58–68
- 645 18 Boyd, R. and Richerson, P.J. (1988) *Culture and the Evolutionary Process*, University of Chicago Press.
- 19 Tomasello, M. (1999) *The Cultural Origins of Human Cognition*, Harvard University Press.
- 20 Dean, L.G. *et al.* (2014) Human cumulative culture: a comparative perspective. *Biol. Rev.* 89, 284–301
- 650 21 Aoki, K. *et al.* (2011) Rates of cultural change and patterns of cultural accumulation in stochastic models of social transmission. *Theor. Popul. Biol.* 79, 192–202
- 22 Henrich, J. (2004) Demography and Cultural Evolution: How Adaptive Cultural Processes can Produce Maladaptive Losses - the Tasmanian case. *Am. Antiq.* 69, 197–
- 655 214
- 23 Fogarty, L. *et al.* (2015) Cultural Evolutionary Perspectives on Creativity and Human Innovation. *Trends Ecol. Evol.* 30, 736–754
- 24 Caldwell, C.A. and Millen, A.E. (2008) Studying cumulative cultural evolution in the laboratory. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 363, 3529–39

- 660 25 Caldwell, C.A. and Millen, A.E. (2010) Human cumulative culture in the laboratory:
Effects of (micro) population size. *Learn. Behav.* 38, 310–318
- 26 Dean, L.G. *et al.* (2012) Identification of the social and cognitive processes underlying
human cumulative culture. *Science* 23, 1114–1118
- 27 Surowiecki, J. (2004) *The Wisdom of Crowds*, Random House Inc.
- 665 28 Sumpter, D.J.T. *et al.* (2008) Consensus Decision Making by Fish. *Curr. Biol.* 18,
1773–1777
- 29 Ward, A.J.W. *et al.* (2008) Quorum decision-making facilitates information transfer in
fish shoals. *Proc. Natl. Acad. Sci. U. S. A.* 105, 6948–6953
- 30 Dell’Ariccia, G. *et al.* (2008) Flock flying improves pigeons’ homing: GPS track
670 analysis of individual flyers versus small groups. *Anim. Behav.* 76, 1165–1172
- 31 Pettit, B. *et al.* (2015) Speed Determines Leadership and Leadership Determines
Learning during Pigeon Flocking. *Curr. Biol.* 25, 3132–3137
- 32 Faria, J.J. *et al.* (2009) Navigation in human crowds; testing the many-wrongs
principle. *Anim. Behav.* 78, 587–591
- 675 33 Sumpter, D.J.. and Pratt, S.C. (2009) Quorum responses and consensus decision
making. *Philos. Trans. R. Soc. B Biol. Sci.* 364, 743–753
- 34 Simons, A.M. (2004) Many wrongs: The advantage of group navigation. *Trends Ecol.
Evol.* 19, 453–455
- 35 Ward, A.J.W. *et al.* (2011) Fast and accurate decisions through collective vigilance in
680 fish shoals. *Proc. Natl. Acad. Sci. U. S. A.* 108, 2312–5
- 36 Schaerf, T.M. *et al.* (2013) Do small swarms have an advantage when house hunting?
The effect of swarm size on nest-site selection by *Apis mellifera*. *J. R. Soc. Interface*
10, 20130533
- 37 Seeley, T.D. *et al.* (2012) Stop Signals Provide Cross Inhibition in Collective
685 Decision-Making by Honeybee Swarms. *Science* 335, 108–111
- 38 Chittka, L. *et al.* (2009) Speed-accuracy tradeoffs in animal decision making. *Trends
Ecol. Evol.* 24, 400–7

- 39 Franks, N.R. *et al.* (2003) Speed versus accuracy in collective decision making. *Proc. R. Soc. B Biol. Sci.* 270, 2457–2463
- 690 40 Krause, J. and Ruxton, G.D. (2002) *Living in Groups*, OUP Oxford.
- 41 Ballerini, M. *et al.* (2008) Interaction ruling animal collective behavior depends on topological rather than metric distance: evidence from a field study. *Proc. Natl. Acad. Sci. U. S. A.* 105, 1232–7
- 42 Conradt, L. and Roper, T.J. (2005) Consensus decision making in animals. *Trends Ecol. Evol.* 20, 449–456
- 695 43 Biro, D. *et al.* (2006) From Compromise to Leadership in Pigeon Homing. *Curr. Biol.* 16, 2123–2128
- 44 Nesterova, A.P. *et al.* (2014) Resolution of navigational conflict in king penguin chicks. *Anim. Behav.* 93, 221–228
- 700 45 Bousquet, C.A.H. and Manser, M.B. (2011) Resolution of experimentally induced symmetrical conflicts of interest in meerkats. *Anim. Behav.* 81, 1101–1107
- 46 Strandburg-Peshkin, A. *et al.* (2015) Shared decision-making drives collective movement in wild baboons. *Science* 348, 1358–1361
- 47 Franks, N.R. *et al.* (2013) Speed-cohesion trade-offs in collective decision making in ants and the concept of precision in animal behaviour. *Anim. Behav.* 85, 1233–1244
- 705 48 Bill, R.G. and Herrnkind, W.F. (1976) Drag reduction by formation movement in spiny lobsters. *Science* 193, 1146–8
- 49 Liao, J.C. *et al.* (2003) Fish exploiting vortices decrease muscle activity. *Science* 302, 1566–1569
- 710 50 Williams, T.M. *et al.* (1992) Travel at low energetic cost by swimming and wave-riding bottlenose dolphins. *Nature* 355, 821–3
- 51 Noren, S.R. *et al.* (2008) Hitching a ride: the formation locomotion strategy of dolphin calves. *Funct. Ecol.* 22, 278–283
- 52 Weimerskirch, H. *et al.* (2001) Energy saving in flight formation. *Nature* 413, 697–8
- 715 53 Fish, F.E. (1995) Kinematics of Ducklings Swimming in Formation - Consequences of

- Position. *J. Exp. Zool.* 273, 1–11
- 54 Kurvers, R.H.J.M. *et al.* (2015) Self-organized flexible leadership promotes collective intelligence in human groups. *R. Soc. Open Sci.* 2, 150222
- 55 Lusseau, D. and Conradt, L. (2009) The emergence of unshared consensus decisions in
720 bottlenose dolphins. *Behav. Ecol. Sociobiol.* 63, 1067–1077
- 56 Stroeymeyt, N. *et al.* (2010) Improving decision speed, accuracy and group cohesion through early information gathering in house-hunting ants. *PLoS One* 5, 10
- 57 Oster, G.F. and Wilson, E.O. (1978) *Caste and Ecology in the Social Insects*, Princeton University Press.
- 725 58 Couzin, I.D. (2009) Collective cognition in animal groups. *Trends Cogn. Sci.* 13, 36–43
- 59 Couzin, I.D. *et al.* (2002) Collective memory and spatial sorting in animal groups. *J. Theor. Biol.* 218, 1–11
- 60 Portugal, S.J. *et al.* (2014) Upwash exploitation and downwash avoidance by flap
730 phasing in ibis formation flight. *Nature* 505, 399–402
- 61 Voelkl, B. *et al.* (2015) Matching times of leading and following suggest cooperation through direct reciprocity during V-formation flight in ibis. *Proc. Natl. Acad. Sci. U. S. A.* 112, 2115–20
- 62 Langridge, E.A. *et al.* (2004) Improvement in collective performance with experience
735 in ants. *Behav. Ecol. Sociobiol.* 56, 523–529
- 63 Caldwell, C.A. and Millen, A.E. (2008) Experimental models for testing hypotheses about cumulative cultural evolution. *Evol. Hum. Behav.* 29, 165–171
- 64 Mueller, T. *et al.* (2013) Social learning of migratory performance. *Science* 341, 999–1002
- 740 65 Petit, D.R. and Bildstein, K.L. (1986) Development of formation flying in juvenile white ibises (*Eudocimus albus*). *Auk* 103, 244–246
- 66 Pitman, R.L. and Durban, J.W. (2012) Cooperative hunting behavior, prey selectivity and prey handling by pack ice killer whales (*Orcinus orca*), type B, in Antarctic

- Peninsula waters. *Mar. Mammal Sci.* 28, 16–36
- 745 67 Wallraff, H.G. (2005) *Avian Navigation: Pigeon Homing as a Paradigm*, Springer
Science & Business Media.
- 68 Biro, D. *et al.* (2004) Familiar route loyalty implies visual pilotage in the homing
pigeon. *Proc. Natl. Acad. Sci. U. S. A.* 101, 17440–3
- 69 Biro, D. *et al.* (2006) From Compromise to Leadership in Pigeon Homing. *Curr. Biol.*
750 16, 2123–2128
- 70 Nagy, M. *et al.* (2010) Hierarchical group dynamics in pigeon flocks. *Nature* 464,
890–893
- 71 Flack, A. *et al.* (2015) Modelling group navigation: transitive social structures
improve navigational performance. *J. R. Soc. Interface* 12, 20150213
- 755 72 Bode, N.W.F. *et al.* (2011) The impact of social networks on animal collective motion.
Anim. Behav. 82, 29–38
- 73 Pratt, S.C. *et al.* (2002) Quorum sensing, recruitment, and collective decision-making
during colony emigration by the ant *Leptothorax albipennis*. *Behav. Ecol. Sociobiol.*
52, 117–127
- 760 74 Sasaki, T. *et al.* (2013) Ant colonies outperform individuals when a sensory
discrimination task is difficult but not when it is easy. *Proc. Natl. Acad. Sci.* 110,
13769–13773
- 75 Sasaki, T. and Pratt, S.C. (2011) Emergence of group rationality from irrational
individuals. *Behav. Ecol.* 22, 276–281
- 765 76 Langridge, E.A. *et al.* (2007) How experienced individuals contribute to an
improvement in collective performance in ants. *Behav. Ecol. Sociobiol.* 62, 447–456
- 77 Sasaki, T. and Pratt, S.C. (2013) Ants learn to rely on more informative attributes
during decision-making. *Biol. Lett.* 9, 20130667
- 78 May, R.M. (1979) Flight formations in geese and other birds. *Nature* 282, 778–780
- 770 79 Lissaman, P.B.S. and Shollenberger, C.A. (1970) Formation flight of birds. *Science*
168, 1003–1005

- 80 Andersson, M. and Wallander, J. (2004) Kin selection and reciprocity in flight
formation? *Behav. Ecol.* 15, 158–162
- 81 Lack, D.L. (1954) *The natural regulation of animal numbers*, Clarendon Press.
- 775 82 Usherwood, J.R. *et al.* (2011) Flying in a flock comes at a cost in pigeons. *Nature* 474,
494–497
- 83 Bailey, I. *et al.* (2013) Group hunting within the Carnivora: physiological, cognitive
and environmental influences on strategy and cooperation. *Behav. Ecol. Sociobiol.* 67,
1–17
- 780 84 Bednarz, J.C. (1988) Cooperative Hunting Harris' Hawks (*Parabuteo unicinctus*).
Science 239, 1525–1527
- 85 Gazda, S.K. *et al.* (2005) A division of labour with role specialization in group-
hunting bottlenose dolphins (*Tursiops truncatus*) off Cedar Key, Florida. *Proc. Biol.*
Sci. 272, 135–140
- 785 86 Tardin, R.H.O. *et al.* (2011) Coordinated feeding tactics of the Guiana dolphin, *Sotalia*
guianensis (Cetacea: Delphinidae), in Ilha Grande Bay, Rio de Janeiro, Brazil. *Zool.*
(Curitiba, Impresso) 28, 291–296
- 87 Neil, D.T. (2002) Cooperative fishing interactions between Aboriginal Australians and
dolphins in eastern Australia. *Anthrozoos A Multidiscip. J. Interact. People Anim.* 15,
3–18
- 790 88 Lopez, J.C. and Lopez, D. (1985) Killer Whales (*Orcinus orca*) of Patagonia, and
Their Behavior of Intentional Stranding While Hunting Nearshore. *J. Mammal.* 66,
181–183
- 89 Guinet, C. (1991) Intentional stranding apprenticeship and social play in killer whales
(*Orcinus orca*). *Can. J. Zool. Can. Zool.* 69, 2712–2716
- 795 90 Visser, I.N. *et al.* (2008) Antarctic peninsula killer whales (*Orcinus orca*) hunt seals
and a penguin on floating ice. *Mar. Mammal Sci.* 24, 225–234
- 91 Rendell, L. and Whitehead, H. (2001) Culture in whales and dolphins. *Behav. Brain*
Sci. 24, 309–82

- 800 92 Cantor, M. and Whitehead, H. (2013) The interplay between social networks and
culture: theoretically and among whales and dolphins. *Philos. Trans. R. Soc. B Biol.*
Sci. 368, 20120340
- 93 Connor, R.C. (2001) Individual foraging specializations in marine mammals: Culture
and ecology. *Behav. Brain Sci.* 24, 329–330
- 805 94 Pruitt, J.N. and Pinter-Wollman, N. (2015) The legacy effects of keystone individuals
on collective behaviour scale to how long they remain within a group. *Proc. R. Soc. B*
Biol. Sci. 282, 20151766

Figure 1




Figure I (Box 1)



Figure II (Box 2)



Figure III (Box 3)




Figure IV (Box 4)

