

Background invertebrate herbivory on dwarf birch (*Betula glandulosa-nana* complex) increases with temperature and precipitation across the tundra biome

Isabel C BARRIO^{1*}, Elin LINDÉN², Mariska TE BEEST², Johan OLOFSSON², Adrian ROCHA³, Eeva M SOININEN⁴, Juha M ALATALO⁵, Tommi ANDERSSON⁶, Ashley ASMUS⁷, Julia BOIKE⁸, Kari Anne BRÅTHEN⁴, John P BRYANT⁹, Agata BUCHWAL^{10,11}, C Guillermo BUENO¹², Katherine S CHRISTIE¹³, Yulia V DENISOVA¹⁴, Dagmar EGELKRAUT², Dorothee EHRICH⁴, LeeAnn FISHBACK¹⁵, Bruce C FORBES¹⁶, Maite GARTZIA¹⁷, Paul GROGAN¹⁸, Martin HALLINGER¹⁹, Monique M P D HEIJMANS²⁰, David S HIK²¹, Annika HOFGAARD²², Milena HOLMGREN²³, Toke T HØYE²⁴, Diane C HUEBNER²⁵, Ingibjörg Svala JÓNSDÓTTIR^{1,26}, Elina KAARLEJÄRVI^{2,27}, Timo KUMPULA²⁸, Cynthia Y M J G LANGE²⁹, Jelena LANGE³⁰, Esther LÉVESQUE³¹, Juul LIMPENS²⁰, Marc MACIAS-FAURIA³², Isla MYERS-SMITH³³, Erik J van NIEUKERKEN³⁴, Signe NORMAND³⁵, Eric S POST³⁶, Niels Martin SCHMIDT³⁷, Judith SITTERS^{2,38}, Anna SKORACKA³⁹, Alexander SOKOLOV^{40,41}, Natalya SOKOLOVA^{40,41}, James D M SPEED⁴², Lorna E STREET⁴³, Maja K SUNDQVIST^{2,44}, Otso SUOMINEN⁶, Nikita TANANAEV⁴⁵, Jean-Pierre TREMBLAY⁴⁶, Christine URBANOWICZ⁴⁷, Sergey A UVAROV⁴⁸, David WATTS⁴⁹, Martin WILMKING³⁰, Philip A WOOLEY⁴², Heike H ZIMMERMANN⁸, Vitali ZVEREV⁵⁰, Mikhail V KOZLOV⁵⁰

¹Department of Life and Environmental Sciences, University of Iceland, Sturlugata 7 101 Reykjavík, Iceland

²Department of Ecology and Environmental Science, Umeå University, SE-901 87 Umeå, Sweden

³Department of Biological Sciences and the Environmental Change Initiative, University of Notre Dame, Notre Dame, IN 46556, USA

⁴Department of Arctic and Marine Biology, UiT – The Arctic University of Norway, NO-9037, Tromsø, Norway

⁵Qatar University, Department of Biological and Environmental Sciences, College of Arts and Sciences, Doha, Qatar

⁶Kevo Subarctic Research Institute, Biodiversity Unit, University of Turku, FI-20014, Turku, Finland

⁷Department of Biology, The University of Texas at Arlington, Arlington, Texas, USA

⁸Alfred Wegener Institute for Polar and Marine Research, Telegrafenberg A43, 14473 Potsdam, Germany

⁹Institute of Arctic Biology, University of Alaska Fairbanks, 997775-7000 Fairbanks, Alaska, USA

¹⁰Adam Mickiewicz University, Institute of Geocology and Geoinformation, Dziejelowa 21, 61-680 Poznan, Poland

¹¹University of Alaska Anchorage, Department of Biological Sciences, 3151 Alumni Loop, 99508 Anchorage, Alaska, USA

¹²Institute of Ecology and Earth Sciences, Department of Botany, University of Tartu, Lai 40, Tartu, 51005, Estonia

¹³Science Department, Alaska SeaLife Center, 301 Railway Ave., 99664 Seward, Alaska, USA

¹⁴Nenets Agrarian-Economic Technical School, Studencheskaya 1, 166000 Naryan-Mar, Russia.

¹⁵Churchill Northern Studies Centre, PO Box 610, R0B 0E0 Churchill, Manitoba, Canada

¹⁶Arctic Centre, University of Lapland, Box 122, FI-96101 Rovaniemi, Finland

¹⁷Pyrenean Institute of Ecology (CSIC). Avda. Nuestra Señora de la Victoria s/n, 22700 Jaca, Spain

¹⁸Department of Biology, Queen's University, Kingston, Ontario K7L 3N6, Canada

¹⁹Swedish Agricultural University, Ullsväg 16, 75651 Uppsala, Sweden

²⁰Plant Ecology and Nature Conservation Group, Wageningen University & Research, Droevendaalsesteeg 3, NL-6708 PB Wageningen, the Netherlands

²¹Department of Biological Sciences, University of Alberta, Edmonton, Alberta, T5N 0R5, Canada

²²Norwegian Institute for Nature Research, NO-7485 Trondheim, Norway

²³Resource Ecology Group, Wageningen University & Research, Droevendaalsesteeg 3, 6708 PB Wageningen, The Netherlands

²⁴Arctic Research Centre and Department of Bioscience, Aarhus University, Grenåvej 14, DK-8410 Rønde, Denmark

²⁵Department of Biology & Wildlife, University of Alaska Fairbanks, 982 N Koyukuk Drive, 101 Murie, 99775-6100 Fairbanks, USA

²⁶University Centre in Svalbard (UNIS), Longyearbyen NO-9171, Norway

²⁷Department of Biology, Vrije Universiteit Brussel (VUB), Pleinlaan 2, 1050 Brussels, Belgium

²⁸Department of Geographical and Historical Studies, University of Eastern Finland, 80101 Joensuu, Finland

²⁹Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Wageningen, The Netherlands

³⁰Institute of Botany and Landscape Ecology, University Greifswald, 17487 Greifswald, Germany

³¹Université du Québec à Trois-Rivières and Centre d'études nordiques, Trois-Rivières, Québec, G9A 5H7, Canada

³²School of Geography & the Environment, University of Oxford, OX1 3QY Oxford, United Kingdom

³³School of GeoSciences, King's Buildings, West Mains Road, University of Edinburgh, Edinburgh, EH9 3FF, United Kingdom

³⁴Naturalis Biodiversity Center, PO Box 9517, 2300 RA Leiden, The Netherlands
³⁵Section for Ecoinformatics and Biodiversity, Department of Bioscience, Aarhus University, Ny Munkegade 114, DK-8000 Aarhus C, Denmark
³⁶Department of Wildlife, Fish, and Conservation Biology, University of California, Davis, USA
³⁷Arctic Research Centre, Department of Bioscience, Aarhus University, 4000 Roskilde, Denmark
³⁸Ecology and Biodiversity, Department Biology, Vrije Universiteit Brussel, 1050 Brussels, Belgium
³⁹Population Ecology Lab, Institute of Environmental Biology, Faculty of Biology, Adam Mickiewicz University in Poznań, Umultowska 89, 61–614 Poznań, Poland
⁴⁰Arctic Research Station, Institute of Plant and Animal Ecology, Ural Branch of Russian Academy of Sciences, Zelenaya Gorka Str., 21, 629400, Labytnangi, Russia
⁴¹Arctic Research Center of the Yamal-Nenets Autonomous District, Salekhard, Russia
⁴²Department of Natural History, NTNU University Museum, Norwegian University of Science and Technology, 7491 Trondheim, Norway
⁴³Environmental Sciences, School of Life Sciences, Heriot-Watt University, Edinburgh EH14 4AS, UK
⁴⁴The Center for Macroecology, Evolution and Climate, The Natural History Museum of Denmark, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen Ø, Denmark
⁴⁵Melnikov Permafrost Institute, Siberian Branch, Russian Academy of Sciences, Yakutsk, Russia
⁴⁶Department of Biology, Centre for Nordic Studies and Centre for Forest Research, Université Laval, G1V 0A6, Québec, QC, Canada
⁴⁷Department of Biological Sciences, Dartmouth College, Hanover, NH, USA
⁴⁸Nenets Museum of Local History, Pobedy 5, 166000 Naryan-Mar, Russia
⁴⁹Intercollege Graduate Degree Program in Ecology and Department of Biology, The Pennsylvania State University, University Park, Pennsylvania 16802, USA
⁵⁰Section of Ecology, Department of Biology, University of Turku, FI-20014, Turku, Finland

*corresponding author: ICB icbarrio@gmail.com

ABSTRACT

Chronic, low intensity herbivory by invertebrates, termed background herbivory, has been understudied in tundra, yet its impacts are likely to increase in a warmer Arctic. The magnitude of these changes is however hard to predict as we know little about the drivers of current levels of invertebrate herbivory in tundra. We assessed the intensity of invertebrate herbivory on a common tundra plant, the dwarf birch (*Betula glandulosa-nana* complex), and investigated its relationship to latitude and climate across the tundra biome. Leaf damage by defoliating, mining and gall-forming invertebrates was measured in samples collected from 192 sites at 56 locations. Our results indicate that invertebrate herbivory is nearly ubiquitous across the tundra biome but occurs at low intensity. On average, invertebrates damaged 11.2% of the leaves and removed 1.4% of total leaf area. The damage was mainly caused by external leaf feeders, and most damaged leaves were only slightly affected (12% leaf area lost). Foliar damage was consistently positively correlated with mid-summer (July) temperature and, to a lesser extent, precipitation in the year of data collection, irrespective of latitude. Our models predict that, on average, foliar losses to invertebrates on dwarf birch are likely to increase by 6-7% over the current levels with a 1 °C increase in summer temperatures. Our results show that invertebrate herbivory on dwarf birch is small in magnitude but given its prevalence and dependence on climatic variables, background invertebrate herbivory should be included in predictions of climate change impacts on tundra ecosystems.

- 93 **Keywords:** background insect herbivory, climate change, externally feeding defoliators, Latitudinal Herbivory
- 94 Hypothesis, leaf damage, leaf miners, gall-makers, macroecological pattern

INTRODUCTION

The role of invertebrate herbivores in tundra ecosystems has been understudied (Haukioja 1981). Admittedly, the proportion of herbivore taxa among invertebrates is lower in Arctic regions than at lower latitudes (Danks 1986), and invertebrate herbivores generally occur at relatively low abundances in tundra (Haukioja 1981). However, outbreaks of invertebrate herbivores have been well documented in the forest-tundra ecotone (Jepsen et al. 2008; Kaukonen et al. 2013) and occasionally in tundra (Post and Pedersen 2008). These massive defoliation events have large impacts on subarctic birch forests, enhancing resource turnover through deposition of frass and carcasses to the soil (Kaukonen et al. 2013) and causing vegetation shifts from forested to open conditions with consequences for ecosystem functioning and trophic interactions (Jepsen et al. 2013; Olofsson et al. 2013; Parker et al. 2016). In contrast, chronic leaf consumption by invertebrate herbivores when they occur at low densities, termed background herbivory (Kozlov and Zvereva 2017), has long been assumed to be unimportant, especially in the Arctic (Batzli et al. 1980; Haukioja 1981). However, recent studies in boreal forest indicate that chronic invertebrate herbivory can have stronger impacts on plant growth in the long term than infrequent bouts of severe damage (Zvereva et al. 2012), and can play a major role in ecosystem-level nutrient cycling (Metcalf et al. 2016). Yet, the extent, drivers and consequences of background herbivory across the tundra biome remain unquantified (Kozlov et al. 2015b).

The Latitudinal Herbivory Hypothesis (LHH) suggests that the intensity of herbivory should decrease with increasing latitude (Coley and Aide 1991; Johnson and Rasmann 2011). In its original formulation, it was argued that a wider diversity of specialist herbivores in tropical areas, together with warmer temperatures and a longer growing season, could lead to increased herbivory rates at lower latitudes. Empirical evidence has found support for LHH at the global scale, demonstrating that invertebrate herbivory of woody plants is generally lower in the polar regions than in temperate and tropical zones (Kozlov et al. 2015a). However, the generality of the LHH across plant functional types, invertebrate species and spatial scales is still debated (Moles et al. 2011; Anstett et al. 2016; Zhang et al. 2016). Latitudinal patterns with increased levels of herbivory at lower latitudes have been described for some species of plants and groups of insect herbivores (Kozlov 2008; Pennings et al. 2009; Moreira et al. 2015; Kozlov et al. 2016), but lack of latitudinal trends (Andrew and Hughes 2005; Kozlov 2008; Pennings et al. 2009), higher levels of herbivory at higher latitudes (Moreira et al. 2015; Kozlov et al. 2016), or even dome-shaped patterns (Kozlov et al. 2015a) have been described for others. Moreover, such studies have typically been conducted in temperate regions; whether latitudinal patterns can be found within the tundra biome remains unknown. Similarly, the mechanisms behind these macroecological patterns in background herbivory are poorly understood. Biotic and abiotic factors vary with latitude and this variability may promote variation in herbivory. For example, it has been suggested that, along with direct effects of climate, latitudinal gradients in herbivory may be shaped by changes in plant defensive chemistry (Moles et al. 2011), in predator pressure (Björkman et al. 2011) or by variations in leaf toughness (Onoda et al. 2011).

Climatic variables are usually considered one of the main drivers of latitudinal patterns in the intensity of biotic interactions. Temperature directly affects the performance and abundance of invertebrate herbivores (Bale et al. 2002), since their physiology, population growth and dynamics are generally controlled by temperature (Hodkinson and Bird 1998). Temperature could also affect invertebrate herbivores indirectly, through changes in the palatability or availability of their host plants (Bale et al. 2002). Warmer temperatures have been associated with increased levels of herbivory in the fossil record (Wilf and Labandeira 1999; Wilf et al. 2001) and in experimental field studies (Richardson et al. 2002; Roy et al. 2004). Temperature was also found to explain latitudinal patterns in background herbivory in northern boreal forests (Kozlov 2008), as well as annual variations in this pattern (Kozlov et al. 2013), with higher temperatures associated with increased levels of herbivory. The effects of precipitation on the levels and types of invertebrate herbivory have been studied less systematically (Bale et al. 2002). Temperature was found to be a better predictor of herbivory than precipitation at a global scale (Kozlov et al. 2015a; Zhang et al. 2016), yet precipitation has a stronger influence on the global patterns of leaf traits related to resistance to damage, with more resistant leaves in sites with lower annual precipitation (Onoda et al. 2011). Broad gradients of temperature and precipitation exist across the tundra biome; we therefore hypothesize that patterns of invertebrate herbivory will also be influenced by climatic gradients in this region.

Patterns of herbivory and their underlying mechanisms may also differ between feeding guilds of herbivores with contrasting life history traits (Hiura and Nakamura 2013; Anstett et al. 2014). Some studies have found that geographical patterns in the abundance of different herbivores or the intensity of herbivory were driven by different climatic variables, suggesting that variation in the sensitivity of feeding guilds to climate could lead to disparate predictions under climate change (Leckey et al. 2014; Moreira et al. 2015). For example, externally feeding defoliators are more exposed to abiotic variables and may respond to them directly, whereas internally feeding herbivores like leaf miners and gallers may be affected by abiotic variables indirectly through their effects on leaf traits (e.g. Andrew and Hughes 2005, Sinclair and Hughes 2008). For instance, precipitation affects the intensity of herbivory by leaf miners and gall-makers (Leckey et al. 2014), possibly through its effects on leaf toughness (Onoda et al. 2011). Tougher leaves are well defended against external herbivores but may favour internally feeding herbivores as they provide safer shelter against pathogens and reduce levels of desiccation (Carneiro et al. 2005). We propose that the same distinction between external and internal feeders will drive differences in the patterns of invertebrate herbivory in tundra.

Temperatures and precipitation are predicted to continue increasing in the Arctic (Cook et al. 2014), and warming in tundra is expected to occur at a higher rate than the global average (IPCC 2013). The rapid pace of environmental changes in the Arctic underscores the urgency of studying the responses of fundamental ecological processes, such as herbivory, to varying climatic conditions. Insects living at higher latitudes are highly responsive to climate changes (Hodkinson and Bird 1998), and warming-induced increases in insect herbivory are expected to be stronger at higher latitudes (Wolf et al. 2008; Kozlov et al. 2015a). Experimental studies in tundra have shown

that the intensity of invertebrate herbivory increases with warming (Barrio et al. 2016; Birkemoe et al. 2016), but the lack of knowledge on current levels of background herbivory across the tundra biome constrains any meaningful predictions.

The objective of this study is to assess the intensity of background invertebrate herbivory and characterise its relationships with latitude and climatic variables in tundra. To achieve this goal, we measured leaf damage by tissue-feeding invertebrate herbivores on a common tundra plant with a circumpolar distribution, the dwarf birch (*Betula glandulosa-nana* complex), across a large number of sites spanning nearly 24° of latitude. Dwarf birch is a main food plant of many tundra herbivores (Koponen 1984; Bryant et al. 2014) and it is an important component in shrub tundra plant communities. The wide distribution of dwarf birch facilitates comparisons within a single host plant across a latitudinal gradient (Anstett et al. 2016); further, the range and abundance of dwarf birch are predicted to expand in response to warming (Euskirchen et al. 2009; Myers-Smith et al. 2011). We test the following hypotheses: (i) background invertebrate herbivory within the tundra biome is greatest at lower latitudes, consistent with the LHH, or where summer temperature and precipitation are highest; and (ii) the patterns of invertebrate herbivory by different feeding guilds will correspond with different climatic variables, given their sensitivity to different environmental cues. Specifically, we expect leaf damage by externally-feeding defoliators to be more strongly associated with summer temperature than damage by internally feeding herbivores (leaf miners and galls), and conversely that the latter will be more affected by climatic variables that determine leaf toughness, such as precipitation.

METHODS

Focal plant species and leaf sampling

Dwarf birch is a taxonomic complex with several closely related and hybridizing species. The main taxonomic units that we identified are *Betula glandulosa* Michx., *B. nana* subsp. *nana* L. and *B. nana* subsp. *exilis* (Sukaczew) Hultén. Species identification was conducted by collectors in the field and verified based on distribution maps (Figure 1; Bryant et al. 2014). *Betula glandulosa* is distributed throughout the northern regions of North America, from Alaska to Newfoundland, as well as the southern part of Greenland (Feilberg 1984). *Betula nana* is distributed throughout the Arctic regions of Eurasia and North America, with *B. nana* subsp. *nana* occurring from Greenland through northern Europe to Western Siberia, and *B. nana* subsp. *exilis* occurring from Eastern Siberia to Alaska and into northern Canada (Bryant et al. 2014). These three taxonomic units differ in leaf chemistry: *B. nana* subsp. *exilis* and *B. glandulosa* have higher concentrations of phenolic glycosides, condensed tannins, and triterpenes, have lower leaf nitrogen, and are less palatable to vertebrate herbivores than *B. nana* subsp. *nana* (Bryant et al.

2014; DeAngelis et al. 2015). However, the importance of these chemical differences for invertebrate herbivory is not known.

Plant material was collected in the summers of 2008-2013 (summarized by Kozlov et al. 2015b), 2014 and 2015 between June 12 and September 17. Although different protocols were followed in different years (**Table 1**), samples were collected in a way that allowed for spatial comparisons to evaluate background herbivory and its relationship to abiotic conditions. The protocol used in 2008-2013 (see Appendix S2 to Kozlov et al. 2015b) aimed at sampling plant foliage for measurements of insect herbivory at the global scale. The 2014 protocol was designed to assess defensive chemistry of dwarf birches across the Arctic (**Online Resource S1**); for the purpose of this study we only used one of the two top shoot samples collected in 2014 (both long and short shoots). We disregarded the samples specifically collected from short shoots (primary growth), as they may not be representative of herbivory on the whole plant. The 2015 protocol was aimed at measuring background invertebrate herbivory in tundra at the plant community level (http://herbivory.biology.ualberta.ca/files/2016/11/background_herbivory_tundra.pdf); here we report only the results for dwarf birch.

Briefly, all sampling protocols requested collection of birch leaves from several individuals per site (2-5 individuals were sampled in 2008-2013, 10 individuals in 2014, and 3 individuals in 2015). The samples included 71-500 leaves per site, where sites were defined as circular areas of approximately 10 m radius, at least 100 m apart. The number of sites within a location (i.e. spatially distinct 'study areas', at least 10 km apart) varied between 1 and 16 (for the 2008-2013 and 2014 protocols, a minimum of 1 site per location was requested, while for the 2015 protocol, the minimum number of sites per location was 5). Samples were collected from 192 sites in 56 locations (**Table 1**), spanning 23.7° of latitude across the tundra biome (including non-Arctic locations in alpine areas), from 55.2° N to 78.9° N (**Figure 1; Online Resource S2**). The geographical distribution of locations was not designed in advance: the requests for sampling were distributed across the research community, and all samples received were included in our study.

Leaf damage assessment

Each leaf was inspected on both sides for leaf damage with a light source against the leaf to detect damage on the surface, and then with the light through the leaf, to detect damage inside the leaf. We distinguished damage caused by three different feeding guilds of invertebrate herbivores: external leaf feeders (chewing or skeletonization) and internally feeding miners and gall makers (**Online Resource S3**). External feeding damage on dwarf birch is caused primarily by sawflies (Hymenoptera: Tenthredinidae) and by moth larvae (Lepidoptera) (Koponen 1984). Leaf miners can include larvae of Lepidoptera and Coleoptera (Viramo 1962), whereas galls are formed by insects and eriophyid mites. Other herbivores, such as phloem-feeders, also occur on dwarf birch (Koponen 1984), but damage imposed by these insects could not be measured from leaf samples.

Following a widely used methodology (Kozlov and Zvereva 2017, and references therein), each leaf was assigned to a damage class according to the each type of damage (multiple damage types on the same leaf occurred in less than 1% of leaves analysed) and to the visually estimated percentage of the leaf area damaged by invertebrates: intact leaves, 0.01-1, 1-5, 5-25, 25-50, 50-75 and 75-100% (Kozlov 2008). Samples from the three protocols were evaluated by three scorers (2008-2013 protocol by MVK, 2014 by EL, 2015 by ICB); assessments made by the three scorers on 6-10 training samples (100 leaves each) indicated no statistically significant effect of the scorer on the estimates of foliar damage (pairwise t-test: all $p > 0.1$; intra-class correlation coefficients ranged between 0.56 and 0.88).

Data analysis

The intensity of invertebrate herbivory at each site was calculated using three complementary measures: 1) percentage of leaves damaged: the percentage of leaves that showed signs of invertebrate herbivory at a site; 2) percentage of leaf area damaged: the percentage of leaf area consumed or otherwise damaged by invertebrate herbivores over the total number of leaves inspected in a sample; and 3) average damage per damaged leaf: the average leaf area consumed or otherwise damaged by invertebrate herbivores per damaged leaf. The percentage of leaves damaged indicates the distribution of damage within a site; the percentage of leaf area damaged gives an approximation of foliar loss per site as a measure of herbivory; and the average damage per damaged leaf reflects how much of the leaf area is affected, once a leaf is damaged (Kozlov 2008; Kozlov et al. 2015b). To calculate the latter two variables, the number of leaves in each damage class was multiplied by the corresponding median value of damage (i.e. 0 for intact leaves, 0.5% for the 0.01-1% class, 3% for the 1-5% class, 15% for the 5-25% class, 37.5% for the 25-50% class, 62.5% for the 50-75% class, and 87.5% for the 75-100% class) and summed for all damage classes. These values were divided by the total number of leaves to obtain an estimate of the percentage of total leaf area damaged, and by the number of damaged leaves to obtain the average damage per damaged leaf (Kozlov et al. 2015a; Kozlov et al. 2015b). All variables were calculated for all invertebrate herbivores and for the three different feeding guilds separately (**Online Resource S2**); given the low occurrence of mines and galls, only the percentage of leaves damaged was included in the models for these groups.

To investigate the effects of latitude and climatic variables on invertebrate herbivory in tundra we built Linear Mixed Effects Models for total herbivory and for each feeding group separately. In all models, sampling protocol (2008-2013, 2014 or 2015) was included as a random effect to account for potential confounding effects of year of sampling, person scoring leaf damage and/or protocol design. Nearly half of the sampling locations (25 out of 62) sampled one site only, so location could not be included in the models as a random factor; therefore, measurements of invertebrate herbivory for locations with more than one site were averaged across sites, and the number of sites sampled at each location was included as weights in the models to account for differences in sampling effort.

We extracted the following indices as potential predictors of background herbivory: mean July temperature in the year of sampling, annual temperature, temperature seasonality (standard deviation of annual temperature), maximum temperature of the warmest month, minimum temperature of the coldest month, mean temp of warmest quarter (Jun-Aug), total July precipitation in the year of sampling, annual precipitation and precipitation seasonality (coefficient of variation). Initial correlation analyses indicated that mean July temperature and total July precipitation in the year of sampling had the highest correlations with measures of herbivory (**Online Resource S4**), so these two climate variables and latitude were included as predictors in the models. These two climate variables were extracted from the Global Historical Climatology Network-Monthly (GHCN-M v3.2.1, Lawrimore et al. 2011). This dataset provides monthly mean temperatures and precipitation as a spatial raster (0.5 degree resolution) based on weather station data. We extracted July temperature and precipitation for the year of sample collection, because current-season weather is more relevant to invertebrate herbivory than multi-year averages (Kozlov et al. 2013). Climate variables and latitude were included as predictors in the models. Correlations between continuous predictor variables were low ($r < 0.4$; **Online Resource S4**) and Variance Inflation Factors indicated no strong multicollinearity ($VIF < 1.2$), so all three variables were included simultaneously in the models. Dwarf birch exhibits patterns of regional and taxonomic variation in defense against browsing by vertebrates (Bryant et al. 2014) that may also affect the observed patterns of invertebrate herbivory, so we included dwarf birch taxon (*B. glandulosa*, *B. nana* subsp. *exilis* and *B. nana* subsp. *nana*) as a fixed effect variable in our analyses. We also included collection date, measured as day-of-year, as a fixed covariate in the models because foliar damage accumulates over the growing season, but damaged leaves tend to abscise prematurely (Torp et al. 2010; Kozlov et al. 2016). For the two locations where no invertebrate leaf damage was detected (Svalbard and Beringa Island), we assigned the lowest possible value for herbivory (0.1% for the percentage of leaves damaged by all herbivores and defoliators, 0.01% in the case of leaf miners and gallers, and 0.01% for leaf area damaged). The percentage of damaged leaves and the percentage leaf area affected by herbivory were then log-transformed to comply with model assumptions of normality and homogeneity of variances, which were checked by visually inspecting model residuals (Zuur et al. 2009). All analyses were run in R 3.2.3 (R Development Core Team 2015) using the *lme4* package to build Linear Mixed Effects Models (Bates et al. 2015).

RESULTS

Feeding marks of invertebrate herbivores were found on 3,949 of the 30,817 leaves examined (12.8%). Damage by invertebrate herbivores affected between 0 and 46.0% of leaves at each location (mean \pm SE: $11.2 \pm 1.3\%$; median = 8.8%, $n=62$; **Online Resource S5**). On average, $1.4 \pm 0.2\%$ of leaf area was damaged at each location (median = 1.1%), and most damaged leaves were only slightly affected ($12.0 \pm 0.9\%$ of leaf area damaged; median = 11.8%). The vast majority of damage (98.6%) was caused by defoliators. Damage by internally feeding herbivores (leaf

miners and galls) was found on relatively few leaves: 31 were mined by larvae of several moth species and only 24 bore galls (see **Online Resource S6** for identification of mines and galls).

Total herbivory

Both the percentage of leaves with signs of invertebrate damage and the percentage of total leaf area damaged were positively associated with July temperature and precipitation (**Table 2a**), but there was no relationship with latitude. The models estimated linear increases of the log-transformed values of herbivory with increasing July temperature and precipitation (**Table 2a**), which implies smaller absolute increases in herbivory at locations with lower values of July temperature and precipitation, relative to locations with warmer and wetter summers (**Figure 2**). For instance, the model estimates indicated a 0.4% absolute increase in the percentage of leaves damaged for every degree C increase in mean July temperature for the coldest mean July temperatures measured in our study (4.7° C; **Figure 2a**). In contrast, the model estimates indicated a 1.7% absolute increase per degree C in the percentage of leaves damaged by invertebrate herbivores at locations with the highest mean July temperatures (16.7° C; **Figure 2a**). Relative to current levels of herbivory (11.2% leaves damaged), these estimated values of absolute increase at locations with July temperature values within the centre of the observed temperature range (absolute increase 0.93%), represent relative increases in the percentage of leaves damaged of 8.3%. Similarly, the percentage of total leaf area damaged was estimated to increase 0.04% per degree C in locations with colder summers and 0.17% per degree C in locations with warmer summers (**Figure 2b**). Relative to observed current levels of leaf area damaged (1.4%), these figures imply predicted increases of 6.7% in leaf area damaged by invertebrate herbivores per degree C increase in mean July temperature, at locations with July temperature values within the centre of the observed temperature range (mean July temperature observed across sites = 11.4° C).

The potential effects of increased precipitation followed similar trends, albeit a much weaker modelled effect than temperature. With a 10 mm increase in July precipitation, the percentage of leaves damaged by invertebrate herbivores increased by 0.3% in locations with the lowest observed precipitation (10.8 mm). In contrast, at locations with the highest observed mean July precipitation measured in our study (136.3 mm), the model estimated a 0.6% absolute increase in the percentage of leaves damaged (**Figure 2c**); the increase in the percentage of leaf area damaged ranged between 0.05% and 0.12% in locations with drier and wetter summers (**Figure 2d**). Relative to current levels of invertebrate herbivory, at sites with intermediate observed levels of July precipitation (mean total July precipitation observed across sites = 53.2 mm), the models predicted a 3.6% relative increase in the percentage of leaves damaged and 4.5% increase in percentage of leaf area damaged per 10 mm of increased precipitation.

None of the covariates (birch taxa or collection date) included in the models for total herbivory were associated with the percentage of leaves damaged at each location or with the percentage leaf area damaged (**Table 2a**). The

average damage per damaged leaf was not associated with latitude, temperature, precipitation, collection date or birch taxa (**Table 2a**).

Externally feeding defoliators

The distribution of damage by free-living defoliators within a site, as measured with the percentage of leaves damaged was associated with higher July temperature and precipitation (**Table 2b**), but no latitudinal pattern was apparent. When looking at foliar loss, the percentage of leaf area affected by defoliators was positively, albeit weakly, related to July precipitation and temperature (**Table 2b**). None of the covariates explained variation in on the percentage of leaves damaged by defoliators at each location or the percentage of leaf area damaged (**Table 2b**). On average, defoliators consumed $11.09 \pm 1.26\%$ of leaf area on damaged leaves (median = 8.56%), and this value was not associated with latitude, temperature, precipitation, collection date or birch taxa (**Table 2b**).

Internally feeding herbivores (leaf miners and galls)

The mean percentage of leaves damaged by leaf miners at each location was $0.06 \pm 0.02\%$ and, when present, leaf miners affected on average $11.77 \pm 3.05\%$ of leaf area. Galls were found on $0.08 \pm 0.05\%$ leaves per location, and affected $35.78 \pm 8.29\%$ of the leaf area of damaged leaves (excluding 2 galled leaves with petiole galls). The percentage of leaves damaged by leaf miners increased with July precipitation and collection date (**Table 3a**) and the percentage of leaves damaged by galls was associated with birch taxa, with *B. glandulosa* having a greater percentage of leaves damaged compared to *B. nana* subsp. *nana* (**Table 3b**).

DISCUSSION

Our study is the first to provide a quantitative analysis of background invertebrate herbivory across the tundra biome. Our analysis of leaf damage on a common tundra shrub in 56 locations across the circumpolar North showed that background invertebrate herbivory is nearly ubiquitous in tundra but occurs at low intensity. On average, invertebrate herbivores consumed 1.4% of leaf area of dwarf birch and affected 11.2% of leaves. The variation in background invertebrate herbivory in tundra showed no latitudinal pattern, but both foliar losses to external feeders and the percentage of leaves damaged by internal leaf-feeders were greater at sites with higher summer temperature and precipitation. Contrary to our expectations, all feeding guilds responded to the different climatic variables in a similar way.

General patterns of herbivory in our study were dominated by external feeding herbivores, which caused over 98% of the damage. Similar to previous studies in northern areas (Kozlov 2008; Kozlov et al. 2015b) and also in the fossil record (Wilf et al. 2001), the incidence of galls and leaf miners relative to that of defoliators was very small.

Although previous studies suggested that different groups of herbivores may respond differently to climate variables (Leckey et al. 2014; Moreira et al. 2015), our results do not support this view for the tundra; rather, our findings suggest a generalized increase in herbivory by all feeding guilds with increased temperature and precipitation. Such positive associations with temperature have already been described for these three groups of herbivores in northern Europe (chewers, Kozlov et al. 2015a; leaf miners, Kozlov et al. 2013; gallers, Kozlov et al. 2016). Similarly, studies in other ecosystems have also found increased herbivory with increased precipitation for leaf chewers (Kozlov 2008; Moreira et al. 2015), leaf miners (Leckey et al. 2014) and the occurrence of galls (Leckey et al. 2014).

Actual summer weather in the year of data collection was found to be a better predictor of herbivory than the latitude of the study site. This is not surprising, because the relationships between climate and latitude in the Arctic are weak. For example, the position of the 10°C July isotherm varies from 50°N near Aleutian Islands to 70°N in Scandinavia (CAVM Team 2003); accordingly, July temperature in our locations showed no correlation with latitude. Thus, although latitude appeared as a good predictor of invertebrate herbivory at the global scale according to the LHH (Pennings et al. 2009; Kozlov et al. 2015a), these trends do not hold within the tundra biome. Our study shows that biome-wide patterns of invertebrate herbivory are associated with proximal environmental cues (i.e. climatic variables) rather than with latitude, and warns against the use of latitudinal gradients as analogues for climate change in the Arctic (e.g. Hodkinson and Bird 1998), unless they do really represent a climatic gradient.

The average damage per damaged leaf (12.1%) is comparable to previous studies (6.9%, Kozlov et al. 2015a). These relatively low levels suggest that herbivores shift their feeding sites after even low levels of damage, possibly as a result of decreases in leaf palatability in response to damage or as a strategy to avoid detection by natural enemies (Fisher et al. 1999; Greyson-Gaito et al. 2016). Reductions in leaf palatability may be related to the production of secondary chemical compounds in response to herbivory (Nykänen and Koricheva 2004). In northern dwarf birch the production of secondary compounds both as induced or constitutive anti-herbivore defence shows local and regional variation (Graglia et al. 2001; Torp et al. 2010). For example, local topography can influence patterns of foliar concentrations of nitrogen and phenolic compounds in *B. nana* subsp. *nana* through its effects on snow accumulation and plant phenology (Torp et al. 2010). This spatial variation in defensive chemistry of birch potentially accounts for the relatively constant values of damage on damaged leaves over a range of climatic conditions. However, concentrations of plant secondary metabolites appear to be poor predictors of the extent of plant damage caused by insects under natural conditions (Carmona et al. 2011). The use of broad metrics, such as total phenolics, to measure resistance against herbivores or comparisons across unrelated species may obscure the importance of secondary compounds (Agrawal and Weber 2015; Anstett et al. 2015). This topic deserves further investigation, especially considering that climate change will also affect the defensive chemistry of plants. For example, phenolic compounds are expected to decrease with increased

temperature (Stark et al. 2015), but the response to warming might differ depending on what type and combination of secondary metabolites plants have (Graglia et al. 2001).

Our models predicted that changes in invertebrate herbivory in response to temperature and precipitation will differ along the range of climates sampled. It must be kept in mind that our approach represents a space-for-time substitution, where we infer changes in herbivory from locations with different climatic variables. Despite its limitations, this approach provides the best solution given the virtual lack of long-term trend data in patterns of invertebrate herbivory in tundra over time. Given that climate models project warming of 6-10 degree C over the next 100 years (IPCC 2013), the influence of temperatures on invertebrate background herbivory could be important. According to the logarithmic relationship indicated by our models, increases in invertebrate herbivory in locations with higher summer temperatures would be more pronounced than at locations with colder summers. The effect of precipitation followed similar trends but was not as pronounced and did not differ as much between the ends of the precipitation gradient. Precipitation is predicted to increase in the Arctic as a result of climate change (Cook et al. 2014), so these modest increases could, however, also be important. Differential climate sensitivities to temperatures and/or soil moisture have been also described for the phenology (Prev  y et al. 2017), community composition (Elmendorf et al. 2012) and growth (Myers-Smith et al. 2015) of tundra plants. For example, the growth of tundra shrubs was found to be more responsive to climate in wetter than in drier regions (Myers-Smith et al. 2015). Herbivory itself may also interact with climate to determine tundra plant performance. This has been observed for vertebrate herbivory in tundra (Speed et al. 2011; Speed et al. 2013) and suggested for invertebrate herbivores (Barrio et al. 2016). In addition, climate warming has been linked to increased growth (Bret-Harte et al. 2001) and decreased investment in defense of *B. nana* subsp. *nana* (Stark et al. 2015), so while invertebrate herbivory may be expected to increase in prevalence in a warmer climate, the net outcome of climate-herbivore-plant interactions is less certain.

The percentage of leaves damaged on dwarf birch by invertebrate herbivores in each location varied between 0 and 49% (median value was 8.7%). This has direct implications for the design of sampling protocols for detecting invertebrate damage. At least 33 leaves have to be collected to find a damaged one with a 95% probability, and 76 leaves to increase this probability to 99.9%. Consequently, for damage detection on dwarf birch we would recommend that at least 100 leaves per sampling site are collected. This was the sample size that we recommended in our collection protocols and we were able to detect invertebrate herbivory in 185 out of 187 sites. Larger sample sizes (~500 leaves) might be recommended to estimate leaf area losses to invertebrate herbivores with an adequate level of accuracy (Kozlov and Zvereva 2017) and to resolve the role of contrasting feeding guilds more robustly.

Our study supports the idea that background invertebrate herbivory could increase with current climatic changes in the tundra biome. Thus, there is an urgent need to understand how increases in the intensity of background

herbivory due to rising temperatures and precipitation will affect plant performance and ecosystem functioning in tundra. Measures of leaf damage alone may not directly reflect the cost of herbivory to the plant (Lim et al. 2015), as the impact of a given amount of herbivory depends on many other factors, including the cost of production of new leaves, resource availability and plant tolerance to herbivory (Kotaniemi and Rosenthal 2000). The extent to which increased background invertebrate herbivory may alter tundra communities will require a comprehensive analysis of foliar damage sustained by a wide variety of species and observations over longer periods, characterization of the invertebrate herbivore fauna, and manipulative field experiments (e.g. Barrio et al. 2016).

ACKNOWLEDGEMENTS

This study is a joint contribution of the Herbivory Network (<http://herbivory.biology.ualberta.ca>) and the Network for Arthropods of the Tundra (NeAT; <https://tundraarthropods.wordpress.com/>). Dwarf birch distribution maps were kindly provided by Kyle Joly. Sample collection during 2014 was facilitated by INTERACT (<http://www.eu-interact.org/>). ICB was supported by a postdoctoral fellowship funded by the Icelandic Research Fund (Rannsóknasjóður, grant nr 152468-051) and AXA Research Fund (15-AXA-PDOC-307); MtB and EK were supported by the Nordic Centre of Excellence TUNDRA, funded by the Norden Top-Level Research Initiative “Effect Studies and Adaptation to Climate Change”; EMS and KAB were supported by COAT (Climate-ecological Observatory of the Arctic Tundra); AB was supported by MOBILITY PLUS (1072/MOB/2013/0) and the Polish-American Fulbright Commission; CGB was supported by IUT 20-28, *EcolChange* Center of Excellence; BCF and TK were supported by the Academy of Finland (project 256991); MMPDH was supported by The Netherlands Organization for Scientific Research (NWO-ALW, VIDI grant 864.09.014); DSH was supported by the Natural Sciences and Engineering Research Council of Canada; AH was supported by the Research Council of Norway (grant 244557/E50); JL was funded by the German Research Foundation DFG (project WI 2680/8-1); MM-F was supported by a NERC IRF fellowship NE/L011859/1; SN was supported by the Villum foundation’s Young Investigator Programme (VKR023456); JS was supported by Kempefondstiftelsen and the Research Foundation Flanders (FWO); AS and NS were supported by the grant of RFBR (project 16-44-890108), grant of UB of RAS (project 15-15-4-35) and IEC “Arctic” of Yamal Government Department of Science and Innovation; LES and PAW were supported by the UK Natural Environment Research Council (NERC) grant NE/K000284/1; MVK and VZ were supported by the Academy of Finland (project 276671).

REFERENCES

- Agrawal A, Weber M (2015) On the study of plant defence and herbivory using comparative approaches: how important are secondary plant compounds. *Ecol Lett* 18:985–991
- Andrew NR, Hughes L (2005) Herbivore damage along a latitudinal gradient: relative impacts of different feeding guilds. *Oikos* 108:176–182
- Anstett D, Ahern J, Glinos J, et al (2015) Can genetically based clines in plant defence explain greater herbivory at higher

latitudes? *Ecol Lett* 18:1376–1386

Anstett DN, Naujokaitis-Lewis I, Johnson MTJ (2014) Latitudinal gradients in herbivory on *Oenothera biennis* vary according to herbivore guild and specialization. *Ecology* 95:2915–2923

Anstett DN, Nunes KA, Baskett C, Kotanen PM (2016) Sources of controversy surrounding latitudinal patterns in herbivory and defense. *Trends Ecol Evol* 31:789–802

Bale JS, Masters GJ, Hodkinson ID, et al (2002) Herbivory in global climate change research: direct effects of rising temperature on insect herbivores. *Glob Chang Biol* 8:1–16

Barrio IC, Bueno CG, Hik DS (2016) Warming the tundra: reciprocal responses of invertebrate herbivores and plants. *Oikos* 125:20–28

Bates D, Maechler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models using lme4. *J Stat Softw* 67:1–48

Batzli GO, White RG, MacLean SF, et al (1980) The herbivore-based ecosystem. In: Brown J, Miller PC, Tieszen LL, Bunnell FL (eds) *An arctic ecosystem: the coastal tundra at Barrow, Alaska*. pp 335–410

Birkemoe T, Bergmann S, Hasle TE, Klanderud K (2016) Experimental warming increases herbivory by leaf-chewing insects in an alpine plant community. *Ecol Evol* 6:6955–6962

Björkman C, Berggren A, Bylund H (2011) Causes behind insect folivory patterns in latitudinal gradients. *J Ecol* 99:367–369

Bret-Harte MS, Shaver GR, Zoerner JP, et al (2001) Developmental plasticity allows *Betula nana* to dominate tundra subjected to an altered environment. *Ecology* 82:18–32

Bryant JP, Joly K, Chapin FS, et al (2014) Can antibrowsing defense regulate the spread of woody vegetation in arctic tundra? *Ecography (Cop)* 37:204–211

Carmona D, Lajeunesse MJ, Johnson MT (2011) Plant traits that predict resistance to herbivores. *Funct Ecol* 25:358–367

Carneiro MAA, Fernandes GW, De Souza OFF (2005) Convergence in the variation of local and regional galling species richness. *Neotrop Entomol* 34:547–553

CAVM Team (2003) Circumpolar Arctic Region Bioclimate Subzones. Scale 1:7,500,000.

Coley P, Aide T (1991) Comparison of herbivory and plant defenses in temperate and tropical broad-leaved forests. In: Price P, Lewinsohn T, Fernandes G, Benson W (eds) *Plant-animal interaction: evolutionary ecology in tropical and temperate regions*. Wiley, New York, USA, pp 25–49

Cook BI, Smerdon JE, Seager R, Coats S (2014) Global warming and 21st century drying. *Clim Dyn* 43:2607–2627

Danks H V (1986) Insect plant interactions in arctic regions. *Rev d'Entomologie du Québec* 31:52–75

DeAngelis DL, Bryant JP, Liu R, et al (2015) A plant toxin mediated mechanism for the lag in snowshoe hare population recovery following cyclic declines. *Oikos* 124:796–805

Elmendorf SC, Henry GHR, Hollister RD, et al (2012) Global assessment of experimental climate warming on tundra vegetation: heterogeneity over space and time. *Ecol Lett* 15:164–175

Euskirchen ES, McGuire AD, Chapin FS, et al (2009) Changes in vegetation in Northern Alaska under scenarios of climate change, 2003–2100: implications for climate feedbacks. *Ecol Appl* 19:1022–1043

Feilberg J (1984) A phytogeographical study of South Greenland. Vascular plants. *Meddelelser om Grønland, Biosci vol* 15 70

Fisher AE., Hartley S., Young M (1999) Behavioural responses of the leaf-chewing guild to the presence of *Eriocrania* mines on silver birch (*Betula pendula*). *Ecol Entomol* 24:156–162

Graglia E, Julkunen-Tiitto R, Shaver GR, et al (2001) Environmental control and intersite variations of phenolics in *Betula nana* in tundra ecosystems. *New Phytol* 151:227–236

Greyson-Gaito CJ, Barbour MA, Rodriguez-Cabal MA, et al (2016) Freedom to move: Arctic caterpillar (Lepidoptera) growth rate increases with access to new willows (Salicaceae). *Can Entomol* 148:673–682

Haukioja E (1981) Invertebrate herbivory at tundra sites. In: Bliss LC, Heal OW, Moore JJ (eds) *Tundra ecosystems: a comparative analysis*. Cambridge University Press, New York, USA, pp 547–555

Hiura T, Nakamura M (2013) Different mechanisms explain feeding type-specific patterns of latitudinal variation in herbivore damage among diverse feeding types of herbivorous insects. *Basic Appl Ecol* 14:480–488

Hodkinson ID, Bird J (1998) Host-specific insect herbivores as sensors of climate change in Arctic and alpine environments. *Arct Alp Res* 30:78–83

IPCC [Intergovernmental Panel on Climate Change] (2013) *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York, US

Jepsen JU, Biuw M, Ims RA, et al (2013) Ecosystem impacts of a range expanding forest defoliator at the forest-tundra ecotone. *Ecosystems* 16:561–575

Jepsen JU, Hagen SB, Ims RA, Yoccoz NG (2008) Climate change and outbreaks of the geometrids *Operophtera brumata* and *Epirrita autumnata* in subarctic birch forest: evidence of a recent outbreak range expansion. *J Anim Ecol* 77:257–264

Johnson MTJ, Rasmann S (2011) The latitudinal herbivory- defence hypothesis takes a detour on the map. *New Phytol* 191:589–592

Kaukonen M, Ruotsalainen AL, Wali P, et al (2013) Moth herbivory enhances resource turnover in subarctic mountain birch forests? *Ecology* 94:267–272

- Koponen S (1984) Abundance of herbivorous insects on dwarf birch near the treeline in Alaska. Reports Kevo Subarctic Research Station 19:19–24
- Kotanan PM, Rosenthal JP (2000) Tolerating herbivory: does the plant care if the herbivore has a backbone? *Evol Ecol* 14:537–549
- Kozlov MVM, Skoracka A, Zverev V, et al (2016) Two birch species demonstrate opposite latitudinal patterns in infestation by gall-making mites in Northern Europe. *PLoS One* 11:e0166641
- Kozlov M V. (2008) Losses of birch foliage due to insect herbivory along geographical gradients in Europe: a climate-driven pattern? *Clim Change* 87:107–117
- Kozlov M V., Lanta V, Zverev V, Zvereva EL (2015a) Global patterns in background losses of woody plant foliage to insects. *Glob Ecol Biogeogr* 24:1126–1135
- Kozlov M V., van Nieuwerkerken EJ, Zverev V, Zvereva EL (2013) Abundance and diversity of birch-feeding leafminers along latitudinal gradients in northern Europe. *Ecography* 36:1138–1149
- Kozlov M V, Filippov BY, Zubrij NA, Zverev V (2015b) Abrupt changes in invertebrate herbivory on woody plants at the forest-tundra ecotone. *Polar Biol* 38:967–974
- Kozlov M V, Zvereva EL (2017) Background insect herbivory: impacts, patterns and methodology. In: Cánovas FM, Lüttge U, Matyssek R (eds) *Progress in Botany*. vol 79. doi: 10.1007/124_2017_4
- Lawrimore JH, Menne MJ, Gleason BE, et al (2011) An overview of the global historical climatology network monthly mean temperature data set, version 3. *J Geophys Res* 116:D19121
- Leckey EH, Smith DM, Nufio CR, Fornash KF (2014) Oak-insect herbivore interactions along a temperature and precipitation gradient. *Acta Oecologica* 61:1–8
- Lim JY, Fine PVA, Mittelbach GG (2015) Assessing the latitudinal gradient in herbivory. *Glob Ecol Biogeogr* 1–7
- Metcalf DB, Crutsinger GM, Kumordzi BB, Wardle DA (2016) Nutrient fluxes from insect herbivory increase during ecosystem retrogression in boreal forest. *Ecology* 97:124–132
- Moles AT, Bonser SP, Poore AGB, et al (2011) Assessing the evidence for latitudinal gradients in plant defence and herbivory. *Funct Ecol* 25:380–388
- Moreira X, Abdala-Roberts L, Parra-Tabla V, Mooney KA (2015) Latitudinal variation in herbivory: Influences of climatic drivers, herbivore identity and natural enemies. *Oikos* 124:1444–1452
- Myers-Smith IH, Elmendorf SC, Beck PSA, et al (2015) Climate sensitivity of shrub growth across the tundra biome. *Nat Clim Chang*. 5:887–891
- Myers-Smith IH, Forbes BC, Wilmking M, et al (2011) Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities. *Environ Res Lett* 6:45509
- Nykänen H, Koricheva J (2004) Damage-induced changes in woody plants and their effects on insect herbivore performance: a meta-analysis. *Oikos* 104:247–268
- Olofsson J, Beest M te, Ericson L (2013) Complex biotic interactions drive long-term vegetation dynamics in a subarctic ecosystem. *Philos Trans R Soc London B* 368:20120486
- Onoda Y, Westoby M, Adler PB, et al (2011) Global patterns of leaf mechanical properties. *Ecol Lett* 14:301–312
- Parker TC, Sadowsky J, Dunleavy H, et al (2017) Slowed biogeochemical cycling in sub-arctic birch forest linked to reduced mycorrhizal growth and community change after a defoliation event. *Ecosystems* 20: 316
- Pennings SC, Ho C, Salgado CS, et al (2009) Latitudinal variation in herbivore pressure in Atlantic coast salt marshes. *Ecology* 90:183–195
- Post E, Pedersen C (2008) Opposing plant community responses to warming with and without herbivores. *Proc Natl Acad Sci U S A* 105:12353–8
- Prevéy J, Vellend M, Rüger N, et al (2017) Greater temperature sensitivity of plant phenology at colder sites: implications for convergence across northern latitudes. *Glob Chang Biol*. doi: DOI: 10.1111/gcb.13619
- R Development Core Team (2015) R: A language and environment for statistical computing.
- Richardson SJ, Press MC, Parsons AN, Hartley SE (2002) How do nutrients and warming impact on plant communities and their insect herbivores? A 9-year study from a sub-Arctic heath. *J Ecol* 90:544–556
- Roy BA, Gusewell S, Harte J (2004) Response of plant pathogens and herbivores to a warming experiment. *Ecology* 85:2570–2581
- Sinclair RJ, Hughes L (2008) Incidence of leaf mining in different vegetation types across rainfall, canopy cover and latitudinal gradients. *Austral Ecol* 33:353–360
- Speed JD, Austrheim G, Hester AJ, Myrnerud A (2011) Browsing interacts with climate to determine tree-ring increment. *Funct Ecol* 25:1018–1023
- Speed JDM, Austrheim G, Hester AJ, Myrnerud A (2013) The response of alpine *Salix* shrubs to long-term browsing varies with elevation and herbivore density. *Arctic, Antarct Alp Res* 45:584–593
- Stark S, Väisänen M, Yläne H, et al (2015) Decreased phenolic defence in dwarf birch (*Betula nana*) after warming in subarctic tundra. *Polar Biol* 38:1993–2005
- Torp M, Witzell J, Baxter R, Olofsson J (2010) The effect of snow on plant chemistry and invertebrate herbivory: experimental

573 manipulations along a natural snow gradient. *Ecosystems* 13:741–751
 574 Viramo J (1962) Über die an der Zwergbirke (*Betula nana* L.) minierenden Insektenarten. *Ann Entomol Fenn* 28:118–126
 575 Wilf P, Labandeira CC (1999) Response of plant-insect associations to Paleocene-Eocene warming. *Science* 284:2153–2156
 576 Wilf P, Labandeira CC, Johnson KR, et al (2001) Insect herbivory, plant defense and early Cenozoic climate change. *Proc Natl*
 577 *Acad Sci* 98:6221–6226
 578 Wolf A, Kozlov M V, Callaghan T V (2008) Impact of non-outbreak insect damage on vegetation in northern Europe will be
 579 greater than expected during a changing climate. *Clim Chang* 87:91–106
 580 Zhang S, Zhang Y, Ma K, Shefferson R (2016) Latitudinal variation in herbivory: hemispheric asymmetries and the role of climatic
 581 drivers. *J Ecol* 104:1089–1095
 582 Zuur AF, Ieno EN, Walker NJ, et al (2009) *Mixed effects models and extensions in ecology with R*. Springer, New York
 583 Zvereva EL, Zverev V, Kozlov M V. (2012) Little strokes fell great oaks: minor but chronic herbivory substantially reduces birch
 584 growth. *Oikos* 121:2036–2043
 585

FIGURES

Figure 1. Sampling locations across the tundra biome. Size of points indicates number of sites per location, and colour indicates sampling protocol used: 2008-2013 (black), 2014 (grey) or 2015 (white). Distribution of dwarf birch taxa is indicated after Bryant et al. (2014).




Figure 2. Relationships between the intensity of total background invertebrate herbivory and July temperatures (a,b) and precipitation (c,d): predicted values (sizes of data points are proportional to the number of samples at each location), fitted lines and 95% confidence intervals. The intensity of herbivory was measured as the percentage of leaves damaged by invertebrates (a,c) and the percentage of leaf area affected out of all leaves examined (b,d). Dashed vertical line indicate mean observed July temperature and precipitation values across sites, and the box around them represents the range over which the absolute increases in herbivory were calculated (1°C in a and b, 10 mm in c and d).




Table 1. Summary of sampling protocols, indicating the aim of the protocol and where it can be found, the dwarf birch taxa targeted and the number of sampling locations and sites where samples were collected across locations. Note that some of the 56 study areas were sampled in different years and/or targeted different dwarf birch taxa, and are kept as separate 'sampling locations', so the number of locations presented in the table exceeds the number of study areas. For more details on each location see Online Resource S2.

Protocol	Aim of protocol and accessibility	Dwarf birch taxon	Number of locations (sites)
2008-2013	Measure insect herbivory at the global scale. Available as Online Resource S2 to Kozlov et al. 2015b.	<i>Betula nana exilis</i>	2(2)
		<i>Betula nana nana</i>	7(7)
2014	Determine the level of anti-browsing defence in dwarf birch across the Arctic. This protocol was not aimed at measuring invertebrate herbivory. Only one of the samples collected in 2014, consisting of top shoots (both long and short shoots) of <i>Betula</i> , was used in the present study. The protocol is available in Online Resource S1 (this study).	<i>Betula glandulosa</i>	8(18)
		<i>Betula nana exilis</i>	6(20)
		<i>Betula nana nana</i>	29(99)
2015	Assess variability of background invertebrate herbivory in tundra at the plant community level. Only the results for dwarf birch were used in the present study. The protocol is available at: http://herbivory.biology.ualberta.ca/files/2016/11/background_herbivory_tundra.pdf	<i>Betula nana exilis</i>	1(5)
		<i>Betula nana nana</i>	9(41)
Total			62(192)

Table 2. Factors explaining variation in different measures of foliar damage in dwarf birch (Linear Mixed Effect Model results) by all herbivores (a) and only defoliators (b), based on 62 samples from 56 locations across the tundra biome (some locations were sampled in different years or targeted different *Betula* taxa (see Table 1) and these locations are kept separate in the analyses). Baseline for species comparisons is *Betula nana* subsp. *nana*. The percentage of leaves affected and leaf area affected were log-transformed before analyses. Estimates in bold indicate that 95% confidence interval does not include zero. Sampling protocol was included as a random effect in the models, and sample sizes at each location were included as weights. Random effects are presented as standard deviations; n indicates the number of sampling protocols, and % refer to the percentage of residual variance assigned to sampling protocol.

a. All herbivores						
Explanatory variables	Percentage of leaves damaged		Percentage of leaf area damaged		Area damaged per damaged leaf	
<i>Fixed effects</i>	Estimate	95% CI	Estimate	95% CI	Estimate	95% CI
Intercept	-2.993	-5.864, -0.122	-2.460	-5.989, 1.069	40.419	-5.593, 86.432
Latitude	0.031	-0.002, 0.064	0.022	-0.018, 0.063	-0.246	-0.781, 0.288
Temperature	0.051	0.016, 0.086	0.053	0.010, 0.096	-0.050	-0.601, 0.501
Precipitation	0.005	0.002, 0.009	0.006	0.001, 0.011	0.022	-0.040, 0.085
Species – <i>B. glandulosa</i>	0.289	-0.067, 0.646	0.179	-0.259, 0.618	-2.629	-8.322, 3.063
Species – <i>B. n. exilis</i>	-0.254	-0.562, 0.054	-0.258	-0.637, 0.120	0.570	-4.362, 5.502
Collection date	0.004	-0.002, 0.010	0.000	-0.007, 0.007	-0.056	-0.148, 0.037
<i>Random effects</i>	SD		SD		SD	
Sampling protocol	0.239	(n=3, 13.1%)	0.279	(n=3, 12.0%)	2.409	(n=3, 5.6%)
Residual	0.615		0.756		9.889	
b. Defoliators						
Explanatory variables	Percentage of leaves damaged		Percentage of leaf area damaged		Area damaged per damaged leaf	
<i>Fixed effects</i>	Estimate	95% CI	Estimate	95% CI	Estimate	95% CI
Intercept	-2.847	-5.724, 0.029	-2.468	-6.008, 1.071	37.042	-9.318, 83.042
Latitude	0.030	-0.004, 0.063	0.023	-0.018, 0.064	-0.218	-0.756, 0.321
Temperature	0.050	0.015, 0.085	0.051	0.008, 0.094	-0.093	-0.644, 0.458
Precipitation	0.005	0.001, 0.009	0.006	0.001, 0.011	0.021	-0.041, 0.084
Species – <i>B. glandulosa</i>	0.271	-0.087, 0.628	0.169	-0.271, 0.608	-2.425	-8.148, 3.297
Species – <i>B. n. exilis</i>	-0.264	-0.573, 0.044	-0.295	-0.675, 0.084	-0.278	-5.245, 4.690
Collection date	0.004	-0.002, 0.010	0.000	-0.007, 0.007	-0.047	-0.140, 0.045
<i>Random effects</i>	SD		SD		SD	
Sampling protocol	0.233	(n=3, 12.5%)	0.266	(n=3, 10.9%)	2.124	(n=3, 4.3%)
Residual	0.616		0.759		9.971	

616

617

Table 3. Factors explaining variation in the percentage of leaves damaged by leaf miners (a) and gall makers (b) on dwarf birch (Linear Mixed Effect Model results), based on 62 samples from 56 locations across the tundra biome. Baseline for species comparisons is *Betula nana* subsp. *nana*. The percentage of leaves affected was log-transformed before analyses. Estimates in bold indicate that 95% confidence interval does not include zero. Sampling protocol was included as a random effect in the models, and sample sizes at each location were included as weights. Random effects are presented as standard deviations; n indicates the number of sampling protocols, and % refer to the percentage of residual variance assigned to sampling protocol.

Explanatory variables	a. Leaf miners		b. Gall makers	
<i>Fixed effects</i>	Estimate	95% CI	Estimate	95% CI
Intercept	-7.465	-12.596, -2.334	-1.587	-6.252, 3.078
Latitude	0.035	-0.024, 0.094	0.008	-0.046, 0.062
Temperature	0.034	-0.021, 0.089	0.013	-0.040, 0.066
Precipitation	0.009	0.002, 0.016	0.005	-0.001, 0.012
Species – <i>B. glandulosa</i>	-0.167	-0.782, 0.448	0.702	0.134, 1.270
Species – <i>B. n. exilis</i>	-0.108	-0.656, 0.439	0.357	-0.142, 0.856
Collection date	0.014	0.004, 0.023	-0.006	-0.015, 0.003
<i>Random effects</i>	SD		SD	
Sampling protocol	0.000	(n=3, 0%)	0.112	(n=3, 1.2%)
Residual	1.107		1.005	

ONLINE RESOURCES

Online Resource S1. Instructions sent to participants to collect dwarf birch samples in the summer 2014

Circumpolar variation in antibrowsing defense in *Betula nana* and *B. glandulosa*

Mariska te Beest¹, Dagmar Egelkraut¹, Maja Sundqvist¹, Judith Sitters¹, Johan Olofsson¹, et al.

¹Department of Ecology and Environmental Science, Umeå University, SE-901 87 Umeå, Sweden

Aim. Determine the level of antibrowsing defense in dwarf birches at a global scale

Methods. We will collect samples from *B. nana* and *B. glandulosa* across the Arctic regions of the world and assess the samples for total N and C, total phenols, condensed tannins and triterpenes.

Protocol. For each site (area of ~ 10m radius), collect top shoots (branch with leaves) from 10 individuals in one paper bag and additional leaves from the same 10 individuals in a second paper bag.

For each *Betula* individual collect:

- Height (in cm)
- 10 top shoots (the top 10 cm of the long shoots (Fig. 1) – keep the leaves attached to the branch)
- An additional sample of appr. 50 leaves from the leaves of short shoots (clusters of 4 leaves directly attached to the main stem, see fig. 1).

Individual samples should be combined per site, resulting in 2 bags per site – a top shoot sample and a leaf sample.

	N per individual	N per site (final bags)
Top shoots	10	100
Leaves	~50	~500

Provide a rough description of the site, including:

- GPS location and altitude
- Collection date
- Dominant species/vegetation type
- Whether the site is dry or wet
- If possible, take a picture
- Report presence and signs of herbivory at the site (browsing, faeces, observations etc.)
- List the major herbivores that you think are important at the site.

Allow the samples to dry at air temperature. If possible collect in 2 (or more) contrasting habitats at each location, for example wet vs. dry tundra or in different study areas, if applicable.




Figure 1. *Betula* makes 2 kinds of shoots, long shoots and short shoots. Long shoots elongate during the season, with younger and smaller leaves towards the top. Short shoots do not elongate and are made up of a cluster of usually 4 leaves that are all of the same age. It is the top 10 cm of the long shoots we are interested for the top shoot samples, and the short shoot leaves that we are interested in for the additional leaf sample (Source: www.biolib.de).

Online Resource S2. List of locations (ordered by taxon, from higher to lower latitude), indicating taxon of dwarf birch sampled and year of sampling (corresponding to the three sampling protocols: 2008-2013, 2014 and 2015). Location characteristics (**Table S2.1**) and herbivory measurements (**Table S2.2**) are shown, first three columns are common to both tables. Separate rows are provided for locations sampling more than one birch taxa or sampled in different years.

Table S2.1. Characteristics of locations including collection date (day-of-year) and number of sites and leaves sampled per location, geographical position (latitude and longitude in degrees) and climatic variables (July temperature and precipitation for the year of sampling).

Location	Taxon	Sampling year	Collection date	Number of sites / leaves	Latitude (°N)	Longitude (°E)	July temperature (°C)	July precipitation (mm)
Herschel Island	<i>B. glandulosa</i>	2014	209	1/100	69.60	-139.90	8.7	19.8
Trail Valley Cr	<i>B. glandulosa</i>	2014	208	2/200	68.74	-133.50	12.7	50.9
Fairbanks	<i>B. glandulosa</i>	2014	192	2/195	64.90	-147.95	16.0	136.3
Deception Bay	<i>B. glandulosa</i>	2014	191	2/200	62.10	-74.31	8.5	43.1
Pika Camp	<i>B. glandulosa</i>	2014	205	5/500	61.21	-138.28	9.6	91.2
Narsarsuaq	<i>B. glandulosa</i>	2014	234	1/100	61.16	-45.38	7.9	57.6
Kluane	<i>B. glandulosa</i>	2014	191	2/200	61.00	-138.40	8.3	68.9
Churchill	<i>B. glandulosa</i>	2014	208	3/300	58.74	-93.83	13.5	67.7
Lena Delta	<i>B. n. exilis</i>	2014	222	3/300	72.37	126.34	10.4	59.1
Tit Ary	<i>B. n. exilis</i>	2014	221	1/100	71.97	127.10	10.9	57.5
Kytalyk	<i>B. n. exilis</i>	2014	204	3/300	70.83	147.48	10.5	68.0
Toolik B	<i>B. n. exilis</i>	2014	184	3/300	68.96	-150.25	7.9	109.4
Toolik A	<i>B. n. exilis</i>	2014	178	5/1679	68.64	-149.57	9.8	36.9
Toolik A	<i>B. n. exilis</i>	2015	212	9/871	68.64	-149.57	7.3	106.1
Fairbanks	<i>B. n. exilis</i>	2014	158	1/100	64.87	-147.86	16.1	126.1
Beringowskij	<i>B. n. exilis</i>	2010	221	1/263	62.70	178.90	10.6	29.0
Beringa Island	<i>B. n. exilis</i>	2009	259	1/92	55.20	166.00	8.6	58.0
Svalbard	<i>B. n. nana</i>	2009	210	1/94	78.90	16.00	5.8	15.0
Zackenberga	<i>B. n. nana</i>	2014	218	4/400	74.47	-20.58	5.3	40.7
Nordkapp	<i>B. n. nana</i>	2009	223	1/180	71.20	25.80	9.8	66.0
Seiland	<i>B. n. nana</i>	2014	190	2/200	70.51	23.53	14.7	12.4
Varanger	<i>B. n. nana</i>	2014	195	16/1600	70.44	27.46	14.5	15.4
Ifjordfjellet	<i>B. n. nana</i>	2014	195	3/300	70.29	29.00	14.6	16.8
Mordy-Yakha	<i>B. n. nana</i>	2014	199	1/100	70.10	68.35	4.7	42.4
Njallavaara	<i>B. n. nana</i>	2015	231	5/1490	70.05	27.60	10.2	26.9
Kevo A	<i>B. n. nana</i>	2014	191	1/100	70.02	27.93	15.2	23.9
Ailigas	<i>B. n. nana</i>	2015	224	5/1619	69.89	27.07	10.2	26.9
Ailigas	<i>B. n. nana</i>	2014	192	1/100	69.82	27.01	15.2	23.9
Kevo B	<i>B. n. nana</i>	2009	221	1/242	69.80	27.00	12.5	33.0
Joatka	<i>B. n. nana</i>	2014	195	6/600	69.75	24.00	14.5	45.7
Reisa	<i>B. n. nana</i>	2014	228	5/500	69.53	29.32	16.0	23.2
Kilpisjärvi	<i>B. n. nana</i>	2014	202	2/200	69.04	20.86	14.0	61.1
Murmansk A	<i>B. n. nana</i>	2013	229	1/500	68.90	33.70	12.6	39.0
Murmansk A	<i>B. n. nana</i>	2015	222	3/1047	68.87	34.54	10.0	42.0
Tobseda	<i>B. n. nana</i>	2014	204	3/300	68.59	52.32	7.9	20.5
Ostojeaggi	<i>B. n. nana</i>	2014	219	1/100	68.48	19.74	14.5	46.5
Kärkevagge	<i>B. n. nana</i>	2014	239	1/100	68.41	18.32	12.0	39.5
Abisko	<i>B. n. nana</i>	2014	237	1/100	68.34	18.82	12.9	39.5
Tavva	<i>B. n. nana</i>	2014	208	1/100	68.34	21.18	14.7	97.8
Paddus	<i>B. n. nana</i>	2014	163	5/500	68.32	18.86	12.9	39.5

Erkuta	<i>B. n. nana</i>	2014	215	1/100	68.23	69.15	9.0	49.8
Erkuta	<i>B. n. nana</i>	2015	216	5/1801	68.23	69.15	10.7	62.7
Latnjajaure	<i>B. n. nana</i>	2015	216	1/369	68.21	18.29	6.9	79.8
Lovozero	<i>B. n. nana</i>	2013	216	1/300	67.90	34.40	13.6	40.0
Ritsem	<i>B. n. nana</i>	2014	229	6/600	67.77	17.54	11.5	60.9
Laborovaya	<i>B. n. nana</i>	2009	226	1/248	67.70	67.60	14.3	36.0
Severnaya	<i>B. n. nana</i>	2015	224	5/1509	67.65	54.05	8.9	46.0
Naryan-Mar	<i>B. n. nana</i>	2013	216	1/200	67.60	53.10	13.3	36.0
Igarka	<i>B. n. nana</i>	2014	221	1/100	67.50	86.54	15.4	46.0
Padjelanta	<i>B. n. nana</i>	2015	213	5/1633	67.31	16.69	7.5	90.1
Padjelanta	<i>B. n. nana</i>	2014	181	3/300	67.27	16.66	13.0	47.7
Kangerlussuaq	<i>B. n. nana</i>	2014	201	2/200	67.13	-50.15	9.9	10.8
Nanoq	<i>B. n. nana</i>	2014	181	3/300	67.10	50.29	10.3	48.9
Labytnangi	<i>B. n. nana</i>	2014	200	1/100	66.85	66.30	10.6	91.5
Ammarnäs	<i>B. n. nana</i>	2014	220	4/400	66.02	16.02	15.5	85.6
Usinsk	<i>B. n. nana</i>	2014	216	1/100	66.00	57.67	11.0	28.7
Theistareykir	<i>B. n. nana</i>	2015	213	5/1566	65.90	-17.08	7.9	45.8
Audkuluheidi	<i>B. n. nana</i>	2015	215	5/1551	65.13	-19.67	7.4	26.3
Dovre	<i>B. n. nana</i>	2014	230	2/200	62.30	9.62	11.9	80.4
Sångfjället	<i>B. n. nana</i>	2014	195	9/900	62.29	13.49	15.5	90.2
Långfjället	<i>B. n. nana</i>	2014	208	11/1100	62.10	12.26	16.7	83.8
Setesdal	<i>B. n. nana</i>	2014	192	2/200	59.05	7.16	16.0	83.5


Table S2.2 Measurements of herbivory at each location (percentage of leaves damaged, percentage of leaf area affected and average damage) for all herbivores and different feeding guilds separately (defoliators, miners and galls). When more than one site was sampled at each location, means and SE are provided.

			all herbivores			defoliators			miners		galls	
Location	Taxon	Sampling year	percentage leaves	percentage leaf area	average damage	percentage leaves	percentage leaf area	average damage	percentage leaves	percentage leaf area	percentage leaves	percentage leaf area
Herschel Island	<i>B. glandulosa</i>	2014	2.00	0.04	1.75	2.00	0.04	1.75	0.00	0.00	0.00	0.00
Trail Valley Cr	<i>B. glandulosa</i>	2014	12.50±2.50	0.68±0.19	5.37±0.47	9.50±0.50	0.65±0.16	7.01±2.11	0.00±0.00	0.00±0.00	3.00±3.00	0.03±0.03
Fairbanks	<i>B. glandulosa</i>	2014	10.34±3.34	1.09±0.05	11.91±4.37	10.34±3.34	1.09±0.05	11.91±4.37	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Deception Bay	<i>B. glandulosa</i>	2014	1.50±0.50	0.34±0.19	20.62±5.62	1.50±0.50	0.34±0.19	20.62±5.62	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Pika Camp	<i>B. glandulosa</i>	2014	19.40±5.28	1.84±0.67	8.73±1.06	18.80±5.30	1.84±0.67	9.10±1.06	0.20±0.20	0.00±0.00	0.40±0.40	0.00±0.00
Narsarsuaq	<i>B. glandulosa</i>	2014	31.00	5.61	18.08	31.00	5.61	18.08	0.00	0.00	0.00	0.00
Kluane	<i>B. glandulosa</i>	2014	28.50±5.50	3.56±0.18	12.85±1.85	28.50±5.50	3.56±0.18	12.85±1.85	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Churchill	<i>B. glandulosa</i>	2014	16.33±2.33	1.20±0.24	7.22±0.71	16.00±2.08	0.99±0.23	6.24±1.22	0.00±0.00	0.00±0.00	0.33±0.33	0.21±0.21
Kytalyk	<i>B. n. exilis</i>	2014	4.00±1.53	0.11±0.07	2.38±0.96	4.00±1.53	0.11±0.07	2.38±0.96	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Lena Delta	<i>B. n. exilis</i>	2014	3.00	0.68	22.50	3.00	0.68	22.50	0.00	0.00	0.00	0.00
Tit Ary	<i>B. n. exilis</i>	2014	9.33±2.33	1.99±0.64	20.76±3.14	9.33±2.33	1.99±0.64	20.76±3.14	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Toolik B	<i>B. n. exilis</i>	2014	4.67±1.33	0.76±0.60	13.25±9.67	4.67±1.33	0.76±0.60	13.25±9.67	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Toolik A	<i>B. n. exilis</i>	2014	6.25±0.98	0.65±0.13	11.69±2.31	5.76±0.99	0.55±0.14	10.33±2.20	0.11±0.11	0.00±0.00	0.38±0.26	0.10±0.07
Toolik A	<i>B. n. exilis</i>	2015	3.69±0.61	0.63±0.18	17.42±5.05	3.45±0.69	0.52±0.17	15.26±4.84	0.12±0.08	0.03±0.02	0.12±0.12	0.08±0.08
Fairbanks	<i>B. n. exilis</i>	2014	3.00	0.45	15.00	3.00	0.45	15.00	0.00	0.00	0.00	0.00
Beringowskij	<i>B. n. exilis</i>	2010	14.45	2.90	20.05	14.45	2.90	20.05	0.00	0.00	0.00	0.00
Beringa Island	<i>B. n. exilis</i>	2009	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Svalbard	<i>B. n. nana</i>	2009	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Zackenbergl	<i>B. n. nana</i>	2014	3.75±1.49	0.32±0.16	5.82±2.23	3.50±1.32	0.32±0.16	6.24±2.55	0.25±0.25	0.00±0.00	0.00±0.00	0.00±0.00
Nordkapp	<i>B. n. nana</i>	2009	6.67	0.66	9.88	6.67	0.66	9.88	0.00	0.00	0.00	0.00
Seiland	<i>B. n. nana</i>	2014	13.50±1.50	3.51±1.31	25.21±6.92	13.50±1.50	3.51±1.31	25.21±6.92	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Varanger	<i>B. n. nana</i>	2014	4.75±1.68	0.50±0.22	6.10±1.72	4.75±1.68	0.50±0.22	6.10±1.72	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Ifjordfjellet	<i>B. n. nana</i>	2014	8.67±3.18	1.06±0.57	9.55±3.78	8.67±3.18	1.06±0.57	9.55±3.78	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Mordy-Yakha	<i>B. n. nana</i>	2014	3.00	0.56	18.50	3.00	0.56	18.50	0.00	0.00	0.00	0.00
Njallavaara	<i>B. n. nana</i>	2015	17.38±2.67	2.62±0.57	14.64±1.43	17.11±2.71	2.58±0.57	14.72±1.52	0.27±0.13	0.04±0.04	0.00±0.00	0.00±0.00
Kevo A	<i>B. n. nana</i>	2014	8.00	2.63	32.94	8.00	2.63	32.94	0.00	0.00	0.00	0.00
Ailigas	<i>B. n. nana</i>	2015	21.34±3.59	1.29±0.15	6.70±1.22	21.02±3.59	1.28±0.15	6.78±1.27	0.31±0.02	0.01±0.01	0.00±0.00	0.00±0.00
Ailigas	<i>B. n. nana</i>	2014	3.00	0.21	7.00	3.00	0.21	7.00	0.00	0.00	0.00	0.00
Kevo B	<i>B. n. nana</i>	2009	3.72	0.15	4.00	3.72	0.15	4.00	0.00	0.00	0.00	0.00
Joatka	<i>B. n. nana</i>	2014	21.33±4.57	3.40±0.91	15.14±1.39	21.00±4.65	3.33±0.94	14.86±1.75	0.17±0.17	0.00±0.00	0.17±0.17	0.06±0.06
Reisa	<i>B. n. nana</i>	2014	10.40±1.75	0.88±0.24	7.76±1.61	9.40±1.44	0.76±0.22	7.32±1.64	1.00±0.45	0.13±0.06	0.00±0.00	0.00±0.00


ELECTRONIC SUPPLEMENTARY MATERIALS – Barrio et al. Background invertebrate herbivory on dwarf birch (*Betula glandulosa-nana* complex) increases with temperature and precipitation across the tundra biome

Kilpisjärvi	<i>B. n. nana</i>	2014	39.50±9.50	5.06±0.60	13.21±1.64	39.50±9.5	5.06±0.60	13.21±1.64	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Murmansk A	<i>B. n. nana</i>	2013	2.60	0.84	32.31	2.60	0.84	32.31	0.00	0.00	0.00	0.00
Murmansk A	<i>B. n. nana</i>	2015	46.00±9.93	7.87±3.30	14.88±2.88	46.00±9.93	7.87±3.30	14.88±2.88	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Tobseda	<i>B. n. nana</i>	2014	4.00±0.58	0.30±0.05	7.56±0.42	4.00±0.58	0.30±0.05	7.56±0.42	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Ostojeaggi	<i>B. n. nana</i>	2014	17.00	1.82	10.74	17.00	1.82	10.74	0.00	0.00	0.00	0.00
Kärkevagge	<i>B. n. nana</i>	2014	10.00	1.55	15.55	10.00	1.55	15.55	0.00	0.00	0.00	0.00
Abisko	<i>B. n. nana</i>	2014	6.00	0.10	1.75	6.00	0.10	1.75	0.00	0.00	0.00	0.00
Tavva	<i>B. n. nana</i>	2014	23.00	1.59	6.91	23.00	1.59	6.91	0.00	0.00	0.00	0.00
Paddus	<i>B. n. nana</i>	2014	4.80±1.24	0.87±0.36	13.75±5.63	4.80±1.24	0.87±0.36	13.75±5.63	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Erkuta	<i>B. n. nana</i>	2014	16.00	1.60	10.03	16.00	1.60	10.03	0.00	0.00	0.00	0.00
Erkuta	<i>B. n. nana</i>	2015	15.04±3.49	1.86±0.36	12.98±1.54	14.54±3.62	1.68±0.42	11.69±2.06	0.11±0.07	0.04±0.03	0.4±0.25	0.13±0.10
Latnjajaure	<i>B. n. nana</i>	2015	5.96	1.05	17.64	5.96	1.05	17.64	0.00	0.00	0.00	0.00
Lovozero	<i>B. n. nana</i>	2013	11.33	2.81	24.79	11.33	2.81	24.79	0.00	0.00	0.00	0.00
Ritsem	<i>B. n. nana</i>	2014	12.17±4.64	1.59±0.45	15.81±2.06	11.83±4.46	1.54±0.42	15.79±2.07	0.33±0.21	0.05±0.03	0.00±0.00	0.00±0.00
Laborovaya	<i>B. n. nana</i>	2009	33.06	2.03	6.13	33.06	2.03	6.13	0.00	0.00	0.00	0.00
Severnaya	<i>B. n. nana</i>	2015	16.00±3.57	1.05±0.23	7.53±1.49	15.81±3.59	1.02±0.23	7.62±1.73	0.20±0.13	0.03±0.03	0.00±0.00	0.00±0.00
Naryan-Mar	<i>B. n. nana</i>	2013	3.50	0.20	5.71	3.50	0.20	5.71	0.00	0.00	0.00	0.00
Igarka	<i>B. n. nana</i>	2014	34.00	2.42	7.10	34.00	2.42	7.10	0.00	0.00	0.00	0.00
Padjelanta	<i>B. n. nana</i>	2015	14.30±3.78	2.66±0.76	18.42±1.93	14.30±3.78	2.66±0.76	18.42±1.93	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Padjelanta	<i>B. n. nana</i>	2014	5.33±0.88	0.33±0.05	6.17±0.32	5.33±0.88	0.33±0.05	6.17±0.32	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Kangerlussuaq	<i>B. n. nana</i>	2014	5.00±1.00	0.12±0.08	2.23±1.10	5.00±1.00	0.12±0.08	2.23±1.10	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Nanoq	<i>B. n. nana</i>	2014	5.67±1.45	1.29±0.96	18.19±10.94	5.67±1.45	1.29±0.96	18.19±10.94	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Labytnangi	<i>B. n. nana</i>	2014	4.00	0.19	4.75	4.00	0.19	4.75	0.00	0.00	0.00	0.00
Ammarnäs	<i>B. n. nana</i>	2014	11.50±3.01	1.19±0.34	10.95±2.94	11.25±3.04	1.18±0.34	11.24±3.06	0.25±0.25	0.01±0.01	0.00±0.00	0.00±0.00
Usinsk	<i>B. n. nana</i>	2014	13.00	0.28	2.19	13.00	0.28	2.19	0.00	0.00	0.00	0.00
Theistareykir	<i>B. n. nana</i>	2015	0.96±0.36	0.12±0.04	21.69±11.45	0.96±0.36	0.12±0.04	21.69±11.45	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Audkuluheidi	<i>B. n. nana</i>	2015	3.90±1.21	0.73±0.24	16.24±4.74	3.90±1.21	0.73±0.24	16.24±4.74	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Dovre	<i>B. n. nana</i>	2014	3.00±0.00	0.21±0.00	7.00±0.00	3.00±0.00	0.21±0.00	7.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Sångfjället	<i>B. n. nana</i>	2014	8.89±2.35	1.12±0.35	11.93±2.61	8.44±2.35	0.98±0.35	10.68±2.39	0.22±0.15	0.00±0.00	0.22±0.22	0.14±0.14
Långfjället	<i>B. n. nana</i>	2014	13.91±2.58	1.78±0.35	13.98±2.15	13.82±2.59	1.78±0.35	14.06±2.15	0.09±0.09	0.00±0.00	0.00±0.00	0.00±0.00
Setesdal	<i>B. n. nana</i>	2014	11.5±0.50	1.76±1.40	14.76±11.58	11.5±0.50	1.76±1.40	14.76±11.58	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00


Online Resource S3. Leaf damage on dwarf birch (*Betula glandulosa-nana* complex) leaves caused by invertebrate herbivores: external leaf feeders (chewing (A, E, I) or skeletonized (B, F, J), leaf miners (C, G, K) or galls (D, H, L).



Online Resource S4. Correlations between one of our measures of herbivory, the proportion of leaves damaged at each location (“herbiv”, log-transformed, as in our analyses), latitude (°N; “lat”) and climatic variables, including mean July temperature in the year of sampling (°C; “temp”) and total July precipitation in the year of sampling (mm; “prec”), together with climatic variables obtained from BioCLIM: annual temperature (“Tannual”; BIO1), temperature seasonality (“Tseason”; BIO4), maximum temperature of the warmest month (“Tmax”; BIO5), minimum temperature of the coldest month (“Tmin”; BIO6), mean temperature of the warmest quarter (“Tmean wQ”; BIO10), annual precipitation (“Pannual”; BIO12) and precipitation seasonality (“Pseason”; BIO15). The upper diagonal shows Pearson correlation coefficients with size proportional to its magnitude; the lower diagonal shows pairwise plots indicating the relationships between pairs of variables, with a LOESS line to aid visual interpretation.



Appendix S5. Distribution of the percentage of leaves affected by invertebrate herbivory for all sampling locations, for all invertebrate herbivores (a) and each feeding guild separately: defoliators (b), leaf miners (c) and gallers (d). Vertical lines indicate median values



Appendix S6. Identification of leaf mines and galls on different dwarf birch taxa. Leaf miner taxa were identified by Erik van Nieuwerkerken, based on mine morphology and larval remains when present. Eriophyid galling mites were identified by Anna Skoracka.

Location	Type of damage	Latitude (°N)	Longitude (°E)	Dwarf birch taxon	Collection date	Number of leaves	Identification
Churchill	gall	58.7506	-93.8596	<i>B. glandulosa</i>	2014-07-28	1	unidentified insect larva
Kluane	mine	61.2016	-138.2595	<i>B. glandulosa</i>	2014-07-25	1	<i>Stigmella</i> sp.
Kluane	gall	68.7414	-133.4973	<i>B. glandulosa</i>	2014-07-25	2	unidentified insect larva
Trail Valley Creek	gall	68.7414	-133.4973	<i>B. glandulosa</i>	2014-07-28	6	eriophyoid mite <i>Eriophyes leionotus</i> (Nalepa)
Toolik A	mine	68.6434	-149.5706	<i>B. n. exilis</i>	2015-08-01	1	<i>Coleophora</i> sp.
Toolik A	mine	68.6426	-149.5753	<i>B. n. exilis</i>	2015-08-01	1	<i>Stigmella</i> sp.
Toolik A	gall	68.6434	-149.5706	<i>B. n. exilis</i>	2015-08-01	2	unidentified insect larva
Toolik A	gall	68.6426	-149.5753	<i>B. n. exilis</i>	2014-06-26	2	unidentified insect larva
Ailigas	mine	69.8953	27.0665	<i>B. n. nana</i>	2015-08-13	1	<i>Stigmella tristis</i>
Ailigas	mine	69.8949	27.0640	<i>B. n. nana</i>	2015-08-13	1	<i>Stigmella tristis</i>
Ailigas	mine	69.8953	27.0665	<i>B. n. nana</i>	2015-08-13	1	<i>Stigmella tristis</i>
Ailigas	mine	69.8943	27.0716	<i>B. n. nana</i>	2015-08-13	1	<i>Stigmella tristis</i>
Ailigas	mine	69.8935	27.0695	<i>B. n. nana</i>	2015-08-13	1	<i>Stigmella tristis</i>
Ammarnäs	mine	66.0187	16.0195	<i>B. n. nana</i>	2014-08-08	1	unknown
Erkuta	mine	68.2267	69.1509	<i>B. n. nana</i>	2015-08-05	1	<i>Phyllonorycter</i> sp.
Erkuta	mine	68.2267	69.1509	<i>B. n. nana</i>	2015-08-05	1	<i>Coleophora</i> sp.
Erkuta	gall	68.2267	69.1509	<i>B. n. nana</i>	2015-08-05	5	unidentified insect larva
Erkuta	gall	68.2267	69.1509	<i>B. n. nana</i>	2015-08-05	1	unidentified insect larva
Erkuta	gall	68.2267	69.1509	<i>B. n. nana</i>	2015-08-05	1	unidentified insect larva
Erkuta	gall	68.2267	69.1509	<i>B. n. nana</i>	2015-08-05	3	unidentified insect larva
Joatka	gall	69.75306	24.0028	<i>B. n. nana</i>	2014-07-15	1	unidentified insect larva
Njallavaara	mine	70.0462	27.6040	<i>B. n. nana</i>	2015-08-20	1	<i>Stigmella tristis</i>
Njallavaara	mine	70.0467	27.6015	<i>B. n. nana</i>	2015-08-20	2	<i>Stigmella tristis</i>
Njallavaara	mine	70.0483	27.6046	<i>B. n. nana</i>	2015-08-20	1	<i>Phyllonorycter</i> sp.
Reisa	mine	69.5256	29.3207	<i>B. n. nana</i>	2014-08-17	5	<i>Coleophora</i> sp.
Reisa	mine	69.5256	29.3207	<i>B. n. nana</i>	2014-08-17	4	<i>Coleophora</i> sp.

Ritsem	mine	67.7731	17.5361	<i>B. n. nana</i>	2014-08-18	31	Gracillariidae
Ritsem	mine	67.7731	17.5361	<i>B. n. nana</i>	2014-08-18	4	<i>Coleophora</i> sp.
Sånfjället	mine	62.2826	13.4752	<i>B. n. nana</i>	2014-07-14	1	<i>Coleophora</i> sp.