Section: forum article

A review of the use of black soldier fly larvae, *Hermetia illucens* (L., 1758) (Diptera: Stratiomyidae), to compost organic waste in tropical regions

Gardenio Diogo Pimentel da Silva [a]*; Thomas Hesselberg [b]

[a] *Energy Planning Program, Institute for Graduate Studies and Research in Engineering at the Federal University of Rio de Janeiro, Rio de Janeiro, Brazil.*

[b] *Department of Zoology, University of Oxford, Oxford, UK.*

E-mail for the correspondent author * gardenio.diogo01@ppe.ufrj.br or engambgardenio@gmail.com

A review of the use of black soldier fly larvae, *Hermetia illucens* (L., 1758) (Diptera: Stratiomyidae), to compost organic waste in tropical regions

Abstract: *Hermetia illucens* (L., 1758) is a fly of the family Stratiomyidae frequently found in tropical zones. Adults flies are not considered pathogens as they incapable of biting and feeding thus transmitting sicknesses to humans. The larval stage feeds off organic matter and offer a rich protein source naturally consumed by animals. The use of black soldier fly (BSF) larvae to treat organic waste is growing around the world. This is especially true for tropical low- and middle-income countries as their favourable climate conditions means that the BSF technology has significant potential to solve existing problems associated with a poorly developed sanitation infrastructure. In this study, we evaluate the feasibility, benefits, and limitations of implementing BSF projects in tropical regions using Belem, in Brazil, as a case study. Black soldier fly prepupae, arising from the waste reduction process, can be used as animal feed. It therefore offers potential to promote regional development, create jobs, and dispose of organic waste locally. Legal requirements as outlined in the Brazilian National Policy on Waste offer further incentives. However, more studies are needed to compare BSF waste reduction efficiency and prepupae yield to other technologies such as traditional composting and vermiculture, which can inform the decision making for implementation of organic waste treatment facilities.

Keywords: biological waste reduction, animal feed, regulations, prepupae yield, green disposal, sanitation

1. Introduction

The field of applied entomology has been growing in importance in recent years. While it has traditionally focused on using insect predators for biological control of pest species in agro-ecological systems (Cock *et al* 2016; Van Lenteren *et al* 2018), more emphasis is now also placed on using insects for developing biologically inspired technologies (Gorb 2011; Lenau *et al* 2018), as alternative food sources for production animals and humans (DeFoliart 1989; Van Huis 2013) and for reuse organic waste (Diener *et al* 2011b, 2011a). In terms of the latter, black soldier fly (BSF) larvae, *Hermetia illucens (L., 1758) (Diptera: Stratiomyidae)*, are particularly interesting as they offer a cost-effective potential alternative for recycling biological waste. The flies can be found tropical, subtropical as well as some temperature zones. Adults flies do not have mouthparts,

stingers or digestive organs and are thus incapable of attacking and feeding (Park 2015; Al-Qazzaz *et al* 2016). This in turn means that even if black soldier flies get in contact with food, they, unlike house flies, do not transmit pathogens from waste to humans (Banks *et al* 2014; Kenis *et al* 2014; Dortmans 2015).

Many studies have pointed out BSF's potential to convert municipal solid waste, pig manure, kitchen waste, human faeces, fruit and vegetable, rice straw, and other organic materials into a stable and valuable product (Diener *et al* 2009b, 2011a, b; Lalander *et al* 2013; Nguyen *et al* 2015; Manurung *et al* 2016; Dortmans *et al* 2017). Treatment with black soldier fly larvae can also reduce the half-life of pharmaceutical substances (Lalander *et al* 2016) and nutrients such as nitrogen and phosphor from waste (Myers *et al* 2008). The fly's prepupal stage, i.e. the last larval stage, is a nutritious source of proteins that can be exploited as animal feed for poultry, pigs, fish, and amphibians, and thereby generate additional income (ST-Hilaire *et al* 2007a, b; Diener *et al* 2011a, b; Nguyen *et al* 2015; Park 2015; Spranghers *et al* 2016). There have been studies on food security pointing to the usage of insects, such as BSF larvae, as an important source of protein for some countries in Africa and Asia (DeFoliart 1989). Insects, such as BSF, meets nutritional requirements for both animals and humans as they are rich in vitamins, micronutrients, saturated and polyunsaturated fatty acids (Rumpold and Schlüter 2013; Van Huis 2013).

However, the BSF technology is not yet a well-established technology for treating organic matter and transforming it into a useful product. Even though substantial initial investments are not needed (Manurung *et al* 2016). The three most used organic conversion processes are composting, vermicomposting, and biomethanation (organic material is microbiologically converted under anaerobic conditions to biogas) (Diener *et al* 2009a, 2011a). The BSF technology has the potential to outperform these methods in tropical regions, but more information on the bioconversion efficiency of agricultural waste, municipal solid waste (the organic fraction), excreta

and faecal sludge in low and middle income countries is required (Lohri *et al* 2017). The potential benefits of the BSF technology is particularly large in developing countries in the tropics as these have climates favouring maggot development, while often also having poor waste disposal systems with unregulated waste disposal in streets, dumping sites, rivers and other areas (Lohri *et al* 2017). Unregulated waste disposal leads to environmental pollution such as disagreeable odour, greenhouse gas emissions, toxic substances leaching to soil and groundwater, and attraction of vector borne diseases. Poor management of municipal solid waste may affect an entire population due to contamination of surface and underground water supplies. Inadequate solid waste disposal can also influence the local economy. For example, if waste management systems function properly, they can generate job opportunities and local economic development (Diener *et al* 2009a).

Brazil is a good example of a developing country that can benefit from usage of the BSF technology, because most of its territory is situated in the tropics and black soldier flies are naturally occurring in the wild. Furthermore, Brazil does not have an efficient traditional waste disposal system. These conditions are especially true of the Northern Region of the country (the Amazon region). The state of Para, the most populated State in the Amazon region accounting for about 8.3 million habitants throughout 144 municipalities, faces several challenges regarding waste disposal management (IBGE- Brazilian Institute of Geography & Statistics 2015). A total of 7 tonnes of waste are produced daily in the entire state, but only 5.4 tonnes/day are collected. Moreover, 35% is disposed in open dumping, 37% goes to controlled landfill dumping, whereas only 28% receives proper disposal management in sanitary landfill (Abrelpe 2015).

Despite the fact that the black soldier fly can be found in northern Brazil (Pujol-Luz *et al* 2008), there has, to our knowledge, only been a few limited studies using BSF larvae to treat organic waste on isolated farms in Brazil (Silva *et al* 2018; Teixeira Filho 2018). There is a need for more studies on the potential to treat organic waste from urban and large rural communities in

Brazil. Thus, the present review aims to investigate the feasibility, benefits and limitations for a BSF project implementation in tropical regions using Belem as a case study.

2. Black soldier fly larvae as animal feed source

The intense utilisation of soy meal or fishmeal have significant negative impacts on the environment. Hence, insects may play an important role in replacing them as a protein source (Sánchez-Muros *et al* 2014). Black soldier fly larvae are rich in lipids and oils which can be utilised to feed animals (Makkar *et al* 2014; Henry *et al* 2015; Spranghers *et al* 2016). A previous study demonstrated the feasibility of using BSF larvae to convert poultry waste into feedstuff for tilapia and catfish (Bondari & Sheppard 1981). There are other studies pointing out the feasibility of utilising prepupae as a source of fishmeal rich in protein. In the Republic of Guinea, for instance, BSF larvae have been reported to digest proteins and fats locked inside palm kernels. The proteins in these palms are not directly accessible to fish. However, if they are processed through BSF treatment, the resultant larvae can be used as fish feed. The authors also state that results from experiments open up possibilities to use BSF to compost other seed waste (Hem *et al* 2008). A study-case presented in (Mohd-Noor *et al* 2017) uses waste from coconut endosperm to feed BSF larvae and enrich its lipid and protein content. This is particularly interesting as the Para State is the largest producer of açaí palm, *Euterpe oleracea Martius*, in the country. In the process of extracting the açaí pulp, only 15% of the fruit is used in the production of açaí vine, while the remaining 85%, which is composed of fibres and seeds, is discarded (Melo *et al* 2017). As a result, thousands of tonnes of daily açaí palm waste is not correctly disposed of. There is, therefore, great potential to exploit the application of BSF larvae to compost açaí palm seeds and generate a new source of investment in the State. Tropical regions are rich in biodiversity of palm seeds thus other fruit can be used to convert palm kernel into animal ration.

Tropical countries, including Brazil, have a rich diversity of fish species, which can be cultivated in tanks (Junk *et al* 2007). Pirarucu fish, *Arapaima gigas* (Cuvier 1817), for example, could be fed through a mixed diet of fishmeal and BSF larvae (Imbiriba 2001; Oliveira *et al* 2012; Queiroz-de Oliveira *et al* 2013). Using BSF prepupae to supplement fishmeal do not have a negative impact on fish development (Bondari & Sheppard 1981; Kroeckel *et al* 2012). For example, replacing fishmeal with a diet that contained up to 15 % BSF larvae (protein content) did not have any negative effects on fishes' feed conversion ratio. Moreover, the same diet allowed a reduction of fishmeal from 36% to 27% and fish oil content from 13% to 8%. (ST-Hilaire *et al* 2007b). Their results are significant because they might represent an alternative to decrease the expenses associated with fish food and make fish farming more attractive to rural communities.

Black soldier fly larvae can also be used to replace or supplement the diet of poultry and livestock, though more research still has to be done to estimate the quantities required to sustain growth (Teguia & Beynen 2005; Veldkamp & Bosch 2015). Two recent studies found that using BSF larvae as feed source for laying chicken can increase eggshell thickness, egg yolk, and egg albumin (Barbosa-Filho *et al* 2018; Kawasaki *et al* 2019). Secci *et al* (2018) analysed the impact of replacing laying hen's soya-based diet with BSF larvae diet. The authors claim that BSF can be used as total substitute of traditional soya ration. However, this is only true when egg quality is the target. Other studies should address the effects on the entire animal. In addition, egg yolk from hen fed on BSF larvae presented lower cholesterol and similar fatty acids. Similar results were obtained with quails (Dalle Zotte *et al* 2019).

Black soldier fly larvae have been used as supplement food for swine (Newton *et al* 1977), where it can be used to both feed the animals and digest the manure produced (Newton *et al* 2005a, b; Veldkamp *et al* 2012). When used to treat manure, BSF larvae can reduce *Escherichia coli* (E. 1885) content from the compost (Liu *et al* 2008) although thermal treatment has to be given to the

prepupae before they can be fed to animals. However, it is important to note that there are still concerns regarding manure composting and using the residual larvae to feed animals. Another benefit associated with a BSF larvae diet is the reduced use of land, water consumption and lower emission of greenhouse gases in comparison to livestock derived products such as soy bean and maize (Rumpold & Schlüter 2013). Terrence & Popa (2012) compliments by affirming that the ammonia (NH_4) concentration can be 5-6 times higher than in the initial substrate, which could be advantageous for waste processing facilities as substances such as NH_4^+ and NH_3^- can easily be absorbed by plants and soil or recovered.

3. Black soldier fly system design and insect maintenance

A BSF system is not complex to build and can be adapted according to local requirements. There are two main parts of the system, the larvero, which is the place where larvae grow, and the fly house where adult flies will live and reproduce (Fig 1). The initial step is to acquire BSF eggs, which can be obtained in the market or captured in the wild of most tropical regions. In terms of Brazil, there are two studies which captured BSF's eggs in Belem (Silva *et al* 2018) and Amazonas (Teixeira Filho 2018), both in the Amazon region. However, BSF can also be found in other states outside the amazon region such as Minas Gerais and in the northeast region.

[insert Fig 1 here]

For capturing eggs in the wild, an "egg trap" can be set up. An egg trap consists of a rearing box (plastic box), food source (vegetables, fruit, manure, or other organic matter), a nylon screen, and corrugated cardboard pieces. Basically, one should place the rearing box filled with a food source out in the open to attract BSF females. The nylon net is used to allow spreading odour and block potential parasitic and predatory insects from entering the system. The cardboard pieces are

placed on top of the nylon net, close to the food source (Sripontan *et al* 2017). The female flies will deposit their eggs on corrugated cardboard where they can be extracted from and placed in the larvero system (Boaru *et al* 2019). Sripontan *et al* (2017) analysed different food diets to attract BSF females and found that oviposition was higher near fruit waste traps in comparison to animal manure. However, many types of decaying organic diets can be used to attract BSF females.

Middle to large scale or urban facilities might have their own nursery unit to provide eggs for the treatment facility. The adult flies are usually kept in a screened cage separately from the larvero. Cicková and co-workers (2012), for example, utilised two experimental metal cages, one with dimensions of 30 x 30x 30 cm with a capacity of 1000-1500 pupae and another with dimensions of 60 x 80 x 145 cm capable of holding 25,000 pupae. Their oviposition scheme used a food diet chosen to attract females to deposit their eggs on a cardboard structure. However, in order to avoid problems associated with inbreeding, self-contained facilities should occasionally be supplemented with wild-caught eggs or larvae.

The recommendation for the larvero is to maintain a regular food supply, a drainage system for leachates, and a ramp where the larvae can crawl out for the prepupae stage, which also ensures easy separation of pupae from larvae (Diener *et al* 2011a). The mass of BSF larvae required per system can be estimated by equation 1 (Dortmans *et al* 2017):

$$\frac{M_{system} = L_{system} * TM_{cont}}{TL_{system}} \qquad (1)$$

Where: M_{system}: mass of BSF larvae required per composting system;

L_{system}: number of larvae needed per system; this number is roughly 600-800 larvae/Kg of wet waste;

TM_{cont} = total mass of larvae in the rearing container, which is simply the weight of larvae in the composting box;

TL_{system} = total number of larvae in the system. This can be determined multiplying the TM_{cont} by number of larvae in a given sample then dividing it by the mass of the sample.

For household sized systems, the growing of larvae might be done in plastic buckets, which has to be adapted to include a ramp and drainage. Even a traditional three bucket composting bin scheme can be used to grow BSF larvae in the feeding phase, as used in a rural system reported by (Silva *et al* 2018). For middle to large sized composting facilities, concrete basins with a ramp can be used too, but they are usually set for manure management (Sheppard *et al* 1994; Newton *et al* 2005a). The size varies with the amount of waste added to the system and the number of larvae within the system. The retention period depends on factors such as surrounding temperatures (Sheppard *et al* 2002), age at inoculation, and type of material (Dortmans 2015). Anaerobic conditions can be avoided if the material height inside the container does not exceed 7.5-10 cm (Čičková *et al* 2015).

Dortmans (2015) used a 6.8 litre bucket (0.25m x 0.55m x 0.5m) in his experiment to compost 132 kg of waste in 47 days, but usually the larvae are ready to be harvested after 12 days (Dortmans *et al* 2017). Other authors adopted different volumes such as a 72 litre bucket (Mutafela 2015) and a 37 litre BSF reactor (Lalander *et al* 2014). For a small to medium scale project, it will be necessary to have containers with a volume above 75 litres (UNC Institute for the Environment 2013). In addition, a recent study demonstrated that black soldier fly larvae strains from different parts of the world show distinct phenotypic plasticity (different development time and waste conversion efficiency) (Zhou *et al* 2013). These differences, therefore, must be accounted for before selection and establishment of the waste management system. We have identified two pilot

studies using BSF larvae to compost in the Brazilian Amazon region, one bucket experiment in the State of Para (Silva *et al* 2018) and another lab scale system in the Federal University of Amazonas, State of Amazonas (Teixeira Filho 2018). Both studies indicate significant potential to use these insects to compost organic matter and produce an added-value by-product. Thus, we recommend conducting a pilot project with BSF larvae to determine its local feasibility as a waste management tool in cities to compost urban wastes.

There are large scale facilities using BSF larvae to produce protein that treat up to 200 tonnes of waste/day (Diener *et al* 2015). However, information on the design and operational procedures of successful large existing BSF treatment facilities are commercially sensitive and not openly shared (Lohri *et al* 2017). A study in Costa Rica estimated a treatment area of a facility to compost 3 tonnes/day of municipal organic waste at about 930 m^2 plus the facility area giving roughly 3.2 kg compost per m^2 per day (Diener *et al* 2009a), compared to vermicomposting with an estimated 2.4 kg compost per m^2 per day based on a much smaller scale plant (Marsh 2009). Recently, Lalander *et al* (2018) compared three strategies for composting food waste and faeces through vermicomposting, BSF larvae, and BSF larvae integrated with an anaerobic digestion. The latter configuration presented the highest final product value due to the combination of biogas, animal feedstuff, and fertilisers whereas vermicomposting had the lowest final product value. However, more studies that directly compare the two methods, when it comes to area requirements, waste conversion, and costs, are needed.

Fig 2 illustrates industrial scale design for the larvero and mating cage whereas Fig 3 shows a house-made larvero. Fig 2 comes from Dortmans *et al*, (2017) who present design systems and procedures for larger facilities based on an operational facility in Indonesia. On this system, the larvae are manually collected and transported to other stages. Other systems might have a type of ramp, so the larvae can craw out larvero by themselves. The design in Fig 3 is similar to the one

proposed by Diener *et al* (2011a). On this design the larvae will migrate to the bucket for pupation (natural process), the prepupae is then collected from the bucket and manually placed in the fly house (some can be diverted to other destinations such as animal feed). Food is added to the larvero manually by opening the lid and placing a layer of approximately 5 cm of waste spread evenly over the area. The main weakness of this type of larvero is the drainage system which requires constant maintenance not to clog up. Alternative designs are easily found online thus constructing the system should not be a problem to small and middle rural communities.

[insert figure 2 and 3 here]

The costs for constructing the BSF system will depend on the scale planned (UNC Institute for the Environment 2013). Nonetheless, there are common requirements that must be accounted for independent of which technology is chosen to process organic waste. Table 1 summarizes these items for an urban organic treatment system. Dortmans *et al* (Dortmans *et al* 2017) provide a wider list of components for BSF systems and equations needed to dimensioning the facility. Rural projects may be simplified without mechanised equipment, huge infrastructure, trucks, office, and other items. Determining the exact dimensions can only be done through a pilot project with detailed field work assessment of local waste production, the local populations' preparedness to separate organic from non-organic waste, environmental conditions and landscape characteristics.

[insert table 1 here]

4. Application of BSF technology in Belem, Brazil

4.1.Brazilian National Solid Waste Policy and black soldier fly technology

One principle established in the Brazilian National Waste Policy (BNWP), law n° 12,305/2010, section VII is the recognition of solid waste, which can be recycled or reutilised, as an economic tool for generating job opportunities, extra income, and social value. Furthermore,

art. 36, section V, of the law determines that the municipality waste management plan must contain alternatives to correctly treat organic content from domestic wastes (Brazil 2010). Waste treatment is not only an environmental concern, it also represents a health and safety issue. Social inclusion and job opportunities for people who work as waste scavengers on dumping sites is also an objective of the Brazilian Policy on Waste (Brazil 2010).

In Costa Rica, as mentioned in the previous section, the BSF method was studied to process municipal organic waste with good results (Diener *et al* 2009a, b, 2011a, b) and BSF has frequently been used in many countries to compost manure and faecal sludge (Sheppard *et al* 1994; Lalander *et al* 2013). It provides opportunities for small businesses and individuals to earn extra income by collecting and treating organic residues (Diener *et al* 2009a, b, 2011a). Environmental licenses (installation and operational licenses), as set out by the National Environmental Council (CONAMA)'s Resolution n°237/1997 for upscaling projects to treat waste, may have simplified the license processes. Small projects might not even need environmental licenses to operate, but this can only be determined by local environmental agencies. Prepupae also have to be analysed according to the standards of the Ministry of Agriculture, Livestock and Supply's (MAPA) normative instruction n°04/2007 about best practices to produce animal food and hygiene conditions of the facilities (MAPA 2007). Residue from BSF larvae can be used as fertiliser for crop production (UNC Institute for the Environment 2013; Lohri *et al* 2017; Boaru *et al* 2018), but this also requires approval from regulations such as the federal decree n°4,954/2004 and the MAPA normative instruction n°25/2009 both of which concerns the usage of biofertilisers in agriculture (Brazil 2004; MAPA 2007). In Sarpong *et al* (2019), BSF composting residue presented good standard quality to be used as a soil fertilizer. The fly's residue is similar to vermicomposting's, which is a mixture of unassimilated material and BSF larvae excrement (Rosmiati *et al* 2017). Thus, it might require drying and screening before application on the soil. All those regulations

must not be viewed as obstacles to implement projects on waste management such as composting, vermiculture, and black soldier fly larvae. Instead they are tools to guarantee the quality of the final product and the safety of operational facilities.

The BSF technology is therefore a novel and potentially valuable tool to achieve the principles and objectives established in the Brazilian National Policy on Solid Waste, but further studies are required to compare its final products (prepupae and residue) to applied regulations regarding solid waste.

4.2.Temperature and humidity

Temperature and humidity are two key factors that influence feeding, development, and the lifecycle of black soldier flies (Dortmans *et al* 2017). The survival temperature for BSF larvae ranges from 0 to 45°C, although the ideal temperature for food consumption is around 30-35°C (UNC Institute for the Environment 2013). A study on bioconversion of human faeces by BSF stated ideal conditions varying from 25 to 30°C (Lalander *et al* 2013). Ideal conditions for reproduction were observed at 32°C, but temperatures ranging from 15 to 47°C were tolerated (Diener *et al* 2011a). However, the majority of studies work with temperature and humidity around 27°C and 67%, respectively (Diener *et al* 2009b; Banks *et al* 2014) or temperatures in the range of 20-35°C and humidity above 30% (Nguyen *et al* 2015) Larvae, reared at temperatures varying between 27 to 32°C, have a final weight 30% greater than those reared at higher or lower temperatures, though they require slightly more hours to develop in these temperatures (27-32°C) (Harnden & Tomberlin 2016). In addition to that, females' weight 17-19% more than males in temperature conditions ranging from 27 to 30°C (Tomberlin *et al* 2009). Humidity affects BSF egg eclosion and an environment with 70% relative humidity (RH) may increase the survival of adult flies by 2 to 3 days, while low RH increase the mortality of larvae (Holmes *et al* 2012).

Northern Brazil has a tropical climate with a relatively stable temperature and humidity throughout the year. The temperature for the State of Para is always greater than 18°C with the annual mean temperature ranging between 23-26°C. Humidity also presents high values above 80% with an annual mean of 90% (Silva & Souza 2016).

Climatic conditions are thus within optimal conditions for reproduction and feeding of these insects, which support the environmental potential to exploit BSF larvae technology in composting organic material in Belem and nearby cities. However, it is important to protect the treatment system (containers and greenhouse) from direct sunlight, because it can increase the temperature of the container and compromise the development and hence compost efficiency of the BSF larvae. Heussler *et al* (2018) demonstrate that indoor rearing with artificial light is also possible. The authors studied the effect of three different light-sources on fly's oviposition and longevity. While light-source did not significantly affect laying of eggs, flies grew under halogen lights experience half-life shorter than flies reared under LED lights. This might be related to the high temperature of halogen lights.

4.3.Organic generation in Belem and estimated waste reduction

The lack of data regarding organic residue generation in Belem associated with the absence of selective collection are two major challenges for projecting and implementing composting techniques to the municipal waste disposal system in the city. For a pilot BSF project to compost organic residues in the city, it is, therefore, necessary to acquire data on the organic matter content of the waste. Restaurants, universities, public markets and other confined areas are preferable starting points to acquire data. Through a literature review, data on waste generation in a university, a restaurant, a public market, and a private school was compiled (see Table 2). The waste generation ranged from 26 kg/day to 27,000 kg/day, at a university and a public market, respectively. Table 2

shows the percentage of the organic fraction that waste represented of the total, and the year in which that data was published.

[insert table 2 here]

The public market, Ver-o-Peso, has the highest organic waste generation and its total daily waste production is comparable to a neighbourhood in Belem with a population of 34,000 (Dos Santos *et al* 2014). The school and the university campus, on the other hand, produced from 26 to 85 kg/day, which suggests that a pilot project to study BSF onsite would be more suitable to be implemented at these sites. Table 3 provides information about the amount of waste fed into different BSF systems, residue from those systems, waste reduction achieved, prepupal yield, feed conversion ratio, and feed source for each project analysed. The organic waste reduction varied according to diet (feed source), the quantity of food added per larva (optimal feed ratio), type of waste (wet or dry matter).

[insert table 3 here]

The final prepupal yield will depend on variables such as temperature and humidity in the system, egg eclosion, reproduction rates, and other parameters. Diener *et al*, (2011a) had on average, in favourable conditions, a prepupae yield of 0.3 kg/m^2/day (wet weight). However, they claim that prepupal yield of 0.8–1.0 kg can be achieved if the system receives a daily diet input of 15 kg of fresh municipal per square meter. Prepupae yield of 46 g/kg from poultry manure has been reported (Čičková *et al* 2012), but generally 1000 kg of manure generates about 200 kg of prepupae (Diener *et al* 2015). In other words, about 20% of organic material can be converted into biomass (see also table 3) that can be used as animal feed and crop fertiliser or can be sold on to other local farms.

Waste conversion into prepupae biomass also depends on the type of waste given to larvae (Tschirner & Simon 2015) (Table 3). The reduction estimated here is, however, important because

it demonstrates the potential to decrease the organic waste produced at a range of sites in Belem including restaurants, organic market, and universities. Organic waste preferentially necessitates prompt processing and treatment due to risks of disease transmission and material putridness. Decentralised organic waste treatment systems using BSF is thus being proposed as an alternative to process organic waste in developing countries (Diener *et al* 2015). The waste treatment facility in Belem is located far from the urban centre, but smaller decentralised facilities distributed throughout the city would minimize the organic waste quantity directed to the inter-municipal sanitary landfill, especially concerning residues from açaí palm kernel whose production is widely distributed throughout the city. Lalander *et al* (2019) also suggest using abattoir waste with mixture of fruit and vegetable scraps as a suitable alternative to feed BSF larvae colonies and reduce urban organic waste. In the light of that, the Ver-o-Peso supermarket and other public markets can be a valuable source of organic waste for a BSF composting facility. This will prolong the lifetime of the waste disposal facility and minimize environmental pollution associated with poor organic waste management. Moreover, through semi-centralised BSF larvae facilities, transportation expenses can be reduced with up to 60% (Diener *et al* 2015).

Onsite pilot studies are necessary to investigate the insects' response to local conditions and to locate areas and potential benefits for construction of a semi-centralised waste management system in Belem.

5. Challenges and future research

Here we have pointed out several benefits of using BSF to compost organic residues including efficient waste reduction, the capacity to degrade diverse types of food, income from selling prepupae as animal feed, creation of jobs and of the resulting financial value of the waste.

However, challenges of using this technology also need to be taken into consideration in order to estimate its true potential for a project of any scale.

First, this technology is based on using living organisms to compost waste. Therefore, controlling the temperature, the heavy metals concentration in the compost and ensuring sufficient oxygen levels in the waste is required to guarantee survival of the insects (Diener *et al* 2011b). The majority of studies using this technology were conducted in developed countries and more research in developing countries is necessary to test its feasibility there (Bonso 2013). Another challenge is the public acceptance of using larvae in agriculture and as animal feed (Westerman and Bicudo 2005). The usage of larvae to feed freshwater fish and poultry is not generally a concern, because insects are a natural feed source for those animals (Van Huis 2013). Heavy metal accumulation in the food chain from prepupae fed to animals is also potentially a major concern and heavy metal contaminated sources must therefore be controlled (Diener *et al* 2011b, a). Future research must focus on developing ways to separate heavy metals from prepupae and residues.

In addition, on a small scale, labour requirements may be high and the market for the by-product is currently unknown. Regulation and nutritional standards for animal feeding or usage as compost on agricultural field may apply too and must be followed (Diener 2010; Mutafela 2015).

There is also a need for additional studies on the capacity of these insects to degrade agricultural waste such as coffee pulp (Lardé 1990), faecal sludge from sanitation facilities and onsite projects including small faecal waste from potential future sustainable dry toilet solutions (Lenau and Hesselberg 2015), municipal solid waste, excreta and all types of disposal management that leads to environmental impacts in developing countries (Diener 2010; Bonso 2013). Up-scaling this technology to a cost-effective system for bioconversion of municipal solid waste on a mass-rearing facility with steady, safe, and reliable production of prepupae/compost will be a challenge (Bonso 2013; Van Huis 2013). Waste segregation is a key factor for accomplishing good

results because it influences the effort required to separate the material and the final product (Diener *et al* 2009a, b). This will be a major challenge for treating municipal waste in Belem, because segregation of material at source is not a common practice in the city. Application of this technology (and any other on composting organic material) on a municipal waste treatment scale must be preceded by public education programmes on segregation and possible economic incentives. Finally, a study comparing installation/operation costs of the BSF system to other types of facilities (vermicomposting, composting, and others) would also be beneficial. Such a study would allow a possible comparison of "unit cost per ton of waste processed per day". Table 4 gives an overview of possible challenges/limitations and benefits/opportunities for BSF systems.

[insert table 4 here]

6. Conclusion

In this study, we have outlined the advantages of using black solider fly larvae to reduce organic waste, while simultaneously producing an income to the local population. For these reasons, interest in using BSF technology for organic waste treatment is growing around the world particularly in low- and middle-income countries due to favourable climate conditions, and existing problems associated with waste disposal. Some developing countries with studies concerning this technology are Costa Rica, Indonesia, Guinea, and China. Most effort is concentrated on manure management and faecal sludge treatment facilities (Sheppard *et al* 1994; Lalander *et al* 2013), though the application of these insects to compost municipal solid waste and agro-industrial waste is rising (Diener *et al* 2011a; Dortmans *et al* 2017). The prepupae resulting from the process can be used as animal feed due to its rich protein and nutrient content. The high prices associated with animal food make the usage of black soldier fly larvae an interesting alternative to traditional

animal feed, but other uses such as biodiesel production and agriculture are also feasible (Sánchez-Muros *et al* 2014). We argue that Brazil, where the black fly soldier fly is found in the wild, has a great potential for exploiting this technology because of its location, climate conditions, land availability, and the political will expressed in the National Policy on Solid Waste. The city of Belem, in particular, has a relative constant temperature and humidity close to the optimum conditions for black soldier fly development and survival. Although the city is quite large without open spaces available to implement this technology near the city centre, decentralised stations could be used to receive and pre-treat the organic waste from public markets, school/universities, and restaurants (Diener *et al* 2015). The next logical step is to set up local pilot projects to explore the feasibility, environmental and financial benefits in more detail.

Acknowledgements

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. We thank AuthorAid for connecting the two authors and making this paper possible.

Author Contribution Statement

GDPS and TH designed, wrote and revised the manuscript. Both authors have done substantial contribution in terms of writing and researching for the final paper.

Conflict of interest statement

The authors declare no conflict of interest.

References

Abrelpe (2015) Panorama dos resíduos sólidos no Brasil 2015, 1st edn. Abrelpe, São Paulo, Brazil

Al-Qazzaz MFA, Ismail D, Akit H, Idris LH (2016) Effect of using insect larvae meal as a complete protein source on quality and productivity characteristics of laying hens. Rev Bras Zootec 45:518–523. doi: 10.1590/S1806-92902016000900003

Banks IJ, Gibson WT, Cameron MM (2014) Growth rates of black soldier fly larvae fed on fresh human faeces and their implication for improving sanitation. Trop Med Int Heal 19:14–22. doi: 10.1111/tmi.12228

Barbosa-Filho J, Vargas TS A, Carneiro AKF, Saito AM, Ogasawara CT, Diniz LT, Simonelli SM, Oba A (2018) Sólidos totais e rendimento dos componentes dos ovos de poedeiras alimentadas com farinha de larvas de mosca de soldado negro (Hermetia illucens). In: XVI Congresso APA - Produção e Comercialização de Ovos. Associação Paulista de Avicultura, Ribeirão Preto, SP, pp 1–4

Boaru A, Vig A, Ladoşi D, Păpuc T, Struţi D, Georgescu B (2019) The use of various oviposition structures for the black soldier fly, Hermetia illucens L. (Diptera: Stratiomydae) in improving the reproductive process in captivity. Adv Agric Bot 11:12–20

Boaru A, Vig A, Ladoşi D, Struţi D, Păpuc T (2018) Studies regarding the fertilizing capacity of poultry manure biocomposted by fly larvae (Diptera : Stratiomyidae). Adv Agric Bot 10:114–121

Bondari K, Sheppard DC (1981) Soldier fly larvae as feed in commercial fish production. Aquaculture 24:103–109. doi: 10.1016/0044-8486(81)90047-8

Bonso NK (2013) Bioconversion of Organic Fraction of Solid Waste Using the. Kwame Nkrumah University of Science and Technology

Brazil (2010) National Solid Waste Policy. http://www.wiego.org/sites/default/files/resources/files/Pereira-Brazilian-Waste-Policy.pdf. Accessed 1 Sep 2019

Brazil (2004) Decree Nº 4.954 de 14 de Janeiro de 2004. Planalto Central, Brazil

BSF Farming (2012) FilterBug. https://www.blacksoldierflyfarming.com/picsvids/109-filterbug-v1. Accessed 26 Jun 2019

Cardoso RSS, Carneiro PFN, Yoshino GH, Correa R dos S (2009) Diagnóstico Dos Resíduos Sólidos Da Cidade Universitária Prof . José Da Silveira Neto Da Universidade Federal Do Pará – Ufpa – Belém - Pará. In: 25º Congresso Brasileiro de Engenharia Sanitária e Ambiental. Engenharia Sanitária e Ambiental, Recife, pp 1–24

Čičková H, Newton GL, Lacy RC, Kozánek M (2015) The use of fly larvae for organic waste treatment. Waste Manag 35:68–80. doi: 10.1016/j.wasman.2014.09.026

Čičková H, Pastor B, Kozánek M, Martínez-Sánchez A, Rojo S, Takáč P (2012) Biodegradation of pig manure by the housefly, musca domestica: A viable ecological strategy for pig manure management. PLoS One 7:. doi: 10.1371/journal.pone.0032798

Cock MJW, Murphy ST, Kairo MTK, Thompson E, Murphy RJ, Francis AW (2016) Trends in the classical biological control of insect pests by insects: an update of the BIOCAT database. BioControl 61:349–363. doi: 10.1007/s10526-016-9726-3

Cristina G, Souza P De (2010) Diagnóstico do manejo dos resíduos sólidos em uma escola da rede de ensino particular de belém, no estado do pará. Universidade de Taubaté

Cunha ER, Carneiro PFN (2007) Diagnóstico e proposta de Gestão Integrada dos Resíduos Sólidos Urbanos da Sede do Município de Curuçá / Pa. Estud Tecnológicos 3:37–46

Dalle Zotte A, Singh Y, Michiels J, Cullere M (2019) Black Soldier Fly (Hermetia Illucens) as Dietary Source for Laying Quails: Live Performance, and Egg Physico-Chemical Quality, Sensory Profile and Storage Stability. Animals 9:115. doi: 10.3390/ani9030115

DeFoliart GR (1989) The human use of insects as food and as animal feed. Am Entomol 35:22–36. doi: 10.1093/besa/35.1.22

Diener S (2010) Valorisation of Organic Solid Waste using the Black Soldier Fly, Hermetia illucens, in Low and Middle-Income Countries. Swiss Federal Institute of Aquatic Science and Technology

Diener S, Lalander C, Zurbruegg C, Vinnerås B (2015) Opportunities and constraints for medium- scale organic waste treatment with fly larvae composting. In: Conference Proceedings Sardinia (2015), Fifteenth International Waste Management and Landfill Symposium. Sardina, Cagliari, Italy

Diener S, Roa Gutiérrez F, Zurbrügg C, Tockner K (2009a) Are Larvae of the Black Soldier Fly - Hermetia illucens - a Financially viable option for Organic Waste Management in Costa Rica? Proc Sardinia 2009, Twelfth Int Waste Manag Landfill Symp 6

Diener S, Studt S, NM, Roa Gutiérrez F, Zurbrügg C, Tockner K (2011a) Biological treatment of municipal organic waste using black soldier fly larvae. Waste and Biomass Valorization 2:357–363. doi: 10.1007/s12649-011-9079-1

Diener S, Zurbrügg C, Gutiérrez FR, Nguyen DH, Morel A, Koottatep T, Tockner K (2011b) Black Soldier Fly Larvae for Organic Waste Treatment – Prospects and Constraints. Proc WasteSafe 2011 – 2nd Int Conf Solid Waste Manag Dev Ctries 52:1–8

Diener S, Zurbrügg C, Tockner K (2009b) Conversion of organic material by black soldier fly larvae: establishing optimal feeding rates. Waste Manag Res 27:603–610. doi: 10.1177/0734242X09103838

Dortmans B (2015) Valorisation of Organic Waste - Effect of the Feeding Regime on Process Parameters in a Continuous Black Soldier Fly Larvae Composting System. Dep Energy Technol MSc:41

Dortmans BMA, Diener S, Verstappen BM, Zurbrügg C (2017) Eawag - Black Soldier Fly Biowaste Processing - Step by step Guide. Eawag – Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

Dos Santos JIN, Da Silva YCR, Pacheco PR, C Lima SRM, Gomes MVCN (2014) Gerenciamento de resíduos sólidos no complexo ver-o- peso no município de Belém-Pará. In: XI Seminário Nacional de Resíduos Sólidos. ABES, Brasília, pp 1–6

Gorb SN (2011) Insect-Inspired Technologies: Insects as a Source for Biomimetics. Springer, Giessen, Germany

Green TR, Popa R (2012) Enhanced ammonia content in compost leachate processed by black soldier fly larvae. Appl Biochem Biotechnol 166:1381–1387. doi: 10.1007/s12010-011-9530-6

Harnden LM, Tomberlin JK (2016) Effects of temperature and diet on black soldier fly, Hermetia illucens (L.) (Diptera: Stratiomyidae), development. Forensic Sci Int 266:109–116. doi: 10.1016/j.forsciint.2016.05.007

Hem S, Toure S, Sagbla C, Legendre M (2008) Bioconversion of palm kernel meal for aquaculture: Experiences from the forest region (Republic of Guinea). African J Biotechnol 7:1192–1198. doi: 10.5897/AJB07.355

Henry M, Gasco L, Piccolo G, Fountoulaki E (2015) Review on the use of insects in the diet of farmed fish: Past and future. Anim Feed Sci Technol 203:1–22. doi: 10.1016/j.anifeedsci.2015.03.001

Heussler CD, Walter A, Oberkofler H, Insam H, Arthofer W, Schlick-Steiner BC, Steiner FM (2018) Influence of three artificial light sources on oviposition and half-life of the black soldier fly, hermetia illucens (diptera: Stratiomyidae): Improving small-scale indoor rearing. PLoS One 13:. doi: 10.1371/journal.pone.0197896

Holmes ALA, Vanlaerhoven SL, Tomberlin JK, Holmes LA, Vanlaerhoven SL (2012) Relative Humidity Effects on the Life History of Hermetia illucens (Diptera : Stratiomyidae). Physiol Ecol 41:971–978

IBGE- Brazilian Institute of Geography & Statistics (2015) Estados. https://cidades.ibge.gov.br/?sigla=pa. Accessed 17 Jan 2017

Imbiriba EP (2001) Potencial de criação de pirarucu em cativeiro. Acta Amaz 31:299–316

Junk WJ, Soares MGM, Bayley PB (2007) Freshwater fishes of the Amazon River basin: Their biodiversity, fisheries, and habitats. Aquat Ecosyst Heal Manag 10:153–173. doi: 10.1080/14634980701351023

Kawasaki K, Hashimoto Y, Hori A, Kawasaki T, Hirayasu H, Iwase S, Hashizume A, Ido A, Miura C, Miura T, Nakamura S, Seyama T, Matsumoto Y, Kasai K, Fujitani Y (2019) Evaluation of Black Soldier Fly (Hermetia illucens) Larvae and Pre-Pupae Raised on Household Organic Waste, as Potential Ingredients for Poultry Feed. Animals 9:98. doi: 10.3390/ani9030098

Kenis M, Koné N, Chrysostome CAAM, Devic E, Koko GKD, Clottey VA, Nacambo S, Mensah GA (2014) Insects used for animal feed in West Africa. Entomologia 2:107–114. doi: 10.4081/entomologia.2014.218

Kroeckel S, Harjes AGE, Roth I, Katz H, Wuertz S, Susenbeth A, Schulz C (2012) When a turbot catches a fly: Evaluation of a pre-pupae meal of the Black Soldier Fly (Hermetia illucens) as fish meal substitute - Growth performance and chitin degradation in juvenile turbot (Psetta maxima). Aquaculture 364–365:345–352. doi: 10.1016/j.aquaculture.2012.08.041

Lalander C, Diener S, Magri ME, Zurbrügg C, Lindström A, Vinnerås B (2013) Faecal sludge management with the larvae of the black soldier fly (Hermetia illucens) - From a hygiene aspect. Sci Total Environ 458–460:312–318. doi: 10.1016/j.scitotenv.2013.04.033

Lalander C, Diener S, Zurbrügg C, Vinnerås B (2019) Effects of feedstock on larval development and process efficiency in waste treatment with black soldier fly (Hermetia illucens). J Clean Prod 208:211–219. doi: 10.1016/j.jclepro.2018.10.017

Lalander C, Nordberg Å, Vinnerås B (2018) A comparison in product-value potential in four treatment strategies for food waste and faeces – assessing composting, fly larvae composting and anaerobic digestion. GCB Bioenergy 10:84–91. doi: 10.1111/gcbb.12470

Lalander C, Senecal J, Gros Calvo M, Ahrens L, Josefsson S, Wiberg K, Vinnerås B (2016) Fate

of pharmaceuticals and pesticides in fly larvae composting. Sci Total Environ 565:279–286. doi: 10.1016/j.scitotenv.2016.04.147

Lalander CH, Fidjeland J, Diener S, Eriksson S, Vinnerås B (2014) High waste-to-biomass conversion and efficient Salmonella spp. reduction using black soldier fly for waste recycling. Agron Sustain Dev 35:261–271. doi: 10.1007/s13593-014-0235-4

Lardé G (1990) Recycling of coffee pulp by Hermetia illucens (Diptera: Stratiomyidae) larvae. Biol Wastes 33:307–310. doi: 10.1016/0269-7483(90)90134-E

Lenau T, Hesselberg T (2015) Dry sanitation concepts with inspiration from nature. J Water, Sanit Hyg Dev 5:330. doi: 10.2166/washdev.2015.178

Lenau TA, Metze A-L, Hesselberg T (2018) Paradigms for biologically inspired design. In: Lakhtakia A (ed) Proc. SPIE 10593, Bioinspiration, Biomimetics, and Bioreplication VIII, 1059302. Denver, Colorado, United States, pp 1–20

Li Q, Zheng L, Cai H, Garza E, Yu Z, Zhou S (2011) From organic waste to biodiesel: Black soldier fly, Hermetia illucens, makes it feasible. Fuel 90:1545–1548. doi: 10.1016/j.fuel.2010.11.016

Liu QL, Tomberlin JK, Brady JA, Sanford MR, Yu ZN (2008) Black Soldier Fly (Diptera: Stratiomyidae) Larvae Reduce Escherichia coli in Dairy Manure. Environ Entomol 37:1525–1530. doi: 10.1603/0046-225x-37.6.1525

Lohri CR, Diener S, Zabaleta I, Mertenat A, Zurbrügg C (2017) Treatment technologies for urban solid biowaste to create value products: a review with focus on low- and middle-income settings. Rev Environ Sci Biotechnol 16:81–130. doi: 10.1007/s11157-017-9422-5

Makkar HPS, Tran G, Heuzé V, Ankers P (2014) State-of-the-art on use of insects as animal feed. Anim Feed Sci Technol 197:1–33. doi: 10.1016/j.anifeedsci.2014.07.008

Manurung R, Supriatna A, Esyanthi RR (2016) Bioconversion of Rice straw waste by black soldier fly larvae (Hermetia illucens L .) : Optimal feed rate for biomass production. J Entomol Zool Stud 4:1036–1041

MAPA (2007) Normative Instruction 4/2007. Planalto Central, Brasília

Marsh L (2009) Composting Your Organic Kitchen Wastes with Worms. Virginia

Melo IM, Pinheiro IS, Freitas JF (2017) Revalorização do caroço de açaí em uma beneficiadora de polpas do município de Ananindeua/PA: proposta de estruturação de um canal reverso orientado pela PNRS e logística reversa. Gestão da Produção, Operações e Sist 12:59–83

Mohd-Noor SN, Wong CY, Lim JW, Mah-Hussin MIA, Uemura Y, Lam MK, Ramli A, Bashir MJK, Tham L (2017) Optimization of self-fermented period of waste coconut endosperm destined to feed black soldier fly larvae in enhancing the lipid and protein yields. Renew Energy 111:646–654. doi: 10.1016/j.renene.2017.04.067

Mutafela RN (2015) High Value Organic Waste Treatment via Black Soldier Fly Bioconversion (Onsite Pilot Study). Royal Institute of Technology, Stockholm.

Myers HM, Tomberlin JK, Lambert BD, Kattes D (2008) Development of Black Soldier Fly (Diptera: Stratiomyidae) Larvae Fed Dairy Manure. Environ Entomol 37:11–15. doi: 10.1093/ee/37.1.11

Newton GL, Booram CV, Barker RW, Hale OM (1977) Dried Hermetia illucens larvae meal as a supplement for swine. J Anim Sci 44:395–400. doi: 10.2527/jas1977.443395x

Newton GL, Sheppard DC, Watson DW, Burtle GJ, Dove CR, Tomberlin JK, Thelen EE (2005a) The Black Soldier Fly, Hermetia Illucens, as a Manure Management / Resource Recovery Tool. Symp State Sci Anim Manure Waste Manag 0–5. doi: http://www.cals.ncsu.edu/waste_mgt/natlcenter/sanantonio/proceedings.htm

Newton L, Shepperd C, Watson DW, Burtle G, Dove R (2005b) Using the black soldier fly, Hermetia illucens, as a value-added tool for the management of swine manure. Tifton

Nguyen TTX, Tomberlin JK, Vanlaerhoven S (2015) Ability of Black Soldier Fly (Diptera: Stratiomyidae) Larvae to Recycle Food Waste. Environ Entomol 44:406–410. doi: 10.1093/ee/nvv002

Oliveira EG, Pinheiro AB, Oliveira VQ, Silva ARM, Moraes MG, Rocha ÍRCB, Sousa RR, Costa FHF (2012) Effects of stocking density on the performance of juvenile pirarucu (Arapaima gigas) in cages. Aquaculture 370–371:96–101. doi: 10.1016/j.aquaculture.2012.09.027

Park HH (2015) Black Soldier Fly Larvae Manual The Black Soldier Fly Larvae Manual. Student Showcase. Paper 14. Amherst

Pujol-Luz JR, Francez PADC, Ururahy-Rodrigues A, Constantino R (2008) The black soldier-fly, Hermetia illucens (Diptera, Stratiomyidae), used to estimate the postmortem interval in a case in Amapá State, Brazil. J Forensic Sci 53:476–478. doi: 10.1111/j.1556-4029.2008.00659.x

Queiroz-de-Oliveira V, Barreto-Matos AR, Araújo-Bezerra T, Campos-Mesquita PE, Queiroz-de Oliveira V, Moreira-de Moraes A, Goncalves-de Oliveira E, Gazzineo-de Moraes M, Rocha-de Dousa R, Farias-Costa FH (2013) Preliminary Studies on the Optimum Feeding Rate for Pirarucu Arapaima gigas Juveniles Reared in Floating Cages. Int J Aquac 3:147–151. doi: 10.5376/ija.2013.03.0025

Rosmiati M, Nurjanah KA, Suantika G, Putra RE (2017) Application of Compost Produced by Bioconversion of Coffee Husk by Black Soldier Fly Larvae (Hermetia Illucens) as Solid Fertilizer to Lettuce (Lactuca Sativa Var . Crispa) : Impact to Growth. In: Proceedings of International Conference on Green Technology. pp 38–44

Rumpold BA, Schlüter OK (2013) Potential and challenges of insects as an innovative source for food and feed production. Innov Food Sci Emerg Technol 17:1–11. doi: 10.1016/j.ifset.2012.11.005

Sánchez-Muros MJ, Barroso FG, Manzano-Agugliaro F (2014) Insect meal as renewable source of food for animal feeding: A review. J Clean Prod 65:16–27. doi: 10.1016/j.jclepro.2013.11.068

Sarpong D, Oduro-Kwarteng S, Gyasi SF, Buamah R, Donkor E, Awuah E, Baah MK (2019) Biodegradation by composting of municipal organic solid waste into organic fertilizer using the black soldier fly (Hermetia illucens) (Diptera: Stratiomyidae) larvae. Int J Recycl Org Waste Agric 2025:. doi: 10.1007/s40093-019-0268-4

Secci G, Bovera F, Nizza S, Baronti N, Gasco L, Conte G, Serra A, Bonelli A, Parisi G (2018)

Quality of eggs from Lohmann Brown Classic laying hens fed black soldier fly meal as substitute for soya bean. Animal 12:2191–2197. doi: 10.1017/s1751731117003603

Sheppard DC, Newton LG, Thompson SA, Savage S (1994) A value added manure management system using the black soldier fly. Bioresour. Technol. 50:275–279

Sheppard DC, Tomberlin JK, Joyce JA, Kiser BC, Sumner SM (2002) Rearing Methods for the Black Soldier Fly (Diptera: Stratiomyidae). J Med Entomol 39:695–698. doi: http://dx.doi.org/10.1603/0022-2585-39.4.695

Silva KR, Costa CA, Oliveira IV, Cruz WP (2018) Creation of larvae of the black soldier fly (Hermetia illucens) organic waste composting. Cad Agroecol An do VI CLAA, X CBA e V SEMDF 13:1–5

Silva GDP, Souza MJR (2016) Análise de variáveis de projeto de sistema solar fotovoltaico utilizando o modelo sam: uma comparação entre Belém, Fortaleza e Brasília. Rev Bras Energias Renov 5:297–312. doi: http://dx.doi.org/10.5380/rber.v5i2.46220

Spranghers T, Ottoboni M, Klootwijk C, Ovyn A, Deboosere S, De Meulenaer B, Michiels J, Eeckhout M, De Clercq P, De Smet S (2016) Nutritional composition of black soldier fly (Hermetia illucens) prepupae reared on different organic waste substrates. J Sci Food Agric 97:2594–2600. doi: 10.1002/jsfa.8081

Sripontan Y, Juntavimon T, Songin S, Chiu C-I (2017) Egg-trapping of black soldier fly, Hermetia illucens (L.) (Diptera: Stratiomyidae) with various wastes and the effects of environmental factors. Khon Kaen Agric J 45:179–184

ST-Hilaire S, Cranfill K, McGuire MA, Mosley EE, Tomberlin JK, Newton L, Sealey W, Sheppard C, Irving S (2007a) Fish offal recycling by the black soldier fly produces a foodstuff high in omega-3 fatty acids. J World Aquac Soc 38:309–313. doi: 10.1111/j.1749-7345.2007.00101.x

ST-Hilaire S, Sheppard C, Tomberlin JK, Irving S, Newton L, McGuire MA, Mosley EE, Hardy RW, Sealey W (2007b) Fly Prepupae as a Feedstuff for Rainbow Trout , Oncorhynchus mykiss. J world Aquac Soc 38:59–67

Teguia A, Beynen AC (2005) Alternative feedstuffs for broilers in Cameroon. Livest Res Rural Dev 17:

Teixeira Filho NP (2018) Devoradores de Lixo: Aspectos biológicos, produtivos e nutricionais da Mosca Soldado Negra Hermetia illucens (L., 1758) (Diptera; Stratiomyidae. Universidade Federal do Amazonas

Tomberlin JK, Adler PH, Myers HM (2009) Development of the Black Soldier Fly (Diptera: Stratiomyidae) in Relation to Temperature. Environ Entomol 38:930–934. doi: 10.1603/022.038.0347

Tschirner M, Simon A (2015) Influence of different growing substrates and processing on the nutrient composition of black soldier fly larvae destined for animal feed. J Insects as Food Feed 1:249–259. doi: 10.3920/JIFF2014.0008

UNC Institute for the Environment (2013) The Black Soldier Fly - How-to-Guide

Van Huis A (2013) Potential of Insects as Food and Feed in Assuring Food Security. Annu Rev Entomol 58:563–583. doi: 10.1146/annurev-ento-120811-153704

Van Lenteren JC, Bolckmans K, Köhl J, Ravensberg WJ, Urbaneja A (2018) Biological control using invertebrates and microorganisms: plenty of new opportunities. BioControl 63:39–59. doi: 10.1007/s10526-017-9801-4

Veldkamp T, Bosch G (2015) Insects: a protein-rich feed ingredient in pig and poultry diets. Anim Front 5:45–50. doi: 10.2527/af.2015-0019

Veldkamp T, Van Duinkerken G, Van Huis A, Lakemond CMM, Ottevanger E, Bosch G, B, Van Boekel MAJS (2012) Insects as a sustainable feed ingredient in pig and poultry diets - a feasibility study. Wageningen UR Livest Res (report 63:62. doi: ISSN 1570 - 8616

Westerman PW, Bicudo JR (2005) Management considerations for organic waste use in agriculture. Bioresour Technol 96:215–221. doi: 10.1016/j.biortech.2004.05.011

Zheng L, Hou Y, Li W, Yang S, Li Q, Yu Z (2012) Biodiesel production from rice straw and restaurant waste employing black soldier fly assisted by microbes. Energy 47:225–229. doi: 10.1016/j.energy.2012.09.006

Zhou F, Tomberlin JK, Zheng L, Yu Z, Zhang J (2013) Developmental and Waste Reduction Plasticity of Three Black Soldier Fly Strains (Diptera: Stratiomyidae) Raised on Different Livestock Manures. J Med Entomol 50:1224–1230. doi: 10.1603/ME13021




Fig 1**.** Flow diagram outlining the process.

Fig 2**.** Medium industrial size scheme for larvero (right) and mating cage (left). Source: (Dortmans *et al* 2017)








Fig 3**.** House-made small-scale larvero (the Filterbug Bin). With permission from Peter Barnard and www.blacksoldierflyfarming.com. Source: (BSF Farming 2012)

Table 1. General items for organic waste treatment system.

General requirements	BSF facility
Land	Residue maturation area (larvero)
Collection truck and fuel	Oven for killing and drying of prepupae
Maintenance of infrastructure/truck	Fly colony
Infrastructure	Fly House
Equipment	
Tool shed	
Sanitary facility	
Sorting area	
Storage of rejects/recyclables	
Screening and bagging of residue	
Office	
Expendables	
Salaries	
Training	
Telephone/water/electricity	

Table 2. Characterization of the data about waste production in Belem or nearby cities.

Location	Organic waste (wet weight)/day	Percentage	Year	Reference
City of Curuça	515 kg	52%	2004	(Cunha & Carneiro 2007)
Private School	85 kg	24.21%	2010	(Cristina & Souza 2010)
UFPA campus[1]	26 Kg	41%	2008	(Cardoso *et al* 2009)
Ver-o-Peso Public Market	27,000 kg[2]	-	2012	(Dos Santos *et al* 2014)

[1] Only a small part of the campus. [2] Mean generation of 27.68 Kg/person, 731 people who worked with organic material. 27.68*731 = 20234.08 Kg/day

Table 3. Information about BSF projects from literature.

Ref.	Local	Total amount of feed (dry weigh)	Total amount of feed (wet weigh)	Residue	Waste reduction (%)	Prepupal Yield	Feed conversion ratio (%)	Feed source
(Diener *et al* 2011a)	Costa Rica	151 Kg		48 kg	68	17.8 kg	14.5	MOW
(Dortmans 2015)	Sweden	-	136 Kg	7.3 kg	85	15.7 kg	19.0*	FHF
(Dortmans 2015)	-	32.9 kg		3.6 kg	71	6.4 kg	19.0*	FHF
(Dortmans 2015)	-	-	136 kg	7.7 kg	85	15.5 kg	18.9*	FHF
(Dortmans 2015)	-	32.9 kg		4.1 kg	71	6.3 kg	18.9*	FHF
(Newton *et al* 2005b)**	USA	67.8 kg		41.6 kg	39	26.2 kg	9.6	PM
(Banks *et al* 2014)	England		658 g	301 g	54.2	108 g	3.4	HF
(Banks *et al* 2014)	-		721 g	327 g	54.6	131 g	33.9	HF
(Banks *et al* 2014)	-		390 g	260.2 g	33.4	8.5 g	15.2	HF
(Banks *et al* 2014)	-		482 g	360.3 g	25.2	11.3 g	10.7	HF
(Banks *et al* 2014)	-		437 g	219.8 g	49.7	65.3 g	3.3	HF
(Banks *et al* 2014)	-		483 g	261.4 g	45.8	110.7 g	2.0	HF
(Mutafela 2015)	-	4.6 kg		2.8 kg	37	-	32.73	Manure
(Mutafela 2015)	-	3.3 kg		2.7 kg	16	-	13.60	Manure
(Mutafela 2015)	Sweden	2.0 kg		0.3 kg	83	-	20.94	Fruits
(Mutafela 2015)	-	1.2 kg		0.3 kg	74	-	13.94	Fruits

MOW: Municipal Organic Waste. PM: Pig Manure. FHF: food waste and human faeces. HF: Human Faeces. *Biomass conversion rate in %, ** calculations was done by (Diener *et al* 2011a).

Table 4. Comparison of benefits/opportunities and limitations/challenges

Benefits and opportunities	Limitations and challenges
Prepupae as an alternative animal feed	Current market for product is unknown
Larvae can produce oil for biodiesel (Li *et al* 2011; Zheng *et al* 2012) and compost to agriculture	Segregation at source needs to be fully established.
Possibility of generating jobs and extra income	Possible regulations about the products
Matches well with the principles and objectives on the Brazilian National Policy on Waste	Necessity for monitoring environmental conditions and breeding parameters.
Waste reduction is faster than composting and minimizes odour problems	Needs more research on market value and enhanced production.
Larvae digest diverse types of residues and are more resistant to environmental changes than worms	Potential build-up of heavy metals contamination on prepupae for animal feeding.
Sludge (faecal) treatment alternative	May require a high initial investment (Mutafela, 2015).
Black soldier flies are not pathogenic and can be found in the wild in Brazil	
Potential to reduce salmonella, viruses, and pharmaceutical substances in the waste	
It can be used to manage animal manure	
Reduce organic waste from organic public markets and increase landfill lifetime	
Reduce nitrogen and phosphor from waste	
Minimize environmental pollution	