

1
2
3 **Title**
4

5 Antibiotic prophylaxis for infective endocarditis in patients undergoing dental
6 procedures: a systematic review and meta-analysis
7

8
9 **Authors**

10
11 Cahill TJ^{*1}, Harrison JL^{*2}, Jewell P³, Onakpoya IJ⁴, Chambers J⁵, Dayer MJ⁶,
12 Lockhart PB⁷, Roberts N⁸, Shanson D⁹, Thornhill MH¹⁰, Heneghan C^{†11}, Prendergast
13 BD^{†12}
14

15 * joint first authorship

16 † to whom correspondence should be addressed
17

18
19 The Corresponding Author has the right to grant on behalf of all authors and does
20 grant on behalf of all authors, an exclusive licence (or non exclusive for government
21 employees) on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees
22 to permit this article (if accepted) to be published in HEART editions and any other
23 BMJ PGL products to exploit all subsidiary rights
24
25

26 **Authors' affiliations**
27

28
29 1. Dr T. J. Cahill
30 Oxford Heart Centre
31 John Radcliffe Hospital
32 Oxford
33 UK
34 thomas.cahill@cardiov.ox.ac.uk
35

36
37 2. Dr J. L. Harrison
38 Department of Cardiology
39 St Thomas' Hospital
40 London
41 U.K.
42 james.harrison@kcl.ac.uk
43

44
45 3. Dr P. Jewell
46 Department of Cardiology
47 John Radcliffe Hospital
48 Oxford
49 UK
50 pauldavidjewell@gmail.com
51

52
53 4. Dr I. J. Onakpoya
54 Centre for Evidence-Based Medicine
55 Nuffield Department of Primary Care Health Sciences
56 University of Oxford
57 Oxford
58 UK
59 igho.onakpoya@phc.ox.ac.uk
60

61
62 5. Professor J. Chambers

1
2
3 Department of Cardiology
4 St Thomas' Hospital
5 London
6 U.K.
7 jboydchambers@aol.com
8

9 6. Dr M. J. Dayer
10 Department of Cardiology
11 Taunton and Somerset NHS Trust
12 Taunton
13 Somerset
14 UK
15 markdayer@me.com
16

17 7. Professor P. Lockhart
18 Department of Oral Medicine
19 Carolinas Medical Center
20 Charlotte
21 NC
22 USA
23 peter.lockhart@carolinashealthcare.org
24

25 8. Ms Nia Roberts
26 Outreach Librarian
27 Knowledge Centre
28 Bodleian Health Care Libraries
29 Oxford
30 nia.roberts@bodleian.ox.ac.uk
31

32 9. Dr D. Shanson
33 Department of Microbiology
34 Great Ormond Street Children's Hospital
35 Great Ormond St
36 London
37 UK
38 davidsha87@hotmail.com
39

40 10. Professor M. H. Thornhill
41 Unit of Oral & Maxillofacial Surgery & Medicine
42 University of Sheffield School of Clinical Dentistry
43 Sheffield
44 UK
45 m.thornhill@sheffield.ac.uk
46

47 11. Professor C. Heneghan
48 Centre for Evidence-Based Medicine
49 Nuffield Department of Primary Care Health Sciences
50 University of Oxford
51 Oxford
52 UK
53 carl.heneghan@phc.ox.ac.uk
54

55 12. Professor B.D. Prendergast
56 Department of Cardiology
57 St Thomas' Hospital
58
59
60

1
2
3 London
4 U.K.
5 bernard.prendergast@gstt.nhs.uk
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Word count (excl title page, abstract, references, figures, tables)

2713

1
2
3 **Abstract**
4
5

6 Objective

7 The use of antibiotic prophylaxis for prevention of infective endocarditis is
8 controversial. In recent years, guidelines to cardiologists and dentists have advised
9 restriction of antibiotic prophylaxis (AP) to high risk groups (in Europe and the USA),
10 or against its use at all (in the UK). The objective of this systematic review was to
11 appraise the evidence for use of AP for prevention of bacteraemia or infective
12 endocarditis in patients undergoing dental procedures.
13
14
15
16

17
18 Methods

19 We conducted electronic searches in Medline, Embase, Cochrane Library and ISI
20 Web of Science. We assessed the methodological characteristics of included studies
21 using the Strengthening the Reporting of Observational Studies in Epidemiology
22 (STROBE) criteria for observational studies, and the Cochrane Risk of Bias Tool for
23 trials. Two reviewers independently determined the eligibility of studies, assessed the
24 methodology of included studies and extracted the data.
25
26
27
28
29

30 Results

31 We identified 178 eligible studies, of which 36 were included in the review. This
32 included 10 time-trend studies, 5 observational studies and 21 trials. All trials
33 identified used bacteraemia as an endpoint, rather than infective endocarditis. One
34 time-trend study suggests that total AP restriction may be associated with a rising
35 incidence of infective endocarditis, while data on the consequences of relative AP
36 restriction are conflicting. Meta-analysis of trials indicates that AP is effective in
37 reducing the incidence of bacteraemia (risk ratio 0.53, 95% CI 0.49-0.57, $p < 0.01$),
38 but case control studies suggest this may not translate to a statistically significant
39 protective effect against infective endocarditis in patients at low risk of disease.
40
41
42
43
44
45
46

47 Conclusions

48 The evidence base for the use of antibiotic prophylaxis is limited, heterogeneous and
49 the methodological quality of many studies is poor. Post-procedural bacteraemia is
50 not a good surrogate endpoint for infective endocarditis. Given the logistical
51 challenges of a randomised trial, high quality case control studies would help to
52 evaluate the role of dental procedures in causing infective endocarditis, and the
53 efficacy of antibiotic prophylaxis in its prevention.
54
55
56
57
58
59
60

1
2
3
4 **Key words**

5 Endocarditis

6 Antibiotic prophylaxis

7 Systematic review
8
9

10 **Abbreviations**

11 ACC/AHA - American College of Cardiology/American Heart Association

12 AP – antibiotic prophylaxis

13 ESC – European Society for Cardiology

14 IE – infective endocarditis

15 NICE – National Institute for Health and Care Excellence (UK)

16 NVE – native valve infective endocarditis

17 PVE – prosthetic valve infective endocarditis

18 RCT – randomized controlled trial

19 UK – United Kingdom

20 USA – United States of America

21 VGS – viridans group streptococci
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Introduction**
4

5
6 Infective endocarditis is a rare but life-threatening disease.[1] Despite trends towards
7 multidisciplinary 'heart team' care and early surgery, 1-year mortality approaches
8 30%.[2] In patients with prosthetic heart valves, rheumatic and congenital heart
9 disease, the risk of acquiring infective endocarditis is thought to be 10-50 fold higher
10 than that of the general population.[3] Effective strategies for prevention of both
11 community and healthcare-acquired infective endocarditis in at-risk groups are
12 required.[4]
13
14
15
16

17
18 The oral cavity was identified as a major portal of entry for bacteria in 1909 by
19 Thomas Horder.[5] Oral streptococci are commensal flora of the oropharynx and
20 account for 10-30% of cases of infective endocarditis, depending on the location, risk
21 factor profile and socio-demographic characteristics of the population studied.[6, 7]
22 Transient bacteraemia, which occurs in the setting of poor oral hygiene and
23 periodontal diseases, dental procedures, or in the course of normal daily activities
24 (e.g. tooth brushing), is thought to be a precursor to the development of some cases
25 of infective endocarditis.[8]
26
27
28
29
30
31

32 For over 50 years, oral antibiotic prophylaxis (AP) was given to patients at risk of
33 infective endocarditis undergoing dental procedures. Between 2007 and 2009,
34 however, the European Society for Cardiology (ESC), American Heart
35 Association/American College of Cardiology (AHA/ACC) and the National Institute for
36 Health and Care Excellence (NICE) recommended restriction of AP to varying
37 degrees.[9, 10, 11] In Europe and the USA there was relative AP restriction to those
38 at highest risk (e.g. patients with previous infective endocarditis, congenital heart
39 disease and rheumatic heart disease, and selected heart transplant recipients)
40 undergoing high-risk dental procedures. In the UK, NICE advised against use of
41 prophylaxis entirely (total AP restriction) in 2008 but softened this stance in July 2016
42 to state that antibiotics should not *routinely* be recommended as prophylaxis for
43 dental procedures.[12]
44
45
46
47
48
49
50

51
52 The rationale for relative or total AP restriction was threefold. First, as medicine
53 shifted towards evidence-based practice there was (and remains) no randomised
54 controlled trial assessing the efficacy of AP for prevention of infective endocarditis.
55 Second, the relative importance of dental procedures as a cause of infective
56 endocarditis remained in doubt, compared with other portals of entry or low grade
57
58
59
60

1
2
3 recurrent bacteraemia occurring in the course of daily life.[8, 13] Third, in moderate
4 risk (and high risk in England) groups, the overall hazards of antibiotic use
5 (particularly anaphylaxis and the development of antibiotic resistance) were felt to
6 weigh against use of AP. The NICE guideline committee also deemed that AP was
7 not cost effective as a result of lack of efficacy and the perceived risks of anaphylaxis.
8
9

10
11 The primary object of this study was to provide a systematic review and synthesis of
12 evidence that directly or indirectly informs clinical use of AP for at-risk patients
13 undergoing dental procedures. The evidence base comprises three types of study:
14 first, trials examining the effect of AP on the incidence of bacteraemia following
15 dental procedures; second, observational studies assessing the efficacy of AP for
16 prevention of infective endocarditis; and finally, time trend studies which examine the
17 effect of changing national or international AP guidelines on the population incidence
18 of infective endocarditis.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Methods

Eligibility and search strategy:

We searched the following databases from inception until 25 February 2016 to identify studies of the efficacy of AP for the prevention of bacteraemia or infective endocarditis in patients undergoing dental procedures: Medline & Medline In-Process (OvidSP) [1946-present], Embase (OvidSP) [1974 to 2016 February 08], Cochrane Central Register of Controlled Trials (Cochrane Library, Wiley) [Issue 1 of 12, January 2016], Cochrane Database of Systematic Reviews (Cochrane Library, Wiley) [Issue 2 of 12, February 2016], Database of Abstracts of Reviews of Effects (Cochrane Library, Wiley) [Issue 2 of 4, April 2015], Science Citation Index Expanded & Conference Proceedings Citation Index- Science (Web of Science Core Collection) [1945-present], Clinicaltrials.gov (<http://clinicaltrials.gov>) and the WHO International Clinical Trials Registry Platform (<http://apps.who.int/trialsearch/Default.aspx>). Search terms used included subject headings and title/abstract keywords for bacterial endocarditis, antibiotics and prophylaxis (see search strategy in Appendix 1). We also searched the reference lists of all included articles. The following categories of study were excluded: studies conducted prior to 1960, studies of AP in patients undergoing cardiac surgery or implantation of cardiac electronic devices, topical therapies, and comparative antibiotic trials with no placebo/control arm.

Data abstraction:

We assessed methodological quality of studies using the Cochrane Risk of Bias tool[14] (for trials) or a checklist adapted from the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) criteria (for observational studies).[15] Two reviewers (TJC and JLH) independently adjudicated the eligibility of studies, assessed the methodology of included studies, and performed data extraction. Disagreement was resolved through consensus.

We extracted data on the study design: for case control studies we extracted baseline characteristics on the cases and the controls; for time trend studies we extracted study population characteristics, the study time period, relevant guideline changes and effects on incidence of IE per 100,000 population. The primary outcome of interest was the incidence of infective endocarditis, incidence of (any) bacteraemia, or for time trend studies, population-adjusted incidence of infective endocarditis.

Where total incidence of bacteraemia was not reported, the time point at which the

1
2
3 highest incidence of bacteraemia was observed in the placebo group was used for
4 comparison.
5
6

7
8 Data analysis:

9 We derived summary tables to report methodological quality and main results of the
10 included studies according to study design. For pooled effects we used a fixed-
11 effects model to generate Forest plots and used odds ratios as the summary
12 measure. We assessed heterogeneity using I-square values, with 25%, 50% and
13 75% representing mild, moderate and substantial heterogeneity, respectively.[16]
14 Forest plots and data summary graphs were compiled using RevMan (Cochrane,
15 UK) and SPSS (SPSS Inc., Chicago, IL, USA), respectively.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

The electronic search identified 3830 articles, after removal of case reports, editorials, animal studies and duplicates (Figure 1). After screening of the title and/or abstract of these, 178 articles were deemed eligible for full text assessment. In total, 36 studies were deemed suitable for inclusion (see Supplementary Table 1 for excluded studies), comprising 10 time-trend studies, 5 observational studies (4 case-control studies and 1 retrospective cohort study) and 21 trials. All identified trials used bacteraemia as a surrogate endpoint for infective endocarditis. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flowchart for study inclusion is shown in Figure 1.

Time trend population studies

We identified 10 studies assessing the effect of changing national and international guidelines concerning the use of AP on the population incidence of infective endocarditis. These included 9 studies of relative AP restriction (from the USA and Europe) and 1 study examining the effect of total AP restriction (from the UK) (Table 1). Changes in the guidelines between 2007 and 2009 by the ESC, ACC/AHA and NICE greatly reduced the use of antibiotic prophylaxis. Annual incidence was reported in two studies[17, 18] and obtained from the authors for two studies.[19, 20] Figure 2 shows the incidence of infective endocarditis per 100,000 population before and after changes in ACC/AHA and NICE guidelines. While only 1 study identified a significant rise in the incidence trend of infective endocarditis, it is important to note that this change was observed in the only population with total AP restriction.

Observational studies

We identified 5 observational studies for inclusion, including 4 case-control studies and 1 retrospective cohort study (Table 2). Data extracted included characteristics of cases and controls (or the two cohorts which were compared[21]), exposures and interventions (i.e. invasive procedures, use of AP), and, where possible, the numbers of patients specifically undergoing dental procedures (Table 2). All studies were assessed to be at high risk of intrinsic methodological bias (Supplementary Table 3). Meta-analysis was conducted on 3 studies with available data concerning the numbers of cardiac patients exposed to dental procedures, use of AP, and infective

1
2
3 endocarditis outcome. Overall, the odds ratio for use of AP in patients with infective
4 endocarditis was 0.59 (95% confidence interval 0.27-1.30, $p = 0.14$, $I^2=48\%$),
5 suggesting no statistically significant difference in exposure to AP between cases
6 (patients with infective endocarditis) and controls. In Van Der Meer *et al*, cases and
7 controls were analysed up to 30 days post-procedure and subgroups combined (first
8 time and recurrent infective endocarditis; definite and possible indications for AP). If
9 patients without a definite AP indication were excluded, this study provided an overall
10 OR of 0.63 (95% CI 0.17-2.36) for AP, modifying the overall meta-analysis to an OR
11 of 0.47 (95% CI 0.21-1.06, $p = 0.07$).
12
13
14
15
16
17
18
19

20 **Bacteraemia trials**

21
22 We included 21 studies investigating the effect of AP on the incidence of
23 bacteraemia (as a surrogate for infective endocarditis) following a dental procedure.
24 All studies reported the incidence of bacteraemia in a placebo group compared to an
25 AP intervention group after a dental procedure. Some studies tested multiple
26 antibiotic regimens (detailed in Supplementary Table 3), and some compared
27 additional endpoints such as the duration or magnitude of bacteraemia, a breakdown
28 of specific organisms grown, or antibiotic sensitivity patterns. A forest plot
29 summarising a total of 35 antibiotic arms against control or placebo is shown in
30 Figure 4. AP was associated with a risk ratio of 0.53 (95% CI 0.49-0.57, $p < 0.01$, $I^2 =$
31 90%) for bacteraemia in patients following dental procedures.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 – Time trend studies examining effect of antibiotic prophylaxis guideline change on the incidence of infective endocarditis

First author, year	PMID	Paper/abstract	Region, Country	Population	Diagnosis	Study time period	Guideline change Level of AP restriction	Increased incidence post-guideline change	Increase in rate of change of incidence post guideline	Guideline time point identified by change point analysis?
Bates 2016[22]	27418041	Paper	USA	Children ≤ 18 yrs identified from Pediatric Health Information System Database (29 hospitals)	All cases – acute and subacute bacterial IE	2003 - 2014	AHA/ACC April 2007 Relative restriction	No	No	NA
Bikdeli 2013[23]	23994421	Paper	USA	Adults ≥ 65 yrs	All cases - principal or secondary discharge dx of IE	1999-2010	AHA/ACC April 2007 Relative restriction	No	NA	NA
Daye 2015[19] & Thornhill 2011[24]*	25467569	Paper	England, UK	All	All cases - primary dx acute or subacute IE	January 1 2000 - March 31 2013	NICE March 2008 Total restriction	Yes	Yes	Yes
DeSimone 2015[20] & De Simone 2012[25]*	26141329	Paper	Olmsted County, Minnesota, USA	Adults ≥ 18 yrs	VGS IE	January 1 1999 to December 31 2013	AHA/ACC April 2007 Relative restriction	No	NA	NA
Duval 2012[26]	22624837	Paper	3 regions of France (Greater Paris, Lorraine, and Rhône-Alpes)	Adults ≥ 20 yrs	All cases of IE and subgroups by causative organism	3 survey years - 1991, 1999, 2008	French guideline restrictions 2002 Relative restriction	No	NA	NA

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Keller 2016[27]	27816113	Paper	Germany	All patients hospitalized with acute or subacute IE	IE due to Streptococcus and Staphylococcus (reported separately)	2005 - 2014	ESC October 2009 Relative restriction	Yes	Yes	No
Mackie 2016[28]	26868840	Paper	Canada (except Quebec and the Northern Territories)	All patients hospitalized with acute or subacute IE as main diagnosis	All hospitalizations with primary dx of IE	April 2002 - March 2013	AHA/ACC April 2007 Relative restriction	Total IE increase, decrease in VGS	Yes	No
Pant 2015[17]	25975469	Paper	USA	Patients in Nationwide Inpatient Sample with ICD codes for IE	All cases of IE and subgroups by causative organism	2000 - 2011	AHA/ACC April 2007 Relative restriction	Yes	Yes	Not performed
Salam 2014[29]	NA	Abstract	Qatar	All patients hospitalized with IE in State of Qatar	All cases	2002 - 2012	AHA/ACC April 2007 Relative restriction	No	NA	NA
van den Brink 2016[18]	Pending	Paper	Netherlands	All patients identified from the national healthcare insurance database	All cases of IE	2005 - 2011	ESC October 2009 Relative restriction	Yes, significant increase in VGS	Yes	NA

* Earlier publications from same research group using same methodology. Study with longer follow-up used for analysis

Table 2 – Observational studies of antibiotic prophylaxis and infective endocarditis

Study	Study design	Cases	Controls	Exposures and/or outcomes analysed	Outcomes for patients undergoing dental interventions	Included in meta-analysis?
Horstkotte 1987[21]	Retrospective cohort study	Group 1: 229 patients with prosthetic heart valves, in whom 287 diagnostic/therapeutic procedures were performed using a prophylactic antibiotic regime	Group 2: 304 patients with prosthetic heart valves, who had undergone "one of the procedures we regard as requiring endocarditis prophylaxis, without having received any antibiotic regime"	Incidence of PVE	Subgroup analysis of patients undergoing dental procedures: 2 cases of PVE from 117 patients undergoing dental procedures without AP. 0 cases (total number of patients undergoing dental procedures is not reported) in those with AP	No – as no denominator provided for total number of dental procedures in the group which received AP
Imperiale 1990[30]	Case control	Patients with a first episode of NVE within 12 weeks of a dental procedure	Patients matched for age, high-risk lesion on echo and frequency of dental visit	Use of antibiotics pre and after procedure	AP used in 1/8 cases, 15/24 controls	Yes
Lacassin 1995[31]	Case control	Adults with definite, probable or possible IE. PVE < 1 year from implantation excluded	Subjects without IE recruited from wards/echocardiography during same period as cases. Matched 1:1 for sex, age, group of underlying cardiac condition	All invasive procedures (not limited to dental) in 3 month period prior to IE, and use of AP	Subgroup analysis of patients undergoing dental procedures with cardiac disease: AP used in 6/26 cases, 6/22 controls	Yes – only patients undergoing dental procedures
Strom 1998[32]	Case control	Persons with community-acquired IE not associated with IV drug use (39 prosthetic valve) 104/273 cases had cardiac disease	Community residents, matched by age, sex, and neighbourhood of residence – i.e. not matched by cardiac risk factors 17/283 had cardiac disease	Survey on risk factors including recent dental treatment, use of AP	Only the patients with cardiac risk factors are relevant Study does not state what proportion of this subgroup received AP	No – details not provided for number of controls with cardiac disease given antibiotic prophylaxis – without this cannot be compared to the cases
Van der Meer 1992[33]	Case control	Patients with NVE and cardiac risk factors and <180 days post procedure (medical or dental) requiring prophylaxis	Cardiac outpatients matched for age and procedures	Survey of recent dental procedures and use of AP Caveat is that not all interventions have 'definite' indication for AP	30-day rate of IE analysed for cases Equivalent time period used for controls follow-up (Table 3) AP used in 5/25 cases, 8/42 controls	Yes, although not possible to separate out the number of patients who underwent dental procedures (cf. other medical procedures) although these account for <10% of both cases and controls

Abbreviations: AP – antibiotic prophylaxis, IE – infective endocarditis, NVE – native valve endocarditis, PVE – prosthetic valve endocarditis

Discussion

In this study we have systematically reviewed the evidence base for the use of AP for prevention of infective endocarditis. This comprises a) population time trend analyses of the effect of changing national and international guidelines on the incidence of infective endocarditis b) focused observational studies, including 4 case control studies and a retrospective cohort, and c) trials of antibiotics after dental procedures, using bacteraemia as a surrogate endpoint for the development of infective endocarditis. No randomised controlled trial (RCT) of AP has been undertaken.

This is the first study to systematically appraise the total evidence base for AP across a range of study designs. We have conducted a comprehensive search and extensively reviewed studies that either directly or indirectly address the question of AP efficacy. All studies have been quality assessed, with risk of bias assessed in a systematic manner. However, our study has some limitations. Our conclusions are limited by the poor methodological quality of included studies (and their heterogeneity) and the lack of randomized trials. Furthermore, we have excluded studies prior to 1960 (in order to maintain relevance to current antimicrobial practice) and have not reviewed the data on use of AP to prevent infective endocarditis in animal models, where some evidence suggests that single dose amoxicillin prophylaxis is effective in preventing streptococcal infective endocarditis.

In total, we identified 10 studies assessing the effect of national and international guideline change on the incidence of infective endocarditis. In all countries where AP is still recommended, there has been no significant change in the overall rate of increase of IE, although in several it is claimed that there has been an increase in the number of streptococcal cases. However, IE rates have increased overall in the only study of total AP cessation from the UK.[19] Although this study was unable to ascertain whether this increase was driven by a rising incidence of streptococcal infective endocarditis, a further study is underway to identify the microbiological aetiology of these additional cases. These studies are intrinsically at high risk of methodological bias (as determined by the STROBE criteria) due to their observational study design and cannot fully account for confounding variables. Studies relying solely on discharge coding may not adequately account for re-admissions or re-coding of historical diagnoses. Pant *et al* included secondary diagnoses of infective endocarditis in their analysis, leading to higher estimates than other studies.[17] Several smaller population studies with validated diagnoses have

1
2
3 provided lower estimates for the incidence of infective endocarditis of fewer than 5
4 cases per 100,000 per year.[7, 34, 35] Finally, even the larger time trend studies may
5 remain underpowered to detect a significant change in infective endocarditis
6 incidence given the limited duration of follow-up. In particular, detection of any small
7 change in incidence in studies of relative AP restriction requires a large population or
8 prolonged duration of follow-up.
9

10
11
12
13 We identified 5 observational studies assessing the efficacy of AP. These were
14 retrospective, of poor methodological quality, varying design (4 case control, 1
15 retrospective cohort) and small sample size. Accordingly, they are at high risk of
16 methodological bias and conclusions should be drawn with caution. With this major
17 caveat, our meta-analysis of 3 observational studies did not show a statistically
18 significant difference in exposure to AP in cases (patients with infective endocarditis
19 after dental procedures) compared to controls. There was a trend towards a
20 protective effect of AP, however, and the lack of statistical significance may reflect
21 the small sample sizes in the primary studies. Furthermore, most of the patients in
22 these studies did not have replacement valves or other high-risk pathology so would
23 not have been considered for AP even according to US or European guidelines. Two
24 studies (Duval; Horstkotte) not included in our meta-analysis examined the protective
25 effect of AP in patients with replacement valves. Both suggested a protective effect
26 from AP. In the study by Horstkotte *et al*, 229 patients with replacement heart valves
27 were followed after 287 diagnostic interventions (including some dental) requiring
28 AP.[21] A group of 304 patients who had undergone invasive procedures without AP
29 was used for comparison. Six cases of infective endocarditis occurred in the group
30 with no AP, compared to 0 in the AP group. In a population study by Duval *et al*,
31 approximately 14 times more IE occurred after unprotected than protected dental
32 procedures in people with replacement valves.[36] The study by Horstkotte *et al* was
33 excluded because the number of dental procedures was not stated and the study by
34 Duval *et al* was excluded due to use of extrapolated rather than absolute numbers.
35
36
37
38
39
40
41
42
43
44
45
46
47

48
49 We identified 21 trials assessing the efficacy of AP in reducing the incidence of
50 bacteraemia after dental procedures. Overall, AP is effective at reducing the
51 incidence of bacteraemia. Other surrogate measures addressed in some studies
52 include the nature of isolated bacteria, the duration of bacteraemia, and its
53 magnitude. However, the relationship between bacteraemia and infective
54 endocarditis is not straightforward. In particular, the relative importance of
55 bacteraemia following dental extraction remains debated, and low level bacteraemia
56
57
58
59
60

1
2
3 occurs commonly in association with daily activities such as tooth brushing,
4 especially in the setting of periodontal disease.[37] As such, its validity as a
5 surrogate endpoint for infective endocarditis is uncertain.
6
7

8
9 An RCT of AP has been debated for several decades but is unlikely to be performed
10 for several reasons. Using infective endocarditis as the primary outcome, such a trial
11 would require several hundred thousand participants with a prolonged duration of
12 recruitment and follow-up. In addition, there may be a lack of equipoise in an RCT,
13 given that the standard of care for high-risk individuals (based on ESC and ACC/AHA
14 guidelines) is to give AP. The size, scale and cost of a government-sponsored trial
15 has been deemed unacceptable to national funding bodies in both the UK and
16 USA.[38]
17
18
19
20
21

22 There is general acceptance that the majority of cases of infective endocarditis
23 caused by oral bacterial species are likely to result from frequent bacteraemia arising
24 from routine daily activities, but this does not exclude the possibility that some cases
25 result from infrequent invasive dental procedures.[13] The focus of clinical research
26 on IE prevention has therefore shifted in recent years from surrogate bacteraemia
27 studies to those examining the role played by inflammation and ulceration of gingival
28 tissues. A large multicentre case control study assessing the associations between
29 poor oral hygiene, dental disease and infective endocarditis is currently underway
30 and may provide the necessary data to permanently shift the focus away from AP as
31 the best strategy to prevent IE.
32
33
34
35
36
37
38

39 As the debate continues, infective endocarditis is changing. Oral streptococci – the
40 target of AP – account for a falling proportion of cases in developed world series.[7]
41 In the absence of high quality evidence and with significant barriers to an RCT,
42 uncertainty is likely to prevail. For cardiologists and dental practitioners faced with
43 high-risk individuals, AP remains a low risk, inexpensive approach that may have
44 benefit.[39, 40] We have previously described a framework for discussion of the
45 risk:benefit balance for high-risk patient groups based on current ESC guidelines.[41]
46 Despite the low quality and limited evidence base, these guidelines (and their
47 counterpart from the ACC/AHA) advising AP for patients at highest risk provide a
48 pragmatic and justified approach.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Figure Legends**
4
5

6 **Figure 1. PRISMA flow chart**
7
8

9 **Figure 2. Annual incidence of infective endocarditis reported in time-trend**
10 **analyses.** The data for annual incidence or prevalence was reported in three
11 studies[17, 18, 27] and obtained from the authors for two studies.[19, 20] The
12 incidence of viridans streptococcal infective endocarditis in DeSimone *et al* was 0 in
13 2009 and 2011. The incidence values for Pant *et al* [17] are higher than other studies
14 due to the inclusion of infective endocarditis as both a primary and secondary
15 diagnosis (included solely as a primary diagnosis in the other studies).
16
17
18
19
20

21 **Figure 3. Meta-analysis of case-control studies testing the association between**
22 **antibiotic prophylaxis (AP) and infective endocarditis after dental procedures.**
23

24 In these studies, cases are patients with infective endocarditis and controls are
25 matched patients at risk (Table 2). The number of 'events' is the use of AP in each
26 group as a proportion of the total number of procedures. Overall, the odds ratio of AP
27 comparing patients with infective endocarditis to those without is 0.59 (95%
28 confidence interval 0.27-1.30, $p = 0.14$), suggesting no statistically significant
29 difference in AP exposure between cases and controls. Abbreviations: AP antibiotic
30 prophylaxis; IE infective endocarditis.
31
32
33
34
35
36

37 **Figure 4. Meta-analysis of trials of antibiotic prophylaxis for prevention of**
38 **bacteraemia after dental procedures.** Where an individual study tested multiple
39 antibiotic regimes, these are represented as (a), (b) etc. and compared against the
40 control/placebo arm. Details of the dental procedure and antibiotic regimen are
41 shown in Supplementary Table 5. Overall, use of AP was associated with a risk ratio
42 for bacteraemia of 0.53 (95% CI 0.49-0.57, $p < 0.01$, $I^2 = 90\%$). Abbreviations: AP
43 antibiotic prophylaxis
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Competing interests

M.J.D. was a non-voting member of the NICE panel that reviewed guidance concerning the use of antibiotic prophylaxis to prevent infective endocarditis in 2015.

Acknowledgements

None

References

- 1 Cahill TJ, Prendergast BD. Infective endocarditis. *Lancet* 2016;**387**:882-93.
- 2 Chambers J, Sandoe J, Ray S, et al. The infective endocarditis team: recommendations from an international working group. *Heart* 2013;**100**:524-7.
- 3 Durack DT. Prophylaxis of Infective endocarditis. In: Bennett JE, Dolin R, Blaser MJ, eds. *Principles and Practice of Infectious Diseases*: Elsevier Health Sciences 2014:1058.
- 4 Cahill TJ, Prendergast BD. Current controversies in infective endocarditis. *F1000Research* 2016:doi: 10.12688/f1000research.6949.1.
- 5 Horder TJ. Infective endocarditis: with an analysis of 150 cases and with special reference to the chronic form of the disease. *QJM* 1909;**os2**:289-324.
- 6 Murdoch DR, Corey G, Hoen B, et al. Clinical presentation, etiology, and outcome of infective endocarditis in the 21st century: The international collaboration on endocarditis—prospective cohort study. *Archives of Internal Medicine* 2009;**169**:463-73.
- 7 Selton-Suty C, Célarde M, Le Moing V, et al. Preeminence of *Staphylococcus aureus* in Infective Endocarditis: A 1-Year Population-Based Survey. *Clinical Infectious Diseases* 2012;**54**:1230-9.
- 8 Delahaye F, M'Hammedi A, Guerpillon B, et al. Systematic Search for Present and Potential Portals of Entry for Infective Endocarditis. *Journal of the American College of Cardiology* 2016;**67**:151-8.
- 9 Habib G, Hoen B, Tornos P, et al. Guidelines on the prevention, diagnosis, and treatment of infective endocarditis (new version 2009): the Task Force on the Prevention, Diagnosis, and Treatment of Infective Endocarditis of the European Society of Cardiology (ESC). Endorsed by the European Society of Clinical Microbiology and Infectious Diseases (ESCMID) and the International Society of Chemotherapy (ISC) for Infection and Cancer. *European Heart Journal*; **30**:2369-413.
- 10 Wilson W, Taubert KA, Gewitz M, et al. Prevention of Infective Endocarditis: Guidelines From the American Heart Association: A Guideline From the American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee, Council on Cardiovascular Disease in the Young, and the Council on Clinical Cardiology, Council on Cardiovascular Surgery and Anesthesia, and the Quality of Care and Outcomes Research Interdisciplinary Working Group. *Circulation* 2007;**116**:1736-54.
- 11 Richey R, Wray D, Stokes T. Prophylaxis against infective endocarditis: summary of NICE guidance. *BMJ* 2008;**336**:770-1.
- 12 Thornhill MH, Dayer M, Lockhart PB, et al. A change in the NICE guidelines on antibiotic prophylaxis. *Br Dent J* 2016;**221**:112-4.
- 13 Roberts JG. Dentists Are Innocent! ``Everyday" Bacteremia Is the Real Culprit: A Review and Assessment of the Evidence That Dental Surgical Procedures Are a Principal Cause of Bacterial Endocarditis in Children. *Pediatric Cardiology* 1999;**20**:317-25.

- 1
2
3 14 Higgins JPT, Altman DG, Gøtzsche PC, et al. The Cochrane Collaboration's
4 tool for assessing risk of bias in randomised trials. *BMJ* 2011;**343**.
5
6 15 STROBE Statement. Strengthening the reporting of observational studies in
7 epidemiology. <http://www.strobe-statement.org/>.
8
9 16 Higgins JPT, Thompson SG, Deeks JJ, et al. Measuring inconsistency in
10 meta-analyses. *BMJ* 2003;**327**:557.
11
12 17 Pant S, Patel NJ, Deshmukh A, et al. Trends in Infective Endocarditis
13 Incidence, Microbiology, and Valve Replacement in the United States From 2000 to
14 2011. *Journal of the American College of Cardiology* 2015;**65**:2070-6.
15
16 18 van den Brink FS, Swaans MJ, Hoogendijk MG, et al. Increased Incidence of
17 Infective Endocarditis after the 2009 European Society of Cardiology Guideline
18 Update: A Nation-wide Study in the Netherlands. *European Heart Journal - Quality of
19 Care and Clinical Outcomes* 2016.
20
21 19 Dayer MJ, Jones S, Prendergast B, et al. Incidence of infective endocarditis
22 in England, 2000–13: a secular trend, interrupted time-series analysis. *The
23 Lancet*;**385**:1219-28.
24
25 20 DeSimone DC, Tleyjeh IM, Correa de Sa DD, et al. Incidence of Infective
26 Endocarditis Due to Viridans Group Streptococci Before and After the 2007 American
27 Heart Association's Prevention Guidelines. *Mayo Clinic Proceedings* 2015;**90**:874-81.
28
29 21 Horstkotte D, Rosin H, Friedrichs W, et al. Contribution for choosing the
30 optimal prophylaxis of bacterial endocarditis. *European Heart Journal* 1987;**8**:379-81.
31
32 22 Bates KE, Hall M, Shah SS, et al. Trends in infective endocarditis
33 hospitalisations at United States children's hospitals from 2003 to 2014: impact of the
34 2007 American Heart Association antibiotic prophylaxis guidelines. *Cardiology in the
35 Young* 2016;**15**:1-5.
36
37 23 Bikdeli B, Wang Y, Kim N, et al. Trends in Hospitalization Rates and
38 Outcomes of Endocarditis Among Medicare Beneficiaries. *Journal of the American
39 College of Cardiology* 2013;**62**:2217-26.
40
41 24 Thornhill MH, Dayer MJ, Forde JM, et al. Impact of the NICE guideline
42 recommending cessation of antibiotic prophylaxis for prevention of infective
43 endocarditis: before and after study. *Bmj* 2011;**342**:d2392.
44
45 25 DeSimone DC, Tleyjeh IM, Correa de Sa DD, et al. Incidence of Infective
46 Endocarditis Caused by Viridans Group Streptococci Before and After Publication of
47 the 2007 American Heart Association's Endocarditis Prevention Guidelines.
48 *Circulation* 2012;**126**:60-4.
49
50 26 Duval X, Delahaye F, Alla F, et al. Temporal Trends in Infective Endocarditis
51 in the Context of Prophylaxis Guideline Modifications: Three Successive Population-
52 Based Surveys. *Journal of the American College of Cardiology* 2012;**59**:1968-76.
53
54 27 Keller K, von Bardleben RS, Ostad MA, et al. Temporal Trends in the
55 Prevalence of Infective Endocarditis in Germany Between 2005 and 2014. *American
56 Journal of Cardiology* 2016.
57
58
59
60

- 1
2
3 28 Mackie AS, Liu W, Savu A, et al. Infective Endocarditis Hospitalizations
4 Before and After the 2007 American Heart Association Prophylaxis Guidelines.
5 *Canadian Journal of Cardiology* 2007;**32**:942-8.
6
- 7 29 Salam AMF, Albinali H, Singh R, et al. Incidence of infective endocarditis
8 before and after the 2007 endocarditis prevention guidelines: a
9 population-based study from Qatar (2002-2012). *European Heart Journal*
10 2014;**35**:908.
11
- 12 30 Imperiale TF, Horwitz RI. Does prophylaxis prevent postdental infective
13 endocarditis? A controlled evaluation of protective efficacy. *The American Journal of*
14 *Medicine* 1990;**88**:131-6.
15
- 16 31 Lacassin F, Hoen B, Leport C, et al. Procedures associated with infective
17 endocarditis in adults. *European Heart Journal* 1995;**16**:1968-74.
18
- 19 32 Strom BL, Abrutyn E, Berlin JA, et al. Dental and Cardiac Risk Factors for
20 Infective Endocarditis: A Population-Based, Case-Control Study. *Annals of Internal*
21 *Medicine* 1998;**129**:761-9.
22
- 23 33 van der Meer JTM, Michel MF, Valkenburg HA, et al. Efficacy of antibiotic
24 prophylaxis for prevention of native-valve endocarditis. *The Lancet* 1992;**339**:135-9.
25
- 26 34 Tleyjeh IM, Abdel-Latif A, Rahbi H, et al. A Systematic Review of Population-
27 Based Studies of Infective Endocarditis. *Chest* 2007;**132**:1025-35.
28
- 29 35 Sy RW, Kritharides L. Health care exposure and age in infective endocarditis:
30 results of a contemporary population-based profile of 1536 patients in Australia.
31 *European Heart Journal* 2010;**31**:1890.
32
- 33 36 Duval X, Alla F, Hoen B, et al. Estimated Risk of Endocarditis in Adults with
34 Predisposing Cardiac Conditions Undergoing Dental Procedures With or Without
35 Antibiotic Prophylaxis. *Clinical Infectious Diseases* 2006;**42**:e102-e7.
36
- 37 37 Lockhart PB, Brennan MT, Sasser HC, et al. Bacteremia Associated With
38 Toothbrushing and Dental Extraction. *Circulation* 2008;**117**:3118-25.
39
- 40 38 Thornhill MH, Lockhart PB, Prendergast B, et al. NICE and antibiotic
41 prophylaxis to prevent endocarditis. *Br Dent J* 2015;**218**:619-21.
42
- 43 39 Franklin M, Wailoo A, Dayer MJ, et al. The cost-effectiveness of antibiotic
44 prophylaxis for patients at risk of infective endocarditis. *Circulation*; **In press**.
45
- 46 40 Thornhill MH, Dayer MJ, Prendergast B, et al. Incidence and nature of
47 adverse reactions to antibiotics used as endocarditis prophylaxis. *Journal of*
48 *Antimicrobial Chemotherapy* 2015;**70**:2382-8.
49
- 50 41 Thornhill MH, Dayer M, Lockhart PB, et al. Guidelines on prophylaxis to
51 prevent infective endocarditis. *Br Dent J* 2016;**220**:51-6.
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1 – PRISMA flow chart




Figure 1. PRISMA flow chart
Figure 1
155x154mm (300 x 300 DPI)




Figure 2. Annual incidence of infective endocarditis reported in time-trend analyses. The data for annual incidence or prevalence was reported in three studies[17, 18, 27] and obtained from the authors for two studies.[19, 20] The incidence of viridans streptococcal infective endocarditis in DeSimone et al was 0 in 2009 and 2011. The incidence values for Pant et al [17] are higher than other studies due to the inclusion of infective endocarditis as both a primary and secondary diagnosis (included solely as a primary diagnosis in the other studies).

Figure 2
31x19mm (600 x 600 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3. Meta-analysis of case-control studies testing the association between antibiotic prophylaxis (AP) and infective endocarditis after dental procedures. In these studies, cases are patients with infective endocarditis and controls are matched patients at risk (Table 2). The number of 'events' is the use of AP in each group as a proportion of the total number of procedures. Overall, the odds ratio of AP comparing patients with infective endocarditis to those without is 0.59 (95% confidence interval 0.27-1.30, p = 0.14), suggesting no statistically significant difference in AP exposure between cases and controls. Abbreviations: AP antibiotic prophylaxis; IE infective endocarditis.

344x62mm (72 x 72 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 4. Meta-analysis of trials of antibiotic prophylaxis for prevention of bacteraemia after dental procedures. Where an individual study tested multiple antibiotic regimens, these are represented as (a), (b) etc. and compared against the control/placebo arm. Details of the dental procedure and antibiotic regimen are shown in Supplementary Table 5. Overall, use of AP was associated with a risk ratio for bacteraemia of 0.53 (95% CI 0.49-0.57, $p < 0.01$, $I^2 = 90\%$). Abbreviations: AP antibiotic prophylaxis

312x242mm (72 x 72 DPI)

SUPPLEMENTARY DATA

Supplementary Appendix 1: Search strategy

MEDLINE

# ▲	Searches
1	exp Endocarditis, Bacterial/ or Endocarditis/
2	((bacter* or infective or inflammatory) adj5 endocarditis) ti,ab.
3	((infect* or inflam*) adj5 endocardium) ti,ab.
4	endocarditis.ti.
5	1 or 2 or 3 or 4
6	exp Anti-Bacterial Agents/
7	(antibiotic? or anti-biotic? or antimicrobial? or anti-microbial? or antibacterial? or anti-bacterial? or anti-infective).ti,ab.
8	(penicillin or amoxicillin or ampicillin or cefazolin or ceftriaxone or cephalixin or clindamycin or azithromycin or clarithromycin or vancomycin).ti,ab.
9	6 or 7 or 8
10	(prophyla* or prevent* or premedicat* or pre-medocat*) ti,ab.
11	9 and 10
12	Antibiotic Prophylaxis/
13	11 or 12
14	5 and 13
15	exp Endocarditis, Bacterial/pc [Prevention & Control]
16	9 and 15
17	(endocarditis and prophyla*).ti.
18	14 or 16 or 17
19	exp animals/ not humans sh.
20	18 not 19

EMBASE

# ▲	Searches
1	exp endocarditis/
2	((bacter* or infective or inflammatory) adj5 endocarditis).ti,ab.
3	((infect* or inflam*) adj5 endocardium) ti,ab.
4	endocarditis.ti.
5	1 or 2 or 3 or 4
6	exp *antibiotic agent/
7	(antibiotic? or anti-biotic? or antimicrobial? or anti-microbial? or antibacterial? or anti-bacterial? or anti-infective).ti,ab.
8	(penicillin or amoxicillin or ampicillin or cefazolin or ceftriaxone or cephalixin or clindamycin or azithromycin or clarithromycin or vancomycin).ti,ab.
9	6 or 7 or 8
10	(prophyla* or prevent* or premedicat* or pre-medocat*) ti,ab.
11	9 and 10
12	Antibiotic Prophylaxis/
13	11 or 12
14	5 and 13
15	(endocarditis and prophyla*).ti.
16	14 or 15
17	(exp animals/ or nonhuman/) not human/

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

18 16 not 17

COCHRANE

ID	Search
#1	endocarditis:ti,ab,kw (Word variations have been searched)
#2	((bacter* or infective or inflammatory) near endocarditis) ti,ab,kw (Word variations have been searched)
#3	abe or sabe ti,ab,kw (Word variations have been searched)
#4	#1 or #2 or #3
#5	MeSH descriptor: [Anti-Bacterial Agents] explode all trees
#6	(antibiotic? or anti-biotic? or antimicrobial? or anti-microbial? or antibacterial? or anti-bacterial? or anti-infective):ti,ab,kw (Word variations have been searched)
#7	penicillin or amoxicillin or ampicillin or cefazolin or ceftriaxone or cephalixin or clindamycin or azithromycin or clarithromycin or vancomycin ti,ab,kw (Word variations have been searched)
#8	#5 or #6 or #7
#9	prophyla* or prevent* or premedicat* or pre-medecat*.ti,ab,kw (Word variations have been searched)
#10	#8 and #9
#11	MeSH descriptor: [Antibiotic Prophylaxis] explode all trees
#12	#10 or #11
#13	#4 and #12
#14	MeSH descriptor: [Endocarditis, Bacterial] explode all trees and with qualifier(s): [Prevention & control - PC]
#15	endocarditis and prophyla* ti (Word variations have been searched)
#16	#13 or #14 or #15

WOK

Set	Results	Save search history and/or create an alertOpen a saved search history
1	29,469	TOPIC: (endocarditis) OR TOPIC: (((infect* or inflam*) NEAR/5 endocardium)) OR TOPIC: (ABE OR SABE)
2	459,418	TS=(antibiotic* or anti-biotic* or antimicrobial* or anti-microbial* or antibacterial* or anti-bacterial* or anti-infective) OR TS=(penicillin or amoxicillin or ampicillin or cefazolin or ceftriaxone or cephalixin or clindamycin or azithromycin or clarithromycin or vancomycin)
3	1,167,210	TOPIC: (prophyla* or prevent* or premedicat* or pre-medecat*)
4	1,506	#3 AND #2 AND #1
5	606	TITLE: (endocarditis AND prophyla*)
6	1,785	#5 OR #4

Supplementary Table 1: Excluded studies

First author	Publication year	PMID/PMCID	Reason excluded
Agha	2005	15951458	Outcome of interest not reported
Aitken	1995	7599015	No control/placebo group
Alexander	2013	Abstract	No population incidence pre/post guideline reported
Al-Karaawi	2001	11119466	Outcome of interest not reported
Al-Omari	2014	24624933	Treatment of infective endocarditis study
Aoki	1996	NA	No abx prophylaxis intervention group
Baddour	2013	NA	Comment/editorial/review
Bain	1985	NA	Comment/editorial/review
Baltch	1988	3365078	No control/placebo group
Bates	2015	Abstract	No population incidence pre/post guideline reported
Bikdeli	2012	Abstract	Dataset/analysis included in later publication
Bolukbasi	2012	22157668	No abx prophylaxis intervention group
Bor	2013	23527296	No population incidence pre/post guideline reported
Bor	1984	6711576	Modelling study/extrapolated data
Brennan	2007	17197406	Dataset/analysis included in later publication
Bublak	2014	25608390	Comment/editorial/review
Bunnell	2011	NA	No control/placebo group
Carretta	1988	3167905	Unobtainable
Casas	1999	10231302	Unobtainable
Cannon	1987	3610906	Outcome of interest not reported
Cecchi	2009	18404005	No control/placebo group
Chen	2015	26512586	Outcome of interest not reported
Cherry	2007	17309589	Study of topical antibiotic/antiseptic
Clark	1969	5192402	Patient population
Connor	1967	5234633	Unobtainable
Cosgrove	2011	Abstract	No population incidence pre/post guideline reported
Couturier	2000	NA	Comment/editorial/review
Crasta	2009	19426179	No abx prophylaxis intervention group
Curran	1976	1075923	Unobtainable
Daly	1997	9153833	No abx prophylaxis intervention group
Daly	2001	11288795	No abx prophylaxis intervention group
Dankert	1982	NA	No abx prophylaxis intervention group
Dankert	1985	NA	Animal study
Darhous	1993	9588119	Unobtainable
Dayer	2010	Abstract	Dataset/analysis included in later publication
Dayer	2014	Abstract	Dataset/analysis included in later publication
Dayer	2011	Abstract	Dataset/analysis included in later publication
DeSimone	2015	26386808	No population incidence pre/post guideline reported
De Aguir	2012	22522720	No abx prophylaxis intervention group
De Leo	1974	4519445	No abx prophylaxis intervention group
Delahaye	2011	Abstract	Dataset/analysis included in later publication
Dell'Atti	2013	Abstract	Outcome of interest not reported
Diz	2013	Abstract	Group sizes not reported
Doshi	2007	Abstract	No population incidence pre/post guideline reported
Durak	1982	6632128	Outcome of interest not reported
Durak	1975	NA	Comment/editorial/review
Duval	2006	16705565	Modelling study/extrapolated data
Edwards	2015	Abstract	No population incidence pre/post guideline reported
Ellervall	2010	20134479	Systematic review
Erichsen	2016	27339641	No population incidence pre/post guideline reported
Esposito	2013	18646101	Systematic review
Feinstein	1961	13698654	Comment/editorial/review
Francioli	1985	3925031	Animal study
Friedlander	2010	20189771	Comment/editorial/review

Garlando	1988	3292185	Patient population
Glenny	2013	241108511	Systematic review
Goker	1992	1308784	Unobtainable
Grattan	2013	26319967	No abx prophylaxis intervention group
Grimard	1987	3680973	Unobtainable
Grimard	1986	NA	No abx prophylaxis intervention group
Guntheroth	1984	6486031	Outcome of interest not reported
Guze	1983	6418063	Animal study
Hakata	2014	NA	No abx prophylaxis intervention group
Harris	1999	10604613	Patient population
Hartzell	2005	15832100	No abx prophylaxis intervention group
Harvey	1961	13712066	Outcome of interest not reported
Hess	1977	NA	Foreign language
Hess	1981	Abstract	No control/placebo group
Hess	1983	6549771	No control/placebo group
Hess	1983	6550789	No control/placebo group
Ho	2015	NA	No population incidence pre/post guideline reported
Kaplan	1971	NA	Comment/editorial/review
Kaplan	1983	6889082	Comment/editorial/review
Kaplan	1983	NA	Unobtainable
Kaneko	1995	8869455	No control/placebo group
Katoh	1992	1300669	No control/placebo group
Kernodle	1993	8494364	Animal study
Kinane	2005	15966875	No abx prophylaxis intervention group
Klug	2007	17724263	Patient population
Kumana	1986	3099944	Outcome of interest not reported
Lador	2012	41122	No control/placebo group
Lee	2007	17761735	Outcome of interest not reported
Lee	2009	NA	Outcome of interest not reported
Lockhart	2009	2770162	No abx prophylaxis intervention group
Longman	1991	1990136	Outcome of interest not reported
Lucas	2000	10769852	Outcome of interest not reported
Oakley	1982	NA	No control/placebo group
Macgregor	1987	NA	Unobtainable
Magelli	1987	NA	Unobtainable
Maharaj	2012	3734757	No abx prophylaxis intervention group
Martin	1964	14244278	Unobtainable
Mazocchi	2007	17954332	No abx prophylaxis intervention group
Messini	1999	10412852	No abx prophylaxis intervention group
Monaco	2009	19531419	Outcome of interest not reported
Mougeot	2015	25758845	Dataset/analysis included in later publication
Murillo	1978	NA	Outcome of interest not reported
Murillo	1979	463947	Outcome of interest not reported
Murphy	2006	16968327	No abx prophylaxis intervention group
Murugiah	2013a	Abstract	No population incidence pre/post guideline reported
Murugiah	2013b	Abstract	No population incidence pre/post guideline reported
Niederrau	1994	7988813	Patient population
Nelson	1965	NA	Patient population
Niwa	2005	1768964	Outcome of interest not reported
Oliver	2004	15106220	Systematic review
Pant	2015b	26338002	Comment/editorial/review
Pasquali	2011	Abstract	No population incidence pre/post guideline reported
Pasquali	2012	22607869	No population incidence pre/post guideline reported
Pasquantonio	2012	23058035	Outcome of interest not reported
Peterson	1976	1253389	No abx prophylaxis intervention group
Pineiro	2010	20701619	Study of topical antibiotic/antiseptic
Piper	2012	Abstract	Outcome of interest not reported
Rahn	1995	NA	Unobtainable

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Rechmann	1989	2639069	Study of topical antibiotic/antiseptic
Rieder	1980	7423172	Patient population
Roberts	1997	8960488	No abx prophylaxis intervention group
Roberts	2002	12572738	No control/placebo group
Roberts	2000	10689771	No abx prophylaxis intervention group
Rogers	2008	18513601	No population incidence pre/post guideline reported
Sasaki	1994	NA	No control/placebo group
Sasaki	1998	NA	No control/placebo group
Sasaki	2001	NA	Foreign language
Santinga	1976	1067349	Outcome of interest not reported
Santinga	1984	NA	No control/placebo group
Sefton	1990	2196261	Duplicate data (Cannell et al)
Schirger	1964	14146015	No control/placebo group
Schlein	1991	2028936	No abx prophylaxis intervention group
Schwartz	2007	17904722	Systematic review
Shanson	1983	6130318	Outcome of interest not reported
Shanson	1984	6334070	Outcome of interest not reported
Sheenchen	1994	8310405	Patient population
Smith	1989	2732123	Outcome of interest not reported
Tempelhof	2012	NA	Systematic review
Thompson	1979	NA	Outcome of interest not reported
Thornhill	2011	21540258	Dataset/analysis included in later publication
Thornhill	2015	25925595	Outcome of interest not reported
Tozer	1966	4159601	Outcome of interest not reported
Tuback	2002	NA	Comment/editorial/review
Tzuckert	1978	285049	Unobtainable
Wong	2011	NA	No population incidence pre/post guideline reported
Yoshimura	1985	2939227	Unobtainable
Yurtman	2010	NA	Patient population
Zhang	2013	23137266	No abx prophylaxis intervention group

Supplementary Table 3: Risk of bias summary – observational studies

	Horstkotte et al 1987[10]	Imperiale et al 1990[11]	Lacassin et al 1995[12]	Strom et al 1998	Van der Meer et al 1992[13]
Definition of cases	Catchment area and time period not specified. Cases defined as '229 patients in whom 287 diagnostic and therapeutic procedures were performed using a prophylactic antibiotic regime correctly administered.	Catchment area and time period specified. Clearly defined inclusion and exclusion criteria. Presence of infective endocarditis determined using the modified Von Reyn criteria.	Catchment area and time period specified. Clearly defined inclusion and exclusion criteria. Quote: 'Definite and probably IE was defined according to Von Reyn's criteria revised with the following modifications to include echocardiographic and macroscopic findings for definite and probably cases. Definite endocarditis was defined on macroscopy or microbiological findings on operation or necropsy; probably endocarditis was defined as (1) persistently positive blood cultures with underlying heart disease plus echocardiographic vegetation or with vascular phenomena plus echocardiographic vegetation or with vascular phenomena plus echocardiographic vegetation or (2) negative or intermittently positive blood cultures with fever, underlying heart disease, vascular phenomena, and echocardiographic vegetation'.	Catchment area and time period specified. Inclusion and exclusion criteria clearly defined. Infective endocarditis not determined using diagnostic criteria. Quote: 'These experts used their own global clinical judgement to classify potential cases as definite, probably or possible cases'.	Catchment area and time period specified. Clearly defined inclusion and exclusion criteria. Presence of infective endocarditis was determined by the Von Reyn criteria.
Definition of controls	Catchment area and time period not specified. 'The second patient group consisted of 304 out of 1898 patients questioned in our outpatient clinic, who gave reliable information that they had undergone one of the procedures we regard as requiring endocarditis prophylaxis, without having received any antibiotic regime'.	Time period specified and catchment area specified and same as cases. Inclusion clearly defined. Exclusion criteria clearly defined and same as cases.	Defined, but different population as cases. Quote: 'Controls were subjects without infective endocarditis who satisfied the same exclusion criteria as the cases. There were all recruited randomly from wards either during a consultation for echo or during hospitalisation in the same period of observations as cases'. 'Each case was matched to one control as regards sex, age and group of underlying cardiac conditions'.	Catchment area and time period not specified. Inclusion poorly defined. Exclusion criteria for controls not defined. Recruited using a modification of the Waksberg random-digit dialling method	Catchment area specified, differing from cases. Time period not specified. Inclusion and exclusion criteria clearly described
Characteristics of each group described	No	Yes: Age, gender, cardiac lesion, history or rheumatic fever, murmur duration, frequency of dental visits, dental clearing at zero time visit, use of diuretics, use of digitalis, penicillin allergy	Yes: Age, sex, valve disease, duration of previous cardiac disease, known cardiac disease	Only age	Yes: Age, procedures, interval between procedure and onset of symptoms
Groups recruited at common stage, in the same manner	Not specified	Cases and controls were recruited during the same time period and from the same catchment area. Both cases and controls were recruited in the same manner, in the form of a letter.	Cases and controls were recruited during the same time period, but catchment area and manner of recruitment not specified	Catchment area and time period for recruitment of controls not specified. Method of recruitment not specified.	Cases and controls were recruited from different catchment areas. Time period of recruitment of controls not specified. Manner of recruitment differed between cases and controls: cases were selective consecutively and interviewed in person; controls were selected randomly, sent a letter and then a telephone call
Sampling strategy	Not specified	Cases: preliminary selected by review of medical and dental records, and completion of telephone interview. Controls: all patients who underwent echocardiographic evaluation during the period covered by the study.	Not specified	Catchment area and time period for recruitment of controls not specified. Sampling strategy varied: controls were randomly recruited from the community; controls were recruited from all in patients with a putative diagnosis of endocarditis.	Cases: Quote 'All patients who were consecutively admitted to hospital in the Netherlands and who were suspected of having bacterial endocarditis'. Comment: defined catchment area over a defined period of time. Controls: sampling strategy not specified.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Is the group representative of the population of interest	In part, but also included procedures other than dental, such as urological and gynaecological	Yes	Yes – only dental subgroup analysed	Yes	Overall yes, although < 10% of procedures were non-dental
Duration of follow-up	No follow-up	No-follow up	No follow-up	No follow-up	No follow-up
Outcome assessment Objective or independently adjudicated?	Outcome: Prosthetic valve endocarditis. Method or criteria for diagnosis of endocarditis is not described.	Telephone interview, use of a standard structured questionnaire not described. Specific purpose of research was not revealed to the interview subjects. Use of independent adjudicator not described.	Cases and controls were interviewed using the same questionnaire. Use of an independent adjudicator, or blinding of the participants to the study purpose is not described	Structured telephone interview where cases and controls completed the same structured questionnaire. Use of an independent adjudicator or blinding of the participants to the study purpose is not described	Telephone interview where cases and controls completed the same structured questionnaire. Use of an independent adjudicator, or blinding of the participants to the study purpose is not described
Overall risk of bias	High	High	High	High	High

Supplementary Table 4 – Risk of bias summary – bacteraemia trials

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Asi 2010	?	?	?	?	+	?	+
Baltch 1982	-	-	?	?	+	+	-
Baltch 1982 (2)	-	-	?	?	+	+	-
Cannell 1991	?	-	?	?	+	+	?
Coulter 1990	-	-	?	?	+	+	-
Diz Dios 2006	+	?	?	?	+	+	?
Duvall 2013	+	?	+	?	?	+	?
Hall 1993	?	?	?	?	?	+	?
Hall 1996	?	?	?	?	?	?	?
Head 1984	-	+	+	+	+	?	?
Khairat 1966	+	?	?	?	+	+	?
Limeres Posse 2016	+	?	?	+	?	?	+
Lockhart 2004	+	+	+	?	+	+	?
Lockhart 2008	+	+	+	+	+	+	?
Maharaj 2012	+	?	?	?	?	+	?
Maskell 1986	?	?	?	?	?	?	?
Roberts 1987	+	?	?	?	?	?	?
Shanson 1978	?	?	-	?	?	?	?
Shanson 1985	?	?	?	?	?	?	?
Shanson 1987	+	-	-	?	?	?	?
Vergis 2001	?	?	?	?	?	?	?

Supplementary Table 4. Risk of bias assessment for trials of antibiotic prophylaxis.

All trials were assessed using the Cochrane risk of bias tool. Symbols:  low risk of bias,  high risk of bias,  unclear risk of bias

Supplementary Table 5: Data extraction – bacteraemia trials

First author, year	PMID	Study design	Population (children, adults, mixed)	Intervention	Comparison arm	Study method	Total no of patients - abx prophylaxis arm 1	Total number of patients - abx prophylaxis arm 2	Total number of patients - abx prophylaxis arm 3	Total number of patients - abx prophylaxis arm 4	Total number of patients – control/placebo arm	Incidence of any bacteraemia in post procedure cultures - prophylaxis group 1	Incidence of any bacteraemia in post procedure cultures - prophylaxis group 2	Incidence of any bacteraemia in post procedure cultures - prophylaxis group 3	Incidence of any bacteraemia in post procedure cultures - prophylaxis group 4	Incidence of any bacteraemia in post procedure cultures - placebo group	Secondary outcome level/duration bacteraemia	Secondary outcome incidence of strep bacteraemia
Baltch, 1982[15]	7148652	TNR Paper USA	Adults (age not specified) 28 with known cardiac abnormality (prophylaxis group) 28 without cardiac abnormality (control group)	Dental cleaning	Control group	Blood cultures taken pre- and 5 and 30 mins post-procedure	28	NA	NA	NA	28	3/28 (10.7%) at 5 minutes 4/26 (15.4%) at 30 minutes	NA	NA	NA	17/28 (60.7%) at 5 minutes 7/28 (25%) at 30 minutes	Not reported	Control group 4/28 (14.3%) at 5 mins, 1/28 (3.6%) at 30 mins. Abx group 1/28 (3.6%) at 5 mins, 1/26 (3.9%) at 30 mins
Asi, 2010[14]	20922074	TNR Paper India	Adults (aged 35-45 years)	Periodontal flap procedure (under LA)	Each patient served as own control - separate quadrants	Blood cultures taken during procedure	30	NA	NA	NA	30	4/30 (13.3%)	NA	NA	NA	14/30 (46.6%)	Not reported	Control group 4/30 (13.3%). Abx group 1/30 (3.3%)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Baltch, 1982b[16]	7044125	TNR	Paper	USA	Adults (range 19 – 83) undergoing dental extraction, 33/128 with known valvular heart disease	Dental extraction (LA)	Aqueous penicillin G 2 million units IV 30 minutes before the procedure and every 4 hours for 72 hours thereafter	Penicillin V orally 0.5-1g 30-45 minutes prior to the procedure	NA	NA	Control group (29 under local anaesthesia used to match antibiotic groups)	Blood cultures pre- and 5 and 30 mins post-procedure	33	36	NA	NA	29	11/32 (34.4%) at 5 minutes 4/33 (12.1%) at 30 minutes	17/36 (47.2%) at 5 minutes 5/36 (13.9%) at 30 minutes	NA	NA	17/29 (58.6%) at 5 minutes 8/29 (26.7%) at 30 minutes	Number of isolates per positive blood culture at 5 minutes IV penicillin 1.6, PO penicillin 2.5, control group 4.1	Proportion with streptococcal bacteraemia at 5 mins IV penicillin 1/32 (3.1%), PO penicillin 1/36 (2.8%), control group 17/30 (56.7%)
Cannell, 1991[17]	1832916	RCT	Paper	UK	Adults (aged 18-65 years)	Dental extraction (anaesthetic not specified)	Erythromycin 1.5g PO, 1-1.5h pre-procedure	Josamycin base 1.5g, 1-1.5h pre-procedure	NA	NA	Placebo group	Blood cultures taken, timing not specified	20	20	NA	NA	20	12/20 (60%)	14/20 (70%)	NA	NA	13/20 (65%)	Not reported	Not reported
Coulter, 1990[18]		TNR	Paper	UK	Children (aged 2-13 years) 26 with known cardiac abnormality (prophylaxis group) 32 with no cardiac abnormality (control group)	Dental extraction (under GA)	Penicillin IM (dose and timing not specified)	Amoxicillin PO (dose and timing not specified)	Amoxicillin IV (dose and timing not specified)	Erythromycin IV (dose and timing not specified)	Control group	Blood cultures taken 1-2 minutes post-extraction	8	8	6	4	32	Overall 9/26 (35%)	NA	NA	NA	20/32 (63%)	Not reported	Number of patients with streptococcal bacteraemia in each group 3/26 with AP 12/32 without AP
Diz Dios, 2006[19]		RCT	Paper	Spain	Adults (aged >=18 years) with learning difficulties	Dental extraction (under GA)	Amoxicillin 2g PO, 1-2h pre-procedure	Clindamycin 600mg PO, 1-2h pre-procedure	Moxifloxacin 400mg PO, 1-2h pre-procedure	NA	Control group	Blood cultures taken at 0s, 30s, 15min and 1h post-extraction	56	202	58	NA	53	46.4% (n=26) at 30s, 10.7% at 15mins, 3.7% at 1h	85.1% (n=46) at 30s, 70.4% at 15mins, 22.2% at 1h	56.9% at 30s (n=33), 24.1% at 15mins, 7.1% at 1h	NA	96.2% (n=51) at 30s, 64.2% at 15mins, 20% at 1h	Not reported	% of cultures +ve for streptococci Control group 63.1%. Amox group 44.4%. Clindamycin group 58.5%. Moxifloxacin 67.7%
Duvall, 2013[20]	23453612	RCT	Paper	USA	Adults (aged 18-29 years)	Dental extraction (under LA)	Amoxicillin 2g PO, 1h pre-procedure	Chlorhexidine rinse. 1h pre-procedure	NA	NA	Placebo group	Blood cultures taken before and 1.5 minutes after 2nd extraction, 1.5 mins after 4th extraction, then at 10 mins after 4th extraction	10	10	NA	NA	10	4/10 (40%)	6/10 (60%)	NA	NA	5/10 (50%)	Magnitude of bacteraemia (SD) placebo group 3.61+/-7.09 CFU/ml amoxicillin 0.63+/-1.33 cfu/ml chlorhex 2.76+/-4.28 cfu/ml	Number of alpha/hemolytic/VGS isolates placebo 5, chlorhexidine 2, amoxicillin 1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Hall, 1993[21]	8399865	RCT	Paper	Sweden	Adults (aged 23-74 years)	Dental extraction (under LA)	Penicillin V 1g PO x2, 1h pre-procedure	Amoxicillin 750mg PO x4, 1h pre-procedure	NA	NA	Placebo group	Blood cultures taken during procedure and 10 minutes post-procedure	20	20	NA	NA	20	Overall 90% (n=18). 90% during procedure, 70% at 10 mins	Overall 85% (n=17). 85% during procedure, 60% at 10 mins	NA	NA	Overall 95% (n=19). 90% during surgery, 80% at 10 mins	Median cfu per ml reported During procedure Pen V 0.66, Amox 1.08, Placebo 0.84 10mins post-procedure Pen V 0.36, Amox 0.24, Placebo 0.36	VGS isolated during procedure Pen V 70%, Amox 55%, Placebo 70% VGS 10mins post-procedure Pen V 25%, Amox 30%, Placebo 40%
Hall, 1996[22]	8894572	RCT	Paper	Sweden	Adults (aged 21-66 years)	Dental extraction (anaesthetic not specified)	Cefaclor 1g PO, 1h pre-procedure	NA	NA	NA	Placebo group	Blood cultures taken pre-, during and 10 minutes post-procedure	19	NA	NA	NA	20	79% (n=15) during procedure, 53% at 10 mins	NA	NA	NA	85% (n=17) during, 47% at 10 mins	See bacteraemia at 10 minutes outcome	Control group 50% during procedure, 30% at 10 mins post-procedure. Cefaclor group 79% during procedure, 26% at 10 mins post-procedure
Head, 1984[23]	6435046	RCT	Paper	Canada	Adults (aged 16-60 years)	Dental extraction (under LA)	Penicillin V 2g PO, 1h pre-procedure	Metronidazole 2g PO, 1h pre-procedure	NA	NA	Placebo group	Blood cultures taken 1 minute post-extraction	25	25	NA	NA	25	20%	52%	NA	NA	84%	Pour plates' were all negative - bacteraemias < 1 microorganism per ml of blood	Reported - need to check
Khairat, 1966[24]	5928605	RCT	Paper	Canada	Not specified	Dental extraction (under LA)	Erythromycin PO, varying dose from 250mg-1000mg at varying time 90-240mins pre-procedure	Tetracycline 275mg IV, 3 mins pre-procedure	NA	NA	Control group	Blood cultures taken, timing not specified	42	100	NA	NA	100	16/42 (38%)	3/100 (3%)	NA	NA	64/100 (64%)	Control group 44 (44%) Pyrrolidino group 2 (2%)	
Limeres Posse, 2016[25]	27029851	RCT	Paper	Spain	Adults (aged >=18 years)	Dental extraction (under GA)	Amoxicillin/clavulanate 1000/200mg IV, after induction	Amoxicillin 2g PO, 1-2h before induction	Clindamycin 600mg PO, 1-2h before induction	Azithromycin 500mg PO, 1-2h before induction	Control group - unmedicated	Blood cultures pre-procedure, 30s post-extraction, and 15 and 60 mins post-procedure	52	50	52	57	55	0/52 (0%)	25/50 (50%)	45/52 (87%)	46/57 (81%)	53/55 (96%)	Results also reported at 15 mins and 1h	Percentage of positive cultures that were VGS Control-50% Amoxicillin/Clavulanate-0% Amoxicillin-31% Clindamycin 56% Azithromycin 53%

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Lockhart, 2004[26]	15173031	RCT	Paper	USA	Children (mean age 3.5 years)	Dental extraction (under GA)	Amoxicillin 50mg/kg PO, 1h pre-procedure	NA	NA	NA	Placebo group	Blood cultures taken pre-, during and 15, 30 and 45 mins post-procedure	49	NA	NA	NA	51	16/49 (33%)	NA	NA	NA	43/51 (84%)	Duration reported at 15 minutes 12% vs 1%, at 30 mins 9% vs 0%, at 45 mins 5% vs 0% - all prophylaxis vs placebo	Numbers of positive cultures for VGS. Control group 45% (n=57). Amoxicillin group 33% (n=8)
Lockhart, 2008[27]	18541739	RCT	Paper	USA	Not specified, (mean age 40 years)	Dental extraction (under LA)	Amoxicillin (dose, route and timing not specified)	Tooth brushing	NA	NA	Placebo group	Blood cultures taken pre-, during, and 20, 40 and 60 mins post-procedure	90	89	NA	NA	89	56%	32%	NA	NA	80%	Duration is reported. All analysed samples were below the detection threshold of 104 CFU per millilitre of blood	Numbers of positive cultures for VGS. Control group 106/151 (70%). Amoxicillin group 23/47 (49%)
Maharaj, 2012[28]	23108516	RCT	Paper	South Africa	Adults (aged 18-66 years), without underlying valve disease or prosthetic valve	Dental extraction (anaesthetic not specified)	Amoxicillin 3g PO, 1h pre-procedure	Clindamycin 600mg PO, 1h pre-procedure	Chlorhexidine rinse, 1h pre-procedure	NA	Control group	Blood cultures taken, timing not specified	40	40	40	NA	40	3/40 (7.5%)	8/40 (20%)	16/40 (40%)	NA	14 (35%)	Not reported	Number of patients with positive VGS cultures control group 5 chlorhexidine 5 amoxicillin 0 clindamycin 0
Maskell, 1986[29]	2941404	TNR	Paper	UK	Not specified	Dental extraction (anaesthetic not specified)	Teicoplanin 200mg IM, 1h pre-procedure	Amoxicillin 3g PO, 1h pre-procedure	NA	NA	Control group	Blood cultures taken during procedure	10	10	NA	NA	10	6/10 (60%)	4/10 (40%)	NA	NA	10 /10 (100%)	NA	Control group 9/10 (90% of patients had oral streptococci isolated 5/10 in teicoplanin group 3/10 in amoxicillin group
Roberts, 1987[30]	NA	Randomised trial	Paper	UK	Children < 16 years	Dental extraction	Amoxicillin 50mg/kg PO, 2h pre-procedure	NA	NA	NA	Control group	Blood cultures pre procedure, post intubation and 2 minutes post extraction	47	NA	NA	NA	47	1/47 (2.13%)	NA	NA	NA	18/47 (38.3%)	NA	Control group - 14/47 patients positive cultures for strep (VGS, non-haemolytic or anaerobic strep) Amoxicillin group - 0/47 positive culture due to strep

Shanson, 1978[31]	99423	RCT	Paper	UK	Adults (age not specified)	Dental extraction (anaesthetic not specified)	Penicillin V 2g PO, 1h pre-procedure	Amoxicillin 2g PO, 1h pre-procedure	NA	NA	Control group - unmedicated	Blood cultures taken 2 mins post-extraction	40	40	NA	NA	40	5/40 (12%)	2/40 (5%)	NA	NA	16/40 (40%)		
Shanson, 1985[32]	3882657	TNR	Paper	UK	Adults (aged 18-71)	Dental extraction (LA)	Erythromycin stearate PO, 1.5g given 1h pre-procedure	NA	NA	NA	Placebo group	Blood cultures 1-2 minutes following extraction	40	NA	NA	NA	42	6/40 (15%)*	NA	NA	NA	18/42 (43%)*	Serial dilutions reported	See primary outcome for this study (only streptococcal bacteraemia reported)
Shanson, 1987[33]	2957348	RCT	Paper	UK	Adults (aged 18-60 years), without underlying valve disease or prosthetic valve	Dental extraction (under GA)	Teicoplanin 400mg IV, 5-10mins before extraction	Amoxicillin 1g IM, 20-30mins before GA	NA		Control group - unmedicated	Blood cultures and antibiotic assay (treatment arms only) 1.5-2mins post-extraction. Antibiotic assay (treatment arms only) at 4h	40	40	NA	NA	40	2.50%	25%	NA	NA	32.50%	Not reported	VGS incidence. Control group 13/40. Teicoplanin group 1/40
Vergis, 2001[34]	11174592	RCT	Paper	USA	Adults (aged 21-79 years), without underlying valve disease or prosthetic valve	Dental extraction (anaesthetic not specified)	2 x 60ml mouthwash applications containing 3g amoxicillin suspension per application, retained for 1-2 mins each (2 hours and 1 hour before extraction)	3g amoxicillin PO (1 hour before extraction)	NA	NA	Control group - unmedicated	Blood cultures taken post-extraction (for 1 extraction, after extraction for 2 extractions, after 2nd tooth extracted for 3 or more extractions, after 3rd or 4th tooth extracted)	10	10	NA	NA	9	6/10 (60%)	1/10 (10%)	NA	NA	8/9 (89%)		

Abbreviations: LA – local anaesthetic, GA – general anaesthetic, IM intramuscular, IV intravenous, NA – not applicable, PO per oral, RCT – randomized controlled trial, TNR – trial, non-randomised
*only viridans streptococcal bacteraemia reported

Supplementary References

- 1 Bates KE, Hall M, Shah SS, et al. Trends in infective endocarditis hospitalisations at United States children's hospitals from 2003 to 2014: impact of the 2007 American Heart Association antibiotic prophylaxis guidelines. *Cardiology in the Young* 2016;**15**:1-5.
- 2 Bikdeli B, Wang Y, Kim N, et al. Trends in Hospitalization Rates and Outcomes of Endocarditis Among Medicare Beneficiaries. *Journal of the American College of Cardiology* 2013;**62**:2217-26.
- 3 Dayer MJ, Jones S, Prendergast B, et al. Incidence of infective endocarditis in England, 2000–13: a secular trend, interrupted time-series analysis. *The Lancet*; **385**:1219-28.
- 4 DeSimone DC, Tleyjeh IM, Correa de Sa DD, et al. Incidence of Infective Endocarditis Due to Viridans Group Streptococci Before and After the 2007 American Heart Association's Prevention Guidelines. *Mayo Clinic Proceedings* 2015;**90**:874-81.
- 5 Duval X, Delahaye F, Alla F, et al. Temporal Trends in Infective Endocarditis in the Context of Prophylaxis Guideline Modifications: Three Successive Population-Based Surveys. *Journal of the American College of Cardiology* 2012;**59**:1968-76.
- 6 Mackie AS, Liu W, Savu A, et al. Infective Endocarditis Hospitalizations Before and After the 2007 American Heart Association Prophylaxis Guidelines. *Canadian Journal of Cardiology* 2007;**32**:942-8.
- 7 Pant S, Patel NJ, Deshmukh A, et al. Trends in Infective Endocarditis Incidence, Microbiology, and Valve Replacement in the United States From 2000 to 2011. *Journal of the American College of Cardiology* 2015;**65**:2070-6.
- 8 Salam AMF, Albinali H, Singh R, et al. Incidence of infective endocarditis before and after the 2007 endocarditis prevention guidelines: a population-based study from Qatar (2002-2012). *European Heart Journal* 2014;**35**:908.
- 9 van den Brink FS, Swaans MJ, Hoogendijk MG, et al. Increased Incidence of Infective Endocarditis after the 2009 European Society of Cardiology Guideline Update: A Nation-wide Study in the Netherlands. *European Heart Journal - Quality of Care and Clinical Outcomes* 2016.
- 10 Horstkotte D, Rosin H, Friedrichs W, et al. Contribution for choosing the optimal prophylaxis of bacterial endocarditis. *European Heart Journal* 1987;**8**:379-81.
- 11 Imperiale TF, Horwitz RI. Does prophylaxis prevent postdental infective endocarditis? A controlled evaluation of protective efficacy. *The American Journal of Medicine* 1990;**88**:131-6.
- 12 Lacassin F, Hoen B, Leport C, et al. Procedures associated with infective endocarditis in adults. *European Heart Journal* 1995;**16**:1968-74.
- 13 van der Meer JTM, Michel MF, Valkenburg HA, et al. Efficacy of antibiotic prophylaxis for prevention of native-valve endocarditis. *The Lancet* 1992;**339**:135-9.

1
2
3 14 Asi KS, Gill AS, Mahajan S. Postoperative bacteremia in periodontal flap
4 surgery, with and without prophylactic antibiotic administration: A comparative study.
5 *J* 2010;**14**:18-22.
6

7 15 Baltch AL, Schaffer C, Hammer MC, et al. Bacteremia following dental
8 cleaning in patients with and without penicillin prophylaxis. *Am Heart J*
9 1982;**104**:1335-9.
10

11 16 Baltch AL, Pressman HL, Hammer MC, et al. Bacteremia Following Dental
12 Extractions in Patients with and without Penicillin Prophylaxis. *The American Journal*
13 *of the Medical Sciences* 1982;**283**:129-40.
14

15 17 Cannell H, Kerawala C, Sefton AM, et al. Failure of two macrolide antibiotics
16 to prevent post-extraction bacteraemia. *British Dental Journal* 1991;**171**:170-3.
17

18 18 Coulter WA, Coffey A, Saunders ID, et al. Bacteremia in children following
19 dental extraction. *J Dent Res* 1990;**69**:1691-5.
20

21 19 Dios PD, Carmona IT, Posse JL, et al. Comparative efficacies of amoxicillin,
22 clindamycin, and moxifloxacin in prevention of bacteremia following dental
23 extractions. *Antimicrobial Agents and Chemotherapy* 2006;**50**:2996-3002.
24

25 20 DuVall NB, Fisher TD, Hensley D, et al. The comparative efficacy of 0.12%
26 chlorhexidine and amoxicillin to reduce the incidence and magnitude of bacteremia
27 during third molar extractions: a prospective, blind, randomized clinical trial. *Oral*
28 *Surgery Oral Medicine Oral Pathology Oral Radiology* 2013;**115**:752-63.
29

30 21 Hall G, Hedstrom SA, Heimdahl A, et al. Prophylactic administration of
31 penicillins for endocarditis does not reduce the incidence of postextraction
32 bacteremia. *Clinical Infectious Diseases* 1993;**17**:188-94.
33

34 22 Hall G, Heimdahl A, Nord CE. Effects of prophylactic administration of
35 cefaclor on transient bacteremia after dental extraction. *European Journal of Clinical*
36 *Microbiology & Infectious Diseases* 1996;**15**:646-9.
37

38 23 Head TW, Bentley KC, Millar EP, et al. A comparative study of the
39 effectiveness of metronidazole and penicillin V in eliminating anaerobes from
40 postextraction bacteremias. *Oral surgery, oral medicine, and oral pathology*
41 1984:152-5.
42

43 24 Khairat O. An effective antibiotic cover for the prevention of endocarditis
44 following dental and other post-operative bacteraemias. *J Clin Pathol* 1966;**19**:561-6.
45

46 25 Limeres Posse J, Álvarez Fernández M, Fernández Feijoo J, et al.
47 Intravenous amoxicillin/clavulanate for the prevention of bacteraemia following dental
48 procedures: a randomized clinical trial. *Journal of Antimicrobial Chemotherapy* 2016.
49

50 26 Lockhart PB, Brennan MT, Kent ML, et al. Impact of amoxicillin prophylaxis
51 on the incidence, nature, and duration of bacteremia in children after intubation and
52 dental procedures. *Circulation* 2004;**109**:2878-84.
53

54 27 Lockhart PB, Brennan MT, Sasser HC, et al. Bacteremia Associated With
55 Toothbrushing and Dental Extraction. *Circulation* 2008;**117**:3118-25.
56
57
58
59
60

- 1
2
3 28 Maharaj B, Coovadia Y, Vayej AC. A comparative study of amoxicillin,
4 clindamycin and chlorhexidine in the prevention of post-extraction bacteraemia.
5 *Cardiovasc* 2012;**23**:491-4.
6
- 7 29 Maskell JP, Carter JLB, Boyd RB, et al. Teicoplanin as a prophylactic
8 antibiotic for dental bacteraemia. *Journal of Antimicrobial Chemotherapy*
9 1986;**17**:651-9.
10
- 11 30 Roberts GJ, Radford P, R. H. Prophylaxis of dental bacteraemia with oral
12 amoxycillin in children. *Br Dent J* 1987;**162**:179-82.
13
- 14 31 Shanson DC, Cannon P, Wilks M. Amoxycillin compared with penicillin for the
15 prophylaxis of dental bacteraemia. *Journal of Antimicrobial Chemotherapy*
16 1978;**4**:431-6.
17
- 18 32 Shanson DC, Akash S, Harris M, et al. Erythromycin stearate, 1|5 g, for the
19 oral prophylaxis of streptococcal bacteraemia in patients undergoing dental
20 extraction: efficacy and tolerance. *Journal of Antimicrobial Chemotherapy*
21 1985;**15**:83-90.
22
- 23 33 Shanson DC, Shehata A, Tadayon M, et al. Comparison of intravenous
24 teicoplanin with intramuscular amoxycillin for the prophylaxis of streptococcal
25 bacteraemia in dental patients. *The Journal of antimicrobial chemotherapy* 1987:85-
26 93.
27
- 28 34 Vergis EN, Demas PN, Vaccarello SJ, et al. Topical antibiotic prophylaxis for
29 bacteremia after dental extractions. *Oral Surgery, Oral Medicine, Oral Pathology,*
30 *Oral Radiology, and Endodontology* 2001;**91**:162-5.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



PRISMA 2009 Checklist

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	5
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	6
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	7
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	NA
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	8
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	8
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Suppl
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	8
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	8
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	8
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	8
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	8-9
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis.	8-9



PRISMA 2009 Checklist

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	9
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	NA
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	10
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	14, 15, Suppl
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	Suppl
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	14, 15, Figures
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	Figures
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	Suppl
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	NA
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	16
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	16
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	17
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	NA

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097