



DATA NOTE

The genome sequence of the Small Magpie moth, *Anania*

hortulata (Linnaeus, 1758)

[version 1; peer review: awaiting peer review]

Douglas Boyes¹⁺, Denise C. Wawman²,
University of Oxford and Wytham Woods Genome Acquisition Lab,
Darwin Tree of Life Barcoding collective,
Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory
team,
Wellcome Sanger Institute Scientific Operations: Sequencing Operations,
Wellcome Sanger Institute Tree of Life Core Informatics team,
Tree of Life Core Informatics collective, Darwin Tree of Life Consortium

¹UK Centre for Ecology & Hydrology, Wallingford, England, UK²Edward Grey Institute, Department of Biology, University of Oxford, Oxford, England, UK

+ Deceased author

V1 First published: 21 Jun 2024, 9:327
<https://doi.org/10.12688/wellcomeopenres.22462.1>
Latest published: 21 Jun 2024, 9:327
<https://doi.org/10.12688/wellcomeopenres.22462.1>

Abstract

We present a genome assembly from an individual female *Anania hortulata* (the Small Magpie; Arthropoda; Insecta; Lepidoptera; Crambidae). The genome sequence is 612.2 megabases in span. Most of the assembly is scaffolded into 32 chromosomal pseudomolecules, including the W and Z sex chromosomes. The mitochondrial genome has also been assembled and is 15.23 kilobases in length.

Keywords

Anania hortulata, Small Magpie moth, genome sequence, chromosomal, Lepidoptera



This article is included in the [Tree of Life](#) gateway.

Open Peer Review

Approval Status AWAITING PEER REVIEW

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Darwin Tree of Life Consortium (mark.blaxter@sanger.ac.uk)

Author roles: **Boyes D:** Investigation, Resources; **Wawman DC:** Writing – Original Draft Preparation, Writing – Review & Editing;

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Wellcome through core funding to the Wellcome Sanger Institute [206194, <https://doi.org/10.35802/206194>] and the Darwin Tree of Life Discretionary Award [218328, <https://doi.org/10.35802/218328>]. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2024 Boyes D *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Boyes D, Wawman DC, University of Oxford and Wytham Woods Genome Acquisition Lab *et al.* **The genome sequence of the Small Magpie moth, *Anania hortulata* (Linnaeus, 1758) [version 1; peer review: awaiting peer review]** Wellcome Open Research 2024, 9:327 <https://doi.org/10.12688/wellcomeopenres.22462.1>

First published: 21 Jun 2024, 9:327 <https://doi.org/10.12688/wellcomeopenres.22462.1>

Species taxonomy

Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Endopterygota; Amphimesenoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Ditrysia; Obtectomera; Pyraloidea; Crambidae; Pyraustinae; *Anania*; *Anania hortulata* (Linnaeus, 1758) (NCBI:txid1594229).

Background

The Small Magpie *Anania hortulata* is one of the larger micro-moths in the family Crambidae, having a wing length of 13–16 mm (Sterling *et al.*, 2023). It is distinctively marked with greyish-black markings on a white background, and bright yellow at the wing base, on the head, and at the tip of the abdomen, with finer yellow bands across the abdomen and thorax.

Anania hortulata is common throughout the United Kingdom where most adults are on the wing from May to September, but it has been recorded from February to November. It is commonly found in gardens and parks, on waste ground, hedgerows, woodland rides and scrub, where it can be easily disturbed during the day and attached to light. The main larval food plant is Common Nettle *Urtica dioica*, but larva sometimes feed on Field Bindweed *Convolvulus arvensis*, Hedge Bindweed *Calystegia sepium*, Black Horehound *Ballota nigra*, White Horehound *Marrubium vulgare*, Mints *Mentha* spp. and Woundworts *Stachys* spp. (Sterling *et al.*, 2023).

Anania hortulata is widely distributed in the Northern Hemisphere, where it is found across Europe and Asia and it has been introduced to North America (Yang & Landry, 2019). An analysis of the DNA of specimens of *A. hortulata* from across its range, using both DNA and detailed morphometric analysis, showed minor differences in the genitalia, corresponding to clustering in the DNA sequences of the mitochondrial cytochrome oxidase I (COI) and nuclear carbamoyl-phosphate synthase domain (CAD) genes, between moths in China and the rest of its range, leading to the Chinese specimens being split into a different species, *Anania sinensis* (Yang & Landry, 2019).

We present a chromosomally complete genome sequence for *Anania hortulata*, based on one female specimen collected at Wytham Woods for the Darwin Tree of Life Project.

Genome sequence report

The genome was sequenced from one female *Anania hortulata* (Figure 1) collected from Wytham Woods, Oxfordshire, UK (51.77, -1.34). A total of 67-fold coverage in Pacific Biosciences single-molecule HiFi long reads was generated. Primary assembly contigs were scaffolded with chromosome conformation Hi-C data. Manual assembly curation corrected 13 missing joins or mis-joins and removed one haplotypic duplication, reducing the scaffold number by 12.73%.



Figure 1. Photograph of the *Anania hortulata* (ilAnaHort2) specimen used for genome sequencing.

The final assembly has a total length of 612.2 Mb in 47 sequence scaffolds with a scaffold N50 of 21.0 Mb (Table 1). The snail plot in Figure 2 provides a summary of the assembly statistics, while the distribution of assembly scaffolds on GC proportion and coverage is shown in Figure 3. The cumulative assembly plot in Figure 4 shows curves for subsets of scaffolds assigned to different phyla. Most (99.81%) of the assembly sequence was assigned to 32 chromosomal-level scaffolds, representing 30 autosomes and the W and Z sex chromosomes. Chromosome-scale scaffolds confirmed by the Hi-C data are named in order of size (Figure 5; Table 2). Chromosomes W and Z were assigned based on read coverage statistics. While not fully phased, the assembly deposited is of one haplotype. Contigs corresponding to the second haplotype have also been deposited. The mitochondrial genome was also assembled and can be found as a contig within the multifasta file of the genome submission.

The estimated Quality Value (QV) of the final assembly is 64.0 with *k*-mer completeness of 100.0%, and the assembly has a BUSCO v5.4.3 completeness of 98.7% (single = 98.4%, duplicated = 0.3%), using the lepidoptera_odb10 reference set (*n* = 5,286).

Metadata for specimens, BOLD barcode results, spectra estimates, sequencing runs, contaminants and pre-curation assembly statistics are given at <https://links.tol.sanger.ac.uk/species/1594229>.

Methods

Sample acquisition and nucleic acid extraction

An adult *Anania hortulata* (specimen ID Ox000595, ToLID ilAnaHort2) was collected from Wytham Woods, Oxfordshire (biological vice-county Berkshire), UK (latitude 51.77, longitude -1.34) on 2020-07-05 using a light trap. The

Table 1. Genome data for *Anania hortulata*, ilAnaHort2.1.

Project accession data		
Assembly identifier	ilAnaHort2.1	
Species	<i>Anania hortulata</i>	
Specimen	ilAnaHort2	
NCBI taxonomy ID	1594229	
BioProject	PRJEB67411	
BioSample ID	SAMEA7701459	
Isolate information	ilAnaHort2, whole organism	
Assembly metrics*		Benchmark
Consensus quality (QV)	64.0	≥ 50
<i>k</i> -mer completeness	100.0%	≥ 95%
BUSCO**	C:98.7%[S:98.4%,D:0.3%], F:0.4%,M:0.9%,n:5,286	C ≥ 95%
Percentage of assembly mapped to chromosomes	99.81%	≥ 95%
Sex chromosomes	W and Z	localised homologous pairs
Organelles	Mitochondrial genome: 15.23 kb	complete single alleles
Raw data accessions		
PacificBiosciences Revio	ERR12120042	
Hi-C Illumina	ERR12121866	
Genome assembly		
Assembly accession	GCA_963576865.1	
Accession of alternate haplotype	GCA_963576775.1	
Span (Mb)	612.2	
Number of contigs	119	
Contig N50 length (Mb)	11.7	
Number of scaffolds	47	
Scaffold N50 length (Mb)	21.0	
Longest scaffold (Mb)	29.16	

* Assembly metric benchmarks are adapted from column VGP-2020 of “Table 1: Proposed standards and metrics for defining genome assembly quality” from [Rhie et al. \(2021\)](#).

** BUSCO scores based on the lepidoptera_odb10 BUSCO set using version v5.4.3. C = complete [S = single copy, D = duplicated], F = fragmented, M = missing, n = number of orthologues in comparison. A full set of BUSCO scores is available at https://blobtoolkit.genomehubs.org/view/Anania_hortulata/dataset/GCA_963576865.1/busco.

specimen was collected and identified by Douglas Boyes (University of Oxford) and preserved on dry ice.

The workflow for high molecular weight (HMW) DNA extraction at the Wellcome Sanger Institute (WSI) Tree of Life Core Laboratory includes a sequence of core procedures: sample preparation; sample homogenisation, DNA extraction,

fragmentation, and clean-up. In sample preparation, the ilAnaHort2 sample was weighed and dissected on dry ice ([Jay et al., 2023](#)). Tissue from the whole organism was homogenised using a PowerMasher II tissue disruptor ([Denton et al., 2023a](#)). HMW DNA was extracted using the Automated MagAttract v2 protocol ([Oatley et al., 2023a](#)). DNA was sheared into an average fragment size of 12–20 kb




Figure 2. Genome assembly of *Anania hortulata*, ilAnaHort2.1: metrics. The BlobToolKit snail plot shows N50 metrics and BUSCO gene completeness. The main plot is divided into 1,000 size-ordered bins around the circumference with each bin representing 0.1% of the 612,213,012 bp assembly. The distribution of scaffold lengths is shown in dark grey with the plot radius scaled to the longest scaffold present in the assembly (29,160,819 bp, shown in red). Orange and pale-orange arcs show the N50 and N90 scaffold lengths (20,978,505 and 12,875,000 bp), respectively. The pale grey spiral shows the cumulative scaffold count on a log scale with white scale lines showing successive orders of magnitude. The blue and pale-blue area around the outside of the plot shows the distribution of GC, AT and N percentages in the same bins as the inner plot. A summary of complete, fragmented, duplicated and missing BUSCO genes in the lepidoptera_odb10 set is shown in the top right. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/Anania_hortulata/dataset/GCA_963576865.1/snail.

in a Megaruptor 3 system with speed setting 31 (Bates *et al.*, 2023). Sheared DNA was purified by solid-phase reversible immobilisation (Oatley *et al.*, 2023b): in brief, the method employs a 1.8X ratio of AMPure PB beads to sample to eliminate shorter fragments and concentrate the DNA. The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer and Qubit dsDNA High Sensitivity Assay kit. Fragment size distribution was evaluated by running the sample on the FemtoPulse system.

Protocols developed by the WSI Tree of Life laboratory are publicly available on protocols.io (Denton *et al.*, 2023b).

Sequencing

Pacific Biosciences HiFi circular consensus DNA sequencing libraries were constructed according to the manufacturers' instructions. DNA sequencing was performed by the Scientific Operations core at the WSI on a Pacific Biosciences Revio instrument. Hi-C data were also generated from tissue of ilAnaHort2 using the Arima v2 kit. The Hi-C sequencing was performed using paired-end sequencing with a read length of 150 bp on the Illumina NovaSeq 6000 instrument.

Genome assembly and curation

Assembly was carried out with Hifiasm (Cheng *et al.*, 2021) and haplotypic duplication was identified and removed with




Figure 3. Genome assembly of *Anania hortulata*, ilAnaHort2.1: BlobToolKit GC-coverage plot. Sequences are coloured by phylum. Circles are sized in proportion to sequence length. Histograms show the distribution of sequence length sum along each axis. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/Anania_hortulata/dataset/GCA_963576865.1/blob.

purge_dups (Guan *et al.*, 2020). The assembly was then scaffolded with Hi-C data (Rao *et al.*, 2014) using YaHS (Zhou *et al.*, 2023). The assembly was checked for contamination and corrected using the TreeVal pipeline (Pointon *et al.*, 2023). Manual curation was performed using JBrowse2 (Diesh *et al.*, 2023), HiGlass (Kerpedjiev *et al.*, 2018) and PretextView (Harry, 2022). The mitochondrial genome was assembled using MitoHiFi (Uliano-Silva *et al.*, 2023), which runs MitoFinder (Allio *et al.*, 2020) or MITOS (Bernt *et al.*, 2013) and uses

these annotations to select the final mitochondrial contig and to ensure the general quality of the sequence.

Final assembly evaluation

The final assembly was post-processed and evaluated with the three Nextflow (Di Tommaso *et al.*, 2017) DSL2 pipelines “sanger-tol/readmapping” (Surana *et al.*, 2023a), “sanger-tol/genomenote” (Surana *et al.*, 2023b), and “sanger-tol/blobtoolkit” (Muffato *et al.*, 2024). The pipeline sanger-tol/readmapping




Figure 4. Genome assembly of *Anania hortulata*, ilAnaHort2.1: BlobToolKit cumulative sequence plot. The grey line shows cumulative length for all sequences. Coloured lines show cumulative lengths of sequences assigned to each phylum using the buscogenes taxrule. An interactive version of this figure is available at https://blobtoolkit.genomehubs.org/view/Anania_hortulata/dataset/GCA_963576865.1/cumulative.

aligns the Hi-C reads with bwa-mem2 (Vasimuddin *et al.*, 2019) and combines the alignment files with SAMtools (Danecek *et al.*, 2021). The sanger-tol/genomenote pipeline transforms the Hi-C alignments into a contact map with BEDTools (Quinlan & Hall, 2010) and the Cooler tool suite (Abdennur & Mirny, 2020), which is then visualised with HiGlass (Kerpedjiev *et al.*, 2018). It also provides statistics about the assembly with the NCBI datasets (Sayers *et al.*, 2024) report, computes *k*-mer completeness and QV consensus quality values with FastK and MerquryFK, and a completeness assessment with BUSCO (Manni *et al.*, 2021).

The sanger-tol/blobtoolkit pipeline is a Nextflow port of the previous Snakemake Blobtoolkit pipeline (Challis *et al.*, 2020). It aligns the PacBio reads with SAMtools and minimap2 (Li, 2018) and generates coverage tracks for regions of fixed size. In parallel, it queries the GoAT database (Challis *et al.*, 2023) to identify all matching BUSCO lineages to run BUSCO (Manni *et al.*, 2021). For the three domain-level BUSCO lineage, the pipeline aligns the BUSCO

genes to the Uniprot Reference Proteomes database (Bateman *et al.*, 2023) with DIAMOND (Buchfink *et al.*, 2021) blastp. The genome is also split into chunks according to the density of the BUSCO genes from the closest taxonomically lineage, and each chunk is aligned to the Uniprot Reference Proteomes database with DIAMOND blastx. Genome sequences that have no hit are then chunked with seqtk and aligned to the NT database with blastn (Altschul *et al.*, 1990). All those outputs are combined with the blobtools suite into a blobdir for visualisation.

All three pipelines were developed using the nf-core tooling (Ewels *et al.*, 2020), use MultiQC (Ewels *et al.*, 2016), and make extensive use of the Conda package manager, the Bioconda initiative (Grüning *et al.*, 2018), the Biocontainers infrastructure (da Veiga Leprevost *et al.*, 2017), and the Docker (Merkel, 2014) and Singularity (Kurtzer *et al.*, 2017) containerisation solutions.

Table 3 contains a list of relevant software tool versions and sources.




Figure 5. Genome assembly of *Anania hortulata*, ilAnaHort2.1: Hi-C contact map of the ilAnaHort2.1 assembly, visualised using HiGlass. Chromosomes are shown in order of size from left to right and top to bottom. An interactive version of this figure may be viewed at <https://genome-note-higlass.tol.sanger.ac.uk/l/?d=j2rxfcQEQRaOp5lXqj2JYg>.

Table 2. Chromosomal pseudomolecules in the genome assembly of *Anania hortulata*, ilAnaHort2.

INSDC accession	Chromosome	Length (Mb)	GC%
OY756388.1	1	25.81	36.5
OY756389.1	2	25.03	36.5
OY756390.1	3	24.84	36.5
OY756391.1	4	24.12	36.5
OY756392.1	5	23.02	36.0
OY756393.1	6	22.94	36.5
OY756394.1	7	22.72	36.0
OY756395.1	8	21.93	36.5
OY756396.1	9	21.75	36.0
OY756397.1	10	21.64	36.0
OY756398.1	11	21.34	36.5
OY756399.1	12	21.08	36.0
OY756400.1	13	20.98	36.5
OY756401.1	14	20.95	36.5
OY756403.1	15	19.84	37.0

INSDC accession	Chromosome	Length (Mb)	GC%
OY756404.1	16	19.48	36.5
OY756405.1	17	19.47	36.5
OY756406.1	18	19.21	36.5
OY756407.1	19	19.2	36.5
OY756408.1	20	17.58	36.5
OY756409.1	21	16.78	36.5
OY756410.1	22	16.42	36.5
OY756411.1	23	14.77	36.5
OY756412.1	24	14.39	36.5
OY756413.1	25	12.88	36.5
OY756414.1	26	12.74	37.0
OY756415.1	27	11.25	39.0
OY756416.1	28	10.48	37.0
OY756417.1	29	9.66	37.5
OY756418.1	30	9.39	37.0
OY756402.1	W	20.22	38.0
OY756387.1	Z	29.16	36.0
OY756419.1	MT	0.02	20.5

Table 3. Software tools: versions and sources.

Software tool	Version	Source
BEDTools	2.30.0	https://github.com/arq5x/bedtools2
Blast	2.14.0	ftp://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/
BlobToolKit	4.3.7	https://github.com/blobtoolkit/blobtoolkit
BUSCO	5.4.3 and 5.5.0	https://gitlab.com/ezlab/busco
bwa-mem2	2.2.1	https://github.com/bwa-mem2/bwa-mem2
Cooler	0.8.11	https://github.com/open2c/cooler
DIAMOND	2.1.8	https://github.com/bbuchfink/diamond
fasta_windows	0.2.4	https://github.com/tolkite/fasta_windows
FastK	427104ea91c78c3b8b8b49f1a7d6bbeaa869ba1c	https://github.com/thegenemyers/FASTK
GoaT CLI	0.2.5	https://github.com/genomehubs/goat-cli
Hifiasm	0.19.5-r587	https://github.com/chhyllp123/hifiasm
HiGlass	44086069ee7d4d3f6f3f0012569789ec138f42b84aa44357826c0b6753eb28de	https://github.com/higlass/higlass
MercuryFK	d00d98157618f4e8d1a9190026b19b471055b22e	https://github.com/thegenemyers/MERQURY.FK
MitoHiFi	3	https://github.com/marcelauliano/MitoHiFi
MultiQC	1.14, 1.17, and 1.18	https://github.com/MultiQC/MultiQC
NCBI Datasets	15.12.0	https://github.com/ncbi/datasets
Nextflow	23.04.0-5857	https://github.com/nextflow-io/nextflow
PretextView	0.2	https://github.com/wtsi-hpag/PretextView
purge_dups	1.2.5	https://github.com/dfguan/purge_dups
samtools	1.16.1, 1.17, and 1.18	https://github.com/samtools/samtools
sanger-tol/genomenote	1.1.1	https://github.com/sanger-tol/genomenote
sanger-tol/readmapping	1.2.1	https://github.com/sanger-tol/readmapping
Seqtk	1.3	https://github.com/lh3/seqtk
Singularity	3.9.0	https://github.com/sylabs/singularity
TreeVal	1.0.0	https://github.com/sanger-tol/treeval
YaHS	1.2a.2	https://github.com/c-zhou/yahs

Wellcome Sanger Institute – Legal and Governance

The materials that have contributed to this genome note have been supplied by a Darwin Tree of Life Partner. The submission of materials by a Darwin Tree of Life Partner is subject to the ‘**Darwin Tree of Life Project Sampling Code of Practice**’, which can be found in full on the Darwin Tree of Life website [here](#). By agreeing with and signing up to the Sampling Code of Practice, the Darwin Tree of Life Partner

agrees they will meet the legal and ethical requirements and standards set out within this document in respect of all samples acquired for, and supplied to, the Darwin Tree of Life Project.

Further, the Wellcome Sanger Institute employs a process whereby due diligence is carried out proportionate to the nature of the materials themselves, and the circumstances under which

they have been/are to be collected and provided for use. The purpose of this is to address and mitigate any potential legal and/or ethical implications of receipt and use of the materials as part of the research project, and to ensure that in doing so we align with best practice wherever possible. The overarching areas of consideration are:

- Ethical review of provenance and sourcing of the material
- Legality of collection, transfer and use (national and international)

Each transfer of samples is further undertaken according to a Research Collaboration Agreement or Material Transfer Agreement entered into by the Darwin Tree of Life Partner, Genome Research Limited (operating as the Wellcome Sanger Institute), and in some circumstances other Darwin Tree of Life collaborators.

Data availability

European Nucleotide Archive: *Anania hortulata* (small magpie). Accession number PRJEB67411; <https://identifiers.org/ena.embl/PRJEB67411> (Wellcome Sanger Institute, 2023). The genome sequence is released openly for reuse. The *Anania hortulata* genome sequencing initiative is part of the Darwin Tree of Life (DTOL) project. All raw sequence data and the assembly have been deposited in INSDC databases. The genome will be annotated using available RNA-Seq data

and presented through the [Ensembl](#) pipeline at the European Bioinformatics Institute. Raw data and assembly accession identifiers are reported in [Table 1](#).

Author information

Members of the University of Oxford and Wytham Woods Genome Acquisition Lab are listed here: <https://doi.org/10.5281/zenodo.7125292>.

Members of the Darwin Tree of Life Barcoding collective are listed here: <https://doi.org/10.5281/zenodo.4893703>.

Members of the Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team are listed here: <https://doi.org/10.5281/zenodo.10066175>.

Members of Wellcome Sanger Institute Scientific Operations: Sequencing Operations are listed here: <https://doi.org/10.5281/zenodo.10043364>.

Members of the Wellcome Sanger Institute Tree of Life Core Informatics team are listed here: <https://doi.org/10.5281/zenodo.10066637>.

Members of the Tree of Life Core Informatics collective are listed here: <https://doi.org/10.5281/zenodo.5013541>.

Members of the Darwin Tree of Life Consortium are listed here: <https://doi.org/10.5281/zenodo.4783558>.

References

- Abdennur N, Mirny LA: **Cooler: scalable storage for Hi-C data and other genomically labeled arrays.** *Bioinformatics.* 2020; **36**(1): 311–316. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Allio R, Schomaker-Bastos A, Romiguier J, et al.: **MitoFinder: efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics.** *Mol Ecol Resour.* 2020; **20**(4): 892–905. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Altschul SF, Gish W, Miller W, et al.: **Basic local alignment search tool.** *J Mol Biol.* 1990; **215**(3): 403–410. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bateman A, Martin MJ, Orchard S, et al.: **UniProt: the Universal protein knowledgebase in 2023.** *Nucleic Acids Res.* 2023; **51**(D1): D523–D531. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bates A, Clayton-Lucey I, Howard C: **Sanger Tree of Life HMW DNA fragmentation: diagenode Megaruptor³ for LI PacBio.** *protocols.io.* 2023. [Publisher Full Text](#)
- Bernt M, Donath A, Jühling F, et al.: **MITOS: improved *de novo* metazoan mitochondrial genome annotation.** *Mol Phylogenet Evol.* 2013; **69**(2): 313–319. [PubMed Abstract](#) | [Publisher Full Text](#)
- Buchfink B, Reuter K, Drost HG: **Sensitive protein alignments at Tree-of-Life scale using DIAMOND.** *Nat Methods.* 2021; **18**(4): 366–368. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challis R, Kumar S, Sotero-Caio C, et al.: **Genomes on a Tree (GoaT): a versatile, scalable search engine for genomic and sequencing project metadata across the eukaryotic Tree of Life [version 1; peer review: 2 approved].** *Wellcome Open Res.* 2023; **8**: 24. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challis R, Richards E, Rajan J, et al.: **BlobToolKit – interactive quality assessment of genome assemblies.** *G3 (Bethesda).* 2020; **10**(4): 1361–1374. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cheng H, Concepcion GT, Feng X, et al.: **Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm.** *Nat Methods.* 2021; **18**(2): 170–175. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- da Veiga Leprevost F, Grüning BA, Alves Aflitos S, et al.: **BioContainers: an open-source and community-driven framework for software standardization.** *Bioinformatics.* 2017; **33**(16): 2580–2582. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Danecek P, Bonfield JK, Liddle J, et al.: **Twelve years of SAMtools and BCFtools.** *GigaScience.* 2021; **10**(2): giab008. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Denton A, Oatley G, Cornwell C, et al.: **Sanger Tree of Life sample homogenisation: PowerMash.** *protocols.io.* 2023a. [Publisher Full Text](#)
- Denton A, Yatsenko H, Jay J, et al.: **Sanger Tree of Life wet laboratory protocol collection V.1.** *protocols.io.* 2023b. [Publisher Full Text](#)
- Di Tommaso P, Chatzou M, Floden EW, et al.: **Nextflow enables reproducible computational workflows.** *Nat Biotechnol.* 2017; **35**(4): 316–319. [PubMed Abstract](#) | [Publisher Full Text](#)
- Diesh C, Stevens GJ, Xie P, et al.: **JBrowse 2: a modular genome browser with views of synteny and structural variation.** *Genome Biol.* 2023; **24**(1): 74. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ewels P, Magnusson M, Lundin S, et al.: **MultiQC: summarize analysis results for multiple tools and samples in a single report.** *Bioinformatics.* 2016; **32**(19): 3047–3048. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ewels PA, Peltzer A, Fillinger S, et al.: **The nf-core framework for community-curated bioinformatics pipelines.** *Nature Biotechnol.* 2020; **38**(3): 276–278. [PubMed Abstract](#) | [Publisher Full Text](#)

- Grüning B, Dale R, Sjödin A, *et al.*: **Bioconda: sustainable and comprehensive software distribution for the life sciences.** *Nat Methods.* 2018; **15**(7): 475–476. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Guan D, McCarthy SA, Wood J, *et al.*: **Identifying and removing haplotypic duplication in primary genome assemblies.** *Bioinformatics.* 2020; **36**(9): 2896–2898. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Harry E: **PretextView (Paired Read Texture Viewer): a desktop application for viewing pretext contact maps.** 2022; [Accessed 19 October 2022]. [Reference Source](#)
- Jay J, Yatsenko H, Narváez-Gómez JP, *et al.*: **Sanger Tree of Life sample preparation: triage and dissection.** *protocols.io.* 2023. [Publisher Full Text](#)
- Kerpedjiev P, Abdennur N, Lekschas F, *et al.*: **HiGlass: web-based visual exploration and analysis of genome interaction maps.** *Genome Biol.* 2018; **19**(1): 125. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kurtzer GM, Sochat V, Bauer MW: **Singularity: scientific containers for mobility of compute.** *PLoS One.* 2017; **12**(5): e0177459. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Li H: **Minimap2: pairwise alignment for nucleotide sequences.** *Bioinformatics.* 2018; **34**(18): 3094–3100. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Manni M, Berkeley MR, Seppy M, *et al.*: **BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.** *Mol Biol Evol.* 2021; **38**(10): 4647–4654. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Merkel D: **Docker: lightweight Linux containers for consistent development and deployment.** *Linux J.* 2014; **2014**(239): 2. [Reference Source](#)
- Muffato M, Butt Z, Challis R, *et al.*: **Sanger-tol/blobtoolkit: v0.3.0 - poliwig.** 2024. [Publisher Full Text](#)
- Oatley G, Denton A, Howard C: **Sanger Tree of Life HMW DNA extraction: automated MagAttract v.2.** *protocols.io.* 2023a. [Publisher Full Text](#)
- Oatley G, Sampaio F, Howard C: **Sanger Tree of Life fragmented DNA clean up: automated SPRI.** *protocols.io.* 2023b. [Publisher Full Text](#)
- Pointon DL, Eagles W, Sims Y, *et al.*: **sanger-tol/treeval v1.0.0 - Ancient Atlantis.** 2023. [Publisher Full Text](#)
- Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing genomic features.** *Bioinformatics.* 2010; **26**(6): 841–842. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rao SSP, Huntley MH, Durand NC, *et al.*: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping.** *Cell.* 2014; **159**(7): 1665–1680. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, McCarthy SA, Fedrigo O, *et al.*: **Towards complete and error-free genome assemblies of all vertebrate species.** *Nature.* 2021; **592**(7856): 737–746. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sayers EW, Cavanaugh M, Clark K, *et al.*: **GenBank 2024 update.** *Nucleic Acids Res.* 2024; **52**(D1): D134–D137. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sterling P, Parsons M, Lewington R: **Field Guide to the Micro-moths of Great Britain and Ireland, Second Edition.** London: Bloomsbury Publishing. 2023. [Reference Source](#)
- Surana P, Muffato M, Qi G: **Sanger-tol/readmapping: sanger-tol/readmapping v1.1.0 - Hebridean Black (1.1.0).** *Zenodo.* 2023a. [Publisher Full Text](#)
- Surana P, Muffato M, Sadasivan Baby C: **Sanger-tol/genomenote (v1.0.dev).** *Zenodo.* 2023b. [Publisher Full Text](#)
- Uliano-Silva M, Ferreira JGRN, Krasheninnikova K, *et al.*: **MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity reads.** *BMC Bioinformatics.* 2023; **24**(1): 288. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vasimuddin M, Misra S, Li H, *et al.*: **Efficient architecture-aware acceleration of BWA-MEM for multicore systems.** In: *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS).* IEEE, 2019; 314–324. [Publisher Full Text](#)
- Wellcome Sanger Institute: **The genome sequence of the Small Magpie, *Anania hortulata* (Linnaeus, 1758).** European Nucleotide Archive, [dataset], accession number PRJEB67411. 2023.
- Yang Z, Landry JF: **Allopatric separation represents an overlooked cryptic species in the *Anania hortulata* species complex (Lepidoptera: Crambidae: Pyraustinae): congruence between genetic and morphological evidence** *Can Entomol.* 2019; **151**(2): 163–186. [Publisher Full Text](#)
- Zhou C, McCarthy SA, Durbin R: **YaHS: yet another Hi-C Scaffolding tool.** *Bioinformatics.* 2023; **39**(1): btac808. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)