

1 **Isotope evidence for agricultural extensification reveals how the world's first**  
2 **cities were fed**

3

4 Amy K. Styring<sup>1\*</sup>, Michael Charles<sup>1</sup>, Federica Fantone<sup>2</sup>, Mette Marie Hald<sup>3</sup>, Augusta  
5 McMahon<sup>4</sup>, Richard H. Meadow<sup>5</sup>, Geoff K. Nicholls<sup>6</sup>, Ajita K. Patel<sup>7</sup>, Mindy C.  
6 Pitre<sup>8</sup>, Alexia Smith<sup>9</sup>, Arkadiusz Sołtysiak<sup>10</sup>, Gil Stein<sup>11</sup>, Jill A. Weber<sup>12</sup>, Harvey  
7 Weiss<sup>13</sup>, Amy Bogaard<sup>1</sup>

8

9 <sup>1</sup> School of Archaeology, University of Oxford, Oxford, UK.

10 Email: [Amy.Styring@arch.ox.ac.uk](mailto:Amy.Styring@arch.ox.ac.uk)

11 <sup>2</sup> Faculty of Archaeology, Leiden University, The Netherlands

12 <sup>3</sup> The National Museum of Denmark, Copenhagen, Denmark

13 <sup>4</sup> Division of Archaeology, University of Cambridge, UK

14 <sup>5</sup> Department of Anthropology and Peabody Museum, Harvard University, USA

15 <sup>6</sup> Department of Statistics, University of Oxford, Oxford, UK

16 <sup>7</sup> Peabody Museum, Harvard University, USA

17 <sup>8</sup> Department of Anthropology, St. Lawrence University, New York, USA

18 <sup>9</sup> Department of Anthropology, University of Connecticut, Connecticut, USA

19 <sup>10</sup> Institute of Archaeology, University of Warsaw, Poland

20 <sup>11</sup> The Oriental Institute, University of Chicago, USA

21 <sup>12</sup> University of Pennsylvania, Philadelphia, Pennsylvania, USA

22 <sup>13</sup> School of Forestry and Environmental Studies, Yale University, USA

23

24 **Abstract**

25 This study sheds light on the agricultural economy that underpinned the emergence of  
26 the first urban centres in northern Mesopotamia. Using  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of crop  
27 remains from the sites of Tell Sabi Abyad, Tell Zeidan, Hamoukar, Tell Brak and Tell  
28 Leilan (6500–2000 cal BC), we reveal that labour intensive practices such as  
29 manuring/middening and water management formed an integral part of the  
30 agricultural strategy from the seventh millennium BC. Increased agricultural  
31 production to support growing urban populations was achieved by cultivation of  
32 larger areas of land, entailing lower manure/midden inputs per unit area—  
33 *extensification*. Our findings paint a nuanced picture of the role of agricultural

34 production in new forms of political centralisation. While the shift towards lower  
35 input farming most plausibly developed gradually at a household level, the increased  
36 importance of land-based wealth constituted a key potential source of political power,  
37 providing the possibility for greater bureaucratic control and contributing to the wider  
38 societal changes that accompanied urbanisation.

39

## 40 **Introduction**

41

### 42 **Strategies to increase crop production**

43 The emergence of the first urban centres represents a pivotal moment in human  
44 history, and much research has focussed on changes in the political, social and  
45 productive economy that accompanied and likely contributed to this change<sup>1,2</sup>. In this  
46 study we consider the stable carbon isotope ( $\delta^{13}\text{C}$ ) and nitrogen isotope ( $\delta^{15}\text{N}$ ) values  
47 of 276 charred cereal grain and 44 pulse seed samples (each comprising 4–25  
48 individual grains/seeds) from the sites of Tell Sabi Abyad, Tell Zeidan, Hamoukar,  
49 Tell Brak and Tell Leilan, located in the Khabur and Balikh drainage basins in  
50 northern Mesopotamia and dating to between 6500 and 2000 BC (Figure 1; Table 1).  
51 This allows us to investigate how the staple economy supported the new population  
52 centres that emerged in the fourth and third millennia BC in northern Mesopotamia,  
53 and thus to reconsider wider debates surrounding the agroecology of early urbanism,  
54 its sustainability and the role of political centralisation in shaping some of the world's  
55 earliest urbanised landscapes.

56

57 A generalised narrative of agricultural 'intensification' has long held sway in  
58 discussion of early urbanisation around the world, in part due to emphasis on  
59 irrigation-based societies<sup>3</sup>. Influential research based in southern Mesopotamia<sup>4,5</sup>,  
60 where irrigation is obligatory and associated with high area yields, has encouraged a  
61 prevailing view that urban civilisation in the rainfed north was likewise supported by  
62 investing higher labour inputs per unit area<sup>2,6–8</sup>. The productive potential of northern  
63 Mesopotamia in recent times, however, has depended on very extensive cultivation,  
64 augmented since World War One by tractors, pump irrigation and agrochemicals,  
65 combined with effective systems of mobilisation and transport<sup>9,10</sup>. When did this  
66 process of extensification begin? Was it initiated by the early cities of northern

67 Mesopotamia in the fourth and third millennia BC, or were these early urban centres  
68 instead dependent on high-intensity land management like their southern  
69 counterparts?

70

71 Although not always explicitly defined in discussion of agricultural practice, here we  
72 refer to agricultural intensity in terms of labour and resource inputs per unit area of  
73 land <sup>11</sup>, placing the emphasis on the *intention* to increase outputs (crop yield) rather  
74 than outputs *per se*. Agricultural *intensification* involves an increase in inputs,  
75 resulting in increased crop yields per unit area of land. Practices that could have  
76 involved high inputs of labour and resources include manuring or middening with  
77 human, animal and/or household waste <sup>8</sup>, controlling weeds through weeding or  
78 turning over the soil and/or decreasing the frequency at which land was left fallow <sup>12</sup>.

79 Management of the water available to crops, e.g. through strategic watering or  
80 planting of less drought-tolerant crops in better watered settings/soils, may have been  
81 another labour- and resource-intensive agricultural strategy in northern Mesopotamia,  
82 given its relatively low (c. 200–500 mm/yr) and highly variable annual rainfall.

83 Increased agricultural production through agricultural *extensification*, by contrast, is  
84 enabled by significant expansion of the land under cultivation, such that reduction of  
85 inputs and yields per unit area are offset by a larger absolute scale of production <sup>13</sup>.

86 Extensification can occur through implementation of labour-saving techniques such as  
87 ploughing, specifically through use of specialized plough animals capable of  
88 preparing a much larger area for sowing than can be achieved manually by a farming  
89 family. Such radical expansion in arable scale requires an additional supply of labour  
90 at harvest time, a system that implies a level of organisation of labour beyond the  
91 immediate household.

92

93 Of course, increasing inputs per unit area or expanding the absolute scale of  
94 cultivation are not mutually exclusive means of increasing production, and a mixture  
95 of the two could be employed to meet the needs of a growing and aggregating  
96 population. Indeed, scatters of abraded sherds (proposed to be derived from household  
97 waste that was spread on fields to improve their fertility) and ‘hollow ways’  
98 (extensive tracks resulting from confining animal movement to areas between  
99 cultivated fields) have been given as evidence for both intensification of manuring  
100 and expansion of cultivated land in the third millennium BC <sup>14,15</sup>, respectively. These

101 sherd scatters and ‘hollow way’ features can be difficult to date, however <sup>16</sup>, and it is  
102 possible that third millennium practices have obscured earlier evidence of manuring.  
103 There is also no way of inferring from this off-site evidence whether cereals, pulses  
104 and oil-seed crops were treated differently and thus how their management mapped  
105 onto these landscapes.

106

### 107 **Directly determining agricultural intensity using crop isotope values**

108 Crop isotope values offer an opportunity to improve our understanding of how  
109 agricultural strategies changed under urbanisation, delivering a complementary  
110 approach to off-site methods that provides *direct* evidence for the growing conditions  
111 of specific crops, better temporal resolution of changing farming practice and more  
112 nuanced insight into the relative importance of intensification and extensification  
113 among sites and contexts. Crop  $\delta^{15}\text{N}$  values largely reflect the  $\delta^{15}\text{N}$  value of the soil in  
114 which they are grown, which in turn is strongly influenced by land use history <sup>17</sup>. In  
115 particular, application of animal manure has been found to increase the  $\delta^{15}\text{N}$  values of  
116 soil and cereals by as much as 10‰, relating to the intensity—amount and  
117 frequency—of manuring <sup>18,19</sup>, as well as to the type of organic matter—compost,  
118 animal manure, household waste—applied <sup>20</sup>. From now on, we use the term  
119 manuring/middening to encapsulate the various means by which organic matter could  
120 have been added to the soil. Intensive manuring/middening requires a high input of  
121 labour—being heavy to transport and spread—and in modern farming studies usually  
122 goes hand-in-hand with other labour-intensive cultivation practices such as weeding  
123 and hoeing, since it enhances the tractability (ease of working) of soil <sup>21</sup>. Crop  $\delta^{15}\text{N}$   
124 values can therefore act as a proxy for the general intensity of agricultural practice, or  
125 labour inputs per unit area. Crop  $\delta^{13}\text{C}$  values reflect the movement of carbon dioxide  
126 through the stomata, which in dry climates is most strongly influenced by the water  
127 status of a crop during its growth period <sup>22</sup>. Since rainfall was relatively low at some  
128 of the sites and during some of the time periods in our study—and thus marginal for  
129 rainfed farming—it is possible that the water status of crops was manipulated in some  
130 way, whether through direct watering or by strategic planting of relatively demanding  
131 crops in areas with greater water availability such as the bottoms or slopes of wadis.  
132 Crop  $\delta^{13}\text{C}$  values can therefore help to elucidate how cultivation was configured in

133 the landscape and identify strategic (and potentially high input) crop management in  
134 relation to water resources.

135

136 In this study we aim to provide a better understanding of how agricultural intensity  
137 changed during two phases of urbanisation in northern Mesopotamia: the Late  
138 Chalcolithic period (4400–3000 cal BC) and the Early Bronze Age (2600–2000 cal  
139 BC). Lawrence and Wilkinson<sup>23</sup> have identified three distinct pathways to urbanism,  
140 characterised by different site types: ‘hub sites’, which grew slowly in areas of  
141 already dense and gradually increasing population (e.g. Tell Brak and Late  
142 Chalcolithic Hamoukar); ‘endogenous upstarts’, which developed rapidly through the  
143 movement of local populations into the urban centre (e.g. Tell Leilan); and  
144 ‘exogenous upstarts’, which also developed rapidly but in areas with little pre-existing  
145 settlement. Within this framework, we can determine whether these contrasting urban  
146 trajectories, with likely different underlying social contexts, entailed different forms  
147 of agricultural practice at Tell Brak, Hamoukar and Tell Leilan. The political and  
148 productive economies of these sizeable population centres are compared to those of  
149 the Late Neolithic settlement of Tell Sabi Abyad (c. 6500–5200 cal BC) and the  
150 Ubaid–Late Chalcolithic 2 town of Tell Zeidan (c. 5300–3850 cal BC). These data  
151 will constrain current models of agricultural intensity and give an unparalleled insight  
152 into changing agricultural practice through time, as settlements expanded and  
153 contracted, and city-states became established. Moreover, by considering *direct*  
154 evidence of crop growing conditions and farming practice, we hope to provide a  
155 counterpoint to top-down ‘elite’ views of agricultural production and move towards a  
156 more ‘bottom-up farmer-centric perspective’ of agricultural change<sup>24</sup>.

157

## 158 **Results**

159

### 160 **Determining manuring intensity at archaeological sites**

161 Aridity can increase plant  $\delta^{15}\text{N}$  values<sup>e.g. 25</sup> and it is therefore necessary to take this  
162 into account when inferring manuring intensity from cereal grain  $\delta^{15}\text{N}$  values. Styring  
163 et al.<sup>19</sup> used the relationship between modern plant  $\delta^{15}\text{N}$  values and rainfall in the  
164 eastern Mediterranean<sup>26</sup> to adjust expected manuring rates based on  $\delta^{15}\text{N}$  values of  
165 cereal grains grown on controlled farming plots in temperate Europe<sup>18,27</sup>. While this

166 allows more accurate (and more conservative) estimates of manuring intensity from  
167 cereal grain  $\delta^{15}\text{N}$  values in semi-arid regions, here we use the cereal grain isotope data  
168 from our studies of modern farming regimes across a wider range of rainfall zones  
169 <sup>18,19,28</sup> to refine this method. Figure 2 shows cereal grain  $\delta^{15}\text{N}$  values from present-day  
170 farming regimes, colour-coded by their known manuring level, plotted against the  
171 natural log of annual rainfall derived from interpolation of average monthly climate  
172 data for 1960–1990, available from the WorldClim database <sup>29</sup>. High manuring  
173 represents annual manuring of crops at rates equivalent to 30+ tonnes manure/ha;  
174 medium manuring represents either annual or biennial manuring of crops at lower  
175 levels (less than 20 tonnes/ha); and low manuring represents no manuring within the  
176 last 3+ years. The lines represent a fitted linear model relating cereal grain  $\delta^{15}\text{N}$   
177 values to mean annual rainfall for each manuring level.

178

179 In this study we use a method called Bayesian multiple imputation <sup>30</sup> to assign  
180 manuring levels to the archaeological samples and simultaneously test for a  
181 relationship between these assigned manuring levels and site-related parameters such  
182 as size and date (see Supplementary Information for more details). In the first  
183 “imputation” stage of multiple imputation, the probability of each cereal grain sample  
184 having each manuring level value is derived. Cereal grain  $\delta^{15}\text{N}$  values, known  
185 manuring levels and rainfall data from our studies of modern farming regimes inform  
186 the regression parameters, and these combine with the  $\delta^{15}\text{N}$  values and past rainfall  
187 ranges of the archaeological material to assign the missing manuring level value (raw  
188 archaeological crop  $\delta^{15}\text{N}$  values in Figure 3 and Supplementary Table 1). We also  
189 allow for uncertainty in past rainfall level values.

190

191 In the second “analysis” stage of multiple imputation we sample multiple assigned  
192 manuring level data sets from the manuring level probability distribution we specified  
193 at the first stage. We analyse each of these data sets and combine the analyses. In this  
194 way our analysis allows for uncertainty in the assigned manuring levels. At this stage  
195 of our analysis, manuring level is an outcome that is potentially explained by site size  
196 and other site-specific covariates, such as date. We therefore regress our assigned  
197 manuring level data on site size and test for a negative (i.e. one sided) effect. Site size  
198 is based on field surveys of concentrated surface sherd scatters (as opposed to the

199 continuous and more sparse scatters of small, abraded sherds that have been  
200 interpreted as manuring) and targeted excavation (see Table 1 for references). We use  
201 a mixed-effects proportional-odds regression model for the three-level ordinal  
202 manuring level outcome (low, medium, high). We include a random effect due to site  
203 in order to model unmeasured site-specific variables that might influence the outcome  
204 and to allow for uncertainty in sample date. Given a particular set of assigned  
205 manuring levels we carry out Bayesian inference using flat priors, so that all remotely  
206 possible parameter values are equally probable *a priori*. The posterior probability  
207 distribution for the site size regression parameter is the average of the posterior  
208 distributions taken over for each of the assigned manuring level data sets. The final  
209 estimated posterior odds (equivalently here, the Bayes Factor) are 5.5 to 1 in favour of  
210 a negative effect of site size on the manuring level of cereal grains (Figure 4). This  
211 estimate incorporates all of the important sources of uncertainty and gives substantial  
212 support for an effect of site size on cereal grain manuring levels.

213

214 The downside of Bayesian multiple imputation is its relative complexity. In order to  
215 make the essential logic of the analysis more accessible, we repeated the analysis  
216 using a very simple single imputation approach, and tests based on the maximum  
217 likelihood estimator (see Supplementary Information for more details). Based on the  
218 fitted linear model relating modern cereal grain  $\delta^{15}\text{N}$  values to mean annual rainfall  
219 for each manuring level (Figure 2), each archaeological sample of cereal grains is  
220 assigned the manuring level that *predicts* a  $\delta^{15}\text{N}$  value as close as possible to the  
221 *observed*  $\delta^{15}\text{N}$  value for that sample. We then regress this assigned manuring level  
222 against site size using the mixed-effects proportional-odds regression model described  
223 above and test for a negative effect due to site size (using a Wald test). Again, we find  
224 clear evidence for an effect ( $p=0.0034$ ), but note that this figure does not allow for  
225 uncertainty in the imputed manuring levels as the multiple imputation approach does.  
226 All our analyses are explained in more detail in the Supplementary Information.

227

### 228 **Spatial variation in agricultural strategy**

229 The variability in the  $\delta^{15}\text{N}$  values of archaeological cereal grain samples demonstrates  
230 that cereals were grown under a range of manuring conditions at each of the  
231 archaeological sites (Figure 3). It seems that we can exclude floodplain cultivation as

232 a potential cause of high  $\delta^{15}\text{N}$  values (denitrification during seasonal flooding can  
233 result in enrichment of soil  $^{15}\text{N}^{31}$ ) because  $\delta^{15}\text{N}$  values of modern barley grains  
234 grown without manure in dry wadi beds that were temporarily flooded following  
235 heavy winter rains in 2014 in the south of Morocco are included in both manuring  
236 level imputation models. However, isotope analysis of more cereals growing in  
237 seasonally flooded settings would be beneficial to test this observation. The large  
238 number of cereal grain samples with high  $\delta^{15}\text{N}$  values (and thus with a low probability  
239 of having a medium or lower manuring level; Figure 4) at Tell Sabi Abyad  
240 demonstrates that manuring/middening formed an integral part of the agricultural  
241 strategy from as early as the seventh millennium cal BC in northern Mesopotamia,  
242 rather than developing later as a reaction to the need to feed a growing population <sup>e.g.</sup>  
243 <sup>12</sup>.

244  
245 Since manure is a heavy and bulky resource to transport, manuring intensity is  
246 generally governed by frictions of distance and is thus likely to be highest in plots  
247 immediately surrounding settlement areas where animal dung from stabled livestock  
248 and/or midden material accumulates <sup>8</sup>. Spatially, then, it is plausible that variable  
249 manuring levels within an archaeological site reflect a spectrum of manuring intensity  
250 radiating out from the settlement—from intensively managed ‘infield’ areas to more  
251 extensively managed fields further away from the urban core. This model mirrors the  
252 ‘halos’ of abraded pot sherd scatters surrounding many third millennium BC urban  
253 centres and the radiating track ways (‘hollow ways’) that extend beyond these scatters  
254 and are believed to delineate the extent of arable cultivation <sup>8</sup>.

255  
256 There is also evidence that individual households had access to cereals grown under a  
257 range of conditions, presumably harvested from plots at varying distances from the  
258 site. Barley and glume wheat (einkorn and emmer) grain samples (n=8) representing  
259 material stored in separate pots in a single household at LC3–4 Tell Brak, cleaned of  
260 weed seeds and sieved for human consumption <sup>32</sup>, have  $\Delta^{13}\text{C}$  values (converted from  
261 determined  $\delta^{13}\text{C}$  values, see Methods section) and  $\delta^{15}\text{N}$  values ranging from 15.1–  
262 18.4‰ and 1.3–8.6‰, respectively. The large range in crop isotope values  
263 demonstrates that household cereal stores derived from plots encompassing a wide  
264 spectrum of agricultural intensity. This is consistent with the household having access

265 to land at a range of distances from the urban centre; higher manure inputs could be  
266 maintained on plots closer to the settlement and better water retention was likely on  
267 soils closer to the Wadi Jaghjagh and Wadi Radd, c. 3 km from the site<sup>33</sup>. There is  
268 also the possibility that some of the crops had come from surrounding villages as a  
269 form of tribute<sup>34</sup> or that crops were farmed on communal land, but while it cannot  
270 necessarily be assumed that this household farmed/owned the disparate plots of land  
271 from which their crops came, its access to such variable production points to a risk-  
272 buffering strategy in the household's own interest<sup>cf. 35</sup>.

273

#### 274 **Changing agricultural practice**

275 Both the single and multiple imputation approaches show that the probability that  
276 cereals received low levels of manure tends to increase as site size increases (Figure  
277 4). Although attempting to relate settlement size to population is fraught with  
278 uncertainty<sup>36</sup>, given the large range in settlement size considered in this study—from  
279 the 1 ha village of Tell Sabi Abyad to the 130 ha sprawl of early fourth millennium  
280 BC Tell Brak—we feel it is valid to treat site size as at least a *general* index of  
281 population. The results of the imputation models predict that as settlements in  
282 northern Mesopotamia expanded and agricultural production increased, cereals were  
283 grown with lower manure inputs per unit area, suggesting that to sustain greater  
284 agricultural production, the area of land under cultivation must have increased  
285 through a process of extensification. Whilst overall effort expended in manuring plots  
286 may well have increased at larger settlements, the crop isotope results demonstrate  
287 that the *bulk* of the increase in cereal production came from expansion of less  
288 intensively manured plots, plausibly those lying beyond the immediate environs of the  
289 urban centre.

290

291 Our results support the findings of Arous et al.<sup>37</sup>, who observed a general trend of  
292 decreasing cereal grain  $\delta^{15}\text{N}$  values through time at various sites in the Near East.  
293 Arous et al.<sup>37</sup> interpreted this general decrease in cereal  $\delta^{15}\text{N}$  values through time as a  
294 decrease in soil fertility that could have been caused by myriad potential factors,  
295 including agricultural overexploitation, cultivation of marginal lands and reduced  
296 manure application. Our model strongly indicates that increasing site size, resulting in  
297 a deliberate change in agricultural practice that involved decreased manuring inputs,

298 was a more important factor in decreasing crop  $\delta^{15}\text{N}$  values than soil degradation  
299 resulting from years of poor agricultural management. This is exemplified by the  
300 higher proportion of low-level manured cereal grain samples at 130 ha Tell Brak (c.  
301 3800–3600 cal BC) compared to the contemporary 15 ha site of Hamoukar (Figure  
302 4a).

303

304 Until now, scatters of abraded sherds dated to the third millennium BC have been  
305 interpreted as the earliest evidence for manuring at Tell Brak, thus reflecting an  
306 intensification of agricultural inputs during the Early Bronze Age <sup>7,8</sup>. Our new results  
307 reveal that the appearance of these sherd scatters does not correlate with an increase in  
308 manuring level, at least of *cereals*. Complementary weed ecological data from Tell  
309 Brak show that fertility levels in cereal fields were also relatively low at this time,  
310 which is consistent with fields receiving low organic matter/manure inputs <sup>38</sup>. Thus,  
311 while sherd scatters are considered to be a visible and persistent sign of the spreading  
312 of organic household waste, perhaps this practice primarily benefited garden crops <sup>8, 14</sup>  
313 and/or was necessitated due to a decrease in the availability of animal manure,  
314 perhaps due to competing demands for its use as fuel <sup>39</sup>. A shift to a highly specialised  
315 pastoral economy focussed on sheep and goat driven by the commodification of  
316 textile production in the third millennium BC <sup>40</sup>, together with the expansion of land  
317 under arable production, would have extended herding into more marginal areas,  
318 thereby reducing the opportunity for manure collection. The relatively high proportion  
319 of cereal grain samples receiving low levels of manure at this time demonstrates that  
320 any manuring of fields by animals allowed to graze on stubble or fallow land was not  
321 comparable with the increase in cereal grain  $\delta^{15}\text{N}$  values that result from deliberate  
322 spreading of stall manure.

323

#### 324 **Crop management in relation to water resources**

325 The archaeological cereal grain and pulse seed  $\delta^{13}\text{C}$  values, which have been  
326 converted into  $\Delta^{13}\text{C}$  values to allow comparison with modern crop studies (see  
327 Methods section), can reveal crop management strategies and thus provide insight into  
328 how arable land was configured to exploit the hydrology of the landscape. Figure 5  
329 shows the  $\Delta^{13}\text{C}$  values of hulled barley, wheat (free-threshing and glume wheats) and  
330 pulses (lentil, pea and grass pea) through time. There are no significant changes in

331 crop  $\Delta^{13}\text{C}$  values across the time period studied, something that would be expected if  
332 the water status of crops was governed solely by the variable rainfall (see  
333 palaeoclimate records from e.g. Lake Van and Soreq Cave<sup>41,42</sup>).

334

335 The lack of significant variation in crop  $\Delta^{13}\text{C}$  values with time suggests that there was  
336 some degree of crop management in relation to water resources at all of the sites. This  
337 observation need not necessarily equate to irrigation or deliberate watering, but  
338 minimally implies that crops were strategically sown in areas with better water  
339 availability, perhaps close to wadis or in soils/areas that retained water, to buffer them  
340 from the effects of low rainfall. A study of barley grain  $\Delta^{13}\text{C}$  values by Riehl et al.<sup>43</sup>  
341 from archaeological sites across the Fertile Crescent observed lower  $\Delta^{13}\text{C}$  values  
342 (indicative of poorer water status) during aridification episodes *only* in the most  
343 marginal settings for agriculture. Thus, our results and those reported by Riehl et al.<sup>43</sup>  
344 reflect the difficulty of using crop  $\delta^{13}\text{C}$  values as evidence of climate change *per se*,  
345 but instead highlight their potential to measure agronomic adaptation to  
346 (independently verified) climate change.

347

348 There is also no significant difference in the  $\Delta^{13}\text{C}$  values of barley, wheat and pulses  
349 *cf.*<sup>44</sup>. Modern studies have shown, however, that if barley is grown in the same  
350 watering conditions as wheat and pulses, it will tend to have a higher  $\Delta^{13}\text{C}$  value;  
351 offsets range from 1‰ in two-row barley to 2‰ in six-row barley *e.g.*<sup>45,46</sup>. There are  
352 indications that the  $\Delta^{13}\text{C}$  values of ancient barley and wheat were also offset, though  
353 the magnitude of this offset may have been smaller: six-row barley grain samples  
354 (n=59) recovered from the archaeological site of Hornstaad-Hörnle IA, Germany  
355 (3909 cal BC) have  $\Delta^{13}\text{C}$  values that are 1.1‰ higher than those of wheat grain  
356 samples (naked wheat and einkorn; n=120) grown in the same year<sup>47</sup>. Since these  
357 cereals were unlikely to have received additional water inputs, this demonstrates that  
358 a c. 1‰ difference in the  $\Delta^{13}\text{C}$  values of barley and wheat grown in the same watering  
359 conditions is observed in these ancient crops. Data from Neolithic Kouphovouno,  
360 Greece are also consistent with an offset between two-row hulled barley and wheat,  
361 though crop remains here do not derive from a single year's harvest, and were  
362 conceivably affected by water management in a Mediterranean zone<sup>48</sup>. When crop  
363  $\Delta^{13}\text{C}$  values are plotted against watering bands adjusted for the physiological

364 differences between barley (mostly two-row) and other crops (Figure 5), the majority  
365 of the barley grain samples fall into the poor and moderately watered bands defined  
366 by Wallace et al. <sup>46</sup>, suggesting that yields were limited by water availability. In  
367 contrast, the majority of wheat and pulse samples fall into the well watered band  
368 (yields are not limited by water availability). The better water status of wheats and  
369 pulses compared to barley further supports an hypothesis of strategic agricultural  
370 practice because, at least today, barley generally tolerates drier conditions better than  
371 wheat, peas and lentils <sup>49</sup>. This strategy would therefore have maximised overall crop  
372 yields in a region where water availability likely presented a key limitation to the  
373 optimal growth of crops.

374

### 375 **The role of cereals in the economy**

376 Plotting carbon and nitrogen isotope values of human and faunal bone collagen  
377 alongside crop isotope values can reveal the importance of crops in the diet of both  
378 humans and animals. Figure 6 shows the isotope values of crops, fauna and humans  
379 from the Late Chalcolithic and Early Bronze Age at Tell Brak and for the Early  
380 Bronze Age at Tell Leilan (faunal and human bone collagen  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values are  
381 in Supplementary Table 2). Shaded ellipses represent the expected distributions (mean  
382  $\pm 2$  standard deviations) of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of individuals consuming various  
383 dietary combinations of cereal grains, pulses and animal products (milk and/or meat).  
384 The determined human isotope values (with their mean  $\pm 2$  standard deviations  
385 distribution outlined in black) clearly overlap more closely with the ellipses  
386 corresponding to lower animal protein consumption in all periods, suggesting that  
387 cereal grains were likely an important part of the human diet (cf. grain ration records  
388 <sup>50</sup>).

389

390 During LC3–4, in particular, there is a shift towards higher faunal  $\delta^{13}\text{C}$  values,  
391 indicating greater  $\text{C}_4$  plant consumption (e.g. *Cyperus*, *Eragrostis*), which is  
392 consistent with grazing on more marginal steppe areas that received lower rainfall.  
393 While consumption of  $\text{C}_3$  plants growing in areas of lower water availability would  
394 also result in higher faunal  $\delta^{13}\text{C}$  values, the particularly high  $\delta^{13}\text{C}$  values of some of  
395 the fauna ( $> -18\text{‰}$ ), can only be due to consumption of  $\text{C}_4$  plants with  $\delta^{13}\text{C}$  values of  
396 c.  $14\text{‰}$  <sup>51</sup>. Tell Brak was 130 ha in extent at this time, and the expansion in cultivation

397 indicated independently by the crop isotope values would plausibly have  
398 complemented this movement of animals away from the settlement. In the Early  
399 Bronze Age at Tell Leilan, the faunal isotope values overlap entirely with the ellipse  
400 for 100% cereal grain consumption, indicating that a significant portion of the  
401 herbivore and pig diet could also have comprised cereal grains. The possibility of  
402 cereal grains being grown as fodder for domestic herbivores has been suggested  
403 previously on the basis of archaeobotanical findings of large quantities of un-cleaned  
404 barley grains at Tell Brak<sup>52</sup> and Tell Leilan<sup>53</sup>, and textual references to allocations of  
405 cereals as animal fodder<sup>50</sup>.

406

## 407 **Discussion**

408

409 The cereal grain isotope values from archaeological sites in the Khabur and Balikh  
410 drainage basins provide a spatial and temporal perspective on changing agricultural  
411 practice prior to and during two phases of urbanisation. The relatively small  
412 settlements of Tell Sabi Abyad and Tell Zeidan (1 and 12 ha, respectively) yielded  
413 cereal grains with relatively high  $\delta^{15}\text{N}$  values that are consistent with high levels of  
414 manuring/middening being practiced from as early as the seventh millennium BC,  
415 thousands of years earlier than the appearance of sherd scatters that have previously  
416 been the primary evidence for manuring<sup>14</sup>. This evidence for early manuring is  
417 contrary to evolutionary models of agricultural development that suggest that highly  
418 labour intensive practices such as manuring were only employed when population  
419 pressure induced such changes<sup>12</sup>. Moreover, we find that strategic crop management  
420 in relation to water resources played a key role in cereal and pulse cultivation at even  
421 the early sites<sup>44,54</sup>, likely as a deliberate means of ensuring adequate production in  
422 such a water-limited region. The centrality of cereals in both human and animal diets,  
423 as seen in bone collagen isotope values, explains this considerable investment.

424

425 Cereal grain nitrogen isotope values reveal that increased agricultural production to  
426 support growing urban populations in northern Mesopotamia was achieved by  
427 cultivation of larger areas of land, using lower manure/midden inputs per unit area—  
428 *extensification*<sup>9</sup>. This evidence for expansion of arable land is in agreement with off-  
429 site survey evidence for extensive arable catchment areas around urban centres

430 (mainly in the third millennium cal BC), indicated by radiating ‘hollow ways’<sup>8</sup> and  
431 regional surveys of site distributions<sup>10,55</sup>. It also aligns with the economies of scale  
432 gained from aggregations of population<sup>56</sup>, since there would have been a supply of  
433 labour at crucial bottlenecks in the agricultural year (such as harvest time) that could  
434 be mobilised from amongst other cadres of society (cf. Sumerian city-states<sup>57</sup>).  
435 Extensification as a means of increasing arable production is in line with the model of  
436 extensive agriculture proposed by Weiss<sup>9,10</sup> for northern Mesopotamia and Halstead  
437<sup>13</sup> for the provisioning of the urban palatial economies of Late Bronze Age southern  
438 Greece, and with evidence for highly intensive management from the initial  
439 establishment of farming in Europe<sup>27</sup> and the Near East<sup>19,58</sup>.

440

441 The relationship between agricultural intensity and settlement size transcends third to  
442 fourth millennium BC differences in social complexity and urban form<sup>23</sup>. Thus, the  
443 shift towards lower input farming at larger urban centres most plausibly developed  
444 gradually at a household level; as households sought to increase production, plots  
445 receiving low labour inputs expanded relative to the more intensively managed plots.  
446 Moreover, while our data are consistent with an overall strategy of extensification,  
447 this broader framework subsumes a range of behavioural variation that is testimony to  
448 a bottom-up as opposed to top-down driver of agricultural change. Individual  
449 households seem to have practiced a nuanced and flexible strategy in which: a) the  
450 crops planted, b) where they were planted, and c) labour and material inputs of water  
451 and/or manure were all fine-tuned to the specific characteristics of the crop, land  
452 and/or soil quality, and the highly variable rainfall circumstances of any given year.  
453 This diversity in agricultural practice makes sense as a household risk-buffering  
454 strategy but not as an elite-controlled share-cropping regime<sup>35</sup>.

455

456 Nonetheless, this extensive agriculture directly ties production to the amount of land  
457 under cultivation—rather than to inputs and therefore to yields per unit area—and  
458 heightens the importance of land-based wealth that can be transferred from generation  
459 to generation<sup>59</sup>. Thus, extensification could fuel inherited wealth inequality as a  
460 potential source of political power. Linking agricultural outputs to land rather than  
461 labour inputs also provides a much more tangible measure upon which to base levels  
462 of taxation/tribute<sup>60</sup>, permitting greater bureaucratic control over surplus, which  
463 could have benefited those in political power. Ultimately, this study reveals that the

464 expansive agricultural economy was integral to the development of these first  
465 northern Mesopotamian cities, driving—as well as being driven by—the wider  
466 societal changes that accompanied this urban trajectory.

467

468

## 469 **Methods**

470

471 **Modern cereal grains.** Carbon and nitrogen isotope analysis was carried out on 273  
472 cereal grain samples (each representing a homogenised batch of 50 cereal grains of  
473 the same taxon) from 14 farming sites/regions. Details of the site locations, mean  
474 annual rainfall, taxon and manuring regimes for each of the crop samples are in Table  
475 2. The grains in each sample were homogenised prior to isotope analysis using a Spex  
476 2760 FreezerMill.

477

478 **Archaeological cereal grains.** Carbon and nitrogen isotope analysis was carried out  
479 on 276 cereal grain and 44 pulse seed samples (each representing a homogenised  
480 batch of 4–25 grains/seeds of the same taxon) from five archaeological sites. Twenty-  
481 seven of the samples from Tell Brak are the same as those whose  $\Delta^{13}\text{C}$  values are  
482 reported in Wallace et al. <sup>44</sup>; these are identified in Table S1. Cereal grains were  
483 recovered in a carbonised state from a range of contexts including storage rooms,  
484 domestic fires, cooking ovens and floors. The chronology of crop samples was based  
485 on stratigraphic dating and radiocarbon ages. Details of the site locations, present-day  
486 mean annual rainfall, estimated past rainfall ranges and settlement size can be found  
487 in Table 1 and isotopic data for each of the crop samples are listed in Table S1.

488

489 **Human and faunal bone collagen.** Carbon and nitrogen isotope analyses were  
490 carried out on bone and dentine collagen isolates of 60 humans and 31 herbivores  
491 (cattle, gazelle, goat and sheep) from LC2, LC3–4 and EBA Tell Brak and 7 humans,  
492 13 herbivores (cattle, gazelle, goat and sheep) and 8 pigs from EBA Tell Leilan. The  
493 occupation periods were selected based on those that had isotope data for crops, fauna  
494 and humans. Details of the archaeological contexts in which bones were found,  
495 chronology and isotopic data for each of the bone collagen isolates are listed in  
496 Supplementary Table 2. Only collagen values with C/N ratios between 2.9 and 3.6

497 were studied, following quality criteria described by DeNiro <sup>61</sup>, and samples with a  
498 collagen yield < 1% were visually inspected to ensure that they looked like collagen.  
499

500 **Preparation of carbonised crop remains for isotope analysis.** Cereal grains and  
501 pulse seeds were examined at ×7–45 magnification for visible surface contaminants,  
502 such as adhering sediment or plant roots; these were removed by gentle scraping.  
503 Around 10% of the total number of samples from each site were scraped clean,  
504 crushed and analysed using Fourier transform infrared spectroscopy with attenuated  
505 total reflectance (FTIR-ATR) to look for the presence of carbonate, nitrate and/or  
506 humic contamination <sup>62</sup>. Peaks characteristic of carbonate contamination (870 and 720  
507 cm<sup>-1</sup>) were observed in three of the FTIR spectra of samples from Tell Brak but in  
508 none of the FTIR spectra from other sites (Supplementary Figure 1). It was therefore  
509 decided to acid pre-treat all of the crop samples from Tell Brak to dissolve any  
510 carbonate <sup>63</sup>. This procedure consists of treatment with 10 ml of 0.5 M hydrochloric  
511 acid at 70°C for thirty to sixty minutes, then rinsing in distilled water three times  
512 before freeze-drying. This acid pre-treatment does not affect the isotope values of  
513 charred grains in the absence of contamination <sup>62</sup>. The samples (both acid pre-treated  
514 and untreated) were crushed using an agate mortar and pestle.

515

516 **Preparation of bones and dentine for isotope analysis.** Between 0.5 and 1 g of  
517 bone/dentine was cleaned of any visible dirt or carbonate crusts using an aluminium  
518 oxide air abrasive. Collagen was isolated using a modified ‘gelatinization method’  
519 based on the methods of Longin <sup>64</sup> and ultrafiltered prior to isotope analysis.

520

521 **Stable carbon and nitrogen isotope analysis.** All samples (modern and  
522 archaeological, crop and bone collagen) were weighed into tin capsules for stable  
523 isotope analysis. The  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of cereal grains and bone collagen were  
524 determined on a SerCon EA-GSL mass spectrometer. The  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of  
525 crops were determined in separate runs due to the low %N in the samples. An internal  
526 alanine standard was used to calculate raw isotope ratios. For  $\delta^{13}\text{C}$  determinations of  
527 crops, two-point normalization to the VPDB scale was carried out using four  
528 replicates each of IAEA-C6 and IAEA-C7, while for  $\delta^{15}\text{N}$  determinations the  
529 standards were caffeine and IAEA-N2. For  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  determinations of bone

530 collagen, two-point normalization was carried out using four replicates each of  
531 caffeine and seal bone collagen. Reported measurement uncertainties are the  
532 calculated combined uncertainty of the raw measurement and reference standards,  
533 after Kragten <sup>65</sup>. The average measurement uncertainties for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of  
534 crops were 0.09‰ and 0.20‰, respectively. The average measurement uncertainties  
535 for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of bone collagen were 0.09‰ and 0.18‰, respectively.  
536 These calculations were performed using the statistical programming language R  
537 (3.0.2). The  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of carbonised crop remains were corrected for the  
538 effect of charring by subtracting 0.11‰ and 0.31‰, respectively, from the determined  
539  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values, which are the average offsets between uncarbonised crop seeds  
540 and those heated for 4, 8 or 24 hours at 215, 230, 245 or 260°C <sup>66</sup>.

541

542 **Conversion of  $\delta^{13}\text{C}$  values to  $\Delta^{13}\text{C}$  values.** The  $\Delta^{13}\text{C}$  values of modern cereal grains  
543 were calculated from the determined  $\delta^{13}\text{C}$  values ( $\delta^{13}\text{C}_{\text{plant}}$ ) and an average  $\delta^{13}\text{C}$  value  
544 of atmospheric  $\text{CO}_2$  ( $\delta^{13}\text{C}_{\text{air}}$ ) determined from air sampled at weekly intervals during  
545 the months that the crops were growing <sup>67</sup>, using the equation below. The  $\Delta^{13}\text{C}$  values  
546 of archaeological cereal grains were calculated from the determined  $\delta^{13}\text{C}$  values  
547 ( $\delta^{13}\text{C}_{\text{plant}}$ ) and a  $\delta^{13}\text{C}_{\text{air}}$  value approximated by the AIRCO2\_LOESS system <sup>45</sup>. The  
548 equation was defined by Farquhar et al. <sup>22</sup>:

549

$$\Delta^{13}\text{C} = \frac{\delta^{13}\text{C}_{\text{air}} - \delta^{13}\text{C}_{\text{plant}}}{1 + \delta^{13}\text{C}_{\text{plant}}/1000}$$

550

551

552 **Estimating past annual rainfall at archaeological sites.** We have used the  
553 difference between past and present-day annual rainfall at Soreq Cave (location in  
554 Figure 1 <sup>68</sup>), estimated from speleothem  $\delta^{18}\text{O}$  values and the present-day calibration  
555 relationship between speleothem  $\delta^{18}\text{O}$  values and rainfall, to adjust present-day  
556 annual rainfall at each of the archaeological sites and thus estimate past rainfall at  
557 200-year intervals (a 1‰ decrease in the  $\delta^{18}\text{O}$  value of precipitation is equivalent to  
558 an increase in annual rainfall of c. 200 mm <sup>68</sup>). The uncertainty associated with these  
559 estimates is accounted for in the multiple imputation model. Recent work has found  
560 that  $\delta^{18}\text{O}$  values of rainfall can be affected by the type of precipitation (convective or  
561 stratiform) as well as by the amount, and so these estimates of past annual rainfall  
562 based on the speleothem  $\delta^{18}\text{O}$  values may well be modified in the future <sup>69</sup>. Although

563 estimating past rainfall from a proxy record located c. 700 km away from the study  
564 sites is problematic, the general trend in climate is similar to that for other lower-  
565 resolution proxy records closer to the region. The  $\delta^{18}\text{O}$  values of sediments from Lake  
566 Mirabad in southwest Iran, Lake Zeribar in western Iran, and Lake Van in southeast  
567 Turkey (locations in Figure 1) also show generally wetter conditions than today  
568 between c. 7000 and 4000 BC followed by a trend towards greater aridity in the third  
569 millennium BC<sup>42,70,71</sup>. Dry phases indicated by higher  $\delta^{18}\text{O}$  values in c. 4500 BC,  
570 3300–3000 BC and 2500–1950 BC are also observed in multiple records<sup>72</sup>.

571

572 **Visualising potential dietary scenarios.** Figure 6 shows the expected distributions  
573 (mean  $\pm$  2 standard deviations) of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of individuals consuming  
574 various dietary combinations of cereal grains, pulses and animal products (milk and/or  
575 meat). The expected  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  value distribution of humans consuming 100%  
576 cereal grains are estimated by adding consumer-diet offsets of 4.8‰<sup>73</sup> and 4‰<sup>74</sup>,  
577 respectively, to the determined cereal grain  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values. The expected  $\delta^{13}\text{C}$   
578 and  $\delta^{15}\text{N}$  value distribution of humans consuming 100% pulses are estimated by  
579 adding consumer-diet offsets of 4.8‰<sup>73</sup> and 4‰<sup>74</sup>, respectively, to the determined  
580 pulse  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values. Since the consumer-diet  $\delta^{13}\text{C}$  value offset is smaller  
581 between carnivores and herbivores than between herbivores and plants<sup>72</sup>, the  
582 expected  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  value distribution of humans consuming 100% animal protein  
583 are estimated by adding consumer-diet offsets of 0.8‰<sup>73</sup> and 4‰<sup>74</sup>, respectively, to  
584 the determined faunal bone collagen  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values. The expected  $\delta^{13}\text{C}$  and  
585  $\delta^{15}\text{N}$  value distributions of humans consuming mixtures of these diets (20, 50 and  
586 80% animal protein, with the remaining proportion of the diet comprising a 50:50 mix  
587 of cereal grains and pulses) are estimated by adding the appropriate consumer-diet  
588 offsets to the  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of the dietary components, and multiplying each  
589 by their proportion in the diet.

590

591 **Data availability.** Raw  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of the archaeological crop samples are  
592 given in Supplementary Table 1. Raw  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of the modern crop  
593 samples used to infer the manuring levels of the archaeological crop samples are  
594 given in Supplementary Information (SI3). Full details of the statistical analysis  
595 (including R files) are available in Supplementary Information (SI1–19). The raw

596  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of the archaeological animal and human bone collagen are given  
597 in Supplementary Table 2.

598

599

600

601

602 **References**

603

- 604 1. Trigger, B. *Understanding Early Civilizations: A Comparative Study*. (Cambridge  
605 University Press, 2003).
- 606 2. Wilkinson, T. J. *et al.* Contextualizing early urbanization: settlement cores, early  
607 states and agro-pastoral strategies in the Fertile Crescent during the fourth and  
608 third millennia BC. *J. World Prehistory* **27**, 43–109 (2014).
- 609 3. *Agricultural Strategies*. (ed. Marcus, J. & Stanish, C.) (Cotsen Institute of  
610 Archaeology, 2006).
- 611 4. Adams, R. *Heartland of Cities: Surveys of Ancient Settlement and Land Use On*  
612 *the Central Floodplain at the Euphrates*. (University of Chicago Press, 1981).
- 613 5. Algaze, G. Initial social complexity in southwestern Asia: the Mesopotamian  
614 advantage. *Curr. Anthropol.* **42**, 199–233 (2001).
- 615 6. Sherratt, A. Water, soil and seasonality in early cereal cultivation. *World*  
616 *Archaeol.* **11**, 313–330 (1980).
- 617 7. Ur, J. & Colantoni, C. in *Inside Ancient Kitchens : New Directions in the Study of*  
618 *Daily Meals and Feasts* (ed. Klarich, E.) (University Press of Colorado, 2010).
- 619 8. Wilkinson, T. J. The structure and dynamics of dry-farming states in upper  
620 Mesopotamia [and comments and reply]. *Curr. Anthropol.* **35**, 483–520 (1994).
- 621 9. Weiss, H. Excavations at Tell Leilan and the origins of north Mesopotamian cities  
622 in the third millennium B.C. *Paléorient* **9**, 39–52 (1983).
- 623 10. Weiss, H. in *The Origins of Cities in Dry-Farming Syria and Mesopotamia in the*  
624 *Third Millennium B.C.* (ed. Weiss, H.) 71–108 (Four Quarters Publishing Co.,  
625 1986).
- 626 11. Morrison, K. D. The intensification of production: archaeological approaches. *J.*  
627 *Archaeol. Method Theory* **1**, 111–159 (1994).
- 628 12. Boserup, E. *The Conditions of Agricultural Growth*. (Aldine Publishing  
629 Company, 1965).
- 630 13. Halstead, P. Plough and power: the economic and social significance of  
631 cultivation with the ox-drawn ard in the Mediterranean. *Bull. Sumer. Agric.* **8**, 11–  
632 22 (1995).
- 633 14. Wilkinson, T. J. The definition of ancient manured zones by means of extensive  
634 sherd-sampling techniques. *J. Field Archaeol.* **9**, 323–333 (1982).

- 635 15. Wilkinson, T. J. Linear hollows in the Jazira, Upper Mesopotamia. *Antiquity* **67**,  
636 548–562 (1993).
- 637 16. Wilkinson, T. J., French, C., Ur, J. A. & Semple, M. The geoarchaeology of route  
638 systems in northern Syria. *Geoarchaeology* **25**, 745–771 (2010).
- 639 17. Peukert, S. *et al.* Understanding spatial variability of soil properties: a key step in  
640 establishing field- to farm-scale agro-ecosystem experiments. *Rapid Commun.*  
641 *Mass Spectrom.* **26**, 2413–2421 (2012).
- 642 18. Fraser, R. A. *et al.* Manuring and stable nitrogen isotope ratios in cereals and  
643 pulses: towards a new archaeobotanical approach to the inference of land use and  
644 dietary practices. *J. Archaeol. Sci.* **38**, 2790–2804 (2011).
- 645 19. Styring, A. K. *et al.* Disentangling the effect of farming practice from aridity on  
646 crop stable isotope values: A present-day model from Morocco and its application  
647 to early farming sites in the eastern Mediterranean. *Anthr. Rev.* **3**, 2–22 (2016).
- 648 20. Szpak, P. Complexities of nitrogen isotope biogeochemistry in plant-soil systems:  
649 implications for the study of ancient agricultural and animal management  
650 practices. *Plant Physiol.* **5**, 288 (2014).
- 651 21. Halstead, P. Traditional and ancient rural economy in Mediterranean Europe: plus  
652 ça change? *J. Hell. Stud.* **107**, 77–87 (1987).
- 653 22. Farquhar, G. D., Ehleringer, J. R. & Hubick, K. T. Carbon isotope discrimination  
654 and photosynthesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **40**, 503–537  
655 (1989).
- 656 23. Lawrence, D. & Wilkinson, T. J. Hubs and upstarts: pathways to urbanism in the  
657 northern Fertile Crescent. *Antiquity* **89**, 328–344 (2015).
- 658 24. Erickson, C. L. in *Agricultural Strategies* (eds. Marcus, J. & Stanish, C.) 334–363  
659 (Cotsen Institute of Archaeology, 2006).
- 660 25. Handley, L. L. *et al.* The  $^{15}\text{N}$  natural abundance ( $\delta^{15}\text{N}$ ) of ecosystem samples  
661 reflects measures of water availability. *Funct. Plant Biol.* **26**, 185–199 (1999).
- 662 26. Hartman, G. & Danin, A. Isotopic values of plants in relation to water availability  
663 in the Eastern Mediterranean region. *Oecologia* **162**, 837–852 (2010).
- 664 27. Bogaard, A. *et al.* Crop manuring and intensive land management by Europe's  
665 first farmers. *Proc. Natl. Acad. Sci.* **110**, 12589–12594 (2013).
- 666 28. Bogaard, A. *et al.* Combining functional weed ecology and crop stable isotope  
667 ratios to identify cultivation intensity: a comparison of cereal production regimes

- 668 in Haute Provence, France and Asturias, Spain. *Veg. Hist. Archaeobotany* **25**, 57–  
669 73 (2016).
- 670 29. Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G. & Jarvis, A. Very high  
671 resolution interpolated climate surfaces for global land areas. *Int. J. Climatol.* **25**,  
672 1965–1978 (2005).
- 673 30. Little, R. J. A. & Rubin, D. B. *Bayes and Multiple Imputation*. (John Wiley &  
674 Sons, Inc., 2002).
- 675 31. Finlay, J. C. & Kendall, C. in *Stable Isotopes in Ecology and Environmental*  
676 *Science* (eds. Michener, R. & Lajtha, K.) 283–333 (Blackwell Publishing Ltd.,  
677 2008).
- 678 32. Hald, M. M. *A Thousand Years of Farming: Late Chalcolithic Agricultural*  
679 *Practices at Tell Brak in Northern Mesopotamia*. (Archaeopress, 2008).
- 680 33. Ur, J. A., Karsgaard, P. & Oates, J. The spatial dimensions of early Mesopotamian  
681 urbanism: the Tell Brak suburban survey, 2003–2006. *Iraq* **73**, 1–19 (2011).
- 682 34. Sallaberger, W. & Ur, J. A. in *Third Millennium Cuneiform Texts from Tell*  
683 *Beydar* (eds. Milano, L., Sallaberger, W., Talon, P. & Van Lerberghe, K.) 51–71  
684 (Brepols Publishers, 2004).
- 685 35. Forbes, H. A. ‘We have a little of everything’: the ecological basis of some  
686 agricultural practices in Methana, Trizinia. *Ann. N. Y. Acad. Sci.* **268**, 236–250  
687 (1976).
- 688 36. Postgate, N. How many Sumerians per hectare? — probing the anatomy of an  
689 early city. *Camb. Archaeol. J.* **4**, 47–65 (1994).
- 690 37. Araus, J. L., Ferrio, J. P., Voltas, J., Aguilera, M. & Buxó, R. Agronomic  
691 conditions and crop evolution in ancient Near East agriculture. *Nat. Commun.* **5**,  
692 3953 (2014).
- 693 38. Bogaard, A. *et al.* From traditional farming in Morocco to early urban  
694 agroecology in northern Mesopotamia: combining present-day arable weed  
695 surveys and crop ‘isoscaples’ to reconstruct past agrosystems in (semi-)arid  
696 regions. *Environ. Archaeol.* (in press).
- 697 39. Smith, A., Dotzel, K., Fountain, J., Proctor, L. & von Baeyer, M. Examining fuel  
698 use in antiquity: archaeobotanical and anthracological approaches in southwest  
699 Asia. *Ethnobiol. Lett.* **6**, 192–195 (2015).
- 700 40. McCorrison, J. The fiber revolution: textile extensification, alienation, and social  
701 stratification in ancient Mesopotamia. *Curr. Anthropol.* **38**, 517–535 (1997).

- 702 41. Bar-Matthews, M. & Ayalon, A. Mid-Holocene climate variations revealed by  
703 high-resolution speleothem records from Soreq Cave, Israel and their correlation  
704 with cultural changes. *The Holocene* **21**, 163–171 (2011).
- 705 42. Wick, L., Lemcke, G. & Sturm, M. Evidence of Lateglacial and Holocene  
706 climatic change and human impact in eastern Anatolia: high-resolution pollen,  
707 charcoal, isotopic and geochemical records from the laminated sediments of Lake  
708 Van, Turkey. *The Holocene* **13**, 665–675 (2003).
- 709 43. Riehl, S., Pustovoytov, K. E., Weippert, H., Klett, S. & Hole, F. Drought stress  
710 variability in ancient Near Eastern agricultural systems evidenced by  $\delta^{13}\text{C}$  in  
711 barley grain. *Proc. Natl. Acad. Sci.* **111**, 12348–12353 (2014).
- 712 44. Wallace, M. P. *et al.* Stable carbon isotope evidence for Neolithic and Bronze Age  
713 crop water management in the eastern Mediterranean and southwest Asia. *PLoS*  
714 *ONE* **10**, e0127085 (2015).
- 715 45. Ferrio, J. P., Araus, J. L., Buxó, R., Voltas, J. & Bort, J. Water management  
716 practices and climate in ancient agriculture: inferences from the stable isotope  
717 composition of archaeobotanical remains. *Veg. Hist. Archaeobotany* **14**, 510–517  
718 (2005). [http://web.udl.es/usuaris/x3845331/AIRCO2\\_LOESS.xls](http://web.udl.es/usuaris/x3845331/AIRCO2_LOESS.xls).
- 719 46. Wallace, M. *et al.* Stable carbon isotope analysis as a direct means of inferring  
720 crop water status and water management practices. *World Archaeol.* **45**, 388–409  
721 (2013).
- 722 47. Styring, A. K. *et al.* Centralisation and long-term change in farming regimes:  
723 comparing agricultural practices in Neolithic and Iron Age south-west Germany.  
724 *Proc. Prehist. Soc.* (in press).
- 725 48. Vaiglova, P. *et al.* An integrated stable isotope study of plants and animals from  
726 Kouphovouno, southern Greece: a new look at Neolithic farming. *J. Archaeol.*  
727 *Sci.* **42**, 201–215 (2014).
- 728 49. Riehl, S. Archaeobotanical evidence for the interrelationship of agricultural  
729 decision-making and climate change in the ancient Near East. *Quat. Int.* **197**, 93–  
730 114 (2009).
- 731 50. Sallaberger, W. in *Administrative Documents from Tell Beydar (Seasons 1993-*  
732 *1995)* (eds. Ismail, F., Sallaberger, W., Talon, P. & Van Lerberghe, K.) 89–106  
733 (Brepols Publishers, 1996).
- 734 51. O’Leary, M. Carbon isotopes in photosynthesis. *BioScience* **38**, 328–336 (1988).

- 735 52. Charles, M. & Bogaard, A. in *Excavations at Tell Brak II: Nagar in the Third*  
736 *Millennium BC* (eds. Oates, D., Oates, J. & McDonald, H.) 301–326 (McDonald  
737 Institute, 2001).
- 738 53. Smith, A. in *Seven Generations Since the Fall of Akkad* (ed. Weiss, H.) 225–240  
739 (Harrassowitz Verlag, 2012).
- 740 54. Smith, A., Graham, P. J. & Stein, G. J. Ubaid plant use at Tell Zeidan, Syria.  
741 *Paléorient* **41**, 51–69 (2015).
- 742 55. Weiss, H. *et al.* Revising the contours of history at Tell Leilan. *Ann.*  
743 *Archéologiques Arab. Syr.* 59–74 (2002).
- 744 56. Bettencourt, L. M. A. The origins of scaling in cities. *Science* **340**, 1438–1441  
745 (2013).
- 746 57. Maekawa, K. Cultivation methods in the Ur III period. *Bull. Sumer. Agric.* **5**,  
747 115–145 (1990).
- 748 58. Bogaard, A. ‘Garden agriculture’ and the nature of early farming in Europe and  
749 the Near East. *World Archaeol.* **37**, 177–196 (2005).
- 750 59. Borgerhoff Mulder, M. B. *et al.* Intergenerational wealth transmission and the  
751 dynamics of inequality in small-scale societies. *Science* **326**, 682–688 (2009).
- 752 60. Scott, J. C. *Seeing Like a State : How Certain Schemes to Improve the Human*  
753 *Condition Have Failed.* (Yale University Press, 1998).
- 754 61. DeNiro, M. J. Postmortem preservation and alteration of in vivo bone collagen  
755 isotope ratios in relation to palaeodietary reconstruction. *Nature* **317**, 806–809  
756 (1985).
- 757 62. Vaiglova, P., Snoeck, C., Nitsch, E., Bogaard, A. & Lee-Thorp, J. A. Impact of  
758 contamination and pre-treatment on stable carbon and nitrogen isotopic  
759 composition of charred plant remains. *Rapid Commun. Mass Spectrom.* **28**, 2497–  
760 2510 (2014).
- 761 63. Bronk Ramsey, C. Radiocarbon dating: revolutions in understanding.  
762 *Archaeometry* **50**, 249–275 (2008).
- 763 64. Longin, R. New method of collagen extraction for radiocarbon dating. *Nature*  
764 **230**, 241–242 (1971).
- 765 65. Kragten, J. Tutorial review. Calculating standard deviations and confidence  
766 intervals with a universally applicable spreadsheet technique. *Analyst* **119**, 2161–  
767 2165 (1994).

- 768 66. Nitsch, E. K., Charles, M. & Bogaard, A. Calculating a statistically robust  $\delta^{13}\text{C}$   
769 and  $\delta^{15}\text{N}$  offset for charred cereal and pulse seeds. *Sci. Technol. Archaeol. Res.* **1**,  
770 1–8 (2015).
- 771 67. White, J. W. C. & Vaughn, B. H. University of Colorado, Institute of Arctic and  
772 Alpine Research (INSTAAR), Stable isotopic composition of atmospheric carbon  
773 dioxide ( $^{13}\text{C}$  and  $^{18}\text{O}$ ) from the NOAA ESRL carbon cycle cooperative global air  
774 sampling network, 1990-2012, Version: 2013-04-05, Path:  
775 ftp://ftp.cmdl.noaa.gov/ccg/co2c13/flask/event/. (2011).
- 776 68. Bar-Matthews, M. & Ayalon, A. in *Past Climate Variability through Europe and*  
777 *Africa* (eds. Battarbee, R., Gasse, F. & Stickley, C.) 363–391 (Kluwer Academic  
778 Publishers, 2004).
- 779 69. Aggarwal, P. K. *et al.* Proportions of convective and stratiform precipitation  
780 revealed in water isotope ratios. *Nat. Geosci.* **9**, 624–629 (2016).
- 781 70. Stevens, L. R., Wright, H. E. & Ito, E. Proposed changes in seasonality of climate  
782 during the Lateglacial and Holocene at Lake Zeribar, Iran. *The Holocene* **11**, 747–  
783 755 (2001).
- 784 71. Stevens, L. R., Ito, E., Schwab, A. & Wright Jr., H. E. Timing of atmospheric  
785 precipitation in the Zagros Mountains inferred from a multi-proxy record from  
786 Lake Mirabad, Iran. *Quat. Res.* **66**, 494–500 (2006).
- 787 72. Clarke, J. *et al.* Climatic changes and social transformations in the Near East and  
788 North Africa during the ‘long’ 4th millennium BC: A comparative study of  
789 environmental and archaeological evidence. *Quat. Sci. Rev.* **136**, 96–121 (2016).
- 790 73. Fernandes, R., Nadeau, M.-J. & Grootes, P. M. Macronutrient-based model for  
791 dietary carbon routing in bone collagen and bioapatite. *Archaeol. Anthropol. Sci.*  
792 **4**, 291–301 (2012).
- 793 74. Steele, K. W. & Daniel, R. M. Fractionation of nitrogen isotopes by animals: a  
794 further complication to the use of variations in the natural abundance of  $^{15}\text{N}$  for  
795 tracer studies. *J. Agric. Sci.* **90**, 7–9 (1978).
- 796 75. Akkermans, P. M. M. G. (ed.) *Tell Sabi Abyad – The Late Neolithic Settlement:*  
797 *Report on the Excavations of the University of Amsterdam (1998) and the*  
798 *National Museum of Antiquities Leiden (1991–1993) in Syria.* (Nederlands  
799 Instituut voor het Nabije Oosten, 1996).
- 800 76. Reichel, C. Urbanism and warfare: the 2005 Hamoukar, Syria, excavations.  
801 *Orient. Inst. News & Notes* **189**, 1–11 (2006).

802 **Corresponding author.** Correspondence and requests for materials to Amy Styring  
803 or Amy Bogaard.

804

805 **Acknowledgements.** The work reported here was funded by the European Research  
806 Council (AGRICURB project, grant no. 312785, PI Bogaard) and the Natural  
807 Environment Research Council (NERC standard grant NE/E003761/1, PI Bogaard). A  
808 portion of the human isotope data from Tell Brak has been obtained with the financial  
809 support by the Polish National Science Centre, grant No. 2012/06/M/HS3/00272.  
810 Archaeobotanical analyses at Tell Sabi Abyad were funded by the “Consolidating  
811 Empire” project at Leiden University (ERC Starting Grant, no. 282785, PI Düring).  
812 Archaeobotanical analyses at Tell Leilan and Tell Zeidan were funded by an NSF  
813 Early Faculty CAREER Award (1054938) granted to Alexia Smith. We are grateful to  
814 Caroline Montrieux and Edward Wilman for processing archaeobotanical samples and  
815 faunal bone collagen for isotope analysis.

816

817 **Author contributions.** A.B. conceived the study and contributed to data  
818 interpretation and the writing of the manuscript; A.K.S. designed the sampling  
819 protocol, carried out analyses, analysed the data and wrote the paper with A.B.; M.C.,  
820 F.F., M.M.H. and A.S. contributed botanical material and data; A.M., G.S. and H.W.  
821 contributed data and gave permission for analysis of material; R.M., A.K.P. and  
822 J.A.W. contributed faunal material and data; G.K.N. led the statistical analysis and  
823 developed the statistical models; M.C.P. and A.So. contributed human bone and  
824 dentine material and data. All authors discussed the results and implications and  
825 commented on the manuscript at all stages.

826

827

828 **Figures**

829

830 **Figure 1. Geographical location of the study area. (a)** Overview of northern  
831 Mesopotamia, with the locations of the palaeoclimate records of Soreq Cave, Lake  
832 Van, Lake Zeribar and Lake Mirabad marked. **(b)** Location of the archaeological sites  
833 Tell Sabi Abyad, Tell Zeidan, Tell Brak, Hamoukar and Tell Leilan included in this  
834 study. Annual rainfall data are derived from interpolation of average monthly climate  
835 data for 1960–1990, available from the WorldClim database <sup>29</sup>.

836

837 **Figure 2. Modern cereal grain  $\delta^{15}\text{N}$  values plotted against the natural log of**  
838 **mean annual rainfall, colour coded by manuring level.** Lines represent a fitted  
839 linear model relating cereal grain  $\delta^{15}\text{N}$  values to mean annual rainfall for each  
840 manuring level. Annual rainfall data are derived from interpolation of average  
841 monthly climate data for 1960–1990, available from the WorldClim database <sup>29</sup>.

842

843 **Figure 3. Archaeological cereal grain sample  $\delta^{15}\text{N}$  values plotted against date.**  
844 Symbol shape varies with site and symbols are colour coded by crop taxon. Points  
845 outlined in black come from a single storage context. Dating of the crop samples is  
846 based on stratigraphic relationships to radiocarbon-dated contexts. More details of the  
847 samples are given in Supplementary Table 1.

848

849 **Figure 4. The probability of an archaeological cereal grain sample having a**  
850 **manuring level  $m$  or lower plotted against site size, where (a)  $m = \text{low}$  and (b)  $m$**   
851  **$= \text{medium (right)}$ .** Symbol shape varies with site (see Figure 3 for legend). Points  
852 give the posterior probability for a given cereal grain sample to have a manuring level  
853 derived by multiple-imputation that is lower than  $m$ . The points are imputed in the  
854 first phase of the inference using a normal linear model regressing  $\delta^{15}\text{N}$  on the natural  
855 log of rainfall and manuring level. Boxes give the quartiles of the fitted posterior  
856 probabilities in the proportional-odds regression of manuring level on site size. The  
857 fitted values are offset by site-dependent random effects. Lines show the expected  
858 posterior probability that a cereal grain sample at a particular site size has manure  
859 level  $m$  or lower, and displays the decrease in manuring intensity as site size  
860 increases. See Supplementary Information for more detail.

861

862 **Figure 5. Archaeological cereal grain and pulse sample  $\Delta^{13}\text{C}$  values plotted**  
863 **against date. (a)** Barley and *Aegilops* grain samples. **(b)** Wheat grain and pulse seed  
864 samples. Symbol shape corresponds to the site and symbols are colour coded by crop  
865 taxon. Points outlined in black come from a single storage context. Dashed horizontal  
866 lines indicate the suggested 'boundaries' between  $\Delta^{13}\text{C}$  ranges indicative of crops  
867 grown under poorly (low  $\Delta^{13}\text{C}$ ), moderately, and well (high  $\Delta^{13}\text{C}$ ) watered conditions,  
868 based on the analysis of present-day crops <sup>cf. 46</sup>. *Aegilops* are plotted with barley  
869 grains because they were found in a barley store and are therefore assumed to have  
870 grown in the same fields as the barley.

871  
872 **Figure 6. Human and faunal bone collagen and crop  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values plotted**  
873 **in relation to ellipses representing the expected distributions (mean  $\pm$  2 standard**  
874 **deviations) of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of individuals consuming various dietary**  
875 **combinations of cereal grains, pulses and animal products (milk and/or meat).**  
876 **(a)** Late Chalcolithic 2 Tell Brak (c. 4200–3900 BC); **(b)** Late Chalcolithic 3–4 Tell  
877 Brak (c. 3900–3300 cal BC); **(c)** Early Bronze Age Tell Brak (c. 3000–2000 cal BC);  
878 and **(d)** Early Bronze Age Tell Leilan (c. 2600–2000 cal BC).


879  
880  
881  
882  
883  
884  
885  
886


887 **Table 1.** Details of archaeological sites, including location, chronology, settlement size and sample details. Present-day annual rainfall is derived  
888 from interpolation of average monthly climate data for 1960–1990, available from the WorldClim database <sup>29</sup>. The date range of each  
889 archaeological phase is based on stratigraphic dating and radiocarbon ages.


Site	Location (latitude N, longitude E)	Present-day annual rainfall (mm)	Archaeological phase	Date (cal BC)	Settlement size (ha)	Summary of contexts	References
Tell Sabi Abyad	39.09, 36.50	280	Early Pottery Neolithic–Halaf	6700–5850	1	Domestic fills, storage bins	75
Tell Zeidan	35.94, 39.08	182	Ubaid–LC2	5300–3850	12	Pyrotechnic features and domestic contexts	54
Tell Brak	36.67, 41.06	363	LC2	4200–3900	55	Mix of workshops, storage, industrial features and monumental buildings	32, 33
			LC3–4	3900–3600	130	Public building, private households and courtyards	32, 33
			LC4–5	3600–3000	45	Large house with southern Late Uruk ceramics	32
			EJ 0	3000–2900	45	Pit cutting LC4–5 house	52
			EJ III–IV	2500–2100	70	Domestic quarters within a 'high status' household	
			EJ V	2100–2000	45	Potentially 'public building'	
Hamoukar	36.81, 41.96	445	LC	3800–3500	15	Area B: tripartite buildings, large ovens	76
Tell Leilan	36.96, 41.51	446	EJ II (Leilan IIIId)	2700–2600	90	Acropolis northwest public stores; Lower Town South residential buildings	53, 55
			EJ III (Leilan IIa)	2600–2300	90	Lower Town South residential buildings	
			EJ IV (Leilan IIb)	2300–2230	90	Lower Town South residential buildings; Acropolis northwest Akkadian palace	
			EJ V (Leilan IIc)	2230–2200	0.1	Acropolis post-Akkadian four-room house	

890 **Table 2.** Details of site location, mean annual rainfall and manuring regimes for modern crop samples. Annual rainfall is derived from  
 891 interpolation of average monthly climate data for 1960–1990, available from the WorldClim database <sup>29</sup>.


Site	Location	Country	Location (latitude N, longitude E)	Annual rainfall (mm)	Year of collection	Crop species	Manuring regimes	No. of plots
Askov	South Jutland	Denmark	55.53, 9.09	838	2007 and 2008	Two-row hulled barley, bread wheat, emmer, spelt	Low and high	28
Sutton Bonington	Nottinghamshire	UK	52.82, -1.25	632	2007 and 2008	Einkorn, emmer, spelt	Low, medium and high	55
Rothamsted Research	Hertfordshire	UK	51.8, -0.36	655	1852- 2004	Two-row hulled barley, bread wheat	Low and high	22
Bad Lauchstädt	Leipzig-Halle	Germany	51.39, 11.83	503	2007 and 2008	Two-row hulled barley, bread wheat	Low, medium and high	11
Sighisoara region	Transylvania	Romania	46.41, 24.92	641	2008	Bread wheat, einkorn	Low and medium	14
Haute Provence	Lubéron/Sault region	France	44.02, 5.42	897	2013	Bread wheat, einkorn	Low	19
Lena district	Asturias	Spain	43.03, -5.76	964	2007	Spelt	Low, medium and high	16
Kastamonu	Kastomonu	Turkey	41.38, 33.70	570	2008	Durum wheat, einkorn, emmer	Low and medium	8
ICARDA	Aleppo	Syria	36.01, 36.93	419	2008	Bread wheat	Low, medium and high	42
Bellota	Ouezzane	Morocco	34.95, -5.54	703	2014	Two-row hulled barley, bread wheat	Low	6
Wadi ibn Hammad	Kerak	Jordan	31.30, 35.63	186	2007	Durum wheat	Low and medium	10
Tighirt	Sidi Ifni	Morocco	29.35, -9.43	272	2014	Two-row hulled barley	Medium	15
Amtoudi (oasis)	Guelmim	Morocco	29.24, -9.19	194	2014	Two-row hulled barley	High	11
Amtoudi (decrue)	Guelmim	Morocco	29.24, -9.19	194	2015	Two-row hulled barley, bread wheat	Low	16









(a)




(b)




(a)



(b)





- Crops**
- Barley
  - △ Bread wheat
  - Free-threshing wheat
  - × Emmer
  - × Einkorn
  - \* Glume wheat
  - △ Lentil
  - ⊠ Pea
  - ⊠ Grass pea
  - ⊠ Aegilops

- Fauna**
- ◇ Gazelle
  - Cattle
  - Sheep
  - ◇ Goat
  - Pig

- Dietary reconstruction**
- Human
  - 100% animal protein consumption
  - 100% cereal grain consumption
  - 100% pulse consumption
  - ▨ 80% animal protein consumption
  - ▨ 50% animal protein consumption
  - ▨ 20% animal protein consumption