

Fine scales of decay of operator semigroups

Charles J.K Batty Ralph Chill Yuri Tomilov

February 2, 2014

Abstract

Motivated by potential applications to partial differential equations, we develop a theory of fine scales of decay rates for operator semigroups. The theory contains, unifies, and extends several notable results in the literature on decay of operator semigroups and yields a number of new ones. Its core is a new operator-theoretical method of deriving rates of decay combining ingredients from functional calculus, and complex, real and harmonic analysis. It also leads to several results of independent interest.

Contents

1	Introduction	2
2	Preliminaries on some classes of functions	11
3	Functional calculus of sectorial operators	19
4	Some estimates for semigroup asymptotics	27
5	Singularity at infinity	33
6	Singularity at zero: general results	49
7	Singularity at zero: polynomial and regularly varying rates	66
8	Singularities at both zero and infinity	73

Batty: St. John's College, Oxford OX1 3JP, United Kingdom; e-mail: charles.batty@sjc.ox.ac.uk

Chill: Technische Universität Dresden, Institut für Analysis, 01062 Dresden, Germany; e-mail: ralph.chill@tu-dresden.de

Tomilov: Department of Mathematics and Computer Science, Nicolas Copernicus University, ul. Chopina 12/18, 87-100 Torun, Poland and Institute of Mathematics, Polish Academy of Sciences, Śniadeckich str.8, 00-956 Warsaw, Poland; e-mail: tomilov@mat.uni.torun.pl

Mathematics Subject Classification (2010): Primary 47D06; Secondary 34D05 34G10

1 Introduction

1.1 Historical background, motivation and sample results

The study of stability of solutions of the abstract Cauchy problem

$$\begin{cases} \dot{u}(t) + Au(t) = 0, & t \geq 0, \\ u(0) = x, & x \in X, \end{cases}$$

where $-A$ is the generator of a C_0 -semigroup $(T(t))_{t \geq 0}$ on a Banach space X , is a classical subject of functional analysis having numerous applications to partial differential equations. Namely, asymptotic stability (or simply: stability), that is convergence of all orbits of $(T(t))_{t \geq 0}$ to zero, and exponential stability, that is, convergence of all orbits of $(T(t))_{t \geq 0}$ to zero with exponential rate, are two of the main building blocks of stability theory. Since the resolvent of the generator is often easier to compute than the semigroup, it is traditional and efficient to use the resolvent when dealing with both kinds of stability, and actually a number of resolvent criteria for stability are known in this context (see [6, Chapter 5], [20] and [56]).

Recently, the study of various PDEs revealed that the resolvent can also be used successfully in order to treat intermediate rates of convergence, thus distinguishing and quantifying fine modes of convergence to zero. This has become especially transparent for the damped wave equation

$$\begin{aligned} u_{tt} + a(x)u_t - \Delta u &= 0 \quad \text{in } \mathbb{R}_+ \times M, \\ u &= 0 \quad \text{in } \mathbb{R}_+ \times \partial M, \\ u(0, \cdot) &= u_0 \quad \text{in } M, \\ u_t(0, \cdot) &= u_1 \quad \text{in } M, \end{aligned} \tag{1.1}$$

which is one of the basic models in control theory. Here M is a smooth, compact, connected Riemannian manifold, $a \in L^\infty(M)$ and $a \geq 0$. The wave equation can be rewritten as a first order Cauchy problem in the Hilbert space $X = H_0^1(M) \times L^2(M)$ with A defined by

$$\begin{aligned} \text{dom}(A) &= (H^2(M) \cap H_0^1(M)) \times H_0^1(M), \\ A &= \begin{pmatrix} 0 & -I \\ -\Delta & a \end{pmatrix}. \end{aligned}$$

The operator A is invertible, and $-A$ generates a non-analytic contraction semigroup $(T(t))_{t \geq 0}$. Since the natural norm on the Hilbert space X corresponds to the energy of the system, any estimate for the rate of decay of the semigroup is an estimate for the rate of decay of the energy of the system.

Recall that a semigroup is exponentially stable if and only if $\|T(t)\| = O(r(t))$ with $\lim_{t \rightarrow \infty} r(t) = 0$. Hence, one cannot expect a uniform rate of decay for all solutions if the semigroup is only stable but not exponentially stable. Nevertheless, one may have uniform estimates of rates of decay for a dense set of initial values. The study of the

damped wave equation and various similar PDEs has revealed that the resolvent can be used successfully to obtain quantitative rates of convergence of the form

$$\|T(t)A^{-1}\| = O(r(t)), \quad t \rightarrow \infty, \quad (1.2)$$

that is, uniform rates for *classical* solutions.

In his pioneering work [51], Lebeau discovered that the spectrum of A is contained in the strip $\{\lambda \in \mathbb{C} : 0 < \operatorname{Re} \lambda \leq 2\|a\|_\infty\}$, the resolvent $(\lambda + A)^{-1}$ admits an exponential estimate on the imaginary axis and this forces the energy of solutions of (1.1) to decay at least with rate

$$r(t) = \frac{\log(3 + \log(3 + t))}{\log(3 + t)}, \quad t \rightarrow \infty.$$

His method was extended subsequently by Burq [18] and Lebeau and Robbiano [52] to cover similar situations involving local energy decay for (1.1) where $a = 0$ and M is a non-compact manifold (“scattering in an exterior domain”), and global energy decay for (1.1) with Neumann boundary conditions, respectively. Moreover Lebeau’s estimate was improved in [18] and [52] to $1/\log(3 + t)$. These results gave rise to a number of papers treating the rates of decay of solutions to PDEs by abstract semigroup methods. Among them, we would like to mention [9], [16], [19], [21], [22], [53], and especially the very recent paper [4] as samples. This last paper includes a complete historical account, and a detailed discussion of damped wave equations in the resolvent context.

The study of rates was put into the setting of Tauberian theorems for Laplace transforms in [13], by regarding the resolvent as the Laplace transform of the semigroup and partially inverting the Laplace transform. This approach by pure complex analysis unified a number of known results (including those of Burq, Lebeau and others), improved some of them and applied to arbitrary growth of resolvents. In particular, the following result was obtained.

Theorem 1.1. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Banach space X with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R}$ is empty, and define*

$$\begin{aligned} M(s) &= \sup\{\|(ir + A)^{-1}\| : |r| \leq s\}, \quad s \geq 1, \\ M_{\log}(s) &= M(s) (\log(1 + M(s)) + \log(1 + s)). \end{aligned}$$

Then there exist positive constants C, C', c, c' such that

$$\frac{c'}{M^{-1}(C't)} \leq \|T(t)A^{-1}\| \leq \frac{C}{M_{\log}^{-1}(ct)} \quad (1.3)$$

for all sufficiently large t .

Note that by [13, Proposition 1.3] any decay of $\|T(t)A^{-1}\|$ to zero as $t \rightarrow \infty$ implies that $\sigma(A) \cap i\mathbb{R}$ is empty. However, in many situations the spectrum approaches the imaginary axis asymptotically forcing the function M to grow at infinity. So we shall say (slightly loosely) that if $M(s) \rightarrow \infty$ as $s \rightarrow \infty$, then the resolvent, restricted to the imaginary axis, has a *singularity at infinity*.

The upper estimate in (1.3) was established by examining a function-theoretic method first used in unquantified form in [5] (see also [11]). The gap between the lower and upper estimates is, in general, of “logarithmic size” and any improvement of (1.3) should fall within that gap. (Note, however, that if M grows exponentially, then the lower and upper estimates are of the same order.) It was conjectured in [13] that the gap could be bridged in the case of Hilbert spaces, but one cannot expect rates better than $(M_{\log}^{-1}(ct))^{-1}$ for general Banach spaces. This conjecture was partially settled in [17]. It was proved in [17] that if M grows polynomially then it is possible to obtain a characterization of decay of $\|T(t)A^{-1}\|$ which is *optimal* in the sense that M_{\log}^{-1} in (1.3) is replaced by M^{-1} . Namely the following theorem holds.

Theorem 1.2. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Hilbert space X , with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R}$ is empty, and fix $\alpha > 0$. Then the following statements are equivalent:*

- (i) $\|(is + A)^{-1}\| = O(|s|^\alpha), \quad |s| \rightarrow \infty.$
- (ii) $\|T(t)A^{-1}\| = O(t^{-1/\alpha}), \quad t \rightarrow \infty.$

Moreover, it was shown in [17] that the logarithmic gap in (1.3) is unavoidable for semigroups on Banach spaces, even if the growth of the resolvent is polynomial as in Theorem 1.2(i).

The result above has already found a number of applications to the study of concrete PDEs, see e.g. [1], [2], [3], [8], [26], [29], [30], [32], [33], [34], [37], [58], [59], [63], [66]. In view of its importance, Theorem 1.2 (and Theorem 1.1) raised the natural and important problem of obtaining similar results for growth scales which are finer than polynomial ones, or showing that for other scales results of this kind cannot be true.

Recall that the optimal function-theoretical analogue of Theorem 1.1 has a logarithmic correction term as in the right-hand side of (1.3) [17, Theorem 3.8]. Thus, one cannot expect, in general, that a purely function-theoretical approach could produce a version of Theorem 1.2 for more general resolvent bounds, and the need for an alternative approach to the problem of sharpening (1.3) becomes apparent. Such an approach based on operator theory is proposed in this paper. Note that Theorem 1.2 itself is proved in [17] by means of an auxiliary operator construction. The framework of the present paper is different, more general, and in a sense more transparent.

The polynomial scale has a number of special (e.g. algebraic) properties making it comparatively amenable to an operator-theoretical approach. For other scales where these properties are not available, our approach must be much more involved and the task of finding appropriate scales is rather non-trivial. The scale which is closest to the polynomial one is the finer scale of regularly varying functions, which are products of polynomials and slowly varying functions. Such scales have been widely used as natural refinements of polynomial scales in various areas of analysis, including number theory, complex analysis and probability theory (see e.g. [15, Sections 6-8]). This suggests that if there is a generalization of Theorem 1.2 to more general classes of resolvent bounds, the scale of regularly varying functions should be one of the first candidates for such a class.

In this paper, we develop a general operator-theoretical approach to the study of *optimal* decay rates of operator semigroups within the fine scales of regularly varying functions. Since our primary motivation comes from the study of rates in (1.2) (the case of singularity at infinity), we first formulate several partial cases of our results obtained below for the setting of (1.2). We use the fact that the function $(\log s)^\beta$ is slowly varying with de Bruijn conjugate $(\log s)^{-\beta}$ (see Theorem 5.6, Corollary 5.7 and Theorem 5.12 below).

Theorem 1.3. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Hilbert space X , with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R}$ is empty, and fix $\alpha > 0$ and $\beta \geq 0$.*

(a) *The following statements are equivalent:*

- (i) $\|(is + A)^{-1}\| = O(|s|^\alpha (\log |s|)^{-\beta}), \quad |s| \rightarrow \infty.$
- (ii) $\|T(t)A^{-1}\| = O(t^{-1/\alpha} (\log t)^{-\beta/\alpha}), \quad t \rightarrow \infty.$

(b) *If*

$$\|(is + A)^{-1}\| = O(|s|^\alpha (\log |s|)^\beta), \quad |s| \rightarrow \infty,$$

then, for every $\varepsilon > 0$,

$$\|T(t)A^{-1}\| = O(t^{-1/\alpha} (\log t)^{\varepsilon + \beta/\alpha}), \quad t \rightarrow \infty.$$

We do not know whether (b) is true for $\varepsilon = 0$. If so, the implication would become an equivalence.

It is apparent that our methodology goes beyond clarifying (1.2) and it allows one to deal with rates of decay of several other operator families stemming from bounded C_0 -semigroups on Hilbert spaces. Thus it is significant from the point of view of both abstract operator theory and applications.

Observe that if $(T(t))_{t \geq 0}$ is a bounded C_0 -semigroup on a Hilbert space X with generator $-A$, then for each $x \in X$ the function $-T(\cdot)A^{-1}x$ is a primitive of $T(\cdot)x$. Now instead of considering asymptotics of the primitives of $T(\cdot)x$, we may study asymptotics of the derivatives of $T(\cdot)x$. In other words, we may consider the orbits of the form $T(\cdot)Ax$ for $x \in \text{dom}(A)$, and so study long-time regularity of $(T(t))_{t \geq 0}$. This type of asymptotic behaviour of semigroups has not been treated systematically in the literature so far. Some related but very partial results pertaining to orbits of analytic semigroups were obtained in [25]. More results are available in the discrete setting, see e.g. [24], [44], [45] and [57]. In this paper, the decay of $T(\cdot)x$ for $x \in \text{ran}(A)$ is studied systematically and the resulting structure appears to be very similar to the one established in the study of decay rates for the orbits $(T(t))_{t \geq 0}$ starting from the domain of A and discussed above.

Since we are interested in a decay of $T(t)Ax$ to zero that is uniform with respect to $x \in \text{dom}(A)$, the problem which we address in this case is to quantify the decay of $\|T(t)A(I+A)^{-1}\|$. Thus our task is to identify spectral conditions on A and a corresponding ‘‘optimal’’ function r such that

$$\|T(t)A(I+A)^{-1}\| = O(r(t)), \quad t \rightarrow \infty. \quad (1.4)$$

Remark that if $\|T(t)A(I+A)^{-1}\| \rightarrow 0$ as $t \rightarrow \infty$, then the spectrum of A does not meet $i\mathbb{R} \setminus \{0\}$ and the resolvent of A is bounded on $i(\mathbb{R} \setminus (-1, 1))$, see Theorem 6.10 below. Therefore the only singularity of the resolvent on the imaginary axis may be at zero and we have a spectral situation which is in a sense opposite to the situation considered above in the study of (1.2). We describe this situation by saying that the resolvent restricted to the imaginary axis has a *singularity at zero*. Basic examples in the framework of a singularity at zero are provided, in particular, by generators of bounded eventually differentiable semigroups, for example those arising in the study of delay differential equations [12], [27, Section VI.6].

To treat (1.4), it is natural to assume the boundedness of $(\lambda + A)^{-1}$ outside a neighbourhood of zero in $i\mathbb{R}$, and to relate the decay of $\|T(t)A(I+A)^{-1}\|$ to the growth of $(\lambda + A)^{-1}$ near zero. The first problem that we encountered on this way was that the mere convergence of $\|T(t)A(I+A)^{-1}\|$ to zero was wide open. This type of convergence can be considered as an extension of the famous Katznelson-Tzafriri theorem (see Theorem 6.1) with L^1 -functions replaced by certain bounded measures on the real half-line. Using a new technique, we obtain an interesting generalization of the Katznelson-Tzafriri theorem for a class of bounded measures. Moreover, we are able to derive a partial analogue of Theorem 1.1 for the case of a singularity at zero thus equipping our version of the Katznelson-Tzafriri theorem with rates. These results are given below, see also Theorems 6.14 and 6.15.

Theorem 1.4. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Hilbert space X with generator $-A$. Assume that $E := -i\sigma(A) \cap \mathbb{R}$ is compact and of spectral synthesis, and moreover the resolvent of A is bounded on $i(\mathbb{R} \setminus (-\eta, \eta))$ for some $\eta > 0$.*

(a) *If μ is a finite measure on \mathbb{R}_+ such that its Fourier transform vanishes on E , then*

$$\lim_{t \rightarrow \infty} \|T(t)\hat{\mu}(T)\| = 0,$$

where $\hat{\mu}(T)x := \int_0^\infty T(s)x \, d\mu(s)$ for $x \in X$.

(b) *Assume that $(\lambda + A)^{-1}$ has a singularity at zero (so $E = \{0\}$), let*

$$m(s) := \sup\{\|(ir + A)^{-1}\| : |r| \geq s\}, \quad s > 0,$$

and assume that $\lim_{s \rightarrow 0^+} sm(s) = \infty$. Let $\varepsilon \in (0, 1)$. Then there exist positive constants c, C, C_ε such that

$$cm^{-1}(Ct) \leq \|T(t)A(I+A)^{-1}\| \leq C_\varepsilon m^{-1}(t^{1-\varepsilon}),$$

for all sufficiently large t .

Our technique of employing fine scales of regularly varying functions to the study of orbits decay proves to be efficient also in the situation of a singularity at zero. In particular, we obtain a counterpart of Theorem 1.3(a) in that case, given in Theorem 1.5 (see also Theorem 7.7). Unfortunately, Theorem 1.4(b) is not quite as strong as Theorem 1.1, because the correction term involves t^ε instead of a logarithmic term. As a result,

the theorem below covers only the case when the resolvent grows slightly slower than a power of $|s|^{-1}$ (in analogy with Theorem 1.3(a)). The problem of characterizing the decay of $\|T(t)A(I+A)^{-1}\|$ in the other case when the resolvent grows slightly faster than a power of $|s|^{-1}$ remains open.

Theorem 1.5. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Hilbert space, with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R} \subset \{0\}$, and let $\alpha > 1$, $\beta \geq 0$. The following statements are equivalent.*

- (i) $\|(is + A)^{-1}\| = \begin{cases} O(|s|^{-\alpha}(\log(1/|s|))^{-\beta}), & s \rightarrow 0, \\ O(1), & |s| \rightarrow \infty. \end{cases}$
- (ii) $\|T(t)A(I+A)^{-1}\| = O(t^{-1/\alpha}(\log t)^{-\beta/\alpha}), \quad t \rightarrow \infty.$

Note that Theorem 1.5 for $\beta = 0$ provides a resolvent characterization of polynomial rates of decay for $\|T(t)A(I+A)^{-1}\|$. This characterization holds for $\alpha = 1$, too. The case when $0 < \alpha < 1$ does not arise since $\|(is + A)^{-1}\| \geq |s|^{-1}$ if $0 \in \sigma(A)$. For other results revealing properties of decay rates for $\|T(t)A(I+A)^{-1}\|$ we refer to Sections 6 and 7.

The operator-theoretical approach of the present paper can also be used successfully to treat decay rates of orbits of $(T(t))_{t \geq 0}$ starting in $\text{dom}(A) \cap \text{ran}(A)$, combining the situations of a singularity at infinity and a singularity at zero considered above. Since $\text{dom}(A) \cap \text{ran}(A) = \text{ran}(A(I+A)^{-2})$ (Proposition 3.10) and since we are interested in uniform rates of decay, the task of characterizing such rates can be considered as the task of characterizing the property

$$\|T(t)A(I+A)^{-2}\| = O(r(t)), \quad t \rightarrow \infty, \quad (1.5)$$

in resolvent terms.

While (1.5) implies that $\sigma(A) \cap i\mathbb{R} \subset \{0\}$, it does not exclude growth of the resolvent near zero and near infinity (along the imaginary axis). At the same time, (1.5) imposes certain restrictions on the resolvent growth (see Theorem 8.1) which might serve as a starting point for obtaining *optimal* decay rates in (1.5). The following result (see Theorem 8.4) illustrates that point. It is a generalization of Theorem 1.2 above.

Theorem 1.6. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Hilbert space X with generator $-A$, and assume that $\sigma(A) \cap i\mathbb{R} = \{0\}$. If there exist $\alpha \geq 1$ and $\beta > 0$ such that*

$$\|(is + A)^{-1}\| = \begin{cases} O(|s|^{-\alpha}), & s \rightarrow 0, \\ O(|s|^\beta), & |s| \rightarrow \infty, \end{cases} \quad (1.6)$$

then

$$\|T(t)A^\alpha(I+A)^{-(\alpha+\beta)}\| = O(t^{-1}), \quad t \rightarrow \infty,$$

and

$$\|T(t)A(I+A)^{-2}\| = O(t^{-1/\gamma}), \quad (1.7)$$

where $\gamma = \max(\alpha, \beta)$.

Conversely, if (1.7) holds for some $\gamma > 0$, then (1.6) holds for $\alpha = \max(1, \gamma)$ and $\beta = \gamma$.

After this review of the main results of this paper, we note that we derive as by-products a number of results of independent interest. These include abstract converses to interpolation inequalities (Theorem 4.3), estimates of decay rates for semigroup orbits with bounded local resolvents (Theorem 4.7), an extension of the Katznelson-Tzafriri theorem to the setting of measure algebras (Theorem 6.14), and lower bounds for orbit decay in the Banach space setting (Corollaries 6.11 and 8.2).

1.2 Strategy

One of the main novelties of the paper is its operator-theoretical method for deriving estimates for rates of decay which in many cases happen to be sharp. Let us describe the method in some more detail.

If $(T(t))_{t \geq 0}$ is a bounded C_0 -semigroup on a Hilbert space X with generator $-A$, then we look for decay rates of $T(t)B$ for a bounded operator B which takes one of the three forms: A^{-1} , $A(I + A)^{-1}$ and $A(I + A)^{-2}$. Given a (upper) bound M for the resolvent on the imaginary axis, we start by establishing lower bounds for decay rates of $T(t)B$ in terms of M . Such bounds are known in the case of a singularity at infinity (see (1.3)), and they are obtained in this paper in the case of a singularity at zero and in the case of singularities at both zero and infinity (Corollaries 6.11 and 8.2).

To obtain an upper bound for decay rates of $\|T(t)B\|$ matching the lower bound mentioned above we then proceed in three steps, of which only the second is restricted to Hilbert spaces. *First*, we show that if the resolvent $(\lambda + A)^{-1}$ grows regularly on the imaginary axis, then it is bounded in the right half-plane when restricted to the range of an associated operator W involving a fractional power and a Bernstein function f of A , see Theorems 5.5 and 7.3. *Second*, in Theorem 4.7, we prove that if $\|(\lambda + A)^{-1}W\|$ is bounded in the right half-plane then $\|T(t)W\|$ decays like t^{-1} as $t \rightarrow \infty$. *Third*, using new abstract converses to interpolation inequalities for Bernstein functions (Theorem 4.3), we deduce that if $\|T(t)W\|$ decays like t^{-1} , then on the domain of A , or the range of A , or the intersection of the two, the decay of $(T(t))_{t \geq 0}$ is expressed in terms of f . In the context of Theorem 1.2 this step reduces to an application of the moment inequality for fractional powers. For the finer scales of rates of decay, the argument is much more subtle and its effect is to improve some a priori bounds for the semigroup. The decay obtained in this way matches the lower bound in many cases and then it is optimal (apart from Theorem 5.12 where the upper bound differs from the lower bound by an arbitrarily small power of a logarithm).

Operator Bernstein functions were used successfully in [31] to deal with rates in mean ergodic theorems for bounded C_0 -semigroups on Banach spaces. While there are formal similarities between [31] and the present paper, the problems treated here are much more involved and technically and ideologically demanding. While the mean ergodic theorem for C_0 -semigroups is a comparatively simple statement, the majority of stability conditions for C_0 -semigroups are deep results with tricky proofs. This extends to the framework of the study of rates.

Our approach establishes a certain structure for dealing with rates in three cases, when the resolvent of the semigroup generator has singularities at infinity, or at the origin, or at both of them. While the structure has many elements in common between the three

cases, there are several essential differences between them which have to be addressed separately.

In this paper the approach is applied to the class of regularly varying rates which are close to polynomial rates. When the resolvent grows rapidly and fairly regularly, the upper and lower bounds in Theorem 1.1 are of the same order. However there are some rates which grow regularly but are intermediate between polynomial and exponential, where the optimal rate of decay of $\|T(t)A^{-1}\|$ in Theorem 1.1 is not established for semigroups on Hilbert space. For very irregular (but arbitrarily fast) rates M of growth of the resolvent it is not possible to improve Theorem 1.1 by replacing M_{\log}^{-1} by M^{-1} , even for semigroups of normal operators on Hilbert spaces (see Propositions 5.1 and 6.13).

1.3 Notation and conventions

In this paper, X will be a complex Banach space, and will often be specified to be a Hilbert space. We let $\mathcal{L}(X)$ denote the space of all bounded linear operators on Banach space X , and the identity operator will be denoted by I . If A is a linear operator on X , we denote the domain of A by $\text{dom}(A)$, the range of A by $\text{ran}(A)$, the spectrum of A by $\sigma(A)$ and the resolvent set by $\rho(A)$. If A is closable, its closure is written as \overline{A} .

If B is another linear operator on X , then we take $A + B$ and AB to be the operators with

$$\begin{aligned} \text{dom}(A + B) &= \text{dom}(A) \cap \text{dom}(B), \\ \text{dom}(AB) &= \{x \in \text{dom}(B) : Bx \in \text{dom}(A)\}. \end{aligned}$$

A complex variable may be denoted by either z or λ . We shall use the symbol ι to denote the identity function on domains in \mathbb{C} . The closure of a subset E of \mathbb{C} will be denoted by \overline{E} .

For $\varphi \in (0, \pi]$ we shall let $\Sigma_\varphi := \{\lambda \in \mathbb{C} : |\arg \lambda| < \varphi\}$ be the sector of angle φ in \mathbb{C} . Note that Σ_π is the slit plane $\mathbb{C} \setminus (-\infty, 0]$. We may write

$$\begin{aligned} \mathbb{C}_+ &:= \Sigma_{\pi/2} = \{\lambda \in \mathbb{C} : \text{Re } \lambda > 0\}, \\ \mathbb{R}_+ &:= [0, \infty). \end{aligned}$$

We shall consider integrals of functions f with respect to positive Radon measures μ over $(0, \infty)$ or $[0, \infty)$. We shall write such integrals as

$$\int_{0+}^{\infty} f(s) \, d\mu(s) \quad \text{or} \quad \int_0^{\infty} f(s) \, d\mu(s),$$

respectively. If g is an increasing right-continuous function on $(0, \infty)$ and μ is the associated Lebesgue-Stieltjes measure, we shall write

$$\int_{0+}^{\infty} f(s) \, dg(s)$$

instead of

$$\int_{0+}^{\infty} f(s) \, d\mu(s).$$

Let $M^b(\mathbb{R})$ denote the space of all complex Borel measures of bounded variation on \mathbb{R} . For $a \in \mathbb{R}$, we let δ_a be the Dirac measure at a .

We consider the spaces $L^1(\mathbb{R}_+)$ and $M^b(\mathbb{R}_+)$ as (closed) subspaces of $L^1(\mathbb{R})$ and $M^b(\mathbb{R})$, respectively, by extending functions or measures on \mathbb{R}_+ by 0 on $(-\infty, 0)$. In addition, we view $L^1(\mathbb{R})$ as a closed subspace of $M^b(\mathbb{R})$. The standard convolution of two measures $\mu_1, \mu_2 \in M^b(\mathbb{R})$ will be denoted by $\mu_1 * \mu_2$.

The Fourier transform of a function $f \in L^1(\mathbb{R}; X)$ is defined by

$$\mathcal{F}f(\xi) := \int_{\mathbb{R}} e^{-i\xi t} f(t) dt, \quad \xi \in \mathbb{R}.$$

This definition of the Fourier transform holds in particular for f in the Schwartz space $\mathcal{S}(\mathbb{R})$ of scalar-valued test functions and it then induces a Fourier transform for all vector-valued, tempered distributions. The Fourier transform of $\mu \in M^b(\mathbb{R})$ is therefore given by

$$\mathcal{F}\mu(\xi) := \int_{\mathbb{R}} e^{-i\xi t} d\mu(t), \quad \xi \in \mathbb{R}.$$

When X is a Hilbert space, we shall also use the symbol \mathcal{F} to denote the Fourier transform induced on the Hilbert space $L^2(\mathbb{R}; X)$, so that $(2\pi)^{-1/2}\mathcal{F}$ is a unitary operator.

We are interested in asymptotic properties of functions defined on intervals of the form $[a, \infty)$ for some $a > 0$, with values in $(0, \infty)$. We shall say that f and g are *asymptotically equivalent*, and we write $f \sim g$, or $f(s) \sim g(s)$, if

$$\lim_{s \rightarrow \infty} \frac{f(s)}{g(s)} = 1.$$

This defines an equivalence relation on such functions and we shall in effect be working with equivalence classes of functions. This viewpoint provides a justification for sometimes not including precise statements about the domains of our functions (provided that each domain contains some interval of the form (a, ∞)) or repeatedly saying that an inequality holds for all sufficiently large s .

Where we use the notation $f(s) \sim g(s)$, it will mean asymptotic equivalence as $s \rightarrow \infty$ unless otherwise specified. Occasionally we shall use the corresponding notation as $s \rightarrow 0+$, but then it will be specified. Thus, for positive functions f and g defined on $(0, a]$, the notation

$$f(s) \sim g(s), \quad s \rightarrow 0+,$$

means

$$\lim_{s \rightarrow 0+} \frac{f(s)}{g(s)} = 1.$$

For an increasing function $f : [a, \infty) \rightarrow (0, \infty)$ such that $\lim_{s \rightarrow \infty} f(s) = \infty$, the notation f^{-1} may denote the inverse function, or more generally a *right inverse*, of f , defined on the range of f , so that $f(f^{-1}(t)) = t$ for all t in the range. It may also denote an *asymptotic inverse* of f , defined on an interval $[b, \infty)$, such that

$$f^{-1}(f(s)) \sim s, \quad f(f^{-1}(s)) \sim s. \quad (1.8)$$

It should be clear from the context which notion of inverse function is involved.

We shall occasionally use the notation f^α to denote function $s \mapsto f(s)^\alpha$ when $\alpha \neq -1$, but we shall use $1/f$ to denote the reciprocal function of f , in order to avoid confusion with any inverse function. Similarly fg or $f.g$ will denote a pointwise product of two functions f and g , and $f \circ g$ will denote composition.

We shall use C and c to denote (strictly) positive constants, whose values may change from place to place.

2 Preliminaries on some classes of functions

In this section we review various classes of functions on $(0, \infty)$ with emphasis on the properties that we shall need.

2.1 Bernstein functions, complete Bernstein functions and Stieltjes functions

In this subsection, we recall the definitions and some properties of complete Bernstein functions and Stieltjes functions. Most of this material can be found in [64]. In Section 3 we shall review the operator functional calculus associated with these functions, and that will be used in later sections of the paper.

Recall that a function $f \in C^\infty(0, \infty)$ is *completely monotone* if

$$(-1)^n f^{(n)}(\lambda) \geq 0 \quad \text{for every } n \in \mathbb{N} \cup \{0\}, \lambda \in (0, \infty).$$

By Bernstein's theorem [64, Theorem 1.4], every completely monotone function f is the Laplace transform of a positive Radon measure on \mathbb{R}_+ , and f extends to a holomorphic function in the right half-plane. A function $f \in C^\infty(0, \infty)$ is called a *Bernstein function* if

$$f \geq 0 \quad \text{and} \quad f' \text{ is completely monotone.}$$

Clearly, every Bernstein function also extends to a holomorphic function in the right half-plane. By the Lévy-Khintchine representation theorem [64, Theorem 3.2], a function f is a Bernstein function if and only if there exist constants $a, b \geq 0$ and a positive Radon measure μ_{LK} on $(0, \infty)$ such that

$$\begin{aligned} \int_{0+}^{\infty} \frac{s}{s+1} d\mu_{\text{LK}}(s) < \infty, \quad \text{and} \\ f(\lambda) = a + b\lambda + \int_{0+}^{\infty} (1 - e^{-\lambda s}) d\mu_{\text{LK}}(s), \quad \lambda > 0. \end{aligned} \quad (2.1)$$

The triple (a, b, μ_{LK}) is uniquely determined by the corresponding Bernstein function f and is called the *Lévy-Khintchine triple* of f .

The class of Bernstein functions is rather large and to ensure good algebraic and function-theoretic properties of Bernstein functions it is convenient, and also sufficient for many purposes, to consider the subclass consisting of complete Bernstein functions. A function $f \in C^\infty(0, \infty)$ is called a *complete Bernstein function* if it is a Bernstein

function and the measure μ_{LK} in the Lévy-Khintchine triple has a completely monotone density with respect to Lebesgue measure [64, Definition 6.1]. By [64, Theorem 6.2], every complete Bernstein function admits a representation of the form

$$f(\lambda) = a + b\lambda + \int_{0+}^{\infty} \frac{\lambda}{s + \lambda} d\mu(s), \quad \lambda > 0, \quad (2.2)$$

for some constants $a, b \geq 0$ and some positive Radon measure μ on $(0, \infty)$ satisfying

$$\int_{0+}^{\infty} \frac{d\mu(s)}{s + 1} < \infty. \quad (2.3)$$

Remark 2.1. Complete Bernstein functions admit various representations different from (2.2). In particular, the following formula is used in some papers related to Bernstein functions (for example in [40], [41], [42], [43], [62]):

$$f(\lambda) = a + \int_0^{\infty} \frac{\lambda}{1 + \lambda t} d\nu(t) = a + \nu(\{0\})\lambda + \int_{0+}^{\infty} \frac{\lambda}{1 + \lambda t} d\nu(t), \quad (2.4)$$

where ν is a positive Radon measure on \mathbb{R}_+ satisfying

$$\int_0^{\infty} \frac{d\nu(t)}{1 + t} < \infty,$$

and the pair (a, ν) is unique.

The representations (2.4) and (2.2) are equivalent by the change of variable $s = 1/t$, with ν being the push-forward measure of μ combined with an atom of mass b at 0, and vice versa.

There are striking dualities between complete Bernstein functions and another class known as Stieltjes functions. A function $h \in C^\infty(0, \infty)$ is a *Stieltjes function* if there exist constants $a, b \geq 0$ and a positive Radon measure μ on $(0, \infty)$ satisfying (2.3) such that

$$h(\lambda) = \frac{a}{\lambda} + b + \int_{0+}^{\infty} \frac{d\mu(s)}{s + \lambda}, \quad \lambda > 0. \quad (2.5)$$

The representation formulas (2.2) (for complete Bernstein functions) and (2.5) (for Stieltjes functions) are unique, and they are called the *Stieltjes representations* for f and h , respectively; see e.g. [64, Chapter 2]. We write $f \sim (a, b, \mu)$ and $h \sim (a, b, \mu)$, and we call (a, b, μ) the *Stieltjes triple*, and μ the *Stieltjes measure* for f and h , respectively. Note that

$$a = \lim_{\lambda \rightarrow 0+} f(\lambda) = \lim_{\lambda \rightarrow 0+} \lambda h(\lambda), \quad b = \lim_{\lambda \rightarrow \infty} \frac{f(\lambda)}{\lambda} = \lim_{\lambda \rightarrow \infty} h(\lambda).$$

We shall be particularly interested in the Stieltjes (and complete Bernstein) functions with Stieltjes representations of the form $(0, 0, \mu)$. When μ is the Lebesgue-Stieltjes measure associated with an increasing right-continuous function g we shall denote the Stieltjes function with representation $(0, 0, \mu)$ by S_g , and we shall call it the *Stieltjes function associated with g* .

Note that every (complete) Bernstein function f is increasing and every Stieltjes function is decreasing. Comparison of (2.2) and (2.5) shows that if $h \in C^\infty(0, \infty)$ is a Stieltjes function then $f(\lambda) := \lambda h(\lambda)$ is a complete Bernstein function, and conversely if $f \in C^\infty(0, \infty)$ is a complete Bernstein function then $h(\lambda) := f(\lambda)/\lambda$ is a Stieltjes function. The classes of complete Bernstein functions and Stieltjes functions are preserved under various operations [64, Theorem 6.2, Theorem 7.3, Corollary 7.4, Corollary 7.6]. We present here some which will be used in the paper.

Theorem 2.2. *Let f be a non-zero function on $(0, \infty)$.*

- (a) *f is a complete Bernstein function if and only if $1/f(\lambda)$ is a Stieltjes function.*
- (b) *If f is a complete Bernstein function, then $\lambda/f(\lambda)$ and $\lambda f(1/\lambda)$ are complete Bernstein functions. Conversely, if $\lambda/f(\lambda)$ or $\lambda f(1/\lambda)$ is a complete Bernstein function, then f is a complete Bernstein function.*
- (c) *If f and g are both complete Bernstein functions or both Stieltjes functions, then $g \circ f$ is a complete Bernstein function.*

Many examples of complete Bernstein functions, and hence of Stieltjes functions, are given in [64, Chapter 15]. We give here a few of the most elementary examples that will be relevant in this paper.

Example 2.3. (a) The function $h(\lambda) := \lambda^{-\gamma}$ ($\gamma \in (0, 1)$) is a Stieltjes function with the Stieltjes representation

$$h(\lambda) = \frac{\sin \pi \gamma}{\pi} \int_0^\infty \frac{1}{s + \lambda} \frac{ds}{s^\gamma}, \quad \lambda > 0.$$

Accordingly, $f(\lambda) = \lambda^{1-\gamma} = \lambda h(\lambda)$ is a complete Bernstein function.

(b) For $\alpha \in (0, 1)$, the function $f(\lambda) := \lambda(\lambda + 1)^{-\alpha}$ is a complete Bernstein function with the Stieltjes representation

$$f(\lambda) = \frac{\sin \pi \alpha}{\pi} \int_1^\infty \frac{\lambda}{s + \lambda} \frac{ds}{(s - 1)^\alpha}.$$

Moreover, $\lambda(\lambda + 1)^{-1}$ is a complete Bernstein function with Stieltjes triple $(0, 0, \delta_1)$.

(c) It follows from (a) and (b), together with Theorem 2.2(c), that $f(\lambda) := \lambda^\alpha(1 + \lambda)^{-\beta}$ is a complete Bernstein function whenever $0 \leq \beta \leq \alpha \leq 1$.

Any Stieltjes function or complete Bernstein function can be extended to a holomorphic function on the slit plane Σ_π by means of the formula (2.5) or (2.2), respectively. We shall often regard the functions as being defined on the slit plane in this way, without explicit comment. The rate of decay or growth of such functions at infinity in sectors is determined by the rate on $(0, \infty)$, as shown by the following known fact [61, Lemma 2], [49, Proposition 2.21 (c)].

Proposition 2.4. *Let g be either a Stieltjes function or a complete Bernstein function with Stieltjes representation $(0, 0, \mu)$. Let $\lambda \in \Sigma_\pi$ and $\varphi = \arg \lambda$. Then*

$$\cos(\varphi/2) g(|\lambda|) \leq |g(\lambda)| \leq \sec(\varphi/2) g(|\lambda|).$$

Proof. The second inequality follows from (2.5) or (2.2) and the elementary inequality:

$$|s + \lambda|^2 \geq \frac{(1 + \cos \varphi)}{2} (s + |\lambda|)^2 = (\cos(\varphi/2) (s + |\lambda|))^2, \quad s \in (0, \infty).$$

For the first inequality, we can assume that $g \neq 0$. By Theorem 2.2, $1/g$ is a complete Bernstein function or a Stieltjes function. Hence

$$\frac{1}{|g(\lambda)|} \leq \frac{1}{\cos(\varphi/2) g(|\lambda|)}. \quad \square$$

2.2 Slowly and regularly varying functions

Most of the material in this subsection is standard (see [15, Chapter 1], [47, Sections IV.1-9] or [65]).

Definition 2.5. Let ℓ be a strictly positive measurable function defined on $[a, \infty)$ for some $a \in \mathbb{R}$ and satisfying

$$\lim_{s \rightarrow \infty} \frac{\ell(\lambda s)}{\ell(s)} = 1$$

for every $\lambda > 0$. Then ℓ is said to be *slowly varying*.

Clearly the value of a is not important in this definition. By defining $\ell(s) = \ell(a)$ for $s \in [0, a]$, we may assume that $a = 0$.

Example 2.6. (a) Standard examples of slowly varying functions include

$$\begin{aligned} &\text{iterated logarithms} \quad \log_k(s) := \log \dots \log s, \quad k \in \mathbb{N}, \\ &\exp \{ (\log s)^{\alpha_1} (\log_2(s))^{\alpha_2} \dots (\log_k(s))^{\alpha_k} \}, \quad \alpha_i \in (0, 1), 1 \leq i \leq k, \\ &\exp \left(\frac{\log s}{\log \log s} \right). \end{aligned}$$

(b) It is a straightforward consequence of Definition 2.5 that the sum and product of two slowly varying functions is slowly varying. Moreover, if ℓ is slowly varying then the following are also slowly varying.

$$\begin{aligned} \ell^\alpha &: s \mapsto \ell(s)^\alpha, \quad \alpha \in \mathbb{R}, \\ \ell_\alpha &: s \mapsto \ell(s^\alpha), \quad \alpha > 0, \\ \ell \cdot \log &: s \mapsto \ell(s) \log s. \end{aligned}$$

A proof of the following representation theorem, originally due to Karamata for continuous functions ℓ , may be found in [15, Theorem 1.3.1] or [47, Section IV.3].

Theorem 2.7. *The function ℓ is slowly varying if and only if it is of the form*

$$\ell(s) = c(s) \exp \left\{ \int_a^s \frac{\varepsilon(t)}{t} dt \right\}, \quad s \geq a,$$

for some $a > 0$, where c and ε are measurable functions, $c(s) \rightarrow c > 0$ and $\varepsilon(s) \rightarrow 0$ as $s \rightarrow \infty$.

The following corollary is easily deduced from Theorem 2.7; see [15, Theorem 1.5.6], [65, p.18].

Corollary 2.8. *Let ℓ be a slowly varying function, and $\gamma > 0$.*

(a) *There are positive constants C, c such that*

$$c \left(\frac{s}{t}\right)^\gamma \leq \frac{\ell(t)}{\ell(s)} \leq C \left(\frac{t}{s}\right)^\gamma$$

for all sufficiently large s, t with $t \geq s$.

(b) *As $s \rightarrow \infty$,*

$$s^\gamma \ell(s) \rightarrow \infty, \quad s^{-\gamma} \ell(s) \rightarrow 0. \tag{2.6}$$

On the other hand there are slowly varying functions such that

$$\liminf_{s \rightarrow \infty} \ell(s) = 0, \quad \limsup_{s \rightarrow \infty} \ell(s) = \infty.$$

Definition 2.9. A positive function f is called *regularly varying with index $\alpha \in \mathbb{R}$* if there is a slowly varying function ℓ such that

$$f(s) = s^\alpha \ell(s), \quad s \geq a.$$

Such a function has a representation

$$f(s) = s^\alpha c(s) \exp \left\{ \int_a^s \frac{\varepsilon(t)}{t} dt \right\}, \quad s \geq a,$$

where c and ε are as in Theorem 2.7.

If f is regularly varying of index $\alpha > 0$, there is a strictly increasing, regularly varying function g which is asymptotically equivalent to f [15, Theorem 1.5.3]. One can also arrange that g is smooth [15, Theorem 1.8.2]. Moreover, f has an asymptotic inverse in the sense of (1.8). For example, one may take the asymptotic inverse of f to be the usual inverse of a strictly increasing, continuous, function which is asymptotically equivalent to f . The asymptotic inverse is regularly varying, it depends only on the asymptotic equivalence class of f , and it is unique up to asymptotic equivalence. Indeed, the asymptotic equivalence classes of regularly varying functions with positive index form a group under composition; they also form a semigroup under pointwise multiplication [15, Theorem 1.8.7].

A convenient way to handle asymptotic inverses of regularly varying functions involves the *de Bruijn conjugate* $\ell^\#$ of the slowly varying function ℓ [15, Section 1.5.7]. This is a slowly varying function $\ell^\#$ such that

$$\ell(s)\ell^\#(s\ell(s)) \rightarrow 1 \quad \text{and} \quad \ell^\#(s)\ell(s\ell^\#(s)) \rightarrow 1 \quad \text{as } s \rightarrow \infty.$$

One can take

$$\ell^\#(s) = \frac{(\iota.\ell)^{-1}(s)}{s},$$

where $(\iota.l)(s) = s\ell(s)$. For this choice of $\ell^\#$, it is easy to see that if ℓ is increasing (resp., decreasing), $\ell^\#$ is decreasing (resp., increasing). The group structure of the asymptotic equivalence classes of regularly varying functions immediately implies that one-sided asymptotic inverses are unique up to asymptotic equivalence. Hence, if k is a slowly varying function and either $\ell(s)k(s\ell(s)) \rightarrow 1$ or $k(s)\ell(sk(s)) \rightarrow 1$ as $s \rightarrow \infty$, then $k \sim \ell^\#$.

Example 2.10. A method for finding many de Bruijn conjugates is given in [15, Appendix 5], including the following cases.

1. If $\ell(s) = (\log s)^\beta$ where $\beta \in \mathbb{R}$, then $\ell^\#(s) \sim (\log s)^{-\beta}$.
2. Let $\ell(s) = \exp((\log s)^\beta)$ where $0 < \beta < 1$.
 - (a) If $0 < \beta \leq 1/2$, then $\ell^\#(s) \sim \exp(-(\log s)^\beta)$.
 - (b) If $\frac{1}{2} \leq \beta < \frac{2}{3}$, then

$$\begin{aligned}\ell^\#(s) &\sim \exp(-(\log s)^\beta + \beta(\log s)^{2\beta-1}), \\ (1/\ell)^\#(s) &\sim \exp((\log s)^\beta + \beta(\log s)^{2\beta-1}).\end{aligned}$$

For values of β between $2/3$ and 1 , there are longer formulas of this type.

Proposition 2.11. *Let ℓ be slowly varying, and let $\alpha > 0$. Then*

- (a) $\ell^{\#\#} \sim \ell$.
- (b) *If $f(s) \sim s^\alpha \ell(s^\alpha)$, then $f^{-1}(s) \sim s^{1/\alpha} \ell^\#(s)^{1/\alpha}$.*
- (c) *If $g : (0, a] \rightarrow (0, \infty)$ and $g(s) \sim s^\alpha / \ell(s^{-\alpha})$ as $s \rightarrow 0+$, then $g^{-1}(s) \sim s^{1/\alpha} / \ell^\#(1/s)^{1/\alpha}$ as $s \rightarrow 0+$.*

Proof. The statements (a) and (b) are in [15, Section 1.5] and [65, Section 1.6]. For (c), note that $g(s) \sim 1/f(1/s)$ as $s \rightarrow 0+$, where f is as (b). Using the regular variation, one can easily deduce that $g^{-1}(s) \sim 1/f^{-1}(1/s)$ as $s \rightarrow 0+$. \square

The following lemma describes a common situation in which $\ell^\#$ has a particularly simple form. Parts of the lemma appear in [15, Section 1.5] and [65, Section 1.6].

Lemma 2.12. *Let ℓ be a slowly varying function. The following are equivalent:*

- (i) $\ell^\# \sim 1/\ell$.
- (ii) $\ell(sl(s)) \sim \ell(s)$.
- (iii) $\ell(s/\ell(s)) \sim \ell(s)$.

If ℓ is monotonic and $\alpha > 0$, these properties are equivalent to each of the following:

- (iv) $\ell(sl(s)^\alpha) \sim \ell(s)$.
- (v) $(\ell_\alpha)^\# \sim 1/\ell_\alpha$, where $\ell_\alpha(s) = \ell(s^\alpha)$.

Proof. The statements (ii) and (iii) are equivalent to the statements

$$\ell(s)k(sl(s)) \rightarrow 1, \quad k(s)\ell(sk(s)) \rightarrow 1$$

respectively, when $k = 1/\ell$. So their equivalence to (i) follows from the one-sided uniqueness properties of $\ell^\#$ set out before Example 2.10.

Now assume that (ii) holds and ℓ is monotonic. Replacing s by $sl(s)$ gives

$$\ell\left(sl(s)\ell(sl(s))\right) \sim \ell(sl(s)) \sim \ell(s).$$

Moreover, by (ii),

$$sl(s)^2 \sim sl(s)\ell(sl(s)),$$

so

$$\ell(sl(s)^2) \sim \ell\left(sl(s)\ell(sl(s))\right) \sim \ell(s)$$

by the Uniform Convergence Theorem [15, Theorem 1.2.1]. Iterating this argument gives

$$\ell(sl(s)^\alpha) \sim \ell(s)$$

whenever α is a power of 2. Then the monotonicity of ℓ gives it for all $\alpha > 0$. Thus (ii) implies (iv). The converse follows by replacing $\ell(s)$ by $\ell(s)^{1/\alpha}$.

Finally, let $k_\alpha = 1/\ell_\alpha$. Assume that (iv) holds. Replacing s by s^α in (iv) gives

$$\ell_\alpha(s)k_\alpha(sl_\alpha(s)) = \frac{\ell(s^\alpha)}{\ell(s^\alpha\ell(s^\alpha)^\alpha)} \rightarrow 1.$$

Then (v) follows from the one-sided uniqueness property of $(\ell_\alpha)^\#$. Replacing α by $1/\alpha$ shows that (v) implies (iv). \square

We shall say that a monotonic, slowly varying, function ℓ is *dB-symmetric* when the conditions of Lemma 2.12 are satisfied. Example 2.10 shows that the following functions ℓ are dB-symmetric:

$$\ell(s) = (\log s)^\beta, \text{ for any } \beta \in \mathbb{R},$$

$$\ell(s) = \exp((\log s)^\beta), \text{ if } 0 < \beta < \frac{1}{2}.$$

If ℓ is dB-symmetric, it is clear from Lemma 2.12 that the following functions are also dB-symmetric:

$$\ell_\alpha : s \mapsto \ell(s^\alpha), \text{ for any } \alpha > 0,$$

$$\ell^\alpha : s \mapsto \ell(s)^\alpha, \text{ for any } \alpha \in \mathbb{R}.$$

It is not difficult to show that the product of two dB-symmetric functions is dB-symmetric.

For a regularly varying function f , let $f.\log$ be the following regularly varying function:

$$(f.\log)(s) = f(s) \log s.$$

We shall need the following relations.

Lemma 2.13. *Let $f(s) = s^\alpha \ell(s)$ be a regularly varying function with $\alpha > 0$, and let $\delta > 1/\alpha$. The following hold for some constant C .*

$$(i) \quad (f.\log) \left(\frac{s}{(\log f(s))^\delta} \right) \leq C f(s).$$

$$(ii) \quad f^{-1}(s) \leq C (f.\log)^{-1}(s) (\log s)^\delta.$$

When ℓ is increasing, these statements are also true for $\delta = 1/\alpha$.

Proof. (i). By Corollary 2.8, $\log f(s) \sim \alpha \log s$. Also,

$$\begin{aligned} (f.\log) \left(\frac{s}{(\log f(s))^\delta} \right) &= f(s) \frac{\log \left(\frac{s}{(\log f(s))^\delta} \right) \ell \left(\frac{s}{(\log f(s))^\delta} \right)}{(\log f(s))^{\alpha\delta} \ell(s)} \\ &\leq C f(s) (\log s)^{1-\alpha\delta} \frac{\ell \left(\frac{s}{(\log f(s))^\delta} \right)}{\ell(s)} \\ &\leq C f(s) (\log s)^{1-\alpha\delta+\delta\gamma} \end{aligned}$$

for any $\gamma > 0$ by Corollary 2.8, and for $\gamma = 0$ if ℓ is increasing.

(ii). Replacing s by $f^{-1}(s)$ in (i) gives

$$(f.\log) \left(\frac{f^{-1}(s)}{(\log s)^\delta} \right) \leq C s.$$

The claim follows since $(f.\log)^{-1}$ is increasing and regularly varying. \square

Finally in this section, we consider Stieltjes functions associated with regularly varying functions.

Example 2.14. Let ℓ be a slowly varying function on \mathbb{R}_+ , $\alpha \geq 0$, and assume that $g(s) := s^\alpha \ell(s)$ is increasing. The associated Stieltjes function

$$S_g(\lambda) := \int_{0+}^{\infty} \frac{dg(s)}{s + \lambda} = \int_0^{\infty} \frac{s^\alpha \ell(s)}{(s + \lambda)^2} ds,$$

is defined if either integral is finite [68, p.7]. This occurs if $\alpha < 1$, by (2.6), or if $\alpha = 1$ and $\int_0^{\infty} \frac{\ell(s)}{s+1} ds$ is finite.

We shall need the following abelian/Tauberian theorem of Karamata. For the main results we shall need only the abelian parts (i) \Rightarrow (ii) and (iii) \Rightarrow (iv).

Theorem 2.15 (Karamata). *Let g be an increasing function on \mathbb{R}_+ , and let S_g be the associated Stieltjes function. Let $0 < \sigma \leq 1$, and ℓ be slowly varying on \mathbb{R}_+ .*

(a) *The following are equivalent:*

$$(i) \quad g(s) \sim s^{1-\sigma} \ell(s) \quad s \rightarrow \infty;$$

$$(ii) \quad S_g(\lambda) \sim \Gamma(\sigma) \Gamma(2 - \sigma) \lambda^{-\sigma} \ell(\lambda), \quad \lambda \rightarrow \infty.$$

(b) *The following are equivalent:*

- (iii) $g(s) \sim s^{1-\sigma} \ell(1/s), \quad s \rightarrow 0+;$
- (iv) $S_g(\lambda) \sim \Gamma(\sigma)\Gamma(2-\sigma)\lambda^{-\sigma} \ell(1/\lambda), \quad \lambda \rightarrow 0+.$

Proof. The first statement is proved in [15, Theorem 1.7.4] (the Tauberian implication is proved in [65, Theorem 2.5]). The proof of the second statement is very similar, using the same preliminary results from [15, Sections 1.5,1.7]. \square

3 Functional calculus of sectorial operators

In this section we recall basic properties of functional calculus of sectorial operators based on complete Bernstein and Stieltjes functions, originally due to Hirsch and extended to include fractional powers. We concentrate on those properties which are needed for our main purposes in the later sections of this paper. A much fuller account of the calculus for Bernstein functions can be found in the monograph [64], and of the extended calculus in [36].

3.1 Sectorial operators

Many parts of the paper will involve the notion of a sectorial operator which we recall now.

Definition 3.1. A densely defined, linear operator A on a Banach space X is called *sectorial* if $(-\infty, 0) \subset \rho(A)$ and there exists $C > 0$ such that

$$\|\lambda(\lambda + A)^{-1}\| \leq C, \quad \lambda > 0. \tag{3.1}$$

Some authors require a sectorial operator to be injective, and some do not require it to be densely defined. Some of the operators that we consider will not be injective.

Note that any sectorial operator A is closed and by the Neumann series expansion, $(\lambda + A)^{-1}$ is defined and (3.1) holds on the sector Σ_φ , for some $\varphi \in (0, \pi]$. Moreover,

$$\lambda(\lambda + A)^{-1}x \rightarrow x, \quad \lambda \rightarrow \infty, \quad x \in X, \tag{3.2}$$

[36, Proposition 2.1.1,d)].

If $-A$ generates a bounded C_0 -semigroup, then A is sectorial. The following standard lemma establishes that certain auxiliary operators which play important roles in this paper are also sectorial even though they may not be negative generators of bounded semigroups.

Lemma 3.2. *Let A be a sectorial operator on a Banach space X . Then the operators A^{-1} (if A is invertible), $A(I + A)^{-1}$ and $A(I + A)^{-2}$ are sectorial.*

Proof. Sectoriality of A^{-1} and $A(I + A)^{-1}$ follows from the identities

$$\begin{aligned} (\lambda + A^{-1})^{-1} &= \lambda^{-1} - \lambda^{-2}(\lambda^{-1} + A)^{-1}, \\ (\lambda + A(I + A)^{-1})^{-1} &= \frac{1}{\lambda + 1} + \frac{1}{(\lambda + 1)^2} \left(\frac{\lambda}{\lambda + 1} + A \right)^{-1}, \end{aligned}$$

which hold for $\lambda > 0$ (see [36, Proposition 2.1.1,b]) and [60, Lemma 3.1]). To prove sectoriality of $A(I + A)^{-2}$, we note the identity

$$\lambda(\lambda + A(I + A)^{-2})^{-1} = I - \lambda^{-1}(\mu + A)^{-1} \left(I - \mu^{-1}(\mu^{-1} + A)^{-1} \right),$$

where $\mu > 1$ and μ^{-1} are the roots of $\mu^2 - (\lambda^{-1} + 2)\mu + 1 = 0$. Since $\mu > \lambda^{-1}$, sectoriality of A implies that $\|\lambda^{-1}(\mu + A)^{-1}\|$ and $\|\mu^{-1}(\mu^{-1} + A)^{-1}\|$ are bounded for $\lambda > 0$, hence $A(I + A)^{-2}$ is sectorial. \square

Other information about sectorial operators may be found in [36] and [55] (in the latter sectorial operators are called non-negative operators).

3.2 Hirsch functional calculus

We now define complete Bernstein functions of sectorial operators and review those basic properties that we need in the sequel. There are several different definitions of functional calculus of sectorial operators, and we shall describe some properties which cross over between the different definitions. We try to present the ideas of functional calculus in a way which reveals the heuristics of our subsequent arguments, and to give the later proofs in ways which do not rely on any unjustified assumptions about compatibility of different definitions.

Let f be a complete Bernstein function with Stieltjes representation (a, b, μ) , and let A be a sectorial operator on a Banach space X . The next definition was essentially given in [41, p. 255].

Definition 3.3. Define an operator $f_0(A) : \text{dom}(A) \rightarrow X$ by

$$f_0(A)x = ax + bAx + \int_{0+}^{\infty} A(\lambda + A)^{-1}x \, d\mu(\lambda), \quad x \in \text{dom}(A). \quad (3.3)$$

By (2.3), this integral is absolutely convergent and $f_0(A)(I + A)^{-1}$ is a bounded operator on X , extending $(I + A)^{-1}f_0(A)$. Hence $f_0(A)$ is closable as an operator on X . Define

$$f(A) = \overline{f_0(A)}.$$

We call $f(A)$ a *complete Bernstein function of A* .

Actually Hirsch used the representation (2.4) for a complete Bernstein function f and then defined the corresponding operator $f(A)$ as

$$f^H(A) = \overline{f_0^H(A)},$$

where f_0^H is defined on $\text{dom}(A)$ by

$$f_0^H(A)x := a + \nu(\{0\})Ax + \int_{0+}^{\infty} A(I + \lambda A)^{-1}x \, d\nu(\lambda), \quad x \in \text{dom}(A). \quad (3.4)$$

By Remark 2.1 the representations (2.2) and (2.4) describe exactly the same classes of complete Bernstein functions. To see that the resulting operators $f(A)$ and $f^H(A)$ coincide, it suffices to change variables in the same way as in Remark 2.1, considering vector-valued integrals instead of scalar ones. Thus we can use Hirsch's results even if we think of complete Bernstein functions f as being represented by (2.2).

When $-A$ generates a bounded C_0 -semigroup, one can define the operators $f(A)$ for arbitrary Bernstein functions f by adapting the Lévy-Khintchine formula (2.1). A detailed discussion of this approach as well as of various properties of $f(A)$ can be found in [64, Section 12] (where A denotes the generator of the semigroup). One can prove that the definition of complete Bernstein functions of semigroup generators in [64] is consistent with Hirsch's definition (Definition 3.3), but we shall not go into details. We shall refer to some results in [64] for complete Bernstein functions even though they start from the Lévy-Khintchine representation.

By Definition 3.3, $\text{dom}(A)$ is a core for $f(A)$. Using (3.2) and the fact that $(\lambda + A)^{-1}$ commutes with $(I + A)f_0(A)(I + A)^{-1}$, we infer that $\text{dom}(A)$ is a core for $(I + A)f_0(A)(I + A)^{-1}$ as well. Since $(I + A)f_0(A)(I + A)^{-1}$ is closed as a product of a closed operator and a bounded operator, it follows that

$$f(A) = (I + A)f_0(A)(I + A)^{-1}.$$

Remark 3.4. If f is a bounded complete Bernstein function then, in the Stieltjes representation of f , $b = 0$ and Fatou's lemma implies that the measure μ is finite (see [64, Corollary 3.7(v)]). In this case, $f_0(A)$ is bounded on $\text{dom}(A)$, and therefore $f(A)$ is a bounded operator on X . It is also straightforward to see that $f(A)$ is bounded for any complete Bernstein function f if A is bounded (see [64, Corollary 12.7]).

Remark 3.5. In the sequel, (3.3) will usually be used for measures μ which are Lebesgue-Stieltjes measures associated with increasing, regularly varying functions g . Then we may think of the integrals as being vector-valued Riemann-Stieltjes integrals rather than Bochner integrals. The theory of vector-valued Riemann-Stieltjes integrals is presented in [6, Section 1.9], [39, Section III.3.3], [68, Section 1]. One can define complete Bernstein functions and operator Bernstein functions initially by means of Riemann-Stieltjes integrals. However, in the relevant literature, including [41], [43], [62] and [64], (2.2) and (3.3), or (2.4) and (3.4), are standard ways to define complete Bernstein functions, and we have chosen to follow an established route.

Complete Bernstein functions of sectorial operators possess a number of properties which allow one to create a partial functional calculus for A . However the set of complete Bernstein functions is not closed under pointwise multiplication, so the multiplicative properties of this process are restricted. The subject was thoroughly investigated by Hirsch in the 1970s, and he proved the following properties of operator Bernstein functions in [41, Théorème 1-3] and [43, Théorème 1] (see also [43, pp. 200-201], [62] and [40]).

Theorem 3.6. *Let A be a sectorial operator on a Banach space X , and let f and g be complete Bernstein functions. Then the following statements hold.*

(i) *The operators $f(A)$ and $g(A)$ are sectorial.*

(ii) *The composition rule holds:*

$$f(g(A)) = (f \circ g)(A).$$

(iii) *If fg is also a complete Bernstein function, then the product rule holds:*

$$f(A)g(A) = (fg)(A).$$

For $\alpha \in (0, 1)$, Example 2.3(a) shows that z^α is a complete Bernstein function, and we write A^α for the corresponding complete Bernstein function of A . These fractional powers coincide with the standard fractional powers which are extensively studied in [55]. If A is sectorial and $\alpha, \beta \in (0, 1)$ then Theorem 3.6 implies that $(A^\alpha)^\beta = A^{\alpha\beta}$, and moreover $A^\alpha A^\beta = A^{\alpha+\beta}$ if $\alpha + \beta \leq 1$. In this paper we shall need these properties for a larger range of α and β . This is standard theory, but we put it in a broader context here.

To this aim, we shall use an extended holomorphic functional calculus, which has become a standard tool to deal with functions of sectorial operators. To keep the presentation within reasonable limits we give only a very brief sketch of part of the extended holomorphic calculus and refer for further details to [36, Chapters 1,2] and [48, Sections II.9, II.15].

Let H be the algebra of functions which are holomorphic in $\Sigma_\pi = \mathbb{C} \setminus (-\infty, 0]$. Let \mathcal{H} stand for the set of all $f \in H$ such that for any $\varphi \in (0, \pi)$ there exist $C > 0$ and $\alpha > 0$ (both depending on φ) satisfying

$$|f(z)| \leq C|z|^\alpha, \quad z \in \Sigma_\varphi. \quad (3.5)$$

Let $\tilde{\mathcal{H}}$ denote the set of all $f \in H$ such that for any $\varphi \in (0, \pi)$ one has

$$|f(z)| \leq C \max(|z|^\alpha, |z|^{-\alpha}), \quad z \in \Sigma_\varphi, \quad (3.6)$$

for some $C, \alpha > 0$. It follows easily from (2.2), (2.3) and (2.5) that $\tilde{\mathcal{H}}$ contains all complete Bernstein functions and all Stieltjes functions, and one can set $\alpha = 1$ for all $\varphi \in (0, \pi)$.

Let $\mathcal{C}(X)$ denote the set of all closed, densely defined, linear operators on X .

Theorem 3.7. *Let A be a sectorial operator on a Banach space X . Then there exists a well-defined mapping*

$$\begin{aligned} \mathcal{H} &\mapsto \mathcal{C}(X), \\ f &\mapsto f(A), \end{aligned}$$

called an extended holomorphic functional calculus, such that

(i) $1(A) = I$ and $z(A) = A$;

- (ii) If $T \in \mathcal{L}(X)$ and $TA \subset AT$, then $Tf(A) \subset f(A)T$;
- (iii) $f(A) + g(A) = (f + g)(A)$, if $g(A) \in \mathcal{L}(X)$;
- (iv) $f(A)g(A) = (fg)(A)$, if $g(A) \in \mathcal{L}(X)$;
- (v) if $g \in \mathcal{H}$ is such that $g(A)$ is sectorial and there exists $\vartheta : (0, \pi) \rightarrow (0, \pi)$ such that $\lim_{\varphi \rightarrow \pi^-} \vartheta(\varphi) = \pi$ and $g(\Sigma_{\vartheta(\varphi)}) \subset \Sigma_{\varphi}$ for all $\varphi \in (0, \pi)$, then $(f \circ g)(A) = f(g(A))$.

If A is injective then there exists a mapping $\tilde{\mathcal{H}} \mapsto \mathcal{C}(X)$, $f \mapsto f(A)$, satisfying the properties (i)-(v) above.

Remark 3.8. We note the following facts about the extended holomorphic functional calculus described in Theorem 3.7.

- (i) If $f(A) \in \mathcal{L}(X)$ and $\lambda \in \rho(A)$, then $f(A)(\lambda + A)^{-1} = (\lambda + A)^{-1}f(A)$.
- (ii) If f is a rational function whose poles all lie in $(-\infty, 0]$, then $f(A)$ as defined in the extended holomorphic functional calculus of Theorem 3.7 agrees with the natural definition of $f(A)$.
- (iii) By [14], if $f \in \mathcal{H}$ is a complete Bernstein function, then $f(A)$ as in the calculus of Theorem 3.7 agrees with $f(A)$ as defined in Definition 3.3. The same holds if A is injective and f is a complete Bernstein function belonging to $\tilde{\mathcal{H}}$.
- (iv) The composition rule Theorem 3.7(v) applies in particular when $g(z) = z^{-1}$ (if A is invertible), $z(1 + z)^{-1}$, or $z(1 + z)^{-2}$. In each case, it suffices to note Lemma 3.2 and to use the fact that the mapping $z \mapsto 1/z$ preserves sectors.
- (v) In particular, if A is invertible, $g \sim (0, 0, \nu)$ is a Stieltjes function and f is the complete Bernstein function given by $f(z) = g(1/z)$, then

$$g(A) = \int_{0+}^{\infty} (\lambda + A)^{-1} d\nu(\lambda) = f(A^{-1}).$$

It is easy to see that $\{z^\alpha : \alpha > 0\} \subset \mathcal{H}$ and $\{z^\alpha : \alpha \in \mathbb{R}\} \subset \tilde{\mathcal{H}}$. Thus if A is sectorial, then the “fractional powers” A^α for $\alpha > 0$ (and for all $\alpha \in \mathbb{R}$, if A is invertible) are well-defined by means of the extended holomorphic functional calculus and, as we shall see below, they behave very similarly to the scalar functions z^α . These fractional powers coincide with the fractional powers considered in [36, Chapters 3,4], [48, Section II.15] and [54, Chapter 2] where all properties of fractional powers of operators needed in this paper can be found. For the reader’s convenience we recall and summarize them in the following statement.

Theorem 3.9. *Let A be a sectorial operator on a Banach space X . Then for any $\alpha > 0$ and $\beta > 0$ the following statements hold.*

- (i) If $T \in \mathcal{L}(X)$ and $TA \subset AT$ then $TA^\alpha \subset A^\alpha T$.

- (ii) *The semigroup law holds: $A^{\alpha+\beta} = A^\alpha A^\beta$.*
- (iii) $\text{dom}(A^\beta) = \text{dom}((I + A)^\beta) = \text{ran}((I + A)^{-\beta})$.
- (iv) *If in addition $\alpha \in (0, 1)$, then A^α is sectorial, and the composition law holds: $(A^\alpha)^\beta = A^{\alpha\beta}$.*

If A is invertible then $A^{-\alpha} = (A^{-1})^\alpha$ for $\alpha > 0$, (ii) is true for $\alpha, \beta \in \mathbb{R}$, and (iv) holds for $\alpha \in (-1, 1)$ and $\beta \in \mathbb{R}$.

The following facts are elementary when α and β are integers (Lemma 3.2 and [48, Proposition 9.4(d)]) and they may be known for fractional powers, but we were unable to find them in the literature. The fact that $(-\infty, 0)$ is contained in the resolvent set of $A^\alpha(I + A)^{-1}$ when $0 < \alpha < 1$ follows from [35, Theorem 4.1], but the method given there does not establish the sectoriality estimate. We are very grateful to Alexander Gomilko for providing the proof of that estimate.

Proposition 3.10. *Let A be a sectorial operator on a Banach space X , and let $\alpha, \beta \in [0, 1]$. The following statements hold:*

- (i) $\text{ran}(A^\alpha(I + A)^{-(\alpha+\beta)}) = \text{ran}(A^\alpha) \cap \text{dom}(A^\beta)$.
- (ii) $A^\alpha(I + A)^{-\beta}$ is sectorial.

Proof. To prove (i), we shall use Theorem 3.9 and the consequential facts that $A^\alpha(I + A)^{-\alpha}$ and $(I + A)^{-\beta}$ are bounded commuting operators, and $\text{ran}(A^\alpha) = \text{ran}(A^\alpha(I + A)^{-\alpha})$. Hence

$$\begin{aligned} \text{ran}(A^\alpha) \cap \text{dom}(A^\beta) &= \text{ran}(A^\alpha(I + A)^{-\alpha}) \cap \text{ran}((I + A)^{-\beta}) \\ &\supseteq \text{ran}(A^\alpha(I + A)^{-(\alpha+\beta)}). \end{aligned}$$

Conversely, let $x \in \text{ran}(A^\alpha) \cap \text{dom}(A^\beta)$. Then $x = A^\alpha y_1$ and $x = (I + A)^{-\beta} y_2$ for some $y_1 \in \text{dom}(A^\alpha)$, $y_2 \in X$. Since $A^\alpha y_1 = (I + A)^{-\beta} y_2 \in \text{dom}(A^\beta)$, one has $y_1 \in \text{dom}(A^{\alpha+\beta}) = \text{dom}((I + A)^{\alpha+\beta})$. Let $y_3 = (I + A)^{\alpha+\beta} y_1$. Then

$$x = A^\alpha y_1 = A^\alpha(I + A)^{-(\alpha+\beta)} y_3.$$

For (ii), first consider the case when $0 < \beta \leq \alpha \leq 1$. By Example 2.3(b), $f(\lambda) := \lambda^\alpha(1 + \lambda)^{-\beta}$ is a complete Bernstein function. By Theorem 3.6 (i),(ii), $A^\alpha(I + A)^{-\beta} = f(A)$ which is sectorial.

Now consider the case when $0 < \alpha < \beta = 1$. Let $\lambda > 0$, and

$$f_\alpha(z) = \frac{z^\alpha}{1 + z},$$

and

$$g_{\alpha,\lambda}(z) = \frac{1}{\lambda} - \frac{1}{f_\alpha(z) + \lambda} = \frac{f_\alpha(z)}{\lambda(f_\alpha(z) + \lambda)} = \frac{z^\alpha}{\lambda(z^\alpha + \lambda(1 + z))},$$

for $z \in \Sigma_\pi$. For $\varphi \in (0, \pi)$, let γ_φ be the contour given by

$$\gamma_\varphi := \{te^{-i\varphi} : t \geq 0\} \cup \{te^{i\varphi} : t \geq 0\},$$

taken in the downward direction. Then

$$\int_{\gamma_\varphi} |g_{\alpha,\lambda}(z)| \frac{|dz|}{|z|} < \infty, \quad (3.7)$$

and

$$g_{\alpha,\lambda}(\mu) = \frac{1}{2\pi i} \int_{\gamma_\varphi} \frac{g_{\alpha,\lambda}(z)}{\mu - z} dz, \quad |\arg \mu| < \varphi.$$

Letting $\varphi \rightarrow \pi-$, we obtain

$$g_{\alpha,\lambda}(\mu) = \frac{\sin \pi\alpha}{\pi} \int_0^\infty \frac{(1-t)t^\alpha}{|e^{\pi i\alpha}t^\alpha + \lambda(1-t)|^2} (\mu + t)^{-1} dt. \quad (3.8)$$

Letting $\mu \rightarrow 0+$ we also have

$$\int_0^\infty \frac{(1-t)t^{\alpha-1}}{|e^{\pi i\alpha}t^\alpha + \lambda(1-t)|^2} dt = 0. \quad (3.9)$$

By the product rule of Theorem 3.7(iv), $f_\alpha(A) = A^\alpha(I + A)^{-1}$. From (3.7) and [36, Section 2.3],

$$g_{\alpha,\lambda}(A) = \frac{1}{2\pi i} \int_{\gamma_\varphi} g_{\alpha,\lambda}(z)(z + A)^{-1} dz,$$

for $\varphi \in (0, \pi)$ sufficiently large. Here the integral is absolutely convergent, so $g_{\alpha,\lambda}(A) \in \mathcal{L}(X)$. Since

$$(\lambda + f_\alpha(z)) (\lambda^{-1} - g_{\alpha,\lambda}(z)) = 1,$$

the product rule implies that

$$\lambda^{-1} - g_{\alpha,\lambda}(A) = (\lambda + A^\alpha(I + A)^{-1})^{-1}. \quad (3.10)$$

The same arguments as for (3.8) show that

$$g_{\alpha,\lambda}(A) = \frac{\sin \pi\alpha}{\pi} \int_0^\infty \frac{(1-t)t^\alpha}{|e^{\pi i\alpha}t^\alpha + \lambda(1-t)|^2} (t + A)^{-1} dt.$$

From this, (3.1) and (3.9) we obtain

$$\begin{aligned} \|g_{\alpha,\lambda}(A)\| &\leq \frac{C \sin \pi\alpha}{\pi} \int_0^\infty \frac{|1-t|t^{\alpha-1}}{|e^{\pi i\alpha}t^\alpha + \lambda(1-t)|^2} dt \\ &= \frac{2C \sin \pi\alpha}{\pi} \int_0^1 \frac{(1-t)t^{\alpha-1}}{|e^{\pi i\alpha}t^\alpha + \lambda(1-t)|^2} dt. \end{aligned}$$

For $t \in (0, 1)$,

$$\begin{aligned} |e^{\pi i\alpha}t^\alpha + \lambda(1-t)|^2 &= t^{2\alpha} + (1-t)^2\lambda^2 + 2\cos(\pi\alpha)\lambda t^\alpha(1-t) \\ &\geq c_\alpha (t^{2\alpha} + (1-t)^2\lambda^2), \end{aligned}$$

for some $c_\alpha > 0$. Hence

$$\|g_{\alpha,\lambda}(A)\| \leq C_\alpha \int_0^1 \frac{(1-t)t^{\alpha-1}}{t^{2\alpha} + (1-t)^2\lambda^2} dt.$$

Now

$$\begin{aligned} \int_0^{1/2} \frac{(1-t)t^{\alpha-1}}{t^{2\alpha} + (1-t)^2\lambda^2} dt &\leq \int_0^{1/2} \frac{t^{\alpha-1}}{t^{2\alpha} + (\lambda/2)^2} dt \\ &\leq \frac{1}{\alpha} \int_0^\infty \frac{d\tau}{\tau^2 + (\lambda/2)^2} = \frac{\pi}{\alpha\lambda}, \end{aligned}$$

and

$$\begin{aligned} \int_{1/2}^1 \frac{(1-t)t^{\alpha-1}}{t^{2\alpha} + (1-t)^2\lambda^2} dt &\leq 2^{1-\alpha} \int_{1/2}^1 \frac{(1-t)}{2^{-2\alpha} + (1-t)^2\lambda^2} dt \\ &= 2^{1-\alpha} \int_0^{1/2} \frac{\tau}{2^{-2\alpha} + \tau^2\lambda^2} d\tau \\ &= \frac{1}{2^\alpha\lambda^2} \log \left(1 + \left(\frac{2^\alpha\lambda}{2} \right)^2 \right) \leq \frac{1}{\lambda}, \end{aligned}$$

since

$$\log(1 + s^2) \leq 2 \log(1 + s) \leq 2s, \quad s > 0,$$

Thus

$$\|\lambda g_{\alpha,\lambda}(A)\| \leq C_\alpha, \quad \lambda > 0.$$

It follows from (3.10) that $A^\alpha(I + A)^{-1}$ is sectorial when $0 < \alpha < 1$.

When $0 \leq \alpha < \beta < 1$, we have

$$A^\alpha(I + A)^{-\beta} = (A^{\alpha/\beta}(I + A)^{-1})^\beta,$$

and this is sectorial, by the previous case together with Theorem 3.9 (iv). \square

Let A be a sectorial operator and f be a complete Bernstein function. Then one has

$$\|f(A)x\| \leq C\|x\|f\left(\frac{\|Ax\|}{\|x\|}\right), \quad x \in \text{dom}(A), x \neq 0, \quad (3.11)$$

where C is a constant independent of x (and f). This is shown in [62] for sectorial operators, and in [64, Corollary 12.8] where $-A$ is assumed to be the generator of a bounded C_0 -semigroup but f may be *any* Bernstein function.

If A is invertible, we also have

$$\|f(A^{-1})x\| \leq C\|x\|f\left(\frac{\|A^{-1}x\|}{\|x\|}\right), \quad x \in X, x \neq 0. \quad (3.12)$$

This follows since A^{-1} is also sectorial. Alternatively, one can easily pass between (3.11) and (3.12) by considering $\varphi(z) := zf(1/z)$. Then φ is also a complete Bernstein function

by Theorem 2.2, and $\varphi(A) = Af(A^{-1})$ by the product rule and composition rule for $g(z) = z^{-1}$, in Theorem 3.7 and Remark 3.8(v). Applying (3.11) with f replaced by φ gives

$$\|f(A^{-1})Ax\| \leq C\|Ax\|f\left(\frac{\|x\|}{\|Ax\|}\right), \quad x \in \text{dom}(A), x \neq 0.$$

Setting $Ax = y$ we obtain (3.12). Conversely, applying (3.12) with f replaced by φ gives

$$\|f(A)A^{-1}x\| \leq C\|A^{-1}x\|f\left(\frac{\|x\|}{\|A^{-1}x\|}\right), \quad x \in X, x \neq 0.$$

Setting $A^{-1}x = y$ (so $y \in \text{dom}(A)$) we obtain (3.11).

When $f(z) = z^\alpha$ ($0 < \alpha < 1$), we recover the classical moment inequality for fractional powers in the forms

$$\|A^\alpha x\| \leq C\|x\|^{1-\alpha}\|Ax\|^\alpha, \quad x \in \text{dom}(A), \quad (3.13)$$

$$\|A^{-\alpha}x\| \leq C\|x\|^{1-\alpha}\|A^{-1}x\|^\alpha, \quad x \in X. \quad (3.14)$$

4 Some estimates for semigroup asymptotics

In this section we give some results relating different types of asymptotic estimates for semigroups. In Subsection 4.1 we present some inequalities which are related to (3.12) and (3.13) but apply to generators of bounded semigroups. In Subsection 4.2 we show how certain resolvent estimates can be transferred to semigroup estimates in the case of bounded semigroups on Hilbert space.

4.1 Moment and interpolation inequalities

We start by recalling the full moment inequality for sectorial operators, extending (3.13) and (3.14).

Proposition 4.1. *Let B be a sectorial operator, and $0 \leq \alpha < \beta < \gamma$. There is a constant C such that*

$$\|B^\beta x\| \leq C\|B^\alpha x\|^{(\gamma-\beta)/(\gamma-\alpha)}\|B^\gamma x\|^{(\beta-\alpha)/(\gamma-\alpha)}, \quad x \in \text{dom}(B^\gamma). \quad (4.1)$$

Hence if $S : X \rightarrow \text{dom}(B^\gamma)$ is a linear operator and $B^\gamma S \in \mathcal{L}(X)$, then

$$\|B^\beta S\| \leq C\|B^\alpha S\|^{(\gamma-\beta)/(\gamma-\alpha)}\|B^\gamma S\|^{(\beta-\alpha)/(\gamma-\alpha)}. \quad (4.2)$$

If B is invertible, then (4.1) and (4.2) hold whenever $\alpha < \beta < \gamma$.

Proof. For $\alpha = 0$, (4.1) is the standard inequality (3.13) [48, Theorem 15.14], [54, Corollary 5.1.13]. The more general cases follow by replacing β by $\beta - \alpha$, γ by $\gamma - \alpha$ and x by $B^\alpha x$. Then (4.2) follows on replacing x by Sx . When B is invertible the range of the inequalities can be extended by replacing x or S by $B^{-n}x$ or $B^{-n}S$. \square

Next we deduce an inequality of interpolation type which was proved in a slightly less general form (and with a slightly different proof) in [9, Proposition 3.1].

Lemma 4.2. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Banach space X , and let $B \in \mathcal{L}(X)$ be a sectorial operator commuting with $(T(t))_{t \geq 0}$. Let $\gamma, \delta > 0$. Then there exist positive constants C, c such that*

$$c\|T(Ct)B^\gamma\|^\delta \leq \|T(t)B^\delta\|^\gamma \leq C\|T(ct)B^\gamma\|^\delta, \quad t > 0. \quad (4.3)$$

In particular, the following statements are equivalent:

$$(i) \|T(t)B^\gamma\| = O(t^{-1}), \quad t \rightarrow \infty.$$

$$(ii) \|T(t)B\| = O(t^{-1/\gamma}), \quad t \rightarrow \infty.$$

Proof. Take $n \in \mathbb{N}$ such that $n\gamma \geq \delta$, and apply Proposition 4.1 with $\alpha = 0$, $\beta = \delta/n$ and $S = T(t/n)$. Then

$$\|B^{\delta/n}T(t/n)\| \leq C\|T(t/n)\|^{1-\delta/(n\gamma)}\|B^\gamma T(t/n)\|^{\delta/(n\gamma)} \leq C\|B^\gamma T(t/n)\|^{\delta/(n\gamma)}.$$

Now

$$\|T(t)B^\delta\|^\gamma \leq \|T(t/n)B^{\delta/n}\|^{n\gamma} \leq C\|T(t/n)B^\gamma\|^\delta.$$

This gives the second inequality in (4.3), and the first follows by interchanging γ and δ . The final statement follows by taking $\delta = 1$. \square

Our next result gives more interpolation properties for the generator of a bounded C_0 -semigroup. We shall need them for our main results in Sections 5 and 7. To simplify the presentation here and in Section 7 we introduce the shorthand notation:

$$B(A) := A(I + A)^{-1}$$

when A is a sectorial operator. By Lemma 3.2, $B(A)$ is sectorial. Thus the fractional powers $B(A)^\alpha, \alpha > 0$, are well-defined, and by the product and composition rules in Theorem 3.7(iv) and Remark 3.8(iv),

$$B(A)^\alpha = A^\alpha(I + A)^{-\alpha}. \quad (4.4)$$

Theorem 4.3. *Let $-A$ be the generator of a bounded C_0 -semigroup $(T(t))_{t \geq 0}$ on a Banach space X , and let $B(A) = A(I + A)^{-1}$. Assume that $T(t)A \neq 0$ for each $t > 0$. There exists a constant $c > 0$ such that the following hold.*

(a) *If A is invertible, f is a complete Bernstein function, $\gamma \leq 1$, and $A^\gamma f(A^{-1})$ is a bounded operator, then*

$$\|T(t_1)A^\gamma f(A^{-1})\| \geq c \frac{\|T(t_1 + t_2)A^{\gamma-1}\|}{\|T(t_2)A^{-1}\|} f(\|T(t_2)A^{-1}\|) \quad (4.5)$$

for all $t_1, t_2 \geq 0$.

(b) If f is a bounded complete Bernstein function and $\gamma > 0$, then

$$\|T(t_1)B(A)^\gamma f(A)\| \geq c \frac{\|T(t_1+t_2)B(A)^{\gamma+1}\|}{\|T(t_2)B(A)\|} f(\|T(t_2)B(A)\|)$$

for all $t_1, t_2 \geq 0$.

Remark 4.4. In our applications of Theorem 4.3, we shall take $t_1 = t_2$ and choose specific values of $\gamma > 0$, but the applications are rather delicate. For example, the ratio

$$\frac{\|T(2t)A^{\gamma-1}\|}{\|T(t)A^{-1}\|}$$

tends to 0 as $t \rightarrow \infty$ in all the cases in which we are interested in Section 5. The inequality (4.5) will be used to improve other estimates for the rate of decay of $\|T(t)A^{-1}\|$.

Proof. (a) We can assume that f is non-zero, and we let φ be the complete Bernstein function given by $\varphi(z) = z/f(z)$, $z > 0$ (Theorem 2.2). Since f , φ and the identity function z are complete Bernstein functions, the product rule (Theorem 3.6(iii)) yields

$$f(A^{-1})\varphi(A^{-1}) = A^{-1}.$$

Then

$$\begin{aligned} \|T(t_1+t_2)A^{\gamma-1}\| &= \|T(t_1+t_2)A^\gamma f(A^{-1})\varphi(A^{-1})\| \\ &\leq \|T(t_1)A^\gamma f(A^{-1})\| \|T(t_2)\varphi(A^{-1})\|. \end{aligned}$$

Thus

$$\|T(t_1)A^\gamma f(A^{-1})\| \geq \frac{\|T(t_1+t_2)A^{\gamma-1}\|}{\|T(t_2)\varphi(A^{-1})\|}. \quad (4.6)$$

We now estimate $\|T(t)\varphi(A^{-1})\|$ from above for $t > 0$. Let φ have Stieltjes representation (a, b, μ) . By (3.3),

$$T(t)\varphi(A^{-1}) = aT(t) + bT(t)A^{-1} + \int_{0+}^{\infty} T(t)A^{-1}(\lambda + A^{-1})^{-1} d\mu(\lambda).$$

Let $\tau = \|T(t)A^{-1}\|$. Then

$$\left\| \int_{0+}^{\tau} T(t)A^{-1}(\lambda + A^{-1})^{-1} d\mu(\lambda) \right\| \leq C \int_{0+}^{\tau} d\mu(\lambda)$$

since A^{-1} is sectorial and $(T(t))_{t \geq 0}$ is bounded. Moreover,

$$\begin{aligned} \left\| \int_{\tau+}^{\infty} T(t)A^{-1}(\lambda + A^{-1})^{-1} d\mu(\lambda) \right\| &\leq \tau \int_{\tau+}^{\infty} \|(\lambda + A^{-1})^{-1}\| d\mu(\lambda) \\ &\leq C\tau \int_{\tau+}^{\infty} \frac{d\mu(\lambda)}{\lambda}. \end{aligned}$$

Let $K = \sup_{t \geq 0} \|T(t)\|$. Then

$$\begin{aligned} \|T(t)\varphi(A^{-1})\| &\leq aK + b\tau + C \int_{0+}^{\tau} d\mu(\lambda) + C\tau \int_{\tau+}^{\infty} \frac{d\mu(\lambda)}{\lambda} \\ &\leq aK + b\tau + 2C \int_{0+}^{\tau} \frac{\tau}{\lambda + \tau} d\mu(\lambda) + 2C \int_{\tau+}^{\infty} \frac{\tau}{\lambda + \tau} d\mu(\lambda) \\ &\leq 2C\varphi(\tau). \end{aligned}$$

Putting $t = t_2$, (4.6) gives

$$\begin{aligned} \|T(t_1)A^\gamma f(A^{-1})\| &\geq c \frac{\|T(t_1 + t_2)A^{\gamma-1}\|}{\varphi(\|T(t_2)A^{-1}\|)} \\ &= c \frac{\|T(t_1 + t_2)A^{\gamma-1}\|}{\|T(t_2)A^{-1}\|} f(\|T(t_2)A^{-1}\|). \end{aligned}$$

(b) The proof is similar to (a). We now use the product rule in the form

$$f(A)\varphi(A) = A.$$

Observe also that $f(A)$ is bounded (Remark 3.4), $\varphi(A)$ is closed and $\text{dom}(A) \subset \text{dom}(\varphi(A))$ (Definition 3.3), and hence the operator $\varphi(A)(I + A)^{-1}$ is bounded. So

$$\begin{aligned} \|T(t_1 + t_2)B(A)^{\gamma+1}\| &= \|T(t_1 + t_2)B(A)^\gamma f(A)\varphi(A)(I + A)^{-1}\| \\ &\leq \|T(t_1)B(A)^\gamma f(A)\| \|T(t_2)\varphi(A)(I + A)^{-1}\|. \end{aligned}$$

Thus

$$\|T(t_1)B(A)^\gamma f(A)\| \geq \frac{\|T(t_1 + t_2)B(A)^{\gamma+1}\|}{\|T(t_2)\varphi(A)(I + A)^{-1}\|}.$$

Now

$$\begin{aligned} T(t)\varphi(A)(I + A)^{-1} &= aT(t)(I + A)^{-1} + bT(t)A(I + A)^{-1} + \int_{0+}^{\infty} T(t)A(\lambda + A)^{-1}(I + A)^{-1} d\mu(\lambda). \end{aligned}$$

Let $\tau = \|T(t)A(I + A)^{-1}\| = \|T(t)B(A)\|$. Estimating as in a) gives

$$\|T(t)\varphi(A)(I + A)^{-1}\| \leq 2C\varphi(\tau).$$

The claim follows as in (a). □

Remark 4.5. The inequality (3.12) easily implies that

$$\|f(A^{-1})\| \leq Cf(\|A^{-1}\|),$$

for a constant C independent of the complete Bernstein function f . On the other hand, taking $t_1 = t_2 = \gamma = 0$ in (4.5) we obtain a reversed inequality

$$\|f(A^{-1})\| \geq cf(\|A^{-1}\|).$$

The two inequalities together form an operator counterpart to Proposition 2.4.

Remark 4.6. Theorem 4.3 can be generalized in various ways. For example, if B is any bounded sectorial operator commuting with $T(t)$ for all $t > 0$ (in particular, if $B = B(A)$), then the following version of (4.5) holds:

$$\|T(t_1)B^\gamma f(B)\| \geq c \frac{\|T(t_1+t_2)B^{\gamma+1}\|}{\|T(t_2)B\|} f(\|T(t_2)B\|)$$

for $\gamma > 0$. It also holds for $\gamma \geq -1$ if B is injective and $B^\gamma f(B)$ is a bounded operator (in particular, if $B = A^{-\delta}$ for $0 < \delta \leq 1$).

4.2 Transference from resolvents to semigroups

The final estimate of this section is for bounded semigroups on Hilbert space. The following result shows how the effect of cancelling resolvent growth can be transferred to an estimate for the semigroup itself. When $B = A^{-\alpha}$, the result was obtained in [17, Theorem 2.4]. In our applications the operator B will be a function of the generator A , such as the operator $W_{\alpha,\beta,\ell}(A)$ of Subsection 5.2.

Theorem 4.7. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Hilbert space X , with generator $-A$, and let $B : \text{dom}(A) \rightarrow X$ be a linear operator which is bounded for the graph norm on $\text{dom}(A)$, and such that $T(t)Bx = BT(t)x$ for all $t \geq 0$ and $x \in \text{dom}(A)$. Assume that*

$$\sup \{ \|B(\lambda + A)^{-1}\| : \lambda \in \mathbb{C}_+ \} < \infty. \tag{4.7}$$

Then $T(t)B$ extends to a bounded linear operator (also denoted by $T(t)B$) on X for each $t > 0$, and $\|T(t)B\| = O(t^{-1})$ as $t \rightarrow \infty$.

Proof. Let $x \in \text{dom}(A)$. For a fixed $\tau > 0$ define

$$f_\tau(t) = \begin{cases} T(t)x, & 0 \leq t \leq \tau, \\ 0, & t > \tau, \end{cases}$$

and

$$\varphi_\tau(t) = (BT * f_\tau)(t) = \begin{cases} tBT(t)x, & 0 \leq t \leq \tau, \\ \tau BT(t)x, & t > \tau. \end{cases}$$

Let \widehat{f}_τ and $\widehat{\varphi}_\tau$ be the Laplace transforms of these functions, so that $\widehat{\varphi}_\tau(\lambda) = (\lambda + A)^{-1} \widehat{f}_\tau(\lambda)$ for $\lambda \in \mathbb{C}_+$. By Plancherel's theorem, for $a > 0$,

$$\begin{aligned} \int_{\mathbb{R}} \|\widehat{\varphi}_\tau(a + is)\|^2 ds &= 2\pi \int_{\mathbb{R}} e^{-2at} \|\varphi_\tau(t)\|^2 dt \\ &\geq 2\pi \int_0^\tau t^2 e^{-2at} \|BT(t)x\|^2 dt. \end{aligned}$$

Letting C be the finite supremum in (4.7), we have

$$\begin{aligned} \int_{\mathbb{R}} \|\widehat{\varphi}_\tau(a + is)\|^2 ds &= \int_{\mathbb{R}} \|B(a + is + A)^{-1} \widehat{f}_\tau(a + is)\|^2 ds \\ &\leq C^2 \int_{\mathbb{R}} \|\widehat{f}_\tau(a + is)\|^2 ds \\ &= C^2 2\pi \int_0^\tau e^{-2at} \|T(t)x\|^2 dt, \end{aligned}$$

again by Plancherel's theorem. These two inequalities imply that

$$C^2 \int_0^\tau e^{-2at} \|T(t)x\|^2 dt \geq \int_0^\tau t^2 e^{-2at} \|BT(t)x\|^2 dt.$$

Letting $a \rightarrow 0+$ one gets

$$C^2 \int_0^\tau \|T(t)x\|^2 dt \geq \int_0^\tau t^2 \|BT(t)x\|^2 dt.$$

Let $K = \sup_{t \geq 0} \|T(t)\|$. Then

$$C^2 K^2 \|x\|^2 \geq \frac{1}{\tau} \int_0^\tau t^2 \|BT(t)x\|^2 dt. \quad (4.8)$$

Hence, for any $y \in X$,

$$\begin{aligned} |\langle \tau T(\tau) Bx, y \rangle| &= \left| \frac{2}{\tau} \int_0^\tau t \langle T(t) Bx, T^*(\tau - t)y \rangle dt \right| \\ &\leq \left\{ \frac{2}{\tau} \int_0^\tau t^2 \|T(t) Bx\|^2 dt \right\}^{\frac{1}{2}} \left\{ \frac{2}{\tau} \int_0^\tau \|T^*(\tau - t)y\|^2 dt \right\}^{\frac{1}{2}} \\ &\leq \left\{ \frac{2}{\tau} \int_0^\tau t^2 \|T(t) Bx\|^2 dt \right\}^{\frac{1}{2}} \sqrt{2} K \|y\| \\ &\leq 2CK^2 \|x\| \|y\|. \end{aligned}$$

This implies that $T(\tau)B$ has a bounded extension to X with norm at most $2CK^2/\tau$. \square

Remark 4.8. We do not know whether there is a converse of Theorem 4.7, for example whether (4.7) holds whenever B is a bounded operator on X , commuting with $T(t)$ and satisfying $\|T(t)B\| = O(t^{-1})$ as $t \rightarrow \infty$. There is a result in function theory [38, Lemma 2.5] which says that under these assumptions the boundary function of $\langle B(\cdot + A)^{-1}x, y \rangle$ on $i\mathbb{R}$ lies in $\text{BMO}(i\mathbb{R})$.

The crux of the proof of Theorem 4.7 is the estimate (4.8) showing that the Cesàro means of the scalar function $t \mapsto t^2 \|BT(t)x\|^2$ are bounded. For a positive measurable function boundedness of its Abel means is equivalent to boundedness of its Cesàro means. On Hilbert space, the Abel means of this function are

$$a \int_{\mathbb{R}_+} t^2 e^{-at} \|BT(t)x\|^2 dt = \frac{\alpha}{\pi} \int_{\mathbb{R}} \|B(\alpha + is + A)^{-2}x\|^2 ds, \quad a = 2\alpha > 0,$$

by Plancherel's theorem. Thus the assumption (4.7) can be replaced by

$$\alpha \int_{\mathbb{R}} \|B(\alpha + is + A)^{-2}x\|^2 ds \leq C\|x\|^2, \quad \alpha > 0, x \in X.$$

On Hilbert space, the bounded operator-valued function $B(\cdot + A)^{-1}$ is an $L^2(\mathbb{R}_+; X)$ -Laplace multiplier, as shown by means of Plancherel's Theorem. Theorem 4.7 holds for a C_0 -semigroup on a Banach space X provided that $B(\cdot + A)^{-1}$ is an $L^p(\mathbb{R}_+; X)$ -Laplace multiplier for some $p \in [1, +\infty)$, in the sense that the convolution operator $f \mapsto BT * f$ is bounded on $L^p(\mathbb{R}_+, X)$. The proof is as in Theorem 4.7 with an application of Hölder's inequality replacing the Cauchy-Schwarz inequality.

5 Singularity at infinity

In this section we shall consider a bounded C_0 -semigroup $(T(t))_{t \geq 0}$, with generator $-A$, on a Hilbert space X under the assumption that $\sigma(A) \cap i\mathbb{R}$ is empty. First we recall and elaborate Theorem 1.1, where X is any Banach space but the other assumptions are the same.

The spectral assumption that $\sigma(A) \cap i\mathbb{R}$ is empty is equivalent to the property that

$$\lim_{t \rightarrow \infty} \|T(t)(\omega + A)^{-1}\| = 0 \tag{5.1}$$

for any $\omega \in \rho(-A)$ [6, Theorem 4.4.14]. We choose to take $\omega = 0$.

The rate of decay in (5.1) is closely related to the growth of the resolvent of A on $i\mathbb{R}$. Let M be a function such that

$$\|(is + A)^{-1}\| \leq M(s), \quad s \in \mathbb{R}. \tag{5.2}$$

We shall always make the natural assumption that M is even, so we shall consider M as being a function on \mathbb{R}_+ . It is also natural to assume that M is increasing and continuous.

Define also

$$M_{\log}(s) = M(s)(\log(1 + M(s)) + \log(1 + s)), \quad s \geq 0. \tag{5.3}$$

It is shown in [13, Theorem 1.5] (see also [6, Theorem 4.4.14]) that

$$\|T(t)A^{-1}\| = O\left(\frac{1}{M_{\log}^{-1}(ct)}\right), \quad t \rightarrow \infty, \tag{5.4}$$

for any $c \in (0, 1)$.

The smallest function M satisfying (5.2) and our other assumptions is given by

$$M(s) = \sup \{ \|(ir + A)^{-1}\| : |r| \leq s \}, \quad s \geq 0. \tag{5.5}$$

For this choice of M it is a simple consequence of [13, Proposition 1.3] (see also [6, Theorem 4.4.14]) that there exist constants $c, C > 0$ such that

$$\|T(t)A^{-1}\| \geq \frac{c}{M^{-1}(Ct)} \tag{5.6}$$

for all sufficiently large t . Here we assume that $\lim_{s \rightarrow \infty} M(s) = \infty$ and M^{-1} may be any right inverse of M .

The estimates (5.4) and (5.6) are both valid for bounded semigroups on any Banach space. They raise the question whether, or when, it is possible to improve (5.4) to

$$\|T(t)A^{-1}\| = O\left(\frac{1}{M^{-1}(ct)}\right), \quad t \rightarrow \infty. \quad (5.7)$$

In some cases, for example if $M(s) = e^{\alpha s}$ for $\alpha > 0$, (5.4) and (5.7) are equivalent. In many cases, each estimate is independent of c . In the case when $M(s) = s^\alpha$ for $\alpha > 0$, the two estimates differ by a logarithmic factor. In this case, (5.4) is optimal for arbitrary Banach spaces, but (5.7) holds when X is a Hilbert space [17]. However, for some M one cannot make this improvement even for normal operators on Hilbert space [6, Example 4.4.15]. We give a more detailed analysis of normal semigroups in Subsection 5.1.

In later subsections, we consider cases when X is a Hilbert space and M is regularly varying. In Subsection 5.2 we shall show that if $-A$ generates a bounded C_0 -semigroup on a Banach space X and $\sigma(A) \cap i\mathbb{R}$ is empty, then for regularly varying functions M , the property (5.2) is equivalent to

$$\|(\lambda + A)^{-1}f_M(A)\| \leq C, \quad \lambda \in \mathbb{C}_+, \quad (5.8)$$

where $f_M(A)$ is defined by the extended functional calculus of Theorem 3.7 for an appropriate function f_M related to the classes of Bernstein functions and Stieltjes functions discussed in Subsection 2.1. In Subsections 5.3 and 5.4, we pass from (5.8) towards (5.7).

5.1 Normal semigroups

The following result gives the precise condition on M which is both necessary (if M is defined by (5.5)) and sufficient for (5.7) to be valid for a semigroup of normal operators on Hilbert space. In fact, it holds more generally for any bounded C_0 -semigroup for which the norms of all the associated bounded operators are determined by $\sigma(A)$, that is,

$$\|T(t)r(A)\| = \sup \{|e^{-\lambda t}r(\lambda)| : \lambda \in \sigma(A)\}$$

for every rational function r whose poles are outside $\sigma(A)$ and which is bounded at infinity. This includes multiplication semigroups on L^p -spaces and spaces of continuous functions. We call such C_0 -semigroups *quasi-multiplication* semigroups.

Proposition 5.1. *Let $(T(t))_{t \geq 0}$ be a quasi-multiplication semigroup on a Banach space X with generator $-A$. Assume that $\sigma(A) \subset \mathbb{C}_+$ and $\inf\{\operatorname{Re} \lambda : \lambda \in \sigma(A)\} = 0$. Let M be defined by (5.5) and let $c > 0$. The following are equivalent:*

(i) *There exists C such that*

$$\|T(t)A^{-1}\| \leq \frac{C}{M^{-1}(ct)}, \quad t \geq c^{-1}M(0); \quad (5.9)$$

(ii) There exists B such that

$$\frac{M(\tau)}{M(s)} \geq c \log \left(\frac{\tau}{s} \right) - B, \quad \tau \geq 0, s \geq 1. \quad (5.10)$$

Proof. Note first that $\|T(t)\| = 1$ for all $t \geq 0$, and

$$M(s)^{-1} = \min \{ |\mu - ir| : \mu \in \sigma(A), |r| \leq s \} \rightarrow 0, \quad s \rightarrow \infty.$$

Now (5.9) is equivalent to

$$\frac{e^{-t\alpha}}{|\mu|} \leq \frac{C}{M^{-1}(ct)}, \quad \mu = \alpha + i\beta \in \sigma(A), t \geq c^{-1}M(0).$$

This may be rewritten as

$$t\alpha \geq \log \left(\frac{M^{-1}(ct)}{C|\mu|} \right) \quad (5.11)$$

for all such μ and t .

Assume that (i) holds. Let $t \geq c^{-1}M(0)$, and put $\tau = M^{-1}(ct)$. From (5.11),

$$M(\tau) \geq \frac{c}{\alpha} \log \left(\frac{\tau}{C|\mu|} \right).$$

Given $s \geq 1$, take $\mu = \alpha + i\beta \in \sigma(A)$ such that $M(s)^{-1} = |\mu - ir|$ for some $|r| \leq s$. Then $\alpha \leq k$, where $k = M(0)^{-1}$, and $|\beta| \leq s + k$. So

$$\frac{M(\tau)}{M(s)} \geq \frac{c|\alpha + i(\beta - r)|}{\alpha} \log \left(\frac{\tau}{C(s + 2k)} \right) \geq c \log \left(\frac{\tau}{C(s + 2k)} \right).$$

Since $\log(C(s + 2k)) - \log s \leq \log((2k + 1)C)$ for $s \geq 1$, it follows that (5.10) holds whenever τ is in the range of M^{-1} . For other values of τ one can apply the above with τ replaced by $\tau_n := M^{-1}(M(\tau) + n^{-1}) > \tau$, and let $n \rightarrow \infty$.

Now assume that (ii) holds. Given $t \geq c^{-1}M(0)$ and $\mu = \alpha + i\beta \in \sigma(A)$ with $|\beta| > 1$, take

$$\tau = M^{-1}(ct), \quad s = |\beta|.$$

By (5.10),

$$\frac{ct}{M(|\beta|)} \geq c \log \left(\frac{M^{-1}(ct)}{|\beta|} \right) - B.$$

Rearranging this, using $\alpha M(|\beta|) \geq 1$ and $|\mu| \geq |\beta|$, and putting $C = \exp(B/c)$ gives (5.11), provided that $|\beta| \geq 1$.

If there exist $\mu = \alpha + i\beta \in \sigma(A)$ with $|\beta| \leq 1$, let

$$\varepsilon = \inf \{ \alpha : \alpha + i\beta \in \sigma(A), |\beta| \leq 1 \} > 0.$$

Putting $s = \sqrt{\tau}$ and then $s = 1$ in (5.10) shows that

$$M(\tau) \geq \left(\frac{c}{2} \log \tau - B \right) M(\sqrt{\tau}) \geq \left(\frac{c}{2} \log \tau - B \right)^2 M(1) \geq \frac{c^2 (\log \tau)^2}{5\varepsilon} \quad (5.12)$$

for all sufficiently large τ . Putting $\tau = M^{-1}(ct)$ shows that

$$ct \geq \frac{c^2(\log M^{-1}(ct))^2}{5\varepsilon},$$

and hence

$$\varepsilon t \geq \log M^{-1}(ct)$$

for all sufficiently large t . We can then choose C sufficiently large that

$$\varepsilon t \geq \log \left(\frac{M^{-1}(ct)}{C\varepsilon} \right)$$

whenever $t \geq c^{-1}M(0)$. Then (5.11) holds for $\mu = \alpha + i\beta \in \sigma(A)$ with $|\beta| \leq 1$ and $t \geq c^{-1}M(0)$. Hence (5.9) holds. \square

It is clear that (5.10) implies that M grows at least logarithmically. This corresponds to the elementary fact that, for a quasi-multiplication semigroup, $\|T(t)A^{-1}\|$ cannot decrease faster than exponentially. The estimate (5.12) shows that (5.10) implies that $M(\tau)$ grows at least as fast as $(\log \tau)^2$. More generally, any slowly varying function M fails to satisfy (5.10). Given B and $c > 0$, choose $\lambda = e^{(B+2)/c}$. Then (5.10) implies that

$$\frac{M(\lambda s)}{M(s)} \geq 2, \quad s \geq 1,$$

so M is not slowly varying. In particular, the rate of decay of $\|T(t)A^{-1}\|$ for a quasi-multiplication semigroup cannot be given by (5.9) unless the rate is slower than $\exp(-ct^{1/n})$ for all n .

The following example shows the rate of decay for normal semigroups when M grows logarithmically.

Example 5.2. Let $(T(t))_{t \geq 0}$ be a quasi-multiplication semigroup on a Banach space X , with generator $-A$ such that

$$\sigma(A) = \left\{ \frac{1}{\log s} + is : s \geq 2 \right\}.$$

Then

$$\begin{aligned} \log s \leq \|(is - A)^{-1}\| &= M(s) \leq \log(s+1), & s \geq 2, \\ \frac{1}{M^{-1}(ct)} &\sim e^{-ct}. \end{aligned}$$

However,

$$\|T(t)A^{-1}\| = \sup \left\{ \frac{\exp(-t/\log s)}{s} : s \geq 2 \right\} = e^{-2\sqrt{t}}, \quad t \geq (\log 2)^2.$$

If $M(s) = (1+s)^\alpha$ for some $\alpha > 0$, then (5.10) holds. More generally, if M is regularly varying with index $\alpha > 0$, then M satisfies (5.10).

On the other hand, rapid growth of M does not on its own imply that (5.10) holds. See [6, Example 4.4.15].

5.2 Cancelling resolvent growth

Here we shall show how regularly varying growth of $\|(is + A)^{-1}\|$ as $|s| \rightarrow \infty$ can be cancelled by restricting to the range of a suitable operator. In the case of purely polynomial growth this was achieved by taking a (negative) fractional power of A [50, Lemma 3.2], [17, Lemma 2.3], but we shall need a more complicated function of A .

The following lemma of Phragmén-Lindelöf type may be known, but we have not been able to trace it in the literature. The formulation stated here is more general than is needed for this section, but we shall need the stronger form, or variants of it, when we consider singularities at zero in Sections 7 and 8.

Lemma 5.3. *Let Y be a Banach space and $f : \overline{\mathbb{C}_+} \setminus \{0\} \rightarrow Y$ be a function which is continuous on $\overline{\mathbb{C}_+} \setminus \{0\}$, holomorphic in \mathbb{C}_+ , and bounded on $i\mathbb{R} \setminus \{0\}$. If there exists $C > 0$ such that*

$$\|f(z)\| \leq \frac{C}{\operatorname{Re} z}, \quad z \in \mathbb{C}_+,$$

then f is bounded in $\overline{\mathbb{C}_+} \setminus \{0\}$.

Proof. Let $K = \sup\{\|f(z)\| : z \in i\mathbb{R}, z \neq 0\}$. For fixed r, R with $0 < r < 1 < R$, let

$$A_{r,R} = \{z \in \overline{\mathbb{C}_+} : r \leq |z| \leq R\}.$$

Consider

$$g_{r,R}(z) := \frac{z}{1+z} \left(1 + \frac{z^2}{R^2}\right) \left(1 + \frac{r^2}{z^2}\right) f(z), \quad z \in A_{r,R}.$$

Let

$$\begin{aligned} \gamma_r &= \{z \in \mathbb{C}_+ : |z| = r\}, & \Gamma_R &= \{z \in \mathbb{C}_+ : |z| = R\}, \\ J_{r,R} &= \{is : r \leq |s| \leq R\}, \end{aligned}$$

so that

$$\partial A_{r,R} = \gamma_r \cup \Gamma_R \cup J_{r,R}.$$

If $z \in \gamma_r$ then

$$\left| \frac{z}{1+z} \right| \leq \frac{r}{1-r}, \quad \left| 1 + \frac{z^2}{R^2} \right| \leq 2, \quad \left| 1 + \frac{r^2}{z^2} \right| = \frac{2 \operatorname{Re} z}{r},$$

so that

$$\|g_{r,R}(z)\| \leq \frac{r}{1-r} \frac{4 \operatorname{Re} z}{r} \frac{C}{\operatorname{Re} z} = \frac{4C}{1-r}.$$

Similarly, if $z \in \Gamma_R$ then

$$\left| \frac{z}{1+z} \right| \leq \frac{R}{R-1}, \quad \left| 1 + \frac{z^2}{R^2} \right| = \frac{2 \operatorname{Re} z}{R}, \quad \left| 1 + \frac{r^2}{z^2} \right| \leq 2,$$

thus

$$\|g_{r,R}(z)\| \leq \frac{4C}{R-1}.$$

Finally, if $z \in J_{r,R}$ then

$$\|g_{r,R}(z)\| \leq 4K.$$

Applying the maximum principle to $g_{r,R}$ over $A_{r,R}$, letting $r \rightarrow 0+$ and $R \rightarrow \infty$ and setting $C' = 4 \max(C, K)$, we infer that

$$\sup_{z \in \mathbb{C}_+} \left\| \frac{z}{1+z} f(z) \right\| \leq C'.$$

In particular, f is bounded on $\{z \in \mathbb{C}_+ : |z| \geq 1\}$ and

$$\|f(z)\| \leq \frac{2C'}{|z|}, \quad |z| \leq 1.$$

Now, by applying a standard Phragmén-Lindelöf principle for half-planes [23, Corollary VI.4.2] to $h(z) := f(z^{-1})$ for $z \in \mathbb{C}_+$, we conclude that f is bounded on $\{z \in \mathbb{C}_+ : |z| \leq 1\}$ and this completes the proof. \square

Definition 5.4. Let $-A$ be the generator of a bounded C_0 -semigroup, and assume that A is invertible. Let $\beta \in (0, 1]$, and let ℓ be a slowly varying function such that $g : s \mapsto s^{1-\beta}\ell(s)$ is increasing on \mathbb{R}_+ . Let S_g be the Stieltjes function associated with g . For $\alpha \geq \beta$, define

$$W_{\alpha,\beta,\ell}(A) := A^{-(\alpha-\beta)} \int_{0+}^{\infty} (s+A)^{-1} d(s^{1-\beta}\ell(s)) = A^{-(\alpha-\beta)} S_g(A). \quad (5.13)$$

See Example 2.14 and Remark 3.8, (iii) and (v), for the definition of S_g , the convergence of the integral and compatibility with the extended holomorphic functional calculus.

For $0 < \alpha < \beta$, define

$$W_{\alpha,\beta,\ell}(A) := \int_{0+}^{\infty} A^{\beta-\alpha} (s+A)^{-1} d(s^{1-\beta}\ell(s)) \quad (5.14)$$

$$= \int_0^{\infty} A^{\beta-\alpha} (s+A)^{-2} s^{1-\beta}\ell(s) ds. \quad (5.15)$$

Since A is sectorial and invertible,

$$\|(s+A)^{-1}\| \leq \frac{C}{1+s}, \quad \|A(s+A)^{-1}\| \leq C.$$

The moment inequality (4.2) gives

$$\|A^{\beta-\alpha} (s+A)^{-1}\| \leq \frac{C}{(1+s)^{1+\alpha-\beta}}.$$

Then convergence of the integrals in (5.14) and (5.15) follows from the discussion of Example 2.14. In particular, $W_{\alpha,\beta,\ell}(A)$ is a bounded operator on X , and $A^{-(\beta-\alpha)} W_{\alpha,\beta,\ell}(A) = S_g(A)$, by the product rule (Theorem 3.7(iv)).

When $\ell(s) = 1$ for all $s > 0$, $W_{\alpha,\beta,\ell}(A) = A^{-\alpha}$ for all $\beta \in (0, 1)$. The following result is already known in that special case [50, Lemma 3.2], [17, Lemma 2.3].

Theorem 5.5. *Let $-A$ be the generator of a bounded C_0 -semigroup $(T(t))_{t \geq 0}$ on a Banach space X such that $i\mathbb{R} \subset \rho(A)$. Let ℓ be a slowly varying function on \mathbb{R}_+ , let $\alpha > 0$ and $\beta \in (0, 1]$. Assume that $g : s \mapsto s^{1-\beta}\ell(s)$ is increasing. The following statements are equivalent:*

$$(i) \quad \|(is + A)^{-1}\| = O\left(\frac{|s|^\alpha}{\ell(|s|)}\right), \quad |s| \rightarrow \infty;$$

$$(ii) \quad \sup_{z \in \mathbb{C}_+} \|(z + A)^{-1}W_{\alpha, \beta, \ell}(A)\| < \infty.$$

Proof. For $z \in \rho(-A)$ and $k \in \mathbb{N}$, observe that

$$(z + A)^{-1}A^{-k} = \frac{1}{(-z)^k}(z + A)^{-1} - \sum_{i=0}^{k-1} \frac{1}{(-z)^{i+1}}A^{-(k-i)}. \quad (5.16)$$

We shall show that (i) is equivalent to

$$\|A^{-(\alpha-\beta)}(is + A)^{-1}\| = O\left(\frac{|s|^\beta}{\ell(|s|)}\right), \quad |s| \rightarrow \infty. \quad (5.17)$$

Write $\alpha - \beta = m + \gamma$ where $m \in \mathbb{Z}$, $m \geq -1$ and $0 \leq \gamma < 1$.

Assume that (i) holds. Then

$$\|A(is + A)^{-1}\| = \|I - is(is + A)^{-1}\| = O\left(\frac{|s|^{\alpha+1}}{\ell(|s|)}\right), \quad |s| \rightarrow \infty.$$

By the moment inequality (4.2),

$$\|A^{1-\gamma}(is + A)^{-1}\| = O\left(\frac{|s|^{\alpha+1-\gamma}}{\ell(|s|)}\right), \quad |s| \rightarrow \infty.$$

Using (5.16) for $k = m + 1$, it follows that

$$\begin{aligned} \|A^{-(\alpha-\beta)}(is + A)^{-1}\| &= \|A^{1-\gamma}A^{-(m+1)}(is + A)^{-1}\| \\ &= O\left(\frac{|s|^{\alpha+1-\gamma-(m+1)}}{\ell(|s|)}\right) \\ &= O\left(\frac{|s|^\beta}{\ell(|s|)}\right), \quad |s| \rightarrow \infty. \end{aligned}$$

Conversely, if $\alpha \geq \beta$, (5.17) implies that

$$\|A^{-(\alpha-\beta-1)}(is + A)^{-1}\| = O\left(\frac{|s|^{\beta+1}}{\ell(|s|)}\right), \quad |s| \rightarrow \infty.$$

Since $m = (1 - \gamma)(\alpha - \beta) + \gamma(\alpha - \beta - 1)$ and $(1 - \gamma)\beta + \gamma(\beta + 1) = \alpha - m$, the moment inequality (4.2) gives

$$\|A^{-m}(is + A)^{-1}\| = O\left(\frac{|s|^{\alpha-m}}{\ell(|s|)}\right), \quad |s| \rightarrow \infty.$$

Then (5.16) for $k = m$ gives (i).

If $\alpha < \beta$, then $m = -1$ and (5.17) implies that

$$\begin{aligned} \|A^{-(\alpha-\beta+1)}(is + A)^{-1}\| &= \frac{\|A^{-(\alpha-\beta+1)} - A^{-(\alpha-\beta)}(is + A)^{-1}\|}{|s|} \\ &= O\left(\frac{|s|^{\beta-1}}{\ell(|s|)}\right), \quad |s| \rightarrow \infty. \end{aligned}$$

Since $0 = \gamma(\alpha - \beta) + (1 - \gamma)(\alpha - \beta + 1)$ and $\gamma\beta + (1 - \gamma)(\beta - 1) = \alpha$, the moment inequality (4.2) gives (i). This completes the proof that (i) is equivalent to (5.17).

Next, we consider the Stieltjes function S_g associated with g , defined as in Example 2.14. By Karamata's Theorem 2.15(a) with $\sigma = \beta$,

$$S_g(\lambda) = O(\lambda^{-\beta}\ell(\lambda)), \quad \lambda \rightarrow \infty. \quad (5.18)$$

By Proposition 2.4,

$$|S_g(-is)| = O(|s|^{-\beta}\ell(|s|)), \quad |s| \rightarrow \infty. \quad (5.19)$$

Now we observe that

$$\begin{aligned} (is + A)^{-1}W_{\alpha,\beta,\ell}(A) &= \int_{0+}^{\infty} (is + A)^{-1}A^{-(\alpha-\beta)}(\lambda + A)^{-1}d(\lambda^{1-\beta}\ell(\lambda)) \\ &= A^{-(\alpha-\beta)}(is + A)^{-1} \int_{0+}^{\infty} \frac{d(\lambda^{1-\beta}\ell(\lambda))}{\lambda - is} \\ &\quad - \int_{0+}^{\infty} \frac{1}{\lambda - is}A^{-(\alpha-\beta)}(\lambda + A)^{-1}d(\lambda^{1-\beta}\ell(\lambda)) \\ &= S_g(-is)A^{-(\alpha-\beta)}(is + A)^{-1} \\ &\quad - \int_{0+}^{\infty} \frac{1}{\lambda - is}A^{-(\alpha-\beta)}(\lambda + A)^{-1}d(\lambda^{1-\beta}\ell(\lambda)). \end{aligned} \quad (5.20)$$

The last integral is bounded, uniformly for $|s| \geq 1$, by the same argument as for (5.14) together with the fact that $|\lambda - is| \geq 1$.

Now assume that (i) holds, so (5.17) holds. By (5.17) and (5.19),

$$\|S_g(-is)A^{-(\alpha-\beta)}(is + A)^{-1}\| \leq C, \quad |s| \geq 1.$$

Hence $\|(is + A)^{-1}W_{\alpha,\beta,\ell}(A)\|$ is bounded for $|s| \geq 1$, and then for all real s . We now apply Lemma 5.3, with

$$f(z) := (z + A)^{-1}W_{\alpha,\beta,\ell}(A),$$

and we deduce that (ii) holds.

Conversely, assume that (ii) holds. Letting $z \rightarrow is$ and using (5.20) shows that $\|S_g(-is)A^{-(\alpha-\beta)}(is + A)^{-1}\|$ is bounded for $s \in \mathbb{R}$, $|s| > 1$. Proposition 2.4 and Theorem 2.15(a) show that (5.17) holds. Then (i) follows. \square

5.3 Resolvent growth slower than s^α

In this and the next subsection we consider cases when $-A$ generates a bounded C_0 -semigroup $(T(t))_{t \geq 0}$ on a Hilbert space X , $\sigma(A) \cap i\mathbb{R}$ is empty and

$$\|(is + A)^{-1}\| = O\left(\frac{|s|^\alpha}{\ell(|s|)}\right), \quad |s| \rightarrow \infty, \quad (5.21)$$

where ℓ is slowly varying and monotonic. The upper bound for $\|T(t)A^{-1}\|$ in (5.4) is then valid for $M(s) = Cs^\alpha/\ell(s)$. Then

$$M_{\log}(s) \sim C(1 + \alpha) \frac{s^\alpha \log s}{\ell(s)} = C(1 + \alpha)(M.\log)(s),$$

so we may replace M_{\log} by $M.\log$ in (5.4).

If we put $k(s) = 1/\ell(s^{1/\alpha})$, then Proposition 2.11 gives

$$(M.\log)^{-1}(t) \sim t^{1/\alpha}(k.\log)^\#(t)^{1/\alpha}.$$

Thus we obtain that

$$\|T(t)A^{-1}\| = O\left(\left(t(k.\log)^\#(t)\right)^{-1/\alpha}\right), \quad t \rightarrow \infty. \quad (5.22)$$

When $k.\log$ is dB-symmetric, this becomes

$$\|T(t)A^{-1}\| = O\left(\frac{(\log t)^{1/\alpha}}{t\ell(t^{1/\alpha})^{1/\alpha}}\right), \quad t \rightarrow \infty. \quad (5.23)$$

On the other hand, if we have

$$\|(is + A)^{-1}\| \geq c \frac{|s|^\alpha}{\ell(|s|)}$$

for large $|s|$, then (5.6) and Proposition 2.11(b) give

$$\|T(t)A^{-1}\| \geq \frac{c}{tk^\#(t^{1/\alpha})^{1/\alpha}}.$$

Thus, assuming (5.21), the optimal result would be to establish that

$$\|T(t)A^{-1}\| = O\left(\left(tk^\#(t)\right)^{-1/\alpha}\right), \quad t \rightarrow \infty, \quad (5.24)$$

or, assuming that ℓ is dB-symmetric,

$$\|T(t)A^{-1}\| = O\left(\left(t\ell(t^{1/\alpha})\right)^{-1/\alpha}\right), \quad t \rightarrow \infty. \quad (5.25)$$

We now give one of our main results showing that (5.25) holds when ℓ is increasing, i.e., $\|(is + A)^{-1}\|$ grows slightly slower than $|s|^\alpha$. The case when ℓ is decreasing will be considered in Subsection 5.4.

Theorem 5.6. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Hilbert space X , with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R}$ is empty, and that*

$$\|(is + A)^{-1}\| = O\left(\frac{|s|^\alpha}{\ell(|s|)}\right), \quad |s| \rightarrow \infty,$$

where $\alpha > 0$ and ℓ is increasing and slowly varying. Then

$$\|T(t)A^{-1}\| = O\left(\left(t\ell(t^{1/\alpha})\right)^{-1/\alpha}\right), \quad t \rightarrow \infty. \quad (5.26)$$

Proof. We can assume that $T(t) \neq 0$ for each $t > 0$. First we note a known upper bound for $\|T(t)A^{-1}\|$. Since $\|(is + A)^{-1}\| = O(|s|^\alpha)$, [17, Theorem 2.4] gives

$$\|T(t)A^{-1}\| \leq \frac{C}{t^{1/\alpha}}. \quad (5.27)$$

By Theorem 5.5,

$$\|(\lambda + A)^{-1}W_{\alpha,1,\ell}(A)\| \leq C, \quad \lambda \in \mathbb{C}_+.$$

By Theorem 4.7,

$$\|T(t)W_{\alpha,1,\ell}(A)\| \leq \frac{C}{t}, \quad t > 0.$$

Let S_ℓ be the Stieltjes function associated with ℓ (Example 2.14) and let f_ℓ be the complete Bernstein function defined by

$$f_\ell(s) = S_\ell(1/s), \quad s > 0.$$

Using (5.13) and Remark 3.8(v), we have

$$\|T(t)A^{-(\alpha-1)}f_\ell(A^{-1})\| \leq \frac{C}{t}, \quad t > 0.$$

By Theorem 4.3(a), with $\gamma = 1 - \alpha$,

$$f_\ell(\|T(t)A^{-1}\|) \leq \frac{C\|T(t)A^{-1}\|}{t\|T(2t)A^{-\alpha}\|}.$$

Letting $f_{\alpha,\ell}(s) = s^{\alpha-1}f_\ell(s)$, we have

$$f_{\alpha,\ell}(\|T(t)A^{-1}\|) \leq \frac{C\|T(t)A^{-1}\|^\alpha}{t\|T(2t)A^{-\alpha}\|}. \quad (5.28)$$

By Theorem 2.15(a) with $g = \ell$ and $\sigma = 1$, $S_\ell(s) \sim s^{-1}\ell(s)$ as $s \rightarrow \infty$. Hence $f_{\alpha,\ell}(s) \sim s^\alpha\ell(1/s)$ as $s \rightarrow 0+$. Let $k(s) = 1/\ell(s^{1/\alpha})$. Then

$$f_{\alpha,\ell}(s) \sim \frac{s^\alpha}{k(s^{-\alpha})}, \quad s \rightarrow 0+,$$

By Proposition 2.11(c),

$$f_{\alpha,\ell}^{-1}(s) \sim \left(\frac{s}{k^\#(1/s)} \right)^{1/\alpha}, \quad s \rightarrow 0+.$$

If the right-hand side of (5.28) is sufficiently small, it follows that

$$\|T(t)A^{-1}\| \leq \frac{C\|T(t)A^{-1}\|}{(tL(t)\|T(2t)A^{-\alpha}\|)^{1/\alpha}}, \quad (5.29)$$

where

$$L(t) = k^\# \left(t \frac{\|T(2t)A^{-\alpha}\|}{\|T(t)A^{-1}\|^\alpha} \right). \quad (5.30)$$

Let $\psi(s) = (sk^\#(s))^{1/\alpha}$. Since ψ is regularly varying with positive index, we can choose $k^\#$ so that ψ is strictly increasing and continuous (see the remarks following Definition 2.9; in fact, we can choose $k^\#$ to be increasing, since k is decreasing). Then

$$\begin{aligned} \frac{1}{\|T(t)A^{-1}\|} &\geq ct^{1/\alpha} \frac{\|T(2t)A^{-\alpha}\|^{1/\alpha}}{\|T(t)A^{-1}\|} k^\# \left(t \frac{\|T(2t)A^{-\alpha}\|}{\|T(t)A^{-1}\|^\alpha} \right)^{1/\alpha} \\ &= c\psi \left(t \frac{\|T(2t)A^{-\alpha}\|}{\|T(t)A^{-1}\|^\alpha} \right). \end{aligned} \quad (5.31)$$

If the right-hand side of (5.28) is bounded away from zero, then (5.31) also holds for some $c > 0$, since $\|T(t)A^{-1}\|$ is bounded and ψ is bounded on bounded intervals. Hence (5.31) holds for all $t > 0$, for some $c > 0$.

Since ψ is (asymptotically equivalent to) an increasing function,

$$t \frac{\|T(2t)A^{-\alpha}\|}{\|T(t)A^{-1}\|^\alpha} \leq \psi^{-1} \left(\frac{C}{\|T(t)A^{-1}\|} \right).$$

By Proposition 2.11(b),

$$\psi^{-1}(s) \sim s^\alpha k^{\#\#}(s^\alpha) \sim \frac{s^\alpha}{\ell(s)}, \quad s \rightarrow \infty,$$

so

$$\|T(2t)A^{-\alpha}\| \leq \frac{C}{t\ell(\|T(t)A^{-1}\|^{-1})}, \quad (5.32)$$

for large t and then for all $t > 0$. Then (5.27) gives

$$\|T(2t)A^{-\alpha}\| \leq \frac{C}{t\ell(t^{1/\alpha})}.$$

Applying Lemma 4.2 with $B = A^{-1}$, $\gamma = \alpha$ and $\delta = 1$, it follows that

$$\|T(t)A^{-1}\| \leq C\|T(ct)A^{-\alpha}\|^{1/\alpha} \leq \frac{C}{(t\ell(t^{1/\alpha}))^{1/\alpha}}.$$

□

Corollary 5.7. *In addition to the assumptions of Theorem 5.6, assume that ℓ is dB-symmetric. Then*

$$\|T(t)A^{-1}\| = O\left(\frac{1}{M^{-1}(t)}\right), \quad t \rightarrow \infty,$$

where M^{-1} is any asymptotic inverse of $s^\alpha/\ell(s)$.

Proof. By Proposition 2.11(b),

$$M^{-1}(t) \sim t^{1/\alpha} k^\#(t)^{1/\alpha},$$

where $k(t) = 1/\ell(t^{1/\alpha})$. By Lemma 2.12, $k^\#(t) \sim \ell(t^{1/\alpha})$ if (and only if) ℓ is dB-symmetric. \square

Remark 5.8. Corollary 5.7 establishes the optimal estimate (5.7) when ℓ is dB-symmetric. However the upper bound (5.26) is not as sharp as (5.7) when ℓ is not dB-symmetric. Under the assumptions of Theorem 5.6 the proof takes the upper bound (5.27) from [17] and improves it to the upper bound (5.26). Given an estimate

$$\|T(t)A^{-1}\| = O\left((tm(t))^{-1/\alpha}\right), \quad t \rightarrow \infty, \quad (5.33)$$

where m is increasing and slowly varying, under the same assumptions the same argument shows that

$$\|T(t)A^{-1}\| = O\left(\left(t\ell(t^{1/\alpha}m(t)^{1/\alpha})\right)^{-1/\alpha}\right), \quad t \rightarrow \infty.$$

This process can be iterated, so one obtains (5.33) for each of the following functions m :

$$m(t) = 1, \ell(t^{1/\alpha}), \ell(t^{1/\alpha}\ell(t^{1/\alpha})^{1/\alpha}), \dots$$

In some cases this process stabilises (up to asymptotic equivalence) after a finite number of iterations at the optimal estimate in Corollary 5.7. This is analogous to Békéssy's method of finding the de Bruijn conjugate of many slowly varying functions [15, Proposition 2.3.5] (see Example 5.10).

Remark 5.9. An alternative to the known upper bound (5.27) is the estimate (5.22). In the context of Theorem 5.6, Lemma 2.13(ii) shows that this gives

$$\|T(t)A^{-1}\| = O\left(\left(\frac{\log t}{tk^\#(t)}\right)^{1/\alpha}\right), \quad t \rightarrow \infty,$$

where $k(s) = 1/\ell(s^{1/\alpha})$. This is (5.33) with

$$m(t) = \frac{k^\#(t)}{\log t}.$$

Following Remark 5.8, we obtain

$$\|T(t)A^{-1}\| = O\left(\left(t\ell\left(\left(\frac{tk^\#(t)}{\log t}\right)^{1/\alpha}\right)\right)^{-1/\alpha}\right), \quad t \rightarrow \infty.$$

In many cases, this is asymptotically equivalent to the optimal estimate (5.24).

Example 5.10. We take $\alpha = 1$ (for simplicity of presentation) and $\ell(s) = \exp((\log s)^\beta)$ where $0 < \beta < 1$ (see Example 2.10). In this case, the process in Remark 5.8, starting from $m(t) = 1$, stabilises at the optimal estimate

$$\|T(t)A^{-1}\| \leq \frac{C}{M^{-1}(t)} \sim \frac{C}{t(1/\ell)^\#(t)}. \quad (5.34)$$

When $1/2 < \beta < 3/4$, the process stabilises after two iterations at the optimal estimate:

$$\|T(t)A^{-1}\| \leq \frac{C}{t} \exp(-((\log t)^\beta + \beta(\log t)^{2\beta-1})).$$

When $(n-1)/n < \beta < n/(n+1)$, n iterations are needed.

Starting from

$$m(t) = \frac{(1/\ell)^\#(t)}{\log t},$$

as in Remark 5.9, the process stabilises at the optimal estimate (5.34) after one iteration for every $\beta \in (0, 1)$.

Remark 5.11. If (5.6) and (5.7) both hold for some regularly varying function M of index $\alpha > 0$, then the ratio

$$\frac{\|T(2t)A^{-1}\|}{\|T(t)A^{-1}\|} \quad (5.35)$$

is bounded away from zero (it is bounded above, for any bounded semigroup).

On the other hand, if (5.35) is bounded away from zero, then so is

$$\frac{\|T(2t)A^{-\alpha}\|}{\|T(t)A^{-1}\|^\alpha},$$

by Lemma 4.2. Then the function L in the proof of Theorem 5.6 is asymptotically equivalent to $k^\#$, and one can then pass easily from (5.30) to (5.7). This argument would not require the assumption that ℓ is dB-symmetric, and it would not use Theorem 4.3 or (5.4). However we do not see any way to prove directly that (5.35) is bounded away from zero.

The assumptions of Theorem 5.6 are not sufficient to ensure that (5.35) is bounded away from zero. However, we may assume in addition that

$$\|(is + A)^{-1}\| \geq cM(|s|). \quad (5.36)$$

If one can prove that (5.7) holds under this additional assumption, then it can be deduced that (5.7) holds without the additional assumption. This follows by means of a direct sum argument which is used in the proof of Theorem 5.12 below.

Under the assumptions of Theorem 5.6 and (5.36), we can obtain from (5.4) and (5.6) that

$$\frac{\|T(2t)A^{-\alpha}\|}{\|T(t)A^{-1}\|^\alpha} \geq \frac{c}{(\log t)^\alpha}.$$

It then follows from (5.29) that

$$\|T(t)A^{-1}\| = O\left(\left(tk^\#\left(\frac{t}{(\log t)^\alpha}\right)\right)^{-1/\alpha}\right), \quad t \rightarrow \infty.$$

The direct sum argument can then be used to show that this estimate holds under the assumptions of Theorem 5.6, without (5.36). In general this estimate is slightly worse than (5.26), but in many cases, for example $\ell(s) = (\log s)^\beta$, one can recover the estimate of Corollary 5.7.

5.4 Resolvent growth faster than s^α

Now we consider the case when (5.21) holds with ℓ decreasing so $\|(is + A)^{-1}\|$ grows slightly faster than $|s|^\alpha$. Unfortunately, our result is not quite optimal in this case as a logarithmic term still appears in (5.37) below. However it does improve (5.22) and (5.23) as the logarithm has an arbitrarily small power. If ℓ is dB-symmetric, then $k^\#(t) = \ell(t^{1/\alpha})$ (see the proof of Corollary 5.7), so (5.37) is directly comparable with (5.23) and (5.25).

Theorem 5.12. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Hilbert space X , with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R}$ is empty, and that*

$$\|(is + A)^{-1}\| = O\left(\frac{|s|^\alpha}{\ell(|s|)}\right), \quad |s| \rightarrow \infty,$$

where $\alpha > 0$ and ℓ is decreasing and slowly varying. Then, for any $\varepsilon > 0$,

$$\|T(t)A^{-1}\| = O\left(\frac{(\log t)^\varepsilon}{(tk^\#(t))^{1/\alpha}}\right), \quad t \rightarrow \infty, \quad (5.37)$$

where $k(t) = 1/\ell(t^{1/\alpha})$ and $k^\#$ is the de Bruijn conjugate of k .

Proof. Much of the proof is similar to Theorem 5.6. We can assume that $\|T(t)\| \neq 0$ for each $t > 0$. Given $\varepsilon \in (0, \alpha^{-2})$, let $\beta = 1 - \varepsilon\alpha^2 \in (0, 1)$. By replacing ℓ by an asymptotically equivalent, decreasing function, we may assume, without loss, that $g : s \mapsto s^{1-\beta}\ell(s)$ is increasing on \mathbb{R}_+ . Let S_ℓ be the Stieltjes function associated with ℓ , and let

$$f_\ell(s) = S_\ell(1/s), \quad s > 0.$$

Then f_ℓ is a complete Bernstein function, and

$$W_{\alpha, \beta, \ell}(A) = A^{-(\alpha-\beta)} f_\ell(A^{-1}).$$

By Theorem 5.5,

$$\sup_{z \in \mathbb{C}_+} \|(z + A)^{-1} W_{\alpha, \beta, \ell}(A)\| < \infty.$$

By Theorem 4.7,

$$\|T(t)A^{-(\alpha-\beta)} f_\ell(A^{-1})\| \leq \frac{C}{t}, \quad t > 0.$$

By Theorem 4.3(b), with $\gamma = \beta - \alpha$,

$$f_\ell(\|T(t)A^{-1}\|) \leq \frac{C\|T(t)A^{-1}\|}{t\|T(2t)A^{-(\alpha-\beta+1)}\|}.$$

Letting $f_{\alpha,\ell}(s) = s^{\alpha-\beta}f_\ell(s)$, we have

$$f_{\alpha,\ell}(\|T(t)A^{-1}\|) \leq \frac{C\|T(t)A^{-1}\|^{\alpha-\beta+1}}{t\|T(2t)A^{-(\alpha-\beta+1)}\|}. \quad (5.38)$$

By Theorem 2.15(a) with $g(\lambda) = \lambda^{1-\beta}\ell(\lambda)$ and $\sigma = \beta$, $S_\ell(s) \sim s^{-\beta}\ell(s)$ as $s \rightarrow \infty$. Hence

$$f_{\alpha,\ell}(s) \sim s^\alpha\ell(1/s) = \frac{s^\alpha}{k(s^{-\alpha})}, \quad s \rightarrow 0+.$$

By Proposition 2.11(c),

$$f_{\alpha,\ell}^{-1}(s) \sim \left(\frac{s}{k^\#(1/s)}\right)^{1/\alpha}, \quad s \rightarrow 0+.$$

If the right-hand side of (5.38) is small, it follows that

$$\|T(t)A^{-1}\| \leq \frac{C\|T(t)A^{-1}\|^{1+(1-\beta)/\alpha}}{(tL(t)\|T(2t)A^{-(\alpha-\beta+1)}\|)^{1/\alpha}},$$

where

$$L(t) = k^\# \left(t \frac{\|T(2t)A^{-(\alpha-\beta+1)}\|}{\|T(t)A^{-1}\|^{\alpha-\beta+1}} \right).$$

Let $\psi(s) = (sk^\#(s))^{1/\alpha}$. Since ψ is regularly varying with positive index, we can choose $k^\#$ so that ψ is strictly increasing and continuous (see the remarks following Definition 2.9; in fact, we can also choose $k^\#$ to be decreasing, since k is increasing). Then

$$\begin{aligned} \frac{1}{\|T(t)A^{-1}\|} &\geq ct^{1/\alpha} \frac{\|T(2t)A^{-(\alpha-\beta+1)}\|^{1/\alpha}}{\|T(t)A^{-1}\|^{1+(1-\beta)/\alpha}} L(t)^{1/\alpha} \\ &= c\psi \left(t \frac{\|T(2t)A^{-(\alpha-\beta+1)}\|}{\|T(t)A^{-1}\|^{\alpha-\beta+1}} \right). \end{aligned} \quad (5.39)$$

If the right-hand side of (5.38) is bounded away from zero, then (5.39) also holds for some $c > 0$, since $\|T(t)A^{-1}\|$ is bounded and ψ is bounded on bounded intervals. Hence (5.39) holds for all $t > 0$, for some $c > 0$.

Since ψ is strictly increasing,

$$t \frac{\|T(2t)A^{-(\alpha-\beta+1)}\|}{\|T(t)A^{-1}\|^{\alpha-\beta+1}} \leq \psi^{-1} \left(\frac{C}{\|T(t)A^{-1}\|} \right).$$

By Proposition 2.11(b),

$$\psi^{-1}(s) \sim s^\alpha k^{\#\#}(s^\alpha) \sim \frac{s^\alpha}{\ell(s)},$$

so

$$\|T(2t)A^{-(\alpha-\beta+1)}\| \leq \frac{C\|T(t)A^{-1}\|^{1-\beta}}{t\ell(\|T(t)A^{-1}\|^{-1})}, \quad (5.40)$$

for large t . By Lemma 4.2 with $B = A^{-1}$, $\gamma = 1$ and $\delta = \alpha - \beta + 1$,

$$\|T(2Kt)A^{-1}\|^{1+(1-\beta)/\alpha} \leq C\|T(2t)A^{-(\alpha-\beta+1)}\|^{1/\alpha}$$

for some K . Hence

$$\|T(2Kt)A^{-1}\| \leq \frac{C}{(t\ell(\|T(t)A^{-1}\|^{-1}))^{1/\alpha}} \left(\frac{\|T(t)A^{-1}\|}{\|T(2Kt)A^{-1}\|} \right)^{\frac{1-\beta}{\alpha}}.$$

Now we shall temporarily assume that

$$cM(s) \leq \|(is + A)^{-1}\| \leq M(s), \quad (5.41)$$

where $M(s) = s^\alpha/\ell(s)$ for large s . We know from (5.4) and (5.6) that

$$\frac{c}{M^{-1}(Ct)} \leq \|T(t)A^{-1}\| \leq \frac{C}{(M.\log)^{-1}(ct)}.$$

Since ℓ is decreasing and slowly varying and M^{-1} is regularly varying,

$$\frac{1}{\ell(\|T(t)A^{-1}\|^{-1})} \leq \frac{1}{\ell(c^{-1}M^{-1}(Ct))} \sim \frac{1}{\ell(M^{-1}(t))}.$$

Hence

$$\|T(2Kt)A^{-1}\| \leq \frac{C}{(t\ell(M^{-1}(t)))^{1/\alpha}} \left(\frac{M^{-1}(Ct)}{(M.\log)^{-1}(ct)} \right)^{\frac{1-\beta}{\alpha}}$$

for large t . Since M is regularly varying of index α , M^{-1} and $(M.\log)^{-1}$ are both regularly varying of index $1/\alpha$. Since ℓ is decreasing, we can apply Lemma 2.13(ii), with ℓ replaced by $1/\ell$ and $\delta = \alpha$, to obtain that for $a, b > 0$ there exists $c_{a,b} > 0$ such that

$$\frac{(M.\log)^{-1}(at)}{M^{-1}(bt)} \geq \frac{c_{a,b}}{(\log t)^{1/\alpha}}$$

for large t . Moreover,

$$M^{-1}(t) \sim t^{1/\alpha} k^\#(t)^{1/\alpha},$$

and $k(tk^\#(t)) \sim (k^\#(t))^{-1}$, by definition of $k^\#$, so

$$\ell(M^{-1}(t)) \sim \ell(t^{1/\alpha} k^\#(t)^{1/\alpha}) \sim k^\#(t).$$

Hence

$$\|T(2Kt)A^{-1}\| \leq \frac{C(\log t)^\varepsilon}{(tk^\#(t))^{1/\alpha}}$$

for large t . Replacing t by $t/(2K)$ and changing the value of C we obtain

$$\|T(t)A^{-1}\| \leq \frac{C(\log t)^\varepsilon}{(tk^\#(t))^{1/\alpha}}$$

for large t .

Now we no longer assume (5.41), and we obtain the same result by means of an artificial device. Let $(S(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Hilbert space Y , with generator $-B$ which does satisfy (5.41). For example, we can take a normal semigroup on a Hilbert space with

$$\sigma(B) = \{2M(s)^{-1} + is : s \geq 2\}.$$

We then consider the C_0 -semigroup

$$T(t) \oplus S(t)$$

on $X \oplus Y$. This satisfies the assumption (5.41), so we find that

$$\|T(t)A^{-1}\| \leq \|T(t)A^{-1} \oplus S(t)B^{-1}\| \leq \frac{C(\log t)^\varepsilon}{(tk^\#(t))^{1/\alpha}}.$$

□

Remark 5.13. A variation of the proof above proceeds from (5.32) by the method of Remark 5.9. This uses the estimate (5.22) in the form

$$\|T(t)A^{-1}\|^{-1} \geq ct^{1/\alpha}(k \cdot \log)^\#(t)^{1/\alpha}.$$

Since the function $s \mapsto s^{1-\beta}\ell(s)$ is asymptotically equivalent to an increasing function,

$$\frac{\ell(\|T(t)A^{-1}\|^{-1})}{\|T(t)A^{-1}\|^{1-\beta}} \geq ct^{(1-\beta)/\alpha}(k \cdot \log)^\#(t)^{(1-\beta)/\alpha} \ell(t^{1/\alpha}(k \cdot \log)^\#(t)^{1/\alpha}).$$

By (5.40),

$$\|T(2t)A^{-(\alpha-\beta+1)}\| \leq \frac{C}{t^{(\alpha-\beta+1)/\alpha}(k \cdot \log)^\#(t)^{(1-\beta)/\alpha} \ell(t^{1/\alpha}(k \cdot \log)^\#(t)^{1/\alpha})}.$$

Applying Lemma 4.2 with $B = A^{-1}$, $\gamma = \alpha - \beta + 1$ and taking β arbitrarily close to 1, it follows that

$$\|T(t)A^{-1}\| \leq \frac{C_\varepsilon}{\left[t(k \cdot \log)^\#(t)^\varepsilon \ell(t^{1/\alpha}(k \cdot \log)^\#(t)^{1/\alpha})^{1+\varepsilon}\right]^{1/\alpha}}.$$

We do not expect that this gives better estimates than the simpler (5.37).

6 Singularity at zero: general results

6.1 Preliminary background

In this section we treat the rates of decay of the derivatives $-T(t)Ax$ for $x \in \text{dom}(A)$, where $(T(t))_{t \geq 0}$ is a bounded C_0 -semigroup on a Banach space or a Hilbert space X , with generator $-A$. In other words, we study the orbits of $(T(t))_{t \geq 0}$ starting at points $y = Ax$ in the range of A . We shall see that such decay corresponds to properties of the resolvent of A permitting a singularity at 0 but requiring boundedness at infinity on the imaginary axis.

If the decay of such orbits is uniform for $\|x\| \leq 1$, the semigroup must be eventually differentiable, i.e., for sufficiently large t , $T(t)$ maps X into $\text{dom}(A)$ and $AT(t) \in \mathcal{L}(X)$.

For an eventually differentiable, bounded C_0 -semigroup on a Banach space, Arendt and Prüss [7, Theorem 3.10] (see [6, Theorem 4.4.16]) showed that

$$\lim_{t \rightarrow \infty} \|AT(t)\| = 0 \quad (6.1)$$

if and only if

$$\sigma(A) \cap i\mathbb{R} \subset \{0\}. \quad (6.2)$$

In fact the Arendt-Prüss theorem follows from (the corollary of) the Katznelson-Tzafriri theorem for C_0 -semigroups which we recall below (see also [67], [28], [10] and the survey [20]). Let E be a closed subset of \mathbb{R} . A function $f \in L^1(\mathbb{R})$ is said to be of *spectral synthesis with respect to E* if there is a sequence (f_n) in $L^1(\mathbb{R})$ such that $\lim_{n \rightarrow \infty} \|f - f_n\|_{L^1} = 0$ and, for each n , $\mathcal{F}f_n$ vanishes in a neighbourhood of E .

The closed subset $E \subset \mathbb{R}$ is said to be of *spectral synthesis* if every function $f \in L^1(\mathbb{R})$ whose Fourier transform $\mathcal{F}f$ vanishes on E is of spectral synthesis with respect to E . Any countable closed subset of \mathbb{R} is of spectral synthesis [46, p. 230].

Theorem 6.1. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Banach space X , with generator $-A$. Let $f \in L^1(\mathbb{R}_+)$ be of spectral synthesis with respect to $E := i\sigma(-A) \cap \mathbb{R}$. Then*

$$\lim_{t \rightarrow \infty} \|T(t)\hat{f}(T)\| = 0, \quad (6.3)$$

where

$$\hat{f}(T)x = \int_0^\infty f(t)T(t)x \, dt, \quad x \in X.$$

Conversely, if $f \in L^1(\mathbb{R}_+)$ and (6.3) holds, then $\mathcal{F}f = 0$ on E .

Corollary 6.2. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Banach space X with generator $-A$. The following statements are equivalent.*

- (i) $\lim_{t \rightarrow \infty} \|T(t)A(I + A)^{-2}\| = 0$;
- (ii) $\sigma(A) \cap i\mathbb{R} \subset \{0\}$.

Proof. This follows from applying the Katznelson-Tzafriri theorem, with

$$f(t) = e^{-t} - te^{-t}, \quad t \geq 0.$$

Then

$$\mathcal{F}f(s) = is(1 + is)^{-2}, \quad s \in \mathbb{R}; \quad \hat{f}(T) = A(I + A)^{-2}.$$

Since $\mathcal{F}f(0) = 0$ and $\{0\}$ is a set of spectral synthesis, f is of spectral synthesis with respect to $\{0\}$. \square

Remark 6.3. If $(T(t))_{t \geq 0}$ is eventually differentiable, then $(1 + A)^2T(\tau)$ is a bounded operator for some $\tau > 0$. Moreover,

$$AT(t + \tau) = (I + A)^2T(\tau)T(t)A(I + A)^{-2},$$

so Corollary 6.2(i) is equivalent to (6.1) in this case. Thus the Arendt-Prüss theorem follows from the Katznelson-Tzafriri theorem.

The range of $A(I + A)^{-2}$ is $\text{ran}(A) \cap \text{dom}(A)$ (Proposition 3.10(i)), and Corollary 6.2 describes decay of orbits on that space in an appropriate uniform sense. We defer consideration of the rate of convergence in Corollary 6.2 (i) until Section 8. In this and the next section, we consider instead the question whether the decay of orbits is uniform with respect to the graph norm on $\text{dom}(A)$, i.e., whether

$$\limsup_{t \rightarrow \infty} \{ \|T(t)Ax\| : x \in \text{dom}(A), \|x\|_{\text{dom}(A)} = 1 \} = 0.$$

This can be reformulated as

$$\lim_{t \rightarrow \infty} \|T(t)A(\omega + A)^{-1}\| = 0, \tag{6.4}$$

for any $\omega \in \rho(-A)$. This property is intermediate between (6.1) and the statements of Corollary 6.2, and it is independent of the choice of ω . When A is invertible, (6.4) is equivalent to the much studied notion of exponential stability, i.e., $\lim_{t \rightarrow \infty} \|T(t)\| = 0$, for which the rate of decay is always at least exponential (see [6, Chapter 5]). Thus we are interested only in cases when $0 \in \sigma(A)$.

Our goal in this section and Section 7 is to develop a framework for decay of the form (6.4) in terms of the spectrum and resolvent of the generator in similar form to the one from Section 5. However, the decay given by (6.4) is much less studied in the literature. Thus we shall prove several statements clarifying the limitations imposed by (6.4) and the consequences following from the decay of orbits in this sense.

In the later subsections, we shall assume that the semigroup is bounded, but first it is instructive to make a few remarks relating to that assumption. We begin by observing that (6.4) implies (6.2), without the assumption of boundedness.

Proposition 6.4. *Let $(T(t))_{t \geq 0}$ be a C_0 -semigroup on a Banach space X , with generator $-A$. Assume that*

$$\lim_{t \rightarrow \infty} \|T(t)A(\omega + A)^{-1}\| = 0$$

for some $\omega \in \rho(-A)$. Then, for each $\eta > 0$,

$$\inf \{ \text{Re } \lambda : \lambda \in \sigma(A), |\lambda| > \eta \} > 0.$$

In particular,

$$\sigma(A) \cap \{ \lambda \in \mathbb{C} : \text{Re } \lambda \leq 0 \} \subset \{0\}.$$

Proof. We may assume that $\omega > 0$ and $\|T(t)\| \leq Me^{(\omega-1)t}$ for all $t > 0$. Define $f_t \in L^1(\mathbb{R}_+) \subset M^b(\mathbb{R}_+)$ and $\mu_t \in M^b(\mathbb{R}_+)$ by

$$\begin{aligned} f_t(s) &= \begin{cases} \omega e^{-\omega(s-t)}, & s \geq t, \\ 0 & 0 \leq s < t, \end{cases} \\ \mu_t &= \delta_t - f_t. \end{aligned}$$

Here δ_t is the unit mass at t and $f_t \in L^1(\mathbb{R}_+)$ is regarded as an absolutely continuous measure. Then

$$T(t)A(\omega + A)^{-1} = \int_0^\infty T(s) d\mu_t(s).$$

By the spectral mapping (inclusion) theorem for the Hille-Phillips functional calculus [39, Theorem 16.3.5],

$$\{e^{-\lambda t} \lambda(\omega + \lambda)^{-1} : \lambda \in \sigma(A)\} \subset \sigma(T(t)A(\omega + A)^{-1}), \quad (6.5)$$

and then

$$\|T(t)A(\omega + A)^{-1}\| \geq \sup \{e^{-\operatorname{Re} \lambda t} |\lambda| |\omega + \lambda|^{-1} : \lambda \in \sigma(A)\}.$$

If $\lambda \in \sigma(A)$ and $|\lambda| \geq \eta > 0$, then $|\lambda| |\omega + \lambda|^{-1} \geq \eta(\omega + \eta)^{-1}$. Taking t such that $\|T(t)A(\omega + A)^{-1}\| \leq \eta/(e(\omega + \eta))$, it follows that $e^{-\operatorname{Re} \lambda t} \leq 1/e$ and $\operatorname{Re} \lambda \geq 1/t$ for all such λ . \square

We shall see in Theorem 6.10 that $\|(is + A)^{-1}\|$ is bounded for $|s| \geq \eta > 0$ if the assumptions of Proposition 6.4 are satisfied and $(T(t))_{t \geq 0}$ is bounded.

The following simple example shows that (6.4) does not imply that the semigroup is bounded.

Example 6.5. Let $X = L^2([0, 1])$. Define a C_0 -semigroup $(T(t))_{t \geq 0}$ on X by

$$(T(t)f)(s) = e^{-(s^{3/2} + is)t} f(s), \quad t \geq 0, \quad f \in L^2([0, 1]), \quad s \in [0, 1].$$

The semigroup $(T(t))_{t \geq 0}$ has a bounded generator $-A$ given by $(-Af)(s) = -(s^{3/2} + is)f(s)$ for $s \in [0, 1]$, and it is easy to check that

$$\|AT(t)\| = O(t^{-2/3}), \quad t \rightarrow \infty.$$

Consider then the semigroup $(\mathcal{T}(t))_{t \geq 0}$, with generator $-\mathcal{A}$, on the space $X \oplus X$ given by the operator matrix

$$\mathcal{T}(t) = \begin{pmatrix} T(t) & -tAT(t) \\ 0 & T(t) \end{pmatrix}, \quad t \geq 0.$$

Then a simple calculation shows that there exist $c, C > 0$ such that

$$\|\mathcal{T}(t)\| \geq ct^{1/3} \quad \text{and} \quad \|\mathcal{A}\mathcal{T}(t)\| \leq Ct^{-1/3}, \quad t > 0.$$

For semigroups of normal operators on Hilbert space, (6.4) does imply boundedness. We omit the easy proof of the following proposition.

Proposition 6.6. Let $(T(t))_{t \geq 0}$ be a C_0 -semigroup of normal operators on a Hilbert space X , with generator $-A$. The following are equivalent:

- (i) For some/all $\omega \in \rho(-A)$, $\lim_{t \rightarrow \infty} \|T(t)A(\omega + A)^{-1}\| = 0$.
- (ii) $\sigma(A) \subset \mathbb{C}_+ \cup \{0\}$ and $\inf\{\operatorname{Re} \lambda : \lambda \in \sigma(A), |\lambda| > \eta\} > 0$ for some/all $\eta > 0$.
- (iii) $(T(t))_{t \geq 0}$ is bounded, $\sigma(A) \cap i\mathbb{R} \subset \{0\}$ and $\sup_{|s| \geq \eta} \|(is + A)^{-1}\| < \infty$ for some/all $\eta > 0$.

Proposition 6.6 shows that there are bounded semigroups of normal operators on Hilbert spaces, with $\sigma(A) \cap i\mathbb{R} \subset \{0\}$, for which property (i) of Proposition 6.6 does not hold. The next example shows that there are bounded semigroups on Hilbert space which satisfy property (ii), but not property (i).

Example 6.7. Let X_0 and X_1 be Hilbert spaces and consider the Hilbert space $X = X_0 \oplus X_1$. Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on X given by $T(t) = T_0(t) \oplus I$, where $(T_0(t))_{t \geq 0}$ is a bounded, but not exponentially stable, C_0 -semigroup on X_0 with generator $-A_0$ such that $\sigma(A_0) \subset \mathbb{C}_+$. If $-A$ is the generator of $(T(t))_{t \geq 0}$ then $\sigma(A) \cap i\mathbb{R} = \{0\}$. Moreover, (6.4) does not hold; if it did, since A_0 is invertible it would follow that $\|T_0(t)\| \rightarrow 0$, a contradiction.

We may take $(T_0(t))_{t \geq 0}$ to be a bounded semigroup on a Hilbert space, which is not exponentially stable, with $\inf\{\operatorname{Re} \lambda : \lambda \in \sigma(A)\} > 0$ (see [6, Example 5.1.10]). Then $(T(t))_{t \geq 0}$ satisfies (ii) of Proposition 6.6, but (i) does not hold.

6.2 Rates of decay

For the rest of Section 6 and throughout Section 7, we shall consider bounded semigroups. For simplicity of notation, we shall set $\omega = 1$ in (6.4).

Our main objective is to deduce (6.4) as a consequence of suitable spectral assumptions. We shall need to assume strong conditions which are consistent with Proposition 6.6 and which also exclude examples such as Example 6.7. In fact, we shall show that the property (iii) of Proposition 6.6 implies (6.4) for all semigroups on Hilbert space, and not only for normal semigroups. This will be deduced from Theorem 6.14 which is a version of the Katznelson-Tzafriri Theorem for semigroups on Hilbert spaces for certain measures which are not absolutely continuous. This extension is of independent interest, and it will be the subject of Subsection 6.3. In this subsection we shall consider other aspects of the rates of decay in (6.4).

Remark 6.8. For bounded semigroups on Banach spaces the property (6.2) implies that $\lim_{t \rightarrow \infty} T(t)A(I+A)^{-1} = 0$ in the strong operator topology. This was explicitly shown in [10, Example, p.802] using a Tauberian theorem for vector-valued Laplace-Stieltjes transforms. It follows from Corollary 6.2 using the uniform boundedness of $T(t)A(I+A)^{-1}$ and the density of the range of $(I+A)^{-1}$.

We consider a bounded C_0 -semigroup $(T(t))_{t \geq 0}$, with generator $-A$, on a Banach space or Hilbert space X . We assume that $\sigma(A) \cap i\mathbb{R} = \{0\}$ and that, for some/all $\eta > 0$, $\sup_{|s| \geq \eta} \|(is + A)^{-1}\| < \infty$. We aim to exhibit the possible rates of decay of $\|T(t)A(I+A)^{-1}\|$ in terms of the growth of $\|(is + A)^{-1}\|$ as $|s| \rightarrow 0$. For this purpose, let m and N be decreasing functions on $(0, \infty)$ such that

$$\|(is + A)^{-1}\| \leq m(|s|), \quad s \neq 0, \tag{6.6}$$

$$\|T(t)A(I+A)^{-1}\| \leq N(t), \quad t > 0. \tag{6.7}$$

The smallest possible functions are given by

$$m(s) = \sup\{\|(ir + A)^{-1}\| : |r| \geq s\}, \quad s > 0, \tag{6.8}$$

$$N(t) = \sup\{\|T(\tau)A(I+A)^{-1}\| : \tau \geq t\}, \quad t > 0. \tag{6.9}$$

This function m is continuous, and we shall always assume continuity of m so that m has a right inverse m^{-1} defined on an interval of the form $[a, \infty)$.

The function N defined by (6.9) may not be continuous, but it is lower semicontinuous and right-continuous. Assuming that $\lim_{t \rightarrow \infty} N(t) = 0$, we define

$$N^*(s) = \min\{t \geq 0 : N(t) \leq s\}, \quad s > 0, \quad (6.10)$$

so that $N(N^*(s)) \leq s$ for all $s > 0$, and $N(N^*(s)) = s$ if s is in the range of N .

Since we assume that $0 \in \sigma(A)$, elementary theory of resolvents implies that $m(s) \geq s^{-1}$ for all $s > 0$. The hypothesis that there might be a corresponding lower bound for N is too naive, because examples of the type considered in Example 6.7 show that N can decay arbitrarily slowly even for semigroups of normal operators on Hilbert space. On the other hand the next result shows that examples where N decays faster than t^{-1} must be of the form considered in Example 6.7.

Theorem 6.9. *Let $(T(t))_{t \geq 0}$ be a C_0 -semigroup on a Banach space X with generator $-A$. If $0 \in \sigma(A)$, then at least one of the following two properties holds:*

- (i) $\limsup_{t \rightarrow \infty} t \|T(t)A(I + A)^{-1}\| > 0$;
- (ii) *There are closed T -invariant subspaces X_0, X_1 of X such that*
 - (a) $X = X_0 \oplus X_1$,
 - (b) $T(t)x = x$ for all $t \geq 0, x \in X_1$, and
 - (c) *the generator $-A_0$ of the restriction of T to X_0 is invertible.*

Proof. Assume first that 0 is a limit point of $\sigma(A)$. Then there exists $\{\lambda_n : n \geq 1\} \subset \sigma(A) \setminus \{0\}$ such that $\lambda_n \rightarrow 0$ as $n \rightarrow \infty$. Setting $t_n = (\operatorname{Re} \lambda_n)^{-1}$, and using (6.5), we infer that

$$\limsup_{t \rightarrow \infty} t \|T(t)A(I + A)^{-1}\| \geq \lim_{n \rightarrow \infty} \left(\frac{t_n e^{-\operatorname{Re} \lambda_n t_n} \operatorname{Re} \lambda_n}{|1 + \lambda_n|} \right) = \frac{1}{e}.$$

So (i) holds.

Now assume that 0 is not a limit point of $\sigma(A)$. Then X can be decomposed into the direct sum of T -invariant subspaces $X = X_0 \oplus X_1$ such that $A_1 := A|_{X_1} \in \mathcal{L}(X_1)$, $\sigma(A_1) = \{0\}$, and for $A_0 := A|_{X_0}$ one has $\sigma(A_0) = \sigma(A) \setminus \{0\}$. If (i) is false, then

$$\begin{aligned} \liminf_{t \rightarrow \infty} t \|A_1 e^{-tA_1}\| &\leq \limsup_{t \rightarrow \infty} t \|A_1 e^{-tA_1} (I + A_1)^{-1}\| \|I + A_1\| \\ &\leq \limsup_{t \rightarrow \infty} t \|T(t)A(I + A)^{-1}\| \|I + A_1\| = 0. \end{aligned}$$

By [44, Theorem 2.1], $A_1 = 0$. Thus (ii) holds. \square

Thus the rates are of interest only in the case when $N(t)$ decreases no faster than t^{-1} as $t \rightarrow \infty$, and $m(s)$ increases at least as fast as s^{-1} as $s \rightarrow 0+$. In that case the bound in (6.12) below gives $m(s) = O(N^*(cs))$ for small $s > 0$.

Theorem 6.10. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Banach space X , with generator $-A$. Assume that*

$$\lim_{t \rightarrow \infty} \|T(t)A(I + A)^{-1}\| = 0. \quad (6.11)$$

Let N be a decreasing function such that (6.7) holds and $\lim_{t \rightarrow \infty} N(t) = 0$, and let N^ denote any function such that $N(N^*(s)) \leq s$ for all $s \in (0, 1)$. Then $\sigma(A) \cap i\mathbb{R} \subset \{0\}$ and, for any $c \in (0, 1)$,*

$$\|(is + A)^{-1}\| = \begin{cases} O(N^*(c|s|) + |s|^{-1}), & s \rightarrow 0, \\ O(1), & |s| \rightarrow \infty. \end{cases} \quad (6.12)$$

Proof. Proposition 6.4 shows that $\sigma(A) \cap i\mathbb{R} \subset \{0\}$. Alternatively, the arguments which follow show that A has no approximate eigenvalues in $i\mathbb{R} \setminus \{0\}$. Since $-A$ generates a bounded semigroup, $\sigma(A) \subset \overline{\mathbb{C}}_+$ and $\sigma(A) \cap i\mathbb{R}$ consists only of approximate eigenvalues.

Let $K = \sup_{t \geq 0} \|T(t)\|$. Let $s \in \mathbb{R} \setminus \{0\}$, $t > 0$ and $x \in \text{dom}(A)$. We use the formula

$$\begin{aligned} ise^{ist}x &= ise^{ist} \int_0^t e^{-is\tau} T(\tau)(is + A)x \, d\tau + isT(t)x \\ &= ise^{ist} \int_0^t e^{-is\tau} T(\tau)(is + A)x \, d\tau + T(t)(I + A)^{-1}(is + A)x \\ &\quad - (1 - is)T(t)A(I + A)^{-1}x. \end{aligned}$$

Since $\|T(t)(I + A)^{-1}\| \leq K$, this gives

$$|s| \|x\| \leq K(|s|t + 1)\|(is + A)x\| + |1 - is|N(t)\|x\|.$$

Hence

$$(|s| - |1 - is|N(t))\|x\| \leq K(|s|t + 1)\|(is + A)x\|. \quad (6.13)$$

Set $t = N^*(c|s|)$. For $|s|$ sufficiently small,

$$|s| - |1 - is|N(t) \geq |s|(1 - |1 - is|c) > 0.$$

For any $K' > K$, (6.13) gives

$$\|(is + A)^{-1}\| \leq \frac{K(|s|N^*(c|s|) + 1)}{|s|(1 - |1 - is|c)} \leq \frac{K'}{1 - c} (N^*(c|s|) + |s|^{-1}), \quad (6.14)$$

for $|s|$ sufficiently small.

Since $\lim_{t \rightarrow \infty} N(t) = 0$, we may set $t = \tau$ with $N(\tau) < 1$. For $|s|$ sufficiently large,

$$|s| - |1 - is|N(\tau) > 1,$$

so (6.13) gives

$$\|(is + A)^{-1}\| \leq \frac{K(|s|\tau + 1)}{|s| - |1 - is|N(\tau)} = O(1), \quad |s| \rightarrow \infty. \quad \square$$

Theorem 6.10 is analogous to [13, Proposition 1.3]. It allows one to get lower bounds for the decay of $\|T(t)A(I+A)^{-1}\|$ in terms of the growth of the resolvent near the origin, analogous to (5.6).

Corollary 6.11. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Banach space X , with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R} = \{0\}$ and that*

$$\lim_{s \rightarrow 0} \max (\|s(is + A)^{-1}\|, \|s(-is + A)^{-1}\|) = \infty. \quad (6.15)$$

Let m be the function defined by (6.8) and m^{-1} be any right inverse for m . Then there exist $c > 0$ and $c' > 0$ such that

$$\|T(t)A(I+A)^{-1}\| \geq cm^{-1}(c't)$$

for all sufficiently large t .

Proof. We may assume that (6.11) holds. Let N and N^* be defined by (6.9) and (6.10) respectively. By Theorem 6.10,

$$m(s) \leq C \left(N^*(cs) + \frac{1}{s} \right), \quad 0 < s \leq 1.$$

Rearranging this and using the assumption (6.15),

$$N^*(cs) > \frac{m(s)}{2C}$$

for all sufficiently small $s > 0$. For t sufficiently large, put $s = c^{-1}N(t)$. Then $N^*(cs) \leq t$. Hence

$$m(m^{-1}(2Ct)) = 2Ct \geq 2CN^*(cs) > m(s).$$

Since m is decreasing,

$$m^{-1}(2Ct) < s = c^{-1}N(t) \leq c^{-1}K\|T(t)A(I+A)^{-1}\|,$$

for all sufficiently large t . □

Remark 6.12. The assumption (6.15) in Corollary 6.11 cannot be omitted, because of the trivial semigroup where $A = 0$, for example. For a C_0 -semigroup of contractions, it can be weakened to the assumption that

$$\liminf_{s \rightarrow 0} \max (\|s(is + A)^{-1}\|, \|s(-is + A)^{-1}\|) > 1.$$

This follows from (6.14).

The ideal counterpart to Corollary 6.11 would be to show that if (6.6) holds, then

$$\|T(t)A(I+A)^{-1}\| = O(m^{-1}(ct)), \quad t \rightarrow \infty, \quad (6.16)$$

for some $c > 0$. However for some m it is not possible to get this sharper estimate, even for semigroups of normal operators on Hilbert space, as the following proposition shows. The

proposition gives the necessary (when m is defined by (6.8)) and sufficient condition on m for (6.16) to be true for semigroups of normal operators on Hilbert spaces. It is presented in the more general context of quasi-multiplication semigroups introduced in Section 5 and it should be compared to Proposition 5.1. The conditions (5.10) for M and (6.18) for m are clearly dual to each other: M satisfies (5.10) if and only if $m(s) := M(1/s)$ satisfies (6.18). Later we shall consider more general semigroups on Hilbert space, and we shall establish (6.16) when $m(s) = s^{-\alpha}$ for some $\alpha \geq 1$, and weaker estimates for more general m , see Theorems 7.6, 7.7 and 6.15.

Proposition 6.13. *Let $(T(t))_{t \geq 0}$ be a quasi-multiplication semigroup on a Banach space X with generator $-A$. Assume that*

$$0 \in \sigma(A) \subset \mathbb{C}_+ \cup \{0\} \quad \text{and} \quad \|(is + A)^{-1}\| = O(1), \quad |s| \rightarrow \infty.$$

Let $c > 0$, m be defined by (6.8), and m^{-1} be any right inverse for m . Then the following are equivalent:

(i) *There exists C such that*

$$\|T(t)A(I + A)^{-1}\| \leq Cm^{-1}(ct), \quad t \geq c^{-1}m(1); \quad (6.17)$$

(ii) *There exists B such that*

$$\frac{m(\tau)}{m(s)} \geq c \log \left(\frac{s}{\tau} \right) - B, \quad 0 \leq \tau, s \leq 1. \quad (6.18)$$

Proof. Note first that the assumptions imply that $\|T(t)\| = 1$ for all $t \geq 0$,

$$m(s)^{-1} = \min \{ |\mu + ir| : \mu \in \sigma(A), |r| \geq s \} \leq s, \quad s > 0,$$

and there exists $\varepsilon > 0$ such that $\operatorname{Re} \mu \geq \varepsilon$ for all $\mu \in \sigma(A)$ with $|\mu| \geq 1$. Moreover (6.17) is equivalent to

$$e^{-t\alpha} \left| \frac{\mu}{1 + \mu} \right| \leq Cm^{-1}(ct), \quad \mu = \alpha + i\beta \in \sigma(A) \setminus \{0\}, \quad t \geq c^{-1}m(1),$$

and hence to

$$t\alpha \geq \log \left(\frac{1}{Cm^{-1}(ct)} \left| \frac{\mu}{1 + \mu} \right| \right) \quad (6.19)$$

for all such μ and t .

Assume that (i) holds. Let $t \geq c^{-1}m(1)$, and put $\tau = m^{-1}(ct)$. From (6.19),

$$m(\tau) \geq \frac{c}{\alpha} \log \left(\frac{|\mu|}{C\tau|1 + \mu|} \right) \geq \frac{c}{\alpha} \log \left(\frac{|\mu|}{2C\tau} \right),$$

if $\mu = \alpha + i\beta \in \sigma(A)$ and $|\mu| \leq 1$.

Given $s > 0$ with $s < \varepsilon$, take $\mu = \alpha + i\beta \in \sigma(A)$ such that $m(s)^{-1} = |\mu + ir|$ for some $|r| \geq s$. Then $\alpha \leq |\mu + ir| \leq s < \varepsilon$, so $|\mu| \leq 1$. If $|\mu| \geq s/2$, then

$$\frac{m(\tau)}{m(s)} \geq \frac{c|\alpha + i(\beta + r)|}{\alpha} \log\left(\frac{s}{4C\tau}\right) \geq c \log\left(\frac{s}{\tau}\right) - c \log(4C).$$

If $|\mu| < s/2$, then $m(s)^{-1} \geq |r| - |\mu| \geq s/2$. Since $m(\tau) \geq 1/\tau$,

$$\frac{m(\tau)}{m(s)} \geq \frac{s}{2\tau} \geq c \log\left(\frac{s}{\tau}\right) - B,$$

for some $B \geq c \log(4C)$.

It follows that (6.18) holds whenever $0 \leq s < \varepsilon$ and τ is in the range of m^{-1} . For other values of τ one can apply the above with τ replaced by $m^{-1}(m(\tau) - n^{-1})$, and let $n \rightarrow \infty$. If $\varepsilon \leq s \leq 1$, one can use

$$\frac{m(\tau)}{m(s)} \geq \frac{m(\tau)}{m(\varepsilon)} \geq c \log\left(\frac{\varepsilon}{\tau}\right) - B \geq c \log\left(\frac{s}{\tau}\right) - B + c \log \varepsilon.$$

Now assume that (ii) holds. Given $t \geq c^{-1}m(1)$ and $\mu = \alpha + i\beta \in \sigma(A)$ with $|\beta| \leq 1$, take

$$\tau = m^{-1}(ct), \quad s = |\beta|.$$

By (6.18),

$$\frac{ct}{m(|\beta|)} \geq c \log\left(\frac{|\beta|}{m^{-1}(ct)}\right) - B.$$

Rearranging this, using $\alpha m(|\beta|) \geq 1$, and putting $C = 2 \exp(B/c)$ gives (6.19), provided that $|\mu|/2 \leq |\beta| \leq 1$. If $|\beta| < |\mu|/2$, then $\alpha > (\sqrt{3}/2)|\mu|$, and

$$t\alpha = \frac{\alpha}{c} m(m^{-1}(ct)) > \frac{\sqrt{3}|\mu|}{2cm^{-1}(ct)} \geq \log\left(\frac{\sqrt{3}|\mu|}{2cm^{-1}(ct)}\right).$$

So we establish (6.19) in this case also with $C = 2c/\sqrt{3}$. Taking the maximum of two values of C we have established (6.19) whenever $|\beta| \leq 1$.

Now consider $\mu = \alpha + i\beta \in \sigma(A)$ with $|\mu| \geq 1$, so $\alpha \geq \varepsilon$. Proceeding in a similar way to the last part of the proof of Proposition 5.1 (or applying the same arguments to the function $M(s) := m(1/s)$), one sees that there exists C such that

$$\varepsilon t \geq \log\left(\frac{1}{Cm^{-1}(ct)}\right) \geq \log\left(\frac{1}{Cm^{-1}(ct)} \left| \frac{\mu}{1 + \mu} \right| \right)$$

whenever $t \geq c^{-1}m(1)$. Then (6.19) holds for $\mu = \alpha + i\beta \in \sigma(A)$ with $|\beta| \geq 1$ and $t \geq c^{-1}m(1)$. Hence (i) holds. \square

6.3 An extension of the Katznelson-Tzafriri Theorem

The following result is an extension of the Katznelson-Tzafriri Theorem 6.1 in the case of semigroups on Hilbert spaces to some measures which are not absolutely continuous.

Theorem 6.14. *Let $-A$ be the generator of a bounded C_0 -semigroup $(T(t))_{t \geq 0}$ on a Hilbert space X . Assume that $E := i\sigma(-A) \cap \mathbb{R}$ is compact and of spectral synthesis. Assume in addition that, for some $\eta \geq 0$,*

$$\sup_{|s| \geq \eta} \|(is + A)^{-1}\| < +\infty. \quad (6.20)$$

Let $\mu \in M^b(\mathbb{R}_+)$ be such that $\mathcal{F}\mu$ vanishes on E . Then

$$\lim_{t \rightarrow \infty} \|T(t)\hat{\mu}(T)\| = 0,$$

where

$$\hat{\mu}(T)x = \int_0^\infty T(s)x \, d\mu(s), \quad x \in X.$$

Proof. Let $\varphi \in \mathcal{S}(\mathbb{R})$ be such that $\mathcal{F}\varphi$ has compact support and such that $\mathcal{F}\varphi = 1$ on a neighbourhood of E . We decompose the measure μ as follows

$$\mu = \mu * \varphi + \mu * (\delta_0 - \varphi) =: \mu_0 + \mu_1,$$

and we note that the measure μ_0 is absolutely continuous with respect to Lebesgue measure: let $f \in L^1(\mathbb{R})$ be the density function for μ_0 .

Consider the function

$$F(t) := \int_{\mathbb{R}} T(t+s) \, d\mu(s), \quad t \in \mathbb{R},$$

where we have extended the semigroup T by 0 on $(-\infty, 0)$ and the integral is convergent in the strong operator topology. Since μ is supported in \mathbb{R}_+ , we have

$$F(t) = T(t)\hat{\mu}(T), \quad t \geq 0.$$

Following the decomposition of μ , we define also the functions

$$F_0(t) := \int_{\mathbb{R}} T(t+s)f(s) \, ds, \quad t \in \mathbb{R},$$

and

$$F_1(t) := \int_{\mathbb{R}} T(t+s) \, d\mu_1(s), \quad t \in \mathbb{R},$$

so that $F = F_0 + F_1$. In these two definitions, we have also extended the semigroup T by 0 on $(-\infty, 0)$.

Note that

$$\mathcal{F}\mu_0(\xi) = \mathcal{F}f(\xi) = \mathcal{F}\mu(\xi) \mathcal{F}\varphi(\xi), \quad \xi \in \mathbb{R}.$$

Since $\mathcal{F}\mu$ vanishes on E , the Fourier transform $\mathcal{F}f$ also vanishes on E . Since E is of spectral synthesis, there exists a sequence $(f_n)_{n \geq 2} \subset L^1(\mathbb{R})$ such that each Fourier transform $\mathcal{F}f_n$ vanishes on a neighbourhood of E and such that $\lim_{n \rightarrow \infty} \|f_n - f\|_{L^1} = 0$. Furthermore, we may assume that each Fourier transform $\mathcal{F}f_n$ has compact support. Set

$$F_n(t) := \int_{\mathbb{R}} T(t+s)f_n(s) ds, \quad n \geq 2, t \in \mathbb{R}.$$

We have, by Parseval's formula,

$$\begin{aligned} F_n(t) &= \lim_{\alpha \rightarrow 0^+} \int_0^\infty e^{-\alpha s} T(s) f_n(s-t) ds \\ &= \lim_{\alpha \rightarrow 0^+} \frac{1}{2\pi} \int_{\mathbb{R}} (\alpha + i\xi + A)^{-1} \mathcal{F}f_n(-\xi) e^{i\xi t} d\xi \\ &= \frac{1}{2\pi} \int_{\mathbb{R}} (i\xi + A)^{-1} \mathcal{F}f_n(-\xi) e^{i\xi t} d\xi, \quad t \in \mathbb{R}. \end{aligned}$$

Here the function under the integral is well defined since $\mathcal{F}f_n$ vanishes on a neighbourhood of E . Moreover, the function $\xi \mapsto (i\xi + A)^{-1} \mathcal{F}f_n(-\xi)$ is continuous and has compact support. Hence, by the Riemann-Lebesgue theorem,

$$\lim_{|t| \rightarrow \infty} \|F_n(t)\| = 0, \quad n \geq 2. \quad (6.21)$$

Since

$$\lim_{n \rightarrow \infty} \sup_{t \in \mathbb{R}} \|F_n(t) - F_0(t)\| = 0,$$

we therefore obtain

$$\lim_{|t| \rightarrow \infty} \|F_0(t)\| = 0. \quad (6.22)$$

Let us now examine the function F_1 . Take $x \in X$. The function $t \mapsto e^{-t}T(t)x$ is in $L^2(\mathbb{R}_+; X)$. Extending this function by zero on $(-\infty, 0)$, Plancherel's theorem implies that

$$\int_{\mathbb{R}} \|(1 + i\xi + A)^{-1}x\|^2 d\xi \leq C^2 \|x\|^2. \quad (6.23)$$

For $\alpha \in (0, 1)$, let

$$F_{1,\alpha,x}(t) = \int_{\mathbb{R}} e^{-\alpha(t+s)} T(t+s)x d\mu_1(s), \quad t \in \mathbb{R}.$$

Then

$$\lim_{\alpha \rightarrow 0^+} F_{1,\alpha,x}(t) = F_1(t)x. \quad (6.24)$$

Moreover, $F_{1,\alpha,x} \in L^1(\mathbb{R}; X)$ and its Fourier transform is

$$\begin{aligned} &(\mathcal{F}F_{1,\alpha,x})(\xi) \\ &= \mathcal{F}\mu(\xi)(1 - \mathcal{F}\varphi(-\xi))(\alpha + i\xi + A)^{-1}x \\ &= \mathcal{F}\mu(\xi)(1 - \mathcal{F}\varphi(-\xi)) (I + (1 - \alpha)(\alpha + i\xi + A)^{-1}) (1 + i\xi + A)^{-1}x, \end{aligned}$$

by the resolvent identity. The assumption (6.20) extends by the Neumann series to boundedness of $\|(\alpha + i\xi + A)^{-1}\|$ for small $\alpha > 0$ and $|\xi| \geq R$, and then for all $\alpha > 0$ and $|\xi| \geq R$ since the semigroup is bounded. Since $\mathcal{F}\varphi = 1$ in a neighbourhood of E , there is a constant C (independent of α , ξ and x) such that

$$\|(\mathcal{F}F_{1,\alpha,x})(\xi)\| \leq C\|(1 + i\xi + A)^{-1}x\|, \quad 0 < \alpha < 1, \xi \in \mathbb{R}. \quad (6.25)$$

Moreover,

$$\begin{aligned} & \lim_{\alpha \rightarrow 0^+} (\mathcal{F}F_{1,\alpha,x})(\xi) \\ &= \begin{cases} \mathcal{F}\mu(\xi)(1 - \mathcal{F}\varphi(-\xi))(i\xi + A)^{-1}x & \text{if } \mathcal{F}\varphi(-\xi) \neq 1, \\ 0 & \text{otherwise.} \end{cases} \\ &=: G_{1,x}(\xi). \end{aligned}$$

Using (6.23), (6.25), and the Dominated Convergence Theorem, it follows that $G_{1,x} \in L^2(\mathbb{R}; X)$, $\|G_{1,x}\|_{L^2(\mathbb{R}; X)} \leq C\|x\|$ and

$$\lim_{\alpha \rightarrow 0^+} \|\mathcal{F}F_{1,\alpha,x} - G_{1,x}\|_{L^2(\mathbb{R}; X)} = 0.$$

Using Plancherel's theorem again shows that

$$\lim_{\alpha \rightarrow 0^+} \|F_{1,\alpha,x} - \mathcal{F}^{-1}G_{1,x}\|_{L^2(\mathbb{R}; X)} = 0,$$

where $\mathcal{F}^{-1}G_{1,x}$ is inverse L^2 -Fourier transform of $G_{1,x}$. From this and (6.24), we deduce that

$$F_1(t)x = (\mathcal{F}^{-1}G_{1,x})(t) \quad (6.26)$$

for almost all $t > 0$. By Plancherel's theorem once more,

$$\int_{\mathbb{R}} \|F_1(t)x\|^2 dt \leq C^2 \|x\|^2, \quad x \in X.$$

Let $t \geq 0$ and $x \in X$. We compute:

$$\begin{aligned} & \int_0^t T(t-s)F_1(s)x ds \\ &= \int_0^t T(t-s) \int_{\mathbb{R}} T(s+r)x d\mu_1(r) ds \\ &= \int_0^t \int_{\mathbb{R}} T(t+r)x d\mu_1(r) ds - \int_0^t \int_{-t}^{(-s)^-} T(t+r)x d\mu_1(r) ds \\ &= t F_1(t)x + \int_{-t}^0 rT(t+r)x d\mu_1(r), \end{aligned}$$

so that

$$F_1(t)x = \frac{1}{t} \int_0^t T(t-s)F_1(s)x ds - \int_{-t}^0 \frac{r}{t} T(t+r)x d\mu_1(r). \quad (6.27)$$

We estimate the first term on the right-hand side:

$$\left\| \frac{1}{t} \int_0^t T(t-s)F_1(s)x \, ds \right\| \leq \frac{1}{t} \int_0^t K \|F_1(s)x\| \, ds \leq KC \frac{1}{\sqrt{t}} \|x\|, \quad (6.28)$$

where $K = \sup_{t \geq 0} \|T(t)\|$. For the second term on the right-hand side of (6.27), we note that

$$\lim_{t \rightarrow \infty} \int_{-t}^0 \left\| \frac{\tau}{t} T(t+\tau) \right\| \, d\mu_1(\tau) = 0,$$

by the Bounded Convergence Theorem for the bounded measure μ_1 . Thus we have

$$\lim_{t \rightarrow \infty} \|F_1(t)\| = 0.$$

From this and (6.22), we obtain

$$\lim_{t \rightarrow \infty} \|F(t)\| = \lim_{t \rightarrow \infty} \|T(t)\hat{\mu}(T)\| = 0,$$

which is the claim. \square

Now we return to the situation of Subsection 6.2. We assume that $\sigma(A) \cap i\mathbb{R} = \{0\}$ and that $\sup_{|s| \geq \eta} \|(is + A)^{-1}\| < +\infty$ for some $\eta > 0$. Since one-point sets are of spectral synthesis, we can apply Theorem 6.14 with $\mu \in M^b(\mathbb{R}_+)$ given by

$$\mu = \delta_0 - e_1, \quad e_1(s) = e^{-s}, \quad s \geq 0. \quad (6.29)$$

Then $\mathcal{F}\mu(0) = 0$ and $\hat{\mu}(T) = I - (I + A)^{-1} = A(I + A)^{-1}$, so the conclusion of Theorem 6.14 is that

$$\lim_{t \rightarrow \infty} \|T(t)A(I + A)^{-1}\| = 0.$$

The following result includes an estimate of the rate of decay of $\|T(t)A(I + A)^{-1}\|$ in terms of the growth of $\|(is + A)^{-1}\|$ as $s \rightarrow 0$.

Theorem 6.15. *Let $(T(t))_{t \geq 0}$ be a C_0 -semigroup on a Hilbert space X with generator $-A$, and assume that $\sigma(A) \cap i\mathbb{R} = \{0\}$ and that $\sup_{|s| \geq \eta} \|(is + A)^{-1}\| < +\infty$ for some $\eta > 0$. Then*

$$\lim_{t \rightarrow \infty} \|T(t)A(I + A)^{-1}\| = 0. \quad (6.30)$$

More precisely, let $m : (0, 1) \rightarrow (1, \infty)$ be a continuous increasing function such that

$$\|(is + A)^{-1}\| = O(m(|s|)) \quad s \rightarrow 0.$$

Let $\varepsilon \in (0, 1)$. Then

$$\|T(t)A(I + A)^{-1}\| = O(m^{-1}(t^{1-\varepsilon})), \quad t \rightarrow \infty. \quad (6.31)$$

Proof. We have already shown how (6.30) follows from Theorem 6.14. To establish (6.31), we follow the proof of that theorem with μ given by (6.29) and $E = \{0\}$. We take $\varphi \in \mathcal{S}(\mathbb{R})$ so that $\mathcal{F}\varphi$ has compact support and $\mathcal{F}\varphi = 1$ on $[-1, 1]$. Let

$$\mu_0 = \mu * \varphi, \quad \mu_1 = \mu - \mu_0.$$

Then μ_0 is absolutely continuous with density function

$$f(s) = \varphi(s) - \int_0^\infty \varphi(s - \tau) e^{-\tau} d\tau,$$

and

$$\mathcal{F}f(\xi) = \frac{i\xi}{1 + i\xi} \mathcal{F}\varphi(\xi).$$

Take a C^∞ -function ψ such that $0 \leq \psi \leq 1$, $\psi = 1$ on $[-1, 1]$ and the support of ψ is contained in $[-2, 2]$. For $0 < r \leq 1/2$, let g_r be the Schwartz function such that

$$\mathcal{F}g_r(\xi) = \mathcal{F}f(\xi)\psi_r(\xi) = \left(\frac{i\xi}{1 + i\xi} \right) \psi\left(\frac{\xi}{r}\right).$$

The last equality holds because $\psi(\xi/r) = 0$ if $|\xi| \geq 1$ and $(\mathcal{F}\varphi)(\xi) = 1$ if $|\xi| \leq 1$. Then $\mathcal{F}g_r(\xi) = 0$ if $|\xi| \geq 2r$, and, for $|\xi| \leq 2r$ and $j \geq 1$,

$$(\mathcal{F}g_r)^{(j)}(\xi) = \sum_{n=0}^{j-1} \frac{(-1)^{j-n+1} i^{j-n} j!}{(1 + i\xi)^{j-n+1} n!} \frac{1}{r^n} \psi^{(n)}\left(\frac{\xi}{r}\right) + \left(\frac{i\xi}{1 + i\xi} \right) \frac{1}{r^j} \psi^{(j)}\left(\frac{\xi}{r}\right).$$

Hence

$$\begin{aligned} |\mathcal{F}g_r(\xi)| &\leq |\xi|, & \|\mathcal{F}g_r\|_{L^1} &\leq 4r^2, \\ |(\mathcal{F}g_r)^{(j)}(\xi)| &\leq \frac{C_j}{r^{j-1}} \left(1 + \frac{|\xi|}{r}\right), & \|(\mathcal{F}g_r)^{(j)}\|_{L^1} &\leq \frac{8C_j}{r^{j-2}}. \end{aligned}$$

Now, for $s \in \mathbb{R}$,

$$\begin{aligned} |g_r(s)| &= \left| \frac{1}{2\pi} \int_{-2r}^{2r} \mathcal{F}g_r(\xi) e^{i\xi s} d\xi \right| \leq \frac{1}{2\pi} \int_{-2r}^{2r} |\xi| d\xi = \frac{2r^2}{\pi}, \\ |s^2 g_r(s)| &= \left| \frac{1}{2\pi} \int_{-2r}^{2r} (\mathcal{F}g_r)''(\xi) e^{i\xi s} d\xi \right| \leq C, \end{aligned}$$

so

$$\|g_r\|_{L^1} \leq \int_{|s| \leq 1/r} \frac{2r^2}{\pi} ds + \int_{|s| \geq 1/r} \frac{C}{s^2} ds \leq Cr.$$

Let $f_r = f - g_r$. Then $f_r \in L^1(\mathbb{R})$, $\mathcal{F}f_r = 0$ on $[-r, r]$, and

$$\|f - f_r\|_{L^1} \leq Cr, \quad \|\mathcal{F}f_r\|_{L^1} \leq C, \quad \|(\mathcal{F}f_r)^{(j)}\|_{L^1} \leq \begin{cases} C_j & j = 0, 1, \\ \frac{C_j}{r^{j-2}}, & j \geq 2. \end{cases}$$

These functions f_r (as $r \rightarrow 0+$) replace the functions f_n (as $n \rightarrow \infty$) in the proof of Theorem 6.14. Accordingly, we define

$$F_r(t) = \int_{\mathbb{R}} T(t+s)f_r(s) ds, \quad 0 < r < 1/2, t > 0.$$

Instead of applying the Riemann-Lebesgue theorem as we did in the proof of Theorem 6.14 to obtain (6.21), we integrate by parts k times, and we obtain that

$$\begin{aligned} F_r(t) &= \frac{i^k}{2\pi t^k} \int_{\mathbb{R}} \left(\frac{d}{d\xi} \right)^k ((i\xi + A)^{-1} \mathcal{F}f_r(-\xi)) e^{i\xi t} d\xi \\ &= \frac{1}{2\pi i^k t^k} \int_{\mathbb{R}} \sum_{j=0}^k \binom{k}{j} i^{k-j} (k-j)! (i\xi + A)^{-(k-j+1)} (\mathcal{F}f_r)^{(j)}(-\xi) e^{i\xi t} d\xi. \end{aligned}$$

Since $m(r) \geq r^{-1} \geq 2$, we have

$$\|F_r(t)\| \leq \frac{C}{t^k} \left(m(r)^{k+1} + m(r)^k + \sum_{j=2}^k \frac{m(r)^{k-j+1}}{r^{j-2}} \right) \leq \frac{Cm(r)^{k+1}}{t^k}.$$

Now

$$\|F_0(t)\| \leq \|F_r(t)\| + K\|f - f_r\|_{L^1} \leq C \left(\frac{m(r)^{k+1}}{t^k} + r \right),$$

where $K = \sup_{t \geq 0} \|T(t)\|$.

Now we want to show that

$$\|F_1(t)\| = O(t^{-1}), \quad t \rightarrow \infty. \quad (6.32)$$

The estimate (6.28) is not adequate for this purpose, and we use a different argument.

Let $x \in X$. First note that the resolvent identity yields

$$\|(i\xi + A)^{-1}x\| \leq (C+1)\|(i\xi + 1 + A)^{-1}x\|, \quad |\xi| \geq 1, \quad (6.33)$$

where $C := \sup_{|\xi| \geq 1} \|(i\xi + A)^{-1}\|$. Similarly,

$$\|(-i\xi + A^*)^{-1}x\| \leq (C+1)\|(-i\xi + 1 + A^*)^{-1}x\|, \quad |\xi| \geq 1. \quad (6.34)$$

Set

$$g(\xi) := \mathcal{F}\mu(\xi)(1 - \mathcal{F}\varphi(-\xi)) = \frac{i\xi}{1+i\xi}(1 - \mathcal{F}\varphi(-\xi)), \quad \xi \in \mathbb{R},$$

and note that g is zero on $(-1, 1)$ and $g' \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$. Let $x \in X$. It follows from (6.26) that $F_1(\cdot)x$ is the inverse Fourier transform (in the L^2 -sense) of the function $G_{1,x}$ which has derivative

$$G'_{1,x}(\xi) = -g(\xi)(i\xi + A)^{-2}x + g'(\xi)(i\xi + A)^{-1}x. \quad (6.35)$$

Thus $G'_{1,x} \in L^2(\mathbb{R}; X)$, and $G_{1,x}$ belongs to the first-order vector-valued Hilbert-Sobolev space $H^1(\mathbb{R}; X)$. Hence

$$tF_1(t)x = -i\mathcal{F}^{-1}(G'_{1,x})(t).$$

Since $g' \in L^1(\mathbb{R})$, the second term on the right-hand side of (6.35) is in $L^1(\mathbb{R}, X)$, and its inverse Fourier transform is bounded by $(2\pi)^{-1}\|g'\|_1 C\|x\|$. To handle the first term, take $y \in X$. By the Cauchy-Schwarz inequality, (6.33) and (6.34) and Plancherel's theorem,

$$\begin{aligned} & \frac{1}{2\pi} \int_{\mathbb{R}} |\langle g(\xi)(i\xi + A)^{-2}x, y \rangle| d\xi \\ & \leq \frac{\|g\|_{\infty}}{2\pi} \left(\int_{\mathbb{R} \setminus (-1,1)} \|(i\xi + A)^{-1}x\|^2 d\xi \right)^{1/2} \left(\int_{\mathbb{R} \setminus (-1,1)} \|(-i\xi + A^*)^{-1}y\|^2 d\xi \right)^{1/2} \\ & \leq \frac{(C+1)^2\|g\|_{\infty}}{2\pi} \left(\int_{\mathbb{R}} \|(1+i\xi + A)^{-1}x\|^2 d\xi \right)^{1/2} \left(\int_{\mathbb{R}} \|(1-i\xi + A^*)^{-1}y\|^2 d\xi \right)^{1/2} \\ & = (C+1)^2\|g\|_{\infty} \left(\int_0^{\infty} \|e^{-\tau}T(\tau)x\|^2 d\tau \right)^{1/2} \left(\int_0^{\infty} \|e^{-\tau}T^*(\tau)y\|^2 d\tau \right)^{1/2} \\ & \leq (C+1)^2\|g\|_{\infty}K^2\|x\| \|y\|. \end{aligned}$$

It then follows that

$$t|\langle F_1(t)x, y \rangle| \leq (C+1)^2\|g\|_{\infty}K^2\|x\| \|y\| + \frac{\|g'\|_1 C}{2\pi} \|x\| \|y\|$$

for almost all t . Since F_1 is continuous, this estimate holds for all t , so (6.32) follows.

Overall we have that

$$\|T(t)A(1+A)^{-1}\| = \|F_0(t) + F_1(t)\| \leq C \left(\frac{m(r)^{k+1}}{t^k} + \frac{1}{t} + r \right).$$

For a given $\varepsilon \in (0, 1)$, we take $r = m^{-1}(t^{1-\varepsilon})$, for sufficiently large t . Then we obtain

$$\|T(t)A(1+A)^{-1}\| \leq C (t^{1-(k+1)\varepsilon} + t^{-1} + m^{-1}(t^{1-\varepsilon})).$$

We may choose k so that $(k+1)\varepsilon \geq 2$. Since $m(r) \geq r^{-1}$, we have $m^{-1}(t^{1-\varepsilon}) \geq t^{\varepsilon-1} \geq t^{-1}$. Then we obtain (6.31). \square

Example 6.16. (a) Consider the case when $m(s) = s^{-\alpha}$ where $\alpha \geq 1$. Then (6.31) gives

$$\|T(t)A(I+A)^{-1}\| = O(t^{-\gamma}), \quad t \rightarrow \infty,$$

for any $\gamma < 1/\alpha$. We shall see in Theorem 7.6 that this also holds with $\gamma = 1/\alpha$.

(b) Now consider the case when $m(s) = e^{\alpha/s}$ where $\alpha > 0$. Then

$$m^{-1}(t^{1-\varepsilon}) = \frac{\alpha}{(1-\varepsilon)\log t},$$

so

$$\|T(t)A(I+A)^{-1}\| = O((\log t)^{-1}) = O(m^{-1}(t)), \quad t \rightarrow \infty.$$

This rate of decay is sharp in this case (Corollary 6.11).

7 Singularity at zero: polynomial and regularly varying rates

In this section we consider the situation of Subsection 6.2 and Theorem 6.15 in cases when X is a Hilbert space and the function $m(1/s)$ is regularly varying. For simplicity of presentation, we shall write $B(A) = A(I + A)^{-1}$.

Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup with generator $-A$ such that $\sigma(A) \cap i\mathbb{R} = \{0\}$ and

$$\|(is + A)^{-1}\| = \begin{cases} O\left(\frac{1}{|s|^{\alpha}\ell(1/|s|)}\right), & s \rightarrow 0, \\ O(1), & |s| \rightarrow \infty, \end{cases}$$

where ℓ is slowly varying and monotonic. Then the optimal estimate (6.16) on the decay of $\|T(t)B(A)\|$ under the assumption of dB-symmetry of ℓ would be

$$\|T(t)B(A)\| = O\left(\frac{1}{(t\ell(t^{1/\alpha}))^{1/\alpha}}\right), \quad t \rightarrow \infty.$$

We shall show in Theorem 7.7 that this estimate does hold if ℓ is increasing, i.e., $\|(is + A)^{-1}\|$ grows slightly slower than $|s|^{-\alpha}$ as $s \rightarrow 0$. Its proof requires a series of steps similar to those used in the case of a singularity at infinity in Section 5.

7.1 Cancelling resolvent growth

Definition 7.1. Let $\beta \in (0, 1]$, and let ℓ be a slowly varying function such that $g : s \mapsto s^{1-\beta}\ell(s)$ is increasing on \mathbb{R}_+ . Let S_g be the Stieltjes function associated with g (Example 2.14), so

$$S_g(\lambda) = \int_{0+}^{\infty} \frac{d(s^{1-\beta}\ell(s))}{s + \lambda}, \quad \lambda > 0.$$

Let

$$f_g(\lambda) := S_g(1/\lambda), \quad \lambda > 0, \quad (7.1)$$

$$f_m(\lambda) := \frac{f_g(\lambda)}{1 + f_g(\lambda)}, \quad \lambda > 0. \quad (7.2)$$

Since f_g is a complete Bernstein function, $1/f_g$ is Stieltjes by Theorem 2.2, and then $f_m = 1/(1 + 1/f_g)$ is also a complete Bernstein function. Thus the operator $f_m(A)$ is well-defined, either by Definition 3.3 or by the extended functional calculus of Theorem 3.7 (see Remark 3.8(iii)). Moreover $f_m(A) = f_g(A)(I + f_g(A))^{-1}$, by the composition rule in Theorem 3.6(ii). In particular $f_m(A)$ is bounded (see Remark 3.4 and [64, Corollary 12.7]).

Now we make a definition analogous to (5.13).

Definition 7.2. For $\alpha \geq 1$ and $\beta \in (0, 1]$, define

$$V_{\alpha,\beta,\ell}(A) := B(A)^{\alpha-\beta} f_m(A) = B(A)^{\alpha-\beta} f_g(A)(I + f_g(A))^{-1},$$

so that $V_{\alpha,\beta,\ell}(A) \in \mathcal{L}(X)$ by the above.

The next statement shows that this operator cancels resolvent growth in the case of a singularity at zero. It is a counterpart of Theorem 5.5 dealing with singularities at infinity. In the proof we again use Karamata's Theorem 2.15, Proposition 2.4 on domination properties of complete Bernstein functions and complex analysis arguments shown in Lemma 5.3.

Theorem 7.3. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Banach space X , with generator $-A$ such that*

$$0 \in \sigma(A) \subset \mathbb{C}_+ \cup \{0\} \quad \text{and} \quad \|(is + A)^{-1}\| = O(1), \quad |s| \rightarrow \infty. \quad (7.3)$$

Let ℓ be a slowly varying function on \mathbb{R}_+ such that $g : s \mapsto s^{1-\beta}\ell(s)$ is increasing for some $\beta \in (0, 1]$. If there exists $\alpha > 1$ such that

$$\|(is + A)^{-1}\| = O\left(\frac{1}{|s|^{\alpha}\ell(1/|s|)}\right), \quad s \rightarrow 0, \quad (7.4)$$

then

$$\sup_{\lambda \in \mathbb{C}_+} \|(\lambda + A)^{-1}V_{\alpha,\beta,\ell}(A)\| < \infty.$$

Remark 7.4. We are interested only in the case when $0 \in \sigma(A)$, so $\|(is + A)^{-1}\| \geq |s|^{-1}$. Thus the assumption that $\alpha > 1$ restricts generality only slightly. Moreover, if $\alpha = 1$, then the function ℓ can be increasing only if it is bounded. In this case, (7.4) is equivalent to the same estimate with $\ell \equiv 1$. This situation is settled by Corollary 7.5 below. Hence if $\alpha = 1$ we are interested only in the case when ℓ is decreasing and $\lim_{\lambda \rightarrow \infty} \ell(\lambda) = 0$, and then necessarily $\beta \in (0, 1)$. The following proof shows that Theorem 7.3 is also true in that case, provided that $\sum_n \ell(2^n)$ converges.

Proof. By the assumption (7.3) and Lemma 5.3 it suffices to prove that

$$\sup \left\{ \|(is + A)^{-1}V_{\alpha,\beta,\ell}(A)\| : s \in \mathbb{R}, 0 < |s| \leq 1 \right\} < \infty. \quad (7.5)$$

To this aim, observe first that if $k \in \mathbb{N}$, then

$$(\lambda + A)^{-1}B(A)^k = \sum_{i=0}^{k-1} (-\lambda)^{k-1-i} A^i (I + A)^{-k} + (-\lambda)^k (\lambda + A)^{-1} (I + A)^{-k}. \quad (7.6)$$

Let $\alpha - \beta = m + \gamma$ where $m \in \mathbb{N} \cup \{0\}$, and $0 \leq \gamma < 1$. By the assumption (7.4),

$$\|A(is + A)^{-1}\| = \|I - is(is + A)^{-1}\| = O\left(\frac{1}{|s|^{\alpha-1}\ell(1/|s|)}\right), \quad s \rightarrow 0.$$

By the moment inequality (4.2),

$$\|A^\gamma(is + A)^{-1}\| = O\left(\frac{1}{|s|^{\alpha-\gamma}\ell(1/|s|)}\right), \quad s \rightarrow 0.$$

Using (7.6) for $k = m$, it follows that

$$\begin{aligned} \|B(A)^{\alpha-\beta}(is + A)^{-1}\| &= \|A^\gamma(is + A)^{-1}B(A)^m(I + A)^{-\gamma}\| & (7.7) \\ &= O\left(\frac{1}{|s|^{\alpha-\gamma-m}\ell(1/|s|)}\right) \\ &= O\left(\frac{1}{|s|^\beta\ell(1/|s|)}\right), \quad s \rightarrow 0. \end{aligned}$$

Let f_g and f_m be the complete Bernstein functions defined by (7.1) and (7.2). Then

$$\lim_{\lambda \rightarrow 0^+} f_g(\lambda) = \lim_{\lambda \rightarrow 0^+} f_m(\lambda) = 0,$$

and in addition f_m is bounded. Hence f_m has Stieltjes representation of the form $(0, 0, \nu)$. Moreover, by (5.18),

$$|f_g(s)| = O(|s|^\beta\ell(1/|s|)), \quad s \rightarrow 0. \quad (7.8)$$

By Proposition 2.4 and (7.8),

$$|f_m(is)| \leq C f_m(|s|) = \frac{C f_g(|s|)}{1 + f_g(|s|)} = O(|s|^\beta\ell(1/|s|)), \quad s \rightarrow 0, \quad (7.9)$$

Now $f_m(A)$ is a bounded operator, and, by (3.3), for every $x \in \text{dom}(A)$,

$$f_m(A)x = \int_{0^+}^{\infty} A(\lambda + A)^{-1}x \, d\nu(\lambda).$$

We shall show that there exists $C > 0$ such that

$$\|(is + A)^{-1}B(A)^{\alpha-\beta}f_m(A)x\| \leq C\|x\|, \quad x \in \text{dom}(A), \quad 0 < |s| \leq 1. \quad (7.10)$$

Since $\text{dom}(A)$ is dense in X , this will imply that (7.10) holds for all $x \in X$, thus giving us (7.5) and completing the proof.

If $x \in \text{dom}(A)$ and $s \in \mathbb{R} \setminus \{0\}$, then

$$\begin{aligned} &(is + A)^{-1}f_m(A)x \\ &= \int_{0^+}^{\infty} (is + A)^{-1}A(\lambda + A)^{-1}x \, d\nu(\lambda) \\ &= \int_{0^+}^{\infty} \left(\frac{\lambda}{\lambda - is}(\lambda + A)^{-1}x - \frac{is}{\lambda - is}(is + A)^{-1}x \right) d\nu(\lambda) \\ &= f_m(-is)(is + A)^{-1}x + \int_{0^+}^{\infty} \frac{\lambda}{\lambda - is}(\lambda + A)^{-1}x \, d\nu(\lambda). \end{aligned}$$

Thus

$$\begin{aligned} &(is + A)^{-1}f_m(A)B(A)^{\alpha-\beta}x & (7.11) \\ &= f_m(-is)(is + A)^{-1}B(A)^{\alpha-\beta}x + \int_{0^+}^1 \frac{\lambda}{\lambda - is}(\lambda + A)^{-1}B(A)^{\alpha-\beta}x \, d\nu(\lambda) \\ &\quad + \int_{1^+}^{\infty} \frac{\lambda}{\lambda - is}(\lambda + A)^{-1}B(A)^{\alpha-\beta}x \, d\nu(\lambda). \end{aligned}$$

we shall estimate each of the three summands above separately.

To bound the first summand, note that by (7.7) and (7.9), there exists $C > 0$ such that

$$|f_m(-is)| \|(is + A)^{-1} B(A)^{\alpha-\beta} x\| \leq C \|x\|, \quad x \in X, \quad 0 < |s| \leq 1.$$

To estimate the second summand in (7.11), we take into account that $\alpha > \beta$ and consider the following two cases.

If $\alpha - \beta \in (0, 1)$ then (4.1) yields

$$\|B(A)^{\alpha-\beta}(\lambda + A)^{-1}\| = \|(I + A)^{\beta-\alpha} A^{\alpha-\beta}(\lambda + A)^{-1}\| \leq C \lambda^{\alpha-\beta-1},$$

by Proposition 4.1. Then we obtain

$$\left\| \int_{0+}^1 \frac{\lambda}{\lambda - is} B(A)^{\alpha-\beta}(\lambda + A)^{-1} x \, d\nu(\lambda) \right\| \leq C \int_{0+}^1 \lambda^{\alpha-\beta-1} \, d\nu(\lambda) \|x\|.$$

We have to show that the integral here is finite. For this, let I_n be the interval $(2^{-(n+1)}, 2^{-n}]$ for $n \in \mathbb{N} \cup \{0\}$. Then

$$\int_{I_n} d\nu(\lambda) \leq \int_{I_n} \frac{2^{-(n-1)}}{2^{-n} + \lambda} \, d\nu(\lambda) \leq 2f_m(2^{-n}) \leq C 2^{-n\beta} \ell(2^n),$$

by (7.9). Hence

$$\int_{I_n} \lambda^{\alpha-\beta-1} \, d\nu(\lambda) \leq 2^{-n(\alpha-\beta-1)+1} \int_{I_n} d\nu(\lambda) \leq C 2^{-n(\alpha-1)} \ell(2^n).$$

For $\alpha > 1$, it follows from (2.6) with $\gamma = (\alpha - 1)/2$ that

$$\int_{0+}^1 \lambda^{\alpha-\beta-1} \, d\nu(\lambda) = \sum_{n=0}^{\infty} \int_{I_n} \lambda^{\alpha-\beta-1} \, d\nu(\lambda) < \infty.$$

The same conclusion holds if $\alpha = 1$ and $\sum_{n=0}^{\infty} \ell(2^n)$ converges.

If $\alpha - \beta \geq 1$, then

$$\|B(A)^{\alpha-\beta}(\lambda + A)^{-1}\| \leq \|B(A)^{\alpha-\beta-1}\| \|(I + A)^{-1} (I - \lambda(\lambda + A)^{-1})\| \leq C,$$

and

$$\left\| \int_{0+}^1 \frac{\lambda}{\lambda - is} B(A)^{\alpha-\beta}(\lambda + A)^{-1} x \, d\nu(\lambda) \right\| \leq C \int_{0+}^1 d\nu(\lambda) \|x\|.$$

The integral here is finite, by (2.3).

For the third summand in (7.11) we have for any $s \in \mathbb{R}$,

$$\begin{aligned} \left\| \int_{1+}^{\infty} \frac{\lambda}{\lambda - is} B(A)^{\alpha-\beta}(\lambda + A)^{-1} x \, d\nu(\lambda) \right\| &\leq K \int_{1+}^{\infty} \frac{d\nu(\lambda)}{\lambda} \|B(A)^{\alpha-\beta}\| \|x\| \\ &= C \|x\|, \end{aligned}$$

where $K = \sup_{t \geq 0} \|T(t)\|$, and C is finite by (2.3).

Finally, summarizing all the estimates above we get (7.10), and thus the proof of the theorem is complete. \square

7.2 Resolvent growth equal to $s^{-\alpha}$

Here we consider the case when we assume that the resolvent grows (at most) like $s^{-\alpha}$. We note first a corollary from the arguments of the proof of Theorem 7.3 that will be useful in the next subsection and also covers the case $\alpha = 1, \ell \equiv 1$ of Theorem 7.3 excluded in Remark 7.4.

Corollary 7.5. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Banach space X with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R} = \{0\}$ and there exists $\alpha \geq 1$ such that*

$$\|(is + A)^{-1}\| = \begin{cases} O(|s|^{-\alpha}), & s \rightarrow 0, \\ O(1), & |s| \rightarrow \infty. \end{cases}$$

Then

$$\sup_{\lambda \in \mathbb{C}_+} \|(\lambda + A)^{-1} B(A)^\alpha\| < \infty.$$

Proof. The proof follows the same lines as the proof of Theorem 7.3, but it is much simpler. It follows immediately from (7.8) (now for $\beta = 0$ and $\ell = 1$) and Lemma 5.3. \square

The following result, which is analogous to [17, Theorem 2.4], gives the optimal result in the case of exactly polynomial growth of the resolvent for a singularity at zero.

Theorem 7.6. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Hilbert space X with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R} = \{0\}$, and let $\alpha \geq 1$. The following are equivalent:*

$$(i) \quad \|(is + A)^{-1}\| = \begin{cases} O(|s|^{-\alpha}), & s \rightarrow 0, \\ O(1), & |s| \rightarrow \infty. \end{cases}$$

$$(ii) \quad \|T(t)B(A)^\alpha\| = O(t^{-1}), \quad t \rightarrow \infty.$$

$$(iii) \quad \|T(t)B(A)\| = O(t^{-1/\alpha}), \quad t \rightarrow \infty.$$

Proof. By Corollary 7.5, the property (i) implies that

$$\sup_{\lambda \in \mathbb{C}_+} \|(\lambda + A)^{-1} B(A)^\alpha\| < \infty.$$

Then Theorem 4.7 yields (ii).

Properties (ii) and (iii) are equivalent by Lemma 4.2. Property (iii) implies (i) by Theorem 6.10. \square

7.3 Resolvent growth slower than $s^{-\alpha}$

Here we give a counterpart of Theorem 5.6.

Theorem 7.7. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Hilbert space X , with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R} = \{0\}$ and*

$$\|(is + A)^{-1}\| = \begin{cases} O\left(\frac{1}{|s|^{\alpha\ell}(1/|s|)}\right), & s \rightarrow 0, \\ O(1), & |s| \rightarrow \infty, \end{cases} \quad (7.12)$$

where $\alpha > 1$ and ℓ is increasing and slowly varying. Then

$$\|T(t)A(I + A)^{-1}\| = O\left(\frac{1}{(t\ell(t^{1/\alpha}))^{1/\alpha}}\right), \quad t \rightarrow \infty.$$

Proof. We can assume that $T(t)A \neq 0$ for each $t > 0$. Since (7.12) implies property (i) of Theorem 7.6, we obtain from property (iii) of that theorem that

$$\|T(t)B(A)\| = O(t^{-1/\alpha}), \quad t \rightarrow \infty. \quad (7.13)$$

We use the notation of Definitions 7.1 and 7.2, and apply Theorem 7.3, with $\beta = 1$ and $g = \ell$. We conclude that there exists $C > 0$ such that

$$\|(\lambda + A)^{-1}B(A)^{\alpha-1}f_m(A)\| \leq C, \quad \lambda \in \mathbb{C}_+.$$

Then, by Theorem 4.7, we have

$$\|T(t)B(A)^{\alpha-1}f_m(A)\| \leq \frac{C}{t}, \quad t > 0,$$

Hence by Theorem 4.3(b), with $B = B(A)$, $\gamma = \alpha - 1$, and $t_1 = t_2 = t$,

$$f_m(\|T(t)B(A)\|) \leq \frac{C\|T(t)B(A)\|}{t\|T(2t)B(A)^\alpha\|}.$$

Putting $f_{\alpha,m}(s) = s^{\alpha-1}f_m(s)$ we then obtain that

$$f_{\alpha,m}(\|T(t)B(A)\|) \leq \frac{C}{t} \frac{\|T(t)B(A)\|^\alpha}{\|T(2t)B(A)^\alpha\|}. \quad (7.14)$$

By Theorem 2.15 with $g = \ell$ and $\rho = \sigma = 1$, $S_\ell(t) \sim t^{-1}\ell(t)$ as $t \rightarrow \infty$. The function $f_m(s)$ has the same decay at zero as $f_\ell(s)$, hence $f_{\alpha,m}(s) \sim s^\alpha\ell(1/s)$ as $s \rightarrow 0+$. Let $k(s) = 1/\ell(s^{1/\alpha})$. Then

$$f_{\alpha,m}(s) \sim \frac{s^\alpha}{k(s^{-\alpha})}, \quad s \rightarrow 0+.$$

By Proposition 2.11(c),

$$f_{\alpha,m}^{-1}(s) \sim \left(\frac{s}{k^\#(1/s)}\right)^{1/\alpha}, \quad s \rightarrow 0+.$$

If the right-hand side of (7.14) is sufficiently small,

$$\|T(t)B(A)\| \leq \frac{C\|T(t)B(A)\|}{(t\ell(t)\|T(2t)B(A)^\alpha\|)^{1/\alpha}},$$

where

$$L(t) = k^\# \left(t \frac{\|T(2t)B(A)\|}{\|T(t)B(A)\|^\alpha} \right).$$

Let $\psi(s) = (sk^\#(s))^{1/\alpha}$. Since ψ is regularly varying with positive index, we can choose $k^\#$ so that ψ is strictly increasing and continuous (see the remarks following Definition 2.9; in fact, we can choose $k^\#$ to be increasing, since k is decreasing). Then

$$\begin{aligned} \|T(t)B(A)\|^{-1} &\geq ct^{1/\alpha} \frac{\|T(2t)B(A)^\alpha\|^{1/\alpha}}{\|T(t)B(A)\|} L(t)^{1/\alpha} \\ &= c\psi \left(t \frac{\|T(2t)B(A)^\alpha\|}{\|T(t)B(A)\|^\alpha} \right). \end{aligned} \quad (7.15)$$

If the right-hand side of (7.14) is bounded away from 0, then (7.15) also holds for some $c > 0$, since $\|T(t)B(A)\|$ is bounded and ψ is bounded on bounded intervals. So (7.15) holds for all $t > 0$, for some $c > 0$.

Since ψ is strictly increasing,

$$t \frac{\|T(2t)B(A)^\alpha\|}{\|T(t)B(A)\|^\alpha} \leq \psi^{-1}(C\|T(t)B(A)\|^{-1}),$$

for some constant $C > 0$. By Proposition 2.11(b),

$$\psi^{-1}(s) \sim s^\alpha k^{\#\#}(s^\alpha) \sim \frac{s^\alpha}{\ell(s)}, \quad s \rightarrow \infty,$$

so

$$\|T(2t)B(A)^\alpha\| \leq \frac{C}{t\ell(\|T(t)B(A)\|^{-1})}.$$

Then (7.13) yields

$$\|T(2t)B(A)^{-\alpha}\| \leq \frac{C}{t\ell(t^{1/\alpha})},$$

for sufficiently large t . Since $B(A)$ is sectorial, we can apply Lemma 4.2 with $B = B(A)$, $\gamma = \alpha$ and $\delta = 1$ and we obtain

$$\|T(t)B(A)\| \leq \frac{C}{(t\ell(t))^{1/\alpha}},$$

for large t , and the proof is finished. \square

Now we can formulate a counterpart of Corollary 5.7, showing that the optimal estimate (6.16) holds when ℓ is increasing and dB-symmetric.

Corollary 7.8. *In addition to the assumptions of Theorem 7.7, assume that ℓ is dB-symmetric. Then*

$$\|T(t)B(A)\| = O(m^{-1}(t)), \quad t \rightarrow \infty,$$

where m^{-1} is any asymptotic inverse of $s^\alpha/\ell(s)$.

Proof. The proof is completely analogous to the proof of Corollary 5.7. □

The discussion following the proof of Corollary 5.7 and addressing the situation when ℓ may not be dB-symmetric applies to the current setting as well. We omit the details.

Remark 7.9. Note the following duality between singularities at zero and at infinity. If $(T(t))_{t \geq 0}$ is a C_0 -semigroup of contractions on a Hilbert space X with generator $-A$ and the range of A is dense in X , then A is injective and the operator $-A^{-1}$, with dense domain $\text{ran}(A)$, also generates a C_0 -semigroup of contractions on X , by a direct application of the Lumer-Phillips theorem. Thus the case of a singularity at infinity for A corresponds to the case of a singularity at zero for A^{-1} . Conversely, the case of a singularity at zero for A corresponds, in general, to the case of singularities at both zero and at infinity for A^{-1} . This case will be studied in Section 8.

Remark 7.10. We are unable to give a result corresponding to Theorem 5.12 because of the lack of an initial estimate for the rate of decay that is sufficiently close to (6.16) for our technique to work. When $\|(is + A)^{-1}\|$ grows slightly faster than $|s|^{-\alpha}$, our best estimate is

$$\|T(t)B(A)\| = O(t^{-1/\beta}), \quad t \rightarrow \infty,$$

for each $\beta > \alpha$ (see Example 6.16(a) and Theorem 7.6). Our technique does not improve this.

8 Singularities at both zero and infinity

In this section we study the rates of decay in the context of Corollary 6.2 and similar situations, so we are interested in the property

$$\lim_{t \rightarrow \infty} \|T(t)A(I + A)^{-2}\| = 0, \tag{8.1}$$

for a bounded C_0 -semigroup $(T(t))_{t \geq 0}$ on a Banach space or Hilbert space. Note that the range of $A(I + A)^{-2}$ is $\text{ran}(A) \cap \text{dom}(A)$ (Proposition 3.10(i)). Thus we consider orbits starting in $\text{ran}(A) \cap \text{dom}(A)$ or similar spaces.

We shall assume that (8.1) holds, or equivalently that $\sigma(A) \cap i\mathbb{R} \subset \{0\}$ (see Corollary 6.2). We wish to examine the relation between the rate of decay in (8.1) and the rate of growth of $\|(is + A)^{-1}\|$ as $s \rightarrow 0$ or as $|s| \rightarrow \infty$. Let $M : [1, \infty) \rightarrow \mathbb{R}_+$ be a continuous increasing function, and $m : (0, 1] \rightarrow \mathbb{R}_+$ be a continuous decreasing function, such that

$$\|(is + A)^{-1}\| \leq \begin{cases} m(s), & 0 < |s| < 1, \\ M(s), & |s| \geq 1. \end{cases}$$

We assume that $0 \in \sigma(A)$ and that $\lim_{s \rightarrow \infty} M(s) = \infty$ (otherwise we are in the situation of Section 5, or Sections 6 and 7, respectively). By standard theory, $m(s) \geq 1/s$.

For example, M and m can be defined by modified versions of (5.5) and (6.8):

$$M(s) = \sup \{ \|(ir + A)^{-1}\| : 1 \leq |r| \leq s \}, \quad s \geq 1, \quad (8.2)$$

$$m(s) = \sup \{ \|(ir + A)^{-1}\| : s < |r| \leq 1 \}, \quad 0 < s < 1. \quad (8.3)$$

Let N_2 be a continuous, decreasing function such that

$$\|T(t)A(I + A)^{-2}\| \leq N_2(t), \quad t \geq 0.$$

For example, we can take

$$N_2(t) = \sup \{ \|T(\tau)A(I + A)^{-2}\| : \tau \geq t \}, \quad t \geq 0.$$

By Corollary 6.2, our assumptions imply that $\lim_{t \rightarrow \infty} N_2(t) = 0$. Let N_2^{-1} denote any right inverse for N_2 .

Under our assumptions, Martínez [55, Corollary 3.3] has shown that

$$\|T(t)A(I + A)^{-2}\| \leq C \max \left(m_{\log}^{-1}(c't), \frac{1}{M_{\log}^{-1}(C't)} \right), \quad (8.4)$$

where M_{\log} is defined by (5.3) and

$$m_{\log}(s) = m(s) \log \left(\frac{1 + m(s)}{s} \right), \quad 0 < s \leq 1.$$

We have omitted a term t^{-1} which appears in [55], because $m^{-1}(t) \geq t$ when $0 \in \sigma(A)$.

The upper bound (8.4) is analogous to the upper bound for $\|T(t)A^{-1}\|$ given in Theorem 1.1 and there is also a lower bound for $\|T(t)A(I + A)^{-2}\|$. Theorem 6.9 extends to our present situation, with an almost unchanged proof. If $\limsup_{t \rightarrow \infty} t \|T(t)A(I + A)^{-2}\| = 0$, then X splits as in Theorem 6.9(ii).

The following is an analogue of Theorem 6.10, with a similar proof. The term $|s|^{-1}$ in the estimate (8.5) is relevant only when splitting occurs and then $\|(is + A)^{-1}\| = |s|^{-1}$ when $|s|$ is sufficiently small.

Theorem 8.1. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Banach space X , with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R} = \{0\}$, and let N_2 be as above. Then, for any $c \in (0, 1)$,*

$$\|(is + A)^{-1}\| = \begin{cases} O(N_2^{-1}(c|s|) + |s|^{-1}), & s \rightarrow 0, \\ O(N_2^{-1}(c|s|^{-1})) & |s| \rightarrow \infty. \end{cases} \quad (8.5)$$

Proof. Let $K = \sup_{t \geq 0} \|T(t)\|$. Let $s \in \mathbb{R} \setminus \{0\}$, $t > 0$ and $x \in \text{dom}(A)$. We use the formula

$$\begin{aligned} ise^{ist}x &= ise^{ist} \int_0^t e^{-is\tau} T(\tau)(is + A)x \, d\tau + isT(t)x \\ &= ise^{ist} \int_0^t e^{-is\tau} T(\tau)(is + A)x \, d\tau + T(t)(1 + A)^{-1}(is + A)x \\ &\quad - (1 - is)T(t)A(I + A)^{-2}((is + A) + (1 - is))x. \end{aligned}$$

Since $\|T(t)(I + A)^{-1}\| \leq K$, this gives

$$|s| \|x\| \leq K(|s|t + 1)\|(is + A)x\| + |1 - is|N_2(t) (\|(is + A)x\| + |1 - is|\|x\|).$$

Hence

$$(|s| - |1 - is|^2N_2(t)) \|x\| \leq (K(|s|t + 1) + |1 - is|N_2(t))\|(is + A)x\|. \quad (8.6)$$

Set $t = N_2^{-1}(c|s|)$. For $|s|$ sufficiently small,

$$|s| - |1 - is|^2N_2(t) = |s| (1 - c(1 + s^2)) > 0.$$

For any $K' > K$, (8.6) gives

$$\begin{aligned} \|(is + A)^{-1}\| &\leq \frac{K(|s|N_2^{-1}(c|s|) + 1) + |1 - is|c|s|}{|s|(1 - c(1 + s^2))} \\ &\leq \frac{K'}{1 - c} (N_2^{-1}(c|s|) + |s|^{-1}), \end{aligned}$$

for $|s|$ sufficiently small.

Now set $t = N_2^{-1}(c/|s|)$. For $|s|$ sufficiently large,

$$|s| - |1 - is|^2N_2(t) = |s| (1 - c(1 + s^{-2})) > 0.$$

So we obtain

$$\begin{aligned} \|(is + A)^{-1}\| &\leq \frac{K (|s|N_2^{-1}(c|s|^{-1}) + 1) + |1 - is|c|s|^{-1}}{|s|(1 - c(1 + s^{-2}))} \\ &= O(N_2^{-1}(c|s|^{-1})), \quad |s| \rightarrow \infty. \end{aligned}$$

□

As in Corollary 6.11 and (5.6), we obtain the following lower bound.

Corollary 8.2. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Banach space X , with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R} = \{0\}$ and that*

$$\lim_{s \rightarrow 0} \max (\|s(is + A)^{-1}\|, \|s(-is + A)^{-1}\|) = \infty.$$

Define M and m by (8.2) and (8.3) respectively. Then there exist $c, c', C' > 0$ such that

$$\|T(t)A(I + A)^{-2}\| \geq c \max \left(m^{-1}(c't), \frac{1}{M^{-1}(C't)} \right)$$

for all sufficiently large t .

Remark 8.3. (a) For contraction semigroups, the assumption in Corollary 8.2 can be weakened in the same way as in Remark 6.12.

(b) If $(T(t))_{t \geq 0}$ is bounded and eventually differentiable, then $(I + A)^2 T(\tau)$ is a bounded operator for some $\tau > 0$. Moreover,

$$AT(t + \tau) = (I + A)^2 T(\tau) T(t) A (I + A)^{-2},$$

so $\lim_{t \rightarrow \infty} \|AT(t + \tau)\| = 0$ if $\sigma(A) \cap i\mathbb{R} = \{0\}$. By Theorem 6.10, $\|(is + A)^{-1}\|$ is bounded for $|s| \geq 1$. Then (8.4) gives

$$\|AT(t)\| = O(m_{\log}^{-1}(c't)).$$

Now we return to upper bounds, and we consider polynomially growing m and M . In this case we can omit the logarithmic terms in (8.4) in a similar way to Theorem 7.6.

Theorem 8.4. *Let $(T(t))_{t \geq 0}$ be a bounded C_0 -semigroup on a Hilbert space X with generator $-A$. Assume that $\sigma(A) \cap i\mathbb{R} = \{0\}$ and that there exist $\alpha \geq 1$ and $\beta > 0$ such that*

$$\|(is + A)^{-1}\| = \begin{cases} O(|s|^{-\alpha}), & s \rightarrow 0, \\ O(|s|^\beta), & |s| \rightarrow \infty. \end{cases} \quad (8.7)$$

Then

$$\|T(t)A^\alpha(I + A)^{-(\alpha+\beta)}\| = O(t^{-1}), \quad t \rightarrow \infty, \quad (8.8)$$

and

$$\|T(t)A(I + A)^{-2}\| = O(t^{-1/\gamma}), \quad t \rightarrow \infty, \quad (8.9)$$

where $\gamma = \max(\alpha, \beta)$.

Conversely, if (8.9) holds for some $\gamma > 0$, then (8.7) holds for $\alpha = \max(\gamma, 1)$ and $\beta = \gamma$.

Proof. The argument which establishes (7.7) in the proof of Theorem 7.3 (where one puts $\beta = 0$ and $\ell \equiv 1$) shows that

$$\sup \{ \|(is + A)^{-1}(A(I + A)^{-1})^\alpha\| : 0 < |s| \leq 1 \} < \infty.$$

Similarly, the argument that (i) implies (5.17) in the proof of Theorem 5.5 shows that

$$\sup \{ \|(is + A)^{-1}(I + A)^{-\beta}\| : |s| \geq 1 \} < \infty.$$

Using product and composition rules similarly to (4.4),

$$\sup \{ \|(is + A)^{-1}A^\alpha(I + A)^{-(\alpha+\beta)}\| : s \neq 0 \} < \infty.$$

Now we can apply Lemma 5.3 to the function

$$f(z) = (z + A)^{-1}A^\alpha(I + A)^{-(\alpha+\beta)},$$

and we deduce that

$$\sup \{ \|\lambda(I + A)^{-1}A^\alpha(I + A)^{-(\alpha+\beta)}\| : \lambda \in \mathbb{C}_+ \} < \infty.$$

By Theorem 4.7,

$$\|T(t)A^\alpha(I + A)^{-(\alpha+\beta)}\| = O(t^{-1}), \quad t \rightarrow \infty.$$

Since $A^{\gamma-\alpha}(I + A)^{-(2\gamma-\alpha-\beta)}$ is a bounded operator,

$$\|T(t)(A(I + A)^{-2})^\gamma\| = O(t^{-1}), \quad t \rightarrow \infty.$$

Since $A(I + A)^{-2}$ is sectorial, by Lemma 4.2

$$\|T(t)A(I + A)^{-2}\| = O(t^{-1/\gamma}), \quad t \rightarrow \infty,$$

for $\gamma = \max(\alpha, \beta)$.

The converse statement follows from Theorem 8.1. □

In Theorem 8.4, the case $\gamma > 1$ can occur only if the space X splits as described before Theorem 8.1.

Remark 8.5. Proposition 3.10 (i) and (8.8) show that $\|T(t)x\| = O(t^{-1})$ for all $x \in \text{ran}(A^\alpha) \cap \text{dom}(A^\beta)$. Moreover, Proposition 3.10 (ii) shows that $A^a(I + A)^{-1}$ is sectorial whenever $0 < a < 1$, in particular for $a = \alpha/(\alpha + \beta)$. By Lemma 4.2 one obtains that

$$\|T(t)A^{-\alpha\gamma}(I + A)^{-(\alpha\gamma+\beta\gamma)}\| = O(t^{-\gamma}), \quad t \rightarrow \infty,$$

for every $\gamma > 0$. Hence $\|T(t)x\| = O(t^{-\gamma})$ for all $x \in \text{ran}(A^{\alpha\gamma}) \cap \text{dom}(A^{\beta\gamma})$.

We do not consider regularly varying rates of resolvent growth in the context of this section. It is not easy to find a single operator which cancels resolvent growth at both zero and infinity simultaneously, without losing information about the fine scale of the behaviour.

Finally we make some remarks about quasi-multiplication semigroups. In many cases (including normal semigroups on Hilbert spaces, and multiplication semigroups on L^p -spaces) the space X can be split into a direct sum of two closed invariant subspaces X_0 and X_1 so that the generator of the semigroup is bounded when restricted to X_0 and invertible when restricted to X_1 . Then the rate of decay on X is the maximum of the rates on X_0 and X_1 , so upper and lower bounds on X can be deduced from those on X_0 and X_1 . However knowing only the rate of decay on X is not sufficient to detect whether it is controlled on X_0 and X_1 . Consequently we do not think it is possible to formulate a succinct result such as Propositions 5.1 and 6.13 in terms of $\|(is + A)^{-1}\|$ in this case. If $\|(is + A)^{-1}\|$ dominates $\|(is^{-1} + A)^{-1}\|$ whenever $|s| > 1$, then the rate of decay is determined by the behaviour on X_1 , and one can apply Proposition 5.1. If $\|(is^{-1} + A)^{-1}\|$ dominates $\|(is + A)^{-1}\|$ whenever $|s| > 1$, then the behaviour on X_0 dominates and Proposition 6.13 is applicable.

Acknowledgements. The research described in this paper was supported by the Leverhulme Trust and by the EPSRC grant EP/J010723/1. The third author was also partially supported by the NCN grant DEC-2011/03/B/ST1/00407 and by the EU IRSES grant PIRSES-GA-2012-318910, AOS.

References

- [1] F. Alabau-Boussouira, P. Cannarsa, and R. Guglielmi, *Indirect stabilization of weakly coupled systems with hybrid boundary conditions*, *Math. Control Relat. Fields* **1** (2011), 413–436.
- [2] K. Ammari, E. Feireisl, S. Nicaise, *Polynomial stabilization of some dissipative hyperbolic systems*, see arXiv:1208.4485.
- [3] K. Ammari, D. Mercier, V. Régnier, and J. Valein, *Spectral analysis and stabilization of a chain of serially connected Euler-Bernoulli beams and strings*, *Commun. Pure Appl. Anal.* **11** (2012), 785–807.
- [4] N. Anantharaman and M. Léautaud, *Decay rates for the damped wave equation on the torus*, preprint, with appendix by S. Nonnenmacher, *Anal. PDE*, to appear; see arXiv:1210.6879.
- [5] W. Arendt and C. J. K. Batty, *Tauberian theorems and stability of one-parameter semigroups*, *Trans. Amer. Math. Soc.* **306** (1988), 837–852.
- [6] W. Arendt, C. J. K. Batty, M. Hieber, and F. Neubrander, *Vector-valued Laplace Transforms and Cauchy Problems*, 2nd ed., *Monographs in Mathematics*, vol. 96, Birkhäuser, Basel, 2011.
- [7] W. Arendt and J. Prüss, *Vector-valued Tauberian theorems and asymptotic behavior of linear Volterra equations*, *SIAM J. Math. Anal.* **23** (1992), 412–448.
- [8] M. Bassam, D. Mercier, S. Nicaise, and A. Wehbe, *Stabilisation frontière indirecte du système de Timoshenko*, *C. R. Math. Acad. Sci. Paris* **349** (2011), 379–384.
- [9] A. Bátkai, K.–J. Engel, J. Prüss and R. Schnaubelt, *Polynomial stability of operator semigroups*, *Math. Nachr.* **279** (2006), 1425–1440.
- [10] C. J. K. Batty, *Tauberian theorems for the Laplace-Stieltjes transform*, *Trans. Amer. Math. Soc.* **322** (1990), 783–804.
- [11] C. J. K. Batty, *Asymptotic behaviour of semigroups of operators*, in: *Functional analysis and operator theory (Warsaw, 1992)*, *Banach Center Publ.* **30**, Polish Acad. Sci., Warsaw, 1994, pp. 35–52.
- [12] C. J. K. Batty, *Differentiability of perturbed semigroups and delay semigroups*, in: *Perspectives in operator theory*, *Banach Center Publ.* **75**, Polish Acad. Sci., Warsaw, 2007, 39–53.
- [13] C. J. K. Batty and T. Duyckaerts, *Non-uniform stability for bounded semi-groups on Banach spaces*, *J. Evol. Eq.* **8** (2008), 765–780.
- [14] C. J. K. Batty, A. Gomilko and Yu. Tomilov, *Product formulas in functional calculi*, see arXiv:1309.0318.

- [15] N. H. Bingham, C. M. Goldie, and J. L. Teugels, *Regular variation*, Encyclopedia of Mathematics and its Applications **27**, Cambridge University Press, Cambridge, 1987.
- [16] J.-F. Bony and V. Petkov, *Resolvent estimates and local energy decay for hyperbolic equations*, Ann. Univ. Ferrara Sez. VII Sci. Mat. **52** (2006), 233–246.
- [17] A. Borichev and Yu. Tomilov, *Optimal polynomial decay of functions and operator semigroups*, Math. Ann. **347** (2010), 455–478.
- [18] N. Burq, *Décroissance de l'énergie locale de l'équation des ondes pour le problème extérieur et absence de résonance au voisinage du réel*, Acta Math. **180** (1998), 1–29.
- [19] N. Burq and M. Hitrik, *Energy decay for damped wave equations on partially rectangular domains*, Math. Res. Letters **14** (2007), 35–47.
- [20] R. Chill and Yu. Tomilov, *Stability of operator semigroups: ideas and results*, in: Perspectives in operator theory, Banach Center Publ. Polish Acad. Sci. **75**, Warsaw, 2007, 71–109.
- [21] H. Christianson, *Applications of the cutoff resolvent estimates to the wave equation*, Math. Res. Letters **16** (2009), 577–590.
- [22] H. Christianson, E. Schenck, A. Vasy, and J. Wunsch, *From resolvent estimates to damped waves*, J. Anal. Math., to appear.
- [23] J.B. Conway, *Functions of one complex variable, I*. Second edition. Graduate Texts in Mathematics, **11**, Springer-Verlag, New York-Berlin, 1978.
- [24] N. Dungey, *On time regularity and related conditions for power-bounded operators*, Proc. London Math. Soc. **97** (2008), 97–116.
- [25] N. Dungey, *Difference estimates for continuous and discrete operator semigroups*, Semigroup Forum **78** (2009), 226–237.
- [26] M. Eller and P. Toundykov, *Carleman estimates for elliptic boundary value problems with applications to the stabilization of hyperbolic systems*, Evolution Equations and Control Theory **1**(2012), 271–296.
- [27] K.-J. Engel and R. Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, Graduate Texts in Mathematics **194**, Springer, New York, 2000.
- [28] J. Esterle, E. Strouse, and F. Zouakia, *Stabilité asymptotique de certains semi-groupes d'opérateurs et idéaux primaires de $L_1(\mathbb{R}^+)$* , J. Operator Theory **28** (1992), 203–227.
- [29] L. Fatori and R. Monteiro, *The optimal decay rate for a weak dissipative Bresse system*, Appl. Math. Lett. **25** (2012), 600–604.

- [30] H. Fernandez Sare and J. Munoz Rivera, *Optimal rates of decay in 2-d thermoelasticity with second sound*, J. Math. Phys. **53** (2012), 073509.
- [31] A. Gomilko, M. Haase, and Yu. Tomilov, *Bernstein functions and rates in mean ergodic theorems for operator semigroups*, J. Anal. Math. **118** (2012), 545–576.
- [32] M. Grobbelaar-van Dalsen, *Strong stabilization of models incorporating the thermoelastic Reissner-Mindlin plate equations with second sound*, Appl. Anal. **90** (2011), 1419–1449.
- [33] M. Grobbelaar-van Dalsen, *On the dissipative effect of a magnetic field in a Mindlin-Timoshenko plate model*, Z. Angew. Math. Phys. **63** (2012), 1047–1065.
- [34] M. Grobbelaar-van Dalsen, *Stabilization of a thermoelastic Mindlin-Timoshenko plate model revisited*, Z. Angew. Math. Phys. **64** (2013), 1305–1325.
- [35] M. Haase, *Spectral mapping theorems for holomorphic functional calculi*, J. London Math. Soc. (2) **71** (2005), 723–739.
- [36] M. Haase, *The Functional Calculus for Sectorial Operators. Operator Theory: Advances and Applications* **169**, Birkhäuser, Basel, 2006.
- [37] J. Hao and Z. Liu, *Stability of an abstract system of coupled hyperbolic and parabolic equations*, Z. Angew. Math. Phys. **64** (2013), 1145–1159.
- [38] Z. Harper, *Laplace transform representations and Paley-Wiener theorems for functions on vertical strips*, Doc. Math. **15** (2010), 235–254.
- [39] E. Hille and R. S. Phillips, *Functional Analysis and Semi-Groups*, 3rd printing of rev. ed. of 1957, Colloq. Publ. **31**, Amer. Math. Soc., Providence, 1974.
- [40] F. Hirsch, *Transformation de Stieltjes et fonctions opérant sur les potentiels abstraits*, in: Théorie du potentiel et analyse harmonique (Journées Soc. Math. France, Inst. Recherche Math. Avancée, Strasbourg, 1973), Lecture Notes in Math. **404**, Springer, Berlin, 1974, 149–163.
- [41] F. Hirsch, *Intégrales de résolvantes et calcul symbolique*, Ann. Inst. Fourier (Grenoble) **22** (1972), 239–264.
- [42] F. Hirsch, *Familles d'opérateurs potentiels*, Ann. Inst. Fourier (Grenoble) **25** (1975), 263–288.
- [43] F. Hirsch, *Domaines d'opérateurs représentés comme intégrales de résolvantes*, J. Funct. Anal. **23** (1976), 199–217.
- [44] N. Kalton, S. Montgomery-Smith, K. Oleszkiewicz, and Yu. Tomilov, *Power-bounded operators and related norm estimates*, J. London Math. Soc. **70** (2004), 463–478.

- [45] N. Kalton and P. Portal, *Remarks on ℓ_1 and ℓ_∞ -maximal regularity for power-bounded operators*, J. Aust. Math. Soc. **84** (2008), 345–365.
- [46] Y. Katznelson, *An introduction to harmonic analysis*, Wiley, New York, 1968.
- [47] J. Korevaar, *Tauberian Theory. A century of developments*, Springer, Berlin, 2004.
- [48] P.C. Kunstmann and L. Weis, *Maximal L_p -regularity for parabolic equations, Fourier multiplier theorems and H^∞ -functional calculus*, in: Functional analytic methods for evolution equations, Lecture Notes in Math. **1855**, Springer, Berlin, 2004, 65–311.
- [49] M. Kwaśnicki, *Spectral analysis of subordinate Brownian motions on the half-line*, Studia Math. **206** (2011), 211–271.
- [50] Yu. Latushkin and R. Shvydkoy, *Hyperbolicity of semigroups and Fourier multipliers*, in: Systems, approximation, singular integral operators, and related topics (Bordeaux, 2000), Oper. Theory Adv. Appl., **129**, Birkhäuser, Basel, 2001, 341–363.
- [51] G. Lebeau, *Équation des ondes amorties*, in: Algebraic and geometric methods in mathematical physics (Kaciveli, 1993), Math. Phys. Stud. **19**, Kluwer Acad. Publ., Dordrecht, 1996, 73–109.
- [52] G. Lebeau and L. Robbiano, *Stabilisation de l'équation des ondes par le bord*, Duke Math. J. **86** (1997), 465–491.
- [53] Z. Liu and B. Rao, *Characterization of polynomial decay rate for the solution of linear evolution equation*, Z. Angew. Math. Phys. **56** (2005), 630–644.
- [54] C. Martínez Carracedo and M. Sanz Alix, *The theory of fractional powers of operators*, North-Holland Mathematics Studies **187**, North-Holland, Amsterdam, 2001.
- [55] M. Martínez, *Decay estimates of functions through singular extensions of vector-valued Laplace transforms*, J. Math. Anal. Appl. **375** (2011), 196–206.
- [56] J. M. A. M. van Neerven, *The asymptotic behaviour of semigroups of linear operators*, Operator Theory: Advances and Applications **88**, Birkhäuser, Basel, 1996.
- [57] O. Nevanlinna, *Convergence of iterations for linear equations*, Lectures in Mathematics, ETH Zürich, Birkhäuser, Basel, 1993.
- [58] S. Nicaise, *Stabilization and asymptotic behavior of dispersive medium models*, Systems Control Lett. **61** (2012), 638–648.
- [59] S. Nicaise and J. Valein, *Stabilization of non-homogeneous elastic materials with voids*, J. Math. Anal. Appl. **387** (2012), 1061–1087.
- [60] N. Okazawa, *Logarithms and imaginary powers of closed linear operators*, Integral Equations Operator Theory **38** (2000), 458–500.

- [61] E. I. Pustyl'nik, *Some inequalities in the theory of self-adjoint operators*, Izv. Uchebn. Zaved. Mat. **1979**, 52–57 (in Russian).
- [62] E. I. Pustyl'nik, *Functions of a positive operator*, Mat. Sb. (N.S.) **119** (1982), 32–47 (in Russian), Math. USSR Sbornik **47**(1984), 27–42.
- [63] M. L. Santosa, D. S. Almeida Júniora, and J.E. Munoz Rivera, *The stability number of the Timoshenko system with second sound* J. Diff. Eq. **253** (2012), 2715–2733.
- [64] R. Schilling, R. Song, and Z. Vondraček, *Bernstein functions*, de Gruyter Studies in Mathematics **37**, Walter de Gruyter, Berlin, 2010.
- [65] E. Seneta, *Regularly varying functions*, Lecture Notes in Math. **508**, Springer, Berlin, 1976.
- [66] L. Tebou, *Energy decay estimates for some weakly coupled Euler-Bernoulli and wave equations with indirect damping mechanisms*, Math. Control Relat. Fields **2** (2012), 45–60.
- [67] Vu Quôc Phóng, *Theorems of Katznelson-Tzafriri type for semigroups of operators*, J. Funct. Anal. **103** (1992), 74–84.
- [68] D. V. Widder, *The Laplace Transform*, Princeton Mathematical Series **6**, Princeton University Press, Princeton, 1941.