

ON THE RECOGNITION OF RIGHT-ANGLED ARTIN GROUPS

MARTIN R. BRIDSON

ABSTRACT. There does not exist an algorithm that can determine whether or not a group presented by commutators is a right-angled Artin group.

1. INTRODUCTION

In [4] Day and Wade introduced an elegant new homology theory for subspace arrangements and related it to recognition problems concerning *right-angled Artin groups* (RAAGs). In setting the context for their work, they asked if there is an algorithmic procedure for recognizing RAAGs among groups given by presentations whose only relations are commutators ([4] Question 1.2) and speculated that the answer was likely to be no. The purpose of this note is to confirm this speculation.

Theorem 1.1. *There does not exist an algorithm that can determine whether or not a group presented by commutators is a RAAG.*

In more detail, there is no algorithm that, given 22 words u_i in the free group $F(a_1, a_2, a_3, a_4)$ can determine whether or not the group with presentation

$$\langle a_1, a_2, a_3, a_4, t \mid [a_1, a_3], [a_1, a_4], [a_2, a_3], [a_2, a_4], [t, u_1], \dots, [t, u_{22}] \rangle$$

is a RAAG. Nor is there an algorithm that can determine whether or not such a group is commensurable with a RAAG or quasi-isometric to a RAAG.

2. FIBRE PRODUCTS AND TRIVIALITY FOR 2-GENERATOR GROUPS

In the aftermath of the construction by Novikov [9] and Boone [2] of finitely presented groups with unsolvable word problem, many other decision problems for groups were proved to be unsolvable through subtle work by many authors. We shall appeal to two results that come from the work of C.F. Miller III.

Let K be a group given by a presentation with n generators and m relations. Following Miller, one can associate to each word w in the generators of K a presentation with 2 generators and $n + m + 3$ relations – this is derived from the presentation in Lemma 3.6 of [7] by making Tietze moves to remove unnecessary generators. The group given by this presentation is trivial if $w = 1$ in K , but it contains K if $w \neq 1$.

The most concise finite presentation that is known for a group with unsolvable word problem is the one constructed by Borisov [3] half a century ago – it has 5 generators and 12 relations. By applying Miller’s construction to words in the

1991 *Mathematics Subject Classification.* 20F36, 20F10.

generators of Borisov's example, we obtain a recursive sequence of 2-generator presentations $\mathcal{P}_n = \langle a_1, a_2 \mid S_n \rangle$ with $|S_n| = 20$ such that the group presented is either trivial or else has an unsolvable word problem, and there is no algorithm that can determine the set of integers n for which each alternative holds.

We exploit these examples in the manner of Mihailova [8] and Miller ([6] p.39).

Lemma 2.1. *Let F be a free group of rank 2 with generators $\{a_1, a_2\}$. If $k \geq 20$, then there exists a recursive sequence (S_n) of subsets of F , each of cardinality k , such that there is no algorithm to determine whether or not $F \times F$ is generated by $U_n = \{(a_1, a_1), (a_2, a_2), (s, 1) : s \in S_n\}$. Moreover, if $\langle U_n \rangle$ is a proper subgroup then $H_2(\langle U_n \rangle, \mathbb{Z})$ is not finitely generated and there is no algorithm to determine which words in the generators of $F \times F$ determine elements of $\langle U_n \rangle$.*

Proof. Associated to any 2-generator finite presentation $\mathcal{P} = \langle a_1, a_2 \mid r_1, \dots, r_M \rangle$ one has the fibre product $P < F \times F$ consisting of pairs (u, v) such that $u = v$ in the group $G(\mathcal{P})$ presented by \mathcal{P} . It is easy to check that P is generated by $U = \{(a_1, a_1), (a_2, a_2), (r_1, 1), \dots, (r_M, 1)\}$. If $G(\mathcal{P})$ is trivial, $P = F \times F$. But if $G(\mathcal{P})$ is infinite, Baumslag and Roseblade's Theorem A [1] shows that $H_2(P, \mathbb{Z})$ is not finitely generated. Moreover, if the word problem is unsolvable in $G(\mathcal{P})$, the membership problem of $P < F \times F$ is unsolvable, because deciding if $(w, 1) \in P$ is equivalent to deciding if $w = 1$ in $G(\mathcal{P})$. Consideration of the presentations $\mathcal{P}_n = \langle a_1, a_2 \mid S_n \rangle$ from the discussion preceding the lemma completes the proof. \square

We shall apply the following lemma with $D = F \times F$ and $C = \langle U_n \rangle$.

Lemma 2.2. *For any HNN extension of the form $\Gamma = D *_C$, if $H_2(D, \mathbb{Z})$ is finitely generated but $H_2(C, \mathbb{Z})$ is not, then $H_3(\Gamma, \mathbb{Z})$ is not finitely generated.*

Proof. The Mayer-Vietoris sequence for the HNN extension contains the exact sequence

$$H_3(\Gamma, \mathbb{Z}) \rightarrow H_2(C, \mathbb{Z}) \rightarrow H_2(D, \mathbb{Z}).$$

\square

3. PROOF OF THEOREM 1.1

Given 20 words $S = \{r_1, \dots, r_{20}\}$ in the free group $F = F(a_1, a_2)$, we denote by $\Gamma(S)$ the group with generators a_1, a_2, a_3, a_4, t and relations

$$[a_1, a_3], [a_1, a_4], [a_2, a_3], [a_2, a_4], [t, a_1 a_3], [t, a_2 a_4], [t, r_1], \dots, [t, r_{20}].$$

Note that this presentation is of the type described in Theorem 1.1. The group $\Gamma(S)$ is an HNN extension of $F \times F$ with a stable letter t that commutes with the fibre product $P < F \times F$ associated to the presentation $\mathcal{P} = \langle a, b \mid S \rangle$.

As in the proof of Lemma 2.1, we have a dichotomy: if $G(\mathcal{P}) = 1$ then $\Gamma(S) = F \times F \times \mathbb{Z}$ is a RAAG; but if $G(\mathcal{P})$ is infinite then $\Gamma(S)$ is not a RAAG, because RAAGs have finite classifying spaces [5] whereas $H_3(\Gamma(S), \mathbb{Z})$ is not finitely generated, by Lemma 2.2, because $H_2(P, \mathbb{Z})$ is not finitely generated.

Lemma 2.1 tells us that there is no algorithm to determine which of the possibilities in this dichotomy holds. Moreover, by choosing sets (S_n) as in the lemma, we can arrange that when $\Gamma(S_n)$ is not equal to $F \times F \times \mathbb{Z}$, it will have an unsolvable word problem: given a word w in the generators a_1, a_2 , Britton's Lemma implies that $[t, w] = 1$ in $\Gamma(S_n)$ if and only if $w = 1$ in $G(\mathcal{P}_n)$, and this is undecidable.

Since being of type FP_3 and having a solvable word problem are both invariants of commensurability and quasi-isometry, there is no algorithm that can determine whether $\Gamma(S_n)$ is commensurable with or quasi-isometric to a RAAG. \square

REFERENCES

- [1] G. Baumslag and J. Roseblade, *Subgroups of direct products of free groups*, J. London Math. Soc. (2) **30** (1984) 44–52.
- [2] W.W. Boone, *Certain simple unsolvable problems in group theory*, I, II, III, IV, V, VI, Nederl. Akad. Wetensch. Proc. **57** (1954) 231–237, 492–497; **58** (1955) 252–256, 571–577; **60** (1957) 22–27, 227–232.
- [3] V. Borisov, *Simple examples of groups with unsolvable word problem*, Math. Zametki **6** (1969) 521–532; English transl., Math. Notes **6** (1969) 768–775.
- [4] M. Day and R. Wade, *Subspace arrangements, BNS invariants, and pure symmetric outer automorphisms of right-angled Artin groups*, Groups Geom. Dyn. **12** (2018) 173–206.
- [5] K.H. Kim and F.W. Roush, *Homology of certain algebras defined by graphs*, J. Pure Appl. Algebra **17** (1980) 179–186.
- [6] C.F. Miller III, *On group-theoretic decision problems and their classification*, Annals of Mathematics Studies, No. 68, Princeton University Press, 1971.
- [7] C.F. Miller III, *Decision problems for groups: survey and reflections*, In: “Algorithms and Classification in Combinatorial Group Theory”, G. Baumslag, C. F. Miller III (eds), MSRI Publications No. 23, Springer-Verlag, 1992, pp. 1–59.
- [8] K.A. Mihailova, *The occurrence problem for direct products of groups*, Dokl. Akad. Nauk SSSR **119** (1958) 1103–1105.
- [9] P.S. Novikov, *On the algorithmic unsolvability of the word problem in group theory*, Trudy Mat. Inst. Steklov **44** (1955) 1–143.

MATHEMATICAL INSTITUTE, ANDREW WILES BUILDING, OXFORD OX2 6GG, EU
E-mail address: `bridson@maths.ox.ac.uk`