



Focal precentral gyrus involvement in osmotic demyelination

Martin R Turner,¹ John Jacob,^{1,2} Lucy Matthews,³ Fintan Sheerin³

¹Nuffield Department of Clinical Neurosciences, Oxford University, John Radcliffe Hospital, Oxford, UK

²Department of Neurology, Milton Keynes University Hospital, Milton Keynes, UK

³Department of Neurology, John Radcliffe Hospital, Oxford University Hospitals NHS Foundation Trust, Oxford, Oxfordshire, UK

Correspondence to

Professor Martin R Turner, Nuffield Department of Clinical Neurosciences, John Radcliffe Hospital, Oxford University, OX3 9DU, Oxford, UK; martin.turner@ndcn.ox.ac.uk

Accepted 24 July 2017

Published Online First

19 August 2017




Figure 1 MR scan of the brain. Axial fluid-attenuated inversion recovery (FLAIR) images show symmetrical pontine (A), basal ganglia (B) and a 'ribbon' of cortical T2 hyperintensity in the posterior precentral gyri (C). The precentral gyral changes are strikingly similar to a focal pattern of MRI changes described in cases of amyotrophic lateral sclerosis.

THE CASE

A previously well 59-year-old woman was admitted to a regional hospital with subacute encephalopathy and found to have severe hyponatraemia (101 mmol/L). She had taken indapamide for hypertension over the preceding 3 months. Before referral to neurological services, she had received intravenous hypertonic saline. Her serum sodium concentration normalised within a few days, but was associated with a deteriorating conscious level. She was referred to our tertiary neurological centre with suspected osmotic demyelination where she presented in a 'locked in' state, involving quadriplegia and absent vertical eye movements. MR scan of the brain showed typical symmetrical

pontine and basal ganglia changes, but also focal hyperintensity of the precentral gyri (figure 1). Nearly 6 months after the initial admission to hospital, the patient had made some recovery of limb strength, but with persistent loss of fine dexterity and unable to stand independently.

Hyponatraemia secondary to thiazide diuretics is well recognised, and cautious correction of chronic hyponatraemia is generally advised to minimise the risk of osmotic demyelination. This unusual extension of MRI changes in extrapontine myelinolysis is uncommon,¹ and our case also shared the bilateral hyperintensity in the posterolateral nuclei of the



To cite: Turner MR, Jacob J, Matthews L, et al. *Pract Neurol* 2017;**17**:410–411.

thalami. The strikingly selective involvement of the posterior border of the precentral gyrus is more commonly described in cases of amyotrophic lateral sclerosis in relation to upper motor neurone degeneration,^{2 3} where iron deposition has been inferred. Corticospinal tract MRI white matter changes occurring in patients with fulminant hepatic failure also overlap in their appearance with those seen in cases of amyotrophic lateral sclerosis.⁴ These observations might reflect a hitherto unrecognised selective anatomical connectivity of subcortical structures with this region of the motor cortex, or a shared focal metabolic susceptibility.

Contributors MRT conceived and drafted the manuscript. JJ, LM and FS edited the manuscript.

Disclaimer MRT, JJ, LM and FS report no disclosures.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed. This paper was reviewed by Roswell Martin, Gloucester, UK.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Harsha KJ, Parameswaran K. Newly described additional sites of extrapontine myelinolysis along with typical pontine and extrapontine myelinolysis. *Neurol India* 2017;65:181–2.
- 2 Hecht MJ, Fellner F, Fellner C, *et al.* Hyperintense and Hypointense MRI signals of the Precentral gyrus and Corticospinal tract in ALS: a follow-up examination including FLAIR images. *J Neurol Sci* 2002;199:59–65.
- 3 Costagli M, Donatelli G, Biagi L, *et al.* Magnetic susceptibility in the deep layers of the primary motor cortex in Amyotrophic Lateral Sclerosis. *Neuroimage Clin* 2016;12:965–9.
- 4 Turner MR. Response to: Reversible diffusion MRI abnormalities and transient mutism after liver transplantation. *Neurology* 2005;64:177.



Focal precentral gyrus involvement in osmotic demyelination

Martin R Turner, John Jacob, Lucy Matthews and Fintan Sheerin

Pract Neurol 2017 17: 410-411 originally published online August 19, 2017

doi: 10.1136/practneurol-2017-001682

Updated information and services can be found at:
<http://pn.bmj.com/content/17/5/410>

These include:

References

This article cites 4 articles, 0 of which you can access for free at:
<http://pn.bmj.com/content/17/5/410#BIBL>

Open Access

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See:
<http://creativecommons.org/licenses/by/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections
[Open access](#) (33)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>